1

The diazepam binding inhibitor's modulation of the GABA-A receptor is subunit-dependent

Jennifer S. Borchardt[1], Lucas M. Blecker[2], Kenneth A. Satyshur[2], and Cynthia Czajkowski[2]*

[1] Cellular and Molecular Biology Graduate Program, University of Wisconsin-Madison, Madison, WI, United States of America

[2] Department of Neuroscience, University of Wisconsin-Madison, Madison, WI, United States of America

* Corresponding author

Email: cmczajko@wisc.edu

Abstract

First synthesized in the 1950s, benzodiazepines are widely prescribed drugs that exert their anxiolytic, sedative and anticonvulsant actions by binding to GABA-A receptors, the main inhibitory ligand-gated ion channel in the brain. Scientists have long theorized that there exists an endogenous benzodiazepine, or endozepine, in the brain. While there is indirect evidence suggesting a peptide, the diazepam binding inhibitor, is capable of modulating the GABA-A receptor, direct evidence of the modulatory effects of the diazepam binding inhibitor is limited.

Here we take a reductionist approach to understand how purified diazepam binding inhibitor interacts with and affects GABA-A receptor activity. We used two-electrode voltage clamp electrophysiology to study how the effects of diazepam binding inhibitor vary with GABA-A receptor subunit composition, and found that GABA-evoked currents from α3-containing GABA-A receptors are weakly inhibited by the diazepam binding inhibitor, while currents from α5-containing receptors are positively modulated. We also used *in silico* protein-protein docking to visualize potential diazepam binding inhibitor/GABA-A receptor interactions that revealed diazepam binding inhibitor bound at the benzodiazepine α/γ binding site interface, which provides a structural framework for understanding diazepam binding inhibitor effects on GABA-A receptors. Our results provide novel insights into mechanisms underlying how the diazepam binding inhibitor modulates GABA-mediated inhibition in the brain.

Introduction

Since their discovery over fifty-five years ago, benzodiazepines (BZDs) have been some of the most widely prescribed drugs in the world, and have remained on the World Health Organization's list of essential drugs since the list's inception in 1977 [1]. BZDs are used to treat a variety of conditions, ranging from epilepsy to insomnia to anxiety [2-4], and mediate their

effects by binding to the main inhibitory neurotransmitter receptor in the brain, the gamma-

aminobutyric acid-A receptor (GABAR) [5]. BZDs modulate GABAR activity, and thus alter

neuronal signaling.

GABARs are heteropentamers comprised from nineteen possible subunits (α1-6, β1-3,

γ1-3, δ, ε, θ, π and ρ1-3) that are expressed in distinct brain regions, with distinct

pharmacological properties. There are two GABA binding sites in the extracellular interfaces

between β and α subunits, while BZDs bind in an extracellular pocket at the α/γ subunit

interface. BZDs, such as diazepam or flurazepam (FZM), can act as positive allosteric

modulators (PAMs), which increase GABA-mediated currents, or negative allosteric modulators

(NAMs) like methyl-6,7-dimethoxy-4-ethyl-beta-carboline-3-carboxylate (DMCM) which

decrease GABA-elicited currents. Zero-modulators, such as flumazenil (FLZ, Ro15-1788),

occupy the BZD site but do not alter GABA-elicited currents. The effects of some BZDs like

Ro15-4513 depend on GABAR subunit composition. Ro15-4513 acts as PAM when interacting

with $\alpha_4\beta_2\gamma_2$ and $\alpha_6\beta_2\gamma_2$ GABARs, but is a NAM when bound to $\alpha_1\beta_2\gamma_2$ receptors[6].

The presence a binding site on the GABAR for synthetically manufactured BZDs has

long suggested the existence of an endogenous synthesized molecule that binds to the same

site. In 1983, a candidate peptide was isolated from rat brain homogenates, which displaced

tritiated diazepam in a radioligand binding assay [7]. This 10kD, 87 amino-acid protein was

named the diazepam binding inhibitor (DBI). Recent DBI knock-down and over-expression

studies in the thalamic reticular nucleus demonstrate that DBI works *in vivo* as a PAM where it

potentiates GABA-mediated currents and suppresses epileptic activity[8]. In contrast, DBI

knock-down in the subventricular zone of the lateral ventricles and in the hippocampal

subgranular zone demonstrate that DBI and one of its peptide fragments, ODN, inhibit GABA-

induced currents [9, 10], indicating that in these brain regions DBI and ODN work as NAMs of

GABARs. We hypothesized that the differences in DBI's modulatory effects are due to regional

differences in GABAR subunit expression and GABAR subunit composition.

Here, we used two-electrode voltage clamping and measured DBI effects on GABA-

activated currents from GABARs expressed in *Xenopus laevis* oocytes. Given DBI is highly

expressed in the hippocampus and thalamus [11, 12], we focused on $\alpha_5\beta_3\gamma_{2L}$ and $\alpha_3\beta_3\gamma_{2L}$

GABAR subtypes which are enriched in these regions [13]. We found that DBI positively

modulates $\alpha_5\beta_3\gamma_{2L}$ receptors, whereas DBI is a weak negative modulator of $\alpha_3\beta_3\gamma_{2L}$ GABARs. We

used *in silico* protein-protein docking to visualize potential DBI-GABAR interactions.

Materials and Methods

DBI expression and purification for functional testing

We used cDNA for human DBI isoform3 (GeneBank ID CAG33237.1) with a C-terminal

histidine tag in a pET28a vector. The plasmid was chemically transformed into BL21 DE3

competent cells, plated on LB-Kanamycin (Kan) agar plates, and grown overnight at 37°C. A

single colony was picked and grown overnight in a 5mL culture of LB with 250µg Kan while

shaking at 37°C. This culture was used to inoculate 500mL LB + Kan and the culture was grown

until OD_{600} reached 0.6-0.8. Protein expression was induced with IPTG (500µM) for 4 hours at

37°C. After induction, cells were pelleted by spinning at 10,000xg for 10min at 4°C. The

bacterial cell pellet was frozen at -80°C. Cell pellet was thawed and resuspended in 50mL

Buffer A (sodium phosphate, 50mM, pH 7.4; NaCl, 300mM) and 25mM imidazole (IDA) with a

protease inhibitor cocktail (pepstatin, aprotinin, leupeptin). After lysis using an Emulsiflex (ATA

Scientific, Taren Point, NSW Australia), samples were spun at 16800xg for 20min, 4°C. The

supernatant was bound to an Ni-NTA column which was pre-equilibrated in column buffer

(Buffer A and 10mM imidazole, pH 7.4). His-tagged DBI was eluted using elution buffer (Buffer

A and 250mM imidazole). A PD10 desalting column was used to remove imidazole, DBI was

eluted in Buffer A. DBI molecular weight was verified via 15% SDS-PAGE stained with

Coomasie blue and protein level quantified using a Nanodrop 2000 (Thermo Fisher Scientific,

Waltham, MA), using an absorption coefficient of 1.68. DBI with the histidine tag attached has a

mass of 10.87kDa. All data shown used purified His-tagged DBI, whose effects were similar to

untagged DBI (data not shown).

DBI purification for NMR

DBI was transformed into BL21 DE3 cells, grown in [15]N-labeled M9 media, and purified

as described above. A ^{1}H-^{15}N HSQC spectrum was obtained using a Bruker Avance III 600 MHz

spectrometer, with 5-10% D_2O added to the protein sample. The spectrum was collected at

25°C. Data were processed using NMRPipe [14] and analyzed using CcpNmr analysis [15].

Expression in *Xenopus laevis* oocytes

Rat cDNA encoding the GABAR α3, α5, β3 and γ2L subunits subcloned into the pUNIV

vector [16] were used. cRNA was transcribed from NotI-digested cDNA using the mMessage T7

kit (Ambion, Austin, TX). *Xenopus laevis* oocytes were harvested and prepared as described

previously [17]. Oocytes were injected with 27-54nL of GABAR subunit cRNA at 10ng/μL α,

10ng/μL β and 100ng/μL γ2L following previously described methods [18], stored at 16°C in

ND96 buffer (96mM NaCl, 2mM KCl, 1mM MgCl2, 1.8mM CaCl2, and 5mM HEPES, pH 7.2)

supplemented with 100μg/ml bovine serum albumin and gentamycin and recorded from 2-7

days after injection.

Two electrode voltage clamp

Electrophysiological recordings were performed as described previously [19]. Oocytes

expressing GABARs were held at -40 to -80 mV under a two-electrode voltage clamp and

continuously perfused with ND96 at 5mL/min in a 200 μL volume chamber. Boroscilicate glass

electrodes (0.4-1.0mΩ, Warner Instruments, Hamden, CT) were filled with 3M KCl. Data were

collected at room temperature using two different volatage clamps. One was a GeneClamp 500

(Molecular Devices, Sunnyvale, CA) interfaced to a computer via a Digidata 1200 device

(Molecular Devices) with data recorded using Whole Cell Program, version 3.6.7 (J. Dempster,

University of Strathclyde, Glasgow, UK). The other was an Oocyte Clamp OC-725A (Warner

Instruments) with a Digidata 1440A (Molecular Devices) with data recorded using AxoScope,

pCLAMP 10 (Molecular Devices). Data and results obtained from the clamps were

indistinguishable.

A stock solution of 1 M GABA (Sigma-Aldrich, St Louis, MO) was made in water, stored

at -20°C and thawed before use. GABA dilutions for experiments were prepared fresh daily in

ND96. A stock solution of 10 mM FZM (Sigma/RBI, Natick, MA) in water was diluted in ND96 for

working concentrations daily. A stock solution of 10mM DMCM (Sigma/RBI, Natick, MA) was

prepared in dimethyl sulfoxide and diluted daily in ND96 for working concentrations in which the

final concentration of dimethyl sulfoxide (≤0.1%) did not affect GABAR function.

129 Concentration-response analysis

GABA concentration-response curves were determined as described previously [19].

Seven concentrations of GABA (1µM, 3µM, 10µM, 30µM, 100µM, 1mM and 10mM) were used

to determine GABA EC50 values. Concentration-response data were fit using Prism version 8.4

(GraphPad Software Inc., San Diego, CA) to the equation: $I = Imax/[1 + (EC50/[A]nH)]$, in which

I is the peak current response to a given GABA concentration, Imax is the maximal amplitude of

GABA activated current, EC50 is the GABA concentration that produces the half-maximal

response, [A] is test GABA concentration, and nH is the Hill coefficient.

137 Drug modulation

Drug modulation was measured at GABA EC_{20} and evaluated as $I_{GABA+Drug}/I_{GABA}$, where

$I_{GABA+Drug}$ is the GABA-mediated current in the presence of drug and I_{GABA} is the GABA-mediated

current in the absence of drug. To measure drug modulation, a brief 5 sec application of GABA

EC_{20} was applied and followed immediately by a 5 sec application of GABA EC_{20}+Drug. When

multiple drugs were applied to a single oocyte, GABA EC_{20} was applied following drug treatment

until current amplitudes returned to initial GABA EC_{20} level to ensure complete washout of drugs

between different drug treatments.

Protein-protein docking

We made $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABAR homology models based on the cryoEM structure

of the $\alpha_1\beta_3\gamma_{2L}$ GABAR obtained in the presence of the BZD alprazolam (PDB 6HUO) [20]. The

homology models were built on Sybyl (Tripos Inc). The GABAR α3 or α5 sequences were

manually threaded on to the cryoEM model of the $\alpha_1\beta_3\gamma_{2L}$ GABAR based on their sequence

alignment with the α1 subunit using the graphical interface of the Sybyl program. After resolving

steric clashes, either manually or by using focused minimization, the resulting structures were

subject to whole molecule energy minimization using the Tripos force field. Gasteiger-Huckel

charges were added to the atoms, and the dielectric function set at 1.0. After Simplex initial

minimization, the Powell function in Tripos was used to minimize to a termination gradient of

0.05 kcal/(mol x angstrom). The GABAR model images were developed using PyMOL

(Schrödinger, LLC, New York). BZD alprazolam was removed from models prior to docking.

Protein-protein interactions between the $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABAR models and the

unliganded human DBI crystal structure (PDB 2FJ9) [21] were identified using ClusPro, a web-

based program for computational docking of proteins [22-24]. ClusPro analyzes 70,000 rotations

of the ligand and generates up to 30 highly populated clusters of ligand structures docked in

similar locations on the receptor with low energy scores. Clusters are ranked by population size

rather than an energy score. Docking clusters selected for analysis were based on their

proximity to the BZD binding site. We analyzed the hydrogen bonding between the GABAR

alpha subunits and DBI and the GABAR gamma subunit and DBI using Pymol's hydrogen

bonding feature.

Statistical analysis

All data were from at least three different oocytes from at least two different frogs. Data are represented as mean ± SD. Significant differences in drug modulation between subunit compositions were calculated via Kruskal-Wallis test with a Dunn's multiple comparisons (Prism v8.4, GraphPad Software Inc, San Diego, CA). This test was selected due to the nonparametric distribution of data. Statistical differences in GABA+drug versus GABA alone currents elicited from the same oocyte were calculated using a paired t-test using Prism software.

The effects of DBI on $\alpha_5\beta_3\gamma_{2L}$ GABA-elicited currents were more variable than effects observed with FZM, a BZD (coefficient of variation for FZM $\alpha_5\beta_3\gamma_{2L}$ is 17.2%, DBI is 36.0%). The variation was not correlated with DBI purification batch nor oocyte/frog. We hypothesize that flexibility and conformation dynamics of DBI (11kDa) may contribute to the increased variability of its effects as compared to a small BZD drug.

Results

In order to study the effects of DBI on the GABAR receptor, we needed milligram quantities of pure, non-aggregated DBI folded predominantly in a single conformation. We used a histidine-tagged human DBI construct (His-DBI) inserted into a pET28A vector, which is optimized for inducible expression of protein in bacterial cells [25]. We expressed the His-DBI in BL21 *E. coli* cells and purified the protein as described in Methods. DBI purity and and size was evaluated using 15% SDS-PAGE (Fig 1A). Following purification, a single coomassie blue stained band was observed with a molecular weight of 11kDa. DBI with the histidine tag attached has a mass of 10.87kDa. Heteronuclear single quantum correlation (HSQC) nuclear magnetic resonance (NMR) of ^{15}N-labeled DBI demonstrated that our DBI purification process yielded pure, non-aggregated and folded protein in a single conformation, as the number of

peaks were very close to the total number of residues in the His-tagged DBI, spectral dispersion

was good, and the linewidths were relatively uniform (Fig 1B).

Fig 1 - Purified DBI is non-aggregated and folded in a single conformation. (A) 15% SDS-

PAGE showing coomassie blue stained molecular weight marker proteins at 25, 20, 15 and

10kDa, and purified histidine-tagged DBI running at its expected size 11kD. (B) ^{1}H-^{15}N HSQC

spectrum from DBI grown in ^{15}N M9 media shows close to expected number of peaks (93

expected, 96 detected), good spectral dispersion and low noise, indicating purified DBI is folded

and non-aggregated.

200   We expressed $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs in *Xenopus laevis* oocytes and used two-

electrode voltage-clamping to measure and compare the effects of a BZD PAM (FZM), a BZD

NAM (DMCM) and DBI on GABA-elicited currents. Initially, we measured GABA concentration

responses from oocytes expressing $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs. GABA EC$_{50}$ values were

47±11.8 µM, n=5 for $\alpha_3\beta_3\gamma_{2L}$ GABARs (Fig 2A) and 34±8.6 µM, n=3 for $\alpha_5\beta_3\gamma_{2L}$ GABARs (Fig

2B), which are consistent with previously published results [26, 27].

Fig 2 – GABA dose response curves for $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs. Current responses

from oocytes expressing $\alpha_3\beta_3\gamma_{2L}$ (A) and $\alpha_5\beta_3\gamma_{2L}$ (B) GABARs were fit by non-linear regression

as described in Methods. Data were normalized to maximal GABA current obtained from curve

fits and averaged from three to five different oocytes, with error bars showing SD.

212   To evaluate and compare the effects of the positive BZD modulator, FZM, the negative

BZD modulator, DMCM and DBI, we applied a low GABA EC20 concentration and then co-

applied FZM (10µM), DMCM (10µM) or DBI (50µM) with GABA EC20 (Fig 3A). FZM (10µM)

significantly increased GABA currents from oocytes expressing $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs

2.7- and 2.1 fold, respectively (Fig. 3; paired t-test p<0.0001 for both GABAR subtypes). The

effects of FZM on α3-containing receptors compared to α5-containing receptors were not

significantly different (Fig 3B, Dunn's p>0.999). DMCM (10µM) significantly inhibited GABA

currents from both $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs (Fig 3) and the effects of DMCM on α3-

containing versus α5-containing GABARs were not significantly different (Fig 3B, Dunn's

p>0.999).

Fig 3 – DBI modulates $\alpha_3\beta_3\gamma_{2L}$ differently than $\alpha_5\beta_3\gamma_{2L}$ GABARs. (A) Sample current traces

from oocytes expressing $\alpha_3\beta_3\gamma_{2L}$ (top) and $\alpha_5\beta_3\gamma_{2L}$ (bottom) GABARs showing co-application of

FZM (10µM) with GABA EC20 increased GABA current and co-application of DMCM (10µM)

decreased current. DBI (50µM) slightly decreased currents from $\alpha_3\beta_3\gamma_{2L}$ GABARs and increased

currents from $\alpha_5\beta_3\gamma_{2L}$ GABARs. (B) Effects of FZM, DMCM and DBI are plotted as

$I_{Drug+GABA}/I_{GABA}$. Data from individual occytes are shown as filled circles, bar graphs represent

mean +/- SD. Dotted line at 1 represents no effect of drug. DBI effects on $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$

GABARs are significantly different from each other, while the effects of FZM and DMCM are

similar for both GABAR subtypes (Kruskal-Wallis test). (C) Table displaying mean +/- SD of

FZM, DMCM and DBI effects on currents from $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABAR, n = number of

oocytes. Currents in the presence of drug are significantly different than GABA currents alone,

(paired t-tests p<0.0001).

We used the same methods to evaluate the effects of purified DBI (50µM) on GABA

EC20 currents from $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs. DBI weakly inhibited GABA currents from

$\alpha_3\beta_3\gamma_2$ GABARs (Fig. 3). Current amplitudes in the presence of DBI were significantly different

from currents elicited by GABA alone (Fig. 3C, paired t-test p<0.0001). In contrast, DBI

significantly potentiated GABA currents from $\alpha_5\beta_3\gamma_{2L}$ GABARs 1.3-fold (Fig 3, paired t-test

p<0.0001). The data indicate that DBI acts as a very weak NAM for α3-containing receptors and

a PAM for α5-containing receptors (Fig 3).

α3 and α5 are homologous GABAR subunit isoforms with many conserved residues

(73% identity) [28]. To explore potential mechanisms underlying DBI's actions on α3-containing

versus α5-containing GABARs, we constructed $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABAR homology models

and examined protein-protein interactions with DBI using ClusPro, a web-based program for

computational docking of proteins [22-24]. This program generates clusters which represent

populations of structures where the ligand has docked with low energy at similar locations on

the protein. Clusters are ranked based on population size.

For both $\alpha_3\beta_3\gamma_{2L}$ and $\alpha_5\beta_3\gamma_{2L}$ GABARs , ClusPro generated many highly populated

clusters with DBI bound at the BZD α/γ binding site interface (Fig.4). A flexible loop between the

second and third α helices of DBI penetrates under Loop C of the BZD binding site. Residues in

this loop code for a DBI proteolytic peptide call octadecaneuropeptide (ODN) that has been

shown to displace [3]H-BZD binding [29]. We selected two clusters, which showed the closest

ODN-BZD site interactions, for further analysis. Fig 4A shows DBI docked at GABAR α_3/γ_{2L} and

α_5/γ_{2L} subunit interfaces. Fig 4B highlights differences in residues in the α3 and α5 GABAR

subunits, located at the BZD binding site interface, that are within H-bonding distance to DBI.

Fig 4C highlights differences in DBI residues that interact with the α3 and α5 GABAR subunits.

Residues in the GABAR γ2L subunit interacting with DBI and residues in DBI interacting with

the γ2L subunit are displayed in Figs. 4D and E, respectively.

Fig 4 – Computational docking reveals DBI can bind at the GABAR α/γ BZD binding site

interface. (A) Side views of α_3/γ_{2L} and $\alpha_5\gamma_{2L}$ GABAR subunits (α-yellow, γ-magenta) from

homology models based on $\alpha_1\beta_3\gamma_{2L}$ cryoEM model (PDB 6HUO) showing DBI in blue (PDB

2FJ9) bound at the BZD α/γ binding site interface with flexible ODN loop under Loop C of the

BZD binding site. *In silico* dockings performed using ClusPro as described in Methods. B)

Extracellular domains of α3 and α5 GABAR subunits with residues predicted to H-bond to DBI

shown in stick and colored red. α3 and α5 GABAR sequence alignments shown below, with H-

bonding residues highlighted in red and non-identical residues underlined. C) DBI with residues

predicted to H-bond to $α_3$ and $α_5$ subunits shown in stick and colored in green. DBI sequences

shown below with H-bonding residues colored green. D) Extracellular domain of γ GABAR

subunit with residues predicted to H-bond to DBI shown in stick and colored blue. E) DBI with

residues predicted to H-bond to γ subunit shown in stick and colored grey. DBI sequences

shown below. For the highlighted hydrophobic residues such as DBI G46, M47, and L48, H-

bonding is via the backbone, not the residue side-chain.

When comparing DBI bound to α3 and α5-containing receptors, several differences in

interactions between DBI and the GABAR were observed. The GABAR α5 subunit had more

residues predicted to H-bond to DBI than the α3 subunit. For example, the α3 histidine at

position 142 in Loop7 is not predicted to H-bond with DBI but the aligned α5 glutamine does. In

Loop C, the α3 Ile202 does not H-bond with DBI but the aligned α5 asparagine does. In

addition, DBI had more H-bond interactions with the γ subunit of α5-containing GABARs.

Overall, the computational dockings provide support for the idea that DBI can bind at the BZD

binding site interface of GABARs with its flexible ODN loop buried in the BZD binding pocket.

The dockings provide a framework for revealing structural mechanisms underlying DBI effects

on GABARs.

Discussion

Despite the fact that DBI was identified as a putative endozepine in the mid-1980s [7],

the mechanisms underlying DBI's actions are still unclear. Due to DBI's intracellular roles in

long-chain fatty acid metabolism, and its multiple biologically active cleavage products [30-33],

direct effects of DBI on GABAR function have not been extensively studied. The use of

flumazenil, Ro15-1788 (FLZ, a BZD binding site antagonist) provides indirect evidence for the

existence of an endogenous GABAR modulator. Application of this BZD zero-modulator

increases the decay of GABA-evoked inhibitory postsynaptic currents in neurons in the

hippocampus [34], dentate gyrus [35], neocortex [36] and the thalamic reticular nucleus [8],

suggesting that FLZ is displacing an endogenous GABAR modulator. Recording from cells in

the thalamic reticular nucleus suggest that DBI positively modulates GABARs [8] while studies

in the subventricular zone and hippocampal subgranular cells suggest DBI is a NAM [9, 10]. In

this study, using purified DBI and heterologous expression of GABARs, we found that DBI is a

PAM for α5-containing GABARs and a weak NAM for α3-containing receptors demonstrating

that effects of DBI on GABAR activity are regulated by receptor subunit composition.

Recordings from cells in the hippocampal subgranular zone suggest that DBI or its

peptide cleavage product, ODN, act as a NAM [10]. Given α5 subunits are highly expressed in

the hippocampus [37], we expected that heterologously expressed α5-containing receptors

would be negatively modulated by DBI versus positively modulated, which we observed (Fig. 3).

Similarly, patch clamp recordings from the thalamic reticular nucleus where α3 is highly

expressed [38] suggested that these receptors are positively modulated by DBI [8], while our

experiments with heterologously expressed $\alpha_3\beta_3\gamma_{2L}$ GABARs demonstrated that DBI acts as

weak NAM (Fig. 3).

The differences between data reported here and previously reports are likely due to

experimental systems used. In neurons, GABARs are likely associated with accessory subunits

such as GARLH or Shisa7 [39, 40], which may alter DBI actions. Furthermore, α3 and α5 are

not the only α subunits expressed in the thalamic reticular nucleus and hippocampus,

respectively. In native cells, the presence of GABARs comprised of other subunits that may be

modulated by DBI make it difficult to assign DBI's effects to one specific GABAR subtype. Here,

using a reductionist approach, we demonstrate that DBI is a PAM of $\alpha_5\beta_3\gamma_{2L}$ GABARs and a

weak NAM of $\alpha_3\beta_3\gamma_{2L}$ GABARs.

318 $\alpha_5\beta_3\gamma_{2L}$ GABARs are highly expressed in the hippocampus and can be found both

319 synaptically and extrasynaptically[41]. α5-containing GABARs play a large role in hippocampal-

320 dependent forms of learning and memory [42-45]. Our data demonstrating that DBI acts as a

321 PAM for α5-containing receptors suggests that DBI should have effects on learning and

322 memory. Consistent with this idea, DBI-knockout mice display a disruption in spatial learning

323 and memory [46] and social behavior [47]. While somatic IPSP recordings from hippocampal

324 CA1 and dentate gyrus regions found that the DBI knockout did not change the effects of FLZ

325 on mIPSPs [48], these fast IPSPs recorded from the soma of pyramidal neurons are unlikely to

326 capture the effects of DBI on the α5-containing receptors that are expressed primarily in the

327 distal dendritic regions, which are involved in hippocampal learning and memory [49, 50].

328 Dysregulation of GABA-mediated signaling is implicated in a wide variety of neurological

329 diseases and disorders, including Alzheimer's [51, 52], anxiety [4], Parkinson's [53] and epilepsy

330 [2]. Our findings demonstrating that DBI's effects are dependent on GABAR subunit composition

331 lay an important foundation for understanding how inhibition in the brain is regulated.

332 Supporting Information

333 **S1 – Original 15% SDS-PAGE of Fig 1A.** Lanes 1 and 10 show Biorad's Precision Protein

334 Plus All Blue protein standards ranging from 10 kD to 75 kD. Lane 2 shows protein content of

335 the supernatant after emulsion and final high-speed spin. Proteins contained in flow-through,

336 first and second washes are observed in lanes 3, 4 and 5, respectively. Sample was then eluted

337 from the NiNTA column with a single band observed at the expected size of 11kD (lane 6). Lane

338 7 is the flow-through when the sample is loaded onto the PD10 desalting column. Lanes 8 and 9

339 are 6 and 3µL of the PD10 elution, which represents the final, purified and desalted DBI.

340 Overall, we observed the purification of a single ~11 kDa protein, which is the expected size of

341 His-DBI. Boxed area indicates which region of the gel was shown in Fig 1A.

S2 Table – Raw data for GABA dose-response experiments in Fig 2. Data represent GABA-

elicted currents in nA.

S3 Table – Table of data from Fig 3. Data represent $I_{GABA+Drug}/I_{GABA}$.

Acknowledgments

We thank the lab of Dr. Henzler-Wildman, University of Wisconsin - Madison for their

help with NMR. This study made use of the National Magnetic Resonance Facility at Madison,

which is supported by NIH grant P41GM136463, old number P41GM103399 (NIGMS) and

P41RR002301. Equipment was purchased with funds from the University of Wisconsin-

Madison, the NIH P41GM103399, S10RR02781, S10RR08438, S10RR023438, S10RR025062,

S10RR029220), the NSF (DMB-8415048, OIA-9977486, BIR-9214394), and the USDA.

Funding for this manuscript from NIH-R21NS099991-01 and NIH-F31NS101804.

References:

1.   Organization WECotSoEDWH. The selection of essential drugs : report of a WHO expert
committee [meeting held in Geneva from 17 to 21 October 1977]. Geneva: 1977.

2.   Palma E, Ruffolo G, Cifelli P, Roseti C, Vliet EAV, Aronica E. Modulation of GABAA
Receptors in the Treatment of Epilepsy. Curr Pharm Des. 2017;23(37):5563-8. Epub
2017/08/12. doi: 10.2174/1381612823666170809100230. PubMed PMID: 28799512.

3.   Wisden W, Yu X, Franks NP. GABA Receptors and the Pharmacology of Sleep. Handb
Exp Pharmacol. 2019;253:279-304. Epub 2017/10/11. doi: 10.1007/164_2017_56. PubMed
PMID: 28993837.

4.   Engin E, Benham RS, Rudolph U. An Emerging Circuit Pharmacology of GABA A
Receptors. Trends in Pharmacological Sciences. 2018;39(8):710-32. doi:
10.1016/j.tips.2018.04.003.

5.   Mohler H. The rise of a new GABA pharmacology. Neuropharmacology. 2011;60(7-
8):1042-9. doi: 10.1016/j.neuropharm.2010.10.020. PubMed PMID: ISI:000290648900004.

6.   Knoflach F, Benke D, Wang Y, Scheurer L, Lüddens H, Hamilton BJ, et al.
Pharmacological modulation of the diazepam-insensitive recombinant gamma-aminobutyric
acidA receptors alpha 4 beta 2 gamma 2 and alpha 6 beta 2 gamma 2. Mol Pharmacol.
1996;50(5):1253-61.

7.   Guidotti A, Forchetti CM, Corda MG, Konkel D, Bennett CD, Costa E. Isolation,
characterization, and purification to homogeneity of an endogenous polypeptide with agonistic
action on benzodiazepine receptors. Proc Natl Acad Sci U S A. 1983;80(11):3531-5. Epub
1983/06/01. doi: 10.1073/pnas.80.11.3531. PubMed PMID: 6304714; PubMed Central PMCID:
PMCPMC394079.

8.   Christian CA, Herbert AG, Holt RL, Peng K, Sherwood KD, Pangratz-Fuehrer S, et al.
Endogenous Positive Allosteric Modulation of GABA(A) Receptors by Diazepam binding
inhibitor. Neuron. 2013;78(6):1063-74. doi: 10.1016/j.neuron.2013.04.026. PubMed PMID:
ISI:000321026900012.

9.   Alfonso J, Le Magueresse C, Zuccotti A, Khodosevich K, Monyer H. Diazepam Binding
Inhibitor Promotes Progenitor Proliferation in the Postnatal SVZ by Reducing GABA Signaling.
Cell Stem Cell. 2012;10(1):76-87. doi: 10.1016/j.stem.2011.11.011. PubMed PMID:
ISI:000299449500012.

10.   Dumitru I, Neitz A, Alfonso J, Monyer H. Diazepam Binding Inhibitor Promotes Stem Cell
Expansion Controlling Environment-Dependent Neurogenesis. Neuron. 2017;94(1):125-37.e5.
doi: 10.1016/j.neuron.2017.03.003.

11.   Liu X, Li Y, Zhou L, Chen H, Su Z, Hao W. Conditioned place preference associates with
the mRNA expression of diazepam binding inhibitor in brain regions of the addicted rat during
withdrawal. 2005;137(1-2):47-54. doi: 10.1016/j.molbrainres.2005.02.021.

12.   Costa E, Guidotti A. Diazepam Binding Inhibitor (Dbi) - a Peptide with Multiple Biological
Actions. Life Sci. 1991;49(5):325-44. doi: Doi 10.1016/0024-3205(91)90440-M. PubMed PMID:
ISI:A1991FV18900001.

13.   Mortensen M, Patel B, Smart TG. GABA potency at GABA(A) receptors found in
synaptic and extrasynaptic zones. Front Cell Neurosci. 2012;6. doi: ARTN 1

10.3389/fncel.2012.00001. PubMed PMID: ISI:000299807800001.

14.   Delaglio F, Grzesiek S, Vuister G, Zhu G, Pfeifer J, Bax A. NMRPipe: A multidimensional
spectral processing system based on UNIX pipes. J Biomol Nmr. 1995;6(3). doi:
10.1007/bf00197809.

15.   Vranken WF, Boucher W, Stevens TJ, Fogh RH, Pajon A, Llinas M, et al. The CCPN
data model for NMR spectroscopy: Development of a software pipeline. Proteins: Structure,
Function, and Bioinformatics. 2005;59(4):687-96. doi: https://doi.org/10.1002/prot.20449.

16.   Venkatachalan SP, Bushman JD, Mercado JL, Sancar F, Christopherson KR, Boileau
AJ. Optimized expression vector for ion channel studies in Xenopus oocytes and mammalian
cells using alfalfa mosaic virus. Pflugers Arch. 2007;454(1):155-63. Epub 2006/12/05. doi:
10.1007/s00424-006-0183-1. PubMed PMID: 17146677.

17.   Boileau AJ, Kucken AM, Evers AR, Czajkowski C. Molecular dissection of
benzodiazepine binding and allosteric coupling using chimeric gamma-aminobutyric acidA
receptor subunits. Mol Pharmacol. 1998;53(2):295-303. Epub 1998/03/14. doi:
10.1124/mol.53.2.295. PubMed PMID: 9463488.

18.   Boileau AJ, Baur R, Sharkey LM, Sigel E, Czajkowski C. The relative amount of cRNA
coding for gamma2 subunits affects stimulation by benzodiazepines in GABA(A) receptors
expressed in Xenopus oocytes. Neuropharmacology. 2002;43(4):695-700. Epub 2002/10/09.
doi: 10.1016/s0028-3908(02)00036-9. PubMed PMID: 12367615.

19.   Hanson SM, Czajkowski C. Structural mechanisms underlying benzodiazepine
modulation of the GABA(A) receptor. J Neurosci. 2008;28(13):3490-9. Epub 2008/03/28. doi:
10.1523/jneurosci.5727-07.2008. PubMed PMID: 18367615; PubMed Central PMCID:
PMCPMC2410040.

20.   Masiulis S, Desai R, Uchański T, Serna Martin I, Laverty D, Karia D, et al. GABAA
receptor signalling mechanisms revealed by structural pharmacology. Nature.
2019;565(7740):454-9. doi: 10.1038/s41586-018-0832-5.

21.   Taskinen JP, Van Aalten DM, Knudsen J, Wierenga RK. High resolution crystal
structures of unliganded and liganded human liver ACBP reveal a new mode of binding for the
acyl-CoA ligand. 2006;66(1):229-38. doi: 10.1002/prot.21124.

22.   Kozakov D, Beglov D, Bohnuud T, Mottarella SE, Xia B, Hall DR, et al. How good is
automated protein docking? 2013;81(12):2159-66. doi: 10.1002/prot.24403.

23.   Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, et al. The ClusPro web
server for protein–protein docking. Nature Protocols. 2017;12(2):255-78. doi:
10.1038/nprot.2016.169.

24.   Vajda S, Yueh C, Beglov D, Bohnuud T, Mottarella SE, Xia B, et al. New additions to the
ClusPro server motivated by CAPRI. Proteins: Structure, Function, and Bioinformatics.
2017;85(3):435-44. doi: 10.1002/prot.25219.

25.   Shilling PJ, Mirzadeh K, Cumming AJ, Widesheim M, Köck Z, Daley DO. Improved
designs for pET expression plasmids increase protein production yield in Escherichia coli.
Communications Biology. 2020;3(1). doi: 10.1038/s42003-020-0939-8.

26.   Karim N, Wellendorph P, Absalom N, Johnston GA, Hanrahan JR, Chebib M. Potency of
GABA at human recombinant GABA(A) receptors expressed in Xenopus oocytes: a mini review.
Amino Acids. 2013;44(4):1139-49. Epub 2013/02/07. doi: 10.1007/s00726-012-1456-y. PubMed
PMID: 23385381.

27.   Nimmich ML, Heidelberg LS, Fisher JL. RNA editing of the GABA(A) receptor alpha3
subunit alters the functional properties of recombinant receptors. Neurosci Res. 2009;63(4):288-
93. doi: 10.1016/j.neures.2009.01.003. PubMed PMID: 19367790.

28.   Duvaud S, Gabella C, Lisacek F, Stockinger H, Ioannidis V, Durinx C. Expasy, the Swiss
Bioinformatics Resource Portal, as designed by its users. Nucleic Acids Research.
2021;49(W1):W216-W27. doi: 10.1093/nar/gkab225.

29.   Ferrero P, Santi MR, Conti-Tronconi B, Costa E, Guidotti A. Study of an
octadecaneuropeptide derived from diazepam binding inhibitor (DBI): biological activity and
presence in rat brain. Proc Natl Acad Sci U S A. 1986;83(3):827-31. Epub 1986/02/01. doi:
10.1073/pnas.83.3.827. PubMed PMID: 3456171; PubMed Central PMCID: PMCPMC322958.

30.   Knudsen J. Acyl-Coa-Binding and Transport, an Alternate Function for Diazepam
Binding Inhibitor (Dbi), Which is Identical with Acyl-Coa-Binding Protein. Neuropharmacology.
1991;30:1405-10.

31.   Bouyakdan K, Taib B, Budry L, Zhao SG, Rodaros D, Neess D, et al. A novel role for
central ACBP/DBI as a regulator of long-chain fatty acid metabolism in astrocytes. J
Neurochem. 2015;133(2):253-65. doi: 10.1111/jnc.13035. PubMed PMID:
ISI:000352796000010.

32.   Slobodyansky E, Guidotti A, Wambebe C, Berkovich A, Costa E. Isolation and
characterization of a rat brain triakontatetraneuropeptide, a posttranslational product of
diazepam binding inhibitor: specific action at the Ro 5-4864 recognition site. J Neurochem.
1989;53(4):1276-84. Epub 1989/10/01. doi: 10.1111/j.1471-4159.1989.tb07425.x. PubMed
PMID: 2769267.

33.   Tonon MC, Vaudry H, Chuquet J, Guillebaud F, Fan J, Masmoudi-Kouki O, et al.
Endozepines and their receptors: Structure, functions and pathophysiological significance.
Pharmacol Ther. 2020;208:107386. Epub 2019/07/10. doi: 10.1016/j.pharmthera.2019.06.008.
PubMed PMID: 31283949.

34.   King GL, Knox JJ, Dingledine R. Reduction of inhibition by a benzodiazepine antagonist,
Ro15-1788, in the rat hippocampal slice. 1985;15(2):371-8. doi: 10.1016/0306-4522(85)90219-
2.

35.   Leroy C, Poisbeau P, Keller AF, Nehlig A. Pharmacological plasticity of GABA A
receptors at dentate gyrus synapses in a rat model of temporal lobe epilepsy. 2004;557(2):473-
87. doi: 10.1113/jphysiol.2003.059246.

36.   Ali AB, Thomson AM. Synaptic 5 Subunit-Containing GABAA Receptors Mediate IPSPs
Elicited by Dendrite-Preferring Cells in Rat Neocortex. Cerebral Cortex. 2008;18(6):1260-71.
doi: 10.1093/cercor/bhm160.

37.   Sequeira A, Shen K, Gottlieb A, Limon A. Human brain transcriptome analysis finds
region- and subject-specific expression signatures of GABAAR subunits. Communications
Biology. 2019;2(1). doi: 10.1038/s42003-019-0413-7.

38.   Hörtnagl H, Tasan RO, Wieselthaler A, Kirchmair E, Sieghart W, Sperk G. Patterns of
mRNA and protein expression for 12 GABAA receptor subunits in the mouse brain.
Neuroscience. 2013;236:345-72. Epub 2013/01/23. doi: 10.1016/j.neuroscience.2013.01.008.
PubMed PMID: 23337532; PubMed Central PMCID: PMCPMC3605588.

39.   Yamasaki T, Hoyos-Ramirez E, Martenson JS, Morimoto-Tomita M, Tomita S. GARLH
Family Proteins Stabilize GABA(A) Receptors at Synapses. Neuron. 2017;93(5):1138-52.e6.
doi: 10.1016/j.neuron.2017.02.023. PubMed PMID: 28279354.

40.   Han W, Li J, Pelkey KA, Pandey S, Chen X, Wang Y-X, et al. Shisa7 is a
GABA$_A$ receptor auxiliary subunit controlling benzodiazepine actions. Science.
2019;366(6462):246-50. doi: 10.1126/science.aax5719.

41.   Mortensen M, Patel B, Smart TG. GABA Potency at GABAA Receptors Found in
Synaptic and Extrasynaptic Zones. Front Cell Neurosci. 2012;6. doi: 10.3389/fncel.2012.00001.

42.   Collinson N, Atack JR, Laughton P, Dawson GR, Stephens DN. An inverse agonist
selective for alpha5 subunit-containing GABAA receptors improves encoding and recall but not
consolidation in the Morris water maze. Psychopharmacology (Berl). 2006;188(4):619-28. Epub
2006/04/25. doi: 10.1007/s00213-006-0361-z. PubMed PMID: 16633803.

43.   Joksimović S, Divljaković J, Van Linn ML, Varagic Z, Brajković G, Milinković MM, et al.
Benzodiazepine-induced spatial learning deficits in rats are regulated by the degree of
modulation of α1 GABAA receptors. European Neuropsychopharmacology. 2013;23(5):390-9.
doi: https://doi.org/10.1016/j.euroneuro.2012.05.003.

44.   Timić T, Joksimović S, Milić M, Divljaković J, Batinić B, Savić MM. Midazolam impairs
acquisition and retrieval, but not consolidation of reference memory in the Morris water maze.
Behavioural Brain Research. 2013;241:198-205. doi: https://doi.org/10.1016/j.bbr.2012.12.014.

45.   Crestani F, Keist R, Fritschy J-M, Benke D, Vogt K, Prut L, et al. Trace fear conditioning
involves hippocampal α5 GABAA receptors. Proceedings of the
National Academy of Sciences. 2002;99(13):8980-5. doi: 10.1073/pnas.142288699.

46.   Ujjainwala AL, Courtney CD, Wojnowski NM, Rhodes JS, Christian CA. Differential
impacts on multiple forms of spatial and contextual memory in diazepam binding inhibitor
knockout mice. Journal of Neuroscience Research. 2019. doi: 10.1002/jnr.24393.

47.   Ujjainwala AL, Courtney CD, Rhoads SG, Rhodes JS, Christian CA. Genetic loss of
diazepam binding inhibitor in mice impairs social interest. Genes, Brain and Behavior.
2018;17(5):e12442. doi: 10.1111/gbb.12442.

48.   Courtney CD, Christian CA. Subregion-Specific Impacts of Genetic Loss of Diazepam
Binding Inhibitor on Synaptic Inhibition in the Murine Hippocampus. Neuroscience.
2018;388:128-38. doi: 10.1016/j.neuroscience.2018.07.012.

49.   Rodgers FC, Zarnowska ED, Laha KT, Engin E, Zeller A, Keist R, et al. Etomidate
Impairs Long-Term Potentiation In Vitro by Targeting 5-Subunit Containing GABAA Receptors
on Nonpyramidal Cells. J Neurosci. 2015;35(26):9707-16. doi: 10.1523/jneurosci.0315-15.2015.

50.   Groen MR, Paulsen O, Pérez-Garci E, Nevian T, Wortel J, Dekker MP, et al.
Development of dendritic tonic GABAergic inhibition regulates excitability and plasticity in CA1
pyramidal neurons. Journal of Neurophysiology. 2014;112(2):287-99. doi:
10.1152/jn.00066.2014. PubMed PMID: 24760781.

51.   Calvo-Flores Guzmán B, Vinnakota C, Govindpani K, Waldvogel HJ, Faull RLM,
Kwakowsky A. The GABAergic system as a therapeutic target for Alzheimer's disease. J
Neurochem. 2018;146(6):649-69. doi: 10.1111/jnc.14345.

52.   Govindpani K, Calvo-Flores Guzmán B, Vinnakota C, Waldvogel HJ, Faull RL,
Kwakowsky A. Towards a Better Understanding of GABAergic Remodeling in Alzheimer's
Disease. International Journal of Molecular Sciences. 2017;18(8):1813. PubMed PMID:
doi:10.3390/ijms18081813.

53.   Brickley SG, Mody I. Extrasynaptic GABA(A) receptors: their function in the CNS and
implications for disease. Neuron. 2012;73(1):23-34. doi: 10.1016/j.neuron.2011.12.012. PubMed
PMID: 22243744.

Figure

Figure

Figure

Figure