Niche theory for within-host parasite dynamics: Analogies to food web modules via feedback loops

Ashwini Ramesh[1*] and Spencer R Hall[1]

[1]Department of Biology, Indiana University, Bloomington, Indiana, USA – 47405

*Corresponding author email: aramesh@iu.edu

Co-author email: sprhall@iu.edu

Mailing address: Biology Building 239, 1001 E 3rd Street, IN 47405 USA

Phone number: + 1 812-855-6009

Statement of authorship: AR and SRH conceptualized the paper. AR performed the analysis and wrote the first draft of the manuscript, and SRH contributed to subsequent revisions.

Data accessibility statement: All Mathematica code will be published in a public repository upon acceptance

Competing Interest: Authors declare no competing interests

ABSTRACT

Why do parasites exhibit a wide dynamical range within their hosts? For instance, why can a parasite only sometimes successfully infect its host? Why do some parasites exhibit large fluctuations? Why do two parasites coinfect, exclude each other, or win only sometimes over another (via priority effects)? For insights, we turn to food webs. An omnivory model (IGP) blueprints one parasite competing with immune cells for host energy (PIE), and a competition model (keystone predation, KP) mirrors a new coinfection model (2PIE). We then draw analogies between models using feedback loops. We translate those loops into the intraspecific direct (DE) and indirect effects (IE) that create various dynamics. Three points arise. First, a prey or parasite can flip between stable and oscillatory coexistence with their enemy with weakening IE and strengthening DE. Second, even with comparable loop structure, a parasite cannot exhibit priority effects seen in IGP due to constraints imposed by production of immune cells. Third, despite simpler loop structure, KP predicts parallel outcomes in the two-parasite model due to comparable structure of interactions between competing victims and their resources and enemies. Hence, food web models offer powerful if imperfect analogies to feedbacks underlying the dynamical repertoire of parasites within hosts.

Key words: intraguild predation, keystone predation, coexistence, coinfection, priority effects, within-host competition, feedback loops, inverse Jacobian matrices

INTRODUCTION

Parasites can show a variety of dynamics within hosts. These within-host dynamics can significantly affect health of individual hosts (Råberg *et al* 2009; Cressler *et al* 2014; Bashey 2015) and alter population-level disease outbreaks (Mideo 2008; Råberg *et al* 2012; Gorisch *et al* 2018). For instance, exposure to parasite propagules often causes infection of some but not all hosts despite seemingly similar starting conditions (Merrill & Cáceres 2018). The success or failure of invasion and persistence of a parasite within a host determines its infection status or even death (Duneau *et al* 2017; Merrill & Cáceres 2018). Once infected, the stability or instability of within-host dynamics can further impact both host health and transmission to others. For instance, periods of high abundance of a cycling gut parasite can both decrease foraging and increase transmission to bee hosts (Otterstatter & Thomson 2006). Similarly, coinfection (*i.e.,* within-host coexistence of parasites) can reduce fecundity of hosts more than single infection (Lohr *et al* 2010). At the between-host scale, coinfection can reduce or increase outbreaks compared to singly infected populations (Clay *et al* 2019; Susi *et al* 2015). Consequently, various dynamics of parasites can arise within hosts, likely mediated by immune systems and nutrient/energy availability (Graham 2008; Cressler *et al* 2014). But which mechanisms determine those dynamics?

To answer this question, two classic food web modules from community ecology provide analogies. In intraguild predation (IGP), an omnivorous predator and its prey compete for a shared resource (Holt & Polis 1997; Verdy & Amarasekare 2010). Similarly, immune cells and a parasite compete for shared energy within a host (Fig. 1A - C; Cressler *et al* 2014). Hence, mechanisms that capture the repertoire of dynamics in IGP (stable coexistence, oscillations, priority effects, exclusion) could also produce those observed in single-parasite studies (Otterstatter & Thomson 2006; Merrill & Cáceres 2018). Additionally, in the diamond-keystone predation (KP) model, two prey species share a resource (exploitative competition) and a predator (apparent competition: Holt *et al.* 1994, Leibold 1996). Similarly, two parasites can compete for a shared resource while facing

immune attack (Fig. 1D - F). In such a scenario, the KP model could anticipate the tradeoffs and

niche dimensions needed for coexistence, exclusions, and priority effects of coinfecting parasites

(Natsopoulou *et al* 2015). The underlying hope, then, is that food web modules might provide apt

blueprints, because they share basic consumer-resource structure with their within-host analogues.

However, a major complication arises: the enemy is generated differently in food webs vs.

within a host (Wodarz 2006). Predators attack and assimilate prey for reproduction into new

predators. Immune cells also attack parasites, but they simultaneously require host energy to

produce immune cells (Cressler *et al* 2014). These "consumer-resource like" interactions can

involve products of immune cells, parasites, and energy. Additionally, hosts can allocate a baseline

level of energy straight to immune cells, a pipeline not enjoyed by even omnivorous predators in

typical food webs. Therefore, these fundamentals of immune function – even when highly

simplified - create more interaction links (Fig. 1). Those additional links might alter feedbacks

underpinning the range of stability outcomes (Metcalf *et al* 2020), possibly undermining analogies

from food webs or creating new outcomes altogether. Given these issues (Wodarz 2006; Alizon *et*

al 2008; Fenton & Perkin 2010), how could we usefully compare and contrast stability outcomes

in food webs (IGP, KP) to their within-host analogues?

We tackle this challenge by using a feedback loop approach. Feedback loops link the

strength of consumer-resource interactions to stability (Puccia & Levins 1991). Indeed, feedback

loops characterize the biology behind stability and enable comparison of structurally similar but

biologically distinct systems. Every *n*-dimensional system consists of *n* levels of feedback (F_n).

The simplest level of feedback, level 1 (F_1), involves intraspecific direct effects (DE). Here, a

small increase of a species impacts its own growth rate – positively or negatively - via interaction

with itself and from those directly consuming it. Then, longer loops (from 2 through *n* dimensions)

involve combinations of intra- and inter-specific direct effects (calculated from Jacobian matrices).

For instance, in a two-species loop, an increase in a resource's density could increase growth rate

of its consumer. Increased density of its consumer, in turn, could then depress growth rate of that

resource (hence, binary consumer-resource interactions create negative F_2). With more

dimensions, the connections between interactors become more complex and varied, potentially

introducing both positive and negative feedbacks. The sign and strength of feedback at all relevant

levels, then, governs stability. Additionally, key ratios of those longer loops determine

intraspecific indirect effects (IE; developed in Box 1). As we then show, feedback at certain levels,

translated into intraspecific direct and indirect effects, determines stability of the interactions.

Using intraspecific direct and indirect feedback loops, we first revisit the foundational IGP

and KP models. We then apply these loop-based lessons to analogous parasite-immune-energy

models (PIE and 2PIE; Fig. 1). To study these analogues, we assume that parasite(s) respond to

attack by a common immune system. That assumption surely oversimplifies some examples (*e.g.,*

in vertebrates: Ezenwa & Jolles 2011), but it seems like a reasonable place to start for others

(Graham 2008; Griffits *et al* 2015; Rynkiewicz *et al* 2015). Additionally, we assume that immune

cells and parasites share host energy (a resource, following Griffits *et al* 2015). We then evaluate

how host energy and immune system interact to mediate parasite dynamics within hosts (De Roode

et al 2005; Ezenwa & Jolles 2011; Doublet *et al* 2015).

The analysis involves three parts. First, we translate loops into a scheme framed as

intraspecific direct and indirect effects (DE and IE). This interpretation works most thoroughly for

3D systems (IGP and PIE models), but we extended key ideas to 4D systems (KP and 2PIE

models). Second, using these tools, we analyzed stability outcomes for IGP vs. PIE. We find that

IGP and PIE share some dynamics (coexistence, stably or via oscillations) due to similar loop

structure. However, the nature of the generation of immune cells prevents some outcomes present

in IGP (involving two variations on alternate stable states). Third, we find that despite having

simpler feedback links, KP anticipates coexistence/coinfection vs. priority effects (or alternate

stable states) of competing parasites in 2PIE models. Furthermore, these outcomes arise for

conceptually similar reasons. Specifically, depending on their traits, competing prey or parasites

exhibit symmetries (coexistence/coinfection) or asymmetries (priority effects) in ratios describing

'*effects on*' their enemies and resources vs. how they are '*affected by*' them. Those (a)symmetries

determine the strength of tradeoffs and competitive hierarchies among competitors. They also

underpin the sign of feedback at a key level, hence they govern whether parasites experience

negative or positive indirect effects. Furthermore, allocation of energy to constitutive immunity

enhances negative feedback, thereby shrinking regions of within-host oscillations, coinfection, and

parasite burden. Thus, feedback loops, guided by our interpretation schemes, show when and why

comparable dynamics arise in these food webs and their within-host analogues. However, we

highlight why differences in production of enemies constrain some outcomes within hosts.

METHODS AND RESULTS

Overview of the models

We compare and contrast modules of food web and within-host dynamics. The model with

one prey, predator, and resources (intraguild predation, IGP; Holt & Polis 1997; Verdy &

Amarasekare 2010) parallels a within-host model of one parasite species, immune cells, and host

energy (PIE; modified from Hite & Cressler 2018). Similar, a two prey, predator, resource model

(keystone predation, KP; Leibold 1996) parallels one with the new two parasite species, immune

cells, and host energy model (2PIE). However, while the food web and within-host modules share

similar structure, they create enemies differently. Specifically, predators proliferate via direct

consumption of their prey, while immune cells jointly require both parasites and a shared energy.

Additionally, the within-host modules include two variants that commonly arise in invertebrate

host-parasite systems. In one, only parasites activate production of immune cells (PIE and 2PIE

models). In the other, energy is continuously allocated to maintain baseline immune function, even

without parasites (PIEc and 2PIEc, constitutive immunity). Despite their differences, we compare

across food web and within-host modules using feedback loops. Feedback loops provide a

common metric to unpack the biology underlying stability. Jacobian matrices provide the starting

point (Box 1, Appendix Section 1). Each of Jacobian terms represents the direct effect of species

j on growth rate of species i (J_{ij}), yielding (hereafter) interspecific *positive effects* (black arrow)

and *negative effects* (red arrow) and intraspecific *self-limitation* (red curve) or *self-facilitation*

(black curve; Box 1, Fig. 1 [not seen here]). As shown below, these terms combine into loops at

various levels.

One prey - predator - resource (IGP) | One parasite - immune cells - host energy (PIE & PIEc)

At their heart, both IGP and PIE models hinge on binary consumer-resource like

interactions (Fig.2, Table 1). In these interactions, consumers directly benefit while resources

suffer direct costs. In IGP, omnivorous predators (P) can consume prey (N_1) and resources (R).

An interior equilibrium becomes feasible when each species can maintain positive density ($P^* >$

$0, N_1^* > 0, R^* > 0$). At IGP's interior, these consumer-resource interactions exert positive effects

on the predator but negative ones on prey and resource. Then, prey also indirectly compete with

predators for the shared resource. Consumption of resources benefits the prey and harms resources.

Finally, chemostat-like renewal imposes self-limitation on resources (Fig. 1A). Similarly, in the

within-host models, immune cells (I) "consume" two items, simultaneously killing parasites (N_1)

while taking up host energy (E). At PIE's interior, these consumer-resource-like interactions have

a positive effect on I and negative on N_1 and E. The parasite competes for this shared energy;

consumption of host energy positively effects parasites and negatively effects energy.

Additionally, the donor-controlled renewal of host energy imposes self-limitation on energy

without baseline allocation ($a_b = 0$; Fig. 1B). Fixed energy allocated to immune cells ($a_b > 0$)

creates additional self-limitation on the immune cells in PIEc (Fig. 1C).

Intraspecific direct and indirect effects: an approach using feedback (Box 1 and Fig. 2).

To compare stability of these models, we used feedback loops (Box 1). Any *n*-dimensional

system can be decomposed into *n* levels of feedback that determine stability of an equilibrium

(Box 1, Top Panel). An interior equilibrium can allow stable or unstable coexistence (oscillations)

or produce, potentially, alternate stable states (if it is a saddle). As illustrated by and applied to

IGP and PIE, the three-dimensional system creates three levels of feedback (Box 1, Fig. 2). For

example, level 1 feedback (F_1) is the sum of intraspecific direct effects (Box 1) and is the

Jacobian's trace (Appendix Section 1). At the interior of IGP and PIE, the basal resource / energy

contributes to the direct effects at the feasible interior (Fig. 2, Appendix Section 2). Self-limitation

comes from the chemostat-supplied resource and its consumption; in PIEc, baseline allocation to

immunity adds self-limitation (hence negative level 1 feedback [see below but also Appendix

Section 1]). Level 2 (F_2) feedback sums up two-species loops (Box 1). In PIE models (Fig. 2), this

sum of pairwise consumer resource come from energy-parasite (E-N_1), energy-immune cells (E-

I), and parasite-immune cells (N_1-I, with R-N_1, R-P, and N_1-P analogies in IGP, respectively). At

the interior, all of these loops contribute to negative feedback. That negative feedback intensifies

in PIEc from the product of energy and immune self-limitation. In all 3D models considered here,

F_2 is negative at the interior equilibrium (based just on signs alone): level 2 feedback stabilizes.

Level 3 feedback (F_3) sums three-species loops. It is also the negative of the determinant

of the Jacobian, - |$\mathbf{J}$| (Appendix Sections 1, 2). In PIE models, these loops at the interior are (Fig.

2, *L-R*): (i) "*N_1 is eaten*", a negative loop coming from the stabilizing product of the immune-

parasite loop and energy self-limitation; (ii) "*starving the enemy*", a positive loop where a small

increase in the parasite reduces energy for the immune cells, hence reduces immune attack (i.e.,

$\uparrow N_1 \rightarrow \downarrow E \rightarrow \downarrow I \rightarrow \uparrow N_1$); (iii) "*feeding the enemy*", a negative loop where a small increase of the

parasite 1 stimulates the immune cells which consume energy that then starves the parasite ($\uparrow N_1 \rightarrow$

$\uparrow I \rightarrow \downarrow E \rightarrow \downarrow N_1$). Thus, negative feedback operates when N_1 hurts itself by "feeding" the immune

cells. Similar loops arise in IGP. Finally, in only PIEc, (iv) a negative "N_1 eats" loop arises from

the stabilizing product of energy consumption by the parasite and immune self-limitation.

Combined, these three levels of feedback determine stability of the interior equilibrium.

First, stability requires that each level of feedback is negative (so, $F_1 < 0$, $F_2 < 0$, and $F_3 < 0$). When

$F_3 < 0$, the interior is stable; when $F_3 > 0$, it is a saddle. Additionally, stability requires that $F_1 F_2$

$+ F_3 > 0$, or that upper-level feedbacks (F_3) not be too strong relative to lower-level feedback (F_1

F_2). All of these conditions have analogies to the Routh-Hurwitz criteria, differing only in sign

(Puccia & Levins 1991). Here, we also reframe stability analysis as sums of intraspecific direct

effects (F_1, hereafter DE) and of intraspecific indirect effects ($-F_2 / F_3$, the trace of the inverse

Jacobian, IE: see Appendix Section 1, Box 1). Written this way, IE is a ratio of the longer loops,

where each species interacts with other species in feedback at level 2 (F_2) and level 3 (F_3). It is

also the sum of intraspecific indirect effects of each species N_i (lying on tr[$-\mathbf{J}^{-1}$]; Eq. 5):

$$IE = IE_{N_1} + IE_{N_2} + IE_{N_3} = -F_2/F_3 \qquad\qquad\qquad \text{(Eq. 5)}$$

The numerator of the intraspecific indirect effects of each species (IE_{N_i}) corresponds to specific

combinations of the reciprocal two-species F_2 loops; the denominator for each species is F_3. For

example, in general form, the corresponding indirect effect of N_1, IE_{N_1}, is the sum of (i) the product

of self-limitation of N_2 and N_3 and (ii) pairwise consumer-resource loop of N_2 and N_3, all divided

by F_3 (green shading over gray shading, Box 1). Similar expressions can be derived for species 2

(IE_{N_2}, yellow shading) and 3 (IE_{N_3}, blue shading). With DE and IE in hand, stability then depends

on their sign and the strength of their product. Specifically, *stable coexistence* occurs when direct

and indirect effects are negative (DE < 0, IE < 0) and jointly strong (DE x IE > 1; bottom of Box

1, light orange region; equivalent to $F_1 F_2 > -F_3$). *Oscillations* arise when direct and indirect effects

are jointly weak (dark orange, DE x IE < 1 for DE < 0, or DE > 0; equivalent to $F_1 F_2 < -F_3$).

Finally, *alternate stable states (aka priority effects)* occur with positive indirect effects (IE > 0,

i.e., when $F_3 > 0$ for IGP and PIE, given that $F_2 < 0$ for both at their interiors; Box 1, gray region).

Note: we use the terms alternate stable states and priority effects interchangeably. They both

denotes how initial densities, even offset in time (*e.g.,* sequential infection), can determine

competitive outcomes (depending on whose domain of attraction they move through).

Weakening DE x IE generates oscillations in these models (Fig. 3 and Appendix Section 2)

Assembly of the interior equilibrium: To compare stability outcomes across IGP and PIE

models, we created parallel 2D bifurcation diagrams (Fig 3). These diagrams spread gradients of

parameter space of an environmental factor, supply of resource (S), and trait space of the focal

prey/parasite, feeding rate of N_1 on resource, f_{N_1} (Fig. 3A-C; interpreted with 1D bifurcation

diagrams: Fig. 3G-I). The lines within the 2D bifurcation diagram represent shifts in species

composition or in dynamics (stable coexistence, oscillations, or alternative stable states; see

Appendix Section 2 for full analysis). In IGP (Fig. 3A), when resource supply and feeding rate of

prey are too low to meet the minimal resource requirement of either the prey or predator, only R

is supported in the system (R, yellow). Increases in S allows R to meet the minimum resource

requirement (R^*) of the predator (hence P invades creating the R-P region, green). As feeding rate

of prey increases along a fixed resource supply ($S = 110$), the prey becomes a better competitor

for resources, enabling births, $f_{N_1}R^*$, to exceed deaths, $f_{PN_1} P^* + m_N$ at the R-P boundary. Hence,

the prey can invade; both prey and predator push R^* lower in the R-N_1-P interior (Fig. 3A, Fig.

3G). In this region, R-N_1-P are able to coexist either stably (light orange) or via oscillations (dark

orange). On the other hand, as the feeding rate of prey increases at low S ($S = 5$), prey invade the

resource-only boundary (creating the R-N_1 region, blue). The omnivorous predator can then invade

this R-N_1 boundary when its birth rate, $e_{PN_1} f_{PN_1} N_1^* + f_{PR} R^*$, exceeds its death rate, m_P.

Successful invasion creates stable coexistence (orange; Fig. 3A). Thus, R, R-P, R-N_1, and R-N_1-P

equilibria (oscillatory and stable coexistence) are possible.

The within-host PIE model contains some but not all analogous states. The key differences

stem from how hosts produce immune cells. In PIE (Fig. 3B), immune cells only generate with

parasites. Hence, a host energy-immune (*E-I*) state, analogous to the *R-P* one, is not possible.

Similar to IGP, though, low feeding rate can prevent parasite invasion (creating the *E*-only state

[yellow]). With increasing feeding rates (illustrated at $S = 500$), parasites (N_1) can eventually

invade creating E-N_1 space (light blue; Fig. 3H). Although parasites depress E^*, sufficiently high

parasite density enables production of immune cells, when the immune system's minimal E N_1

requirement is met (Fig. 3H). Hence, at medium to high feeding rates of parasites, E-N_1-I coexist

either stably (light orange) or via oscillations (dark orange). However, at sufficiently high f_{N_1},

parasites compete too strongly: they successfully starve out the immune cells, re-establishing a E-

N_1 space (blue). Thus, PIE exhibits only three equilibria: E, E-N_1, and E-N_1-I (oscillations and

stable coexistence; Fig. 3B; see Appendix Section 3B for details).

In PIEc, the host always allocates energy to the immune system ($a_b > 0$). This biology

eliminates the *E*-alone and *E-N₁* region. Instead, immune cells prevent infection (*E-I* space; low

f_{N_1}; green) or hosts become infected (*E-N₁-I* space; higher f_{N_1}; orange). Parasites successfully

infect when their births, $(f_{N_1} / e_{N_1 E}) E^*$, exceed their deaths at immune-enhanced, $m_I + f_{I N_1} I^*$. Like

IGP, at high feeding rates of parasites, the *E-N₁-I* remains stable (light orange) or oscillates (only

at low a_b [Appendix Section 2]). Thus, in PIEc, immune biology eliminates even more states; only

E-I and *E-I-N₁* (oscillations and stable coexistence) remain possible (Fig. 3C).

Stability outcomes: Despite these differences, oscillations arise for similar reasons in the

food web and within-host models. As noted in Box 1, oscillations require weakening of the joint

product of intraspecific direct effects (DE) and indirect effects (IE). In all three cases, stable

coexistence (DE x IE > 1) envelopes oscillatory regions (DE x IE < 1; Fig. 3A-B; Appendix Fig.

A2). In IGP, weakening of IE at low feeding rate of the prey, f_{N_1}, triggered oscillations despite

strengthening of DE (Fig. 3D). At high f_{N_1}, DE strengthens to regain stability. This general pattern

also arises with increasing feeding rate of parasites in the PIE and PIEc models (Fig. 3E-F).

However, DE differed slightly among models (Fig. 3D-F, black, solid lines), where DE is:

IGP: $-a - (f_{N_1} Q_{N_1} N_1^* + f_{PR} Q_P P^*)$ (Eq. 6A)

PIE: $-r - (f_{N_1} N_1^* + e_{IN_1} f_{IN_1} N_1^* I^*)$ (Eq. 6B)

PIEc: $\underbrace{-r - (f_{N_1} N_1^* + e_{IN_1} f_{IN_1} N_1^* I^*) - a_b}_{DE_E} + \underbrace{\frac{-e_I m_I + e_{IN_1} f_{IN_1} N_1 E}{e_I}}_{DE_I}$ (Eq. 6C)

Here, either the sum (IGP) or sum and product (PIE) of victim and enemy densities strengthens

DE (Eq. 6). Strong enough DE then tips the coexistence equilibrium from oscillations to stability.

In IGP the additive increase in predator and prey densities strengthens DE of resources (Eq. 6A).

Thus, strengthening of DE restores stability at higher f_{N_1} (Fig. 3D, black, solid line). In the PIE

models, however, increase in the (weighted) sum of parasites plus the product of parasite and

immune cells strengthens DE of energy at higher f_{N_1} (Eq. 6B-C; Fig. 3E-F, black, solid line).

Constitutive immunity enhanced stability in PIEc models by creating immune self-

limitation. This self-limitation generates additional loops featured in both intraspecific direct

effects (DE) and indirect (IE; Fig. 2). In PIEc, DE now has two components: DE of energy (DE_E)

and DE of immune cells (DE_I; Eq. 6C). The updated DE_E adds fixed energy allocation towards

baseline immunity ($- a_b$; Fig. 3F, orange, solid line [DE_E]). DE_I is always negative, too (see

Appendix Section 2C). This added immune self-limitation (Fig. 3F, green, solid line [negative

DE_I]) further strengthens DE (as shown in Fig. 3E: black, solid line), potentially enough to help to

eliminate even the possibility of oscillations in PIEc (eliminated in Fig. 3C; see Appendix Section

2C, Fig. A2). Constitutive immunity also elevates density of immune cells (I^*), lowering parasite

burden (N_1^*) while maintaining slightly higher E^* available to hosts for metabolic work (at rE^*;

compare Figs. 3H vs. 3I). Hence, constitutive immunity both stabilized dynamics via self-

limitation in PIEc (Fig. 2) and reduced parasite burden.

At low feeding rate in both IGP and PIE models, weakening of IE of the prey or parasite

triggers oscillations. IE is a ratio of shorter, binary loops (F_2) and longer, three-species loops (F_3)

(Fig. 2). In all three models, F_3 became more negative with feeding rate of the prey/parasite, f_{N_1},

than did F_2 (as parameterized). That helps explain why the ratio $-F_2/F_3$ (IE) became less negative

with increasing f_{N_1} – and hence why weakening IE triggered oscillations (Figs. 2, 3D-F). However,

the precise details varied among the models (so it becomes unfruitful to generalize here).

Positive feedback leads to alternate stable states in IGP alone (Fig. 4 and Appendix Section 2)

301   Despite the parallels described above, IGP model can also produce alterative stable states

not found in the with-host analogues. In fact, using different parameter values (Fig. 4A; see

Appendix Section 2A), a 2D bifurcation diagram shows two forms of alternate stable states in IGP

(Fig. 4A; Alt-SS1 [dark grey] and Alt SS-2 [light grey]; see also Verdy & Amarasekare 2010). In

these regions, initial densities of species determine competitive outcomes (Fig. 4C, D). The simpler

form of alternative stable states (Alt SS – 1; Fig. 4C) yields just the prey (R-N_1) or just the predator

(R-P) with the resource. If it held in PIE, analogous outcomes would lead to exclusion of immune

cells by parasites (yielding stable infection in the E-N_1 state) or vice versa (yielding immune

clearance, a stable E-I), depending on initial parasite dose. However, in PIE, the host cannot

generate immune cells without parasites. Consequently, since the E-I state is both biologically and

mathematically not feasible, the analogy breaks (Appendix Section 2A). In the other type of

alternative stable states in IGP (Alt SS – 2; Fig. 4D), we find either coexistence (R-N_1-P) or a

predator-only (R-P) equilibrium, depending on initial conditions (due to two feasible interior

equilibria, one of which is a saddle). Such outcomes would resemble a host that became infected

(E-N_1-I) or cleared infection (E-I), depending on parasite dose. We could not fine such an outcome

in PIEc (see Appendix Section 2C). Hence, despite the qualitatively similar loop structure, IGP

offered two scenarios for alternative stable states not present in within-host analogues. Alternative

infection states would require some other mechanism to produce positive feedback.

Two prey – predator - resource (KP) | Two parasite - immune cells - host energy (2PIE / 2PIEc)

Model Summary (*Fig. 1, Table 2, Appendix Section 3*)

Both keystone predation (KP) and within-host (2PIE) models feature two competitors

engaged indirectly via consumer-resource like interactions. In both, consumers directly benefit

(positive, black arrow) with direct costs to resources (negative, red circle). In KP, predators (P)

can consume two prey (N_i), leading to positive interspecific direct effects for the predator and

negative effects on prey (Fig. 1D). The two prey themselves compete for a shared resource (R),

creating similar +/- interactions. Additionally, the resource experiences self-limitation at the

interior (four species) equilibrium (Fig. 1D). Unlike in IGP, the predator does not consume the

resource (i.e., it is not omnivorous). Hence, competitors engage only in apparent and exploitative

competition (with P and R, respectively). Similarly, 2PIE and 2PIEc extend the within-host PIE

models. Now, two parasites (N_i) compete for host energy but are attacked by immune cells (Fig.

1E-F). During these attacks, immune cells use host energy to proliferate (as described in the PIE

models). In 2PIEc, hosts also directly allocate energy to production of immune cells ($a_b > 0$),

forging new connections and self-limitation of immune cells (Fig. 1F, Appendix Section 2).

Further, *E-I* links introduce additional components of loops not found in KP (e.g., in F_3; Fig 5).

Therefore, it might seem that KP provides too simple of a comparison to 2PIE/2PIEc. In fact, 2PIE

models best resemble a combination of KP and IGP. Yet, the feedback loops that determine

coexistence versus alternate stable states (F_4) of the prey or parasite are qualitatively similar (Fig.

5). The resemblance arises because both the prey and parasite are similarly *effected by* and have

affects on their resources and their enemy. However, stronger negative feedback at level 3 in 2PIEc

will shrink opportunities for coexistence while lowering parasite burden (see below).

 Intraspecific direct and indirect effects in the 4D KP and 2PIE model (Fig. 5)

Similar to 3D systems, the sign of intraspecific indirect effects determines coexistence vs.

priority effects in 4D systems. But first, in both KP and 2PIE, only self-limitation of the resource

or host energy, respectively, contribute to summed intraspecific direct effects (DE; level 1

feedback, F_1; Fig. 5). 2PIEc has additional contributions from the immune self-limitation. Then,

summed intraspecific indirect effects (IE) involve a ratio of feedback loops with three species (F_3)

and four (F_4; IE = $-F_3/F_4$; Fig 5; Appendix Section 3D). In KP and 2PIE models, F_3 sums the

negative consumer-resource loop between each victim and its enemy times self-limitation of the

resource (analogous to loop [i] in PIE [Fig. 2]). Then, energy-immune interactions add two sets of

additional loops, analogous to positive "*starving the enemy*" and negative "*feeding the enemy*"

loops in PIE, but for each parasite (loops [ii] and [iii], respectively). Finally, immune self-

limitation in 2PIEc adds two more negative feedback loops (involving $N_i - E$ for each parasite;

loops [iv]). At the interior, we find negative F_3 for the interiors evaluated here (algebraically or

numerically). Additionally, the sum of loops involving parasite N_2 provide the numerator of the

intraspecific indirect effect of N_1 (IE_{N_1}; blue shading), while those involving N_1 provide that of N_2

(IE_{N_2}: purple shading; Fig. 5). Unlike in IGP/ PIE, the intraspecific indirect effect of the enemy

and resource now becomes zero. Hence, only the sum of IE_{N_i} determines stability: coexistence

occurs when each competitor N_i indirectly decreases its own growth rate.

Importantly, level 4 feedback loops are qualitatively similar across the KP, 2PIE, and 2PIEc

models. These loops (from *L-R*) are: (i) a positive "*feeding the enemy*" loop, where a small

increase in one parasite stimulates the immune cells which attack the competing parasite, freeing

up energy for the first parasite. In a second positive loop, (ii) "*starving the enemy*", a small

increase in a parasite reduces resources, hence density of the competing parasite, thereby lowering

immune activation, ultimately reducing mortality on the first parasite. Those two positive

(destabilizing) loops then push against two negative (stabilizing) loops, (iii and iv) "*N_i is eaten, N_j*

eats". These loops therefore add the stabilizing consumer-resource interactions within which each

prey/parasite is enmeshed. For instance, a small increase in parasite *i* creates a negative loop with

the immune system while parasite *j* is braked by its loop with the resource. Then, those roles

reverse, i.e., parasite *j* is slowed by the immune system and parasite *i* is by the resource. Summed,

those two loops (iii and iv) jointly determine the amount of negative feedback in the system at

level 4. Combined then, loops (i)-(iv) determine the sign of F_4, the sign of the summed intraspecific

indirect effects ($-F_3/F_4$; since $F_3 < 0$: see above), and stability of a feasible interior equilibrium.

In these 4D systems, stability of the interior can also be understood via symmetries or

asymmetries in two quantities. These quantities emerge upon rearranging the loops comprising the

level 4 feedback (F_4). They reflect ratios of how each prey or parasite has effects on ($-\varepsilon_i$) and is

affected by ($-\alpha_i$) their enemies and resources (Fig 5; Eq. 7):

$$F_4 = -J_{N_1E}\, J_{N_2E}\, J_{EN_1}\, J_{EN_2}\, x \underbrace{\left(\frac{J_{IN_1}}{J_{EN_1}} - \frac{J_{IN_2}}{J_{EN_2}}\right)}_{effects\ on\ (-\varepsilon_i)}\, x \underbrace{\left(\frac{J_{N_1I}}{J_{N_1E}} - \frac{J_{N_2I}}{J_{N_2E}}\right)}_{affected\ by\ (-\alpha_i)} \qquad \text{(Eq. 7)}$$

where J_{ij} are interspecific direct effects (Jacobians) of species *j* on species *i*, $\varepsilon_1 = -J_{IN_1}/J_{EN_1}$, $\alpha_1 =$

$-J_{N_1I}/J_{N_1E}$, *etc.* Stability hinges on (a)symmetry of these ratios. As described elsewhere

(Appendix Sections 3A, B), the difference in *affected by* ratios ensures a tradeoff in traits

influencing resource and apparent competition that permits coexistence. If N_1 is the superior

resource competitor without the enemy, then $\alpha_1 > \alpha_2$ ensures N_2 is sufficiently resistant to attack.

The difference in *effects on* ratios then determines arrangements of competitive hierarchies with

enemies (involving exploitative [resource, energy] and apparent [predator, immune] competition).

Coexistence occurs with symmetry in these ratios (i.e., if $\alpha_1 > \alpha_2$ if $\varepsilon_1 > \varepsilon_2$), guaranteeing that net

negative feedbacks dominate ($F_4 < 0$). In contrast, asymmetry in these ratios, (e.g., $\alpha_1 > \alpha_2$ still but

now $\varepsilon_1 < \varepsilon_2$) leads to net positive feedback ($F_4 > 0$), triggering alternate stable states.

Coexistence (coinfection) and alternate stable states (priority effects) in KP & 2PIE (Fig. 6 and

Appendix Section 3)

Despite differences in biology, KP and 2PIE models share similar four species feedback loops, hence qualitatively similar outcomes for stability of the interior equilibrium. That similarity becomes readily apparent in 2D bifurcation diagrams for each model in parameter space of supply point (S) – N_1's feeding rate on resources (f_{N_1}). In KP (Fig. 6A), at low S-f_{N_1}, only the resource is supported in the system (R, yellow). Increased feeding rate at low resource supply enables prey N_1 to meet its minimum resource requirement (R-N_1; light blue). As S increases, the minimum prey requirement of the predator is met (its N_1^*) allowing predators to invade (R-N_1-P; dark blue). Then, the more resistant N_2 can invade when the R-N_1-P food chain provides enough resources, given predator density (mortality). This invasion enables four species coexistence (R-N_1-N_2-P; light orange). Similar assembly rules apply for invasion of N_2 when feeding rate of N_1 stays low enough to grant N_2 competitive superiority ($R_2^* < R_1^*$). With increasing S, first N_2 invades (R-N_2 space; light purple), then the predator does (R-N_2-P: dark purple). A less typical assembly arises at lower S ($S = 1$) but with increasing f_{N_1}. Here, N_1 and P simultaneously invade creating the jump from 2 (R-N_2) to 4 (R-N_1-N_2-P) dimensional stability (see Appendix Section 3C for details). Overall, similar transitions appear in 2PIE models. Like KP, the possible states in 2PIE models are energy alone (E), just one parasite (E-N_i), addition of immune cells (E-N_i-I), or an interior (E-N_1-N_2-I; Fig. 6B). However, since hosts always allocate energy to immune cells, they are always present in 2PIEc. Hence, 2PIEc only has *E-I, E-I-N_i,* and *E-N_1-N_2-I* states (but not E or E-N_i; Fig. 6C).

In all three models, low or high combinations of supply point (S) and feeding rate (f_{N_1}), create (a)symmetry in *effects on* and *affected by* ratios that, in turn, generate coexistence or alternate stable states for a feasible interior. We assume N_1 competes superiorly for energy without immune cells but is sufficiently more vulnerable to immune attack (enabling $\alpha_1 > \alpha_2$). In the coexistence region (i.e., at lower f_{N_1}- S), N_1 is the superior apparent competitor while more resistant N_2 is the

superior resource competitor (see Appendix Section 3A,B for details and proof). Such a switch in

competitive hierarchy means N_1 exerts stronger effects on immune cells relative to energy ($\varepsilon_1 > \varepsilon_2$;

Eq. 7). It also produces a symmetry in ratios that generates net negative feedback ($F_4 < 0$) and

negative intraspecific indirect effects of the competitors ($IE_i < 0$) enabling coexistence

(coinfection; 6D-F). However, high f_{N_1}- S produces an asymmetry: N_2 now is the superior

apparent competitor, and it has stronger effects on immune cells relative to energy ($\varepsilon_1 < \varepsilon_2$) while

N_1 is the superior resource competitor. This asymmetry in ratios generates net positive feedback

($F_4 > 0$) triggering alternate stable states (or priority effects; Appendix Section 3A,B). Hence, this

interior saddle separates dominance by one or the other parasite (or prey). Furthermore,

intraspecific indirect effects are both positive ($IE_{N_i} > 0$): each competitor now benefits itself through

the feedback loops. With even higher f_{N_1}, N_2 becomes excluded in all three models (Fig. 6G-I).

Notice also how, at the border of that exclusion from the coexistence region, IE_{N_2} becomes

extremely strong (infinitely negative, since $F_4 = 0$; Fig. 6D-F). In other words, the excluded species

suffers extremely strong negative intraspecific indirect effects (i.e., $IE_{N_2} \rightarrow -\infty$).

The 2PIEc model modifies the patterns in 2PIE in qualitative and quantitative ways.

Qualitatively, the range of outcomes in S-f_{N_1} space simplifies (as in PIEc: Fig. 4C). Since baseline

allocation guarantees positive density of immune cells, there are no E or E-N_i regions like in 2PIE

(Fig. 6B vs. 6C). Quantitatively, in the co-infection region, constitutive immunity strengthens

intraspecific indirect effects (because of the added negative feedback in the F_3 loops, the numerator

of IE of each species [Fig. 5]). Stronger IE, in turn, narrows the parameter space permitting

coexistence. Constitutive immunity also elevates density of immune cells (I^*), lowering parasite

burden ($N_1^* + N_2^*$) while maintaining slightly higher E^* (allocated at rE^*) for other metabolic work

(compare Figs. 6H vs. 6I). Hence, constitutive immunity adds to F_3 loop structure of 2PIEc (Fig.

5). That addition squeezes coinfection space while reducing parasite load relative to 2PIE.

DISCUSSION

What within-host factors determine parasite dynamics? To answer this question, we compare two classic food web modules, intraguild predation (IGP) and keystone predation (KP), to their within-host analogues (PIE and 2PIE). On the one hand, the comparison seems apt. Both prey and parasite consume a shared resource while facing mortality from an enemy (predator or immune cell). On the other hand, predator and immune cells are produced rather differently. For instance, predators can eat only prey or the resource or both (in IGP). In contrast, induced proliferation of immune cells requires both energy and parasites simultaneously. Furthermore, hosts can allocate energy to immunity constitutively (a pipeline uncommon to food webs). Given these similarities and differences, we analyze stability of equilibria produced by each model using feedback loops. In particular, we interpret loops as direct and indirect intraspecific effects, and we show how their sign and strength govern stability outcomes (Box 1). With these tools, we make three points. First, both IGP and PIE systems predict stable coexistence / infection or oscillations due to similar shifts in direct and indirect effects (Fig. 2-3). Second, that enemy-generation difference eliminates alternative stable states (priority effects) seen in IGP – the PIE models cannot produce them without inclusion of some other mechanisms (Fig. 4). Third, despite that KP is structurally simpler than 2PIE models, competing prey and parasites coexist (co-infect) or show priority effects for essentially the same reasons (Fig. 5-6). The outcomes hinge on the sum of similar four-dimensional loops that generate positive and negative feedback. We show how those loops, in turn, translate into parallel symmetries (stabilizing) or asymmetries in *effects on* and *affected by* ratios of the two competitors. Overall, food web models offer powerful if imperfect analogies to feedbacks underlying the dynamical repertoire of parasites within hosts.

Feedback loops enabled biologically meaningful comparisons of the stabilizing and destabilizing forces in the models analysed here. Traditionally, one evaluates (local) stability using Routh-Hurwitz criteria (Routh 1877; Hurwitz 1895). By rearranging these criteria, we unpack the

biology of the feedback loops underlying stability (Box 1; Puccia & Levins 1991; Novak *et al*

2016). These loops can involve shorter, direct effects (DE), where increases in intraspecific density

leads to self-limitation (negative feedback) or facilitation (positive). Longer loops involve chains

of connected interactions – here, between two (e.g., binary consumer-resource), three, or four

species (Puccia & Levins 1991). We show that certain ratios of these loops at different feedback

levels, in turn, correspond to intraspecific indirect effects (IE – falling along the trace of the inverse

Jacobian matrix evaluated at the equilibrium; Box 1). Furthermore, we illustrate how negative IE

leads to coexistence (stable or oscillatory [in 3D systems]), and positive IE can produce alternative

stable states. In 3D systems, that transition from stability to oscillations involves weakening of DE

x IE. Such weakening occurs when feedback at longer loops becomes too strong relative to those

at shorter loops. With that framework in mind, the loops then facilitated comparison between

structurally similar but biologically distinct systems (Novak *et al* 2016), *e.g.,* food web and within-

host modules. Indeed, loops can reveal the biology obscured or overlooked in stability analysis.

Single infection dynamics

This loop-based approach to stability analysis revealed that single prey and parasite can persist

– stably or via oscillations – with their resource and enemy for similar reasons. Generally speaking,

like in IGP, coexistence in PIE requires summed negative intraspecific indirect effects (IE). Like

in IGP, weakening of IE ($-F_2/F_3$) triggers oscillations in PIE (due to weakening of F_3 relative to

F_2; Fig. 3D-E). Yet, either the sum (IGP) or sum and product (PIE) of victim and enemy densities

strengthens intraspecific direct effects (DE). Strong enough DE then tips the coexistence

equilibrium from oscillations to stability. Those oscillations matter because they create boom-bust

cycles of parasite and immune response that can harm hosts (Otterstatter & Thomson 2006; Childs

& Buckee 2015). For instance, these oscillations produce rapid ramp up or down of immune

function that can stress host physiology (exacting e.g., oxidative damage, cellular senescence),

making them sicker (Costantini & Mølle 2009). However, in hosts with high constitutive immunity

(PIEc), the strengthening of DE by immune-self limitation can prevent oscillations, potentially

reducing immune damage to host. Additionally, constitutive immunity can improve host health by

lowering parasite burden while maintaining higher energy for other metabolic work.

Despite similarities in loop structure, alternate stable states emerge in IGP but not PIE models.

The reason for this discrepancy hinges on production of the enemies. In principle, when each prey

/ parasite *"starves the enemy"* enough, the resulting positive feedback through the three species

might trigger alternative stable states (alt SS; Fig. 2). In fact, IGP with a semi-chemostat like

resource produces two types of them (Verdy & Amarasekare 2010). In the simpler, more typical

one, either the prey or the predator dominates. In the other less common one, predator and prey

coexist or the prey is excluded (Fig. 4A). If they existed, such alt SS in PIE models might explain

why large infectious doses overwhelm immune clearance, leading to infection (yielding stable E-

N_I or E-N_I-I states), while small doses can be cleared by the immune system (in a stable E-I state;

as seen in experiments, e.g., Merrill & Cáceres 2018). Yet, we found no such alt SS in PIE or PIEc

as formulated (nor did it arise in a similar model with another representation of immune

functioning: Greenspoon *et al.* 2018). Hence, some other mechanism is needed to generate alt SS

within hosts (e.g., effects of parasites on within-host resource supply points: van Leeuwan *et al*.

2019). In the future, such mechanisms could be added to the PIE/PIEc framework. However, they

would likely break the straightforward food web analogues that we strove for here.

Coinfection dynamics

In contrast, KP and 2PIE models produced either coexistence or priority effects of competitors

for similar reasons. At first glance, such parallels might seem surprising since KP more simply

connects species engaged in resource and apparent competition. In contrast, 2PIE has more

complex loop structure (created by those omnivory-like I-E connections). Despite these structural

differences, both models share qualitatively identical four species loops that govern stability (Fig.

5). We translated the net sum of those positive and negative loops into differences in *effects on*

and *affected by* ratios for each competitor. Those ratios then lay out conditions for successful

coinfection *vs.* alternative stable states (priority effects). First, either case requires a sufficient

tradeoff. If one species competes superiorly for host energy, the other must better resist immune

attack. Such a tradeoff anchors a directionality of the *affected by* ratio (immune to energy) between

competing parasites. Coinfection (and coexistence more broadly), then, requires a symmetry: the

parasite with greater *affected by* ratio must also have the greater *effects on* ratio (Fig. 5). Those

conditions arise when the superior energy competitor without immune cells (i.e., the one with

lower minimal energy needs) becomes the superior apparent competitor with them (i.e., the one

that supports and withstands the highest immune density). Meanwhile, the more resistant parasite

becomes the best energy competitor. If true, each parasite indirectly inhibits its own growth and

facilitates its competitor. Coinfection ensues due to net negative feedback. In contrast, if one

parasite always competes superiorly for energy while the superior apparent competitor enjoys

resistance, asymmetry in these ratios ensues. That asymmetry means each competitor indirectly

facilitates its own growth and slows its competitor, either by *"feeding the immune cells"* that attack

the competitor (freeing up energy), or by *"starving the immune cells"* via starving the competitor

(Fig. 5). When strong enough, such interactions ensure priority effects via net positive feedbacks.

These new within-host competition models can also guide future coinfection experiments.

First, the competition models focus attention on traits and mechanism. Traditional coinfection

experiments alter initial densities / order of arrival of parasites within a host (Lohr *et al* 2010;

Natsopoulou *et al* 2015; *reviewed in* Karvonen *et al* 2019). Then, mechanism is inferred from

pattern. Alternatively, with 2PIE-like models in hand now, experimenters can directly measure

and/or fit key host-parasite traits that produce within-host dynamics. Second, they highlight how

multiple niche dimensions govern competitive outcomes within hosts. Some recent empirical

studies emphasized immune- (Ezenwa *et al* 2010; Halliday *et al* 2018) or resource-mediated

(Budischak *et al* 2018) competition within hosts. Yet, competition along a single niche dimension

predicts only competitive exclusion (*sensu stricto*). Instead, measurements of both immune and

energetic niche dimensions together (like in Budischak *et al* 2015) could evaluate the entire range

of outcomes anticipated by 2PIE. Third, the bifurcation maps here suggest joint manipulation of

resource supply to hosts and parasite traits, perhaps leveraging genetic variation in traits among

parasite strains. Such manipulations could then capture a whole array of within-host competitive

outcomes (Budischak *et al* 2018) or explain increasing parasite burden along a resource gradient

(Wale *et al* 2017). Finally, our model predicts that allocation to constitutive immunity strengthens

indirect effects, squeezing parameter space allowing coinfection and reducing parasite burden. One

could test such predictions using strains with immune knockouts (like in Chen *et al* 2020) or host

genotypes differing in allocation (Fuess *et al* 2020).

Future directions and conclusion

This within-host parasite framework could become expanded in the future. First, models could

add niche dimensions. For instance, parasites might compete for multiple within-host resources

(Budishack *et al* 2015; Ezenwa 2021) and/or face multiple immune defenses (Jolles *et al* 2008;

Ezenwa & Jolles 2011). These additional dimensions parallel food webs with multiple resources

and predators (necessitating statements of clear assumptions about specialist vs. generalist

consumption, different possible tradeoffs, etc.: Hulot & Loreau 2006). These niche dimensions

might expand conditions promoting coinfection or priority effects (Graham 2008; Rynkiewicz *et*

al 2015). Second, other mechanisms like relative non-linearities (Armstrong & McGehee 1980),

competitive intransitivity (May & Leonard 1975), or other variation-based mechanisms (Chesson

2000) may predict successful coexistence in our within-host framework. For example, since PIE

can oscillate, two parasites might coexist via oscillations (relative non-linearities). Similarly,

multiple parasites competing intransitively for two or more resources might coexist (Huisman &

Weissing 2001). Finally, environmental variation could alter key within-host traits (*e.g.,*

temperature fluctuations can modulate host immunity and parasite attack rates [Scharsack *et al*

2016]). Such intersections of environment with trait plasticity might then alter or enhance

opportunities for parasite coexistence (Chesson 2000). Finally, the direct (DE) vs. indirect effect

(IE) approach to predicting oscillations may have limits. Perhaps theoreticians can extend it

beyond three to four or more dimensions. Together, these expansions would extend mechanistic

models of within-host dynamics and produce insight into disease and coexistence alike.

In this study, we gleaned insights for within-host parasite dynamics using food web modules

and feedback loops. Feedback loops enabled comparison of stability outcomes in structurally

similar but biological distinct systems. For instance, similarity in loop structure in IGP and PIE

successfully predicted single prey / parasite persistence either stably or via oscillations. Yet,

despite these structural similarities, differences in generation of enemies (predator vs. immune

cells) prevented IGP's two alternate stable states from arising in PIE. In contrast, despite

omnivory-like *I-E* connections in 2PIE, competing prey or parasites coexist (coinfect) or show

priority effects like in KP. Those outcomes arose because competitors within hosts or in food webs

engage in structurally similar resource and apparent competition. Such comparable structure meant

that symmetries in *effects on* and *affected by* ratios (involving the enemy and resource) determined

stability of competition. Hence, dynamical forces governing stability in food-webs can mirror

those within hosts. Such insights can guide more predictive experiments at the within-host scale

but also better population-level models. For instance, fluctuation in food supply could shift

competitive outcomes within hosts that then shifts multi-parasite outbreaks at the population scale.

Hence, further development of resource and immune-explicit frameworks can only enhance

predictive insight into host health and disease outbreaks alike.

ACKNOWLEDGMENTS

AR was supported by the Department of Biology, Indiana University – Bloomington. The study was supported in part by NSF DEB (grant 1655656). We thank F. Bashey-Visser, C.M Lively, and members of the Hall lab for their valuable feedback.

REFERENCES

Alizon, S., & van Baalen, M. (2008). Multiple infections, immune dynamics, and the evolution of virulence. The American Naturalist, 172(4), E150-E168.

Armstrong, R. A., & McGehee, R. (1980). Competitive exclusion. The American Naturalist, 115(2), 151-170.

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the maintenance of parasite diversity. Philosophical Transactions of the Royal Society B: Biological Sciences, 370(1675), 20140301.

Budischak, S. A., Wiria, A. E., Hamid, F., Wammes, L. J., Kaisar, M. M., van Lieshout, L., ... & Graham, A. L. (2018). Competing for blood: the ecology of parasite resource competition in human malaria–helminth co-infections. *Ecology letters*, *21*(4), 536-545.

Budischak, S. A., Sakamoto, K., Megow, L. C., Cummings, K. R., Urban Jr, J. F., & Ezenwa, V. O. (2015). Resource limitation alters the consequences of co-infection for both hosts and parasites. International Journal for Parasitology, 45(7), 455-463.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. Annual review of Ecology and Systematics, 31(1), 343-366.

Chen, C. C., Louie, S., McCormick, B., Walker, W. A., & Shi, H. N. (2005). Concurrent infection with an intestinal helminth parasite impairs host resistance to enteric Citrobacter

rodentium and enhances Citrobacter-induced colitis in mice. *Infection and immunity*, *73*(9),

5468-5481.

Childs, L. M., & Buckee, C. O. (2015). Dissecting the determinants of malaria chronicity: why

within-host models struggle to reproduce infection dynamics. *Journal of The Royal Society*

Interface, *12*(104), 20141379.

Clay, P. A., Cortez, M. H., Duffy, M. A., & Rudolf, V. H. (2019). Priority effects within

coinfected hosts can drive unexpected population-scale patterns of parasite prevalence. *Oikos*,

128(4), 571-583.

Costantini, D., & Møller, A. P. (2009). Does immune response cause oxidative stress in birds? A

meta-analysis. Comparative Biochemistry and Physiology Part A: Molecular & Integrative

Physiology, 153(3), 339-344.

Cressler, C. E., Nelson, W. A., Day, T., & McCauley, E. (2014). Disentangling the interaction

among host resources, the immune system and pathogens. Ecology letters, 17(3), 284-293.

De Roode, J. C., Helinski, M. E., Anwar, M. A., & Read, A. F. (2005). Dynamics of multiple

infection and within-host competition in genetically diverse malaria infections. *The American*

Naturalist, *166*(5), 531-542.

Doublet, V., Natsopoulou, M. E., Zschiesche, L., & Paxton, R. J. (2015). Within-host

competition among the honey bees pathogens Nosema ceranae and Deformed wing virus is

asymmetric and to the disadvantage of the virus. *Journal of invertebrate pathology*, *124*, 31-34.

Duneau, D., Ferdy, J. B., Revah, J., Kondolf, H., Ortiz, G. A., Lazzaro, B. P., & Buchon, N.

(2017). Stochastic variation in the initial phase of bacterial infection predicts the probability of

survival in D. melanogaster. *Elife*, *6*, e28298.

Ezenwa, V. O. (2021). Co-infection and nutrition: integrating ecological and epidemiological

perspectives. In *Nutrition and Infectious Diseases* (pp. 411-428). Humana, Cham.

Ezenwa, V. O., & Jolles, A. E. (2011). From host immunity to pathogen invasion: the effects of

helminth coinfection on the dynamics of microparasites. *Integrative and comparative biology*,

51(4), 540-551.

Ezenwa, V. O., Etienne, R. S., Luikart, G., Beja-Pereira, A., & Jolles, A. E. (2010). Hidden

consequences of living in a wormy world: nematode-induced immune suppression facilitates

tuberculosis invasion in African buffalo. *The American Naturalist*, *176*(5), 613-624.

Fenton, A., & Perkins, S. E. (2010). Applying predator-prey theory to modelling immune-

mediated, within-host interspecific parasite interactions. *Parasitology*, *137*(6), 1027-1038.

Fuess, L. E., Weber, J. N., den Haan, S., Steinel, N. C., Shim, K. C., & Bolnick, D. I. (2020). A

test of the Baldwin Effect: Differences in both constitutive expression and inducible responses to

parasites underlie variation in host response to a parasite. *bioRxiv*.

Gorsich, E. E., Etienne, R. S., Medlock, J., Beechler, B. R., Spaan, J. M., Spaan, R. S., ... &

Jolles, A. E. (2018). Opposite outcomes of coinfection at individual and population scales.

Proceedings of the National Academy of Sciences, *115*(29), 7545-7550.

Graham, A. L. (2008). Ecological rules governing helminth–microparasite coinfection.

Proceedings of the National Academy of Sciences, 105(2), 566-570.

Greenspoon, P. B., Banton, S., & Mideo, N. (2018). Immune system handling time may alter the

outcome of competition between pathogens and the immune system. *Journal of theoretical*

biology, *447*, 25-31.

Griffiths, E. C., Fairlie-Clarke, K., Allen, J. E., Metcalf, C. J. E., & Graham, A. L. (2015).

Bottom-up regulation of malaria population dynamics in mice co-infected with lung-migratory

nematodes. Ecology letters, 18(12), 1387-1396.

Halliday, F. W., Umbanhowar, J., & Mitchell, C. E. (2018). A host immune hormone modifies

parasite species interactions and epidemics: insights from a field manipulation. *Proceedings of*

the Royal Society B, *285*(1890), 20182075.

Hite, J. L., & Cressler, C. E. (2018). Resource-driven changes to host population stability alter

the evolution of virulence and transmission. Philosophical Transactions of the Royal Society B:

Biological Sciences, 373(1745), 20170087.

Holt, R. D., & Polis, G. A. (1997). A theoretical framework for intraguild predation. The

American Naturalist, 149(4), 745-764.

Holt, R. D., Grover, J., & Tilman, D. (1994). Simple rules for interspecific dominance in systems

with exploitative and apparent competition. The American Naturalist, 144(5), 741-771.

Hulot, F. D., & Loreau, M. (2006). Nutrient-limited food webs with up to three trophic levels:

feasibility, stability, assembly rules, and effects of nutrient enrichment. Theoretical Population

Biology, 69(1), 48-66.

Hurwitz, A. (1895). On the conditions under which an equation has only roots with negative real

parts. *Math. Ann.* 46, 273-284 (1895)

Jolles, A. E., Ezenwa, V. O., Etienne, R. S., Turner, W. C., & Olff, H. (2008). Interactions

between macroparasites and microparasites drive infection patterns in free-ranging African

buffalo. *Ecology*, *89*(8), 2239-2250.

Huisman, J., & Weissing, F. J. (2001). Fundamental unpredictability in multispecies

competition. *The American Naturalist*, *157*(5), 488-494.

Karvonen, A., Jokela, J., & Laine, A. L. (2019). Importance of sequence and timing in parasite

coinfections. *Trends in parasitology*, *35*(2), 109-118.

Leibold, M. A. (1996). A graphical model of keystone predators in food webs: trophic regulation

of abundance, incidence, and diversity patterns in communities. The American Naturalist,

147(5), 784-812.

Lohr, J. N., Yin, M., & Wolinska, J. (2010). Prior residency does not always pay off–co-

infections in Daphnia. Parasitology.

May, R. M., & Leonard, W. J. (1975). Nonlinear aspects of competition between three species.

SIAM journal on applied mathematics, 29(2), 243-253.

Merrill, T. E. S., & Cáceres, C. E. (2018). Within-host complexity of a plankton-parasite

interaction. Ecology, 99(12), 2864-2867.

Metcalf, C. J. E., Grenfell, B. T., & Graham, A. L. (2020). Disentangling the dynamical

underpinnings of differences in SARS-CoV-2 pathology using within-host ecological models.

PLoS pathogens, *16*(12), e1009105.

Mideo, N., Alizon, S., & Day, T. (2008). Linking within-and between-host dynamics in the

evolutionary epidemiology of infectious diseases. Trends in ecology & evolution, 23(9), 511-

517.

Natsopoulou, M. E., McMahon, D. P., Doublet, V., Bryden, J., & Paxton, R. J. (2015).

Interspecific competition in honeybee intracellular gut parasites is asymmetric and favours the

spread of an emerging infectious disease. *Proceedings of the Royal Society B: Biological*

Sciences, *282*(1798), 20141896.

Novak, M., Yeakel, J. D., Noble, A. E., Doak, D. F., Emmerson, M., Estes, J. A., ... & Wootton,

J. T. (2016). Characterizing species interactions to understand press perturbations: what is the

community matrix?. *Annual Review of Ecology, Evolution, and Systematics*, *47*, 409-432.

Otterstatter, M. C., & Thomson, J. D. (2006). Within-host dynamics of an intestinal pathogen of

bumble bees. *Parasitology*, *133*(6), 749.

Puccia, C. J., & Levins, R. (1991). Qualitative modeling in ecology: loop analysis, signed

digraphs, and time averaging. In Qualitative simulation modeling and analysis (pp. 119-143).

Springer, New York, NY.

Råberg, L. (2012). Infection intensity and infectivity of the tick-borne pathogen Borrelia afzelii.

Journal of evolutionary biology, *25*(7), 1448-1453.

Råberg, L., Graham, A. L., & Read, A. F. (2009). Decomposing health: tolerance and resistance

to parasites in animals. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

364(1513), 37-49.

Routh, E. J. (1877). *A Treatise on the Stability of a Given State of Motion, Particularly Steady*

Motion: Being the Essay to which the Adams Prize was Adjudged in 1877, in the University of

Cambridge. Macmillan and Company.

Rynkiewicz, E. C., Pedersen, A. B., & Fenton, A. (2015). An ecosystem approach to

understanding and managing within-host parasite community dynamics. *Trends in parasitology*,

31(5), 212-221.

Scharsack, J. P., Franke, F., Erin, N. I., Kuske, A., Büscher, J., Stolz, H., ... & Kalbe, M. (2016).

Effects of environmental variation on host–parasite interaction in three-spined sticklebacks

(Gasterosteus aculeatus). *Zoology*, *119*(4), 375-383.

Susi, H., Barrès, B., Vale, P. F., & Laine, A. L. (2015). Co-infection alters population dynamics

of infectious disease. Nature communications, 6(1), 1-8.

Van Leeuwen, A., Budischak, S. A., Graham, A. L., & Cressler, C. E. (2019). Parasite resource

manipulation drives bimodal variation in infection duration. Proceedings of the Royal Society B,

286(1902), 20190456.

Verdy, A., & Amarasekare, P. (2010). Alternative stable states in communities with intraguild

predation. Journal of Theoretical Biology, 262(1), 116-128.

Wodarz, D. (2006). Ecological and evolutionary principles in immunology. Ecology Letters,

9(6), 694-705.

Wolfram Research, Inc., Mathematica, Version 12.2, Champaign, IL (2020)

TABLES

Table 1. Equations for IGP (Eq. 1) and PIE models (PIE and PIEc constitutive immunity; Eq. 2) with top consumers (predators or immune cells), intermediate consumers (prey or parasite), and basal resource (or host energy). Listed below each model are the variables and parameters, their definition (with units), and corresponding default values used (in Figs. 3,4).

IGP			PIE and PIEc	
$\frac{dP}{dt} = (e_{PN_1} f_{PN_1} N_1 + f_{PR} R - m_P) P$ (Eq. 1A)			$\frac{dI}{dt} = \frac{a_b E + e_{IN_1} f_{IN_1} N_1 E I}{e_I} - m_I I$ (Eq. 2A)	
$\frac{dN_1}{dt} = (f_{N_1} R - f_{PN_1} P - m_N) N_1$ (Eq. 1B)			$\frac{dN_1}{dt} = \left(\frac{f_{N_1} E}{e_{N_1 E}} - f_{IN_1} I - m_N\right) N_1$ (Eq. 2B)	
$\frac{dR}{dt} = a(S - R) - (f_{N_1} Q_{N_1} N_1 + f_{PR} Q_P P) R$ (Eq. 1C)			$\frac{dE}{dt} = \frac{f_E S}{h+S} - rE - (f_{N_1} N_1 E + e_{IN_1} f_{IN_1} N_1 I)E - a_b E$ (Eq. 2C)	

		Val.			Val.
Var	**Definition**	**Fig. 3[a]**	**Fig. 4[a]**		
Var	**Definition**				**Fig. 3**

Var	Definition	Fig. 3[a]	Fig. 4[a]	Var	Definition	Fig. 3
P	Predator density (#·L^{-1})			I	Immune cell density (L^{-1})	
N_1	Prey density (#·L^{-1})			N_1	Parasite density (L^{-1})	
R	Resource density (#·L^{-1})			E	Host energy density (L^{-1})	
e_{PN_1}	Conversion, prey: predator	31.61	0.75	e_{IN_1}	Induction of energy per parasite killed by the immune response (L)	4
f_{PN_1}	Feeding rate, predator on prey, (L·day^{-1})	0.002	0.002	f_{IN_1}	Killing rate of parasite, per immune cell (L . time^{-1})	12
f_{PR}	Feeding rate, prey on resource (L . day^{-1})	0.0005	0.01	m_I	Immune cells background mortality rate (time^{-1})	24
m_P	Mortality rate, predator (day^{-1})	0.05	0.05	e_I	Cost of producing immune cell	3

Symbol	Description			Symbol	Description	
f_{N_1}	Feeding rate, prey on resource (L·day^{-1})	var[b]	var[b]	e_{N_1E}	Cost of producing parasite	30
m_N	Mortality rate, prey (day^{-1})	0.05	0.05	f_{N_1}	Feeding rate per parasite on energy (L . time^{-1})	var[b]
a	Dilution rate (day^{-1})	0.1	0.1	m_N	Mortality rate, parasite (time^{-1})	6
S	Supply point (L^{-1})	var[b]	var[b]	f_E	Host resource ingestion rate (L^{-1} . time^{-1})	44
Q_P	Conversion, resource: predator	1	0.5	S	Food supply point, hosts (L^{-1})	var[b]
Q_{N_1}	Conversion resource: prey	1	2	h	Resource ingestion half-saturation constant, host (L^{-1})	19
				r	Rate of host energy use for non-parasite purposes (time^{-1})	1
				a_b	Baseline allocation to immunity (time^{-1})	PIE: 0 PIE$_c$: 0.03

810
811   [a] Values chosen in Fig. 3 such that $Q_{N_1} < e_{PN_1} Q_P$ (ensuring $F_3 < 0$; eqs. A14, 15). In fig. 4, Q_{N_1}
812    $> e_{PN_1} Q_P$, enabling alternative stable states ($F_3 > 0$ for interior).   [b] See values on figures.
813
814

815   **Table 2.** Equations for KP (Eq. 3) and 2PIE models (2PIE and 2PIEc constitutive immunity; Eq.

816   4) with top consumers (predators or immune cells), intermediate consumers (prey or parasites),

817   and basal resource (or host energy). Listed below each model are the variables and parameters,

818   their definition (with units), and corresponding default values used (in Figs. 6).

KP	2PIE and 2PIEc
$\frac{dP}{dt} = \left(\Sigma e_{PN_i} f_{PN_i} N_i - m_P\right) P$ (Eq. 3A)	$\frac{dI}{dt} = \frac{a_b E + \Sigma e_{IN_i} f_{IN_i} N_i E\, I}{e_I} - m_I I$ (Eq. 4A)
$\frac{dN_i}{dt} = \left(e_{N_i R} f_{N_i} R - f_{PN_i} P - m_N\right) N_i$ (Eq. 3B)	$\frac{d N_i}{dt} = \left(\frac{f_{N_i} E}{e_{N_i E}} - f_{IN_i} I - m_N\right) N_i$ (Eq. 4B)
$\frac{dR}{dt} = a(S - R) - \Sigma(f_{N_i} N_i)\, Q\, R$ (Eq. 3C)	$\frac{dE}{dt} = \frac{f_E S}{h+S} - rE - \left(\Sigma e_{IN_i} f_{IN_i} N_i I + \Sigma f_{N_i} N_i\right)E - a_b E$ (Eq. 4C)

Var	Definition	Values Fig. 6[a]	Var	Definition	Values Fig. 6[a]
P	Predator density (#·L^{-1})		I	Immune cells density (L^{-1})	
N_i	Prey i density (L^{-1})		N_i	Parasite i density (L^{-1})	
R	Resource density (L^{-1})		E	Host energy density (L^{-1})	
e_{PN_i}	Conversion, prey i: predator	2.4, 4.5	e_{IN_i}	Induced energy allocation rate per parasite i killed by the immune response (L)	3, 4.7
f_{PN_i}	Feeding rate, predator on prey i, (L·day^{-1})	9.8, 2.1	f_{IN_i}	Killing rate, immune cell on parasite i (L . time^{-1})	0.9, 0.4
$e_{N_i R}$	Conversion, resource: prey i	2.8, 2.8	m_I	Immune cells background mortality rate (time^{-1})	0.4
m_P	Mortality rate, predator (day^{-1})	0.1	e_I	Cost of producing immune cell	2.5

f_{N_i}	Feeding rate, prey on resource (L·day⁻¹)	var[b], 10.4	e_{N_iE}	Cost of producing parasite i	4, 3.4
m_N	Mortality rate, prey (day⁻¹)[b]	0.1	f_{N_i}	Feeding rate per parasite i on energy (L . time⁻¹)	var[b], 7.2
a	Dilution rate (day⁻¹)	7.9	m_N	Parasite background mortality rate (time⁻¹)	0.8
S	Supply point (L⁻¹)	var[b]	f_E	Host resource ingestion rate L⁻¹ . time⁻¹)	7.8
Q	Stoichiometric conversion, resource: prey	2.8[c]	S	Food supply point, hosts (L⁻¹)	var[a]
			h	Resource ingestion half-saturation constant (L⁻¹)	3.2
			r	Rate of host energy use for non-parasite purposes (time⁻¹)	3.7
			a_b	Baseline energy allocation to immunity (time⁻¹)	2PIE: 0 2PIEc:0.1

819 [a] Values chosen to ensure N_1 is the superior competitor for resources without enemies (prey: $R_1{}^*$
820 $< R_2{}^*$ [eq. A35]; parasite: $E_1{}^* < E_2{}^*$ [eq. A47.a]) while N_2 is more resistant to attack to enemies
821 (prey: $f_{PN_1} > f_{PN_2}$; parasites: $f_{IN_1} > : f_{IN_2}$). Additionally, they guarantee the *effects on* ratio has
822 structure $\alpha_1 > \alpha_2$ (eqs. A34, A46). [b] See values on figures. [c] Identical between prey i.
823

    FIGURE CAPTIONS

Figure 1: *Direct interspecific effects and self-limitation in food web and within-host*

*parasite model*s. *Top row, single prey or parasite models (three-dimension, 3D)*: (A) one prey-

predator-resource (intraguild predation, IGP), (B) one parasite-immune cells-host energy (PIE),

and (C) PIEc with constitutive immunity (i.e., fixed allocation of energy to production of immune

cells). *Bottom row, two prey or two parasite models (four dimension, 4D):* (D) two prey - predator-

resource (keystone predation, KP), (E) two parasite - immune cells - host energy (2PIE), and (F)

2PIEc with constitutive immunity. All direct effects evaluated when densities are positive (i.e., at

a feasible interior equilibrium). Red (black) arrows: negative (positive) interspecific direct effect;

red curve: negative intraspecific specific effect.

Figure 2: *Intraspecific direct and indirect effects: an illustration of levels of feedback in*

IGP, PIE, and PIEc models (Fig. 1A-C). Evaluated at a feasible interior equilibrium, the IGP and

PIE models have similar feedback structure. PIEc adds additional loops involving immune self-

limitation. Level 1 feedback (F_1) is the sum of the shorter intraspecific direct effects (solid curves):

self-limitation (-) of resource/energy and immune cells (PIEc only). In intraspecific indirect effects

(dashed curves), each species limits or facilitates itself via interaction with other species in longer

loops. The intraspecific indirect effect (IE) sums that of energy, E (IE_E; yellow shading), parasite

N_1 (IE_{N_1}; blue), and immune cells I (IE_I; green). IE, in turn, is a ratio of two levels of feedback.

Level 2 feedback (F_2, numerator of IE) sums pairwise consumer-resource loops (-) and the product

of energy and immune self-limitation (-; only in PIEc; these are the numerators of the IE

components). Level 3 feedback (F_3, denominator of IE) sums three-species loops. These loops

from *L-R* are: (i) "N_1 *is eaten*", the I-N_1 loop times E self-limitation (-), (ii) "*starving the enemy*"

(+), (iii) "*feeding the enemy*" (-), and (iv) "N_1 *eats*", the E- N_1 loop times I self-limitation (- ; PIEc

only). This sum of (i)-(iv) determines the sign of F_3 and of IE (since IE $= -F_2/F_3$, and $F_2 < 0$

always). The predator (P, green), prey (N_1, blue), and resources (R, yellow) in IGP are analogous

to immune cells (I), parasite (N_1), and host energy (E), respectively, in the PIE models.

Figure 3: *Weakening of the product of intraspecific direct and indirect effects generates*

oscillations in both IGP and PIE / PIEc models (Fig. 1, Box 1). (A)-(C) *Bifurcation diagrams* over

gradients of supply point (S) and feeding rate of prey or parasite (f_{N_1}): Oscillations (dark orange

region) found in (A) IGP also arise in (B) PIE models of within-host dynamics (see Appendix

Section 2). These oscillatory regions are enveloped within a region of stable coexistence (light

orange). In PIE models, a_b denotes baseline energy allocated to immune cells: $a_b > 0$ in (C) PIEc.

(D)-(F) *Direct and indirect effects (after Fig. 2)*: Oscillations occur with weakening of the product

of intraspecific direct effects (DE) and intraspecific indirect effects (IE). Weakening of IE

primarily triggers oscillations at low f_{N_1} while strengthening of DE restores stability at higher f_{N_1}

in (D) IGP and (E) PIE. In (F) PIEc, self-limitation comes from energy (DE_E, orange) and immune

cells (DE_I, green). Summed DE eliminates oscillations while lowering parasite burden relative to

PIE (see also Fig. A2). (G)-(I) *Equilibrial densities along f_{N_1}*: resource, R^* or energy, E^* (orange);

prey or parasite, N_1^*, (blue); and predator, P^* or immune cells, I^* (green) in (G) IGP, (H) PIE, and

(I) PIEc models. Shifts in these densities weakens or strengthens DE and IE (see text and Appendix

Section 2 for details; Table 1 for default parameters).

Figure 4: *Positive intraspecific indirect feedback leads to two forms of alternate stable*

states (Alt SS-1 and -2) in the IGP model of prey (N_1), predators (P), and resources (R) not found

in PIE models. (A) A bifurcation diagram over gradients of resource supply point (S) and prey

feeding rate (f_{N_1}) and species dynamics at the interior equilibrium (see Fig. A2 for more details).

(B-D) Sample dynamics. (A,B) Stable coexistence ($R - N_1 - P$, light orange) found in IGP also

arises in PIE predicting successful infection (Fig 3; $E - N_1 - I$). (B-D) IGP predicts two forms of

alternate stable states, where initial densities (inset) determine competitive outcomes between N_1

and P. (A, C): In more typical alternative stable states (Alt SS-1; dark grey), either prey (R-N_1) or

predator (R-P) wins. Analogous within-host states would lead to infection (E-N_1) or not (E-I). (A,

F) Another example (Alt SS – 2; light grey) predicts either coexistence (R-N_1-P) or an only-

predator (R-P) equilibrium (analogous to infection [E-N_1-I] or clearance [E-I]). Neither of these

types of alternative states emerge in PIE models (see text, Appendix Section 2).

Figure 5: *Intraspecific direct and indirect effects in four-dimension models: feedback in*

KP, PIE, and 2PIEc (Fig 1). Evaluated at a feasible interior equilibrium, feedbacks for KP and

2PIE are nested within 2PIEc. In 2PIE and 2PIEc, additional loops arise from interactions between

immune cells and host energy ("add loop in 2PIE & 2PIEc"), and immune self-limitation loop

("add loop in 2PIEc"). Level 1 feedback (F_1) sums intraspecific direct effects (solid curves) from

energy and immune self-limitation (- ; only in 2PIEc). The intraspecific indirect effects (dashed

curves) involve a ratio of longer loops, where each species, N_1 (IE_{N_1}; blue shading) and N_2

(IE_{N_2}; purple shading), interacts with others in feedback at level 3 (F_3) and 4 (F_4). F_3 sums three

species loops in PIE models: (i) "N_1 *is eaten*", the I-N_1 loop times E self-limitation (- ; all models),

(ii) "*starving the enemy*" (+ ; from PIE models), (iii) "*feeding the enemy*" (-; from PIE models),

and (iv) "N_1 *eats*", the E-N_1 loop times I self-limitation (- ; only from PIEc). F_4 sums, from L-R,

two destabilizing (+) loops and stabilizing (-) ones. N_i benefits as it (i) "*feeds*" or (ii) "*starves*" the

enemy, but is restrained by (iii) and (iv) "N_i *is eaten*, N_j *eats*" loops (the product of I-N_i and N_j-E

loops). With some rearrangement, F_4 becomes proportional to differences in ratios of how each

competitor has *effects on* ($\varepsilon_1 - \varepsilon_2$) and is *affected by* ($\alpha_1 - \alpha_2$) immune cells and host energy (see

text). Finally, the sign of F_4 determines that of IE (see text). The predator (P, green), prey 1 (N_1,

blue), prey 2 (N_2, purple), and resources (R, yellow) in KP are analogous to immune cells (I),

parasites (N_1 and N_2), and host energy (E) in 2PIE models, respectively.

Figure 6: *KP anticipates coexistence and alternate stable states in 2PIE models.* (A)-(C).

Bifurcation diagrams over gradients of resource supply point (S) and feeding rate of prey or

parasite (f_{N_1}): alternative stable states or priority effects (grey region) and stable coexistence

(orange) found in (A) KP also arise in (B) 2PIE and (C) 2PIE constitutive immunity (2PIEc)

models of within-host dynamics. In 2PIE, a_b denotes baseline energy allocated to immune cells.

Stable coexistence (coinfection) happens at low f_{N_1} and S in all models. (D)-(F) *Indirect effects*

through coexistence/coinfection regions: Strong, negative indirect effects (IE; black, short dash)

from N_1 (IE_{N_1}; blue, long dash) and N_2 (IE_{N_2}; purple, long dash) shift systems from coexistence to

exclusion of N_1 or N_2 at lower or higher f_{N_1} respectively in (D) KP, (E) 2PIE, and (F) 2PIEc. (F)

Immune self-limitation in 2PIEc enhances negative indirect effects and squeezes parameter space

enabling coinfection. (G)-(I) *Equilibrial densities along f_{N_1}*: Resource, R or energy, E (orange);

prey or parasite 1, N_1, (blue); prey or parasite 2, N_2, (purple); and predator, P or immune cells, I

(green) in (G) KP, (H) 2PIE, and (I) 2PIEc models. (See text and Appendix Section 3 for details,

Table 2 for default parameters).

Box 1. *A feedback loop approach to analyze a general 3-dimensional system of equations*

Imagine three species N_1, N_2 and N_3 where the effects of their interactions are captured by a 3D Jacobian matrix (J_3):

$$J_3 = \begin{bmatrix} J_{11} & J_{12} & J_{13} \\ J_{21} & J_{22} & J_{23} \\ J_{31} & J_{32} & J_{33} \end{bmatrix}$$

Characteristic polynomial of J_3
$$\lambda^3 + \lambda^2 F_1 + \lambda F_2 + F_3$$
F_i is level of feedback

Stability requires that ..
$F_1 < 0$
$F_2 < 0$ .. each level of feedback is negative*
$F_3 < 0$

and $F_1 F_2 + F_3 > 0$

We rewrite this criterion as $F_1 \times \left(-\frac{F_2}{F_3}\right) > 1$

to highlight intraspecific direct and indirect effects

Stability as a product of intraspecific direct effects (DE) and indirect effects (IE) (Appendix)

F_1: sum of intra-specific effects

Direct effects   Indirect effects
$$F_1 \quad \times \quad \left(-\frac{F_2}{F_3}\right)$$

$IE = IE_{N_1} + IE_{N_2} + IE_{N_3}$

F_2: sum of two-species loops

F_3: sum of three-species loops

Stability of an equilibrium depends on the sign and strength of the product of DE and IE

Alternate stable states: Indirect effects are positive (DE < 0, IE > 0)

Stable coexistence: Direct and indirect effects are negative (DE < 0, IE < 0) and jointly strong (DE x IE > 1)

Saddle (Alternate stable states)
Direct effects
DE x IE = 1
Stable coexistence   **Oscillations**
Indirect effects

effect on growth rate

Oscillations: Weakening of DE x IE
1. DE < 0 and IE < 0, DE x IE < 1
2. DE > 0 and IE < 0.

*Note: The signs required for stability using Routh Hurwitz criteria (more traditional) are opposite those here for feedback. Thus, F_1 is Tr(J_3) and F_3 is $-|J_3|$

Figure 1

Figure 2

Intraspecific Direct Effects

Level 1 Feedback (F_1)

Sum of self-limitation of each species

IGP, PIE & PIEc   *Add loop in PIEc*

E + I

Intraspecific Indirect Effects

$- \dfrac{\text{Level 2 Feedback } (F_2)}{\text{Level 3 Feedback } (F_3)}$

IE_E IE_{N_1} IE_I

E + N₁ + I

Level 2 Feedback (F_2)
Sum of two-species loops

IGP, PIE, & PIEc models   *Add loop in PIEc*

I   I   N₁   +   +   −   E   I

N₁   E   E

Level 3 Feedback (F_3)
Sum of three-species loops

IGP, PIE, & PIEc models   *Add loop in PIEc*

i   ii   iii   iv

−   +   +   −

I   I   I   I

N₁   N₁   N₁   N₁

E   E   E   E

⊖   ⊕   ⊖   ⊖

P Predator   R Resource   N₁ Prey

I Immune Cells   E Host Energy   Parasite

→ + effect
—● - effect
⌣● - self-limitation

Figure 3

Figure 4

Figure 5

Intraspecific Direct Effects

Level 1 Feedback (F_1)
Sum of self-limitation of each species

In KP, 2PIE & 2PIEc   Add loop in 2PIEc

Intraspecific Indirect Effects

$- \dfrac{\text{Level 3 Feedback }(F_3)}{\text{Level 4 Feedback }(F_4)}$

IE_{N_1} IE_{N_2}

Level 3 Feedback (F_3)
Sum of three-species loop

In KP, 2PIE & 2PIEc   Add loop in 2PIE & 2PIEc   Add loop in 2PIEc

Level 4 Feedback (F_4)
Sum of four-species loops

$\oplus$ $\oplus$ $\ominus$ $\ominus$

F_4 when rearranged is proportional to..

$\dfrac{J_{IN_1}}{J_{EN_1}} - \dfrac{J_{IN_2}}{J_{EN_2}}$ x $\dfrac{J_{N_1I}}{J_{N_1E}} - \dfrac{J_{N_2I}}{J_{N_2E}}$

Effects on ($-\epsilon_i$) E & I   Affected by ($-\alpha_i$) E & I

P Predator   **R** Resource N_i **Prey$_i$**   ⟶ + effect
I Immune Cells   **E** Host Energy   **Parasite$_i$**   — - effect   ⌣ - self-limitation

Figure 6