Drought exerts a greater influence than growth temperature on the temperature response of leaf

day respiration in wheat (*Triticum aestivum*)

Liang Fang[1], Xinyou Yin[1*], Peter E. L. van der Putten[1], Pierre Martre[2], Paul C. Struik[1]

[1] Centre for Crop Systems Analysis, Department of Plant Sciences, Wageningen University &

Research, PO Box 430, 6700 AK Wageningen, The Netherlands

[2] LEPSE, Univ Montpellier, INRAE, Institut Agro SupAgro, Montpellier, France

*Author for correspondence

Dr. Xinyou Yin, Centre for Crop Systems Analysis, Department of Plant Sciences, Wageningen

University & Research, PO Box 430, 6700 AK Wageningen, The Netherlands

Email: xinyou.yin@wur.nl

ORCID IDs

Liang Fang: https://orcid.org/0000-0002-0989-0092

Xinyou Yin: https://orcid.org/0000-0001-8273-8022

Pierre Martre: https://orcid.org/0000-0002-7419-6558

Paul C. Struik: https://orcid.org/0000-0003-2196-547X

Highlight

Leaf day respiration (R_d) acclimated little to growth temperature, but significantly to drought by
reducing its thermal sensitivity. The Kok method underestimates R_d and should no longer be used.

Abstract

We assessed how the temperature response of leaf day respiration (R_d) in wheat responded to
contrasting water regimes and growth temperatures. In Experiment 1, well-watered and drought-
stressed conditions were imposed on two genotypes; in Experiment 2, the two water regimes combined
with high (HT), medium (MT) and low (LT) growth temperatures were imposed on one of the
genotypes. R_d was estimated from simultaneous gas exchange and chlorophyll fluorescence
measurements at six leaf temperatures (T_{leaf}) for each treatment, using the Yin method for non-
photorespiratory conditions and the non-rectangular hyperbolic fitting method for photorespiratory
conditions. The two genotypes responded similarly to growth and measurement conditions. Estimates
of R_d for non-photorespiratory conditions were generally higher than those for photorespiratory
conditions but their responses to T_{leaf} were similar. Under well-watered conditions, R_d and its
sensitivity to T_{leaf} slightly acclimated to LT but did not acclimate to HT. Temperature sensitivities of
R_d were considerably suppressed by drought, and the suppression varied among growth temperatures.
Thus, it is necessary to quantify interactions between drought and growth temperature for reliably
modelling R_d under climate change. Our study also demonstrated that the Kok method, a currently
popular method for estimating R_d, underestimated R_d significantly and should be abandoned.

Keywords: acclimation, climate change, drought, leaf day respiration, photorespiration, temperature,
winter wheat.

Introduction

The ongoing global climate change has resulted in frequent and intense extreme climatic events, such as heat waves, cold snaps and drought spells (Solomon *et al.*, 2009; Lloret *et al.*, 2012; Perkins-Kirkpatrick & Lewis, 2020; IPCC, 2021). Understanding how these climatic events affect crop physiological processes, particularly photosynthesis and respiration, will be critical for global food security and modelling crop productivity as well as for carbon budgets of agricultural ecosystems in response to climate change (Lobell & Gourdji, 2012; Heskel *et al.*, 2013; Yin & Struik, 2017). Respiration plays an essential role in maintaining primary metabolic and physiological functions of plants and costs ca. 40% of gross photosynthetic assimilates of whole plants (Gifford, 1995; Amthor, 2010). Therefore, it strongly affects not only the daily net carbon gain, nutrient acquisition and growth of individual plants, but also the carbon fluxes at ecosystem level (Tcherkez *et al.*, 2017a; Tcherkez & Atkin, 2021), in an ever-changing environment.

Respiration occurring in leaves, the metabolically most active plant organs, accounts for a large part of the whole plant respiration (Atkin *et al.*, 2007). Leaf respiration is sensitive to short-term (minutes to hours) fluctuations in leaf temperature (T_{leaf}) and also acclimates to long-term (days) growth temperature changes (Atkin & Tjoelker, 2003). The response of respiration to short-term changes in temperature is often quantified by the parameter called "activation energy" (E_a) of the Arrhenius model or by the Q_{10} factor (Atkin & Tjoelker, 2003). The thermal acclimation of leaf respiration has been widely investigated (e.g. Atkin *et al.*, 2006; Way *et al.*, 2019; Coast *et al.*, 2020). The degree to which leaf respiration acclimates to growth temperature differs among species and development stages of leaves (Atkin & Tjoelker, 2003; Atkin *et al.*, 2005). Often, leaf respiration acclimates to a sustained warmer growth temperature by decreasing its rate at a reference temperature and/or its thermal sensitivity (i.e. E_a or Q_{10}, the slope of the response curve), while acclimation to cooler temperature increases the values of these parameters (Atkin & Tjoelker, 2003). However, this acclimation is not always observed. A recent study that was conducted on the fast-growing species *Eucalyptus globulus* showed an upregulation in basal rates measured at 25°C and E_a of leaf respiration in warm-grown plants, although the underlying mechanisms are speculative (Crous *et al.*, 2017).

Leaf respiration is also modulated by soil water availability, and drought-induced changes in respiration could be associated with changes in availability of substrates (e.g. soluble sugars and carbohydrates), demand for respiratory products (e.g. ATP and NADH) and capacity of respiratory enzymes (Atkin & Macherel, 2009). The impact of drought stress on leaf respiration varies with species, drought severity and drought duration (Flexas *et al.*, 2005). In ca. two thirds of the studies reviewed

by Atkin & Macherel (2009), leaf respiration was reduced by drought, while in the remaining studies

it was unaffected or occasionally increased. Particularly, resilience of leaf respiration in drought was

observed under cool and moderate measurement temperatures (Gimeno *et al.*, 2010; Gauthier *et al.*,

2014). Furthermore, the effect of drought on leaf respiration can also interact with the effects of other

environmental factors, for example those of short- or long-term temperature changes and elevated

atmospheric CO_2, especially under field conditions (Ayub *et al.*, 2011; Crous *et al.*, 2011, 2012;

Gauthier *et al.*, 2014). These interactions will lead to more unpredictable responses of leaf respiration

to drought.

There is growing evidence that the metabolic pathways of leaf respiration vary between illuminated

and non-illuminated leaves, as a result of the inhibition of respiration in the light (Tcherkez *et al.*,

2017a,b; Tcherkez & Atkin, 2021). Unlike leaf respiration in the dark (R_{dk}), leaf day respiration (R_d)

occurs simultaneously with photosynthetic CO_2 assimilation and other physiological processes, such

as photorespiration, reassimilation and photoinhibition, in the daytime (Yin *et al.*, 2020). R_d is an

important parameter in modelling net photosynthetic CO_2 assimilation (Farquhar *et al.*, 1980) and can

influence the estimation of other key photosynthetic parameters, such as V_{cmax}, the maximum rate of

Rubisco carboxylation (De Kauwe *et al.*, 2016). As the FvCB model (Farquhar *et al.*, 1980) is widely

used as the basic model for predicting leaf photosynthesis that is to be scaled up to the ecosystem level,

R_d is also crucial in modelling ecosystem gross CO_2 efflux. However, the different metabolic pathways

of R_d and R_{dk} may result in different responses to environmental variables (Gulías *et al.*, 2002; Way *et*

al., 2019). Compared with the abundance of studies that have explored the environmental impacts on

R_{dk}, the experimental data on how R_d responds to various environments is generally more lacking,

probably because it is difficult to measure R_d.

Techniques to quantify R_d have been implemented for decades, while there is still debate about the

best technique to quantify R_d (Tcherkez *et al.*, 2017a,b; Tcherkez & Atkin, 2021). Either indirect or

direct techniques have been developed (Kok, 1948; Laisk, 1977; Haupt-Herting *et al.*, 2001; Yin *et al.*,

2009; Gong *et al.*, 2015; Berghuijs *et al.*, 2019). Direct measurement of R_d (e.g. Haupt-Herting *et al.*,

2001; Gong *et al.*, 2015) requires sophisticated devices, which are often unavailable. Indirect

estimation of R_d by gas exchange measurements in ecophysiological studies is mostly based on either

the Kok method (Kok, 1948) or the Laisk method (Laisk, 1977). The Kok method exploits the Kok

effect, that is the abrupt decrease in the slope of photosynthetic CO_2 assimilation rate (A) against

irradiance at around the light compensation point (0 - 40 $\mu mol\ m^{-2}\ s^{-1}$). This abrupt switch is interpreted

as the consequence of light inhibition to leaf respiration, and thus R_d can be calculated as the intercept

of this linear relationship using points above the break point while the intercept of the linear

relationship below the break point is interpreted as R_{dk} (Heskel *et al.*, 2013; Tcherkez *et al.*, 2017b;

Yin *et al.*, 2020). The Laisk method is based on the FvCB model (Farquhar *et al.*, 1980), where the

response of A to low intercellular CO_2 concentration (C_i) can be obtained at several (commonly three)

levels of irradiances. These curves theoretically intersect at a common point, where the value of A

represents R_d and the value of C_i represents the CO_2 compensation point in the absence of respiration.

The R_d estimate by the Kok method is often somewhat lower than the estimate by the Laisk method

(Villar *et al.*, 1994; Yin *et al.*, 2011), probably because the Kok method assumes that the PSII electron

transport efficiency (Φ_2) is constant across the light levels used for the gas exchange measurements.

A modified Kok method, now known as the Yin method (see Tcherkez *et al.*, 2017a) was developed

to overcome this weakness of the Kok method, by incorporating the information from chlorophyll

fluorescence that accounts for the decline of Φ_2 with light intensity (Yin *et al.*, 2009, 2011). The Yin

method gave estimates of R_d that were comparable with those from the Laisk method (Yin *et al.*, 2011),

with a benefit that the measurements are easier and less time-consuming to implement than those with

the Laisk method provided that a fluorometer is available for the concurrent gas exchange and

chlorophyll fluorescence measurements.

Although the Kok method, and sometimes also the Yin method, have been applied to common

photorespiratory (PR) conditions, theoretically both methods require measurements that are

undertaken under nonphotorespiratory (NPR) conditions. This is because both methods using simple

linear regression to estimate R_d implicitly assume that the chloroplast CO_2 partial pressure (C_c) is

constant across light levels whereas a modelling study demonstrated that under PR conditions, C_c

sharply decreased (thus the relative amount of photorespiration increased) with increasing irradiance

(Farquhar & Busch, 2017). The significant change of C_c with irradiance (even if the ambient CO_2 level

is maintained constant) is the result of stomatal and mesophyll regulation of CO_2 diffusion inside the

leaf, ensuring some reassimilation of CO_2 released by photorespiration and respiration. Using leaf

anatomical data combined with 2-D modelling that accounts for CO_2 diffusion, and thus reassimilation,

Berghuijs *et al.* (2019) were able to estimate R_d based on gas exchange and chlorophyll fluorescence

data for PR conditions; they showed that applying the Kok and the Yin methods to PR conditions cause

an underestimation of R_d. If the 2-D modelling of the CO_2 diffusion can reliably estimate R_d, a simpler

method, i.e. the coupled FvCB and g_m (mesophyll conductance) models, can also be explored to

estimate R_d by fitting the coupled model to gas exchange and chlorophyll fluorescence data obtained

under PR conditions, because the coupled model can implicitly consider the reassimilation of

(photo)respired CO_2 (Yin *et al.*, 2021). The coupled model has a non-rectangular hyperbolic (NRH)

form (von Caemmerer, 2000), and this NRH model was exploited by Yin & Struik (2009) to estimate

g_m from combined gas exchange and chlorophyll fluorescence data, which was shown to be more

reliable than the well-known variable J method for estimating g_m. Here, we will test if this model can

well estimate R_d under PR conditions, even if g_m is unknown beforehand.

Wheat (*Triticum aestivum* L.), as one of the most important staple food crops, is grown worldwide.

There is little practical data on how R_d in C_3 Poaceae species like wheat varies with environmental

variables, in contrast to the case for tree species, for which studies under either climate-controlled or

field conditions have been extensively published in the recent past (e.g. Crous *et al.*, 2011, 2012; Way

et al., 2019; Kumarathunge *et al.*, 2020). To date, to our knowledge, no single study provides a set of

data of R_d and its sensitivity to T_{leaf} in response to a combination of contrasting water regimes and

different growth temperatures in wheat. Understanding how R_d responds to environmental variables

(especially to drought) is critical for quantifying wheat productivity in response to current and future

climate change scenarios.

The main objective of this study is to identify the impacts of soil water deficit and growth temperatures

on the instantaneous temperature response of R_d in wheat. We hypothesize that R_d and its sensitivity

to T_{leaf} would acclimate to contrasting water regimes and to different growth temperatures, and the

thermal acclimation of R_d may differ between two contrasting water treatments. We estimated R_d using

the established Yin method that requires measurements under NPR conditions. We also aimed to

estimate R_d for common PR conditions; so, the NRH method was explored to estimate R_d as well. We

then evaluated if acclimation differed between PR and NPR conditions. Moreover, we also compared

the R_d estimated by the Kok, Yin and NRH methods.

Materials and Methods

Plant materials and growth conditions

To determine any interactive effect of water deficit and genotype on R_d of wheat plants, an experiment

was conducted in a climate-controlled glasshouse at Wageningen University in 2019 (EXP2019). Two

genotypes of winter wheat (*Triticum aestivum* L., Thésée and Récital) were used in this experiment.

Four batches of seeds were sown with 10-day intervals, creating four replicates. In each replicate, seeds

were germinated on a moist filter paper in Petri dishes (one night at room temperature followed by 24

h at 4°C), and then the seedlings were transplanted to multi-cell seedling trays in a glasshouse. When

the first leaf fully emerged (ca. 1 week after sowing), plants were moved to a 4°C cold-room (12 h day

length, 50 µmol m^{-2} s^{-1} photon flux density) to vernalize for 7 weeks. After vernalization, plants were

transplanted to 7-L pots (three plants per pot) filled with a mixture of black soil and peat in a 2:1 (v:v)

ratio. The soil mixture was 6 kg per pot and mixed with 1 g N, 1 g P and 1 g K. All potted plants were

then moved to a glasshouse and the position of pots were rotated daily and randomly. The climate

condition in the glasshouse compartment was set as: 400 ± 5 ppm atmospheric CO_2 concentration,

$22/16 \pm 2°C$ day/night air temperature (20°C daily average), $75 \pm 5\%$ relative humidity (corresponding

to a vapour pressure deficit (VPD) of 0.66/0.46 kPa for day/night), 16 h photoperiod, and photon flux

density at > 400 µmol m^{-2} s^{-1} supplied by sunlight plus supplementary sodium lamps. To avoid any

nutrient deficiency, 0.5 g N and 0.25 g N were applied to each pot at the tillering and the stem-

elongating stages, respectively.

Another experiment was conducted in a climate-controlled growth chamber in 2020 (EXP2020) to

examine whether the leaf day respiration can be altered by different water regimes and growth

temperatures. EXP2019 showed no significant difference between the two genotypes in R_d and its

temperature response (see Results); so, only one genotype, Thésée, was used in EXP2020. Considering

varying crop durations in different growth temperature regimes, four batches of seeds were sown with

14-day, 17-day and 20-day intervals, respectively, creating four replicates. The same vernalization

treatment and plant management practices as in EXP2019 were applied. After vernalization, the potted

plants were moved to a climate chamber. The climate settings in the climate chamber were set as:

atmospheric CO_2 concentration, 400 ppm; day/night temperature, 21/17°C; relative humidity, 65%;

VPD, 0.87/0.68 kPa for day/night; photon flux density, ca. 410 µmol m^{-2} s^{-1} at soil level; photoperiod,

16 h.

Experimental treatments

In EXP2019, drought treatment was applied at anthesis in both genotypes. From sowing to anthesis all

pots were constantly irrigated to 90% soil water holding capacity, with a gravimetric soil water content

of ca. 42%. After anthesis, four replicates of each genotype were well irrigated as control plants (well-

watered treatment, WW), whereas the other four replicates of each genotype were subjected to drought

stress (treatment DS) by withholding irrigation until the gravimetric soil water content reduced to ca.

16% (drought-stressed treatment, DS). This drought level was maintained until the end of

measurements (ca. one week).

In EXP2020, combinations of water deficit treatments and growth temperature were applied. At the

booting stage before flag leaves appeared, plants were allocated to three climate chambers with

different day/night air temperature settings: high temperature (HT: 28/24°C, average 26.67°C),

medium temperature (MT: 21/17°C, average 19.67°C) and low temperature (LT: 16/12°C, average

14.67°C). MT is considered as the control treatment since the temperature was the same as the growth

temperature before the temperature treatments started. To minimize any confounding impact of

varying VPD, VPD was set identically across chambers (0.87/0.68 kPa for day/night VPD); as a result,

77%, 65%, and 52% relative humidity were applied for HT, MT, and LT treatments, respectively.

After anthesis, plants from each climate chamber were subjected to the same two soil water treatments

as in EXP2019.

Gas exchange, chlorophyll fluorescence and leaf N measurements

In both experiments, simultaneous gas exchange and chlorophyll fluorescence measurements were

carried out on flag leaves at six leaf temperatures (T_{leaf}; from 15°C to 40°C with 5°C intervals, except

for the LT plants in EXP2020, which were measured at T_{leaf} from 12°C to 35°C), 10 days after the

onset of the drought treatment, by using a portable photosynthetic system (Li-Cor 6800; Li-Cor Inc.,

Lincoln, NE, USA) with an integrated fluorescence chamber head of 6 cm^2. Li-Cor 6800 and plants

were together moved to a climate cabinet during measurement to achieve the desired T_{leaf}. The VPD

in the cuvette increased with an increase in T_{leaf} and ranged from 1.0 kPa (at 15°C) to 3.0 kPa (at 40°C)

for all plants in EXP2019 as well as HT and MT plants in EXP2020. For LT plants in EXP2020, VPD

ranged from 1.0 (at 12°C) kPa to 2.5 kPa (at 35°C). For a given T_{leaf}, incident-irradiance response

curves (A-I_{inc}) were assessed under both photorespiratory (PR, i.e. 21% O_2 combined with 400 ppm

ambient CO_2 (C_a)) and non-photorespiratory (NPR, i.e. 2% O_2 combined with 1000 ppm C_a) conditions

on the same leaf. For the measurements at NPR conditions, a gas cylinder containing a mixture of 2%

O_2 and 98% N_2 was used. Gas from the cylinder was supplied to the Li-Cor 6800 where CO_2 was

blended with the gas. Photon flux densities in the measurement chamber were 200, 150, 120, 90, 60,

40 and 0 µmol m^{-2} s^{-1} (applied in that order; the value of A at 0 µmol m^{-2} s^{-1} of I_{inc} represents R_{dk}) with

5-6 min for each step. The measurements were conducted randomly in each treatment and T_{leaf}. The

operating Φ_2 was determined at each light step as ($1-F_s/F'_m$) (Genty *et al.*, 1989), where F_s is the

steady-state fluorescence, and F'_m is the maximum fluorescence during the saturating light pulse

determined by the multi-phase flash method (Loriaux *et al.*, 2013).

After the measurements, the portion of the flag leaves used for gas exchange and chlorophyll

fluorescence measurements was cut to measure leaf N elemental content. Rectangle area of the leaf

portion was calculated as length multiplied by width which were measured by a vernier caliper. Then

the leaf material was weighed after drying in a forced-air oven at 70°C to a constant weight. The

concentration of total N in leaf material was analyzed using an EA1108 CHN-O Element Analyzer

(Fisons Instruments) based on the micro-Dumas combustion method. From these data, leaf N content

on area basis (N_{area}, g m^{-2}) and mass basis (N_{mass}, mg g^{-1}) were calculated.

Estimation of leaf day respiration

R_d was estimated by the Yin method ($R_{d(Yin)}$) (Yin *et al.*, 2009; 2011), where $R_{d(Yin)}$ is estimated when

A is limited by the light-dependent electron transport rate, at which A can be described as (Yin *et al.*,

2004):

$$A = J_2 \left(1 - \frac{f_{pseudo}}{1-f_{cyc}}\right) \left(\frac{C_c - \Gamma_*}{C_c + 2\Gamma_*}\right)/4 - R_d \qquad (1)$$

where J_2 is the total rate of e$^-$ transport passing PSII, f_{cyc} and f_{pseudo} represent fractions of the total e$^-$

passing PSI that follow cyclic and pseudocyclic pathways respectively, C_c is the chloroplast CO_2

partial pressure, and Γ_* is the C_c-based CO_2 compensation point in the absence of R_d. By definition, J_2

can be replaced by $\rho_2 \beta I_{inc} \Phi_2$, where ρ_2 is the proportion of absorbed irradiance partitioned to PSII, β is

the absorptance by leaf photosynthetic pigments, I_{inc} is the incident irradiance, and Φ_2 is the quantum

efficiency of PSII electron transport. Then Eqn (1) becomes:

$$A = \rho_2 \beta I_{inc} \Phi_2 \left(1 - \frac{f_{pseudo}}{1-f_{cyc}}\right) \left(\frac{C_c - \Gamma_*}{C_c + 2\Gamma_*}\right)/4 - R_d \qquad (2)$$

For NPR conditions, C_c is assumed infinite and/or Γ_* approaches zero, then Eqn (2) becomes:

$$A = s(I_{inc} \Phi_2/4) - R_d \qquad (3)$$

where the lumped parameter calibration factor $s = \rho_2 \beta [1 - f_{pseudo}/(1 - f_{cyc})]$. So, using data of the

electron-transport-limited range (200 to 40 µmol m^{-2} s^{-1}) under NPR conditions, linear regression plots

of A against ($I_{inc}\Phi_2/4$) can be produced, in which Φ_2 is based on chlorophyll fluorescence

measurements. The slope of the regression gives the estimate of a calibration factor s, and the intercept

yields the estimate of $R_{d(Yin)}$ under NPR conditions (Yin *et al.*, 2009). Obviously, this approach requires

that all points are within the linear range of the A vs ($I_{inc}\Phi_2/4$) curves. Here, curves of A against ($I_{inc}\Phi_2/4$)

were inspected to exclude the points at high ends that might deviate from the linear pattern, especially

for drought plants.

Eqn (2) could also be explored to estimate R_d under PR conditions if C_c is maintained constant across

irradiance levels. However, it is practically difficult to control C_c because, before measurements are

actually undertaken, one does not know actual values of photosynthetic rate, stomatal conductance and

mesophyll conductance required for calculating C_c. Here, the aforementioned non-rectangular

hyperbolic (NRH) equation was used to estimate R_d ($R_{d(NRH)}$) for PR conditions. This equation is

obtained by combining the well-known FvCB model (Farquhar *et al.*, 1980) for e$^-$ transport-limited A

with the Fick's first law of diffusion for the relation between A, intercellular CO_2 partial pressure (C_i)

and C_c:

$$A = 0.5\{J/4 - R_d + g_m(C_i + 2\Gamma_*) -$$

$$\sqrt{[J/4 - R_d + g_m(C_i + 2\Gamma_*)]^2 - 4g_m[(C_i - \Gamma_*)J/4 - R_d(C_i + 2\Gamma_*)]}\} \qquad (4)$$

where g_m is the mesophyll conductance, and J is the linear e$^-$ transport rate through PSII, which can be

calculated as: $J = sI_{inc}\Phi_2$ (Yin et $al.$, 2009). The calibration factor s was adopted from the slope value

of the linear regression for the Yin method from the data under NPR conditions (see above). Γ_* can be

calculated from the relation $\Gamma_* = 0.5O/S_{c/o}$ (where O is the concentration of oxygen and $S_{c/o}$ is the

relative CO_2/O_2 specificity factor for Rubisco; Farquhar et $al.$, 1980; von Caemmerer, 2013). Nonlinear

curve fitting based on Eqn (4) was used to estimate g_m in a previous study (Yin & Struik, 2009). Here,

this NRH equation for A was fitted to simultaneously estimate $R_{d(NRH)}$ and g_m under PR conditions.

The data used in the NRH method was within the same range of light levels as used in the Yin method.

Use of Eqn (4) combined with $J = sI_{inc}\Phi_2$ to estimate R_d and g_m requires that the calibration factor s is

obtained from strictly NPR conditions. There is no guarantee that this could be the case at extreme

conditions (especially when high temperature is combined with drought stress), as the stomatal

conductance and g_m are so low under such conditions that Γ_*/C_c could not be maintained at the required

low level to achieve NPR conditions even when C_a was set at 1000 ppm. In other words, if a NPR

condition cannot be ensured, the obtained s is not equal to $\rho_2\beta[1-f_{pseudo}/(1- f_{cyc})]$, but to

$\rho_2\beta[1-f_{pseudo}/(1-f_{cyc})][(C_c-\Gamma_*)/(C_c+2\Gamma_*)]$; so, s would be lowered by a factor $(C_c-\Gamma_*)/(C_c+2\Gamma_*)$. We

introduced dummy variables to Eqn (4) to avoid the confounding effect of measuring under conditions

that were not truly NPR and examined whether or not the calibration factor s was underestimated. The

following identity was verified:

$$R_d = Z_1 R_{d,NPR} + Z_2 R_{d,PR} \qquad (5)$$

where $R_{d,NPR}$ and $R_{d,PR}$ are leaf day respiration under NPR and PR conditions, respectively, and Z_1 and

Z_2 are dummy variables, which were set in such a way that $Z_1 = 1$ and $Z_2 = 0$ correspond to the NPR

condition and $Z_1 = 0$ and $Z_2 = 1$ correspond to the PR condition. Such a procedure allows to

simultaneously estimate the common parameters (s and g_m) as well as the different parameters ($R_{d,NPR}$

and $R_{d,PR}$) from fitting to the combined data obtained under both NPR and PR conditions. This

procedure for estimating the common parameters s and g_m is equivalent to the method of simultaneous

fitting of the calibration factor and g_m (e.g. Pons et $al.$, 2009) if one is not sure whether a NPR state is

reached.

As stated earlier, like the Laisk method, the Kok method is a popular method to estimate R_d indirectly

(e.g. still used recently by Way *et al.*, 2019). As our data also allow an implementation of the Kok

method, here we compare the NRH, Yin and Kok methods in estimating R_d. As the Kok method has

been applied to estimate R_d ($R_{d(Kok)}$) under both PR and NPR conditions (e.g. Tcherkez *et al.*, 2017a),

for the comparative purpose we also apply the Yin method to the PR conditions by fitting a linear

regression to data at the same range of irradiances. It is worthy to note that Eqn (4), upon which the

NRH method is based to estimate R_d, can only be applied to the PR conditions.

The temperature responses of parameters

The thermal response of R_{dk} and R_d was described by the Arrhenius equation normalized with respect

to its value at 25°C:

303
$$X = X_{25} e^{\left(\frac{1}{298} - \frac{1}{273+T_{\text{leaf}}}\right)\frac{E_X}{R}} \tag{6}$$

where X_{25} represents the value of parameters estimated at 25°C (R_{d25} and $S_{c/o25}$), E_X is the activation

energy of relevant parameters to temperature (E_{Rd} and $E_{Sc/o}$; in kJ mol^{-1}), and R is the universal gas

constant (0.008314 kJ K^{-1} mol^{-1}). $S_{c/o}$ and its temperature response are generally considered to be

conserved among C$_3$ species (von Caemmerer *et al.*, 2009), so we adopted the value of Cousins *et al.*

(2010) for $S_{c/o25}$ (3.022 mbar μbar^{-1}), and the value of Bernacchi *et al.* (2002) for $E_{Sc/o}$, which

mathematically equals to the negative of the activation energy for Γ_*: 24.46 kJ mol^{-1}. Then the value

of $S_{c/o}$ at each T_{leaf} could be estimated. Sensitivity analysis showed that unlike g_m, the estimated R_d and

its temperature response varied little with variations of $S_{c/o25}$ and $E_{Sc/o}$ within the physiologically

relevant ranges.

Model analyses and statistics

Simple linear regressions in the Yin method were performed using Microsoft Excel. Non-linear curve

fitting procedures in the NRH method and the Arrhenius equation were carried out using the GAUSS

method in PROC NLIN of SAS (SAS Institute Inc, Cary, NC, USA). The SAS codes can be obtained

upon request to the corresponding author.

Results

The approach combining the NRH method and dummy variables was used to examine if there was a

confounding effect of any under-estimation of the calibration factor s on $R_{d(NRH)}$ for PR conditions.

Using methods either with or without dummy variables, we found that overfitting occurred for a few

T_{leaf} values of drought plants due to the variability of data, which resulted in the failure of estimating

g_m. For such cases, nevertheless, the estimations of s and $R_{d(NRH)}$ were still reasonable. The results of

each parameter (calibration factor s, g_m, $R_{d(NRH)}$ for PR and $R_{d(Yin)}$ for NPR conditions) from approaches

with and without dummy variables were very similar (Fig. S1). On average, s was underestimated only

by ca. 1% (Fig. S1b), suggesting that the gas mixture we used (2% O_2 combined with 1000 ppm C_a)

for estimating the calibration factor did allow to reach a nearly non-photorespiratory state, even for

extreme conditions (drought combined with high temperatures). So, for the sake of simplicity, we only

present and discuss the results obtained from the method without using the dummy variables.

Comparisons of R_d and its inhibition by light estimated by different methods

$R_{d(Kok)}$ was generally lower than $R_{d(Yin)}$ under both NPR (18.2% lower) and PR (13.9% lower)

conditions (Fig. 1A,B). Under PR conditions, values of $R_{d(Kok)}$ were only 66.5% of $R_{d(NRH)}$, and those

of $R_{d(Yin)}$ were 79.6% of $R_{d(NRH)}$ (Fig. 1C,D). These findings suggest that the Kok method

underestimated R_d substantially, and the Yin method, if also applied to PR conditions, also

underestimated R_d, but less so than the Kok method. Therefore, the light inhibition of leaf respiration,

estimated as $(1-R_d/R_{dk})$, depended on the methods used for estimating R_d. Under NPR conditions, the

estimated average light inhibition by the Kok method was 18.7%, whereas that by the Yin method was

only 1.4% (Fig. 2A,B). Under PR conditions, the estimated average light inhibition by the Kok, Yin

and NRH methods was 40.3%, 28.5% and 10.1%, respectively (Fig. 2C,D,E), indicating that light

inhibition was stronger under PR than NPR conditions. In general, values of R_{dk} and R_d under NPR

conditions were greater than under PR conditions (Fig. 3), which was in agreement with results from

previous report where both R_{dk} and R_d were higher at 2% O_2 than at 21% O_2 in mature leaves (Buckley

et al., 2017).

The obtained thermal responses of R_{dk}, $R_{d(Kok)}$ and $R_{d(Yin)}$ under PR and NPR conditions as well as

$R_{d(NRH)}$ under PR conditions were similar (Figs. S2, S3 & S4; Tables S1, S2). Hereafter, we only use

$R_{d(Yin)}$ for NPR conditions and $R_{d(NRH)}$ for PR conditions for further analyses because our study focuses

on R_d and theoretically the Yin method and the NRH method work best to estimate R_d for NPR and

PR conditions, respectively.

Impact of water regimes on R_d and its response to leaf temperature in two wheat genotypes

In EXP2019, R_d was estimated across two genotypes and two water regimes. As expected, for both

NPR and PR conditions, the estimated values of R_d increased with rising T_{leaf} across genotypes and

water regimes (Fig. 4), as was widely found in previous studies (Atkin *et al.*, 2006; Crous *et al.*, 2011;

Yin *et al.*, 2014; Way *et al.*, 2019). The temperature response of R_d did not vary much between Thésée

and Récital but significantly differed between WW and DS conditions. Generally, R_d was suppressed

by drought stress. In agreement with the results of Crous *et al.* (2012), this suppression was slightly or

not significant at lower T_{leaf} (15 and 20°C), but became increasingly pronounced with increasing T_{leaf},

reaching up to 76.3% and 71.5% under NPR conditions and up to 74.2% and 64.7% under PR

conditions in Thésée and Récital, respectively, for $T_{leaf} = 40°C$ (Fig. 4).

The temperature response of R_d was well described by the Arrhenius equation, although the estimated

R_d deviated more under drought conditions as a result of higher variabilities of data among replicated

plants. R_d and its response to T_{leaf} showed appreciable acclimation to drought. Under drought stress,

R_{d25} was apparently reduced, with the reduction ranging from 44.1% to 55.2% across genotypes and

between PR and NPR conditions (Fig. 4, Table 1). The values of E_{Rd} under well-watered conditions

were similar, ranging from 58.87 to 68.64 kJ mol^{-1} in the two genotypes under NPR and PR conditions

(Table 1), which agreed with the estimate (64.18 kJ mol^{-1}) by Yin *et al.* (2014) for tomato. The

estimated E_{Rd} was notably affected by water treatments. Under drought conditions the values of E_{Rd}

decreased to less than 42 kJ mol^{-1} (Table 1), reflecting that R_d was less sensitive to increasing

temperature under water-deficit conditions than in well-watered conditions (Fig. 4).

Impact of the combined water and growth temperature regimes on R_d and its temperature response

In EXP2020, treatments were designed to investigate the interactive effect of growth temperature and

water regime on R_d. The response of R_d to T_{leaf} was described by the Arrhenius equation in each

treatment, although it did not fit well the data for DS plants grown at LT ($r^2 = 0.329$ and 0.395 for

NPR and PR conditions, respectively).

Under WW conditions, R_d estimated at each respective T_{leaf} was rather consistently lower in plants

grown at HT and MT than in plants grown at LT in both NPR and PR conditions (Fig. 5A,B). This

was also reflected in the higher estimation of R_{d25} in LT plants under WW conditions (Table 2).

However, the estimated E_{Rd} for WW plants herein was found to be lowest at LT and highest at HT

(Table 2), although the difference between HT and MT was not significant, especially at PR conditions.

Our results are contradictory to the lower sensitivity of R_d to T_{leaf} at warmer growth temperature

reported previously (Atkin *et al.*, 2005 and references therein).

Again, drought stress reduced R_d across various growth environments and between PR and NPR

conditions and this reduction was more pronounced at higher T_{leaf} (Fig. 5), which led to lower

sensitivity to T_{leaf}, and, thus, lower E_{Rd} in DS plants than in WW plants (Table 1). However, this

drought-induced reduction of R_d differed among various growth regimes. DS plants grown at LT

maintained relatively higher R_d at lower T_{leaf} (below 25°C) than HT- and MT-grown plants, while at

higher T_{leaf} (above 30°C) higher R_d values were observed in DS plants grown at MT as compared with

those grown at HT and LT (Fig. 5C,D). This resulted in a downward shift in the temperature response

curve and a lower estimated R_{d25} (0.78 and 0.51 µmol m^{-2} s^{-1} for NPR and PR, respectively) in DS

plants grown at HT; and a nearly horizontal curve with an extremely low estimate of E_{Rd} (8.10 and

12.93 kJ mol^{-1} for NPR and PR, respectively) in DS plants grown at LT as compared to those grown

at HT and MT (Table 2).

Discussion

The present study investigated the impacts of long-term drought and growth temperature treatments

on the short-term thermal response of R_d in two winter wheat genotypes. Because little difference in

R_d was observed between the two wheat genotypes under the two contrasting water regimes in

EXP2019, only Thésée was used in EXP2020.

Impact of growth temperature on R_d under well-watered conditions

Thermal acclimation of respiration is usually assessed from how changes with growth temperature in

the response of respiration rate to measurement T_{leaf} sustained over time (Crous *et al.*, 2011). In many

previous studies, plants acclimated to a sustained warmer climate by reducing their respiration rate at

a given measurement temperature (e.g. Crous *et al.*, 2011). Moreover, the extent of thermal acclimation

of respiration is often lower in pre-existing, fully-expanded leaves that are shifted to a new growth

temperature than in leaves that develop under various growth temperatures (Atkin & Tjoelker, 2003).

Here, although the plants were subjected to the growth temperature treatment before flag leaves

emerged, our results showed that under WW conditions rates of R_d at any given T_{leaf} were nearly

identical for plants grown at HT and MT (i.e. no thermal acclimation) regardless of estimation methods

or photorespiratory conditions, but R_d slightly acclimated to the LT with an upward shift of the

temperature response curve and a higher R_{d25} (Fig. 5A,B, Table 2). This was consistent with findings

in previous studies across various R_d estimation methods and species, indicating that little or only

partial thermal acclimation of R_d is more common (Atkin *et al.*, 2006; Way *et al.*, 2019).

Plant respiration is sensitive to short-term changes in temperature, while the effect of growth

temperature on the thermal sensitivity of respiration varies among species (Atkin *et al.*, 2005). Previous

works have reported that the sensitivity of respiration to T_{leaf} declines with increasing growth

temperature, as a result of the acclimation of respiration to a warmer temperature (e.g. Cai *et al.*, 2020).

In contrast, we found that the estimates of E_{Rd} were lower in LT plants than in MT and HT plants under

WW conditions (Table 2), implying that plants grown at lower temperatures are likely to be less
sensitive to rising T_{leaf} than those grown at warmer temperatures. Coincidentally, Crous et al. (2017)
also observed a higher value of E_{Rd} in a fast-growing tree species (*Eucalyptus globulus*) grown in a
warmer climate, although the mechanisms behind this remain unclear. The possible explanation of this
upward trend of sensitivity to T_{leaf} in warm climates could be linked to the higher leaf N content on a
mass basis at higher growth temperature (Fig. S5E,F); plant N status is highly associated with factors
such as enzyme capacity, substrate supply and respiratory products that may affect the metabolic
activities in plant tissues (O'Leary et al., 2019 and references therein). Additionally, our data showed
that R_{d25} positively correlated with leaf N content on area basis (Fig. S5A,B).

Impact of drought stress on R_d and its interaction with growth temperature

Accumulated evidence indicates that the response of respiration and its thermal sensitivity to stressed
environments are associated with the alternations in substrate supply, respiratory products demand and
respiration/enzyme capacity (Atkin & Tjoelker, 2003; Flexas et al., 2005; Atkin & Macherel, 2009).
Previous studies suggested that the decreased leaf respiration rate under drought was more likely due
to the reduced demand for respiratory products (Ayub et al., 2011), while the changed thermal
sensitivity was more likely to depend on substrate availability and respiratory enzyme capacity (Atkin
& Tjoelker, 2003). The response of leaf respiration to limited water availability seems to be equivocal
and elusive as decreased (Ayub et al., 2011), unaffected (Gimeno et al., 2010) and even increased
(Gauthier et al., 2014) leaf respiration rate under water deficit have been reported. The various
responses could be linked to difference in species used, the severity and duration of soil dehydration
and/or other environmental factors, e.g. temperature (Flexas et al., 2005).

In our study, inhibition by severe drought on R_d was slight or not significant under lower T_{leaf} across
all treatments (Figs. 4,5), consistent with Gauthier et al. (2014) who reported that the resilience of leaf
respiration in the dark (R_{dk}) was observed in low to moderate ranges of T_{leaf}. This finding might explain
the unaffected and even slightly increased leaf respiration under drought treatment in some cases where
R_d or R_{dk} was measured at a set of common or lower temperature rather than a short-term change of
measurement temperature (Gimeno et al., 2010; Sperlich et al., 2016). At $T_{leaf} > 20°C$, however, our
results showed that the inhibition of R_d by drought was substantial (Figs. 4,5), which was in agreement
with previous studies (Haupt-Herting et al., 2001; Crous et al., 2012; Dahal & Vanlerberghe, 2017).
As a result, there was low sensitivity of R_d to T_{leaf} in drought plants (Figs. 4,5). However, several
previous studies reported contradictory results: drought may exacerbate R_{dk} at moderate or higher
temperature (Zagdańska, 1995; Slot et al., 2008) and even lead to a 'respiratory burst' in an extremely

high (> 40°C) short-term measurement temperature range (Gauthier *et al.*, 2014), although the

underlying mechanisms remain speculative. In some of these studies (e.g. Gauthier *et al.*, 2014), plants

experienced two phases of drought with a rewatering treatment in between, and the temperature

response of leaf respiration was measured at the end of the second period of drought. This means that

the increased leaf respiration under drought stress could be due to drought priming. Moreover, in these

early studies, drought treatment was applied at seedling or sapling stage (Zagdańska, 1995; Bartoli *et*

al., 2005; Gauthier *et al.*, 2014), instead of at post-anthesis stage at which assimilated carbohydrates

and nitrogenous compounds, those related to substrate supply and respiratory capacity (Tjoelker *et al.*,

1999), are being translocated from source (leaves) to sink (grains) (Shao *et al.*, 2021), and energy

demand for sucrose synthesis and/or phloem loading is in decline (Atkin & Macherel, 2009). The

above may explain the difference between their results and our experiments.

Furthermore, our results showed that indeed there was an interactive impact of drought stress and

growth temperature on the response of R_d to T_{leaf}, with a much lower E_{Rd} value in DS plants grown at

LT and a lower R_{d25} in DS plants grown at HT than in WW plants (Table 2). A low E_{Rd} in DS plants

grown at LT means a reduced R_d at higher T_{leaf}. When T_{leaf} was < 25°C, LT treatment appeared to

alleviate the negative impact of drought on R_d rates as compared with HT and MT treatments, whereas

this was not the case in the higher T_{leaf} range (> 25°C) where DS plants grown at MT exhibited the

highest rates of R_d (Fig. 5C,D). The possible explanation for the mitigated drought effect on LT-grown

plants could be linked to the up-regulated alternative oxidase (Searle *et al.*, 2011; Dahal &

Vanlerberghe, 2017) and mitochondrial uncoupling proteins (Nantes *et al.*, 1999; Barreto *et al.*, 2017).

Comparisons of the Kok method, the Yin method and the NRH method

Compared with the Kok method, the Yin method accounts for the additional information from

chlorophyll fluorescence data. Our results showed that $R_{d(Kok)}$ was lower than $R_{d(Yin)}$ under both NPR

and PR conditions (Fig. 1A,B), which was in line with previous studies (e.g. Yin *et al.*, 2011). The

lower estimates of $R_{d(Kok)}$ were due to the neglect of a decrease in Φ_2 with increasing light intensity,

which occurs even with limiting light levels. Theoretically, the Yin method works well under NPR

conditions (Berghuijs *et al.*, 2019). It works for PR conditions only if C_c is maintained constant across

light levels, which is technically difficult to achieve in measurements because g_m is unknown

beforehand. Berghuijs *et al.* (2019) pointed out that the linear regression as used in the Yin method

will underestimate R_d if applied to PR conditions; this is also confirmed by our data (Fig. 1D). The

theoretical underpinning is that under PR conditions C_c is not constant but decreases with increasing

light intensity, leading to an apparent Kok effect (Farquhar & Busch, 2017; Yin *et al.*, 2020). The

combined FvCB and g_m model, Eqn (4), can account for the decline of C_c with increasing irradiance

(Farquhar & Busch, 2017), and thus, in principle, can estimate R_d for PR conditions. This equation

was previously used by Yin & Struik (2009) to estimate g_m, and here we use it to estimate g_m and R_d

simultaneously, by exploring both gas exchange and chlorophyll fluorescence data across a range of

low light intensities after adopting the values of calibration factor s (see Eqn (3)) from data measured

under NPR conditions. The principle is in analogy to the 2-D modelling of Berghuijs *et al.* (2019) that

accounts for the reassimilation of (photo)respired CO_2, but with the benefit that Eqn (4) is considerably

simpler than the 2-D model. It is also in analogy to the procedure of Brooks & Farquhar (1985) that

corrects for the decrease in C_i, and of Ayub *et al.* (2011) that further corrects for the decrease in C_c,

with increasing irradiance, but with the benefit that the NRH fitting method is easier to implement.

The estimation of g_m is very sensitive to measurement errors (Yin & Struik, 2009). Here, we failed to

estimate g_m in some cases due to the variation in data from stressed plants (Table S5). Nevertheless,

the estimates of R_d could still be precise as reflected by the low standard error of estimates of R_d. The

NRH method has the following advantages. First, this method can estimate R_d under PR conditions by

implicitly considering the variation in C_c with irradiance as it corrects for the error of the linear

regression methods assuming a constant C_c under PR conditions. Second, this method only requires

gas exchange and chlorophyll fluorescence data, but does not require sophisticated and expensive

isotopic devices as required by direct R_d measuring methods (e.g. Haupt-Herting *et al.*, 2001; Gong *et*

al., 2015) or leaf anatomical data as required by the method of Berghuijs *et al.* (2019). Third, it

provides additional estimates for parameters g_m, which could be recognized as indicators of plant

physiological processes in response to environmental variables.

Our results showed that light inhibition of R_{dk} was higher under PR conditions than under NPR

conditions (Fig. 2), consistent with the result of Yin *et al*. (2020) that apparent light inhibition of

respiration and thus Kok effect are not obvious under low O_2 or high CO_2 conditions or their

combinations. Moreover, our data also showed that the underestimation of R_d by the Kok and Yin

methods at photorespiratory conditions may lead to an overestimation of light inhibition of leaf

respiration (Fig. 2C,D). Berghuijs *et al.* (2019) suggested that R_d estimated by the Yin and the Kok

methods at NPR conditions cannot represent the real R_d at PR conditions. Here, our results showed

that values of R_d for NPR conditions were generally higher than those for PR conditions (Fig. 3B,C),

which was in agreement with the result of Yin *et al.* (2020) that real light inhibition on respiration

increases with increasing amount of photorespiration. Again, the reason for the greater light

suppression of R_d under PR conditions remains to be elucidated.

Concluding remarks

In contrast to the plethora of studies that have explored the responses of R_{dk} to contrasting environments, we assessed the extent to which R_d of wheat leaves acclimated to drought and growth temperature. We proved a simple method that can estimate R_d for photorespiratory conditions by using gas exchange and chlorophyll fluorescence data. It was demonstrated that as for non-photorespiratory conditions, R_d and its temperature response for photorespiratory conditions acclimated more to drought than to growth temperature. Understanding this acclimation of R_d is needed to support the modelling of R_d, and thus of crop productivity and of carbon cycling in agricultural ecosystems, under future climate change.

Figure 1 Correlations between leaf day respiration estimated by the Kok method ($R_{d(Kok)}$) and estimated by the Yin method ($R_{d(Yin)}$) under non-photorespiratory (NPR; panel A) or photorespiratory (PR; panel B) conditions, and correlations between $R_{d(Kok)}$ (panel C) or $R_{d(Yin)}$ (panel D) and leaf day respiration estimated by the NRH method ($R_{d(NRH)}$) under PR conditions, across wheat genotypes Thésée and Récital, growth temperature and water treatments, and leaf temperatures during measurements. Note that the Yin method suits to estimate R_d for the NPR conditions only; as stated in the text, it is applied also to PR conditions here merely for the comparison purpose. The dashed diagonal represents the 1:1 relationship. The number (%) in each panel is the average of y-axis relative to *x*-axis values.

Figure 2 Correlations between leaf day respiration estimated by the Kok ($R_{d(Kok)}$), Yin ($R_{d(Yin)}$) or NRH ($R_{d(NRH)}$) methods and leaf respiration in the dark (R_{dk}) under non-photorespiratory (NPR; upper panels) and photorespiratory (PR; lower panels) conditions, across wheat genotypes Thésée and Récital, growth temperature and water treatments, and leaf temperatures during measurements. Note that the Yin method suits to estimate R_d for the NPR conditions only; as stated in the text, it is applied also to PR conditions here merely for the comparison purpose. The dashed diagonal represents the 1:1 relationship. The number (%) in each panel is the average of y-axis relative to x-axis values.

Figure 3 Correlations between leaf respiration in the dark (R_{dk}) under NPR conditions versus R_{dk} under PR conditions (panel A), and correlations between leaf day respiration estimated by the Yin method ($R_{d(Yin)}$) under NPR conditions versus $R_{d(Yin)}$ under PR conditions (panel B) or that estimated by the NRH method ($R_{d(NRH)}$) under PR conditions (panel C), across wheat genotypes Thésée and Récital, growth temperature and water treatments, and leaf temperatures during measurements. Note that the Yin method suits to estimate R_d for the NPR conditions only; as stated in the text, it is applied also to PR conditions here merely for the comparison purpose. The dashed diagonal represents the 1:1 relationship. The number (%) in each panel is the average of y-axis relative to *x*-axis values.

Figure 4 Thermal responses of day respiration estimated by the Yin method ($R_{d(Yin)}$) for non-photorespiratory (NPR) conditions (panel A) or by the NRH method ($R_{d(NRH)}$) for photorespiratory (PR) conditions (panel B) in two wheat genotypes (Thésée and Récital) under well-watered (WW) and drought-stressed (DS) conditions in EXP2019. The filled points and solid lines represent the WW plants, the open points and dashed lines represent the DS plants. Black symbols and lines refer to Thésée, while gray symbols and lines refer to Récital. Lines are the Arrhenius equation fitted to the data. Error bars indicate standard error of the means (SE) (n = 4).

Figure 5 Thermal responses of day respiration estimated by the Yin method ($R_{d(Yin)}$) for non-photorespiratory (NPR) conditions (left panels) or by the NRH method ($R_{d(NRH)}$) for photorespiratory (PR) conditions (right panels) in winter wheat Thésée grown at three growth temperatures (HT: high temperature, MT: medium temperature, and LT: low temperature) under well-watered (WW; upper panels) and drought-stressed (DS; lower panels) conditions in EXP2020. The filled points and solid lines represent the well-watered plants, the open points and dashed lines represent the drought-stressed plants. Lines are the Arrhenius equation fitted to the data. Error bars indicate standard error of the means (SE) (n=4).

568 **Table 1** Values of modelled leaf day respiration at 25 °C (R_{d25}) and activation energy for leaf day respiration (E_{Rd}) estimated by the Arrhenius

569 equation for two genotypes of wheat (Thésée and Récital) under well-watered (WW) and drought-stressed (DS) conditions in EXP2019. Data of

570 leaf day respiration used for fitting the Arrhenius equation was estimated by either the Yin method ($R_{d(Yin)}$) under non-photorespiratory (NPR)

571 conditions or the NRH method ($R_{d(NRH)}$) under photorespiratory (PR) conditions. Standard errors of the estimates are between brackets.

	Treatment				
Condition	Genotype	Water regime	R_{d25} (μmol m^{-2} s^{-1})	E_{Rd} (kJ mol^{-1})	r^2
NPR conditions ($R_{d(Yin)}$)	Thésée	WW	1.45 (0.09)	62.29 (3.72)	0.991
		DS	0.81 (0.08)	23.53 (7.56)	0.746
	Récital	WW	1.25 (0.06)	68.64 (3.05)	0.996
		DS	0.56 (0.15)	41.18 (17.74)	0.615
PR conditions ($R_{d(NRH)}$)	Thésée	WW	1.27 (0.04)	58.87 (2.00)	0.997
		DS	0.58 (0.15)	40.90 (16.93)	0.716
	Récital	WW	1.04 (0.05)	60.67 (2.72)	0.995
		DS	0.52 (0.10)	40.94 (12.87)	0.765

574 **Table 2** Values of modelled leaf day respiration at 25 °C (R_{d25}), activation energy for leaf day respiration (E_{Rd}) estimated by the Arrhenius equation for

575 wheat Thésée grown at three growth temperatures (HT: high temperature, AMT: medium temperature and LT: low temperature) under well-watered

576 (WW) and drought-stressed (DS) conditions in EXP2020. Data of leaf day respiration used for fitting the Arrhenius equation was estimated by

577 either the Yin method ($R_{d(Yin)}$) under non-photorespiratory (NPR) conditions or the NRH method ($R_{d(NRH)}$) under photorespiratory (PR) conditions.

578 Standard errors of the estimates are between brackets.

| | | Treatment | | | |
| | | Water | R_{d25} | E_{Rd} | |
Condition	Genotype	regime	(μmol m^{-2} s^{-1})	(kJ mol^{-1})	r^2
NPR conditions ($R_{d(Yin)}$)	HT	WW	1.26 (0.15)	70.69 (7.15)	0.978
		DS	0.78 (0.13)	28.56 (11.86)	0.672
	MT	WW	1.36 (0.16)	64.58 (7.44)	0.969
		DS	1.13 (0.11)	26.96 (7.27)	0.816
	LT	WW	1.60 (0.18)	58.00 (10.35)	0.942
		DS	1.16 (0.08)	8.10 (6.09)	0.329
PR conditions ($R_{d(NRH)}$)	HT	WW	1.22 (0.14)	60.90 (7.39)	0.967
		DS	0.51 (0.08)	47.10 (9.86)	0.902
	MT	WW	1.27 (0.03)	59.17 (1.50)	0.998
		DS	0.95 (0.14)	36.09 (10.35)	0.816
	LT	WW	1.52 (0.14)	55.83 (8.39)	0.944
		DS	0.78 (0.07)	12.93 (8.55)	0.395

580 **Author contributions**

581 XY and LF conceived the study. LF and PELvdP designed the experiment and conducted the
582 measurements. LF and XY analyzed the data. LF wrote the draft and finalized it with significant input
583 from the other coauthors.

References

Amthor JS. **2010**. From sunlight to phytomass: on the potential efficiency of converting solar radiation to phyto-energy. *New Phytologist* **188**: 939–959.

Atkin OK, Bruhn D, Hurry VM, Tjoelker MG. **2005**. The hot and the cold: unravelling the variable response of plant respiration to temperature. *Functional Plant Biology* **32**: 87–105.

Atkin OK, Macherel D. **2009**. The crucial role of plant mitochondria in orchestrating drought tolerance. *Annals of Botany* **103**: 581–597.

Atkin OK, Scheurwater I, Pons T. **2006**. High thermal acclimation potential of both photosynthesis and respiration in two lowland *Plantago* species in contrast to an alpine congeneric. *Global Change Biology* **12**: 500–515.

Atkin OK, Scheurwater I, Pons TL. **2007**. Respiration as a percentage of daily photosynthesis in whole plants is homeostatic at moderate, but not high, growth temperatures. *New Phytologist* **174**: 367–380.

Atkin OK, Tjoelker MG. **2003**. Thermal acclimation and the dynamic response of plant respiration to temperature. *Trends in Plant Science* **8**: 343–351.

Ayub G, Smith RA, Tissue DT, Atkin OK. **2011**. Impacts of drought on leaf respiration in darkness and light in *Eucalyptus saligna* exposed to industrial-age atmospheric CO_2 and growth temperature. *New Phytologist* **190**: 1003–1018.

Barreto P, Yassitepe JECT, Wilson ZA, Arruda P. **2017**. Mitochondrial uncoupling protein 1 overexpression increases yield in nicotiana tabacum under drought stress by improving source and sink metabolism. *Frontiers in Plant Science* **8**: 1–20.

Bartoli CG, Gomez F, Gergoff G, Guiamét JJ, Puntarulo S. **2005**. Up-regulation of the mitochondrial alternative oxidase pathway enhances photosynthetic electron transport under drought conditions. *Journal of experimental Botany* **56**: 1269–1276.

Berghuijs HNC, Yin X, Ho QT, Retta MA, Nicolaï BM, Struik PC. **2019**. Using a reaction-diffusion model to estimate day respiration and reassimilation of (photo)respired CO_2 in leaves. *New Phytologist* **223**: 619–631.

Bernacchi CJ, Portis AR, Nakano H, Von Caemmerer S, Long SP. **2002**. Temperature response of mesophyll conductance. Implications for the determination of Rubisco enzyme kinetics and for limitations to photosynthesis in vivo. *Plant physiology* **130**: 1992–1998.

Brooks A, Farquhar GD. **1985**. Effect of temperature on the CO_2/O_2 specificity of ribulose-1,5-bisphosphate carboxylase/oxygenase and the rate of respiration in the light. *Planta* **165**: 397–406.

Buckley TN, Vice H, Adams MA. **2017**. The Kok effect in Vicia faba cannot be explained solely by changes in chloroplastic CO_2 concentration. *New Phytologist* **216**: 1064–1071.

von Caemmerer S. **2000**. *Biochemical models of leaf photosynthesis*. Collingwood, VIC, Australia: CSIRO Publishing.

von Caemmerer S. **2013**. Steady-state models of photosynthesis. *Plant, Cell & Environment* **36**: 1617–1630.

von Caemmerer S, Farquhar G, Berry J. **2009**. Biochemical model of C_3 photosynthesis. In: Laisk A, Nedbal L, Govindjee, eds. *Photosynthesis in silico: understanding complexity from molecules to ecosystems.* Dordrecht, the Netherlands: Springer Science + Business Media, 209–230.

Cai C, Li G, Di L, Ding Y, Fu L, Guo X, Struik PC, Pan G, Li H, Chen W, *et al.* 2020. The acclimation of leaf photosynthesis of wheat and rice to seasonal temperature changes in T-FACE environments. *Global Change Biology* **26**: 539–556.

Coast O, Posch BC, Bramley H, Gaju O, Richards RA, Lu M, Ruan YL, Trethowan R, Atkin OK. **2020**. Acclimation of leaf photosynthesis and respiration to warming in field-grown wheat. *Plant Cell & Environment*: 1–16.

Cousins AB, Ghannoum O, Von Caemmerer S, Badger MR. **2010**. Simultaneous determination of Rubisco carboxylase and oxygenase kinetic parameters in Triticum aestivum and *Zea mays* using membrane inlet mass spectrometry. *Plant, cell & environment* **33**: 444–452.

Crous KY, Wallin G, Atkin OK, Uddling J, Ekenstam AA. **2017**. Acclimation of light and dark respiration to experimental and seasonal warming are mediated by changes in leaf nitrogen in *Eucalyptus globulus*. *Tree Physiology* **37**: 1069–1083.

Crous KY, Zaragoza-Castells J, Ellsworth DS, Duursma RA, Löw M, Tissue DT, Atkin OK. **2012**. Light inhibition of leaf respiration in field-grown *Eucalyptus saligna* in whole-tree chambers under elevated atmospheric CO_2 and summer drought. *Plant, Cell & Environment* **35**: 966–981.

Crous KY, Zaragoza-Castells J, Löw M, Ellsworth DS, Tissue DT, Tjoelker MG, Barton CVM, Gimeno TE, Atkin OK. **2011**. Seasonal acclimation of leaf respiration in *Eucalyptus saligna* trees: Impacts of elevated atmospheric CO_2 and summer drought. *Global Change Biology* **17**: 1560–1576.

Dahal K, Vanlerberghe GC. **2017**. Alternative oxidase respiration maintains both mitochondrial and chloroplast function during drought. *New Phytologist* **213**: 560–571.

Farquhar GD, Busch FA. **2017**. Changes in the chloroplastic CO_2 concentration explain much of the observed Kok effect: a model. *New Phytologist* **214**: 570–584.

Farquhar GD, von Caemmerer S von, Berry JA. **1980**. A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species. *Planta* **149**: 78–90.

Flexas J, Galmes J, Ribas-Carbo M, Medrano H. **2005**. The effects of water stress on plant respiration. In: Lambers H, Ribas-Carbo M, eds. *Plant respiration: from cell to ecosystem.* Dordrecht, the Netherlands: Springer, 85–94.

Gauthier PPG, Crous KY, Ayub G, Duan H, Weerasinghe LK, Ellsworth DS, Tjoelker MG, Evans JR, Tissue DT, Atkin OK. **2014**. Drought increases heat tolerance of leaf respiration in *Eucalyptus globulus* saplings grown under both ambient and elevated atmospheric [CO_2] and temperature. *Journal of Experimental Botany* **65**: 6471–6485.

Genty B, Briantais JM, Baker NR. **1989**. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta - General Subjects* **990**: 87–92.

Gifford RM. **1995**. Whole plant respiration and photosynthesis of wheat under increased CO_2 concentration and temperature: long-term vs. short-term distinctions for modelling. *Global Change Biology* **1**: 385–396.

Gimeno TE, Sommerville KE, Valladares F, Atkin OK. **2010**. Homeostasis of respiration under drought and its important consequences for foliar carbon balance in a drier climate: insights from two contrasting *Acacia* species. *Functional Plant Biology* **37**: 323–333.

Gong XY, Schäufele R, Feneis W, Schnyder H. **2015**. $^{13}CO_2/^{12}CO_2$ exchange fluxes in a clamp-on leaf cuvette: Disentangling artefacts and flux components. *Plant Cell & Environment* **38**: 2417–2432.

Gulías J, Flexas J, Abadía A, Medrano H. **2002**. Photosynthetic responses to water deficit in six Mediterranean sclerophyll species: possible factors explaining the declining distribution of *Rhamnus ludovici-salvatoris*, an endemic Balearic species. *Tree Physiology* **22**: 687–697.

Haupt-Herting S, Klug K, Fock HP. **2001**. A new approach to measure gross CO_2 fluxes in leaves. Gross CO_2 assimilation, photorespiration, and mitochondrial respiration in the light in tomato under drought stress. *Plant Physiology* **126**: 388–396.

Heskel MA, Atkin OK, Turnbull MH, Griffin KL. **2013**. Bringing the Kok effect to light: A review on the integration of daytime respiration and net ecosystem exchange. *Ecosphere* **4**: 1–14.

IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani (eds.)]. Cambridge University Press. In Press.

De Kauwe MG, Lin YS, Wright IJ, Medlyn BE, Crous KY, Ellsworth DS, Maire V, Prentice IC, Atkin OK, Rogers A, *et al.* 2016. A test of the 'one-point method' for estimating maximum carboxylation capacity from field-measured, light-saturated photosynthesis. *New Phytologist* **210**: 1130–1144.

Kok B. **1948**. *A Critical consideration of the quantum yield of Chlorella-photosynthesis. Proefschrift ter verkrijging van de graad van doctor in de wis-en natuurkunde aan de Rijksuniversiteit te Utrecht... door Bessel Kok...* W. Junk.

Kumarathunge DP, Drake JE, Tjoelker MG, López R, Pfautsch S, Vårhammar A, Medlyn BE. **2020**. The temperature optima for tree seedling photosynthesis and growth depend on water inputs. *Global Change Biology* **26**: 2544–2560.

Laisk AK. **1977**. Kinetics of photosynthesis and photorespiration in C_3-plants.[In Russian] Nauka.

Lloret F, Escudero A, Iriondo JM, Martínez-Vilalta J, Valladares F. **2012**. Extreme climatic events and vegetation: The role of stabilizing processes. *Global Change Biology* **18**: 797–805.

Lobell DB, Gourdji SM. **2012**. The influence of climate change on global crop productivity. *Plant Physiology* **160**: 1686–1697.

Loriaux SD, Avenson TJ, Welles JM, Mcdermitt DK, Eckles RD, Riensche B, Genty B. **2013**. Closing in on maximum yield of chlorophyll fluorescence using a single multiphase flash of sub-saturating intensity. *Plant, Cell & Environment* **36**: 1755–1770.

Nantes IL, Fagian MM, Catisti R, Arruda P, Maia IG, Vercesi AE. **1999**. Low temperature and aging-promoted expression of PUMP in potato tuber mitochondria. *FEBS Letters* **457**: 103–106.

O'Leary BM, Asao S, Millar AH, Atkin OK. **2019**. Core principles which explain variation in respiration across biological scales. *New Phytologist* **222**: 670–686.

Perkins-Kirkpatrick SE, Lewis SC. **2020**. Increasing trends in regional heatwaves. *Nature Communications* **11**: 1–8.

Pons TL, Flexas J, Von Caemmerer S, Evans JR, Genty B, Ribas-Carbo M, Brugnoli E. **2009**. Estimating mesophyll conductance to CO2: methodology, potential errors, and recommendations. *Journal of Experimental Botany* **60**: 2217–2234.

Searle SY, Thomas S, Griffin KL, Horton T, Kornfeld A, Yakir D, Hurry V, Turnbull MH. **2011**. Leaf respiration and alternative oxidase in field-grown alpine grasses respond to natural changes in temperature and light. *New Phytologist* **189**: 1027–1039.

Shao L, Liu Z, Li H, Zhang Y, Dong M, Guo X, Zhang H, Huang B, Ni R, Li G. **2021**. The impact of global dimming on crop yields is determined by the source–sink imbalance of carbon during grain filling. *Global Change Biology* **27**: 689–708.

Slot M, Zaragoza-Castells J, Atkin OK. **2008**. Transient shade and drought have divergent impacts on the temperature sensitivity of dark respiration in leaves of Geum urbanum. *Functional Plant Biology* **35**: 1135–1146.

Solomon S, Plattner GK, Knutti R, Friedlingstein P. **2009**. Irreversible climate change due to carbon dioxide emissions. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 1704–1709.

Sperlich D, Barbeta A, Ogaya R, Sabaté S, Peñuelas J. **2016**. Balance between carbon gain and loss under long-term drought: Impacts on foliar respiration and photosynthesis in *Quercus ilex* L. *Journal of Experimental Botany* **67**: 821–833.

Tcherkez G, Atkin OK. **2021**. Unravelling mechanisms and impacts of day respiration in plant leaves: an introduction to a Virtual Issue. *New Phytologist* **230**: 5–10.

Tcherkez G, Gauthier P, Buckley TN, Busch FA, Barbour MM, Bruhn D, Heskel MA, Gong XY, Crous KY, Griffin K, *et al.* 2017a. Leaf day respiration: low CO_2 flux but high significance for metabolism and carbon balance. *New Phytologist* **216**: 986–1001.

Tcherkez G, Gauthier P, Buckley TN, Busch FA, Barbour MM, Bruhn D, Heskel MA, Gong XY, Crous K, Griffin KL, *et al.* 2017b. Tracking the origins of the Kok effect, 70 years after its discovery. *New Phytologist* **214**: 506–510.

Tjoelker MG, Reich PB, Oleksyn J. **1999**. Changes in leaf nitrogen and carbohydrates underlie temperature and CO_2 acclimation of dark respiration in five boreal tree species. *Plant, Cell & Environment* **22**: 767–778.

Villar R, Held AA, Merino J. **1994**. Comparison of methods to estimate dark respiration in the light in leaves of two woody species. *Plant Physiology* **105**: 167–172.

Way DA, Aspinwall MJ, Drake JE, Crous KY, Campany CE, Ghannoum O, Tissue DT, Tjoelker MG. **2019**. Responses of respiration in the light to warming in field-grown trees: a comparison of the thermal sensitivity of the Kok and Laisk methods. *New Phytologist* **222**: 132–143.

Yin X, Belay DW, Van Der Putten PEL, Struik PC. **2014**. Accounting for the decrease of photosystem photochemical efficiency with increasing irradiance to estimate quantum yield of leaf photosynthesis. *Photosynthesis Research* **122**: 323–335.

Yin X, Busch FA, Struik PC, Sharkey TD. **2021**. Evolution of a biochemical model of steady-state photosynthesis. *Plant, Cell & Environment*.

Yin X, Niu Y, van der Putten PEL, Struik PC. **2020**. The Kok effect revisited. *New Phytologist* **227**: 1764–1775.

Yin X, Van Oijen M, Schapendonk AHCM. **2004**. Extension of a biochemical model for the generalized stoichiometry of electron transport limited C_3 photosynthesis. *Plant, Cell & Environment* **27**: 1211–1222.

Yin X, Struik PC. **2009**. Theoretical reconsiderations when estimating the mesophyll conductance to CO_2 diffusion in leaves of C_3 plants by analysis of combined gas exchange and chlorophyll fluorescence measurements. *Plant, Cell & Environment* **32**: 1513–1524.

Yin X, Struik PC. **2017**. Can increased leaf photosynthesis be converted into higher crop mass production? A simulation study for rice using the crop model GECROS. *Journal of Experimental Botany* **68**: 2345–2360.

Yin X, Struik PC, Romero P, Harbinson J, Evers JB, Van Der Putten PEL, Vos J. **2009**. Using combined measurements of gas exchange and chlorophyll fluorescence to estimate parameters of a biochemical C_3 photosynthesis model: A critical appraisal and a new integrated approach applied to

leaves in a wheat (*Triticum aestivum*) canopy. *Plant, Cell & Environment* **32**: 448–464.

Yin X, Sun Z, Struik PC, Gu J. **2011**. Evaluating a new method to estimate the rate of leaf respiration in the light by analysis of combined gas exchange and chlorophyll fluorescence measurements. *Journal of Experimental Botany* **62**: 3489–3499.

Zagdańska B. **1995**. Respiratory energy demand for protein turnover and ion transport in wheat leaves upon water deficit. *Physiologia Plantarum* **95**: 428–436.