Title: On the nature of the earliest known lifeforms

Dheeraj Kanaparthi[1,2&3*], Frances Westall[4], Marko Lampe[5], Baoli Zhu[2&6], Thomas Boesen[7], Bettina Scheu[3] Andreas Klingl[8], Petra Schwille[1] & Tillmann Lueders[2*]

Affiliations:

[1]Max-Planck Institute for Biochemistry, Munich, Germany

[2]Chair of Ecological Microbiology, BayCeer, University of Bayreuth, Germany.

[3]Earth and environmental sciences, Ludwig Maximilian University, Munich, Germany.

[4]CNRS-Centre de Biophysique Moléculaire, Orléans, France.

[5]Advanced Light Microscopy Facility, European Molecular Biology Laboratory, Heidelberg, Germany.

[6]Key Laboratory of Agro-Ecological Processes in Subtropical Regions, Taoyuan Agroecosystem Research Station, Institute of Subtropical Agriculture, Chinese Academy of Sciences, Changsha, China.

[7]Department of Biosciences, Center for Electromicrobiology, Aarhus, Denmark.

[8]Department of Botany I, Ludwig Maximilian University, Munich, Germany.

*Corresponding authors

Lead contact: kanaparthi@biochem.mpg.de

Abstract

Microfossils from the Paleoarchean Eon are the oldest known evidence of life. Despite their

significance in understanding the history of life on Earth, any interpretation of the nature of

these microfossils has been a point of contention among researchers. Decades of back-and-

forth arguments led to the consensus that reconstructing the lifecycles of Archaean Eon

organisms is the only way of understanding the nature of these microfossils. Here, we

transformed a Gram-positive bacterium into a primitive lipid vesicle-like state and studied it

under environmental conditions prevalent on early Earth. Using this approach, we

successfully reconstructed morphologies and life cycles of Archaean microfossils. In addition

to reproducing microfossil morphologies, we conducted experiments that spanned years to

understand the process of cell degradation and how Archaean cells could have undergone

encrustation minerals (in this case, salt), leading to their preservation as fossilized organic

carbon in the rock record. These degradation products strongly resemble fossiliferous features

from Archaean rock formations. Our observations suggest that microfossils aged between 3.8

to 2.5Ga most likely were liposome-like protocells that have evolved physiological pathways

of energy conservation but not the mechanisms to regulate their morphology. Based on these

observations, we propose that morphology is not a reliable indicator of taxonomy in these

microfossils.

Introduction

The Pilbara Greenstone Belt (PGB), Western Australia, and Barberton Greenstone Belt

(BGB), South Africa, are the oldest known rock formations that have not undergone

significant metamorphic alterations (1,2). Hence, these rock formations have been the subject

of numerous scientific investigations focused on understanding the biology and

biogeochemistry of Archaean Earth (1–5). Over a span of 50 years, these studies have

documented various organic structures within these rock formations that resemble fossilized

cells and their degradation products, with $\delta^{13}C$ composition consistent with biologically

derived organic carbon (6,7). Although these observations suggest that these organic

structures were fossil remnants of Archaean microorganisms, any such interpretation,

together with the biological origin of these structures, has been a point of contention among

researchers (8–11). Two factors currently limit wider acceptance of their biological origin –

the absence of truly analogous microfossil morphologies among extant prokaryotes and an

indication of an ongoing biological process, like cell division, among most microfossils

(4,12). Moreover, most of the described microfossils are larger than present-day prokaryotes

and often exhibit considerable cytoplasmic complexity (3,5,13). These complex

morphologies and relatively larger cell sizes of supposedly primitive Archaean Eon cells is

not in accordance with our current understanding of how biological complexity evolved

through Darwinian evolution (14,15).

Apart from the chemical and mineralogical considerations (13,16–18), one key emphasis of

the studies arguing for and against the biological origin of Archaean Eon organic structures

involves an extensive morphological comparison with extant prokaryotes or abiotically

formed minerals (8,9,19,20). Cell morphology among extant organisms is maintained by a

plethora of intracellular processes and is determined by the information encoded in their

genome (21). In our opinion, drawing parallels between the present-day prokaryotes and

Archaean Eon organisms inherently involves subscribing to the notion that paleo-Archaean

life forms possess all the complex molecular biological mechanisms to regulate their

morphology as present-day cells. Any such presumptions are not in tune with the current

scientific consensus of how life could have originated on early Earth (14,22,23). It is now

widely believed that life evolved in the form of protocells devoid of most molecular

biological complexity (23). These primitive cells are thought to have undergone slow

Darwinian evolution, resulting in present-day cells with intricate intracellular processes

(24,25). Given the unlikelihood of Archaean cells possessing complex molecular biological

processes, we test the possibility that complex morphologies of Archaean microfossils result

from the complete absence of intracellular mechanisms regulating their morphology.

(26).

In this study, we have used an alternative approach to understand early life. We transformed a

Gram-positive bacterium (*Exiguobacterium Strain-Molly*) into its primitive lipid vesicle-like

state (*EM-P*)(27,28). Cells in this state can be described as a simple sack of cytoplasm devoid

of all mechanisms to regulate their morphology and reproduction (27). Although it has not

been empirically demonstrated, some studies have suggested that cells in this lipid vesicle-

like state likely resemble primitive protocells, and their morphology and reproduction are

determined by environmental conditions (27–29). Hence, we subsequently studied the life

cycle of these cells in their native environment (conditions prevalent on early Earth). While

the precise environmental conditions of early Earth remain uncertain, a growing consensus

within the scientific community suggests that surface temperatures on Archaean Earth ranged between 26° and 35°C (30–32). Cells on early Earth, particularly the microfossils relevant to the present study, could also have been well acclimatized to high salt environments due to their prevalence in coastal marine environments (6,33). Hence, to replicate the high salinities of the coastal marine environments, *EM-P* was cultivated in half-strength tryptic soy broth supplemented with 7% (w/v) Dead Sea Salt (TSB-DSS) at 30°C. We chose Dead Sea Salt over pure NaCl to better emulate complex salt compositions of natural environments.

Given that *EM-P*'s life cycle and the biophysical basis of such a reproduction process is extensively discussed in the previous manuscript (27), the primary focus of this manuscript will be restricted to the morphological comparison of *EM-P* cells and Archaean Eon microfossils. Below, we present the morphological comparison between *EM-P* cells and Archaean Eon microfossils. In addition to this morphological comparison, we also conducted experiments that spanned years to understand the process of protocell degradation, how they become encrusted in salt, and how they are preserved as fossilized organic carbon in the rock record.

Results

Morphological comparison of top-down modified cells with fossilized Archaean cells

When cultured under the environmental conditions of early Earth, *EM-P* exhibited cell sizes that were an order of magnitude larger than their original size. They also exhibited complex morphologies and reproduced by a relatively less understood process (27,28). The life cycle of these cells involves reproduction by two methods – via forming internal or a string of external daughter cells (27). *EM-P* reproducing by both these processes bears close morphological resemblance to microfossils reported from the Archaean Eon.

Fig 1. Morphological comparison of the Apex Chert and the Strelley Pool Formation

microfossils with *EM-P*. Images A-D are TEM images of *EM-P* cells forming ICVs and

intracellular daughter cells. The numbered arrows in these images point to different stages of

ICV formation (see Fig. S1). Images E, F, K & L show TEM, SEM, and STED microscope

images of *EM-P* cells with ICVs and surface depressions (black arrows). Cells in image F

were stained with universal membrane stain, FM^{TM}5-95 (red), and DNA stain, PicoGreen

(green). Images G-J & M are spherical microfossils reported from the Apex Chert and the

Strelley Pool Formation, respectively (originally published by Schopf et al., 1987 & Delarue

et al., 2019)(34,35). Cyan arrows in images E-H point to cytoplasm sandwiched between

large hollow vesicles. The arrow in the image I point to the dual membrane enclosing the

microfossil. Morphologically similar images of *EM-P* cells are shown in Fig. S3. Black

arrows in images K-M point to surface depressions in both *EM-P* and the Strelley Pool

Formation microfossils, possibly formed by the rupture of ICV's as shown in D & E (arrows)

(also see Fig. S4-S6). Scale bars: A-D (0.5µm) E, K & L (2 µm), and 5 µm (F).

The first step in reproduction by intracellular daughter cells is the formation of hollow

intracellular vesicles (Fig. 1A). These vesicles were formed by a process that resembles

endocytosis (Fig. 1A). Over time, the number of intracellular vesicles (ICV's) within *EM-P*

gradually increased (Fig. 1A-F). No uniformity was observed in the size of ICVs within a cell

or the number of ICVs among different cells (Fig. 1E, 1F, S1 & S2). *EM-P* cells with such

intracellular vesicles resemble spherical microfossils reported from 3.46 billion-year-old (Ga)

Apex chert (34). Like the Apex-chert microfossils, ICVs of *EM-P* were hollow, and organic carbon (cytoplasm) in these cells is restricted to spaces between the vesicles (Fig. 1F-K & S3).

The three-dimensional STED and SEM images of *EM-P* show numerous surface depressions (Fig. 1K, arrow & S1E). Such depressions are formed either during vesicle formation (Fig. 1A & 1B, arrow) or by the rupture of intracellular vesicles attached to the cell membrane (Fig. 1D, 1E, S1G-K & S4-S6). *EM-P* cells with such surface depressions and intracellular vesicles bear uncanny morphological resemblance to microfossils reported from 3.4 Ga Strelley Pool Formation (SPF) microfossils (35) (Fig. 1M & Fig. S4-S6). Microfossils reported from other sites, such as the Farrel Quarzite (Fig. S7)(36), Turee Creek (Fig. S8)(37), and the Fig Tree Formations (Fig. S9)(38), likely are morphological variants of Archaean *EM-P*-like cells and the Apex Chert microfossils. For instance, *EM-P* with many but relatively smaller intracellular vesicles resemble the Fig Tree microfossils, both in cell size and shape (Fig. S9). On the other hand, *EM-P,* with a relatively larger number of intracellular vesicles squeezed into polygonal shapes, resemble microfossils reported from the SPF (Fig. 1G-I & S4-S6), the Farrel Quartzite microfossils with polygonal alveolar structures (Fig. S7) and the Turee creek microfossils (Fig. S8)(36).

The second step in this method of reproduction involves the formation of daughter cells into the ICVs (Fig. S10)(27). Daughter cells were formed in the intracellular vesicles by a process resembling budding (Fig S10). Over time, these bud-like daughter cells detached from the vesicle wall and were released into the ICV (Movie 1). Due to a gradual loss of cytoplasm to the daughter cells, we observed a gradual reduction of the cytoplasmic volume of the parent *EM-P* cells and a corresponding increase in the number of daughter cells within the ICVs (Fig. S11, Movie 2-4). Over time, *EM-P* cells transformed into hollow vesicles with multiple tiny daughter cells (Fig. S11e-h, Movie 3&4). These intracellular daughter cells were

Fig 2. Morphological comparison between the Mt. Goldsworthy microfossils and *EM-P*.

Images A-E show the process of cell lysis and release of intracellular vesicles in *EM-P*.

Image A shows an intact cell with intracellular vesicles. Images B-E show lysis and gradual

dispersion of these vesicles. Insert in image D shows enlarged images of individual ICVs.

Images F-I show spherical microfossils reported from the Mt. Goldsworthy formation

(originally published by Sugitani et al., 2009)(39). The arrow in this image, A & F, points to

a cell surrounded by an intact membrane. The black arrow in these images points to

filamentous extensions connecting individual vesicles. The boxed region in images D & I

highlights a similar discontinuous distribution of organic carbon in ICVs and microfossils.

Also see Fig. S13-S17. Scale bars: 20μm (A-E) & 20μm (F-I).

released into the surroundings by a two-step process. In the first step, cells underwent lysis to

release the ICVs (Fig. 2, S12 A & B and Movies. 5-9). In the second step, the vesicle

membrane underwent lysis (Fig. S12C-E) to release the daughter cells (Fig. S12F-H). *EM-P*

cells undergoing this method of reproduction closely resemble fossilized microbial cells

discovered from several Archaean Eon rock formations (Fig. 2 & S13-S23)(35,39,40). For instance, microfossils reported from Mt. Goldsworthy Formation exhibit cells with ICVs containing daughter cells, cells that underwent lysis to release these ICVs (Fig. 2 & S13-S17), and the subsequent rupture of the vesicle membrane and release of the daughter cells (Fig. S14D-F)(27).

The microfossils reported from sites like the Farrel Quartzite (Fig. S7)(36), the Strelley Pool Formation (Fig. S18-S20), the Waterfall locality (Fig. S21-S22)(39,42,43), the Turee Creek (Fig. S23)(40), and Dresser Formation (Fig. S24-S28)(10), bear close morphological resemblance with morphologies of *EM-P* cells reproducing by this process. *EM-P* cells exhibited all the distinctive features of the Dresser formation microfossils, like the presence of hollow regions within the cell (Fig.S25) and discontinuous or thick-porous cell walls (Fig. S24 & S27). The step-by-step transformation of *EM-P* cells into these morphologies is shown in Fig. S27. Due to the lysis and release of daughter cells, most late stationary growth phase *EM-P* cells were deflated with numerous surface depressions. The morphology and surface texture of such deflated *EM-P* cells resemble the morphologies of microfossils reported from the Kromberg Formations (Fig. S29)(44).

Reproduction by external daughter cells happens by two different processes. Tiny daughter cells initially appeared as buds attached to the cell membrane (Fig. S30, S31 & Movie 10). These buds subsequently grew in size and detached from the parent cell. Depending on the size of the daughter cells (buds)*, EM-P* cells appear to have been reproducing either by budding or binary fission. *EM-P* cells that appear to have been reproducing by budding resemble microfossils reported from the North-pole formations (Fig. S31). As observed in our incubations, microfossils from this site are a mix of individual spherical cells, spherical cells with pustular protuberances, cells with bud-like structures, and cells undergoing binary fission (Fig. S31). Other *EM-P* morphotypes, like individual spherical cells, hourglass-shaped cells undergoing fission, and cells in dyads, bear close morphological resemblance to microfossils reported from both the Swartkopie (Fig. S32) and the Sheba formations (Fig. S33 & S34)(12,33,46). Like the Sheba Formation microfossils, the cells undergoing binary fission were observed to be in close contact with extracellular organic carbon (clasts of organic carbon in the Sheba Formation). This extracellular organic carbon likely represents the intracellular constituents released during the lysis of cells, as described above (Movies 5-8 & Fig. S34).

In some cases, the above-described buds did not detach from the cell surface but transformed

into long tentacles (Movies 11&12). These initially hollow tentacles (27) gradually received

cytoplasm from the parent cells and gradually transformed into "string-of-spherical daughter

cells" (Fig. 3, S31A-C & S33)(27). Subsequently, these filaments detached from the parent

cell, and due to the constant motion of daughter cells within these filaments, the "string-of-

spherical daughter cells" fragmented into smaller and smaller strings and ultimately into

multiple individual daughter cells (Movies 13-15). Apart from the cells that received

cytoplasm (daughter cells), we also observed hollow spherical structures within these

tentacles that did not receive cytoplasm from the parent cells (Fig. 3F & 3G, black arrows).

Fig 3. Morphological comparison of the Cleaverville microfossils with *EM-P*: Images A-

E are the microfossils reported from Cleverville formation (originally reported by Ueno *et al.,*

2006). Images F-K are the *EM-P* cells that are morphologically analogous to the Cleverville

Formation microfossils. Open arrows in images A, B, F & G point to the membrane tethers

connecting the spherical cells within the filamentous extensions. Red arrows in the images point to the cells that have a similar distribution of organic carbon within the cells. Boxed and magnified regions in images B, F & G highlight the arrangement of cells in the filaments in pairs. The boxed region in image H highlights the cluster of hollow vesicles in *EM-P* incubations similar to the hollow organic structures in the Cleverville Formation, as shown in image c. Images D, E, and I-J show spherical cells that were largely hollow with organic carbon (cytoplasm) restricted to discontinuous patches at the periphery of the cell. Scale bars: 20μm (F-K).

All *EM-P* morphotypes observed undergoing this reproduction process bear close morphological resemblance to microfossils reported from the Cleaverville Formation (Fig. 3). All distinctive features of the Cleaverville microfossils, like the arrangement of cells as pairs within a string (Fig. 3H, 3F & 3G), microfossils with a discontinuous layer of organic carbon at the cell periphery (Fig. 3E, 3D & 3I-K), were also observed in *EM-P*. Several hollow spherical structures devoid of organic carbon were reported from the Cleaverville formation (Fig. 3). As in *EM-P*, these structures could have been the hollow membranous structures that didn't receive the cytoplasm from the parent cells. Similar structures were also reported from other microfossil sites, such as the organic structures reported from the Onverwacht Group (Fig. S35)(6).

In addition to the Cleaverville microfossils, *EM-P* cells in our incubations also resemble filamentous structures with spherical inclusions reported from the Sulphur Spring Formations (Fig. S36-S38)(17). Based on the morphological similarities, we propose that these structures could have been the filamentous extensions with spherical daughter cells observed in *EM-P*. Similar but smaller filamentous structures were reported from the Mt. Grant Formation (Fig. S39)(37) and could have been the shorter fragments of similar "strings-of-daughter cells" (Movie 14). The spherical structures with sparsely distributed organic carbon reported from the Sheba Formations could also have been such cells undergoing fragmentation into smaller and smaller filaments (Fig. S33). In tune with this proposition, the Sheba Formation microfossils, like the *EM-P* cells, exhibited an un-uniform distribution of organic carbon within the cells and filamentous overhangs (Fig. S33, Movie 16).

Five to seven days after the start of the experiment, most cells in the incubations were the daughter cells. We observed three distinct types of daughter cells – a string of daughter cells (Fig. 3, S39 & Movie 14), daughter cells that were still attached to the membrane debris of the parent cell (Fig. 4, 5, S40-S48 & Movie 17) and individual daughter cells (Fig. 3h & Movie 15). All these daughter cell morphotypes resemble – a cluster of tiny spherical globules reported from the SPF (Fig. S18), a string of daughter cells reported from Mt. Grant formation (Fig. S39)(37,39), and spherical daughter cells still attached to membrane debris of parent cell like the ones reported from the Sulphur Spring Formations, Mt.Goldsworthy, the Farrel Quartzite, the Moodies Group, the Dresser Formation, and the SPF (Fig. 4 & S40-S48)(10,39,41,43,49).

The gradual transformation of *EM-P* cells into lamination-like structures and their comparison with Archaean organic structures.

Fossilization and preservation of individual cells is considered unlikely due to the absence of rigid structures. However, recent studies indicate such a process could happen under favorable environmental conditions (50,51). However, the prevailing observations suggest that a significant portion of cell biomass undergoes taphonomic alteration and is preserved as degraded organic matter. To understand the possibility of *EM-P*-like cells forming structures similar to those observed in Archaean rocks, we studied the morphological transformation of individual *EM-P* cells and biofilms over 12 to 30 months. Below, we present the step-by-step transformation of individual cells into organic structures and their morphological resemblance to Archaean organic structures.

Fig. 4. Morphological comparison between *EM-P* and the Mt. Goldsworthy microfossils.

Images A, B & C are organic structures reported from the Mt. Goldsworthy Formation

(Sugitani *et al.,* 2009)(41). Image D shows morphologically analogous film-like membrane

debris observed in *EM-P* incubations. Arrows in images A-D point to either clusters or

individual spherical structures attached to these film-like structures. Scale bar: 50μm (A-C) &

10μm (D).

EM-P grew in our incubations as a biofilm at the bottom of the culture flask (Fig. 6 & Fig.

S49). The rapid biofilm formation by *EM-P* can be attributed to the presence of extracellular

DNA released during the lysis of *EM-P* cells (Fig. S50)(27). DNA released by such processes

is known to promote biofilm formation (52). Over time, increased cell numbers resulted in

biofilms comprising multiple layers of closely packed individual spherical cells (Fig. 6 &

S50). A subsequent increase in the number of cells led to lateral compression and

transformation of spherical cells into a polygonal shape (Fig. 6). By the late stationary growth

phase, most cells underwent such a transformation, resulting in a honeycomb-like biofilm.

The step-by-step transformation of individual spherical cells into these structures is shown in

Fig. 6. Morphologically similar organic structures were reported from several paleo-Archaean

sites, like the North Pole Formation (Fig. S51-S62)(45). A similar taphonomic degradation of

organic matter associated with a biofilm was demonstrated by previous studies (Westall et al., 2006, Figure 10B)(53).

Large aggregations of spherical cells devoid of internal organic carbon were reported from the North-Pole Formation (45) (Fig S52). These structures closely resemble the aggregations of hollow ICVs released after the lysis of *EM-P* cells (Fig. 2A-2E & S22). As observed in *EM-P*, the distribution of organic carbon in the North-Pole Formation microfossils is restricted to the periphery of the spherical cells (Fig. S52). Along with the morphological and organizational similarities, *EM-P* also exhibited all the accessory structures associated with the North Pole formation microfossils, such as the large clots of organic carbon (Fig. S51 arrows), filamentous structures originating from the spherical cells, and spherical clots of organic carbon within these filamentous (Fig. S52-S55)(45). Based on the similarities, we propose that the large clots of organic carbon would have been the membrane debris formed during the lysis and the release of intracellular vesicles (Fig. S51, Movie 5-9). As observed in *EM-P*, the filamentous structures associated with the microfossils could also have formed during the release of intracellular vesicles (Movie 5-9). The organic carbon clots within the filamentous structures could have been fossilized daughter cells (Fig. S53 & S54, arrows).

Apart from the aggregations of hollow spherical cells, honeycomb-like structures were also reported from several microfossil sites, like the SPF, the Nuga Formation, the Buck Reef Chert, the Moodies Group, and the Turee Creek formations (Fig. 6, S57-S62)(40,54–56). As observed in *EM-P*, these structures could have been formed by the lateral compression of cells or hollow vesicles within the biofilm (Fig. 6). In tune with our proposition, Archaean honeycomb-like structures are often closely associated with spherical *EM-P*-like cells (Fig. 6).

Fig. 5. Morphological comparison of the Mt. Goldsworthy and the Sulphur Spring

microfossils with *EM-P*. Image A-C are microfossils reported from the Mt. Goldsworthy

Formation (37)(Sugitani *et al.,* 2007). Image D is the 3D-rendered STED microscope images

of morphologically analogous membrane debris of *EM-P* cell with attached daughter cells

(highlighted region) (also see Fig. S40). Images E & F are microfossils reported from the

Sulphur Spring site (57)(Duck *et al.,* 2007), showing spherical structures attached to

membrane debris. Images G & J are the morphologically analogous structures observed in

EM-P incubations. Images H & I show the magnified regions of G & J showing spherical

EM-P daughter cells attached to membrane debris (also see Fig. S40-S47, Movie 17). Cells

and membrane debris in these images were stained with the membrane stain FM^{TM}5-95

(yellow). Scale bars: A (50μm), G & J (20μm).

Fig 6: Sequential steps involved in the formation of honeycomb-shaped mats:

Images A-C show single *EM-P* cells that gradually transformed from spherical cells with

intracellular vesicles into honeycomb-like structures. Images D-E show a similar

transformation of biofilms composed of individual spherical cells into honeycomb-like

structures. Cells in these images are stained with membrane stain, FM[TM]5-95 (red), and

imaged using a STED microscope. Images G-J are the microfossils reported from the SPF

(originally published by Sugitani et al., 2007)(37). Scale bars: A-F (10µm), G & H (20µm),

and I (50µm).

Spherical microfossils from the Pilbara and Barberton Greenstone Belts were often

discovered within layers of organic carbon (46,49,58,59). Over a period of 2-6 months, we

observed cells in our incubations gradually being enclosed by membrane debris. These

structures were formed by a multi-step process (Fig S63). First, *EM-P* grew as multiple layers

of cells within a biofilm (Fig. S63A). Second, the lysis of these cells led to the formation of a

considerable amount of membrane debris (Fig. S63B, S64 & Movie 17&18). Subsequently,

this membrane debris coalesced to form large fabric-like structures (Fig. S65). These

membrane fabrics were then expelled from the biofilm (Fig. S63D, S63E & Movie 18). Over

time, these expelled membrane fabrics grew in surface area to form a continuous layer of

membrane enclosing a large population of cells (Fig. S65-S69 & Movie 19). This fabric-like

membrane debris enclosing biofilms observed in *EM-P* incubations bear close morphological

resemblance to microfossils reported from Chinaman Creek in the Pilbara (Fig. S69), and Mt.

Goldsworthy Formation (Fig. 4 & 5)(37,41,60).

Parallel layers of organic carbon termed laminations were reported from several Archaean

microfossil sites (33,49,58,61). Structures similar to these laminations were observed in our

incubations. As described above, the reproduction in *EM-P* involves the lysis of cells to

facilitate the release of the intracellular daughter cells, resulting in a considerable amount of

cell debris (Movie 9 & 17). The parallel layers of organic carbon in our incubations (Fig. 7 &

8) are formed by lysis and collapse within individual biofilm layers (Fig. S49). Another way

the organic carbon layers could have formed is by the lateral compression of honeycomb-like

biofilms (Fig. S71-S74). Sequential steps resulting in the formation of such structures are

shown in Fig. S72. Such layers of cell debris closely resemble different types of laminated

structures reported from the Barberton Greenstone Belt and Pilbara Iron Formations, like the

α, and β laminations (Fig. 7, 8 & S69-S75)(49,58,59).

Also similar to the Archaean laminations, we observed layers of cell debris in our incubations

have lenticular gaps (Fig. 8, S69, S70-S71 & S76-S79). Within these lenticular gaps, we

observed intact *EM-P* cells or honeycomb patterns, suggesting that lenticular gaps within

otherwise uniformly parallel laminations were formed due to non-uniform lysis or incomplete

deflation of cells within individual layers of *EM-P* cells (Fig. 7, S75-S79 & Movies 20 & 21).

Although in the case of Archaean laminations, these lenticular gaps were thought to have

been formed by the entrapment of air bubbles (49), based on our results, we argue that there

could have been more than one way such structures could have formed. Other distinctive features of the lamination, like raised mounds or swirls (Fig. S81 - S83)(49,59), were also observed in batch cultures of *EM-P*. Given these morphological similarities, we propose that some of the laminated and other diaphanous filamentous structures could have been formed by the cell debris of the *EM-P*-like cells that inhabited these sites during the Archaean Eon. We will discuss these possibilities in more detail below.

Over a period of 3-12 months, we observed the biofilms solidifying into a solid crust (Fig. S84 & S85). The SEM-EDX characterization of these solidified biofilms showed the presence of potassium and magnesium minerals on the surface, suggesting that these structures were formed by the gradual adsorption of positively charged cations on the negatively charged biofilms (Fig. S84). Most Archaean microfossils are restricted to coastal marine environments. Compared to open oceans, these coastal marine environments harbor higher concentrations of salt due to higher evaporation rates. Hence, these microfossils could have undergone a similar encrustation process as observed in our incubations. Moreover, solidified *EM-P* biofilms resemble the mineral-encrusted structures reported from the Kromberg Formation (Fig. S85)(62). Like the Kromberg Formations structures, solidified *EM-P* biofilms are composed of desiccation cracks and spherical cells beneath the surface (Fig. S85).

When these salt-encrusted cells were transferred into fresh media, we observed a gradual increase in cell numbers (results not shown). However, given that these cells are encrusted in a layer of salts, we observed early growth-phase cells breaking out of the thick salt crust, resulting in stellar morphologies (Fig. S86 & S87). The observed morphologies of these cells closely resemble the morphologies of microfossils reported from the Strelley Pool and other North Pole cherts (Fig. S86 & S87)(39,45). All distinctive features of these microfossils, like the stellar-shaped cells undergoing binary fission and a string of daughter cells extending out of such stellar cells (Movie 22), were also observed in our incubations.

1
2
Fig. 7: Morphological comparison of the Buch Reef Chert β-laminations with *EM-P's*

membrane debris. Image A shows a 3D-rendered image of *EM-P's* membrane debris. Cells

in the image are stained with membrane stain Nile red and imaged using a STED microscope.

Images B & C show β-type laminations reported from Buch Reef Chert (originally published

by Tice *et al.,* 2009)(59). The boxed region in image-a highlights the membrane-forming

rolled-up structures containing spherical daughter cells, as described in the case of BRC

organic structures. Scale bars: 50μm.

10
11
12
13

1
2
Fig. 8: Morphological comparison between laminated structures reported from the

Moodies Group and structures formed by *EM-P*. Images A & B are laminated structures

reported from the Moodies Group (originally published by Homann *et al.,* 2015)(61). They

show filamentous structures with lenticular gaps. These gaps were filled with quartz granules.

Image C is a 3D-rendered confocal image of *EM-P*'s membrane debris. Filamentous

membrane debris bifurcating, forming spherical\lenticular gaps, can be seen in several

regions. Some spherical/lenticular gaps were hollow (box), and some had a honeycomb

pattern within them (arrow), indicating the presence of large spherical *EM-P* cells with

intracellular vesicles (D). 3D rendered STED microscope images of individual

spherical/lenticular gaps with and without *EM-P* cell debris were shown in Fig. S75 & S76,

and S77. Membranes were stained with Nile red, and imaging was done using a STED

microscope. The scales: 50µm.

Discussion:

Advances in microscopic (FIB-SEM) and analytical (NanoSIMS) techniques over the past

few decades have facilitated better imaging and precise determination of chemical and

isotopic compositions of microfossils (13,63). Nevertheless, there is considerable

disagreement among researchers regarding the interpretation of this information (9). Given

the importance of morphology in determining the biogenicity and taxonomic affiliation of the

microfossils, reconstructing the lifecycles of Archaean Eon organisms is considered crucial to

understanding the nature of these microfossils (41). Our study is the first to reconstruct all

known microfossil morphologies and their lifecycles from extant bacteria. Furthermore, we

have shown that many of the taphonomic structures observed in our study closely resemble

the controversial structures observed in rocks of the Palaeo-Mesoarchaean age (3.6-3.0 Ga)

and even in the Neoarchaean (3.0-2.4 Ga). These similarities help us answer long-standing

questions regarding the origin and the nature of Archaean microfossils.

The nature of Archaean organic structures is currently being debated among researchers

(7,9,10). While some studies suggest that these structures could be remnants of Archaean

microorganisms, others suggest that they may have been abiotic minerals that formed due to

volcanic activity (10). The argument for this proposition is based on the fact that these

organic structures share more similarities with inorganic mineral structures than with extant

prokaryotes. To establish the biogenicity of a microfossil, it is essential to either find a

convincing morphological analog among extant bacteria or establish a biogenic process

through which they are formed (3). The biogenicity of microfossils reported from several

sites like the Swartkoppoie Formation, Kitty's Gap Chert, and the Josephdal Formation is

widely accepted among the scientific community due to the discovery of spherical

microfossils in different stages of their lifecycle (Fig. S32)(12,53,62). However, such a step-

by-step biological process through which Archean Eon organic structures could have formed

has never been demonstrated empirically.

The biological origin of microfossils reported from several sites, like the Dresser Formation,

to date, remains a matter of debate (10). Several morphological features of these organic

structures, like the presence of organic carbon only at the periphery, the absence of internal

cell constituents, the presence of pyrite and silicate minerals inside the cells, and the presence

of a thick porous or discontinuous cell wall, were all argued as claims for their abiotic origin

(10). Justifiably, these morphological features have never been observed in any living

organism. Nevertheless, all spherical microfossils reported from the Dresser Formation

resemble *EM-P* cells, especially those with a single large ICV (Fig. S24-S27). What was

thought to have been a thick, porous cell wall in the microfossils could have been the

cytoplasm with tiny ICVs sandwiched between the cell and vesicle membrane (Fig. S24 &

S27). Similarly, the hollow cells with a discontinuous cell wall could either have been the

ICVs released by cell lysis or the late-growth stage cells with little cytoplasm (Fig. S26). In

such cells, the presence of cytoplasm is restricted to discontinuous patches around the

periphery of the cells (Fig. S26). The sequence of steps leading to these cell morphologies

that resemble the Dresser Formation microfossils is shown in Fig. S27. A closer inspection of

the Dresser Formation microfossils shows the ICVs membrane rupture and the daughter cell

release (fig. S28). Morphologies indicating this method of reproduction among microfossils

is not unique to the Dresser Formation. Microfossils with similar morphological features

were reported from sites like the Strelley Pool, the Waterfall region, and Mt. Goldsworthy

Formation (Fig. S3-S14)(3,41). These similarities suggest that microfossil morphologies

observed in the Dresser Formation are in tune with other microfossils of similar geological

time periods, suggesting their biological origin.

Spherical structures half-coated with pyrite are reported from the Dresser Formation

(Fig.S25)(10). These structures could have been the iron-reducing *EM-P*-like cells with

hollow ICV constituting half their volume. The selective co-localization of pyrite and carbon

could be explained by the Fe(III) reduction happening at the cell surface. The Fe(II) produced

from this metabolic reaction could have reacted with environmental sulfide, converted to

insoluble pyrite, and precipitated onto the cell surface. Given the absence of this metabolic

process within the hollow ICV, these structures remained pyrite-free (Fig.S25). In addition to

the Dresser Formation, organic structures coated with pyrite have also been reported from

other microfossil sites like the Sulphur Spring Formations (17). The selective presence of

pyrite on these microfossils could also be explained by a similar mechanism (Fig. S36-S38).

Apart from pyrite, minerals like anatase were reported to have been present within the cells

(10). The presence of anatase within these cells could be explained by the transport of these

minerals into the cells during the ICV formation (Fig. 1B&D). ICVs are formed by a process

similar to endocytosis, which involves the intake of salt-rich media and minerals into the cells

(Fig. S88). Like the Dresser Formation microfossils, we often observed the presence of salts

and minerals within *EM-P*s vesicles (Fig. S88). Moreover, the presence of minerals within

cells is not unique to the Dresser Formation microfossils (10) and was reported previously

from several bonafide microfossils from Gunflint Iron Formations (64). Additionally, we

observed remarkable similarities between the *EM-P* cell debris and all the organic structures

closely associated with the microfossil, such as the wavy lamellar and pumice-like structures

(Fig. S59, S61 & S89). The step-by-step transformation of cell debris into pumice-like

structures is shown in Fig. 6. Based on these morphological similarities between the Dresser

Formation organic structures and *EM-P*, we hypothesize that these organic structures are the

fossil remnants of *EM-P*-like bacteria rather than mineral aggregates.

Morphological similarity between microfossils from far-flung sites like Western Australia

and Southern Africa could be explained by the similarity in the environmental conditions in

both sites (5). This relationship between cell morphology, reproductive processes, and

environmental conditions was discussed extensively in our previous work (27,65). The

experimental conditions that we employed in our study are likely similar to the environmental

conditions faced by Archaean organisms from both these sites at the time of their

fossilization. All sites from which microfossils were reported are shallow intertidal regions.

Evidence for periodic evaporation and flooding with sea water was presented from the

Barberton and Pilbara Greenstone Belts (6,66), suggesting that the original microorganisms

experienced high salinities. The salinities of our experiments are broadly similar to those of

Archaean oceans (5-10% w/v)(31). To our knowledge, the exact salt composition of the

Archaean Ocean has not been elucidated. Hence, we used a complex mixture of salts (DSS)

as a proxy to reproduce these salinities in our experiments. Salts like Mg^{+2}, Ca^{+2}, $Na^{+,}$ and K^+

or their oxides were also reported to be present and constitute 1-5% by weight in both Pilbara

and Barberton greenstone belt microfossil sites (6,66). Moreover, these salts were shown to

be closely associated with microfossils (66). The spatial distribution of these salts resembles

the spatial distribution pattern of organic carbon, possibly indicating the chelation of these

salts to the cell membrane, which is also in agreement with our observations (9). The presence of potassium phyllosilicates and NaCl crystals within the microfossils (66) is also in agreement with our hypothesis that internal structures of the microfossils should have formed by invagination of cell-membrane taking in salt-rich water (Fig. S86). As observed in the microfossils (66), salt crystals on the cell surface, within the membrane invaginations, or cell debris were often observed in *EM-P* (Fig. S89).

The above-presented results suggest that Archaean Eon cells are likely primitive lipid-vesicle-like protocells that lack a cell wall. From a physiological perspective, it would have been unlikely for primitive cells to possess a cell wall given the substantial number of genes required to synthesize individual building blocks, to mediate its assembly, and its constant modification to facilitate cell growth and reproduction (21,67,68). Furthermore, a cell wall could impede the transport of physiologically relevant compounds in and out of the cells. To overcome this limitation, present-day microorganisms (with a cell wall) had to develop extensive molecular biological processes for transporting nutrients and metabolic end products across the cell wall (69,70). This could not have been the case for primitive Archaean life forms. Hence, rather than drawing parallels between the microfossils and life as we know it today, we propose that these microfossils could have been liposome-like protocells, as proposed by the theory of chemical evolution (26). Indeed, it has been recently shown that liposome-like molecules could be produced in some of the hydrothermal settings proposed for the emergence of life (71). To the best of our knowledge, this is the first study to provide a link between theoretical propositions and geological evidence for the existence of protocells on early Earth.

According to the theory of chemical evolution, biological organic compounds are formed by abiotic processes (72). These compounds then self-assembled to form lipid vesicles, which grew in complexity and eventually evolved into self-replicating protocells (73,74). These protocells are believed to have undergone Darwinian evolution, resulting in the emergence of bacteria, archaea, and eukaryotes (75). It was previously thought that the fragility of protocells made it unlikely for them to be preserved in rock formations. However, later studies showed the preservation of cellular features by a rapid encrustation of cells with cationic minerals (50,51). The rapid encrustation and preservation of cells observed in our study (Fig. S84-S85) is in accordance with the proposition that environmental conditions influence the extent of cellular preservation. Our study aligns with the interpretations from

these studies that environmental conditions play a pivotal role in determining the extent of

cellular preservation.

Conclusion

For the first time, our investigations have been able to reproduce morphologies of most

Archaean microfossils from wall-less, extant cells. Apart from reproducing the morphologies,

we also presented a step-by-step biological process by which Archaean organic structures

could have formed. Based on these results, we propose that Archaean microfossils were

likely liposome-like cells, which had evolved mechanisms for energy conservation but not for

regulating cell morphology and replication. In an earlier study, we have shown that the

morphologies of such primitive cells are determined by environmental conditions (27,65)

rather than the information encoded in their genome. Given this lack of intrinsic ability to

regulate their morphology, we argue that morphological features such as cell size, shape, or

cytological complexity are reliable factors in interpreting either the phylogeny or the

physiology of microfossils (at least from Archaean Eon). Rather than attempting to assign

present-day taxonomies to these microfossils, we suggest that these microfossils represent

primitive protocells proposed by the theory of chemical evolution. To the best of our

knowledge, ours is the first study to provide paleontological evidence of the possible

existence of protocells on the Palaeoarchaean Earth.

Methods:

Isolation of cells and their transformation to protoplasts:

Exiguobacterium strain-Molly (*EM*) was isolated from submerged freshwater springs within

the Dead Sea (76). The taxonomic identification of the isolate to the genus *Exiguobacterium*

was determined by 16S rRNA gene sequencing (77,78). *EM* cells were transformed into

protoplasts following a previously documented protocol(28). The resulting *EM-P* cells were

cultured in half-strength TSB with 7% Dead Sea Salt (7%DSS-TSB).

Microscopic observation of *EM-P* cells:

Morphology *EM-P* was routinely assessed using an Axioskop 2plus microscope (Carl Zeiss, Germany) with a Plan-NEOFLUAR 100X/1.3 objective. Images were captured using a Leica DSF9000 camera (Leica Microsystems, Mannheim, Germany). STED microscopy was performed with an inverted TCS SP8 STED 3X microscope (Leica Microsystems, Mannheim Germany) using an 86x/1.2 NA water immersion objective (Leica HC PL APO CS2 - STED White). Fluorophores were excited with 488, 561nm, 594nm, or 633m laser light derived from an 80 MHz pulsed White Light Laser (Leica Microsystems, Mannheim Germany). For stimulated emission, either a pulsed 775 nm laser or a 592nm CW laser (Leica Microsystems, Mannheim, Germany) was used depending on the fluorophore. Photon counting mode and line accumulation were used for image recording, and Huygens Professional (SVI, Hilversum, The Netherlands) performed image deconvolution on selected images and movies.

Spinning Disk Microscopy was performed using an Olympus SpinSR10 spinning disk confocal microscope (Olympus, Tokyo, Japan) equipped with a 100x/NA1.35 silicone oil immersion objective (Olympus UPLSAPO100XS, Tokyo, Japan), a CSU-W1-Spinning Disk-Unit (Yokogawa, Tokyo, Japan) and ORCA☐Flash 4.0 V3 Digital CMOS Camera (Hamamatsu, Hamamatsu City, Japan).

Transmission electron microscopy was conducted utilizing a Zeiss EM 912 (Zeiss, Oberkochen, Germany) equipped with an integrated OMEGA filter, operating at 80 kilovolts (kV). Image acquisition was carried out using a 2k x 2k pixel slow-scan CCD camera (TRS, TrÖndle Restlichtverstrkersysteme, Moorenweis, Germany) with ImageSP software (SysProg, Minsk, Belarus).

Acknowledgments

We want to thank Gabriella Berthal for her excellent technical support and Christian Sibert for providing the Dead Sea samples from which *EM* was isolated. We thank the Advanced Light Microscopy Facility at EMBL, Heidelberg, Ulf Schwartz from Leica Microsystems, and colleagues at the departments of Ecological Microbiology (Bayreuth University) and of Cellular and Molecular Biophysics (Max Planck Institute for Biochemistry) for their support throughout the work.

Competing Interests

The authors declare that there are no conflicts of interest regarding the publication of this article.

4
5
6

Data Availability Statement

Dr. Dheeraj Kanaparthi will share all data, materials, and methods upon reasonable request.

Funding

This research was funded by the European Research Council (ERC) grant agreement 616644 (POLLOX) and by the Deutsche Forschungsgemeinschaft (DFG) grant agreements DFG-TRR174 and Seed funding from Excellence Cluster ORIGINS EXC2094 – 390783311.

12
13
14
15

References:

1. Shields GA. Chapter 7.6 The Marine Carbonate and Chert Isotope Records and Their Implications for Tectonics, Life and Climate on the Early Earth. Dev Precambrian Geol. 2007;15.

2. van Zuilen MA. Chapter 38 - The Significance of Carbonaceous Matter to Understanding Life Processes on Early Earth. In: Van Kranendonk MJ, Bennett VC, Hoffmann JEBTEOR (Second E, editors. Elsevier; 2019. p. 945–63. Available from: https://www.sciencedirect.com/science/article/pii/B9780444639011000381

3. Sugitani K, Mimura K, Takeuchi M, Lepot K, Ito S, Javaux EJ. Early evolution of large micro-organisms with cytological complexity revealed by microanalyses of 3.4 Ga organic-walled microfossils. Geobiology. 2015;13(6):507–21.

4. Hickman-Lewis K, Westall F. A southern African perspective on the co-evolution of early life and environments. South Afr J Geol. 2021;124(1).

5. Oehler DZ, Walsh MM, Sugitani K, Liu MC, House CH. Large and robust lenticular microorganisms on the young Earth. Precambrian Res. 2017;296:112–9.

6. Walsh MW. Microfossils and possible microfossils from the early archean onverwacht group, barberton mountain land, South Africa. Precambrian Res. 1992;54(2–4):271–93.

7. Schopf JW, Kitajima K, Spicuzza MJ, Kudryavtsev AB, Valley JW. SIMS analyses of the oldest known assemblage of microfossils document their taxon-correlated carbon isotope compositions. Proc Natl Acad Sci. 2018;115(1):53–8.

8. Schopf JW, Kudryavtsev AB. Biogenicity of Earth's earliest fossils: a resolution of the controversy. Gondwana Res. 2012;22(3):761–71.

9. Wacey D, Saunders M, Kong C, Brasier A, Brasier M. 3.46 Ga Apex chert 'microfossils' reinterpreted as mineral artefacts produced during phyllosilicate exfoliation. Gondwana Res. 2016;36:296–313.

10. Wacey D, Noffke N, Saunders M, Guagliardo P, Pyle DM. Volcanogenic Pseudo-Fossils from the ~3.48 Ga Dresser Formation, Pilbara, Western Australia. Astrobiology. 2018;

11. Gee H. That's life? Nature. 2002;416(6876):28.

12. Knoll AH, Barghoorn ES. Archean microfossils showing cell division from the Swaziland System of South Africa. Science. 1977;198(4315):396–8.

13. Lepot K, Williford KH, Ushikubo T, Sugitani K, Mimura K, Spicuzza MJ, et al. Texture-specific isotopic compositions in 3.4 Gyr old organic matter support selective preservation in cell-like structures. Geochim Cosmochim Acta. 2013;112:66–86.

14. Adami C, Ofria C, Collier TC. Evolution of biological complexity. Proc Natl Acad Sci. 2000;97(9):4463–8.

15. Wolf YI, Katsnelson MI, Koonin EV. Physical foundations of biological complexity. Proc Natl Acad Sci U S A. 2018;115(37).

16. Malaterre C, ten Kate IL, Baqué M, Debaille V, Grenfell JL, Javaux EJ, et al. Is There Such a Thing as a Biosignature? Astrobiology. 2023 Nov 1;23(11):1213–27.

17. Wacey D, Saunders M, Cliff J, Kilburn MR, Kong C, Barley ME, et al. Geochemistry and nano-structure of a putative~ 3240 million-year-old black smoker biota, Sulphur Springs Group, Western Australia. Precambrian Res. 2014;249:1–12.

18. Hickman-Lewis K, Garwood RJ, Brasier MD, Goral T, Jiang H, McLoughlin N, et al. Carbonaceous microstructures from sedimentary laminated chert within the 3.46 Ga Apex Basalt, Chinaman Creek locality, Pilbara, Western Australia. Precambrian Res. 2016;

19. Schopf JW. Microfossils of the Early Archean Apex chert: new evidence of the antiquity of life. Science. 1993;260(5108):640–6.

20. Wacey D, Saunders M, Kong C. Remarkably preserved tephra from the 3430 Ma Strelley Pool Formation, Western Australia: Implications for the interpretation of Precambrian microfossils. Earth Planet Sci Lett. 2018;

21. Adams DW, Errington J. Bacterial cell division: assembly, maintenance and disassembly of the Z ring. Nat Rev Microbiol. 2009;7(9):642.

22. Westall F, Hickman-Lewis K, Hinman N, Gautret P, Campbell KA, Bréhéret JG, et al. A Hydrothermal-Sedimentary Context for the Origin of Life. Astrobiology. 2018;18(3).

23. Szostak JW. The narrow road to the deep past: in search of the chemistry of the origin of life. Angew Chem Int Ed. 2017;56(37):11037–43.

24. Lane N, Martin W. The energetics of genome complexity. Nature. 2010;

25. Lane N. Bioenergetic constraints on the evolution of complex life. Cold Spring Harb Perspect Biol. 2014;

26. Oparin AI. Chemistry and the origin of life. R Inst Chem Rev. 1969;

27. Kanaparthi D, Lampe M, Hildebrand F, Boesen T, Klingl A, Phapale P, et al. On the reproductive mechanisms of Gram-positive protocells. bioRxiv. 2021 Jan 1;2021.11.25.470039.

28. Kanaparthi D, Lampe M, Krohn JH, Zhu B, Klingl A, Lueders T. The reproduction of gram-negative protoplasts and the influence of environmental conditions on this process. iScience [Internet]. 2023 Nov 17;26(11). Available from: https://doi.org/10.1016/j.isci.2023.108149

29. Errington J. L-form bacteria, cell walls and the origins of life. Open Biol. 2013;3(1):120143.

30. Knauth LP. Temperature and salinity history of the Precambrian ocean: implications for the course of microbial evolution. Palaeogeogr Palaeoclimatol Palaeoecol. 2005;219(1):53–69.

31. Knauth LP. Salinity history of the Earth's early ocean. Nature. 1998;395(6702):554.

32. Catling DC, Zahnle KJ. The Archean atmosphere. Sci Adv. 2020;6(9).

33. Hickman-Lewis K, Westall F, Cavalazzi B. Traces of Early Life From the Barberton Greenstone Belt, South Africa. In: Earth's Oldest Rocks. 2019.

34. Schopf JW, Packer BM. Early Archean (3.3-billion to 3.5-billion-year-old) microfossils from Warrawoona Group, Australia. Science. 1987;237(4810):70–3.

35. Delarue F, Robert F, Derenne S, Tartèse R, Jauvion C, Bernard S, et al. Out of rock: a new look at the morphological and geochemical preservation of microfossils from the 3.46 Gyr-old Strelley Pool Formation. Precambrian Res. 2019;

36. Retallack GJ, Krinsley DH, Fischer R, Razink JJ, Langworthy KA. Archean coastal-plain paleosols and life on land. Gondwana Res. 2016;

37. Sugitani K, Grey K, Allwood A, Nagaoka T, Mimura K, Minami M, et al. Diverse microstructures from Archaean chert from the Mount Goldsworthy–Mount Grant area, Pilbara Craton, Western Australia: microfossils, dubiofossils, or pseudofossils? Precambrian Res. 2007;158(3–4):228–62.

38. Schopf JW, Barghoorn ES. Alga-like fossils from the early Precambrian of South Africa. Science. 1967;156(3774):508–12.

39. Sugitani K, Mimura K, Nagaoka T, Lepot K, Takeuchi M. Microfossil assemblage from the 3400 Ma Strelley Pool Formation in the Pilbara Craton, Western Australia: results form a new locality. Precambrian Res. 2013;226:59–74.

40. Barlow EV, Van Kranendonk MJ. Snapshot of an early Paleoproterozoic ecosystem: Two diverse microfossil communities from the Turee Creek Group, Western Australia. Geobiology. 2018;

41. Sugitani K, Grey K, Nagaoka T, Mimura K, Walter MR. Taxonomy and biogenicity of Archaean spheroidal microfossils (ca. 3.0 Ga) from the Mount Goldsworthy–Mount Grant area in the northeastern Pilbara Craton, Western Australia. Precambrian Res. 2009;173(1–4):50–9.

42. Sugitani K, Mimura K, Takeuchi M, Yamaguchi T, Suzuki K, Senda R, et al. A Paleoarchean coastal hydrothermal field inhabited by diverse microbial communities: the Strelley Pool Formation, Pilbara Craton, Western Australia. Geobiology. 2015;13(6):522–45.

43. Wacey D, Kilburn MR, Saunders M, Cliff J, Brasier MD. Microfossils of sulphur-metabolizing cells in 3.4-billion-year-old rocks of Western Australia. Nat Geosci. 2011;4(10):698–702.

44. Kaźmierczak J, Kremer B. Pattern of cell division in~ 3.4 Ga-old microbes from South Africa. Precambrian Res. 2019;105357.

45. Buick R. Microfossil Recognition in Archean Rocks: An Appraisal of Spheroids and Filaments from a 3500 M.Y. Old Chert-Barite Unit at North Pole, Western Australia. PALAIOS. 1990;

46. Homann M. Earliest life on Earth: Evidence from the Barberton Greenstone Belt, South Africa. Earth-Sci Rev. 2019;196:102888.

47. Ueno Y, Isozaki Y, McNamara KJ. Coccoid-like microstructures in a 3.0 Ga chert from Western Australia. Int Geol Rev. 2006;48(1):78–88.

48. Kazmierczak J, Altermann W, Kremer B, Kempe S, Eriksson PG. Mass occurrence of benthic coccoid cyanobacteria and their role in the production of Neoarchean carbonates of South Africa. Precambrian Res. 2009;173(1–4):79–92.

49. Homann M, Sansjofre P, Van Zuilen M, Heubeck C, Gong J, Killingsworth B, et al. Microbial life and biogeochemical cycling on land 3,220 million years ago. Nat Geosci. 2018;

50. Orange F, Westall F, Disnar JR, Prieur D, Bienvenu N, Le Romancer M, et al. Experimental silicification of the extremophilic archaea pyrococcus abyssi and methanocaldococcus jannaschii: Applications in the search for evidence of life in early earth and extraterrestrial rocks. Geobiology. 2009;7(4).

51. Orange F, Disnar JR, Westall F, Prieur D, Baillif P. Metal cation binding by the hyperthermophilic microorganism, Archaea Methanocaldococcus Jannaschii, and its effects on silicification. Palaeontology. 2011;54(5).

52. Gödeke J, Paul K, Lassak J, Thormann KM. Phage-induced lysis enhances biofilm formation in Shewanella oneidensis MR-1. ISME J. 2011;5(4).

53. Westall F, De Vries ST, Nijman W, Rouchon V, Orberger B, Pearson V, et al. The 3.466 Ga "Kitty's Gap Chert," an early Archean microbial ecosystem. Spec Pap Geol Soc Am. 2006;

54. GAMPER A, HEUBECK C, DEMSKE D, HOEHSE M. Composition and Microfacies of Archean Microbial Mats (Moodies Group, ca. 3.22 Ga, South Africa). In: Microbial Mats in Siliclastic Depositional Systems Through Time. 2012.

55. Schopf JW, Kudryavtsev AB, Osterhout JT, Williford KH, Kitajima K, Valley JW, et al. An anaerobic ~3400 Ma shallow-water microbial consortium: Presumptive evidence of Earth's Paleoarchean anoxic atmosphere. Precambrian Res. 2017;

56. Tice MM, Lowe DR. The origin of carbonaceous matter in pre-3.0 Ga greenstone terrains: A review and new evidence from the 3.42 Ga Buck Reef Chert. Earth-Sci Rev. 2006;

57. Duck LJ, Glikson M, Golding SD, Webb RE. Microbial remains and other carbonaceous forms from the 3.24 Ga Sulphur Springs black smoker deposit, Western Australia. Precambrian Res. 2007;154(3–4):205–20.

58. Tice MM, Lowe DR. Photosynthetic microbial mats in the 3,416-Myr-old ocean. Nature. 2004;

59. Tice MM. Environmental controls on photosynthetic microbial mat distribution and morphogenesis on a 3.42 Ga clastic-starved platform. Astrobiology. 2009;

60. Brasier MD, Green OR, Lindsay JF, McLoughlin N, Steele A, Stoakes C. Critical testing of Earth's oldest putative fossil assemblage from the ~3.5 Ga Apex chert, Chinaman Creek, Western Australia. Precambrian Res. 2005;

61. Homann M, Heubeck C, Airo A, Tice MM. Morphological adaptations of 3.22 Ga-old tufted microbial mats to Archean coastal habitats (Moodies Group, Barberton Greenstone Belt, South Africa). Precambrian Res. 2015;

62. Westall F, de Wit MJ, Dann J, van der Gaast S, de Ronde CEJ, Gerneke D. Early Archean fossil bacteria and biofilms in hydrothermally-influenced sediments from the Barberton greenstone belt, South Africa. Precambrian Res. 2001;106(1–2):93–116.

63. Brasier MD, Antcliffe J, Saunders M, Wacey D. Changing the picture of Earth's earliest fossils (3.5–1.9 Ga) with new approaches and new discoveries. Proc Natl Acad Sci. 2015;112(16):4859–64.

64. Lepot K, Addad A, Knoll AH, Wang J, Troadec D, Béché A, et al. Iron minerals within specific microfossil morphospecies of the 1.88 Ga Gunflint Formation. Nat Commun. 2017;8:14890.

65. Kanaparthi D, Lampe M, Zhu B, Klingl A, Lueders T, Schwille P. On the reproductive mechanism of Gram-negative protocells. bioRxiv. 2021 Jan 1;2021.11.25.470037.

66. Alleon J, Bernard S, Le Guillou C, Beyssac O, Sugitani K, Robert F. Chemical nature of the 3.4 Ga Strelley Pool microfossils. Geochem Perspect Lett. 2018;7:37–42.

67. Vollmer W, Joris B, Charlier P, Foster S. Bacterial peptidoglycan (murein) hydrolases. FEMS Microbiol Rev. 2008;32(2).

68. Egan AJF, Cleverley RM, Peters K, Lewis RJ, Vollmer W. Regulation of bacterial cell wall growth. FEBS J. 2017;

69. Dijkstra AJ, Keck W. Peptidoglycan as a barrier to transenvelope transport. J Bacteriol. 1996;178(19).

70. Prajapati JD, Kleinekathöfer U, Winterhalter M. How to Enter a Bacterium: Bacterial Porins and the Permeation of Antibiotics. Chem Rev. 2021;121(9).

71. Purvis G, Šiller L, Crosskey A, Vincent J, Wills C, Sheriff J, et al. Generation of long-chain fatty acids by hydrogen-driven bicarbonate reduction in ancient alkaline hydrothermal vents. Commun Earth Environ. 2024;5(1):30.

72. Powner MW, Gerland B, Sutherland JD. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. Nature. 2009;459(7244).

73. Zhu TF, Szostak JW. Coupled growth and division of model protocell membranes. J Am Chem Soc. 2009;131(15):5705.

74. Schwille P. Division in synthetic cells. Emerg Top Life Sci. 2019;

75. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: Proposal for the domains Archaea, Bacteria, and Eucarya. Proc Natl Acad Sci U S A. 1990;87(12).

76. Häusler S, Noriega-Ortega BE, Polerecky L, Meyer V, de Beer D, Ionescu D. Microenvironments of reduced salinity harbour biofilms in Dead Sea underwater springs. Environ Microbiol Rep. 2014;

77. Kanaparthi D, Reim A, Martinson GO, Pommerenke B, Conrad R. Methane emission from feather moss stands. Glob Change Biol. 2017;23(11).

78. Kanaparthi D, Conrad R. Role of humic substances in promoting autotrophic growth in nitrate-dependent iron-oxidizing bacteria. Syst Appl Microbiol. 2015;38(3):184–8.

79. Duck LJ, Glikson M, Golding SD, Webb RE. Microbial remains and other carbonaceous forms from the 3.24 Ga Sulphur Springs black smoker deposit, Western Australia. Precambrian Res. 2007;154(3–4):205–20.

80. Köhler I, Heubeck C. Microbial-mat-associated tephra of the Archean Moodies Group,
Barberton Greenstone Belt (BGB), South Africa: Resemblance to potential biostructures
and ecological implications. South Afr J Geol. 2019;

81. Sugitani K, Lepot K, Nagaoka T, Mimura K, Van Kranendonk M, Oehler DZ, et al.
Biogenicity of morphologically diverse carbonaceous microstructures from the ca. 3400
Ma strelley pool formation, in the pilbara craton, Western Australia. Astrobiology.
2010;10(9).