# Title: Functional Connectivity of the World's Protected Areas

**Authors:** A. Brennan*[1,2,3], R. Naidoo[2,3], L. Greenstreet[2], Z. Mehrabi[2,4], N. Ramankutty[2,4] and C. Kremen[1,2,5]

**Affiliations:**

[1]Biodiversity Research Centre, University of British Columbia; Vancouver, Canada.

[2]Institute for Resources, Environment and Sustainability, University of British Columbia; Vancouver, Canada.

[3]World Wildlife Fund; Washington, DC, USA.

[4]School of Public Policy and Global Affairs, University of British Columbia; Vancouver, Canada.

[5]Department of Zoology, University of British Columbia; Vancouver, Canada.

*Corresponding author. Email: angela.brennan@ubc.ca

**Abstract**

Rapid environmental change threatens to isolate the world's wildlife populations and intensify biodiversity loss. Global policies have called for expanding and connecting the world's protected areas (PAs) to curtail the crisis, yet how well PA networks currently support wildlife movement, and where connectivity conservation or restoration is most critical, have never been mapped globally. Here, we map the functional connectivity (how animals move through landscapes) of the world's terrestrial PAs for the first time. Also, going beyond existing global connectivity indices, we quantify national PA-connectedness using an approach that

meaningfully represents animal movement through anthropogenic landscapes. We find that

reducing the human footprint may improve national PA-connectivity more than adding new PAs;

however, both strategies are critical for improving and preserving connectivity in places where

the predicted flow of animal movement is highly concentrated. We show that the majority of

critical connectivity areas (CCAs) (defined as globally important areas of concentrated animal

movements) remain unprotected. Of these, 72% overlap with previously-identified global

conservation priority areas, while 3% of CCAs occur within moderate to heavily modified lands.

Conservation and restoration of CCAs could safeguard connectivity of the world's PAs, and

dovetail with previously identified global conservation priorities.

Our current global system of protected areas (PAs) has been unable to slow biodiversity loss[1,2] or provide adequate coverage for many threatened species and ecosystems[1,3–7]. PAs are constrained in size, ecological representation and governance[8], and 90% or more exist within a matrix of human-dominated, fragmented land[9] that is rapidly eroding[10,11], endangering animal movement[12,13] and survival[14]. As a result, PAs and the animal populations they contain can become isolated, interrupting the flow of vital ecological (e.g., migration, dispersal) and evolutionary (e.g., gene flow) processes that maintain populations, ecosystems and adaptive capacity[14–17]. For these reasons, the Aichi Target 11 of the Convention on Biological Diversity's 2020 Strategic Plan for Biodiversity stipulated expanding the global PA network not only to cover 17% of terrestrial areas but also to ensure connectivity among PAs[18], to facilitate the flow of individuals, genes and processes[19]. While these targets remained unmet by the end of 2020, post-2020 global biodiversity framework discussions continue to champion the importance of connectivity, both as a stand-alone target and integrated into other relevant targets[20,21]. To date, only a few evaluations of the connectedness of the world's PAs exist[9,22,23] and none explicitly map the functional connectivity of PAs. This research gap prevents the identification of globally important areas for wildlife connectivity and the spatially explicit assessment of potential threats to connectivity.

Here, we modelled the functional connectivity of terrestrial PAs (excluding Antarctica) to quantify the connectedness of each PA and map the world's most critical areas for connectivity conservation. Our analysis focuses on the following questions relevant to global conservation practice and policy: (1) Where are the most and least functionally connected PAs in the world located? (2) How does the functional connectivity of national PA-networks compare to previous estimates of national connectivity, and what mechanisms exist for national governments to improve the functional connectivity of their PA-networks? (3) Where are the world's most

important connectivity landscapes? and (4) How do these vulnerable connectivity landscapes

overlap with existing global prioritization schemes for biodiversity conservation? To answer

these questions we first generated a global resistance-to-movement surface from empirical

estimates of animal movement (624 individuals of 48 mammalian species) in response to the

human footprint index (HFI)[12,24], which describes the combined effects of infrastructure, land use

and human access across the planet. By incorporating observed animal movements, our analysis

permits a first-ever assessment of global functional connectivity, reflecting the predicted flow of

mammal movement between the world's PAs.

We applied circuit theory[25–27] to the resistance-to-movement surface using two distinct

approaches that relate the flow of animal movement across a landscape to the flow of electrical

current across a circuit. First, we quantified the 'effective resistance'[26–29] – previously shown to

predict gene flow[27,30] – between each terrestrial PA and all other PAs on the same landmass (i.e.,

continent or island). Effective resistance relates the total resistance between nodes in a circuit to

the degree that each node (in our case, each PA) is isolated from all others. PAs with high

effective resistance are therefore further away from other PAs and/or are more difficult to travel

to, based on the resistance-to-movement surface; we thus use effective resistance as an index of

PA isolation (hereafter referred to as PAI). Second, we mapped the flow of electrical current

across continents. For each continent we used the center grid cell of each PA as a 'ground' to

which electrical current flowed from all other PAs[28], proportional to PA size, with current

flowing in greater density in regions of lower resistance. We summed the results from all PAs to

produce a global map of relative electrical current densities that reflects the net movement

probabilities of randomly moving individuals (i.e., mammal movement probability [MMP])

across all possible land-based travel routes between all terrestrial PAs. To validate our results,

we compared our global map of MMP against independent GPS data from 366 individuals

representing 7 species of mammals (see Methods).

**Protected Area Isolation**

Does our PA isolation (PAI) index reveal new information about the connectedness of the

world's PAs? While our results agree with the expectation that the least isolated PAs occur

within the world's two most intact biomes – boreal forest and tundra[31,32] (Figs. 1 and S1) – we

nonetheless found stark contrasts between our PAI index and several global connectivity

indicators previously developed for monitoring progress towards post-2020 targets[33] (Fig. 2).

Key to these contrasts is that each global indicator employs a different methodological approach

to assess a different component of connectivity (Fig. S2). For example, we found that PAI, used

to assess functional connectivity, varied considerably among countries that were assigned a

connectivity value of zero by the ConnIntact indicator[9], which is used to assess the contiguity of

intact natural land between PAs irrespective of animal movement (i.e., structural connectivity)

(Fig. 2). Indeed, PAI was only weakly correlated with all existing global connectivity indicators

(Pearson's $r$ ranged from 0.36 – 0.55; Fig. 2) and it identified a different set of countries as most

connected (Table S1). For example, our PI index identified Canada, which has the second-largest

area of wilderness after Russia[34], as the 2nd most connected country, while the other three

indicators identified Canada as only the 15th, 52nd or 111th most connected country. Our results

suggest that the PAI index we present here provides a new view of connectivity that

complements how connectivity is evaluated by the existing global indicators.

**Figure 1. Protected area isolation (PAI).** Isolation of the world's terrestrial PAs, as

measured by effective resistance to mammal movement. The most connected PAs are those

categorized as wilderness areas (i.e., IUCN category Ib) and those that occur within the most

intact biomes (boreal forest and tundra), while the most isolated PAs occur in mangroves (a

critically endangered biome with few PAs).

What potential mechanisms exist for national governments to improve their PAI value?

Because restoration[35] and PA expansion[36] are both recognized strategies for biodiversity

conservation, we compared effects on national PAI from decreasing the human footprint (e.g.,

via the restoration of degraded habitats[37]) and increasing PA coverage, using a linear mixed

effects model with continent as a random intercept. We found that reducing a country's

aggregate human footprint would reduce PAI more strongly than would increasing a country's

PA estate (Fig. S3). Although these strategies do not entail equal costs or effort, we found that on

average a 50% reduction in the human footprint would decrease national PAI by 24% (95% CI:

17-45%), whereas a 50% increase in PA coverage would decrease national PAI by only 11%

(95% CI: 9-21%). Utilizing both strategies in combination, however, has the greatest benefit,

decreasing PAI by 42% (95% CI = 27-73%). These results suggest that habitat restoration, and

favorable land management practices that improve the permeability of anthropogenic landscapes

to animal movement[35,37,38], can complement formal protection efforts to improve connectivity.

**Figure 2. National protected area isolation (PAI).** (A) PAI aggregated to the national

level (dark grey bars depict PAI values); higher values reflect greater degree of isolation. Bars

are organized by continent (AS = Asia; NA = North America; EU = Europe; AF = Africa; SA =

South America; OC = Oceania and Australia). Red-labeled countries have the most connected

national PA networks (95th percentile). (B) Comparisons of our national PAI index to three

existing global indicators of connectivity. ConnIntact[9] is a recently updated version of the

protected-connected index[22]; PARC is the PA connectedness index[59]. The PAI and global

indicator values were weakly correlated, as evaluated using Pearson's *r*.

**Global Connectivity Conservation Priorities**

Beyond the connectedness of each of the world's PAs, can we also identify where connectivity is most vulnerable and in need of protection or restoration around the world? We focused on identifying critical concentrations of MMP that typically occur through corridors and pinch points (Fig. 3). Because these areas have few or no alternative paths for animal movement, they can have disproportionate impacts on connectivity if further restricted or destroyed[27]. We identified concentrated flows using MMP grid cells in the 90th percentile, hereafter referred to as critical connectivity areas (CCAs). We further divided CCAs into two categories based on a threshold HFI value of 4 previously used by others[9,11,31]: "modified" CCAs (HFI ≥ 4) and "intact" CCAs (HFI < 4). We found that 97% of CCAs occur in intact areas and 3% are modified (i.e., 9.7% and 0.3% of the terrestrial realm, respectively). Modified CCAs thus represent small but critical pinch points that if conserved or managed to limit further modification (e.g., through conservation easements, payments for ecosystem services, or community-based conservation[38]) could achieve gains in safeguarding connectivity through anthropogenic landscapes. The vast majority of these modified CCAs are in sub-Saharan Africa, Europe and South America, with the greatest total coverage occurring in the Bolivia, Tanzania, Democratic Republic of Congo and Poland (Fig. S4). The greatest total coverage of intact CCAs on the other hand, are in Russia, Brazil, Australia and Canada (Fig. S5).

**Figure 3. Global mammal movement probability (MMP).** Predicted flow of mammal movement between all PAs based on patterns of electrical current flow. Grid cells with the highest MMP reflect areas where mammal movement is predicted to be most concentrated across the globe, flowing through corridors or pinch points (in light yellow) surrounded by less permeable anthropogenic landscape. Areas in orange and purple reflect moderate flows dispersed across many pathways. Areas of concentrated and dispersed flow are both important to

149 connectivity, but with multiple pathways in dispersed areas, there is a lower risk of total loss of

150 connectivity. Black regions depict areas of lower relative flow (low MMP). Boxes highlight

151 example landscapes. Box A: corridors through mountains of western North America (e.g.,

152 Yellowstone to Yukon corridor). Box B: corridors and dispersed flow across sub-saharan

153 Africa's Kavango-Zambezi Transfrontier Conservation Area and coastal deserts of Namibia. Box

154 C: flows through rainforests of Indonesia and Malaysia (e.g., Heart of Borneo conservation area).

155   Where are CCAs unprotected and thus vulnerable to further land-use conversion? We

156 found that two-thirds of all CCAs are currently unprotected (Fig. 4) and that this lack of formal

157 protection is particularly widespread for intact CCAs. We found small but important clusters of

158 unprotected modified-CCAs in parts of Europe, sub-Saharan Africa and North and South

159 America, where wide-ranging species, such as red deer *Cervus elaphus*, African wild dog *Lycaon*

160 *pictus* and jaguar *Panthera onca*, can occur. Further, roughly 24% of all CCAs are both

161 unprotected and occur on land suitable for future agricultural expansion[39] (Fig. S6). The potential

162 for rapid land conversion to agriculture highlights the need to act quickly to secure these regions

163 for long term connectivity, either via formal protection measures or, with working lands

164 conservation practices that focus on managing for permeable agricultural landscapes[37,38]. Such

165 working lands, managed using strategies to enhance biodiversity conservation (e.g., silvo-

166 pastoral, agroforestry and other agroecological management practices) and benefits to humans

167 (e.g., pollination services and pest control), could also represent an important other effective

168 area-based conservation measure (OECM) and as such could provide important contributions to

169 global conservation policy targets[18,21].

170 **Figure 4. Mapping critical connectivity areas globally (CCAs).** (A) Intact CCAs

171 (blue) and modified CCAs (red) were classified based on mammal movement probability (MMP)

172 and human footprint index values (HFI< 4, HFI ≥ 4, respectively). CCAs in green and orange are

covered by protected areas. Black boxes highlight four areas with clusters of modified CCAs, occurring in: (1) Mexico, (2) Poland, Slovenia, Italy, Belarus and Ukraine, (3) Bolivia and Brazil, and (4) Democratic Republic of Congo, Tanzania and Zambia. Pie charts indicate the proportion of each CCA type and protection status in each continent (AS = Asia; NA = North America; EU = Europe [excluding Russia]; AF = Africa; SA = South America; OC = Oceania and Australia) (B) Proportion of CCAs that overlap with expanded conservation areas of the Global Safety Net (GSN). Stacked bars (far left) depict the proportion of protected and unprotected CCAs. Individual bars (right) represent the proportion of CCAs that overlap with each of the priority conservation areas selected for specific purposes under the GSN.

Which of the world's continents, biomes and ecoregions contain the greatest proportion of CCAs? We found that the tropical moist forests of South America, the tropical grasslands and deserts of Africa, and the boreal forests of Asia contain > 80% more CCAs than other biomes (Fig. 4). Globally, these four biomes contain 41% of the world's ecoregions and a majority of the world's species, 50% of which are contained within tropical moist forests[40]. Of the world's 846 ecoregions, the 50 that were recently identified as having the greatest potential to protect biodiversity[36] also contribute disproportionately to connectivity; unprotected portions of these priority ecoregions contain on average two times greater MMP than all other unprotected ecoregions (Fig. S7) and contain > 42% of the unprotected CCAs. Thus, conserving movement corridors and restoring degraded habitat around pinch points in the unprotected portions of these priority ecoregions is likely to benefit biodiversity conservation *and* safeguard the connectivity of currently established PAs, a key goal in the post-2020 global biodiversity framework[21].

**Connectivity-Biodiversity Conservation Synergies**

Where can we further maximize conservation effectiveness by identifying landscapes that simultaneously enhance connectivity and biodiversity? We examined the proportion of CCAs that overlap with the Global Safety Net (GSN) – a proposed global conservation scheme that identifies new priority areas for expanded protection that would boost the world's terrestrial PA coverage to ~ 50%[36]. We found that roughly 72% of unprotected CCAs, and a majority of CCAs suitable for future agricultural expansion, overlap with the GSN priority areas for expanded conservation (Figs. 4 and S7). Thus, the GSN would protect key biodiversity elements as well as more than 81% of the world's most important areas for connectivity. Of the remaining CCAs not covered by the GSN, many are vulnerable to potential disturbance, degradation and/or land-conversion (9% are modified and roughly 30% have the potential for agricultural development; Fig. S7).

We also examined CCA overlap with other global conservation prioritization schemes that were not explicitly incorporated into the GSN, including Frontier Forests[41], Global 200 Ecoregions[42] and High-Biodiversity Wilderness Areas[43]. We found that > 60% of the CCAs overlap with currently-unprotected portions of these global conservation schemes, and that the Global 200 Ecoregions covered the highest percentage of CCAs (~30%; Fig. S8). This overlap between CCAs and global conservation priorities highlights potential synergies that conservation efforts in these areas could deliver via maintaining globally-significant areas for connectivity, while also preserving rare species and assemblages, biologically outstanding ecoregions and intact habitats critical for biodiversity and ecosystem function.

**Conclusions**

Our study identifies key opportunities for conserving globally important areas for connectivity and for monitoring national PA connectivity using circuit theory methods adapted to

the global scale. Our results show that the functional connectivity approach substantially changes

the view of national PA connectivity compared to the existing global-policy indicators that are

more geared towards assessing other aspects of connectivity. A key difference is that our

approach specifically incorporates observed animal responses to attributes of natural and human-

dominated landscapes, which is lacking in the other global indicators. Our global application of

circuit theory also makes it possible for the first time to assess the permeability of both intact and

human-modified landscapes to mammal movement across the terrestrial realm. With this

approach, we identified small areas of modified but still permeable lands, representing only 0.3%

of the terrestrial world, where timely conservation is of utmost importance to prevent further

impediments to the flow of mammals between PAs.

Our study specifically evaluates the functional connectivity of protected areas at the

global scale, and therefore we do not explicitly consider the connectedness of other potential

sources and destinations of movement (e.g., OECMs or other natural areas), nor do we portray

connectivity for more localized movements (e.g., inter-patch movement). In addition, our

connectivity model is informed only by mammal movements, and therefore may not capture

connectivity for other taxa. Indeed, our results illustrate that differences in national PA

connectivity can occur even among mammalian species; we found that PAs appear more isolated

from the perspective of larger mammals (see Methods and Fig. S9). While this suggests that

larger mammals are more sensitive to higher HFI levels, we found that patterns of flow – that is,

where the flow of mammal movement is dispersed versus constricted – remain consistent across

mammalian body sizes. Finally, although our approach is based on mammalian movement, our

results illustrate how CCAs spatially overlap with global conservation priorities that aim to

protect a variety of taxonomic groups.

We found that the vast majority of the world's CCAs currently under threat (i.e.,

unprotected or located on modified land), or likely to be threatened in the future (i.e., located on

land suitable for future food production), would be protected by expanded conservation under the

proposed Global Safety Net (GSN)[36]. If protected, these areas of GSN-CCA overlap could result

in important conservation synergies that slow or prevent biodiversity loss. Since many of these

areas of overlap are also predicted to be suitable for future agricultural expansion, where formal

protections could be contested over livelihoods or food supply needs, alternative strategies for

conserving or restoring connectivity may be needed. Our results suggest that managing the

matrix for more permeability to animal movement (e.g., habitat restoration) can be a better single

strategy for improving connectivity than adding protected areas alone. However, formal

protection and habitat restoration together generated the largest improvements to connectivity,

supporting the need for holistic approaches that combine working lands conservation[37,38] with

protected areas. Such combined strategies can also provide significant benefits to humans[37,38,44],

thus advancing the post-2020 global biodiversity framework vision of "living in harmony with

nature"[21].

**Methods**

We modelled connectivity using Circuitscape, an open-source program implemented in

Julia[45] and widely used in research and conservation planning[25,27,46–51] that relies on principles

from circuit theory to relate the flow of animal movement across a heterogeneous landscape to

the flow of electrical current across a circuit[27]. This approach treats a gridded resistance-to-

movement surface as a conductive layer where neighboring grid cells are connected by resistors,

whose level of resistance represents the friction of the landscape to animal movement (i.e., low-

resistance grid cells are most likely to be traversed). Electrical current, injected at a source node

representing a source of animal movement, flows across all neighboring landscape grid cells

(resistors) in the circuit to a ground node representing the destination of animal movement.

Connectivity is then evaluated in two distinct ways: (1) from the effective resistance of the

circuit, measuring the degree to which the source node is isolated from the ground node and (2)

from the pattern of electrical current flow (i.e., animal movement probabilities) across the

resistance-to-movement surface[27], as outlined below and illustrated as a workflow in Fig. S10.

Two inputs are required for this analysis: the resistance-to-movement surface describing

the friction of the landscape to animal movement and the layer specifying sources and

destinations of animal movement. To model functional connectivity (Fig. S3), we generated a

resistance-to-movement surface using the data and linear mixed effects model of 0.95 quantile

displacement distances from 10-day intervals presented in Tucker et al. (2018)[12] for 624 GPS-

collared individuals from 48 species of mammals. This movement model included as predictor

variables the human footprint index (HFI)[24], normalized difference vegetation index (NDVI),

body weight and dietary guild, and species as a random effect. We focused on the long-distance,

long-time scale movements as those more likely to represent the scale of movement occurring

between protected areas, thus representing movements promoting gene flow, rescue effects or

climate adaptation.

Using the model above, we predicted mammal movement for all values of HFI across

Earth's terrestrial surface after resampling the HFI layer from 1 km$^2$ to ~70 km$^2$ (in Mollweide

equal area projection) to reduce computation time in our Circuitscape analysis from months to

days (using a Google virtual machine with 96 CPUs and 360 GB memory). We rescaled these

predicted movements to between 0 and 1 to represent the landscape's permeability to mammal

movement across the globe. We then used an inverse linear function (e.g., 1-permeability) to

convert the permeability surface to a resistance-to-movement surface[52–54]. Following other

studies, we further modified resistance-to-movement, M, by percent slope to reflect the energetic

costs of moving in steep terrain[28,50,55] using the equation $\widehat{M} = M^{1-\left(\frac{s}{max(s)}\right)}$ , where s = terrain

slope.

We used all terrestrial PAs from the May 2020 World Database on Protected Areas

(WDPA)[56] as the sources and destinations of mammal movement (i.e., source and ground of

electrical current). We combined this WDPA layer with PAs in China from the April 2018

WDPA[56], because China removed all its national PAs from the WDPA after this date. As

recommended in the WDPA best practices guide, we removed any PA that did not report its

area, or with a 'Proposed' status or 'UNESCO-MAB Biosphere Reserve' designation

(https://www.protectedplanet.net/en/resources/calculating-protected-area-coverage). We used PA

centroids as the source and ground of electrical current, rather than entire PA polygons, because

this allowed for estimates of electrical current flow within PAs, rather than assuming flow is

equally good within PA boundaries[28]. We rasterized PA-centroids to match the resolution and

extent of the resistance-to-movement surface. We excluded PA-centroid grid cells for which the

sum total of PA area was less than half the grid cell resolution, 35 km$^2$, to ensure that electrical

source grid cells are primarily represented by a probable source or destination of mammal

movement. By excluding these small PAs (similar to[28,57]), we also avoided overestimating

connectivity from using source and destination nodes much smaller than the resolution of our

analysis and the home ranges of many of the mammals that we examined. Excluded PAs

represented 90% of the total number of individual PAs but, with a mean area of 2.4 $km^2$, they

comprised only ~2% of the total area of the global PA estate (Fig S11 and Table S2). Included

PAs were on average 420 times larger than excluded PAs. Note also that excluded PA-centroids

were removed only as sources and destinations of mammal movement in the model; importantly

their benefits to connectivity were fully incorporated into the resistance-to-movement surface as

areas of lower resistance (i.e., lower HFI). We modelled connectivity by continent, excluding

Antarctica, islands $< 2,500$ $km^2$ and islands with $< 2$ PAs. We also combined Europe and Asia to

avoid boundary effects where the two continents meet.

***Protected Area Isolation***

To obtain a value of effective resistance for all terrestrial PAs (steps 4-5 in Fig. S10), we

applied Circuitscape's one-to-all mode[58] to the input layers. This mode connects a focal PA-

centroid to an electrical current source and connects all remaining PA-centroids to ground,

thereby generating one circuit where electrical current flows from the focal PA source to all

other PAs within the continent. When the circuit is connected, effective resistance, $R_{eff}$, is

calculated as $1/R_{eff} = 1/R_1 + 1/R_2 + 1/R_3 ... 1/R_n$, where each $R_i$ represents a pathway (i.e.,

summed grid cell 'resistors' connected in series) between PAs. Following this central equation of

circuit theory, effective resistance decreases, and thus connectivity increases, as more

connections (i.e., redundant pathways, $R_i$) are added to the circuit[27]. For this process we maintain

the default current injection of 1 Amp, because each grid cell 'resistor' in the landscape has an

assigned resistance value, effective resistance does not vary with the amount of current injected.

We repeated this procedure for each PA and mapped the resulting PA isolation (PAI) values for

all qualified terrestrial PAs (i.e., with summed area within a grid cell of $> 35$ km$^2$) and evaluated

PAI within the world's biomes[42] and continents.

We calculated the median PAI for each country's PA network. Because measures of PAI

were generated from connecting PAs across entire continents, national PAI may be affected by

connections with neighboring cross-border PAs. To ensure that national PAI was not unduly

sensitive to the location and extent of PAs outside the country's borders, we repeated the steps to

model PAI at the country level for 25 randomly selected countries. To do so, we clipped the

resistance-to-movement surface and PA layer to each of the 25 countries, re-applied circuit

theory using the one-to-all approach and re-computed median PAI by country. National PAI

from the continent and country analyses were highly correlated (Pearson's $r = 0.92$; Fig. S12),

suggesting that metrics from our global analysis are suitable proxies for country-level PAI.

We evaluated how hypothetical changes to mean-national HFI and proportion of PA

would affect national PAI. We used a linear mixed effects model with mean HFI and % PA area

as predictor variables scaled by subtracting the mean and dividing by the standard deviation to

evaluate the relative effects of each, with continent as a random intercept. We then used the

estimated coefficients to simulate the effects of changing the proportion of protected area

(including PAs of all sizes) and mean HFI on national PAI.

We compared our national PAI index, which explicitly accounts for mammal movement,

to three proposed indicators of progress towards post-2020 global connectivity targets that lack

measurements of animal movement (Fig. S3): (1) the proportion of country protected, (2)

ConnIntact[9] – an updated version of Protconn[22], the currently proposed global indicator – and (3)

the PA connectedness index (PARC)[59]. ConnIntact is a network-based structural connectivity

approach that accounts for the number of PAs, total area of PAs, total country area and the

contiguity of intact land between PAs. Specifically, this approach assumes PAs are perfectly

connected whenever there is a contiguous pathway of 1 km$^2$ grid cells of intact natural land

directly connecting PA edges or indirectly connecting PAs through connecting intermediate PAs

as stepping stones. PAs are assumed to be completely disconnected when there is no contiguous

pathway of intact land. The PARC index accounts for connectivity between ~1 km$^2$ grid PA grid

cells and between these PA grid cells and ~1 km$^2$ grid cells containing primary vegetation using

a cost-distance approach where the cost of moving between grid cells is assumed to be

negatively related to the proportion of primary vegetation (i.e., the more primary vegetation in a

grid cell, the lower the cost). We used Pearson's *r* to evaluate the correlations between national

PAI and the three global indicators of connectivity.

***Mapping Global Connectivity***

To map patterns of mammal movement probability (MMP) using electrical current

density (steps 6-10 in Fig. S10) we applied Circuitscape's all-to-one mode[58] to the input layers.

Similar to the pairwise mode[58], the all-to-one mode produces patterns of electrical current flow

between each source-ground pair but with greater computational efficiency. It does this by

generating many circuits simultaneously; in our application, one PA-centroid is designated as

ground while all remaining PA-centroids are current sources. At each source, we injected an

electrical current amount proportional to PA area (e.g., we injected 100 Amps into any PA

centroid that represented a 100-km$^2$ PA), to account for the effect of PA size on the number of

animals potentially moving from each PA (i.e., assuming that all else equal, large PAs have a

greater abundance of wildlife and thus more dispersing animals than small PAs[60]). To account

for the size of the destination PA (i.e., all else equal, large PAs will receive more immigrating

animals than small PAs[60]), we multiplied the output electrical current density map – depicting

electrical current flow from all source PAs to one ground PA – by the area of the corresponding

ground PA. We repeated this process for each PA-centroid connected to ground and summed all

maps to obtain a map of cumulative electrical current density (i.e., MMP) per continent.

We found that electrical current densities increased with land mass area. To correct for

this effect and 'normalize' electrical current densities into a common scale – using current

density/km$^2$ – we divided electrical current densities by the area of their corresponding land

mass. Finally, we merged all electrical current density/km$^2$ maps together to generate a

comparable global map.

***Validation and Sensitivity Analyses***

We validated our global map of electrical current density (MMP) using two different

approaches. First, we used independent GPS movement data (> 1.5 million locations) from 366

individuals and 7 mammal species, combined with GPS movement data (> 220,000 locations)

from another 88 individuals and 4 species included in the Tucker et al. (2018)[12] analysis, to

evaluate whether connectivity values were higher where animals were observed moving (see

Table S3 for details and list of data providers; see Fig. S13 for dataset locations). We obtained

known range data for each species (https://www.iucnredlist.org/). Then for each species-dataset

we (1) generated a 95% utilization distribution (UD) (i.e., 95% kernel home range) from the GPS

data using the adehabitatHR package[61] in the R statistical computing environment[62] version

3.6.3, (2) randomly selected 1000 points within that UD to represent locations where those

individuals were likely to occur and (3) extracted electrical current density values for those

points. To compare these values to electrical current density values where the collared

individuals could have been but were not observed, we (4) calculated max displacement distance

using the adehabitatHR package[61] in R and buffered the UD by this distance, (5) clipped the

species range data to that buffer, (6) randomly selected points within that clipped range outside

of the UD and (7) extracted electrical current density values from those random points.

We used a Bayesian hierarchical approach to evaluate the differences in electrical current density extracted from points inside and outside the UDs. We used the package brms[63] in R to construct the model, with log electrical current density as the response variable, inside or outside UD as a categorical predictor variable, and a random intercept and slope by species. We used uninformative prior distributions, ran 4 iterations of the model with 400 chains per iteration, and confirmed that all parameters converged with R-hat values < 1.1. Using the predicted posterior distributions, we calculated the percentage that electrical current densities were higher inside the UDs and provide strong evidence for species moving within areas of higher predicted movement probability (Fig. S14).

Second, using the same data, we evaluated whether animals were selecting for critical connectivity areas (CCAs) (i.e., grid cells having electrical current density values in the 90th percentile). Using a raster of CCAs and the GPS datasets described above, we extracted the number of unique CCA grid cells that were used by each species. We calculated the proportion of used grid cells that were CCAs. We also calculated the proportion of unused grid cells that were CCAs, within the same buffered areas described previously. From this, we calculated the percent that CCAs were used (i.e., inside UDs) relative to what was available and found strong evidence that 9 of the 11 species preferred to move within CCAs (Table S4).

We also evaluated the sensitivity of effective resistance (PAI) and electrical current density (MMP) to species body size[12,64] and continent-specific species attributes. We first created three different resistance-to-movement surfaces using a similar linear mixed effects model that was used in Tucker et al. (2018)[12] and our previous analysis, but instead fitted to: (1) 10-day displacement distances recorded by species whose mean body mass was greater than the mean body mass, ~ 53 kg, of the 48 species considered in our full model, (2) 10-day displacement distances recorded by species whose mean body mass was less than the mean body mass of the

48 species, and (3) 10-day displacement distances recorded by African species only. We repeated

the steps in the workflow described for our main analysis (Fig. S10) and compared effective

resistances and electrical current densities between our full model and the sensitivity test models

(Fig. S9). Effective resistances increased with body size and electrical current densities were

highly correlated across models ($R^2 > 0.9$), suggesting that while larger mammals are more

sensitive to high HFI, the patterns of flow remain consistent across mammalian body sizes.

*Areas of Overlap for Priority Connectivity and Biodiversity Sites*

431            We summarized the total area of both modified and intact categories of critical

connectivity areas (CCAs) by country and identified countries with the largest area (95th

percentile) of CCAs. To assess the protection status of CCAs, we quantified CCA overlap with

PAs using the May 2020 World Database on Protected Areas (WDPA) and the April 2018

WDPA for PAs in China[56]. We used both WDPA polygons and WDPA points files to generate

our PA layer. We converted WDPA points to polygons using circular buffers with areas equal to

their corresponding reported PA area (https://www.protectedplanet.net/en/resources/calculating-

protected-area-coverage), and then we spatially intersected these buffers with their country of

origin to exclude parts of the buffer that occurred in other countries. We compared MMP in

protected and unprotected portions of the 50 ecoregions recently identified as contributing most

to enhancing biodiversity protection[36] and in protected and unprotected portions of all other

ecoregions (https://ecoregions2017.appspot.com/). We also summarized the total area of CCAs

within continents, ecoregions[40] and biomes[42].

444            We quantified the amount of unprotected CCAs that overlapped with lands where future

agricultural expansion is predicted using a global land-systems (GLS) map describing crop

suitability as determined by agro-climatic and agro-edaphic conditions[39]. The crop suitable areas

in the GLS map included those that were categorized as very high to marginally suitable for high

intensity cropland along with low intensity cropland, and thus may overpredict cropland

expansion in some cases[39]. However, suitable areas for tree plantation and livestock expansion

were not included in the GLS map, potentially resulting in an underprediction of CCA overlap

with future agricultural areas.

We examined the proportion of unprotected CCAs that overlapped with four areas

prioritized for expanded conservation under the Global Safety Net (GSN)[36], the most recent

global prioritization scheme for biodiversity conservation. These areas are prioritized in the GSN

because of their disproportionate contribution to (1) important rare and threatened species, (2)

distinct species assemblages, (3) intact assemblages of large mammals and (4) wilderness or

other macrorefugia for wildlife. Conservation of these additional four priority areas, plus climate

stabilization sites, would extend coverage of the PA estate to ~50% of the terrestrial realm.

We also quantified the proportion of unprotected CCAs that overlapped with three other

global conservation prioritization schemes that target other critical ecological conditions or

regions important to biodiversity (which may be partially overlapping with the GSN), including

Frontier Forests[41], Global 200 Ecoregions[65] and High-Biodiversity Wilderness Areas[43]. Frontier

forests represent large contiguous wilderness areas that specifically encompass forests and

conditions important to forest integrity. The Global 200 Ecoregions is based on representation

and biologically outstanding character, and is thus meant to encompass the most biologically

outstanding ecoregions in every biome-realm combination. High Biodiversity Wilderness Areas

are the complement of Biodiversity Hotspots, thus encompassing non-threatened areas of high

biodiversity.

**References**

1. Maxwell, S. L. *et al.* Area-based conservation in the twenty-first century. *Nature* **586**, 217–227 (2020).

2. Butchart, S. H. M. *et al.* Shortfalls and Solutions for Meeting National and Global Conservation Area Targets. *Conserv. Lett.* **8**, 329–337 (2015).

3. Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction, distribution, and protection. *Science* **344**, (2014).

4. Venter, O. *et al.* Bias in protected-area location and its effects on long-term aspirations of biodiversity conventions. *Conserv. Biol.* **32**, 127–134 (2018).

5. Venter, O. *et al.* Targeting Global Protected Area Expansion for Imperiled Biodiversity. *PLoS Biol.* **12**, (2014).

6. Rodrigues, A. S. L. *et al.* Global gap analysis: Priority regions for expanding the global protected-area network. *Bioscience* **54**, 1092–1100 (2004).

7. Terraube, J., Van doninck, J., Helle, P. & Cabeza, M. Assessing the effectiveness of a national protected area network for carnivore conservation. *Nat. Commun.* **11**, 1–9 (2020).

8. Visconti, B. P. *et al.* Protected area targets post-2020. *Science* **364**, 239–241 (2019).

9. Ward, M. *et al.* Only ten percent of the global terrestrial protected area network is connected via intact land. *Nat. Commun.* 1–10 (2020) doi:10.1101/2020.01.28.920488.

10. Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S. & Kiesecker, J. Managing the middle: A shift in conservation priorities based on the global human modification gradient. *Glob. Chang. Biol.* **25**, 811–826 (2019).

11. Watson, J. E. M. *et al.* Persistent Disparities between Recent Rates of Habitat Conversion and Protection and Implications for Future Global Conservation Targets. *Conserv. Lett.* **9**, 413–421 (2016).

494    12.    Tucker, M. A. *et al.* Moving in the Anthropocene: Global reductions in terrestrial

495            mammalian movements. *Science* **359**, 466–469 (2018).

496    13.    Harris, G., Thirgood, S., Hopcraft, J. G. C., Cromsigt, J. P. G. M. & Berger, J. Global

497            decline in aggregated migrations of large terrestrial mammals. *Endanger. Species Res.* **7**,

498            55–76 (2009).

499    14.    Haddad, N. M. *et al.* Habitat fragmentation and its lasting impact on Earth's ecosystems.

500            *Sci. Adv.* **1**, 1–10 (2015).

501    15.    Pacifici, M., Di Marco, M. & Watson, J. E. M. Protected areas are now the last

502            strongholds for many imperiled mammal species. *Conserv. Lett.* **13**, 1–7 (2020).

503    16.    UNEP. Ecological connectivity: a bridge to preserving biodiversity. *Front. 2018/19*

504            *Emerg. Issues Environ. Concern. United Nations Environ. Program.* 24–37 (2019).

505    17.    Schmidt, C., Domaratzki, M., Kinnunen, R. P., Bowman, J. & Garroway, C. J. Continent-

506            wide effects of urbanization on bird and mammal genetic diversity. *Proc. R. Soc. B Biol.*

507            *Sci.* **287**, 20192497 (2020).

508    18.    CBD. *Strategic Plan for Biodiversity 2011-2020*. https://www.cbd.int/sp/ (2011).

509    19.    Taylor, P. D., Fahrig, L., Henein, K. & Merriam, G. Connectivity Is a Vital Element of

510            Landscape Structure Author(s): *Oikos* **68**, 571–573 (1993).

511    20.    CMS. Ecological Connectivity in the Post-2020 Global Biodiversity Framework.

512            https://go.nature.com/38GB6Nf (2019).

513    21.    Secretariat of the United Nations Convention on Biological Diversity. *First Draft of the*

514            *Post-2020 Global Biodiversity Framework*. *Cbd/Wg2020/3/3* (2021).

515    22.    Saura, S. *et al.* Protected area connectivity: Shortfalls in global targets and country-level

516            priorities. *Biol. Conserv.* **219**, 53–67 (2018).

517    23.    Saura, S. *et al.* Global trends in protected area connectivity from 2010 to 2018. *Biol.*

518      *Conserv.* **238**, 108183 (2019).

24.      Venter, O. *et al.* Sixteen years of change in the global terrestrial human footprint and

         implications for biodiversity conservation. *Nat. Commun.* **7**, 1–11 (2016).

25.      Dickson, B. G. *et al.* Circuit-theory applications to connectivity science and conservation.

         *Conserv. Biol.* **33**, 239–249 (2019).

26.      McRae, B. H. Isolation By Resistance. *Evolution (N. Y).* **60**, 1551 (2006).

27.      McRae, B. H., Dickson, B. G., Keitt, T. H. & Shah, V. B. Using circuit theory to model

         connectivity in ecology, evolution, and conservation. *Ecology* **89**, 2712–2724 (2008).

28.      Dickson, B. G. *et al.* Informing Strategic Efforts to Expand and Connect Protected Areas

         Using a Model of Ecological Flow , with Application to the Western United States.

         *Conserv. Lett.* **10**, 564–571 (2017).

29.      Keeley, A. T. H., Beier, P. & Jenness, J. S. Connectivity metrics for conservation planning

         and monitoring. *Biol. Conserv.* **255**, 109008 (2021).

30.      Garroway, C. J., Bowman, J. & Wilson, P. J. Using a genetic network to parameterize a

         landscape resistance surface for fishers, Martes pennanti. *Mol. Ecol.* **20**, 3978–3988

         (2011).

31.      Riggio, J. *et al.* Global human influence maps reveal clear opportunities in conserving

         Earth's remaining intact terrestrial ecosystems. *Glob. Chang. Biol.* **26**, 4344–4356 (2020).

32.      Watson, J. E. M. *et al.* Catastrophic Declines in Wilderness Areas Undermine Global

         Environment Targets. *Curr. Biol.* **26**, 2929–2934 (2016).

33.      Secretariat of the United Nations Convention on Biological Diversity. *Zero Draft of the*

         *Post- 2020 Global Biodiversity*. (2020).

34.      Morales-Hidalgo, D., Oswalt, S. N. & Somanathan, E. Status and trends in global primary

         forest, protected areas, and areas designated for conservation of biodiversity from the

542             Global Forest Resources Assessment 2015. *For. Ecol. Manage.* **352**, 68–77 (2015).

543    35.    Strassburg, B. B. N. *et al.* Global priority areas for ecosystem restoration. *Nature* **586**,

544             724–729 (2020).

545    36.    Dinerstein, E. *et al.* A 'Global Safety Net' to reverse biodiversity loss and stabilize Earth's

546             climate. *Sci. Adv.* **6**, 1–14 (2020).

547    37.    Garibaldi, L. A. *et al.* Working landscapes need at least 20% native habitat. *Conserv. Lett.*

548             1–10 (2020) doi:10.1111/conl.12773.

549    38.    Kremen, C. & Merenlender, A. M. Landscapes that work for biodiversity and people.

550             *Science* **362**, (2018).

551    39.    Kehoe, L. *et al.* Biodiversity at risk under future cropland expansion and intensification.

552             *Nat. Ecol. Evol.* **1**, 1129–1135 (2017).

553    40.    Dinerstein, E. *et al.* An Ecoregion-Based Approach to Protecting Half the Terrestrial

554             Realm. *Bioscience* **67**, 534–545 (2017).

555    41.    Bryant, D., Nielsen, D. & Tangley, L. *The Last Frontier Forests: Ecosystems and*

556             *Economics on the Edge*. (1997).

557    42.    Olson, D. M. *et al.* Terrestrial ecoregions of the world: A new map of life on Earth.

558             *Bioscience* **51**, 933–938 (2001).

559    43.    Mittermeier, R. A. *et al.* Wilderness and biodiversity conservation. *Proc. Natl. Acad. Sci.*

560             *U. S. A.* **100**, 10309–10313 (2003).

561    44.    Díaz, S. *et al.* Assessing nature's contributions to people. *Science* **359**, 270–272 (2018).

562    45.    Hall, K. R. *et al.* Circuitscape in Julia: Empowering Dynamic Approaches to Connectivity

563             Assessment. *Land* **10**, 301 (2021).

564    46.    McClure, M. L., Dickson, B. G. & Nicholson, K. L. Modeling connectivity to identify

565             current and future anthropogenic barriers to movement of large carnivores: A case study

566    in the American Southwest. *Ecol. Evol.* **7**, 3762–3772 (2017).

567    47.    Brennan, A. *et al.* Characterizing multispecies connectivity across a transfrontier

568           conservation landscape. *J. Appl. Ecol.* **57**, 1700–1710 (2020).

569    48.    Naidoo, R. *et al.* Evaluating the effectiveness of local- and regional-scale wildlife

570           corridors using quantitative metrics of functional connectivity. *Biol. Conserv.* **217**, 96–103

571           (2018).

572    49.    Koen, E. L., Bowman, J., Sadowski, C. & Walpole, A. A. Landscape connectivity for

573           wildlife: Development and validation of multispecies linkage maps. *Methods Ecol. Evol.*

574           **5**, 626–633 (2014).

575    50.    Marrec, R. *et al.* Conceptual framework and uncertainty analysis for large-scale, species-

576           agnostic modelling of landscape connectivity across Alberta, Canada. *Sci. Rep.* **10**, 1–15

577           (2020).

578    51.    McRae, B. *et al.* Conserving Nature's Stage: Mapping Omnidirectional Connectivity for

579           Resilient Terrestrial Landscapes in the Pacific Northwest. 47 (2016)

580           doi:10.13140/RG.2.1.4158.6166.

581    52.    Keeley, A. T. H., Beier, P. & Gagnon, J. W. Estimating landscape resistance from habitat

582           suitability: effects of data source and nonlinearities. *Landsc. Ecol.* **31**, 2151–2162 (2016).

583    53.    González-Saucedo, Z. Y., González-Bernal, A. & Martínez-Meyer, E. Identifying priority

584           areas for landscape connectivity for three large carnivores in northwestern Mexico and

585           southwestern United States. *Landsc. Ecol.* **4**, (2021).

586    54.    Brennan, A. *et al.* Examining speed versus selection in connectivity models using elk

587           migration as an example. *Landsc. Ecol.* **33**, 955–968 (2018).

588    55.    Theobald, D. M., Reed, S. E., Fields, K. & Soulé, M. Connecting natural landscapes using

589           a landscape permeability model to prioritize conservation activities in the United States.

*Conserv. Lett.* **5**, 123–133 (2012).

56.  UNEP-WCMC and IUCN. Protected Planet: the World Database on Protected Areas

(WDPA), Cambridge, UK. www.protectedplanet.com.

57.  Barnett, K. & Belote, R. T. Modeling an aspirational connected network of protected areas

across North America. *Ecol. Appl.* 0–2 (2021) doi:10.1002/eap.2387.

58.  McRae, B., Shah, V. & Mohapatra, T. *Circuitscape 4 User Guide*. (2013).

59.  CSIRO. *Protected Area Connectedness Index (PARC-Connectedness)*.

https://www.bipindicators.net/indicators/protected-area-connectedness-index-parc-

connectedness (2019).

60.  Hanski, I. & Gyllenberg, M. Uniting two general patterns in the distribution of species.

*Science* **275**, 397–400 (1997).

61.  Calenge, C. Home Range Estimation in R: the adehabitatHR Package. (2015).

62.  R Core Development Team. R: A language and environment for statistical computing.

(2020).

63.  Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*

*Softw.* **80**, (2017).

64.  Pacifici, M. *et al.* Global correlates of range contractions and expansions in terrestrial

mammals. *Nat. Commun.* **11**, 1–9 (2020).

65.  Olson, D. M. & Dinerstein, E. The global 200: Priority ecoregions for global conservation.

in *Annals of the Missouri Botanical Garden* vol. 89 199–224 (2002).

**Acknowledgements**

We are grateful to Christina Kennedy for her thoughtful comments on the manuscript.

**Competing interests**

The authors declare no competing interests.