Network analysis of ten thousand genomes shed light on *Pseudomonas* diversity and classification

Hemanoel Passarelli-Araujo[1,2,*], Glória Regina Franco[1], Thiago M. Venancio[2,*]

[1]Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

[2]Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, RJ, Brazil.

*Corresponding authors

Av. Alberto Lamego 2000, P5 sala 217; Parque Califórnia

Campos dos Goytacazes, RJ, Brazil

CEP: 28013-602

HPA: hemanuel.passarelli@gmail.com; TMV: thiago.venancio@gmail.com

SUMMARY

The growth of sequenced bacterial genomes has revolutionized the assessment of microbial
diversity. *Pseudomonas* is a widely diverse genus, comprising isolates associated with processes
from pathogenesis to biotechnological applications. However, this high diversity led to historical
taxonomic inconsistencies. Although type strains have been employed to estimate
Pseudomonas diversity, they represent a small fraction of the genomic diversity at a genus level.
We used 10,035 available *Pseudomonas* genomes, including 210 type strains, to build a genomic
distance network to estimate the number of species through community identification. We
identified inconsistencies with several type strains and found that 25.65% of the *Pseudomonas*
genomes deposited on Genbank are misclassified. We retrieved the 13 main *Pseudomonas*
groups and proposed *P. alcaligenes* as a new group. Finally, this work provides new insights on
the phylogenetic boundaries of *Pseudomonas* and highlights that the *Pseudomonas* diversity has
been hitherto overlooked.

Keywords: *Phylogenomics, Pseudomonads, Taxonomy, Community detection*

INTRODUCTION

Biological networks have been an essential analytical tool to better understand microbial diversity and ecology[1, 2]. A network is a set of connected objects, in which objects can be represented as nodes and connections as edges. Networks provide a simple and powerful abstraction to evaluate the importance of individual or clustered nodes in maintaining a given system. Coupled with whole-genome sequencing, it can refine our knowledge about genetic relationships of diverse bacteria such as *Pseudomonas*.

Pseudomonas is a genus within the *Gammaproteobacteria* class, whose members colonize aquatic and terrestrial habitats. These bacteria are involved in plant and human diseases, as well as in biotechnological applications such as plant growth-promotion and bioremediation[3]. The genus *Pseudomonas* was described at the end of the nineteenth century based on morphology, and its remarkable nutritional versatility was recognized thereafter[4]. The metabolic diversity of pseudomonads, combined with biochemical tests to describe species, culminated in a chaotic taxonomic situation[4].

In 1984, the genus was revised and subdivided into five groups based on DNA-DNA and rRNA-DNA hybridization[5], with group I retaining the name *Pseudomonas*. Over the past 30 years, other molecular markers such as housekeeping genes have been used to mitigate the issues of *Pseudomonas* taxonomy[6, 7, 8]. Based on the 16S rRNA gene sequences, the genus is divided into three main lineages represented by *Pseudomonas pertucinogena*, *Pseudomonas aeruginosa*, and *Pseudomonas fluorescens*[9]. These lineages comprise groups of different species – both, lineages and groups receive the name of the representative species. Currently, there are 254 *Pseudomonas* species with validated names according to the List of Prokaryotic Names with Standing in the Nomenclature (LPSN)[10]. However, although the genus division into lineages and groups has facilitated the classification of new species, the remnants of the *Pseudomonas* misclassification still linger in public databases[11, 12].

The explosion in availability of complete genomes for both cultured and uncultured microorganisms has improved the classification of several bacteria, including *Pseudomonas*[8, 13]. One of the gold standards for species circumscription is the digital whole genome comparison by Average Nucleotide Identity (ANI)[14]. Since using only genomes from type strains might bias and provide an unrealistic picture of microbial diversity, we aimed to estimate the *Pseudomonas* diversity using all available genomes through a network approach. Here, we provide new perspectives on *Pseudomonas* diversity by exploring the topology of the genomic distance network and the phylogenetic tree from representative genomes. This work also provides novel insights about the misclassification and phylogenetic borders of *Pseudomonas*.

RESULTS

Dataset collection

We obtained 11,025 genomes from GenBank on June 2020. After evaluating the quality of each genome (see methods for more details) and removing fragmented genomes, 10,035 genomes passed in the 80% quality threshold (Figure S1). We used 238 type strains with available genomes and names validly published according to the *List of Prokaryotic Names with Standing in Nomenclature* in March 2021. According to the NCBI classification, the top four abundant

species in our dataset are *P. aeruginosa* (n = 5,088), *P. viridiflava* (n = 1,509), *Pseudomonas sp.*
(n = 1,083), and *P. syringae* (n = 435) (Table S1). The sizes of the retrieved genome ranged from
3.0 to 9.4 Mb.

Genome-based analyses reveals the presence of synonymous *Pseudomonas* species
The misclassification of some *Pseudomonas* type strains has been reported by several studies [8,]
[15, 16, 17]. Type strains play an important role in taxonomy by anchoring species names as
unambiguous points of reference[18]. In this context, the term "synonym" refers to the situation
where the same taxon receives different scientific names. We used 238 type strain genomes to
evaluate the presence of synonymous species in *Pseudomonas.* The ANI was computed for all
type strains to construct an identity network that was further used to check the linkage between
genomes based on a 95% ANI threshold (Figure 1). Since 95% has been accepted as species
delimitation threshold[14], connections between type strains indicate synonymous names or
subspecies.
We identified 30 connected genomes in the ANI network (Figure 1). Four of these
connected genomes are expected because they represent *P. chlororaphis* and its subspecies. Of
the 26 remaining connected species, 15 have been previously reported, such as that in the group
containing *P. amygdali*, *P. ficuserectae*, and *P. savastanoi*[15, 16]. Here, we observed 11
connections, including the one between *P. panacis* and *P. marginalis* with 97.34% identity,
suggesting that *P. panacis* is a later synonym of *P. marginalis*.

91
Figure 1. Type strain validation based on Average Nucleotide Identity. Each node in the network
represent a type strain genome and nodes are connected if they share at least 95% identity. The left panel
is a magnified representation of the connected nodes, with edges colored according to percent identity
between the nodes.

The *Pseudomonas* genomic distance network is highly structured
In networks, the community structure plays an important role to understand network topology.
We used all 10,035 *Pseudomonas* genomes to construct a distance network to estimate the
number of *Pseudomonas* species from the number of communities detected in this network.
Since alignment-based methods to estimate genome similarity (e.g. ANI) is computationally
expensive due to the algorithm quadratic time complexity[19], it becomes impractical for thousand

genomes. Therefore, we estimated the Mash distance that strongly correlates with ANI and can
be rapidly computed for large datasets[20].

Mash distances are computed by reducing large sequences to small and representative
sketches[20]. We estimated the pairwise Mash distance for all genomes using sketch sizes of 1000
and 5000, which converged to similar distance values (Figure S2a). However, we observed that
the greater the distance between two *Pseudomonas* genomes, the more divergent the distance
estimation (Figure S2b), although the density distribution is similar (Figure S2c). The final
distance between two genomes was given as the average distance value from both sketch sizes.
We used the reciprocal of Mash distance (1 - Mash) to estimate the ANI for all 10,035 genomes.

We generated a weighted *Pseudomonas* distance network considering nodes as
genomes and edges as the identity between two genomes. Although the 95% ANI value has been
widely accepted to delineate species, we evaluated how different thresholds affect network
topology by assessing density, transitivity, and the number of connected components (Figure 2).
The network density, i.e. the ratio of the number of edges and the number of possible edges,
decreased throughout the interval, but stabilized between 90% and 97% ANI, keeping the
network topology almost unchanged (Figure 2a). To estimate how structured was the network
with different ANI thresholds, we also computed the global network transitivity (also called as
global clustering coefficient) (Figure 2b). The global transitivity measures the probability of a
given node having adjacent nodes interconnected. The high transitivity values revealed that the
Pseudomonas network is highly structured (i.e. formed by tightly connected clusters) and
reaches maximum values between 88% and 97%. This structured profile was observed before
for the *P. putida* group network[17], indicating that communities in *Pseudomonas* distance
networks rarely overlap.

127
Figure 2. *Pseudomonas* distance network topology evolution. a) Proportion of present connections
(network density) and b) global transitivity change over different identity (1 – Mash) cut-off values. c)
Number of network components detected with different identity thresholds. Light orange dots represent
the total number of components, whereas the dark dots represent only components with more than ten
nodes.

To decrease the influence overrepresented species (e.g. *P. aeruginosa*) on the
topological network statistics, we also computed the variation in the number of components
(Figure 2c). A connected component in a network is a subset of nodes connected via a path. At
70% identity, we had a single giant connected component. Expectedly, the number of connected

components increased with the identity threshold because of the emergence of smaller
components or even orphan nodes. Interestingly, connected components with more than 10
nodes arose only above 81% identity threshold and stabilizes close to 95%, highlighting that the
95% ANI threshold is an accurate threshold for species demarcation.
We used the *Pseudomonas* network discarding connections lower than 95% identity to
estimate the number of species from the number of communities in the network. We detected
573 communities by using the label propagation algorithm[21]. This number is similar to the
number of connected components at 95% identity threshold (n = 570), further supporting that
the *Pseudomonas* distance network is highly structured, containing non-overlapping
communities. By considering each community as a different *Pseudomonas* species, we
evaluated the distribution of type strains in these communities.
Seventeen communities had more than one type strain in the same cluster, indicating
the existence of later heterotypic synonyms, as shown in Figure 1. For each community, we
assigned only one representative genome (see methods for more detail). For example, in the
community containing *P. amygdali, P. ficuserectae, and P. savastanoi,* we maintained *P.*
amygdali as the representative strain and the others were considered later heterotypic
synonyms, as previously proposed[11]. We observed that only 210 communities (36.64%) had
representative genomes from validly described species, reinforcing the underestimation of the
number of *Pseudomonas* species if only the type strains are considered.
Regarding the community's sizes, *P. aeruginosa* corresponds to the largest community,
which comprises 5116 genomes (Figure 3, Table S2). Most communities had few genomes.
Although large communities tend to have type strains, 61 type strains (29.04%) are single nodes
(Figure 3, Table S2), further demonstrating that estimating the diversity of *Pseudomonas* only
by type strains severely underestimates diversity. For example, the community containing
Pseudomonas spp7 has 122 genomes and is potentially a new genomospecies.
163

Comparison with NCBI classification highlights *Pseudomonas* misclassification
After delimiting the species by the community detection approach, we compared them with the
classification available in NCBI Taxonomy[22]. Briefly, we computed how many genomes were
deposited with a given species name and how many genomes were identified for that species
by our network approach. Of the 10,035 genomes used in this work, 25.65% were misclassified
in NCBI Taxonomy (Table S4). The most poorly classified species were *P. brassicacearum*
(95.65%), *P. fluorescens* (95.23%), *P. stutzeri* (94.58%), and *P. putida* (88.70%). This high rate of
misclassification is linked to the type strain determined for each species. For example, the critical
classification problem of *P. putida* has been recently reported by us[17]. The *P. putida* NBRC 14164[T]
type strain forms an isolated community in the network with only 15 genomes. On the other
hand, the community of *P. alloputida* Kh7[T] harbors 69 genomes, constituting the largest
community in the *P. putida* group. Thus, most of the genomes deposited as *P. putida* are actually
from *P. alloputida*. Regarding the misclassification of *P. stutzeri*, 122 genomes fall into the
community represented by *Pseudomonas spp7*, a potentially new genomospecies mentioned
above.
We also assessed the impact of our approach defining the species-level taxonomy of the
1,083 non-classified *Pseudomonas* genomes available in Genbank (*Pseudomonas sp.*).
Interestingly, 511 *Pseudomonas sp.* genomes (47.18%) were distributed among 97 communities

 containing type strains (Table S5). The species that received the most genomes were *P. glycinae*
183 (n = 35), *P. lactis* (n = 34), and *P. mandelii* (n = 31).

185
186 **Figure 3. *Pseudomonas* community sizes.** Dark and light purple dots represent communities with and
187 without type strains, respectively. The names and number of genomes are displayed in those communities
188 with more than 100 genomes. y-axis is in log scale.

189
190

The *Pseudomonas* phylogeny reveals at least fourteen groups
To reduce the influence of overrepresented species, we used the 573 representative genomes
from each community to retrieve orthologous genes and reconstruct the *Pseudomonas*
phylogeny. The *Cellvibrio japonicus* Ueda 107[T] was used as an outgroup. We identified 31,094
orthogroups, of which 168 were present in all species, including 30 single-copy genes. We used
the single-copy genes to reconstruct the *Pseudomonas* phylogeny and identify the main
Pseudomonas groups (Figure 4).
The main *Pseudomonas* groups have been previously characterized using housekeeping
genes such as 16S rDNA, *gyrB*, *rpoB*, and *rpoD* from type strains[8, 16]. To delineate each group, we
retrieved those representative genomes (species) within previously-described groups (Table S6).
We then tracked the Most Recent Common Ancestor (MRCA) for those species in the
Pseudomonas phylogenetic tree to include uncharacterized representative genomes as well. For
example, the *P. lutea* group comprises three known species: *P. abietaniphila*, *P. graminis*, and *P.*
lutea[8]. By tracking the corresponding MRCA node, we ensured the monophyly and included *P.*
bohemica and 12 uncharacterized species in this group (Table S2). This approach allowed a more
accurate characterization of both recently described type strains and other uncharacterized
species (Figure 4, Table S2). We identified the 13 main *Pseudomonas* groups and one new group
with 10 genomes and three type strains: *P. alcaligenes*, *P. fluvialis*, and *P. pohangensis* (Figure
4, Table S7). Since *P. alcaligenes* is the firstly-described type strain in this group[23], we named
this group as *P. alcaligenes* group.

211
Figure 4. Phylogenetic tree mapping *Pseudomonas* groups. Maximum-likelihood phylogenetic tree using
core single-copy genes in representative genomes from 573 communities detected in the *Pseudomonas*
network. Colors indicate *Pseudomonas* groups. The number of genomes in each group is in parenthesis.
The asterisk highlights the *P. alcaligenes* group described here. The outgroup is *Cellvibrio japonicus* Ueda
107[T].

Lineage and genus boundaries
The genus *Pseudomonas* has three recognized lineages: *P. pertucinogena*, *P. aeruginosa*, and *P.*
fluorescens. The *P. pertucinogena* lineage is composed of a single phylogenetic group. The *P.*
aeruginosa lineage comprises 6 phylogenetic groups (*P. oryzihabitans*, *P. stutzeri*, *P. oleovorans*,
P. resinovorans, *P. aeruginosa,* and *P. linyingensis*). The *P. fluorescens* lineage also comprises 6
phylogenetic groups (*P. fluorescens*, *P. lutea*, *P. syringae*, *P. putida*, *P. anguilliseptica*, and *P.*
straminea); the *P. fluorescens* group is further divided into 8 or 9 phylogenetic subgroups[16]. In
this work, 70.38% of the communities (species) belong to the *P. fluorescens* lineage, 16.72% to
P. aeruginosa, and 4.52% to *P. pertucinogena*; 8.36% were unclassified communities. We
observed that, unlike the *P. pertucinogena* and *P. fluorescens* lineages, the *P. aeruginosa* lineage
is polyphyletic (Figure 5a).
We used the Genome Taxonomy Database (GTDB) approach[13] to evaluate whether
Pseudomonas should be divided into different genera. The GTDB proposes a framework to
classify genomes in higher taxonomic ranks (e.g. genus). By using the GTDB classification,
Pseudomonas should be divided into 17 genera named generically with "Pseudomonas"
followed by a letter (e.g. "*Pseudomonas_A*"), with the *P. aeruginosa* group retaining the name
Pseudomonas. We found a high correspondence between *Pseudomonas* groups and the
proposed genera, with few inconsistencies (Figure 5a, Table S7). According to the GTDB

classification, the *P. fluorescens* lineage, together with the *P. oleovorans* group and the here
described *P. alcaligenes* group, would form a single genus called *Pseudomonas_E* (Figure 5a),
which corresponds to 77.52% of the species (communities) estimated in our study.
239

Figure 4. *Pseudomonas* phylogenetic tree with proposed genus boundaries and Percentage of Conserved Proteins
(POCP) network. a) Phylogenetic tree annotated with *Pseudomonas* lineages. The outer letters indicate the
annotation adopted by the Genome Taxonomy Database (GTDB). The genus proposed to keep the name
Pseudomonas is marked with a red asterisk. Other genera proposed by GTDB adopt the nomenclature "*Pseudomonas*"
followed by a letter (e.g. *Pseudomonas_E*); for clarity, only the letters and those proposed genera with more than five
communities are displayed. b) Network based on POCP index using a 60% threshold. Colors represent lineages. Blue
nodes embedded in the component with genomes of *Pseudomonas aeruginosa* lineage belong to the groups *P.*
anguilliseptica and *P. straminea*; these two groups are marked in the phylogenetic tree with black asterisks.

We also used the Percentage of Conserved Proteins (POCP) index to evaluate the
relationships between lineages (Figure 5b) and complement the GTDB approach. Briefly, the
POCP index measures the proportion of shared proteins between two genomes[24]. The original
proposal is that genomes belong to the same genus if they share at least half of their proteins[24].
By using 50% as a threshold, we observed that only the outgroup *C. japonicus* and other four
genomes do not belong to the main POCP network component with all lineages. However, by
using a 60% threshold to link communities, we observed two main clusters (Figure 5b).
Apart from *P. anguilliseptica* and *P. straminea* groups, the *P. fluorescens* lineage forms
an isolated component in the network (Figure 5b). The *P. pertucinogena* and *P. aeruginosa*
lineages are in the same component, but linked by a few connections, including a bridge via a *P.*
caeni genome. The outgroup *C. japonicus* is an orphan in the network, as well as *P. kirkiae*. The
species *P. boreopolis*, *P. cissicula*, and *P. geniculata* were also isolated. These three species have
already been recognized as belonging to the genus *Xanthomonas*[25]. Yet, they remain classified
as *Pseudomonas* in Genbank and still labeled as validly published with a correct name in LPSN.
265

DISCUSSION

The *Pseudomonas* genus underwent several taxonomic reclassifications over the years. Here,
we used 10,035 *Pseudomonas* genomes to estimate the genus diversity through network
analysis and community detection. We observed that several type strains are later synonyms
and should be officially revised, as also noted elsewhere [8, 16].

Regarding the *Pseudomonas* network, we observed that the number of detected
communities is very close to the number of network components at a 95% identity threshold.
Combined with the stabilization of density and transitivity around this threshold, we conclude
that the *Pseudomonas* network is highly structured. This structured network profile has also
been noted previously reported for the *P. putida* group[17].

Considering each community as a different genomospecies, we identified 573
communities, way more than the 233 *Pseudomonas* species with validly published names.
Moreover, we found 61 orphan type strains in the network, indicating that the diversity
estimated using only type strains is highly underestimated. In addition, this work shows that
25.65% of the *Pseudomonas* genomes are misclassified. This is a matter of concern, as
misclassified genomes in public repositories can introduce noise to pangenome studies, reduce
strain typing accuracy, and propagate labeling errors to several studies, including those
reporting the characterization of new species.

Here, we also showed potential new genomospecies. For example, the community
assigned as *Pseudomonas spp7* contains 122 genomes, and it is a sister group of *P. stutzeri*. The
high misclassification rate of *P. stutzeri* (Table S4) can be explained by the presence of this new
closely related species. Such inconsistencies could be mitigated through a standardized
taxonomic framework, as previously proposed[26]. However, there is still resistance to define
species based solely on genome sequences, even with the massive number of available
genomes[18]. Therefore, isolating and characterizing members from *Pseudomonas spp7*
community will allow the consolidation of this new species.

Although previous works provided insights about what would be considered
Pseudomonas [8, 16, 27], how to delimit the *Pseudomonas* genus remains an open question. We
tried to address this problem by using GTDB classification and POCP index network, two
approaches proposed to delimit genera. The GTDB results indicate that the *P. fluorescens* lineage
and the *P. oleovorans* and *P. alcaligenes* groups would constitute a genus with the generic name
Pseudomonas_E (Figure 4). However, the POCP index network at 60% shows that *P. straminea*
and *P. anguilliseptica* groups are closer to *P. aeruginosa* than to *P. fluorescens* lineage (Figure
4b). Aiming for a parsimonious separation, we propose that the *P. fluorescens* lineage, excluding
the *P. straminea* and *P. anguilliseptica* groups, should be considered a new genus. Furthermore,
by the GTDB results, the *Pseudomonas* groups from *P. aeruginosa* lineage should also be revised
to assess whether they are new genera, as the *P. aeruginosa* lineage itself is polyphyletic.
Prioritizing the GTDB approach here should provide the best approach because it normalizes
taxonomic ranks and ensures group monophyly[13].

CONCLUSION

In this study, we estimated the *Pseudomonas* diversity using a network approach. We show that type strains represent less than half of the estimated number of species, and that many of them are orphans in the network. We discovered new genomospecies and groups, such as *Pseudomonas spp7* and *P. alcaligenes*, respectively. Although genus delineation is somewhat complex, we propose the *Pseudomonas* genus division by combining GTDB classification and POCP index. To fully understand the *Pseudomonas* diversity, it will be important to focus on each group and characterize species from communities without type strains. This study provides a state-of-the-art classification to delimit bacterial species, which we expect to serve as a guide for future studies with *Pseudomonas spp*, reducing the problems caused by misclassified genomes.

METHODS

Dataset collection and annotation

We recovered 11,025 genomes of *Pseudomonas* from Genbank in June 2020. Genome quality was evaluated with BUSCO v4.0.6[28] using the *Pseudomonadales* dataset. We defined completeness as 100% minus the percentage of missing genes, and contamination as the fraction of duplicated genes. Quality was defined as completeness − 5 x contamination[13]. Genomes with more than 400 contigs were removed, and contigs shorter than 500bp were removed from the remaining genomes. We used mash v2.2.2[20] to calculate the pairwise distances between those genomes with quality higher than 80% using sketches of 1000 and 5000. Regarding the type strains, we used all species with available genomes and validated taxonomic names according to the LPSN[10] on March 2021. The pairwise distances between type strains were performed using pyani v0.2.10[29]. We reannotated the genomes with prokka v1.14[30] to allow a systematic large-scale genome comparison.

Network analysis

By using the pairwise Mash distances, we generated the corresponding graph and obtained the topological graph properties such as density, transitivity, and number of components with the igraph package[31]. We used the label propagation algorithm to detect communities[21]. The representative genome for each community was defined based on three conditions: i) if the community has only one type strain, the type strain was considered the representative genome; ii) if the community has more than one type strains, the first described type strain was chosen; iii) else, we randomly chose a genome in a community (seed = 1996) and assigned the community name with the notation *Pseudomonas sppX*, where *X* is the community number.

Phylogeny and POCP index

We used OrthoFinder v2.5.2[32] to obtain the orthogroups from community type genomes. All single-copy genes were aligned with MAFFT v7.467[33] and concatenated to reconstruct the *Pseudomonas* phylogeny with IQ-TREE v2.1.2[34]. The best-fit model detected through ModelFinder[35] was LG+F+I+G4. One thousand bootstrap replicates were generated to assess the

significance of internal nodes. Phylogenetic trees were visualized and annotated using ggtree[36].
We tracked MRCA nodes for *Pseudomonas* groups definitions using treeio[36].

The Percentage of Conserved Proteins (POCP) between two genomes were calculated
using the formula $\frac{C_1+C_2}{T_1+T_2}$, where *C* is the number of conserved proteins and *T* is the total number
of proteins[24]. The number of conserved proteins was obtained from the orthologs matrix A_{ij}
generated by OrthoFinder, where each entry (i, j) is the total number of genes in species *i* that
have orthologues in species *j*. The graphs were generated and visualized using igraph[31] and
ggnetwork v0.5.8[37], respectively. The GTDB classification was obtained in April
2021 (http://gtdb.ecogenomic.org/).
355

DECLARATION OF COMPETING INTEREST

357   The authors declare no conflict of interest.
358

ACKNOWLEDGEMENTS

360   This work was supported by Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do
361   Rio de Janeiro (FAPERJ; grants E-26/203.309/2016 and E-26/203.014/2018), Coordenação de
362   Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES; Finance Code 001), and Conselho
363   Nacional de Desenvolvimento Científico e Tecnológico. The funding agencies had no role in the
364   design of the study and collection, analysis, and interpretation of data and in writing.
365

REFERENCES

367   1.   Coutinho, F.H. *et al.* Niche distribution and influence of environmental
368   parameters in marine microbial communities: a systematic review. *PeerJ* **3**,
369   e1008 (2015).
370

371   2.   Layeghifard, M. *et al.* Microbiome networks and change-point analysis reveal key
372   community changes associated with cystic fibrosis pulmonary exacerbations. *NPJ*
373   *Biofilms Microbiomes* **5**, 4 (2019).
374

375   3.   Silby, M.W., Winstanley, C., Godfrey, S.A., Levy, S.B. & Jackson, R.W.
376   Pseudomonas genomes: diverse and adaptable. *FEMS Microbiol Rev* **35**, 652-680
377   (2011).
378

379   4.   Palleroni, N.J. The Pseudomonas story. *Environ Microbiol* **12**, 1377-1383 (2010).
380

381   5.   Palleroni, N.J. Genus I. Pseudomonas Migula 1894. *Bergey's Manual of*
382   *Systematic Bacteriology* **1**, 59 (1984).
383

384 6. Ait Tayeb, L., Ageron, E., Grimont, F. & Grimont, P.A. Molecular phylogeny of the
385 genus Pseudomonas based on rpoB sequences and application for the
386 identification of isolates. *Res Microbiol* **156**, 763-773 (2005).
387

388 7. Mulet, M. *et al.* Concordance between whole-cell matrix-assisted laser-
389 desorption/ionization time-of-flight mass spectrometry and multilocus sequence
390 analysis approaches in species discrimination within the genus Pseudomonas.
391 *Syst Appl Microbiol* **35**, 455-464 (2012).
392

393 8. Gomila, M., Pena, A., Mulet, M., Lalucat, J. & Garcia-Valdes, E. Phylogenomics
394 and systematics in Pseudomonas. *Front Microbiol* **6**, 214 (2015).
395

396 9. Peix, A., Ramirez-Bahena, M.H. & Velazquez, E. The current status on the
397 taxonomy of Pseudomonas revisited: An update. *Infect Genet Evol* **57**, 106-116
398 (2018).
399

400 10. Parte, A.C., Sarda Carbasse, J., Meier-Kolthoff, J.P., Reimer, L.C. & Goker, M. List
401 of Prokaryotic names with Standing in Nomenclature (LPSN) moves to the DSMZ.
402 *Int J Syst Evol Microbiol* **70**, 5607-5612 (2020).
403

404 11. Gomila, M., Busquets, A., Mulet, M., Garcia-Valdes, E. & Lalucat, J. Clarification
405 of Taxonomic Status within the Pseudomonas syringae Species Group Based on
406 a Phylogenomic Analysis. *Front Microbiol* **8**, 2422 (2017).
407

408 12. Tran, P.N., Savka, M.A. & Gan, H.M. In-silico Taxonomic Classification of 373
409 Genomes Reveals Species Misidentification and New Genospecies within the
410 Genus Pseudomonas. *Front Microbiol* **8**, 1296 (2017).
411

412 13. Parks, D.H. *et al.* A standardized bacterial taxonomy based on genome phylogeny
413 substantially revises the tree of life. *Nat Biotechnol* **36**, 996-1004 (2018).
414

415 14. Bobay, L.M. The Prokaryotic Species Concept and Challenges. In: Tettelin, H. &
416 Medini, D. (eds). *The Pangenome: Diversity, Dynamics and Evolution of
417 Genomes*: Cham (CH), 2020, pp 21-49.
418

419 15. Lalucat, J., Mulet, M., Gomila, M. & Garcia-Valdes, E. Genomics in Bacterial
420 Taxonomy: Impact on the Genus Pseudomonas. *Genes (Basel)* **11** (2020).
421

422 16. Hesse, C. *et al.* Genome-based evolutionary history of Pseudomonas spp. *Environ
423 Microbiol* **20**, 2142-2159 (2018).
424

425 17. Passarelli-Araujo, H., Jacobs, H.S., Franco, G.R. & Venancio, T.M. Phylogenetic
426 and population structure analyses uncover pervasive misclassification and help
427 assessing the biosafety of Pseudomonas alloputida for biotechnological
428 applications. *bioRxiv* (2021).
429

430   18.   Hugenholtz, P., Chuvochina, M., Oren, A., Parks, D.H. & Soo, R.M. Prokaryotic
431   taxonomy and nomenclature in the age of big sequence data. *ISME J* **15**, 1879-
432   1892 (2021).

434   19.   Backurs, A. & Indyk, P. Edit distance cannot be computed in strongly
435   subquadratic time (Unless SETH is False). *SIAM J Comput* **47**, 10 (2015).

437   20.   Ondov, B.D. *et al.* Mash: fast genome and metagenome distance estimation
438   using MinHash. *Genome Biol* **17**, 132 (2016).

440   21.   Raghavan, U.N., Albert, R. & Kumara, S. Near linear time algorithm to detect
441   community structures in large-scale networks. *Phys Rev E Stat Nonlin Soft Matter*
442   *Phys* **76**, 036106 (2007).

444   22.   Schoch, C.L. *et al.* NCBI Taxonomy: a comprehensive update on curation,
445   resources and tools. *Database (Oxford)* **2020** (2020).

447   23.   Monias, B.L. Classification of Bacterium alcaligenes pyocyaneum and
448   fluorescens. *The Journal of Infectious Diseases* **43**, 330-334 (1928).

450   24.   Qin, Q.L. *et al.* A proposed genus boundary for the prokaryotes based on
451   genomic insights. *J Bacteriol* **196**, 2210-2215 (2014).

453   25.   Anzai, Y., Kim, H., Park, J.Y., Wakabayashi, H. & Oyaizu, H. Phylogenetic affiliation
454   of the pseudomonads based on 16S rRNA sequence. *Int J Syst Evol Microbiol* **50**
455   **Pt 4**, 1563-1589 (2000).

457   26.   Hugenholtz, P., Chuvochina, M., Oren, A., Parks, D.H. & Soo, R.M. Prokaryotic
458   taxonomy and nomenclature in the age of big sequence data. *ISME J* (2021).

460   27.   Ozen, A.I. & Ussery, D.W. Defining the Pseudomonas genus: where do we draw
461   the line with Azotobacter? *Microb Ecol* **63**, 239-248 (2012).

463   28.   Seppey, M., Manni, M. & Zdobnov, E.M. BUSCO: Assessing Genome Assembly
464   and Annotation Completeness. *Methods Mol Biol* **1962**, 227-245 (2019).

466   29.   Pritchard, L., Glover, H.R., Humphris, S., Elphinstone, J.G. & Toth, I.K. Genomics
467   and taxonomy in diagnostics for food security: soft-rotting enterobacterial plant
468   pathogens. *Anal. Methods* **8**, 10-24 (2016).

470   30.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**,
471   2068-2069 (2014).

473   31.   Csardi, G. & Nepusz, T. The igraph software package for complex network
474   research. *InterJournal* **Complex Systems**, 1695 (2006).

476   32.   Emms, D.M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for
477   comparative genomics. *Genome Biol* **20**, 238 (2019).
478

479   33.   Katoh, K. & Standley, D.M. MAFFT multiple sequence alignment software version
480   7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-780 (2013).
481

482   34.   Minh, B.Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
483   Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020).
484

485   35.   Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A. & Jermiin, L.S.
486   ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat
487   Methods* **14**, 587-589 (2017).
488

489   36.   Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc
490   Bioinformatics* **69**, e96 (2020).
491

492   37.   Briatte, F. ggnetwork: Geometries to Plot Networks with 'ggplot2'. R package
493   version 0.5.8.; 2020.

SUPPLEMENTARY FIGURES

Network analysis of ten thousand genomes shed light on *Pseudomonas* diversity and classification

Hemanoel Passarelli-Araujo[1,2,*], Glória Regina Franco[1], Thiago M. Venancio[2,*]

[1]Departamento de Bioquímica e Imunologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

[2]Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, RJ, Brazil.

*Corresponding authors

Av. Alberto Lamego 2000, P5 sala 217; Parque Califórnia
Campos dos Goytacazes, RJ, Brazil
CEP: 28013-602
HPA: hemanuel.passarelli@gmail.com; TMV: thiago.venancio@gmail.com

Figure S1. BUSCO estimation for completeness and contamination for all *Pseudomonas* genomes. a) Distribution for all 11,025 *Pseudomonas* genomes. b) Genomes used in this study after discarding genomes based on 80% quality threshold and fragmentation higher than 400 scaffolds (see methods).

Figure S2. Mash distance statistics. a) Comparison of estimated Mash distance using sketches sizes of 1000 and 5000. b) Mash distances restricted to the interval [0.0, 0.3] in both axes. c) Mash distance distribution for each sketch size.

Figure S3. Genome size distribution for *Pseudomonas* communities. Maximum-likelihood phylogenetic tree using core single-copy genes in representative genomes from 573 communities detected in the *Pseudomonas* distance network. Colors indicate the genome size distribution.