Cyclin-G-associated kinase GAK/dAux regulates autophagy initiation via ULK1/Atg1 in glia

Shiping Zhang[1][†], Shuanglong Yi[1,5,†], Linfang Wang[1,5,†], Shuhua Li[1], Honglei Wang[1],

Li Song[2], Jiayao Ou[2], Min Zhang[1], Ruiqi Wang[1], Mengxiao Wang[1], Yuchen Zheng[1],

Kai Yang[3], Tong Liu[4], Margaret S. Ho[1,*]

[1]School of Life Science and Technology, ShanghaiTech University, Shanghai, 201210, China

[2]Department of Anatomy and Neurobiology, Tongji University School of Medicine, Shanghai, 200092, China

[3]International Academic Center of Complex Systems, Advanced Institute of Natural Sciences, Beijing Normal University at Zhuhai, 519087, China

[4]Institute of Neuroscience, State Key Laboratory of Neuroscience, CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai, 200031, China

[5]University of Chinese Academy of Sciences, Beijing, 100049, China

[†]These authors contributed equally

[*]Corresponding author: Margaret S. Ho, Ph.D. Email: margareth@shanghaitech.edu.cn

Abstract

Autophagy is a major means for the elimination of protein inclusions in neurons in neurodegenerative diseases such as Parkinson's disease (PD). Yet, the mechanism of autophagy in the other brain cell type, glia, is less well characterized and remains largely unknown. Here we present evidence that the PD risk factor Cyclin G-associated kinase (GAK)/dAuxilin (dAux) is a new component in glial autophagy. GAK/dAux directly interacts with ULK1/Atg1 via its uncoating domain. Lack of GAK/dAux increases the autophagosome number and size in adult fly glia and mouse microglia, and generally upregulates levels of components in the initiation and PI3K class III complexes including ULK1/Atg1, demonstrating that GAK/dAux regulates the onset of glial autophagy. Consistently, lack of GAK/dAux enhances Atg1 and Atg9 trafficking to autophagosomes, promoting autophagy initiation. On the other hand, lack of GAK/dAux impairs the autophagic flux and blocks substrate degradation, suggesting that GAK/dAux might play additional roles in glial autophagy. Our findings identify a new autophagy factor in glia; considering the pivotal role of glia under pathological conditions, targeting glial autophagy is potentially a therapeutic strategy for PD.

Keywords: Autophagy; Glia; GAK; dAux; Parkinson's disease

Introduction

As the primary catabolic mechanism, macroautophagy (henceforth referred to as autophagy) was initially characterized as a bulk and non-selective degradation pathway induced by nutrient deprivation (1-3). Constitutive autophagy at the basal level, however, contributes to intracellular homeostasis in non-starved cells by degrading cellular materials like damaged organelles or misfolded proteins (4, 5). Interestingly, suppression of basal autophagy in the mouse nervous system results in the accumulation of ubiquitinated proteins and neurodegeneration (6, 7), whereas enhancing basal autophagy in the *Drosophila* nervous system promotes longevity and oxidant resistance (8). Furthermore, disease-associated mutations in core autophagy genes have been identified, supporting the link between autophagy and PD (9, 10). These findings mainly focus on a role for autophagy in neurons, as perturbation of such process leads to catastrophic consequences including pathological protein aggregation, deleterious lesions, and cell death (11). Whether autophagy in glia, the other major brain cell type, is executed via similar interplay using the same components as in neurons or other tissues, and whether it drives or contributes to disease pathology, remain largely unclear.

Studies from yeast and mammals provided fundamental insights into *de novo* autophagosome biogenesis mediated by the evolutionarily conserved autophagy-related (ATG) genes (4, 5, 12). Upon autophagy initiation, stress induces the recruitment of ATG1/UNC-51 like autophagy activating kinase 1 (ULK1) complex to the phagophore assembly site or the pre-autophagosomal structure (PAS). The ATG1/ULK1 complex

facilitates the assembly of phosphatidylinositol 3-kinase (PI3KC3) class III complex

for PI3P production and promote the nucleation of the isolated membranes (also termed

phagophores) (13). Whereas phagophore expansion is achieved by two ubiquitin-like

conjugation steps: the formation of ATG12-ATG5 conjugate and ATG8 lipidation,

additional lipids and proteins contributing to membrane expansion are delivered by the

ATG9-containing vesicles derived from the Golgi apparatus, recycling endosomes or

the plasma membrane (14-18). The expanded phagophores are sealed into a double-

layered autophagosome, which eventually fuses with lysosome to form autolysosome,

the milieu for degrading autophagic cargoes by acidic hydrolases.

Here we present evidence that the PD risk factor Cyclin G-associated kinase

(GAK)/dAuxilin (dAux) is a new component in glial autophagy. GAK/dAux interacts

with ULK1/Atg1 via the J domain with uncoating activity. GAK/dAux regulates the

protein levels of ULK1/Atg1 and other components in the initiation and PI3K

complexes controlling autophagy initiation. *In-vivo*, Lack of GAK/dAux increases the

autophagosome number and size, and regulates the trafficking of ULK1/Atg1 and Atg9

to the autophagosomes. On the other hand, lack of GAK/dAux impairs autophagic flux

and substrate degradation. Taken together, our findings identify a key factor regulating

glial autophagy via controlling the activity of the initiation regulator ULK1/Atg1.

Results

Lack of dAux selectively increases the number of autophagosomes in adult fly glia

Identified in a behavior screen, the *Drosophila* homolog of the PD risk factor GAK,

dAux, has been implicated in a broad spectrum of parkinsonian symptoms (19).

GAK/dAux is a serine/threonine kinase that regulates clathrin-uncoating, cell cycle

progression, and hepatitis C entry (20-23). Whereas another closely-related homolog

for *GAK* (*DNAJC26*), *AUXILIN* (*DNAJC6*), is dominantly expressed in neurons, there

is only one *Drosophila* homolog *dauxilin* (*daux*) that exhibits a higher homology with

GAK (GAK 41.66%, AUXILIN 34.52%). GAK and dAux are both ubiquitiously

expressed and shares a conserved kinase domain. To investigate dAux function at the

cellular level, an RNAi line targeting *Drosophila daux* (*daux*-RNAi, V16182) was

expressed to reduce *daux* expression in neurons or glia by the driver *GMR57C10-Gal4*

or *repo-Gal4*, respectively. qRT-PCR, western blot (WB), and immunostaining analyses

indicated that *daux* expression was efficiently reduced by the RNAi (Figures S1A-S1D).

Under these conditions, the morphology, distribution, and number of the

autophagic organelles labeled by the marker mCherry.Atg8a were analyzed in different

adult brain regions, including the region "Glia" enriched of glial processes, glial nuclei,

and neuronal nuclei, and the region "Antenna Lobe, AL" enriched of axon neuropils in

close proximity to glial processes (Figures 1A and 1B). Interestingly, downregulating

daux expression by *daux*-RNAi pan-neuronally did not cause severe change in the

number or size of the Atg8a-positive puncta (Figures S1E and S1F). Instead,

downregulating *daux* expression in glia caused a significant increase in the number of

Atg8a-positive puncta (Figures 1C and 1D). These results, confirmed by one additional

daux-RNAi (*daux*-RNAi[#2], BL39017), an endogenous protein trap *pmCh.Atg8a*, anti-

Atg8a antibodies, and Mosaic Analysis with a Repressible Cell Marker (MARCM)

clones of *daux* mutant carrying EMS point mutations *I670K*, *L8H*, or *G257E* (24)

(Figures S1G, S1H, and 1E-1J,), indicated an increase in glial autophagic structures in

the absence of dAux. Results from using a different control *UAS-luciferase*-RNAi (*luc*-

RNAi) were also consistent (Figures S1I and S1J). Ultrastructural analysis by

Transmission Electron Microscopy (TEM) also showed an increase in the number of

autophagosomes, which exhibits typical double-membrane structure with a

characteristic cleft in-between and contains non-degraded cytoplasm, upon *daux*-RNAi

expression in glia (pink asterisks, Figures 1K and 1L). Moreover, whereas the Rab7-

positive late endosomes appeared in clusters with puncta number unaffected (Figures

S1K and S1L), the formation of the Rab5-positive early endosomes remained largely

intact upon glial dAux depletion (Figure S1M). Notably, expression of the fly dAux or

human GAK in glia rescued the *daux*-RNAi-induced increase in the autophagosome

number (Figures 1O and 1P). Taken together, these results indicate that dAux regulates

glial autophagosome number and the function of GAK/dAux is conserved across

species.

Lack of Gak increases the number and size of autophagosomes in mouse microglia

In addition, the mouse immortalized microglial cell line (IMG) and the primary mouse

microglia were examined for the effect of lacking Gak on autophagic structures. The

IMG cells were stained positive for ionized calcium-binding adapter molecule 1 (Iba1)

or transmembrane protein 119 (TMEM119), confirming their microglial identity

(Figures S2A and S2B). Consistent with the previously reported expression in microglia

and astrocytes (25, 26), Gak was also detected in the IMG cells (Figures S2C-S2G) (27).

Treatment with either of the two siRNAs targeting *Gak* reduced its expression in IMG

as shown by qRT-PCR, WB, and immunostaining analyses (Figures S2C-S2G). *Gak*

mRNA and protein expression were also reduced down to 50% and 60%, respectively

in *Gak* conditional knockout (cKO) mouse microglia (Figures S2H-S2J). We failed to

collect *Gak* cKO primary microglia for WB analysis of Gak protein levels due to

inefficient growth.

In IMG cells, treatment of either siRNA increased autophagosome size labeled by

the microtubule-associated protein 1A/1B-light chain 3 (LC3) (Figures 1Q and 1R).

Both the intensity and size of LC3 in microglia, and the LC3-Iba1 colocalization

increased in *Gak* cKO microglia, suggesting increased autophagosome number in

microglia (Figures 1S and 1T). Interestingly, the size of autophagosome consistently

increases in the IMG, *Gak* cKO microglia, and fly *daux* mutant MARCM clones, but

not *daux*-RNAi brains, possibly owing to the incomplete knock down by RNAi (Figures

1D, 1J, 1R, and 1T). Taken together, these results suggest that Gak regulates

autophagosome biogenesis in mouse microglia.

Lack of GAK/dAux promotes autophagy initiation

Based on the increase in the autophagosome number, it is conceivable that lack of glial

dAux induces or blocks autophagy. Interestingly, Atg8a-II protein levels in adult fly

brains downregulating glial dAux expression increased, which further increased in the

presence of Bafilomycin A1 (BafA1), an inhibitor blocking lysosome acidification and

autophagosome-lysosome fusion (28, 29) (Figures 2A and 2B). These results indicate

that lack of glial dAux still triggers increase in the autophagosome number even when

blocking the fusion, reinforcing a role for dAux in autophagy initiation. Using the anti-

ULK1 antibodies that also recognize the *Drosophila* Atg1 proteins as validated by *atg1-*

RNAi (Figures S3A and S3B), an increase in the number of ULK1-positive puncta was

detected upon glial dAux depletion (Figures 2C and 2D). In addition, components of

the initiation complex including Atg13, Atg6, and Atg14 were all affected as the number

of mCherry.Atg13-, GFP.Atg6-, and GFP.Atg14-positive puncta increased upon glial

dAux depletion (Figures 2C-2F). Notably, omegasomes labeled by two independent

reporters: GFP.ZFYVE1 (12, 30) and EGFP.DFCP1 (31), exhibited significant increase

in number upon glial *daux*-RNAi expression, with the size of EGFP.DFCP1-positive

omegasomes became significantly larger as fused by multiple smaller rings (Figures 2G

and 2H). Taken together, these results demonstrate the increased formation of

autophagosome precursors in the absence of glial dAux.

To better understand the mechanism of dAux-mediated autophagy initiation, we

turned to a different system, HeLa cells, for further characterization. A knockdown

system was established using two independent siRNAs targeting *GAK* (siRNA[#1] and

siRNA[#2]). Treatment of either siRNA reduced *GAK* expression in HeLa cells as shown

by WB and immunostaining analyses (Figures S3C-S3F). Interestingly, ULK1, ATG13,

and BECLIN1 protein levels significantly increased while ATG14 protein levels

exhibited an increasing trend upon *GAK* siRNA treatment, suggesting a general

upregulation in autophagy initiation (Figures 2I and 2J). In addition, the mTORC1-

mediated phosphorylation on the ULK1 Serine 757 (pS757-ULK1) decreased,

suggesting that the ULK1 activity is enhanced as mTORC1 inhibition is relieved

(Figures 2I and 2J). In addition, a *GAK* KO cell line was created by silencing the *GAK*

expression in HeLa cells using CRISPR-Cas9 (Figures S3G-S3J). Consistent to the

observations in flies, lack of GAK causes an increase in LC3-II protein levels in both

GAK siRNA-treated and KO cells (Figures 2K and 2L, S3K and S3L). LC3-II protein

levels also further increased in the presence of BafA1 in *GAK* siRNA-treated cells,

reinforcing a direct role of GAK in autophagy initiation (Figures 2K and 2L). Taken

together, these results indicate that autophagy initiation is induced in the absence of

GAK/dAux; GAK/dAux regulates the onset of glial autophagy.

GAK/dAux interacts with ULK1/Atg1 and regulates its protein levels

As shown in flies and HeLa cells, lack of GAK/dAux enhances ULK1/Atg1 activity

and induces autophagy initiation. We next tested if ULK1/Atg1 acts downstream of

GAK/dAux in regulating autophagy initiation. Co-immunoprecipitation (Co-IP)

analysis using fly S2 cells expressing Flag-dAux and Myc-Atg1 revealed a direct

interaction between dAux and Atg1 when pulling down Myc-Atg1 with the anti-Myc

beads. Reverse pull down by Flag-tagged dAux also revealed similar interaction (Figure

3A). To determine which dAux domain is responsible for Atg1 interaction, different

dAux deletion variants were made and expressed in flies for testing their interaction

with Atg1. All variants expressed properly except the one lacking PTEN domain

(dAux$^{\Delta PTEN}$, Figures 3B and 3C). Co-IP analysis showed that Atg1 interacts with most

of the dAux variants except the one lacking J domain (dAux$^{\Delta J}$, Figure 3D). In addition,

overexpression of dAux$^{\Delta Kinase}$ or dAux$^{\Delta J}$ enhanced ULK1 levels (Figures 3E and 3F).

Importantly, the early-onset PD SNP mutation in the J domain of AUXILIN (DNAJC6)

(32, 33) is conserved in dAux, and overexpressing the dAux carrying this mutation

dAux^{DNAJC6m} enhanced ULK1 levels (Figures 3E and 3F). These results implicate that

dAux kinase and uncoating function are important for regulating the ULK1/Atg1

protein level.

On the other hand, GAK interacted with ULK1 in HEK293FT cells, but failed to

interact with ULK1 lacking the C-terminal domain (ULK1$^{\Delta CTD}$), suggesting that CTD

is required for GAK-ULK1 interaction (Figures 3G and 3H). Consistent to the enhanced

ULK1 activity, the ULK1 protein level was also elevated in *GAK* siRNA-treated HeLa

cells (Figures 3I-3L). To further test the relationship between dAux and Atg1, epistatic

genetic analysis between dAux and Atg1 was conducted. Silencing *atg1* expression in

glia caused a dramatic decrease in the number of omegasomes and autophagosomes,

reinforcing the master regulatory role of Atg1 in autophagy initiation (Figures 3M-3P).

Co-expression of *atg1*-RNAi and *daux*-RNAi rescued and further suppressed the *daux*-

RNAi-induced increase in the number of omegasome and autophagosome, suggesting

that dAux acts upstream of Atg1 in regulating autophagy initiation (Figures 3M-3P).

Taken together, these results demonstrate that GAK/dAux interacts with ULK1/Atg1

and regulates its protein level.

dAux regulates Atg1 independently of its kinase function

Given that both dAux kinase and J domains are implicated in the Atg1 activation,

phosphoproteomic analysis was subsequently conducted using adult fly head samples

collected from control and *repo>daux*-RNAi flies (Figure S4A). These results revealed

significant differences on the phosphorylation profiles of autophagy-related proteins

(Figure S4B). Nonetheless, no significant changes were detected for Atg1

phosphorylation when downregulating dAux expression in glia. Only minor

alternations on the two Atg1 residues, S282 (RNAi/LacZ ratio 1.131) and T292

(RNAi/LacZ ratio 1.053), were identified (Table S2). In addition, our Co-IP analysis

revealed that dAux$^{\Delta Kinase}$ interacts with Atg1, indicating that the kinase domain is not

essential for the interaction (Figure 3D). Despite that dAux$^{\Delta Kinase}$ expression enhances

ULK1 levels, its impact on autophagy remains similar to dAux when analyzing the

protein levels of Atg8a-II, Ubi, and P62 (Figures S4C-S4F), suggesting that lacking

kinase domain does not hinder dAux function in autophagy. Hence, it is unlikely that

dAux regulates Atg1 in a kinase-dependent manner.

Lack of GAK/dAux promotes ULK1/Atg1 trafficking to autophagosomes

One of the major steps in initiating autophagy is the ULK1/Atg1 transport to the autophagy initiation sites for forming the autophagosomes. We next analyzed the colocalization of ULK1/Atg1 with LC3/Atg8a as an indicator for the ULK1/Atg1 trafficking efficiency. Since an increase in the ULK1/Atg1 protein levels was also detected, we carefully chose the Manders' Correlation method when analyzing the colocalization given that the change in protein level is also considered in this statistical approach (34-36). Consistent to the observations in flies and mice, the number of LC3-positive puncta labeled by the transfected RFP-LC3 increased in the *GAK* siRNA-treated or KO cells (Figures 4A and 4B, S5A and S5B). In addition, the ULK1/Atg1 trafficking to the autophagosomes was enhanced as shown by the increased colocalization of ULK1/Atg1 and LC3/Atg8a in flies expressing *daux*-RNAi, *GAK* siRNA-treated, and KO HeLa cells (Figures 4A-4D, S5A and S5B). In contrast, the colocalization of ULK1 with TGN38-marked trans-Golgi network (TGN) decreased, further suggesting that ULK1 trafficking to the autophagosomes is enhanced when lacking GAK (Figures 4E and 4F, S5C and S5D). In support of these findings, ULK1 trafficking to the omegasomes also increased as the colocalization of ULK1 with transfected GFP-DFCP1 increased in the *GAK* siRNA-treated (Figures S5E and S5F) and KO cells (Figures S5G and S5H). Taken together, these results indicate that GAK/dAux regulates autophagy initiation via ULK1/Atg1 trafficking.

Glial dAux regulates Atg9 trafficking to autophagosomes via Atg1

In addition to promote Atg1 trafficking, lack of dAux increased the glial Atg9-Atg8a

colocalization analyzed by the *mCherry.Atg8a* and *EGFP.Atg9* reporters, indicating that

dAux also regulates Atg9 trafficking to autophagosomes (Figures 5A, 5A', and 5C).

Next, live-cell imaging analysis was conducted, and the dynamic of Atg9 trafficking

was analyzed by Imaris (Figures 5E-H, Supplementary Video 1 and 2). Interestingly,

the speed of both Atg8a- and Atg9-positive puncta decreased upon glial dAux depletion,

suggesting that these puncta are less mobile (Figures 5E and 5H). Notably, the duration

of the Atg9-Atg8a contact was significantly longer, and the percentage of Atg9-positive

puncta being trafficked to the vicinity of Atg8a-positive puncta (with a radius of 0.35

μm) was also significantly greater (Figures 5F and 5G). These results demonstrate that

glial dAux regulates the dynamic of Atg9 trafficking to and its fall off from

autophagosomes.

Consistent to the notion that Atg1 regulates Atg9 phosphorylation and transport

(37-39), lacking Atg1 abolished Atg9 trafficking to the autophagosomes as the Atg9-

Atg8a colocalization decreased significantly (Figures 5A, 5A', and 5C). These results

suggest that dAux and Atg1 regulates Atg9 trafficking in opposite means. Interestingly,

co-expression of *atg1*-RNAi and *daux*-RNAi rescued and further suppressed the dAux-

mediated Atg9 trafficking (Figures 5B, 5B', and 5D), suggesting that dAux regulates

Atg9 trafficking via Atg1. Given that Atg9 is a downstream target of Atg1, it is likely

that dAux-mediated Atg1 activation promotes Atg9 trafficking to the autophagosomes,

hence initiating autophagy.

Lack of GAK/dAux impairs autophagic flux and substrate degradation

To further demonstrate a role for dAux in glial autophagy, the autophagic flux was

analyzed using a dual reporter *UAS-GFP-mCherry-Atg8a* in *Drosophila* adult brains.

Whereas mCherry-positive puncta were GFP-negative and autophagy proceeded

normally in the control animals, autophagic flux was impaired in the absence of glial

dAux as both the GFP- and mCh-positive puncta colocalization and the intensity of

GFP puncta increased significantly upon glial dAux depletion. In the presence of BafA1,

the GFP- and mCh-positive puncta colocalization further increased upon depleting glial

dAux and a similar trend of GFP puncta intensity was detected (Figures 6A and 6B).

As these non-quenched GFP signals could represent either the non-fused

autophgasomes or the fused but dysfunctional autolysosomes, we next checked whether

the fusion of autophagosomes and lysosomes is affected by glial dAux depletion.

Interestingly, analysis of Lamp1-Atg8a colocalization with Stimulated Emission

Depletion (STED) imaging on adult fly brains expressing *daux*-RNAi revealed an

increased number of enlarged and well-colocalized autolysosomes, indicating that

lysosomes are able to fuse with autophagosomes (Figures 6C and 6D). Ultrastructural

TEM analysis also showed an increase in the number of autolysosomes, which have

only one limiting membrane and contain electron dense cytoplasmic materials and/or

organelles, upon *daux*-RNAi expression in glia (blue arrows, Figures 1K and 1N).

Reducing the expression of fusion factors such as *syx17* or *vamp7* caused a block in

fusion (40), and co-expression of either RNAi reverted the *daux*-RNAi-induced

increase in Lamp1-Atg8a colocalization (Figures S6A and S6B). These results suggest

that some successful fusion occurs so that increasing number of enlarged autolysosomes

is detected.

As a consequence, inhibiting dAux expression in glia caused the accumulation of

autophagic substrates such as Ref(2)P (the *Drosophila* ortholog of P62) and Ubiquitin

(Ubi) (Figures 6E-6H). P62 and Ubi were also accumulated in *Gak* siRNA-treated IMG

cells (Figures 6I and 6J), *GAK* siRNA-treated and *GAK* KO HeLa cells (Figures 6K,

6L, S3K, and S3L). Furthermore, increased Ref(2) puncta observed upon *daux*-RNAi

expression in glia were well localized with autophagosomes and lysosomes (Figure

S6C), suggesting that theses fused autolysosomes are able to recognize and encircle

protein substrates, but failed to degrade them. Taken together, these findings suggest

that GAK/dAux regulates glial autophagy via the initiation step but with an ultimate

effect on impaired flux and substrate degradation (Figure 6M).

Discussion

Our findings pinpoint a role for GAK/dAux in autophagy initiation. Despite that the

ultimate consequence for lacking GAK/dAux blocks autophagic substrate degradation

and a potential compensatory mechanism leading to the increased autophagosome

biogenesis might exist, our evidence indicates that GAK/dAux functions directly on

autophagy initiation by modulating ULK1/Atg1 activity and protein level. First, the

dAux-mediated increase in the Atg8a-II/I protein levels is further enhanced by the

presence of BafA1, indicating that autophagy is promoted even when substrate

degradation is blocked. Lack of GAK/dAux also causes an increase in phagophore

nucleation and omegasome formation, suggesting that these earlier steps in autophagy

initiation are all affected. Furthermore, the levels of components in the initiation and

PI3K class III complexes including ULK1/Atg1, ATG13, BECLIN1, and ATG14 all

increase in the absence of GAK, indicating a general upregulation in autophagy

initiation. It has been shown that mTOR senses the nutrient-rich condition and blocks

autophagic induction by phosphorylating ULK1 at S757 to prevent its activation (41).

In the absence of GAK, S757 phosphorylation is significantly decreased, indicating

enhanced ULK1 activity. Thus, GAK/dAux suppresses autophagy initiation; it is

conceivable that GAK/dAux acts as a glial sensor to control the onset of autophagy for

sensing pathological signals like protein inclusions.

ULK1/Atg1 is a master regulator initiating autophagy. In complex with ATG13

and FIP200, ULK1/Atg1 initiates autophagy by phosphorylating a vast array of

downstream components including the factors in the initiation and PI3P class III

complexes (42-44). The ULK1/Atg1 activity undergoes intricate controls by post-

translational modification including phosphorylation, acetylation, or ubiquitination (45,

46). These modifications can either activate or inhibit ULK1/Atg1 activity, thereby

promoting or inhibiting autophagy, respectively. Despite that a direct interaction

between GAK/dAux and ULK1/Atg1 is detected, our phosphoproteomic analysis

reveals no significant alternation on Atg1 phosphorylation upon glial dAux depletion.

In addition, dAux lacking the kinase activity remains associated with Atg1. Thus, it is

unlikely that GAK/dAux regulates ULK1/Atg1 via phosphorylation.

Previous studies reported that inhibiting ULK1 kinase activity, either by ULK1

inhibitors or expressing a kinase-dead ULK1, caused accumulation of stalled early

autophagosomal structures (47, 48). While we observe some phenotypes similar to

stalled early autophagosomes, but with crucial differences. First, instead of lacking

LC3-II flux, we observed increased LC3-II flux with/without BafA1 when

downregulating GAK/dAux both in HeLa cells and fly. Immunostaining of dual

autophagy flux reporter in adult fly brains with BafA1 got similar results. Second, the

accumulated autophagosomes were able to fuse with lysosomes to form autolysosomes

when downregulating glial dAux. Third, we observed substantial co-localization of

autophagosomes, lysosomes, and Ref(2)P, suggesting that theses fused autolysosomes

are able to recognize and encircle protein substrates, but failed to degrade them. Last

but most importantly, we observed decreased S757 phosphorylation in the absence of

GAK, indicating enhanced ULK1 activity.

As we demonstrated that dAux J domain is required for its interaction with Atg1,

it is conceivable that dAux regulates Atg1 via its uncoating function. Expression of

dAux lacking the J domain or dAux carrying a conserved early-onset PD SNP mutation

in the J domain causes an increase in the ULK1 protein level. These results implicate

the relevance of dAux clathrin-uncoating activity with ULK1/Atg1. In addition, it is

also possible that GAK/dAux contributes to the recycling of ULK1/Atg1 from

autophagosomes to TGN, thus leading to similar enhancement on autophagy initiation.

To this end, our live-cell imaging analysis supports the notion that dAux plays dual

roles in the context of Atg9 trafficking to autophagosomes. As increased amount of

Atg9-positive puncta is detected at the vicinity of Atg8a-positive puncta, the contact

duration time between Atg9 and Atg8a is also significantly enhanced in the absence of

dAux. Thus, dAux might be needed for both the anterograde trafficking to and the

recycling from autophagosomes. On the other hand, our results support a competition

model for how GAK initiates autophagy. GAK interacts with the C-terminus of ULK1,

the domain also required for its interaction with Atg13 (49). It is reasonable to

spectulate that GAK competes with Atg13 for ULK1 binding. In the absence of GAK,

increased amounts of ULK1 proteins are relieved from the inhibition for interaction

with Atg13 and together trafficked to the omegasomes/autophagosomes for autophagy

initiation.

Despite an effect on autophagy initiation, autophagic flux and substrate

degradation were impaired in the absence of glial dAux. Recently, two independent

studies reported the involvement of GAK in autophagy and lysosomal homeostasis.

Munson et. al. reported that GAK inhibition alters lysosomal morphology in U2OS cells

(50). After treatment of GAK kinase inhibitor, increased lysosome numbers and large

autolysosome structures were observed, possibly mediated by the increased TFEB

nuclear localization (50). Miyazaki et. al. found that *GAK* KO or GAK kinase inhibitor

treatment could induce autophagosome accumulation and enlarged autolysosomes in

A549 cells, which were caused by impaired autophagosome-lysosome fusion and

autophagic lysosome reformation (ALR) (51). Given that we demonstrated that the

fusion can still occur, it is possible that lysosome function is affected so that flux and

substrate degradation are impaired upon glial dAux depletion. Consistent to this notion,

the number and size of glial Lamp1-positive lysosomes were increased in adult fly

brains expressing the *daux*-RNAi (Figures S6D and S6E). Ultrastructural TEM analysis

also showed an increase in the number and size of lysosome, which exhibits smaller

and electron dense structures with single-membrane, upon *daux*-RNAi expression in

glia (yellow and red arrows, Figures 1K and 1M). Thus, it is feasible that GAK/dAux

regulates autophagy at multiple steps via different mechanisms.

In the present study, we identify a new autophagy factor in glia and elucidate its

regulatory mechanism on autophagy initiation. Our findings provide new insights on

glial autophagy and advance the understanding of PD pathology from a glial

perspective. Future work will be required to investigate how GAK/dAux plays different

roles in glia and neurons, and how it contributes to neurodegeneration in a tissue-

specific manner.

Materials and Methods

All animal experiments were conducted according to proper regulation and guideline

under ShanghaiTech. Details of fly and mouse genotypes are included in SI Appendix

and Table S1. Methods including cell lines and transfections, molecular cloning,

immunohistochemistry, *in-vivo* time lapse analysis, TEM, phosphoproteomic analysis,

biochemistry, confocal microscopy, and statistical analysis are described in the SI

Appendix.

Acknowledgments

We thank Bloomington *Drosophila* Stock Center, Vienna *Drosophila* RNAi Center,

Tsinghua Fly Center, the Core Facility of *Drosophila* Resource and Technology,

Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences,

Developmental Studies Hybridoma Bank, Chao Tong, Henry C. Chang, Lei Xue, Henry

Sun, Tor Erik Rusten, Gabor Juhasz, Patrik Verstreken, Aike Guo, Yijun Liu, and

Gerald M Rubin for fly stocks and antibodies; Chih-Hao Lee for IMG cell line; Yanfeng

Liu for HeLa and HEK293FT cell lines; Kangmin He for HeLa *GAK* KO cell line;

Jiawei Zhou for *CX3CR1-Cre* mice. We also thank the Molecular Imaging Core Facility

(MICF), the Molecular and Cell Biology Core Facility (MCBCF), and the Multi-Omics

Core Facility (MOCF) at the School of Life Science and Technology, ShanghaiTech

University for providing technical support; Yu Kong and Lijun Pan in Electron

Microscopy Facilities of Center for Excellence in Brain Science and Technology,

Chinese Academy of Science for assistance with TEM sample preparation; Wei-Chung

Lee for help on TEM; Ho lab members for discussion and comments. This work was

supported by grants from ShanghaiTech and National Natural Science Foundation of

China (31871039 and 32170962).

Author contributions

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Lack of Gak/dAux increases the number and size of autophagosome in

glia

(**A** and **B**) Illustration (A) and representative images (B) of an adult fly brain showing

the expression pattern of glial cells. *repo-GAL4* drives the expression of *UAS-IVS-*

Syn21-GFP-p10 (*repo>IVS-Syn21-GFP-p10*) that labels both membrane and nucleus

of glia. Dashed squares in yellow and blue indicate the relative positions of "Glia" and

the antenna lobe ("AL") assessed throughout the study. The regions "Glia" are enriched

with GFP-positive glial membrane (green), GFP- and Repo-positive glial nuclei (green

and magenta), as well as Elav-positive neuronal nuclei (red). The regions "AL" are

enriched with axon neuropils and glial processes (green). (**C** and **D**) Representative

images (C) and quantifications (D) of autophagosomes in the Whole brain, "Glia", and

"AL". Glial autophagosomes (red) were labeled with *repo-GAL4* driven *UAS-*

mCherry.Atg8a (*UAS-mCherry.Atg8a; repo-GAL4*). Note that downregulating glial

daux induced an increase in the autophagosome number in both "Glia" and "AL". (**E-**

H) Representative images (E and G) and quantifications (F and H) of the "Glia" region

in adult fly brains carrying an endogenous mCherry protein trap at the promoter of

Atg8a (*pmCh.Atg8a*, E) or stained with the anti-Atg8a antibodies (G). Note that similar

to *UAS-mCh.Atg8a*, the autophagosome number and fluorescent intensities in "Glia"

region increased upon expressing the *daux*-RNAi in glia. (**I** and **J**) Glial MARCM

clones of *daux* mutants ($daux^{L78H}$, $daux^{G257E}$, and $daux^{I670K}$) expressing the *UAS-*

mCh.Atg8a by *repo-GAL4,* and adult fly brain "Glia" region were shown. Dashed lines

encircle the clonal area. Note that the number and size of glial autophagosomes per

clone increased in $daux^{G257E}$ and $daux^{I670K}$ mutants. (**K-N**) Representative TEM images

(K) and quantifications of autophagosome (L), lysosome (M), and autolysosome (N) in

"Glia" region of control and *repo>daux*-RNAi adult fly brains. Note an increase in the

number of autophagosome (pink asterisks), lysosome (yellow arrows) with the presence

of large lysosome-like structure (red arrows), and autolysosome (blue arrows) when

daux-RNAi was expressed in glia. (**O** and **P**) Co-expression of fly dAux or human GAK

(hGAK) rescued the *daux*-RNAi-induced increase in glial autophagosome number in

"Glia" region. (**Q** and **R**) Representative images (Q) and quantifications (R) of LC3-

positive autophagosomes in IMG cells treated with either *Gak* siRNA. Note that LC3-

positive autophagosomes were enlarged upon abolishing *Gak* expression. (**S** and **T**)

Representative images (S) and quantifications (T) of LC3-positive autophagosome in

mouse SN microglia deficient of *Gak* at 36 weeks. In addition to the increase in the

intensity and size of LC3 in Iba1$^+$ microglia, the colocalization of Iba1 with LC3 also

increased upon microglial Gak depletion. Scale bars and the sample number n are

indicated in the Figures. For each experiment, more than three biologically independent

replicates were done. Whereas results were consistent, representative results from one

experiment was shown. A serial confocal Z-stack sections were taken with 0.4 μm each,

and representative single layer images acquired at the similar plane of brains across all

genotypes are shown, except mouse brain slices (S) shown as maximal projection. For

TEM, more than 40 images from at least 5 independent adult fly brains of each genotype

were analyzed. Colocalization is analyzed using the Pearson's Correlation and present

as the Pearson's R value. Quantification figures from the same images but with different

parameters are shown under the same label unless noted otherwise. Data are shown as

mean ± SEM. P-values of significance (indicated with asterisks, ns no significance, *

$p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated by two-tailed unpaired t-test,

Mann-Whitney test, ordinary one-way ANOVA followed by Tukey's multiple

comparisons test, or Kruskal-Wallis tests followed by Dunn's multiple comparisons test.

All genotypes analyzed are listed in Table S1.

Figure 2. Lack of GAK/dAux promotes autophagy initiation

(**A** and **B**) Representative WB images (A) and quantifications (B) of Atg8a-II protein

levels in the control and *repo>daux*-RNAi adult fly brains. Note that lack of glial dAux

increased Atg8a-II protein levels, and BafA1 treatment further increased the Atg8a-II

levels. (**C-H**) Representative images (C, E, G) and quantifications (D, F, H) of ULK1-,

mCh.Atg13-, GFP.Atg6-, GFP.Atg14-, GFP.ZFYVE1-, and EGFP.DFCP1-positive

puncta in "Glia" region of control and *repo>daux*-RNAi adult fly brains. *Drosophila*

Atg1 was detected using the mammal anti-ULK1 antibodies. Note that lack of glial

dAux increased the puncta number of all above, indicating a general upregulation in

autophagy initiation, phagophore nucleation, and omegasome formation. (**I-L**)

Representative WB images (I, K) and quantifications (J, L) of different autophagy

protein levels in *GAK* siRNA-treated HeLa cells. Results from two independent siRNAs,

GAK siRNA[#1] and *GAK* siRNA[#2], were consistent. Note that *GAK* knockdown

significantly increased the protein levels of pS757-ULK1, ATG13, BECN1, and LC3-

II. LC3-II protein level enhanced by *GAK* knockdown was further enhanced in the

presence of BafA1. Scale bars and the sample number n are indicated in the Figures.

For each experiment, more than three biologically independent replicates were done.

Whereas results were consistent, representative results from one experiment was shown.

A serial confocal Z-stack sections were taken with 0.4 μm each, and representative

single layer images acquired at the similar plane of brains across all genotypes are

shown. Colocalization is analyzed using the Pearson's Correlation and present as the

Pearson's R value. Data are shown as mean ± SEM. P-values of significance (indicated

with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated

by two-tailed unpaired t-test, Mann-Whitney test, or ordinary one-way ANOVA

followed by Tukey's multiple comparisons test.

Figure 3. GAK/dAux interacts with ULK1/Atg1 and regulates its protein levels

(**A**) Co-IP analysis using fly S2 cells expressing Flag-dAux, Myc-Atg1, or both

revealed a direct interaction between dAux and Atg1. dAux or Atg1 was detected in the

pull down of Myc-Atg1 or Flag-dAux, respectively. (**B** and **C**) dAux variants lacking

different domains or carrying an early-onset PD SNP mutation (DNAJC6m) were

constructed and expressed in fly, adult heads were used for WB to check the expression

level. Note that dAux$^{\Delta PTEN}$ expressed poorly. (**D**) Different dAux variants were

expressed in S2 cells, and analyzed for their interaction with Atg1. Note that J domain

was required for dAux-Atg1 interaction. (**E** and **F**) Representative images (E) and

quantifications (F) of the ULK1 puncta number in the "Glia" region of adult fly brains

expressing different dAux variants. Note that the ULK1 puncta number increased when

expressing dAux$^{\Delta Kinase}$, dAux$^{\Delta J}$, or dAux^{DNAJC6m}. (**G** and **H**) N-terminal EGFP-fused

ULK1 variants lacking different domains were constructed, expressed in HEK293FT

cells, and analyzed for their interaction with GAK. Note that ULK1 lacking the C-

terminal domain CTD failed to interact with GAK. (**I-L**) The representative

immunostaining (I) and WB (K) images and quantifications (J and L) of the ULK1

protein level in *GAK* siRNA-treated HeLa cells. Note that *GAK* knockdown caused an

increase in the number and size of the ULK1 puncta, also the ULK1 protein level. (**M-**

P) Epistatic genetic analysis revealed that dAux acts upstream of Atg1 in regulating

autophagy initiation. Co-expression of *atg1*-RNAi in glia rescueed and further

suppressed the *daux*-RNAi-induced increase in the number of GFP.ZFYVE1- and

mCh.Atg8a-positive puncta in "Glia" region. Scale bars and the sample number n are

indicated in the Figures. For each experiment, more than three biologically independent

replicates were done. Whereas results were consistent, representative results from one

experiment was shown. A serial confocal Z-stack sections were taken with 0.4 μm each,

and representative single layer images acquired at the similar plane are shown.　　Data

are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no

significance, * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$) are calculated by Mann-Whitney

test, ordinary one-way ANOVA followed by Tukey's multiple comparisons test, or

Kruskal-Wallis tests followed by Dunn's multiple comparisons test.

Figure 4. GAK/dAux regulates ULK1/Atg1 trafficking from TGN to

autophagosomes

(**A** and **B**) HeLa cells transfected with RFP-LC3 in the presence or absence of the *GAK*

siRNA were collected for immunostaining analysis with the anti-ULK1 antibodies.

Note that both the LC3-positive puncta number and the ULK1 (green)-LC3 (red)

colocalization increased upon GAK depletion. (**C** and **D**) ULK1-Atg8a colocalization

was analyzed in "Glia" region of adult fly brains expressing mCh.Atg8a

(*repo>mCh.Atg8a*) to label glial autophagosomes. Note that ULK1 (green)-Atg8a (red)

colocalization increased in the absence of glial dAux. (**E** and **F**) ULK1 (green)-TGN38

(red) colocalization is analyzed in *GAK* siRNA-treated HeLa cells. Note that ULK1-

TGN38 colocalization decreased, suggesting enhanced ULK1 trafficking to the

autophagosomes upon GAK depletion. Areas enclosed by the white dashed squares in

the representative images (A, C, and E) are enlarged in A', C', and E'. HeLa cell nuclei

were stained with DAPI (blue). Scale bars and the sample number n are indicated in the

Figures. For each experiment, more than three biologically independent replicates were

done. Whereas results were consistent, representative results from one experiment was

shown. A serial confocal Z-stack sections were taken with 0.4 μm each, and

representative single layer images acquired at the similar plane are shown.

Colocalization is analyzed using the Manders' Correlation, taking into account the

change in the protein level. Data are shown as mean ± SEM. P-values of significance

(indicated with asterisks, ns no significance, * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$)

are calculated by two-tailed unpaired t-test or ordinary one-way ANOVA followed by

Tukey's multiple comparisons test.

Figure 5. dAux-mediated Atg9 trafficking depend on Atg1

(**A-D**) Atg9 trafficking is investigated by analyzing the Atg9 (green)-Atg8a (red) colocalization in the "Glia" region of adult fly brains expressing the *UAS-EGFP.Atg9* and *UAS-mCh.Atg8a* reporters (*UAS-mCh.Atg8a; repo-GAL4, UAS-EGFP.Atg9*). Note that *daux*-RNAi or *atg1*-RNAi expression in glia caused a significant increase or decrease in the Atg9-Atg8a colocalization, respectively. Co-expression of *daux*-RNAi and *atg1*-RNAi rescued and further suppressed the increased Atg9-Atg8a colocalization. Areas enclosed by the white dashed squares in the representative images (A and B) are enlarged in A' and B'. (**E-H**) Live-cell imaging (E) and quantifications (F-H) of Atg9 trafficking to Atg8a in "Glia" region of adult fly brains. Note the percentage of Atg9-positic puncta being trafficked to the vicinity of Atg8a-positive puncta (with a radius of 0.35 um) was increased (F), the duration of the Atg9-Atg8a contacting was longer (G), and the speed of both Atg8a- and Atg9-positive puncta were decreased upon glial dAux depletion (H). Scale bars and the sample number n are indicated in the Figures. For each experiment, more than three biologically independent replicates were done. Whereas results were consistent, representative results from one experiment was shown. A serial confocal Z-stack sections were taken with 0.4 μm each, and representative single layer images acquired at the similar plane are shown. Colocalization is analyzed using the Manders' Correlation, taking into account the change in the protein level. Data are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated by two-tailed

unpaired t-test, Mann-Whitney test, ordinary one-way ANOVA followed by Tukey's

multiple comparisons test, or Kruskal-Wallis tests followed by Dunn's multiple

comparisons test.

Figure 6. Lack of GAK/dAux impairs autophagic flux and substrate degradation

(**A** and **B**) Autophagic flux was impaired as examined by expressing the flux reporter

UASp-GFP-mCh-Atg8a. Both the GFP- and mCh-positive puncta colocalization and

the intensity of GFP puncta in "Glia" region increased upon depleting glial dAux. In

the presence of BafA1, the GFP- and mCh-positive puncta colocalization further

increased upon depleting glial dAux, a similar trend of GFP puncta intensity was

detected but not statistically significant. Note that GFP- and mCh-positive puncta were

enlarged and potentially dysfunctional due to the non-quenched GFP signals. (**C** and **D**)

Representative confocal/STED images (C) and quantifications (D) of Lamp1- and

Atg8a-positive puncta in "Glia" region of adult fly brains. Note that dAux depletion in

glia increased the number and size of glial autolysosomes and the colocalization of

autophagosomes and lysosomes. (**E-H**) Autophagic substrates such as Ref(2)P (the

Drosophila P62 homolog) and Ubiquitin (Ubi) accumulated in *repo>daux*-RNAi adult

fly brains as revealed by immunostaining ("Glia" region) and WB analyses. (**I** and **J**)

P62 and Ubi protein levels increased upon *Gak* depletion in IMG cells. (**K** and **L**) P62

and Ubi protein levels increased upon *GAK* depletion in HeLa cells. (**M**) A model

illustrating the regulating roles of GAK/dAux in glial autophagy. In *GAK/daux-*

deficient glia, lack of GAK/dAux activates ULK1/Atg1, upregulates protein levels of

components in the initiation and PI3P class III complexes, and enhances Atg1

trafficking to the autophagosomes (thicker black arrows). dAux also regulates Atg9

trafficking to the autophagosomes via Atg1. These actions promote autophagy initiation

and start the glial clearance program for substrates degradation. On the other hand,

despite the benefits of inducing glial autophagy, lack of GAK/dAux also disrupts

autophagic flux and substrate degradation. GAK/dAux regulatory steps are highlighted

in green. Scale bars and the sample number n are indicated in the Figures. For each

experiment, more than three biologically independent replicates were done. Whereas

results were consistent, representative results from one experiment was shown. A serial

confocal Z-stack sections were taken with 0.4 μm each, and representative single layer

images acquired at the similar plane of brains across all genotypes are shown. Data are

shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no

significance, * $p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated by two-tailed t-test

or ordinary one-way ANOVA followed by Tukey's multiple comparisons test.

References

1.   Mizushima N (2018) A brief history of autophagy from cell biology to
physiology and disease. *Nat Cell Biol* 20(5):521-527.
2.   Dikic I & Elazar Z (2018) Mechanism and medical implications of mammalian
autophagy. *Nat Rev Mol Cell Biol* 19(6):349-364.
3.   Mizushima N & Komatsu M (2011) Autophagy: renovation of cells and tissues.
Cell 147(4):728-741.
4.   Mizushima N (2020) The ATG conjugation systems in autophagy. *Curr Opin*
Cell Biol 63:1-10.

5. Galluzzi L, *et al.* (2017) Molecular definitions of autophagy and related
processes. *EMBO J* 36(13):1811-1836.
6. Hara T, *et al.* (2006) Suppression of basal autophagy in neural cells causes
neurodegenerative disease in mice. *Nature* 441(7095):885-889.
7. Komatsu M, *et al.* (2006) Loss of autophagy in the central nervous system
causes neurodegeneration in mice. *Nature* 441(7095):880-884.
8. Simonsen A, *et al.* (2008) Promoting basal levels of autophagy in the nervous
system enhances longevity and oxidant resistance in adult Drosophila.
Autophagy 4(2):176-184.
9. Deng Z, *et al.* (2017) Autophagy Receptors and Neurodegenerative Diseases.
Trends Cell Biol 27(7):491-504.
10. Levine B & Kroemer G (2019) Biological Functions of Autophagy Genes: A
Disease Perspective. *Cell* 176(1-2):11-42.
11. Kulkarni A, Chen J, & Maday S (2018) Neuronal autophagy and intercellular
regulation of homeostasis in the brain. *Curr Opin Neurobiol* 51:29-36.
12. Klionsky DJ, *et al.* (2016) Guidelines for the use and interpretation of assays
for monitoring autophagy (3rd edition). *Autophagy* 12(1):1-222.
13. Russell RC, *et al.* (2013) ULK1 induces autophagy by phosphorylating Beclin-
1 and activating VPS34 lipid kinase. *Nat Cell Biol* 15(7):741-750.
14. Mari M, *et al.* (2010) An Atg9-containing compartment that functions in the
early steps of autophagosome biogenesis. *J Cell Biol* 190(6):1005-1022.
15. Yamamoto H, *et al.* (2012) Atg9 vesicles are an important membrane source
during early steps of autophagosome formation. *J Cell Biol* 198(2):219-233.
16. Orsi A, *et al.* (2012) Dynamic and transient interactions of Atg9 with
autophagosomes, but not membrane integration, are required for autophagy.
Mol Biol Cell 23(10):1860-1873.
17. Puri C, Renna M, Bento CF, Moreau K, & Rubinsztein DC (2014) ATG16L1
meets ATG9 in recycling endosomes: additional roles for the plasma membrane
and endocytosis in autophagosome biogenesis. *Autophagy* 10(1):182-184.
18. Ravikumar B, Moreau K, Jahreiss L, Puri C, & Rubinsztein DC (2010) Plasma
membrane contributes to the formation of pre-autophagosomal structures. *Nat
Cell Biol* 12(8):747-757.
19. Song L, *et al.* (2017) Auxilin Underlies Progressive Locomotor Deficits and
Dopaminergic Neuron Loss in a Drosophila Model of Parkinson's Disease. *Cell
Rep* 18(5):1132-1143.
20. Zhang CX, *et al.* (2005) Multiple roles for cyclin G-associated kinase in
clathrin-mediated sorting events. *Traffic* 6(12):1103-1113.
21. Lee DW, Wu X, Eisenberg E, & Greene LE (2006) Recruitment dynamics of
GAK and auxilin to clathrin-coated pits during endocytosis. *J Cell Sci* 119(Pt
17):3502-3512.
22. He K, *et al.* (2020) Dynamics of Auxilin 1 and GAK in clathrin-mediated traffic.
J Cell Biol 219(3).

23.   Shimizu H, Nagamori I, Yabuta N, & Nojima H (2009) GAK, a regulator of
676   clathrin-mediated membrane traffic, also controls centrosome integrity and
677   chromosome congression. *J Cell Sci* 122(Pt 17):3145-3152.

678   24.   Zhou X*, et al.* (2011) Auxilin is required for formation of Golgi-derived
679   clathrin-coated vesicles during Drosophila spermatogenesis. *Development*
680   138(6):1111-1120.

681   25.   Zhang Y*, et al.* (2014) An RNA-sequencing transcriptome and splicing database
682   of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci*
683   34(36):11929-11947.

684   26.   Zhang Y*, et al.* (2016) Purification and Characterization of Progenitor and
685   Mature Human Astrocytes Reveals Transcriptional and Functional Differences
686   with Mouse. *Neuron* 89(1):37-53.

687   27.   McCarthy RC*, et al.* (2016) Characterization of a novel adult murine
688   immortalized microglial cell line and its activation by amyloid-beta. *J*
689   *Neuroinflammation* 13:21.

690   28.   Mauvezin C, Nagy P, Juhasz G, & Neufeld TP (2015) Autophagosome-
691   lysosome fusion is independent of V-ATPase-mediated acidification. *Nat*
692   *Commun* 6:7007.

693   29.   Yoshimori T, Yamamoto A, Moriyama Y, Futai M, & Tashiro Y (1991)
694   Bafilomycin A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits
695   acidification and protein degradation in lysosomes of cultured cells. *J Biol*
696   *Chem* 266(26):17707-17712.

697   30.   Liu W, Duan X, Fang X, Shang W, & Tong C (2018) Mitochondrial protein
698   import regulates cytosolic protein homeostasis and neuronal integrity.
699   *Autophagy* 14(8):1293-1309.

700   31.   Melani M*, et al.* (2017) Zonda is a novel early component of the autophagy
701   pathway in Drosophila. *Mol Biol Cell* 28(22):3070-3081.

702   32.   Elsayed LE*, et al.* (2016) A Novel Nonsense Mutation in DNAJC6 Expands the
703   Phenotype of Autosomal-Recessive Juvenile-Onset Parkinson's Disease. *Ann*
704   *Neurol* 79(2):335-337.

705   33.   Olgiati S*, et al.* (2016) DNAJC6 Mutations Associated With Early-Onset
706   Parkinson's Disease. *Ann Neurol* 79(2):244-256.

707   34.   Bido S*, et al.* (2021) Microglia-specific overexpression of alpha-synuclein leads
708   to severe dopaminergic neurodegeneration by phagocytic exhaustion and
709   oxidative toxicity. *Nat Commun* 12(1):6237.

710   35.   Nabar NR*, et al.* (2022) LRRK2 is required for CD38-mediated NAADP-Ca(2+)
711   signaling and the downstream activation of TFEB (transcription factor EB) in
712   immune cells. *Autophagy* 18(1):204-222.

713   36.   Winslow AR*, et al.* (2010) alpha-Synuclein impairs macroautophagy:
714   implications for Parkinson's disease. *J Cell Biol* 190(6):1023-1037.

715   37.   Papinski D*, et al.* (2014) Early steps in autophagy depend on direct
716   phosphorylation of Atg9 by the Atg1 kinase. *Mol Cell* 53(3):471-483.

38. Reggiori F, Tucker KA, Stromhaug PE, & Klionsky DJ (2004) The Atg1-Atg13
complex regulates Atg9 and Atg23 retrieval transport from the pre-
autophagosomal structure. *Dev Cell* 6(1):79-90.

39. Zhou C*, et al.* (2017) Regulation of mATG9 trafficking by Src- and ULK1-
mediated phosphorylation in basal and starvation-induced autophagy. *Cell Res*
27(2):184-201.

40. Hegedus K, Takats S, Kovacs AL, & Juhasz G (2013) Evolutionarily conserved
role and physiological relevance of a STX17/Syx17 (syntaxin 17)-containing
SNARE complex in autophagosome fusion with endosomes and lysosomes.
Autophagy 9(10):1642-1646.

41. Kim J, Kundu M, Viollet B, & Guan KL (2011) AMPK and mTOR regulate
autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol* 13(2):132-141.

42. Karanasios E*, et al.* (2013) Dynamic association of the ULK1 complex with
omegasomes during autophagy induction. *J Cell Sci* 126(Pt 22):5224-5238.

43. Papinski D & Kraft C (2016) Regulation of Autophagy By Signaling Through
the Atg1/ULK1 Complex. *J Mol Biol* 428(9 Pt A):1725-1741.

44. Pengo N, Agrotis A, Prak K, Jones J, & Ketteler R (2017) A reversible phospho-
switch mediated by ULK1 regulates the activity of autophagy protease ATG4B.
Nat Commun 8(1):294.

45. Hurley JH & Young LN (2017) Mechanisms of Autophagy Initiation. *Annu Rev*
Biochem 86:225-244.

46. Park JM*, et al.* (2016) The ULK1 complex mediates MTORC1 signaling to the
autophagy initiation machinery via binding and phosphorylating ATG14.
Autophagy 12(3):547-564.

47. Petherick KJ*, et al.* (2015) Pharmacological inhibition of ULK1 kinase blocks
mammalian target of rapamycin (mTOR)-dependent autophagy. *J Biol Chem*
290(18):11376-11383.

48. Zachari M, Longo M, & Ganley IG (2020) Aberrant autophagosome formation
occurs upon small molecule inhibition of ULK1 kinase activity. *Life Sci*
Alliance 3(12).

49. Chan EY, Longatti A, McKnight NC, & Tooze SA (2009) Kinase-inactivated
ULK proteins inhibit autophagy via their conserved C-terminal domains using
an Atg13-independent mechanism. *Mol Cell Biol* 29(1):157-171.

50. Munson MJ*, et al.* (2021) GAK and PRKCD are positive regulators of PRKN-
independent mitophagy. *Nat Commun* 12(1):6101.

51. Miyazaki M*, et al.* (2021) Targeted disruption of GAK stagnates autophagic flux
by disturbing lysosomal dynamics. *Int J Mol Med* 48(4).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6