1 **SNARE-dependent mitochondria-lysosome contacts regulate glial**

2 **mitochondrial dynamics and dopaminergic neuron survival**

5 Honglei Wang[1], Linfang Wang[1], Shuanglong Yi[1], Shiping Zhang[1], Margaret S. Ho[1][*]

7 [1]School of Life Science and Technology, ShanghaiTech University, Shanghai, 201210,

8 China

[*]Corresponding author:

Margaret S. Ho, Ph.D.

#B416, L building,

#230 Haike Road, Pudong New District,

Shanghai, 201210

Phone: +86-21-2068-5424

Email: margareth@shanghaitech.edu.cn

Summary

Mitochondria are dynamic organelles that undergo fission and fusion, enabling swift structural modification to adapt cellular needs. Disturbances in mitochondrial dynamics, frequently defects ascribed to neurons, have been associated with pathological contexts such as neurodegeneration in Parkinson's disease. Nonetheless, the mechanism of glial mitochondrial dynamics contributing to neurodegeneration remains unclear. Here we present evidence that the *Drosophila* R-SNARE VAMP7 regulates glial mitochondrial dynamics and dopaminergic neuron survival via modulating the dynamic of mitochondria-lysosome contact, which determines the mitochondrial fission site. Independent of its characterized role in autophagosome-lysosome fusion, glial VAMP7 depletion causes mitochondrial elongation and dysfunction, increased ROS levels, and production of lipid droplets. These conferred changes in glia in turn affects nearby dopaminergic neuron survival. Glial VAMP7 genetically interacts with the mitochondrial fission/fusion factors Drp1 and Marf1 and controls glial mitochondrial dynamics via regulating the frequency and duration of mitochondria-lysosome contact. Our findings indicate that SNARE proteins, although not direct mediators on mitochondrial fusion, provide spatial cues to modulate glial mitochondrial fission via organelle contacts, impacting on neuron survival in a non-cell-autonomous manner.

Keywords

VAMP7; Glia; Mitochondrial Dynamics; Mitochondria-Lysosome Contact;

Neurodegeneration

Introduction

Mitochondria are dynamic organelles that undergo structural and functional changes

in response to a variety of cellular signals such as energy production, metabolism, or

survival. They constantly remodel and interact with other organelles so that their

morphologies are swiftly modified to adapt the cellular needs. Different aspects of

mitochondrial dynamics have been uncovered including fission (division), fusion, or

shape transition, generating different forms of mitochondria that function in various

cellular contexts (1-3). These dynamic changes in morphology and function are under

tight regulation and have been extensively linked to catastrophic consequences such

as neurodegeneration and death (4, 5). For instance, mitochondrial fission regulated

by dynamin-related protein-1 (Drp1), a GTPase that dynamically associates with the

endoplasmic reticulum (ER) and mitochondria (6-10), suppresses oxidative damage

and safeguards neuron survival (11). It is generally considered that the fission process

generates smaller mitochondria easier to be targeted by mitophagy, the

autophagosomal degradation of damaged mitochondria mediated by Parkinson's

disease (PD) risk genes *PINK1* and Parkin (12-14). On the other hand, elongated

mitochondria due to the lack of fission accumulate oxidative damage and lose

respiratory function, transforming into large spheres resistant to mitophagy.

Interestingly, most of these mitochondrial dynamic defects underlying

neurodegeneration are ascribed to neurons, whereas how mitochondrial dynamics in

glia, the other major brain cell type important for neuron survival, contributes to

neurodegeneration, remains very limitedly explored.

In addition to ER, mitochondrial fission sites are also contacted by lysosomes,

which help to restrict mitochondria motility by recruiting the lysosome GTPase Rab7

(15, 16). It has been shown that Rab7 GTP hydrolysis is crucial for the untethering of

lysosome and mitochondria (16). Upon mitochondria-lysosome contact untethers,

mitochondrial fission occurs. These findings implicate a role for lysosome in

regulating mitochondrial dynamics. In regard to this aspect, whether lysosome and its

related factors play a role in glial mitochondrial fission is largely unknown; how

mitochondrial dynamic is controlled in glia has not been extensively studied.

The *Drosophila* vesicular R-SNARE (soluble N-ethylmaleimide-sensitive factor

attachment protein receptors), VAMP7, is homologous to mammalian VAMP7 and

VAMP8, and the latter exhibits a prominent expression in microglia (17, 18).

Mammalian VAMP7 or VAMP8 localizes on the late endosome and lysosome and

forms a complex with the cytosolic Synaptosomal-associated protein 29 (SNAP-29)

and the autophagosomal Syntaxin-17 (Syx17); together this complex regulates the

autophagosome-lysosome fusion in autophagy (19, 20). Despite a role in vesicle

fusion, whether VAMP7 plays additional roles with respect to its lysosomal origin

remains unclear. Here we present evidence that VAMP7 regulates glial mitochondrial

dynamics via modulating the dynamics of mitochondria-lysosome contacts. Lack of

glial VAMP7 causes mitochondrial elongation independent of its autophagy function,

increased ROS levels and the production of lipid droplets (LDs), and regulates nearby

dopaminergic (DA) neuron survival. Glial VAMP7-mediated mitochondrial

elongation is suppressed by expressing the fission factor Drp1 or downregulating the

expression of the mitochondrial assembly regulatory factor Marf, indicating a role for

glial VAMP7 in mitochondrial fission/fusion. Considering that SNARE proteins

mediate membrane fusion, our findings uncover a new role for a canonical SNARE

factor in glial mitochondrial fission, demonstrating its importance in glial contribution

to neuron survival.

Results

Glia deficient of VAMP7 exhibit mitochondrial elongation

Previously characterized as a fusion factor regulating autophagosome-lysosome

fusion, we first confirm if VAMP7 functions similarly in glia. Two independent RNAi

lines: *vamp7*-RNAi[#38300] and *vamp7*-RNAi[#43543] were used to selectively silence

vamp7 expression. Expression of either RNAi in glia abolished *vamp7* transcription

levels, validating the efficacy of RNAi (Figure S1A). Consistent to previous findings,

glial autophagic flux monitored by a dual reporter *UAS-GFP-mCherry-Atg8a* in the

absence of glial VAMP7 was strikingly impaired. The values of Pearson's coefficient

indicating colocalization of GFP and mCherry fluorescent signals were significantly

increased upon glial VAMP depletion, indicating non-quenched GFP signals due to

impaired fusion (Figure S1B and S1C). These results confirm a role for VAMP7 in

glial autophagosome-lysosome fusion.

Taking advantage of the RNAi lines, we next investigated thoroughly the

consequences of lacking glial VAMP7 in the brains. Striking defects were detected

when analyzing Transmission Electron Microscopy (TEM) images of adult fly brains

expressing *vamp7*-RNAi in glia. In addition to the presence of LDs and mitophagic

vacuoles, mitochondria were elongated, indicating a possible defect in fission or

fusion (yellow arrows, Figure 1A). The morphologies of mitochondria were

categorized and quantified by comparing the values of mitochondrial length over

diameter (L/D). Interestingly, the ratio of mitochondria with a L/D number lower than

2.5 decreased, whereas the ones with a L/D number higher than 2.5 increased upon

glial VAMP7 depletion (Figures 1A and 1B). These results indicate that mitochondria

exhibit elongated morphology in the absence of glial VAMP7. Furthermore, two

independent mitochondrial markers: *UAS-mito-HA-GFP* (*mito.GFP*) and

UAS-mitoOMM-mCherry (*mitoOMM.mCherry*), that expresses a mitochondrial

import signal and labels the outer mitochondrial membrane, respectively, were used

to assess glial mitochondrial dynamics in the absence of VAMP7. Consistent with the

EM results, mitochondrial area and volume were significantly increased when

expressing *vamp7*-RNAi in glia, indicating that mitochondria are elongated (Figures

1C-1E and S1D-S1F). Taken together, these results indicate that lack of glial VAMP7

disrupts mitochondrial dynamics, possibly due to a defect in fission or fusion.

Given that VAMP7 is a characterized fusion factor in autophagy, we next

determine if VAMP7-mediated glial mitochondrial dynamics depends on

autophagosome-lysosome fusion. Interestingly, mitochondrial area and volume in glia

were not altered in the presence of Bafilomycin A1 (BafA1), a drug blocking

autophagosome-lysosome fusion (21, 22) (Figures 1F-1H). Moreover, glial inhibition

of the other factor involved in autophagosome-lysosome fusion, Syx17, did not cause

mitochondrial elongation (Figures S1G-S1I). Taken together, these results indicate

that VAMP7 mediates glial mitochondrial dynamics independent of its function in

autophagosome-lysosome fusion.

Lack of glial VAMP7 leads to mitochondrial dysfunction

To further demonstrate that VAMP7-mediated mitochondrial elongation is

physiologically relevant, the calcium sensor *mitoGCamp5.0* was expressed in glia to

monitor the mitochondrial calcium activity (23). Whereas an increase in the

mitoGCamp5.0 signal was detected in the control young animals, the increase was

blocked when silencing glial *vamp7* expression (Figure 2A). The increased

mitoGCamp5.0 signal was similarly blocked by glial VAMP7 depletion when flies

aged (Figure 2B). These results indicate that glial VAMP7 regulates mitochondrial

calcium activity. As mitochondria are the major sites for ROS production, adult fly

brains of control and *repo>vamp7*-RNAi animals were examined for ROS levels with

the dihydroethidium (DHE) staining. Consistent with previous findings, DHE-positive

fluorescent intensities increased when expressing the RNAi targeting *superoxide*

dismutase 1 (*sod1*), the enzyme that regulates ROS levels by catalyzing the

conversion of superoxide into oxygen and hydrogen peroxide (Figure 2C).

Interestingly, inhibiting *vamp7* expression in glia caused similar increase, indicating

that glial VAMP7 expression regulates cellular ROS levels (Figure 2C). In addition,

the transcriptional levels of genes related to the mitochondrial fatty acid β-oxidation

pathway, including those encoding acyl-coA dehydrogenases (*CG17544* and *CG9527*)

and fatty acid synthase (*FASN*), were increased upon glial *vamp7*-RNAi expression,

indicating altered mitochondrial function (24, 25) (Figure 2D). Notably, the levels of

BODIPY-positive LDs increased in both larval and adult fly brains in the absence of

glial VAMP7 (Figures 2E-2G), with a similar increase in the overall LD

triacylglycerols (TAG) content (Figure 2H). The increased LD number was

suppressed in the presence of α-Lipoic acid, a compound known for suppressing LD

formation (Figures 2I and 2J), indicating that glial VAMP7 regulates LD formation.

Taken together, these results indicate that glial VAMP7 regulates mitochondrial

function, ROS levels, and LD production via modulating mitochondrial dynamics in

glia.

Glial VAMP7 contributes to DA neuron survival

As glia interact with neurons and the glia-neuron crosstalk often contributes to

neuronal function and survival, we next investigated if VAMP7-mediated glial

mitochondrial dynamics contributes to DA neuron survival using the *Drosophila* PD

model established in the lab. DA neurons are located in clusters and named upon their

relative positions in adult fly brains (26, 27). These clusters include protocerebral

posterior lateral (PPL)1, PPL2, protocerebral posterior medial (PPM)1, PPM2, and

PPM3 (Figure 3A). Interestingly, lack of glial VAMP7 causes DA neuron loss at

PPM1/2, PPM3, and PPL1 clusters in an age-dependent manner (Figures 3B-3D).

These results suggest that glial VAMP7 contributes to DA neuron survival. Given that

the degeneration of DA neurons in the PPM1/2 cluster has been implicated in a

Drosophila PD model overexpressing α-Synuclein (28), it is conceivable that glial

VAMP7 plays important roles in regulating DA neuron survival under pathological

conditions.

Glial VAMP7 regulates mitochondrial fission and fusion

Given that lack of glial VAMP7 induces mitochondrial elongation, possibly due to

decreased fission or increased fusion, we tested genetically if glial VAMP7 interacts

with the fission factor Drp1 or the fusion factor Marf. Interestingly, increased Drp1

expression suppressed the mitochondrial elongation in the absence of glial VAMP7

(Figures 3E-3G), suggesting that increasing fission helps to prevent VAMP7-mediated

mitochondrial elongation. Moreover, decreased Marf expression in the presence of

glial *vamp7*-RNAi also suppressed the mitochondrial elongation. These results

indicate that Drp1 and Marf are epistatically downstream of glial VAMP7 in

controlling mitochondrial fission and fusion. Interestingly, endogenous protein levels

of Fission, mitochondrial 1 (Fis1), Drp1, Marf, and Optic astrophy 1 (Opa1) were not

affected by the absence of glial VAMP7, suggesting that VAMP7 does not regulate

glial mitochondrial dynamics via tuning fission/fusion proteins levels (29) (Figures

S2A-S2H). These results suggest that glial VAMP7 acts upstream of these

mitochondrial factors in regulating mitochondrial fission/fusion; mitochondrial

elongation caused by the lack of glial VAMP7 is potentially due to a change in the

dynamic or balance of fission and fusion.

Glial VAMP7 regulates the dynamic of mitochondria-lysosome contact

It has been reported that lysosomes contact mitochondria to restrict its motility and

determine the fission site (16, 30). Given that VAMP7 is localized on the lysosome,

we next investigated whether glial VAMP7 regulates mitochondrial dynamic via the

mitochondria-lysosome contact. Two reporters: *UAS-mitoOMM-mCherry* and

UAS-Lamp1-GFP, were co-expressed to label mitochondria and lysosomes

simultaneously, such that mitochondria-lysosome contact can be visualized under

various conditions. Interestingly, colocalization of mitoOMM and Lamp1 increased

upon glial VAMP7 depletion, indicating an increase in mitochondria-lysosome contact

(Figures 4A and 4B). Importantly, these dual positive contacts were not positive for

the autophagosome marker Atg8a, indicating that these do not represent mitochondria

undergoing mitophagy (Figures S2I-S2K). Whereas the value of Pearson's coefficient

for mitoOMM and Lamp1 increased significantly (Figure S2J), the colocalization

values for Atg8a and mitoOMM decreased upon glial VAMP7 depletion (Figure S2K).

These results indicate that lack of glial VAMP7 increases mitochondria-lysosome

contact and blocks mitophagy, consistent with the notion that elongated mitochondria

are resistant to and slows down mitophagy.

　　In addition, the dynamics of colocalized mitoOMM and Lamp1 were analyzed by

time lapse analysis *in-vivo*. Interestingly, mitochondria and lysosomes contacted each

other and dissociated over a course of 100 seconds, whereas the duration of these

contacts persisted and prolonged up to 400 seconds upon glial VAMP7 depletion

(Figures 4C, 4D, and Movie S1-S3). The frequency of mitochondria and lysosome

contacting each other and the duration of these contacts increased significantly

(Figures 4E and 4F) and a higher number of contacts persisting more than 5 min was

also detected (Figure 4G). Consistently, mitochondria and lysosome were significantly

less mobile in the absence of glial VAMP7 as indicated by their moving speed

(Figures 4H and 4I). Taken together, these results indicate that glial VAMP7 regulates

the dynamics of mitochondria-lysosome contacts.

Discussion

SNARE proteins are canonical regulators of membrane fusion, yet to date no

evidence indicates that they mediate mitochondrial fusion. Mitochondrial fission,

however, has been shown to rely on the activity of SNARE protein Syx17. Upon

starvation-induced autophagy, Syx17-Drp1 interaction is disrupted, causing a block in

mitochondrial fission, hence the elongated mitochondria resistant to mitophagy (31).

Syx17 also mediates fusion between the mitochondrial-derived vesicles (MDVs) and

the late endosome/lysosome under stress conditions (32). These results indicate that

SNARE proteins are possible mediators in mitochondrial dynamics. *Drosophila*

VAMP7 is highly homologous with mammalian VAMP8, which exhibits prominent

expression in microglia. Our findings reveal that lack of glial VAMP7 causes

elongated mitochondria, a phenotype genetically suppressed by increasing the

expression of the fission factor Drp1. These elongated mitochondria are dysfunctional

as mitochondrial calcium activity is significantly disrupted and the levels of ROS and

lipid droplets are increased, endangering the survival of nearby DA neurons. Despite a

role in autophagosome-lysosome fusion, glial VAMP7-mediated mitochondrial

dynamics does not depend on its fusion function as inhibiting

autophagosome-lysosome fusion by BafA1 does not alter mitochondrial length.

Mitochondrial fission site is determined by contacts between mitochondria and

ER or lysosomes. With its lysosomal localization, it is intriguing to speculate that

VAMP7 regulates mitochondria-lysosome contacts, determines the mitochondrial

fission site, in turn controls the fate of mitochondria undergoing fission, fusion, or

inter-mitochondrial clustering. Multiple lines of evidence support this hypothesis: first,

elongated mitochondria were aggregated and exhibit increasing contacts with

lysosomes upon glial VAMP7 depletion, suggesting abnormal tethering among

mitochondria themselves or between mitochondria and lysosomes. These abnormal

clusterings do not represent mitophagy events as no Atg8a-positive autophagosomes

are detected at these sites. Rather, the phenomenon suggests defective

mitochondria-lysosome contacts, which prevent mitochondria from fission, hence the

elongation. Similarly, the critical regulator Rab7 in mitochondria-lysosome contact

also appears to be mislocalized and aggregated in the absence of glial VAMP7, further

suggesting the disruptive formation of mitochondria-lysosome contact (Figure S3A).

It is noteworthy to mention that lysosomes exhibit abnormal acidity in the absence of

glial VAMP7 (Figures S3B and S3C). Moreover, as contact sites between

mitochondria and ER are known as sites for calcium transfer, inhibiting the glial

expression of TRPML, a mucolipin channel on the lysosomal membrane that releases

calcium, does not cause mitochondrial elongation (Figures S3D-S3F). These findings

suggest that lysosomal calcium transfer might not be responsible for the persisted

contacts and mitochondrial elongation induced by glial VAMP7 depletion.

Mitochondria-lysosome contacts might provide a more direct and physical cue to

modulate glial mitochondrial dynamics, via a route different from the calcium transfer

between mitochondria and ER.

Similar to neurons, glia, particularly astrocytes, possess SNARE machineries for

their vesicular trafficking function. These SNARE machineries, some in the same

composition as in neurons but some different, are responsible for slow secretory

exocytosis rather than fast and immediate release of signaling molecules (33). Very

importantly, they regulate synaptic transmission and plasticity, antigen presentation in

neuro-inflammation, and control different types of behavior such as sleep. Our

findings unexpectedly reveal a new role for the canonical SNARE protein VAMP7 in

glial mitochondrial dynamics. Considering that mitochondria are present in astrocytic

processes in close proximity to neurons and contribute to local calcium signaling, ATP

production, and metabolism of signaling molecules, it is conceivable that

VAMP7-mediated glial mitochondrial dynamics subsequently impacts mitochondrial

function and participates in the metabolic process supporting neuron survival. Given

that DA neurodegeneration is one of the hallmarks in PD, our findings illustrate a

SNARE-dependent mechanism controlling organelle contacts, glial mitochondrial

dynamics, and DA neuron survival, providing the mechanistic basis underlying DA

neurodegeneration. Canonical SNARE proteins not only regulate membrane fusion,

but also important mediators in mitochondrial dynamics; future work is warranted to explore additional roles of SNARE proteins in various cellular contexts in both cell-autonomous and non-cell-autonomous fashions.

Acknowledgements

We thank Bloomington *Drosophila* Stock Center, Vienna *Drosophila* RNAi Center, Tsinghua Fly Center, the Core Facility of *Drosophila* Resource and Technology, Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences, Developmental Studies Hybridoma Bank, Chao Tong, Chun-Hong Chen, Henry Sun, and Aike Guo for fly stocks and antibodies. We also thank the Molecular Imaging Core Facility (MICF), the Molecular and Cell Biology Core Facility (MCBCF), and the Multi-Omics Core Facility (MOCF) at the School of Life Science and Technology, ShanghaiTech University for providing technical support; Yu Kong and Lijun Pan in Electron Microscopy Facilities of Center for Excellence in Brain Science and Technology, Chinese Academy of Science for assistance with EM sample preparation. We thank Ho lab members for discussion and comments. This work was supported by grants from ShanghaiTech and National Natural Science Foundation of China (31871039).

Author Contributions

H.W, S.Z, and M.S.H conceived and designed the study. H.W, L.W, and S.Y performed the experiments. H.W and M.S.H analyzed the data. M.S.H wrote the paper

with input from H.W. All authors read and approved the manuscript.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Lack of glial VAMP7 causes mitochondrial elongation (**A** and **B**)

Representative TEM images (A) and quantifications (B) of mitochondrial morphology

in control and *repo>vamp7*-RNAi[43543] adult fly brains of 10-day-old. Note the

presence of lipid droplets (red arrows), mitophagic vacuoles (pink arrows), and

elongated mitochondria (yellow arrows) in the absence of glial VAMP7. Note that the

ratio of mitochondria with a length to diameter (L/D) number lower than 2.5 is

significantly lower while the ratio for L/D number above 2.5 is significantly higher

upon glial VAMP7 depletion. (**C**-**E**) Representative confocal images (C) and

quantifications (D and E) of glial mitochondrial morphology monitored by

UAS-mito.GFP in control and *repo>vamp7*-RNAi[43543] adult fly brains. Note that the

average mitochondrial area and volume quantified by Imaris are significantly larger

than the control animals when expressing *vamp7*-RNAi in glia. (**F**-**H**) Representative

confocal images (F) and quantifications (G and H) of glial mitochondrial morphology

in the absence or presence of Bafilomycin A1 (BafA1). Note that no change in the

mitochondrial area and volume is observed when autophagosome-lysosome fusion is

blocked by BafA1. Insets represent enlarged view of single mitochondria. Scale bar:

500 nm in A, 10 μm in C and F. Mitochondrial area and volume are quantified by

Imaris as detailed in the Materials and Methods. The numbers of brains analyzed and

quantified are indicated in the Figures. For TEM, 6 independent fly brains for each

genotype were analyzed. Data are shown as mean ± SEM. P-values of significance

(indicated with asterisks, ns no significance, * p<0.05, ** p<0.01, and *** p<0.001)

are calculated by two-tailed unpaired t-test. Approximately 20 confocal Z-stack

sections were taken with 0.125 μm each. Merged images are shown. All data in

subsequent figures are present and analyzed as described above unless stated

otherwise.

Figure 2. Lack of glial VAMP7 disrupts mitochondrial function and increases

ROS levels and lipid droplet production (**A** and **B**) The reporter

UAS-mitoGCamp5.0 was expressed together with *UAS-mitoOMM.mCherry* in glia to

monitor mitochondrial calcium activity (*mitoOMM.mCherry; repo-Gal4,*

mitoGCamp5.0). Note that the increased mitochondrial calcium activity is blocked in

both young and old flies when inhibiting glial VAMP7 expression. (**C**) DHE

fluorescence intensities representing cellular ROS levels were significantly increased

upon glial VAMP7 depletion. (**D**) RT-qPCR analysis reveals increased transcription

levels of genes related to mitochondrial function including the ones encoding

acyl-coA dehydrogenases (*CG17544* and *CG9527*) and fatty acid synthase (*FASN*).

(**E-H**) The numbers of LDs labelled by the BODIPY dye in both larval and adult fly

brains (E-G) and the TAG content in the LDs (H) are significantly increased in the

absence of glial VAMP7. (**I** and **J**) Glial VAMP7 regulates LD formation as addition

of α-lipoic acid suppresses LD formation in both control and *repo>vamp7*-RNAi[43543]

adult fly brains of 3-day-old. Scale bar: 100 μm in 3rd instar larvae in E and I, 50 μm

in Day 3 in E. The numbers of brains analyzed and quantified are indicated in the

Figures. For α-lipoic acid treatment, 8 independent fly brains for each genotype were

analyzed. Data are shown as mean ± SEM. P-values of significance (indicated with

asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated by

two-tailed unpaired t-test.

Figure 3. VAMP7 regulates glial mitochondrial dynamics and DA neuron

survival (**A**) An schematic diagram illustrating the DA neuron clusters in adult fly

brains. The black dashed box encloses the area visualized in B. (**B-D**) The numbers of

DA neurons in PPM1/2, PPM3, and PPL1 are significantly decreased (white arrows in

B) in the aged flies lacking glial VAMP7. (**E-G**) Epistatic analysis of VAMP7 and

mitochondrial fission/fusion factors Drp1, Marf, and Opa1 in regulating mitochondrial

dynamics. Note that increasing expression of the fission factor Drp1 or

downregulating the expression of the fusion factor Marf or Opa1 causes increased

fission, opposite to the mitochondrial elongation phenotype in the absence of glial

VAMP7. Co-expression of *Drp1*, *Marf*-RNAi, or *Opa1*-RNAi and *vamp7*-RNAi in

glia all leads to suppression on mitochondrial elongation. Insets represent enlarged

view of single mitochondria. Scale bar: 50 μm in B, 10 μm in E. The numbers of

brains analyzed and quantified are indicated in the Figures. For DA neurons statistics,

15 independent fly brains for each genotype were analyzed (C and D). Data are shown

as mean ± SEM. P-values of significance (indicated with asterisks, ns no significance,

* $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) are calculated by ordinary

one-way ANOVA followed by Tukey's multiple comparisons test.

Figure 4. VAMP7 regulate glial mitochondrial dynamics via

mitochondria-lysosome contacts (**A** and **B**) Two reporters *UAS-mitoOMM.mCherry*

(labels the mitochondrial outer membrane, red) and *UAS-Lamp1* (labels lysosome,

green) are co-expressed to visualize mitochondria-lysosome contact. Note that

significant increase in colocalization of mitoOMM and Lamp1 upon glial VAMP7

depletion analyzed by Imaris. Co-localization traces are illustrated in yellow in A.

(**C-I**) Representative images (C and D) and quantifications (E-I) of the dynamic of

mitochondria-lysosome contact monitored by *in-vivo* time lapse analysis. Note that

mitochondria and lysosome contact each other and separate within the course of 100

seconds, while the contact persists up to 400 seconds upon glial VAMP7 depletion.

Both the frequency and the duration of the contacts increase, while the mobility of

both organelles decrease, when lacking glial VAMP7. Parameters in E-I are quantified

using Imaris and ImageJ. Scale bar: 10 μm in A, 5 µm in A (colocalization panel), 1

µm in C and D. The numbers of brains analyzed and quantified are indicated in the

Figures. For Lamp1.GFP and mitoOMM.mCherry colocalization statistics, 9

independent fly brains for each genotype were analyzed (B). Data are shown as mean

± SEM. P-values of significance (indicated with asterisks, ns no significance, *

$p<0.05$, ** $p<0.01$, and *** $p<0.001$) are calculated by two-tailed unpaired t-test.

Figure S1. Glia deficient of VAMP7 exhibit mitochondrial elongation (**A**)

RT-qPCR analysis indicates that expression of either *vamp7*-RNAi, #38300 or #43543,

in glia reduces the *vamp7* transcription levels significantly. (**B** and **C**) Glial

autophagic flux monitored by the dual reporter *UAS-GFP-mCherry-Atg8a* is impaired

upon glial VAMP7 depletion, suggesting impaired fusion. Note that the Pearson's

coefficient value indicating colocalization of GFP and mCherry increases when

lacking glial VAMP7 (**B**). (**D-F**) Glial mitochondrial dynamic is monitored by

expressing the reporter *UAS-mitoOMM.mCherry* in glia. Consistent with the other

reporter *UAS-mito.GFP*, Imaris analysis reveals an increase in the mitochondrial area

and volume in the absence of glial VAMP7. (**G-I**) Glial inhibition of *syx17* expression

does not cause mitochondrial elongation. Note that mitochondrial area and volume

quantified using the reporter *UAS-mito.GFP* remain unaffected in the presence of

syx17-RNAi. Scale bar: 10 μm in C, D and G. Mitochondrial area and volume are

quantified by Imaris as detailed in the Materials and Methods. The numbers of brains

analyzed and quantified are indicated in the Figures. Data are shown as mean ± SEM.

P-values of significance (indicated with asterisks, ns no significance, * $p<0.05$, **

$p<0.01$, and *** $p<0.001$) are calculated by two-tailed unpaired t-test or ordinary

one-way ANOVA followed by Tukey's multiple comparisons test. Approximately 20

confocal Z-stack sections were taken with 0.125 μm each. Merged images are shown.

All data in subsequent figures are present and analyzed as described above unless

stated otherwise.

Figure S2. VAMP7 regulates glial mitochondrial dynamics via

mitochondria-lysosome contact (**A-H**) Western blot analysis of endogenous

fission/fusion protein levels in the control and *repo>vamp7*-RNAi animals. Knock-in

lines at the original locus of Drp1, Opa1, and Marf created with an insertion of the HA

tag for protein detection and the Fis1 antibodies are gifts from Chun-Hong Chen. (**I-K**)

Two reporters *UAS-mitoOMM.mCherry* (labels the mitochondrial outer membrane,

red) and *UAS-Lamp1* (labels lysosome, green) are co-expressed to visualize

mitochondria-lysosome contact. Note that significant increase in colocalization of

mitoOMM and Lamp1 upon glial VAMP7 depletion analyzed by Imaris (J).

Immunostaining with anti-Atg8a antibodies reveals a decrease in the colocalization of

Atg8a and mitoOMM, suggesting that mitophagy is not induced (K). Note that no

Atg8a signals are detected in contact puncta positive for mitoOMM and Lamp1

(insets in I). Scale bar: 10 μm in I. The numbers of brains analyzed and quantified are

indicated in the Figures. Data are shown as mean ± SEM. P-values of significance

(indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, and *** $p<0.001$)

are calculated by two-tailed unpaired t-test or ordinary one-way ANOVA followed by

Tukey's multiple comparisons test.

Figure S3. VAMP7 regulates the dynamic of mitochondria-lysosome contact (**A**)

Two reporters *UAS-mitoOMM.mCherry* (labels the mitochondrial outer membrane,

red) and *UAS-Rab7.GFP* (labels late endosome and lysosome, green) are

co-expressed to visualize the dynamic of Rab7 contacting mitochondria. Note that

Rab7 exhibit aggregated and mislocalized pattern upon glial VAMP7 depletion. (**B**

and **C**) Lysosomal acidity monitored by the LysoTracker Red dye is significantly

impaired when expressing *vamp7*-RNAi in glia. Note that LysoTracker Red

fluorescent intensities increase upon glial VAMP7 depletion. (**D-F**) Mitochondrial

dynamic monitored by the reporter *UAS-mito.GFP* is significantly affected by

trpml-RNAi expression in glia. Note that mitochondrial area and volume are reduced,

indicating rounder and shorter morphology, opposite to mitochondrial elongation.

Scale bar: 10 μm in A, B and D. The numbers of brains analyzed and quantified are

indicated in the Figures. Data are shown as mean ± SEM. P-values of significance

(indicated with asterisks, ns no significance, * p<0.05, ** p<0.01, and *** p<0.001)

are calculated by two-tailed unpaired t-test or ordinary one-way ANOVA followed by

Tukey's multiple comparisons test.

Movie S1. *In-vivo* time lapse analysis of mitochondria-lysosome contact in the

control flies. Two reporters *UAS-mitoOMM.mCherry* (labels the mitochondrial outer

membrane, red) and *UAS-Lamp1* (labels lysosome, green) are co-expressed to

visualize mitochondria-lysosome contact. A representative contact is shown. A Z-stack

of 10 images was taken at an interval of 19 seconds for 10 minutes. Note that

mitochondria and lysosomes contact and dissociate over the imaging period.

Movie S2. *In-vivo* time lapse analysis of mitochondria-lysosome contact in flies

lacking glial VAMP7. Two reporters *UAS-mitoOMM.mCherry* (labels the

mitochondrial outer membrane, red) and *UAS-Lamp1* (labels lysosome, green) are

co-expressed to visualize mitochondria-lysosome contact. A representative contact is

shown. A Z-stack of 10 images was taken at an interval of 19 seconds for 10 minutes.

Note the significant increase in colocalization of mitoOMM and Lamp1 upon glial

VAMP7 depletion.

Movie S3. *In-vivo* time lapse analysis of mitochondria-lysosome contact in flies

lacking glial VAMP7. Two reporters *UAS-mitoOMM.mCherry* (labels the

mitochondrial outer membrane, red) and *UAS-Lamp1* (labels lysosome, green) are

co-expressed to visualize mitochondria-lysosome contact. A representative contact

different from Movie S2 is shown. A Z-stack of 10 images was taken at an interval of

19 seconds for 10 minutes. Note the significant increase in colocalization of

mitoOMM and Lamp1 upon glial VAMP7 depletion.

Materials and Methods

***Drosophila* genetics**

Flies were raised under standard conditions at 25□ with 70% humidity. All fly crosses

were carried out at 25°C with standard laboratory conditions unless noted otherwise.

All strains were obtained from Bloomington Drosophila Stock Center (BDSC), the

Vienna Drosophila RNAi Center (VDRC), Tsinghua Fly Center, or as gifts from

colleagues. The following fly strains were used in this study: *repo-GAL4* (a pan-glial

driver), *UAS-luciferase-IR* (BL31603), *UAS-lacZ* (BL1777), *UAS-mito-HA-GFP*

(BL8442), *UAS-mito.OMM.mCherry* (BL66532)*, UAS-Lamp1.GFP*, *UAS-vamp7*

RNAi (BL38300 and BL43543), *UAS-marf RNAi* (BL31157), *UAS-opa1 RNAi*

(THU0811), *UAS-sod1 RNAi* (THU3207), *UAS-Rab7.GFP* (BL42705);

UASp-GFP-mCherry-Atg8a (Henry Sun); *UAS-HRP* (Aike Guo); *UAS-drp1* (Chao

Tong); *drp1-HA*, *opa1-HA*, and *marf-HA* (Chun-Hong Chen).

Molecular biology

<u>qRT-PCR</u>: Total RNAs were extracted from adult fly heads using TransZol Up (Cat.

#ET111-01, TransGen, Beijing, China). cDNAs were reversely transcribed using HiScript III

RT SuperMix (Cat. #R323-01, Vazyme, Nanjing, China). qRT-PCR reactions were performed

using ChamQ Universal SYBR qPCR Master Mix (Cat. #Q711-02, Vazyme) and ABI 7500

RT-PCR system. The mRNA levels were normalized to *rp49* , respectively. The expression

levels were analyzed by $\Delta\Delta$CT method. Primers used for amplification:

rp49-F: CCACCAGTCGGATCGATATGC

rp49-R: CTCTTGAGAACGCAGGCGACC

vamp7-F: GGCTCGCAATTGGATAGTAGT

vamp7-R: TGCTTATCGGGGAATGAGCC

CG17544-F: *GTACTATGCTCTGACGCGCT*

CG17544-R: *GGTGGTCATGTGCTTCTCCA*

CG9527-F*: CTGCTTCCCCTGCTGAAGAA*

CG9527-R*: GACGTTTCCATCGCTCATGC*

FASN-F: *CGCTCCACTCCAAGAACTCG*

FASN-R*: CAGGTTTAGTTGTAGGGGCTAGA*

Immunohistochemistry

Adult fly brains were dissected and fixed in 4% formaldehyde for 45 minutes, then

washed with PBT (PBS + 0.3% TX-100) for 3 times and dissected further to remove

additional debris in PBS solution. Clean and fixed brains were blocked in PBT

solution with 5% Normal Donkey Serum (NDS) or 10% Bovine Serum Albumin

(BSA) and subsequently stained with primary antibodies at 4□ overnight and then

secondary antibodies at room temperature for 2 hours. Primary antibodies used:

mouse anti-Atg8a (1:1000, gift from Chao Tong) and rabbit anti-TH (1:1000, Cat.

#NB300-109, Novus). For all samples, secondary antibodies used were from Jackson

ImmunoResearch (West Grove, PA, USA): donkey anti-rabbit Cy3 (1:1000, Cat.

#711-166-152), donkey anti-mouse Cy5 (1:500, Cat. #715-175-150). All samples

were mounted in Vectashield mounting media (Cat. #H-1500-10, Vector Laboratories,

Burlingame, CA, USA).

In-vivo **time lapse analysis**

Live adult fly brains were dissected and kept in PBS solution. All samples were

mounted in PBS, and immediately scanned anteriorly to posteriorly (top to bottom)

using Nikon C2 confocal microscope (Tokyo, Japan) with a 60X oil objective. The

number of contacts were analyzed by Imaris using Autoregressive motion tracking

mode. Duration of contacts were analyzed by ImageJ using Time stamper.

Calcium imaging

Live adult fly brains expressing mitoGCamp5.0 were dissected and kept in PBS

solution. All samples were mounted in PBS, and immediately scanned in a single

focal plane using Nikon A1 confocal microscope (Tokyo, Japan) with a 60X oil

objective. The fluorescence intensity were analyzed by ImageJ using Time series

analyzer V3.

Live fluorescent dye imaging

Live adult fly brains were dissected and kept in PBS solution. For the analysis of

LDs, brains were incubated with 1 μg/mL BODIPY™ 493/503 (Cat. #D3922,

Invitrogen) for 5 minutes at 25°C, then immediately scanned by the Nikon C2

confocal microscope (Tokyo, Japan) with a 20X objective. To analyze LD formation,

brains were also treated with 10 μM α-Lipoic acid (Cat #D118666, Aladdin) for 3

days, followed by 1 μg/mL BODIPY™ 493/503 (Cat. #D3922, Invitrogen) for 5

minutes at 25°C, then immediately scanned by the Nikon C2 confocal microscope

(Tokyo, Japan) with a 20X objective. For the analysis of mitochondria-derived ROS

levels, brains were incubated with 5 μM DHE (Cat. #D11347, Invitrogen) for 5

minutes at 25°C, then immediately scanned using the Nikon A1 confocal microscope

(Tokyo, Japan) with a 20X objective. For the analysis of lysosome acidity, brains

were incubated with LysoTracker Red DND-99 (1:1000, Cat. #40739ES50, Yeasen)

solution in PBS for 5 minutes at 25°C, then immediately scanned using Nikon C2

confocal microscope (Tokyo, Japan) with a 60X oil objective.

Transmission electron microscopy

Adult fly brains were fixed, embedded, stained, and dehydrated according to previous

protocols (34). Ultrathin sections (70 nm) of each brain were cut with a

Diatomediamond knife on a Leica EM UC7 ultra-microtome (Leica Microsystems,

Germany) and collected on copper grids. Images were acquired with Talos L120C

transmission electron microscope (Thermo Fisher Scientific, Waltham, MA, UAS)

operating at an acceleration voltage of 120 kV.

Biochemistry

Adult fly heads were homogenised using a motorised pestle (Cat. #116005500, MP

Biomedicals, Irvine, CA, USA) in lysis buffer (0.4% NP-40, 0.2 mM EDTA, 150 mM

NaCl, 20% glycerol, 100 mM Tris-HCl pH7.5, 2% Tween 20, 0.5 mM

phosphodiesterase inhibitors, and 1 mM PMSF). Proteins were separated on

SDS-PAGE gels and transferred to PVDF membranes (Cat. #IPFL00010, Millipore,

Billerica, MA, USA). The membranes were blocked by 5% fat-free milk diluted in

PBS with 0.1% Tween-20 for 40 minutes. Samples were incubated with the primary

antibodies at 4°C overnight and HRP-conjugated secondary antibodies at room

temperature for 2 hours. Primary antibodies used: rat anti-HA-HRP (1:1000, Cat.

#3F10, Roche), mouse anti-α-Tubulin (1:5000, Cat. #T9026, Sigma). Bands were

visualized by Clarity Western ECL Substrate (Cat. #WBKLS0500, Millipore). Results

are representative of at least three biological replicates. For BafilomycinA1 treatment:

BafA1 (Cat. #BML-CM110-0100, ENZO) was diluted in DMSO (Cat. #D8371-50,

solarbio) in a concentration of 1 mM, then added to 0.5 ml of liquid yeast for a final

concentration of 4 μM. Yeast containing BafA1 was added to the top of the fly food in

standard vials kept overnight to allow the evaporation of residual DMSO. Flies of

different genotypes (20 females and 20 males) were then put in each vial and kept at

25°C for 5 days. After treatment, flies were collected for western blot analysis. The

TAG content in the LD was quantified by an triglyceride assay kit (Cat #E1103,

Applygen Technologies).

Confocal microscopy and statistical analysis

Images of adult fly brains were acquired by scanning a serial Z-stack of average 10-20

sections, each of 0.8 or 0.125 μm thickness, using Nikon A1, C2 confocal microscope

(Tokyo, Japan) with the 20X or 60X oil objective. The whole brain was positioned so

that they can be scanned anteriorly to posteriorly (top to bottom). For statistical

analyses of mitochondrial area and volume, original and unmodified images were

imported into Imaris 9.6 (Oxford Instruments) using Surface to create parameters to

batch data from different sample groups. For statistical analyses of LD number and

intensity, original and unmodified images were imported into ImageJ (National

Institutes of Health), and the intensity threshold for the relevant channel was set so

that maximum number of dots were selected without miscounting two adjacent dots

into one. For colocalization measurements, primary images were imported into

ImageJ (National Institutes of Health) or Imaris 9.6 (Oxford Instruments), and

analyzed automatically using the colocalization plugin with the intensity correlation

tool, showing as Pearson's correlation coefficient (R) or colocalized trace. All data

were imported into GraphPad Prism 7 and shown in scatter plots or column bar graphs.

For calculating the statistical significance, two-tailed unpaired t-test or ordinary

one-way ANOVA followed by Tukey's multiple comparisons test was used. P value

less than 0.05 is considered significant. ns: no significance, $p \geq 0.05$; *: $p<0.05$; **:

$p<0.01$; ***: $p<0.001$; ****: $p<0.0001$.

References

1.   A. R. Fenton, T. A. Jongens, E. L. F. Holzbaur, Mitochondrial dynamics:
Shaping and remodeling an organelle network. *Curr Opin Cell Biol* **68**, 28-36
(2020).
2.   S. Gao, J. Hu, Mitochondrial Fusion: The Machineries In and Out. *Trends Cell*
Biol **31**, 62-74 (2021).
3.   S. Mattie, M. Krols, H. M. McBride, The enigma of an interconnected
mitochondrial reticulum: new insights into mitochondrial fusion. *Curr Opin*
Cell Biol **59**, 159-166 (2019).
4.   D. H. Cho, T. Nakamura, S. A. Lipton, Mitochondrial dynamics in cell death
and neurodegeneration. *Cell Mol Life Sci* **67**, 3435-3447 (2010).
5.   K. Itoh, K. Nakamura, M. Iijima, H. Sesaki, Mitochondrial dynamics in
neurodegeneration. *Trends Cell Biol* **23**, 64-71 (2013).
6.   J. R. Friedman *et al.*, ER tubules mark sites of mitochondrial division. *Science*
334, 358-362 (2011).
7.   R. Kalia *et al.*, Structural basis of mitochondrial receptor binding and
constriction by DRP1. *Nature* **558**, 401-405 (2018).
8.   S. C. Kamerkar, F. Kraus, A. J. Sharpe, T. J. Pucadyil, M. T. Ryan,
Dynamin-related protein 1 has membrane constricting and severing abilities
sufficient for mitochondrial and peroxisomal fission. *Nat Commun* **9**, 5239
(2018).

9.   H. Otera *et al.*, Mff is an essential factor for mitochondrial recruitment of
Drp1 during mitochondrial fission in mammalian cells. *J Cell Biol* **191**,
1141-1158 (2010).
10.   C. S. Palmer *et al.*, Adaptor proteins MiD49 and MiD51 can act independently
of Mff and Fis1 in Drp1 recruitment and are specific for mitochondrial fission.
J Biol Chem **288**, 27584-27593 (2013).
11.   Y. Kageyama *et al.*, Mitochondrial division ensures the survival of postmitotic
neurons by suppressing oxidative damage. *J Cell Biol* **197**, 535-551 (2012).
12.   M. E. Gegg *et al.*, Mitofusin 1 and mitofusin 2 are ubiquitinated in a
PINK1/parkin-dependent manner upon induction of mitophagy. *Hum Mol*
Genet **19**, 4861-4870 (2010).
13.   A. Tanaka *et al.*, Proteasome and p97 mediate mitophagy and degradation of
mitofusins induced by Parkin. *J Cell Biol* **191**, 1367-1380 (2010).
14.   E. Ziviani, R. N. Tao, A. J. Whitworth, Drosophila parkin requires PINK1 for
mitochondrial translocation and ubiquitinates mitofusin. *Proc Natl Acad Sci U*
S A **107**, 5018-5023 (2010).
15.   Y. C. Wong, W. Peng, D. Krainc, Lysosomal Regulation of Inter-mitochondrial
Contact Fate and Motility in Charcot-Marie-Tooth Type 2. *Dev Cell* **50**,
339-354 e334 (2019).
16.   Y. C. Wong, D. Ysselstein, D. Krainc, Mitochondria-lysosome contacts
regulate mitochondrial fission via RAB7 GTP hydrolysis. *Nature* **554**, 382-386
(2018).
17.   Y. Zhang *et al.*, An RNA-sequencing transcriptome and splicing database of
glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**,
11929-11947 (2014).
18.   Y. Zhang *et al.*, Purification and Characterization of Progenitor and Mature
Human Astrocytes Reveals Transcriptional and Functional Differences with
Mouse. *Neuron* **89**, 37-53 (2016).
19.   W. Hong, SNAREs and traffic. *Biochim Biophys Acta* **1744**, 120-144 (2005).
20.   X. Tian, J. Teng, J. Chen, New insights regarding SNARE proteins in
autophagosome-lysosome fusion. *Autophagy*
10.1080/15548627.2020.1823124, 1-9 (2020).
21.   C. Mauvezin, P. Nagy, G. Juhasz, T. P. Neufeld, Autophagosome-lysosome
fusion is independent of V-ATPase-mediated acidification. *Nat Commun* **6**,
7007 (2015).
22.   T. Yoshimori, A. Yamamoto, Y. Moriyama, M. Futai, Y. Tashiro, Bafilomycin
A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification
and protein degradation in lysosomes of cultured cells. *J Biol Chem* **266**,
17707-17712 (1991).
23.   G. A. Smith *et al.*, Glutathione S-Transferase Regulates Mitochondrial
Populations in Axons through Increased Glutathione Oxidation. *Neuron* **103**,
52-65 e56 (2019).
24.   Y. Liu, H. Bao, W. Wang, H. Y. Lim, Cardiac Snail family of transcription
factors directs systemic lipid metabolism in Drosophila. *PLoS Genet* **15**,

e1008487 (2019).

25. R. Senos Demarco, B. S. Uyemura, C. D'Alterio, D. L. Jones, Mitochondrial fusion regulates lipid homeostasis and stem cell maintenance in the Drosophila testis. *Nat Cell Biol* **21**, 710-720 (2019).

26. V. Budnik, K. White, Catecholamine-containing neurons in Drosophila melanogaster: distribution and development. *J Comp Neurol* **268**, 400-413 (1988).

27. D. R. Nassel, K. Elekes, Aminergic neurons in the brain of blowflies and Drosophila: dopamine- and tyrosine hydroxylase-immunoreactive neurons and their relationship with putative histaminergic neurons. *Cell Tissue Res* **267**, 147-167 (1992).

28. M. B. Feany, W. W. Bender, A Drosophila model of Parkinson's disease. *Nature* **404**, 394-398 (2000).

29. P. L. Chen *et al.*, Vesicular transport mediates the uptake of cytoplasmic proteins into mitochondria in Drosophila melanogaster. *Nat Commun* **11**, 2592 (2020).

30. C. M. Deus, K. F. Yambire, P. J. Oliveira, N. Raimundo, Mitochondria-Lysosome Crosstalk: From Physiology to Neurodegeneration. *Trends Mol Med* **26**, 71-88 (2020).

31. K. Arasaki *et al.*, A role for the ancient SNARE syntaxin 17 in regulating mitochondrial division. *Dev Cell* **32**, 304-317 (2015).

32. G. L. McLelland, S. A. Lee, H. M. McBride, E. A. Fon, Syntaxin-17 delivers PINK1/parkin-dependent mitochondrial vesicles to the endolysosomal system. *J Cell Biol* **214**, 275-291 (2016).

33. A. Verkhratsky, M. Matteoli, V. Parpura, J. P. Mothet, R. Zorec, Astrocytes as secretory cells of the central nervous system: idiosyncrasies of vesicular secretion. *EMBO J* **35**, 239-257 (2016).

34. H. Li, Y. Li, Z. Lei, K. Wang, A. Guo, Transformation of odor selectivity from projection neurons to single mushroom body neurons mapped with dual-color calcium imaging. *Proc Natl Acad Sci U S A* **110**, 12084-12089 (2013).

A *HRP; repo-Gal4*

LacZ | *vamp7-RNAi⁴³⁵⁴³*

B *HRP; repo-Gal4* Day 10

Mitochondrial ratio

■ *LacZ*
■ *vamp7-RNAi⁴³⁵⁴³*

<2.5   >2.5

Length/Diameter (L/D)

C *mito.GFP; repo-Gal4*

Luc-RNAi | *vamp7-RNAi³⁸³⁰⁰* | *vamp7-RNAi⁴³⁵⁴³*

GFP

D *mito.GFP; repo-Gal4*

Mitochondrial area (μm²)

Luc-RNAi | vamp7-RNAi³⁸³⁰⁰ | vamp7-RNAi⁴³⁵⁴³
10 | 10 | 10

E *mito.GFP; repo-Gal4*

Mitochondrial volume (μm³)

Luc-RNAi | vamp7-RNAi³⁸³⁰⁰ | vamp7-RNAi⁴³⁵⁴³
10 | 10 | 10

F *mito.GFP; repo-Gal4,Luc-RNAi*

- | Bafilomycin A1 (4 μM)

GFP

G *mito.GFP; repo-Gal4*

Mitochondrial area (μm²)

ns

Bafilomycin A1   -   4 μM
11 | 26

H *mito.GFP; repo-Gal4*

Mitochondrial volume (μm³)

ns

Bafilomycin A1   -   4 μM
11 | 26