VAMP7-dependent mitochondria-lysosome contacts contribute to glial mitochondrial dynamics and dopaminergic neuron survival

Honglei Wang[1,4], Mengxiao Wang[4], Yu-Ting Tsai[2], Yi-Hsuan Cheng[2], Yu-Tung Lin[2], Chia-Ching Lin[2], Yi-Hua Lee[2], Linfang Wang[4,5], Shuanglong Yi[4,6], Shiping Zhang[4,7], Yufeng Pan[1,*], Margaret S. Ho[2,3,4,*]

[1]The Key Laboratory of Developmental Genes and Human Disease, School of Life Science and Technology, Southeast University, Nanjing, 210096, China
[2]Institute of Neuroscience, National Yang Ming Chiao Tung University, Taipei, Taiwan
[3]Brain Research Center, National Yang Ming Chiao Tung University, Taipei, Taiwan
[4]School of Life Science and Technology, ShanghaiTech University, Shanghai, 201210, China
[5]The Institute of Seed Industry, Xianghu Laboratory, Qiantang River International Innovation Belt of the Xiaoshan Economic and Technological Development Zone, Hangzhou, 311200, China
[6]Shanghai Institute of Thoracic Oncology, Shanghai Chest Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200030, China
[7]State Key Laboratory of Medical Neurobiology and MOE Frontiers Center for Brain Science, School of Life Sciences, Fudan University, Shanghai 200438, China

[*]Corresponding authors:
Margaret S. Ho, Ph.D. Email: margaret.ho@nycu.edu.tw
Yufeng Pan, Ph.D. Email: pany@seu.edu.cn

Summary

Disrupted mitochondrial dynamics in neurons are linked to neurodegenerative diseases, but their regulation in glia remains poorly understood. Here, we show that the R-SNARE protein VAMP7 regulates the untethering of mitochondria–lysosome contacts (MLCs) in adult fly glia. Glial-specific knockdown of VAMP7 leads to prolonged MLCs and mitochondrial elongation related to altered fission/fusion dynamics. These VAMP7-deficient mitochondria exhibit hyperpolarized membrane potential, resulting in increased ROS activity, lipid droplet accumulation, and dopaminergic (DA) neuron degeneration. Mechanistically, VAMP7 interacts with the GTPase-activating protein TBC1D15-17 to promote Rab7 GTP hydrolysis. Without VAMP7, TBC1D15-17 remains bound to Rab7 but fails to activate its hydrolysis, resulting in elevated GTP-bound Rab7 and impaired MLCs untethering. Consistently, expression of GTP-locked Rab7^{Q67L} or GAP-dead TBC1D15-17$^{\Delta GAP}$ phenocopies the mitochondrial defects, while GDP-bound Rab7^T22N or wild-type TBC1D15-17 restores the MLC dynamics. Considering that SNARE proteins mediate membrane fusion, our results demonstrate a new role for VAMP7 in glial mitochondrial dynamics via organelle contacts, impacting neuron survival in a non-cell-autonomous manner.

Keywords

VAMP7; Glia; Mitochondrial Dynamics; Mitochondrial Elongation; Mitochondria-Lysosome Contacts; Dopaminergic Neuron Survival; Neurodegeneration

Introduction

Mitochondria are dynamic organelles that undergo structural and functional changes in response to a variety of cellular signals such as energy production, metabolism, or survival. They constantly remodel and interact with other organelles so that their morphologies are swiftly modified to adapt the cellular needs. Different aspects of mitochondrial dynamics have been uncovered including fission (division), fusion, or shape transition, generating different forms of mitochondria that function in various cellular contexts[1, 2, 3]. These dynamic changes in morphology and function are under tight regulation and have been linked to catastrophic consequences such as neurodegeneration and death[4, 5]. For instance, mitochondrial fission regulated by the dynamin-related protein-1 (Drp1), a GTPase that dynamically associates with the endoplasmic reticulum (ER) and mitochondria[6, 7, 8, 9], not only occurs in healthy mitochondria to accommodate physiological needs, but also suppresses oxidative damage and safeguards neuron survival[10]. It is generally considered that the fission process generates smaller mitochondria easier to be targeted by mitophagy, the autophagosomal degradation of damaged mitochondria mediated by the Parkinson's disease risk genes *PINK1* and *Parkin*[11, 12, 13]. On the other hand, elongated mitochondria lacking fission accumulate oxidative damage and lose respiratory function, transforming into large spheres resistant to mitophagy. Interestingly, most of these mitochondrial dynamic defects underlying neurodegeneration are ascribed to neurons, whereas how mitochondrial dynamics in the brain cells "glia" contribute to neurodegeneration remains poorly understood.

In addition to ER, mitochondrial fission sites are also contacted by lysosomes, which help to restrict the mitochondrial motility by recruiting the lysosomal GTPase Rab7[14, 15]. It has been shown that Rab7 GTP hydrolysis, converting the Rab7-GTP to the inactive GDP form, is crucial for the untethering of lysosome and mitochondria, marking the site for mitochondrial fission[15]. Rab7 GTP hydrolysis is regulated by guanine nucleotide exchange factors (GEFs) that stimulate the exchange of bound GDP for GTP, resulting in Rab7 activation, and GTPase activating proteins (GAPs) that facilitate Rab7 GTP hydrolysis, thereby inactivating Rab7[16]. These findings implicate a role for lysosome in regulating mitochondrial dynamics. In regard to this aspect, whether lysosome and its related factors play a role in glial mitochondrial fission is largely unknown; how mitochondrial dynamic is controlled in glia has not been extensively studied.

The *Drosophila* vesicular R-SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptors), VAMP7, is homologous to mammalian VAMP7 and VAMP8, and the latter exhibits a prominent expression in microglia[17, 18]. Mammalian VAMP7 or VAMP8 localizes to the late endosome and lysosome and forms a complex with the cytosolic Synaptosomal-associated protein 29 (SNAP-29) and the autophagosomal Syntaxin-17 (STX17). Together, this complex regulates the autophagosome-lysosome fusion in autophagy[19, 20]. Despite a role in vesicle fusion,

whether VAMP7 plays additional roles with respect to its lysosomal origin remains
unclear. Considering the lysosomal impact on mitochondrial dynamics and a possible
role of VAMP7 in glia, it is worthy of investigating whether the lysosomal VAMP7
regulates mitochondrial dynamics in glia. Our findings reveal that VAMP7 regulates
the untethering of mitochondria-lysosome contacts (MLCs) in adult fly glia. Reduced
VAMP7 expression in glia causes fission/fusion-dependent mitochondrial elongation,
likely owing to the persistent and prolonged MLCs. These elongated mitochondria are
dysfunctional with hyperpolarized membrane potential, increased ROS levels, and the
production of lipid droplets, ultimately leading to dopaminergic (DA)
neurodegeneration in the brain. Mechanistically, VAMP7 interacts with the Rab7
GAP TBC1D15-17, facilitating Rab7 GTP hydrolysis to control the dynamics of
MLCs. Considering that SNARE proteins mediate membrane fusion, our findings
uncover a new role for a canonical SNARE factor in glial mitochondrial dynamics,
demonstrating its importance in glial contribution to neuron survival.
145
Results
Adult fly glia deficient of VAMP7 exhibit mitochondrial elongation
To investigate VAMP7 function in glia, two independent fly RNAi lines:
vamp7-RNAi[38300] and *vamp7*-RNAi[43543] were used to silence *vamp7* expression in
adult fly glia. To assess the efficacy of these lines, 3-day-old adult fly brains were first
collected to analyze *vamp7* mRNA levels upon RNAi expression in glia. RT-qPCR
analyses revealed a significant decrease in *vamp7* mRNA levels, with
vamp7-RNAi[43543] causing a greater reduction (Figure S1A). To examine *vamp7*
mRNA levels only in adult fly glia, 100 adult fly brains expressing *UAS-mCD8.GFP*,
with or without *vamp7*-RNAi, under the control of the pan-glial driver *repo*-GAL4
were dissected and sorted by fluorescence-activated cell sorting (FACS). Consistently,
RT-qPCR analyses revealed significantly decreased *vamp7* mRNA levels in sorted glia
upon expressing *vamp7*-RNAi[43543] (Figure S1B). Hence, *vamp7*-RNAi[43543] was used
as the major RNAi line throughout the study.
We then investigated the consequences of silencing glial *vamp7* expression in the
brains by Transmission Electron Microscopy (TEM). A marker *UAS-HRP* was
expressed in glia (*repo>HRP*), attempting to label the glial processes visualized in
black. Mitochondria were identified for their typical structures of inner and outer
mitochondrial membranes, with the inner membranes folding as cristae and extending
into the matrix (Figure 1A). Compared to the 3-day-old control brains expressing the
Luciferase RNAi (*luc*-RNAi), mitochondria were elongated when expressing
vamp7-RNAi[43543] in glia, indicating a possible defect of fission or fusion (enclosed by
white dashed lines, Figure 1A). The value of mitochondrial length over diameter (L/D)
was significantly increased (Figure 1B). In addition, the ratio of mitochondria with a
L/D number lower than 2.5 (L/D<2.5/Total) decreased, while the ratio of those with a
L/D number higher than 2.5 (L/D>2.5/Total) increased upon glial VAMP7 reduction
(Figure 1C).
To verify these results, two independent mitochondrial reporter lines:
UAS-mito.HA.GFP (*mito.GFP*) and *UAS-mCherry.mitoOMM* (*mCherry.mitoOMM*),

which express a mitochondrial import signal and label the outer mitochondrial
membrane, respectively, were used to assess the mitochondrial morphology in glia
silencing *vamp7* expression. Consistent with the EM results, glial VAMP7-depleted
mitochondria were elongated with significantly increased area in 3-day-old adult fly
brains (Figures 1D-1H). A consistent increase was also detected when using a
different quantification method to assess the mitochondrial volume (Figures S1C and
S1D, please see Methods for quantification details). For consistency, the
dorsal-medial region of the adult fly brain was analyzed throughout the study.
Interestingly, reintroducing VAMP7 by expressing a Myc-tagged *vamp7* transgene in
glia suppressed the mitochondrial elongation upon silencing glial *vamp7* (Figures 1I
and 1J). Interestingly, neuronal mitochondria labeled by expressing
LexAOP-mCherry.mitoOMM under the control of the pan-neuronal driver
GMR57C10-GAL4 remain largely intact with normal area and morphology when
simultaneously expressing *vamp7*-RNAi[43543] in glia (Figures 1K and 1L). Thus, these
results indicate that VAMP7 regulates mitochondrial morphology in adult fly glia.
190
Mitochondrial elongation in VAMP7-deficient adult fly glia depends on the
expression of mitochondrial fission and fusion factors
Given that mitochondrial elongation is linked to altered fission or fusion, we next
asked if glial VAMP7 genetically interacts with the fission factor Dynamin-related
protein 1 (Drp1) or the fusion factors Mitochondrial assembly regulatory factor (Marf)
and Optic atrophy 1 (Opa1). Consistent with previous findings in other cell types,
Drp1 overexpression (*drp*-O/E, increased fission) or downregulated Marf or Opa1
expression by RNAi (*marf*-RNAi or *opa1*-RNAi, decreased fusion) in glia caused a
reduction in the mitochondrial area, indicative of their function in fission and fusion
(Figures 1M and 1N)[21, 22]. Interestingly, *drp*-O/E, *marf*-RNAi, or *opa1*-RNAi
expression significantly suppressed the mitochondrial elongation in the
vamp7-RNAi[43543]-expressing glia (Figures 1O and 1P), suggesting that altered
fission/fusion helps prevent mitochondrial elongation induced by silencing *vamp7*
expression in glia. These results suggest that VAMP7-mediated mitochondrial
dynamics in adult fly glia are fission/fusion dependent.
206
VAMP7 regulates the dynamics of mitochondria-lysosome contacts in adult fly
glia
It has been reported that lysosomes contact mitochondria to restrict their motility and
determine the fission site[15, 23]. Given the lysosomal localization of VAMP7 and the
VAMP7-mediated mitochondrial elongation, we next investigated whether VAMP7
regulates mitochondrial morphology via the mitochondria-lysosome contacts (MLCs)
in adult fly glia. Two reporters: *mCherry.mitoOMM* and *UAS-Lamp1.GFP*
(*Lamp1.GFP*) were co-expressed to label mitochondria and lysosomes, respectively,
so to visualize glial MLCs in live brains. Images acquired during the time-lapse
analysis revealed that mCherry and GFP fluorescent puncta were in close contact and
then dissociated in the control glia at t=120s (white arrow, Figure 2A and Movie S1).
Whereas dissociation between the mCherry and GFP fluorescent puncta was also

detected at t=100s in the control glia in a parallel experiment (white arrow, top panel
in Figure 2B), persistent and prolonged contacts were observed in glia expressing
either *vamp7*-RNAi, up to t=200s (middle and bottom panels in Figure 2B and Movies
S2-S4). The frequency and duration of mCherry-positive mitochondria and
GFP-positive lysosomes contacting each other increased significantly when reducing
glial *vamp7* expression (Figures 2B-2D). Consistent with the observations on
mitochondrial morphology, reintroducing VAMP7 in glia suppressed these contact
deficits. Whereas MLCs in the control glia expressing two *UAS* transgenes (*LacZ*;
luc-RNAi) began to dissociate at t=80s, MLCs in the *vamp7*-RNAi[43543]-expressing
glia remained persistently associated (middle panel in Figure 2E). Upon VAMP7
co-expression in glia, MLCs dissociation was restored and occurred faster at t=170s
(white arrows, bottom two panels in Figure 2E). The frequency and duration of these
MLCs were also suppressed by VAMP7 co-expression (Figures 2E-2G and Movies
S5-S7). Taken together, these results suggest that VAMP7 regulates the dynamics of
MLCs in adult fly glia.
234

GTP-bound Rab7 levels increase upon silencing glial *vamp7* expression
Previous studies have demonstrated that the untethering of MLCs depends on Rab7
GTP hydrolysis[15]. We next checked if VAMP7 regulates the dynamics of MLCs via
Rab7 GTP hydrolysis. In light of the reported GST-Glutathione Transferase-Rab
Interacting Lysosomal Protein (RILP) assay for measuring the GTP-bound Rab7
levels[24], the gene encoding RILP in flies, *RILP-like* (*RILPL*), was fused with a
N-terminal three tandem repeat HA tag (*3xHA-RILPL*) and cloned for transgenic
expression in flies. RILPL preferentially binds to and specifically reflects GTP-bound
Rab7 levels. By co-immunoprecipitation (Co-IP) analysis, 3-day-old adult fly brains
expressing 3xHA-RILPL in glia were dissected and homogenized for further
pull-downs using beads conjugated to the anti-HA antibodies. Validating the approach,
expression of the constitutively active form of Rab7, 3xFlag-Rab7^{Q67L}, resulted in
increased levels of GTP-bound Rab7 as quantified by the ratio of Rab7 in the
3xHA-RILPL pull downs over total Rab7 in the input (Figures 3A and 3B). Notably,
GTP-bound Rab7/total Rab7 ratio also increased upon glial VAMP7 reduction,
suggesting that silencing VAMP7 disrupts Rab7 GTP hydrolysis, leading to increased
GTP-bound Rab7 levels (Figures 3A and 3B).
Previously reported Rab7 effectors include RILP (used in the present study) and
two other related proteins PLEKHM1 and Rubicon[25]. Of these, *Drosophila*
PLEKHM1 (CG6613) was shown to interact with Rab7 in a GTP-dependent manner[26].
Based on these findings, transgenic flies expressing PLEKHM1-GFP were made for
testing the GTP-bound Rab7 levels in flies. Consistent with the RILPL results,
PLEKHM1-GFP levels increased as quantified by GFP intensities and area when
expressing *vamp7*-RNAi[43543] or 3xFlag-Rab7^{Q67L}, indicating more GTP-bound Rab7
(Figures 3C-3E). Interestingly, co-expression of 3xFlag-Rab7^{WT} or the constitutively
active 3xFlag-Rab7^{Q67L} fails to impact the increased GTP-bound Rab7 levels upon
glial VAMP7 reduction, whereas co-expression of the GDP-bound 3xFlag-Rab7^{T22N}
suppressed the increase (Figures 3F-3H). These results suggest that glial VAMP7

regulates Rab7 GTP hydrolysis as reflected by the level of two Rab7 effectors
3xHA-RILPL and PLEKHM1-GFP; supplying more GDP-bound Rab7 helps restore
the Rab7 GTP/GDP balance *in-vivo*.
266

VAMP7 regulates the dynamics of MLCs via Rab7 GTP hydrolysis in adult fly
glia
Given that the untethering of MLCs depends on Rab7 GTP hydrolysis and glial
VAMP7 regulates both the dynamics of MLCs and the GTP-bound Rab7 levels, we
next asked if VAMP7 regulates the untethering of MLCs via Rab7 GTP hydrolysis.
By conducting the *in-vivo* time lapse analysis using *mCherry.mitoOMM* and
Lamp1.GFP, the glial MLCs were monitored under various genetic conditions
(Figures 4A-4C and Movies S8-S12). The impaired untethering of MLCs was
consistently detected in adult fly glia with reduced *vamp7* expression. Interestingly,
only the co-expression of $Rab7^{T22N}$, but not $Rab7^{WT}$ nor $Rab7^{Q67L}$, significantly
restored the prolonged contact dynamics in the absence of glial VAMP7. As the
control MLCs untethered at around t=180s (white arrow in top panel, Figure 4A), the
frequency and duration of the contacts were restored when co-expressing $Rab7^{T22N}$
and *vamp7*-RNAi[43543] (t=140s, white arrows in bottom two panels, Figures 4A-4C).
Taken together, these results suggest that increasing GDP-bound Rab7 levels restores
the disrupted dynamics of MLCs upon glial VAMP7 reduction; VAMP7 regulates the
balance of Rab7 GTP/GDP levels attributing to the untethering of the MLCs in adult
fly glia.
As the untethering of MLCs underlies mitochondrial fission, we also analyzed
mitochondrial morphology when altering Rab7 GTP hydrolysis when silencing *vamp7*
expression in glia. Interestingly, expression of $Rab7^{Q67L}$ (i.e., increased GTP-bound
Rab7 levels), but not $Rab7^{WT}$ nor the GDP-bound $Rab7^{T22N}$, caused mitochondrial
elongation in adult fly glia (Figures 4D and 4E). Furthermore, co-expression of
$Rab7^{T22N}$, but not $Rab7^{WT}$ nor $Rab7^{Q67L}$, restored the *vamp7* RNAi[43543]-induced
mitochondrial elongation in glia (Figures 4F and 4G). These results are consistent
with the observations on the untethering of the MLCs, suggesting that the balance of
Rab7 GTP/GDP levels is crucial for VAMP7-mediated mitochondrial dynamics in
adult fly glia.
295

VAMP7 interacts with the Rab7 GAP TBC1D15-17
Given that VAMP7 regulates GTP-bound Rab7 levels, we next asked if VAMP7
interacts with Rab7 or its associated GTPase-activating protein (GAP) TBC1D15-17,
a member of the TBC (Tre2/Bub2/Cdc16)-domain-containing protein family that
facilitates Rab7 GTP hydrolysis. TBC1D15-17 is a fly protein homologous to
mammalian TBC1D15 and TBC1D17 with overall 30% identity and 63% and 61%
similarity, respectively[27]. Mammalian TBC1D15 has been shown to localize to
mitochondria via binding with the outer mitochondrial membrane protein Fis1, then
drives Rab7 GTP hydrolysis to untether the MLCs[15]. To test interaction among
VAMP7, TBC1D15-17, and Rab7, transgenic flies expressing these proteins with
different   tags   in   tandem   repeats   were   generated.   These   flies   carried

UAS-3xHA-TBC1D15-17, *UAS-6xMyc-vamp7*, or *UAS-3xFlag-Rab7* to express the
corresponding proteins. Different combinations of these proteins were then
co-expressed in flies for Co-IP analyses. Using antibodies targeting different tags, a
strong interaction between VAMP7 and TBC1D15-17 was detected, but not between
VAMP7 and Rab7 (Figures 5A and 5B). While a strong interaction was also detected
between TBC1D15-17 and Rab7, the interaction remains unaffected and strong upon
vamp7-RNAi[43543] expression in glia (Figures 5C and 5D). These results suggest that
VAMP7 physically interacts with TBC1D15-17 and both are within the same protein
complex. It is feasible that VAMP7-TBC1D15-17 interaction might be a prerequisite
for TBC1D15-17 to mediate Rab7 GTP hydrolysis for the untethering of MLCs.
Notably, TBC1D15-17 remains associated with Rab7 upon glial VAMP7 reduction,
suggesting that VAMP7 is not required for their physical interaction; rather, VAMP7
interaction with TBC1D15-17 might promote its GAP activity.
320
Expression of TBC1D15-17 lacking the GAP activity increases GTP-bound Rab7
levels, impairs the untethering of MLCs, and causes mitochondrial elongation in
adult fly glia
TBC1D15-17 contains multiple domains including a TBC domain that mediates GAP
activity and an uncharacterized disordered domain (Figure 5E). Given that mutations
in mammalian TBC1D15 disrupting its GAP activity, like D397A or R400K in the
TBC domain, are not conserved in fly TBC1D15-17[15], transgenic flies expressing the
truncated   TBC1D15-17   lacking   either   the   entire   TBC   domain
(*UAS-3xFlag-TBC1D15-17$^{\Delta GAP}$*)   or   the   uncharacterized   disordered   domain
(*UAS-3xFlag-TBC1D15-17$^{\Delta disordered}$*)   were   generated.   Consistent   with   the
vamp7-RNAi results, TBC1D15-17$^{\Delta GAP}$ expression caused increased intensities and
area of PLEKHM1-GFP in adult fly glia, whereas TBC1D15-17$^{\Delta disordered}$ failed to do
so (Figures 5F-5H). TBC1D15-17 expression also caused an increase in
PLEKHM1-GFP   levels,   suggesting   that   other   regulations   exist   for
TBC1D15-17-mediated Rab7 GTP hydrolysis. Consequently, the dynamics of MLCs
were analyzed upon expression of different TBC1D15-17 variants in adult fly glia.
Whereas   the   control   MLCs   untethered   around   t=110s,   MLCs   expressing
TBC1D15-17^{WT} or TBC1D15-17$^{\Delta disordered}$ untethered at a time scale similar to that of
the control, at t=140s or 170s, respectively (white arrows in top three panels, Figure
5I). On the other hand, TBC1D15-17$^{\Delta GAP}$ expression significantly increased the
frequency and duration of the MLCs (Figures 5I-5K). Consistent with these results,
mitochondrial elongation was also detected upon TBC1D15-17$^{\Delta GAP}$ expression in glia
(Figures 5L and 5M). Taken together, these results suggest that TBC1D15-17 GAP
activity contributes to the untethering of MLCs and mitochondrial dynamics in glia.
345
346 **TBC1D15-17 GAP activity contributes to VAMP7-mediated MLCs untethering**
347 **and mitochondrial dynamics in adult fly glia**
348 We next asked if VAMP7 regulates the dynamics of MLCs and glial mitochondrial
349 elongation via TBC1D15-17 function. While the MLCs expressing two control *UAS*
350 transgenes untethered at around t=130s, the MLCs in VAMP7-deficient glia failed to

untether within the time frame examined (Figures 6A-6C). Interestingly, the MLCs co-expressing *vamp7*-RNAi[43543] and TBC1D15-17[WT] or TBC1D15-17[Δdisordered] started to untether at a similar time scale to the control at around t=90s and 50s, respectively (white arrows in the middle panel, Figure 6A). The increased frequency and duration of the contacts were significantly suppressed upon co-expression, except that the duration of the contacts co-expressing TBC1D15-17[Δdisordered] and *vamp7*-RNAi was decreased without significance (Figures 6B and 6C). Notably, TBC1D15-17[ΔGAP] expression failed to suppress the increase in frequency and duration of the MLCs expressing *vamp7*-RNAi[43543] (Figures 6A-6C). Furthermore, expression of TBC1D15-17[WT] or TBC1D15-17[Δdisordered], but not TBC1D15-17[ΔGAP], significantly suppressed the mitochondrial elongation upon glial VAMP7 reduction (Figures 6D and 6E). Taken together, these results indicate that VAMP7 regulates the dynamics of MLCs and mitochondrial dynamics via interacting with TBC1D15-17 in adult fly glia, with TBC1D15-17 GAP activity contributing to both events.

Reducing *vamp7* expression causes mitochondrial dysfunction in adult fly glia
To further demonstrate that VAMP7 regulation of MLCs and mitochondrial dynamics in glia are functionally relevant, we assessed the levels of various reporters indicative of mitochondrial function. As mitochondria are the major sites for ROS production, control and *repo>vamp7*-RNAi[43543] adult fly brains were examined for ROS levels with the dihydroethidium (DHE) staining. Consistent with previous findings, DHE-positive fluorescent intensities increased when downregulating the expression of superoxide dismutase 1 (SOD1), the enzyme that regulates ROS levels by catalyzing the conversion of superoxide into oxygen and hydrogen peroxide (Figure 7A). Interestingly, *vamp7*-RNAi[43543] expression in glia caused a similar increase, suggesting that glial VAMP7 regulates cellular ROS levels (Figure 7A). In addition, the transcription levels of genes related to the mitochondrial fatty acid β-oxidation pathway, including those encoding acyl-CoA dehydrogenases (*CG17544* and *CG9527*) and fatty acid synthase (*FASN*), were increased in adult fly brains expressing *vamp7*-RNAi[43543] in glia[28, 29] (Figure 7B). Notably, the levels of BODIPY-positive LDs increased in both larval and adult fly brains upon glial VAMP7 reduction (Figures 7C-7E), with a similar increase in the overall LD triacylglycerol (TAG) content (Figure 7F). The increased LD number was reduced upon adding the potent antioxidant α-Lipoic acid in 3-day-old adult fly brains (Figures 7G and 7H). Taken together, these results indicate that glial VAMP7 regulates mitochondrial function, ROS levels, and LD production in the brain.

We also investigated mitochondrial function by analyzing the activity of the reporter *UAS-mito-roGFP2-Orp1*[30]. Interestingly, the 405/408 intensities reflecting the H_2O_2 reduction-oxidation activity in the mitochondrial matrix were significantly increased in 10-day-old adult fly glia expressing Rab7[Q67L] or *vamp7*-RNAi[43543] (Figures 7I and 7K). Staining the brains with the dye mitotrackerCMXROS, indicative of the living mitochondria membrane potential, also revealed increased intensities upon Rab7[Q67L] or *vamp7*-RNAi[43543] expression in glia (Figures 7J and 7L). Results from these quantitative analyses indicate that VAMP7 regulates mitochondrial

function in glia. Importantly, both the 405/408 intensities and the mitotracker dye
levels were suppressed when co-expressing the GDP-bound Rab7^{T22N} or
TBC1D15-17^{WT} in glia (Figures 7M-7P), suggesting that altering Rab7 GTP
hydrolysis by increasing GDP-bound Rab7 or Rab7 GAP activity in VAMP7-deficient
glia reestablishes the balance of Rab7 GTP/GDP levels, hence recovering the
mitochondrial function in adult fly glia.
401
Glial VAMP7 contributes to DA neuron survival
As glia-neuron crosstalk underlies neuronal function and survival, we next
investigated if VAMP7-mediated mitochondrial dynamics in glia contribute to DA
neuron survival. DA neurons are located in clusters and named according to their
relative positions in adult fly brains[31, 32]. These clusters include protocerebral posterior
lateral (PPL)1, PPL2, protocerebral posterior medial (PPM)1, PPM2, and PPM3
(Figure 8A). Interestingly, silencing glial *vamp7* expression caused DA neuron loss at
PPM1/2, PPM3, and PPL1 clusters in an age-dependent manner (white arrows,
Figures 8B-8D). In addition, *vamp7*-RNAi^{43543} expression in glia in the presence of
Elav-GAL80, which inhibits GAL4 expression in neurons, also caused DA neuron
loss in the PPM1/2 cluster, further demonstrating that DA neuron loss is a result
explicitly of glial VAMP7 knockdown (Figures 8E and 8F). Given that DA
neurodegeneration in the PPM1/2 cluster has been implicated in a *Drosophila* PD
model overexpressing α-synuclein[33], it is conceivable that glial VAMP7 plays
important roles in regulating DA neuron survival under pathological conditions.
On the other hand, whereas Rab7^{Q67L} expression in glia caused similar DA neuron
loss in the PPM1/2 cluster (Figures 8G and 8H), co-expression of Rab7^{T22N} or
TBC1D15-17 with *vamp7*-RNAi^{43543} suppressed the degeneration, suggesting that
reestablishing the balance of Rab7 GTP/GDP levels, potentially restoring the
dynamics of MLCs in glia, promotes DA neuron survival (Figures 8I and 8J). Finally,
suppressing the ROS levels by expressing SOD1 in either glia or neurons restored the
DA neurodegeneration in *vamp7*-RNAi^{43543}-expressing brains, suggesting that
modulating ROS levels in either glia or neurons promotes DA neuron survival
(Figures 8K-8N). These results further implicate a role of glial mitochondrial
dynamics in DA neuron survival, as tuning down the ROS levels in glia only also
helps to ameliorate DA neurodegeneration.
428
Discussion
SNARE proteins are canonical regulators of membrane fusion. Our findings uncover a
new role for the SNARE protein VAMP7 in glial mitochondrial dynamics. Silencing
vamp7 expression in glia causes fission/fusion-dependent mitochondrial elongation,
owing to the dysregulated untethering of MLCs. VAMP7 interacts with the Rab7 GAP
TBC1D15-17 and regulates Rab7 GTP/GDP levels, hence modulating the dynamics
of MLCs. Mitochondria in VAMP7-deficient glia are dysfunctional with
hyperpolarized membrane potential, leading to increased ROS activity and lipid
droplet formation in the brain. These deficits endanger nearby DA neurons, linking
glial mitochondrial dynamics to DA neuron survival (Figure 9).

VAMP7 mediates mitochondrial dynamics in glia

As the untethering of MLCs depends on Rab7 GTP hydrolysis, we uncovered that the persistent contact (i.e., dysregulated untethering) induced by decreased *vamp7* expression in glia is linked to a disrupted balance of Rab7 GTP/GDP levels; Rab7 GTP levels are significantly increased upon silencing glial *vamp7* expression. VAMP7 interacts with the Rab7 GAP TBC1D15-17 to promote Rab7 GTP hydrolysis. Without VAMP7, TBC1D15-17 remains associated with Rab7, yet the TBC1D15-17 GAP activity is likely impaired given the increased GTP-bound Rab7 levels. Expression of TBC1D15-17 lacking the GAP activity, like *vamp7*-RNAi, also causes prolonged MLCs and mitochondrial elongation in adult fly glia. These results suggest that VAMP7 interaction with TBC1D15-17 promotes Rab7 GTP hydrolysis, hence regulating the MLCs untethering and mitochondrial dynamics in glia.

VAMP7 mediates glial mitochondrial dynamics independently of autophagosome-lysosome fusion and mitophagy

Consistent to a role of VAMP7 in autophagosome-lysosome fusion, the autophagic flux we monitored using a dual reporter *UAS-GFP-mCherry-Atg8a* when expressing *vamp7*-RNAi in glia was significantly impaired (Figures S2A and S2B). The Pearson's coefficient values for colocalization of GFP and mCherry-positive signals were significantly increased upon glial VAMP7 reduction, indicating non-quenched GFP signals due to impaired fusion (Figure S2B). Nonetheless, the glial mitochondrial area was not altered in the presence of Bafilomycin A1 (BafA1), a drug blocking autophagosome-lysosome fusion (Figures S2C and S2D)[34, 35, 36]. Despite that the STX17-SNAP29-VAMP7 complex regulates autophagosome-lysosome fusion, STX17 inhibition in glia failed to induce mitochondrial elongation (Figures S2E and S2F). DA neuron number also remained unaffected upon STX17 or SNAP29 inhibition in glia (Figures S2G and S2H). Thus, it is likely that VAMP7 regulates the untethering of MLCs independently of its function in the autophagosome-lysosome fusion.

Consistent with the results of disrupting autophagosome-lysosome fusion, inhibiting autophagy by downregulating the expression of *atg1* or *atg8* in glia causes smaller mitochondria, suggesting enhanced mitochondrial fission (Figures S2I and S2J)[37, 38]. Given that *vamp7*-RNAi expression induces mitochondrial elongation in glia, these results reinforce a role for VAMP7 in glial mitochondrial dynamics independent of autophagy. On the other hand, whereas the colocalization of Lamp1.GFP- and mitoOMM.mCherry-positive puncta is significantly increased (as shown throughout the study as prolonged contacts, Figure S2K, S2L, and S2N). Atg8a was very minimally localized to the MLCs upon glial VAMP7 reduction (Figures S2K, S2M, and S2O). These results are consistent with the notion that elongated mitochondria are resistant to and slow down mitophagy; reduced glial *vamp7* expression increases the formation of MLCs yet inhibits mitophagy.

VAMP7-mediated mitochondrial dynamics in glia functionally contribute to DA

neuron survival
Our findings further support the hypothesis that MLCs mark the sites for
mitochondrial fission. Upon reduction in glial *vamp7* expression, the untethering of
MLCs is hindered, disrupting subsequent fission, leading to elongated mitochondria,
which is suppressed by the expression of the fission/fusion factors. Furthermore, these
elongated mitochondria are dysfunctional with hyperpolarized membrane potential,
enhanced ROS activity, and lipid droplet formation in the brain. These changes are at
the whole-brain level and affect DA neuron survival. Reducing the ROS activity by
expressing SOD1 in glia or neurons suppresses the DA neurodegeneration, supporting
the notion that VAMP7-mediated mitochondrial dynamics underlie the mechanism of
glia-neuron crosstalk contributing to neuron survival.
494
SNARE proteins in glial mitochondrial dynamics
To date, no evidence indicates that SNARE proteins mediate mitochondrial fusion. A
recent report reveals that the STX17-SNAP29-VAMP7 complex contributes to the
megamitochondria engulfing lysosome (MMEL) under hypoxia[36]. STX17 has also
been reported to regulate the activity of the fission factor Drp1[39, 40]. Notably, STX17
localizes to the ER-mitochondria contacts to execute its function in mitochondrial
fission[39, 41], while we uncover a role for VAMP7 in the untethering of MLCs. Based
on these findings, it is feasible to speculate that SNARE proteins are significant
players in the contacts of different organelles, in addition to their function in
membrane fusion.
Similar to neurons, glia, particularly astrocytes, possess SNARE machineries for
their vesicular trafficking function. These SNARE machineries, some in the same
composition as in neurons and some different, are responsible for slow secretory
exocytosis rather than fast and immediate release of signaling molecules[42]. Very
importantly, they regulate synaptic transmission and plasticity, antigen presentation in
neuro-inflammation, and control different types of behavior. Our findings
unexpectedly reveal a new role for the canonical SNARE protein VAMP7 in glial
mitochondrial dynamics. Considering that mitochondria are present in astrocytic
processes in close proximity to neurons and contribute to local calcium signaling, ATP
production, and metabolism of signaling molecules, it is conceivable that
VAMP7-mediated glial mitochondrial dynamics participate in the metabolic process
supporting neuron survival. In support of the hypothesis, downregulating *vamp7*
expression in the fly astrocyte-like glia also caused DA neuron loss (Figure S3).
Given that DA neurodegeneration is one of the hallmarks in PD, our findings illustrate
a SNARE-dependent mechanism controlling organelle contacts, glial mitochondrial
dynamics, and DA neuron survival. Canonical SNARE proteins not only regulate
membrane fusion, but also are important mediators in mitochondrial dynamics; future
work is warranted to explore additional roles of SNARE proteins in various cellular
contexts in both cell-autonomous and non-cell-autonomous fashions.
524
Acknowledgments
We thank Bloomington *Drosophila* Stock Center, Vienna *Drosophila* RNAi Center,

Tsinghua Fly Center, the Core Facility of *Drosophila* Resource and Technology,
Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences,
Developmental Studies Hybridoma Bank, Chao Tong, Chun-Hong Chen, Henry Sun,
and Aike Guo for fly stocks and antibodies. We also thank the Molecular Imaging
Core Facility (MICF), the Molecular and Cell Biology Core Facility (MCBCF), and
the Multi-Omics Core Facility (MOCF) at the School of Life Science and Technology,
ShanghaiTech University for providing technical support; Shengxiong Wang in
Shanghai JiaoTong University School of Medicine and Yu Kong and Lijun Pan in
Electron Microscopy Facilities of Center for Excellence in Brain Science and
Technology, Chinese Academy of Science for assistance with TEM sample
preparation; The Electron Microscopy Center in the School of Physical Science and
Technology, ShanghaiTech University for providing TEM technical support; DroBot
Biotechnology for quality fly food supply, delicate fly-keeping service, and
experimental device design; Ho lab members and Yan Zou for discussion and
comments. This work was supported by intramural grants from National Yang Ming
Chiao Tung University, National Natural Science Foundation of China (32170962 to
M.S.H, 32371067 to Y.P), Shenzhen Medical Academy of Research and Translation
(B2402005 to Y.P.), 2030 Cross-Generation International Outstanding Young Scholars
Program National Science and Technology Council Taiwan (113-2628-B-A49-003),
and Brain Research Center National Yang Ming Chiao Tung University from The
Featured Areas Research Center Program within the framework of the Higher
Education Sprout Project by the Ministry of Education (MOE) in Taiwan.
549

550 **Author Contributions**
551 H.W and M.S.H conceived and designed the study. H.W, M.W, C.L, Y.L, L.W, S.Y,
552 and S.Z performed the experiments. H.W, Y.T, Y.C, Y.L, Y.P, and M.S.H analyzed the
553 data. M.S.H wrote the paper with input from H.W and Y.P. All authors read and
554 approved the manuscript.
555

556 **Declaration of Interests**
557 The authors declare no competing interests.
558

Materials and Methods
Drosophila **genetics**
Flies were raised under standard conditions at 25□ with 70% humidity. All fly crosses
were carried out at 25°C with standard laboratory conditions unless noted otherwise.
All strains were obtained from Bloomington *Drosophila* Stock Center (BDSC), the
Vienna *Drosophila* RNAi Center (VDRC), Tsinghua Fly Center, or as gifts from
colleagues. The following fly strains were used in this study: *repo-GAL4* (a pan-glial
driver), *UAS-luciferase-IR* (BL31603), *UAS-LacZ* (BL1777), *UAS-mito.HA.GFP*
(BL8442), *UAS-mCherry.mitoOMM* (BL66532), *UAS-Lamp1.GFP* [43],
UAS-vamp7-RNAi (BL38300 and BL43543), *UAS-marf RNAi* (BL31157), *UAS-opa1*
RNAi (THU0811), *UAS-sod1 RNAi* (THU3207), *UAS-mito-roGFP2-Orp1* (BL67667),
UASp-GFP-mCherry-Atg8a (Henry Sun), *UAS-HRP* (Aike Guo), *UAS-drp1* (Chao

Tong), and transgenic flies made in the present study (*UAS-6xMyc-vamp7*,

UAS-3xFlag-Rab7^{wt},   *UAS-3xFlag-Rab7^{Q67L}*,   *UAS-3xFlag-Rab7^{T22N}*,

UAS-3xHA-TBC1D15-17,   *UAS-3xHA-RILPL*,   *LexAOP-3xFlag-sod1*,

UAS-TBC1D15-17^{wt},   *UAS-3xFlag-TBC1D15-17^{Δdisordered}*,

UAS-3xFlag-TBC1D15-17^{ΔGAP}).

Molecular biology

<u>Plasmid cloning</u>: Plasmids were constructed by the polymerase chain reaction (PCR)

using DNA templates from *Drosophila*. These plasmids express wild type Rab7 (621

bp), Rab7 point mutant variants including Rab7^{Q67L} (621 bp, Glutamine 67 to

Leucine), and Rab7^{T22N} (621 bp, Threonine 22 to Asparagine), VAMP7,

TBC1D15-17, RILPL, and PLEKHM1. The corresponding DNA sequences were

subcloned into pUAST-attB vector containing a 3xFlag, 3xHA, or GFP epitope tag by

GenScript (Nanjing, China). *sod1* gene was subcloned into the LexAOP vector

containing a 3xFlag epitope tag by PCR. Primers used for amplification:

Sod1-F: CGACGACAAACTCGAGGTGGTTAAAGCTGTCTGCGET

Sod1-R: ACAAAGATCCTCTAGATTAGACCTTGGCAATGCCAATAACG

Fly microinjection was conducted by the *Drosophila* Core Facility, Institute of

Biochemistry and Cell Biology, Chinese Academy of Sciences.

<u>RT-qPCR</u>: Total RNAs were extracted from adult fly heads using TransZol Up (Cat.

#ET111-01, TransGen, Beijing, China). cDNAs were reversely transcribed using

HiScript III RT SuperMix (Cat. #R323-01, Vazyme, Nanjing, China). RT-qPCR

reactions were performed using ChamQ Universal SYBR qPCR Master Mix (Cat.

#Q711-02, Vazyme) and ABI 7500 RT-PCR system. The mRNA levels were

normalized to *rp49*. The expression levels were analyzed by $\Delta\Delta CT$ method. Primers

used for amplification:

rp49-F: CCACCAGTCGGATCGATATGC

rp49-R: CTCTTGAGAACGCAGGCGACC

vamp7-F: GGCTCGCAATTGGATAGTAGT

vamp7-R: TGCTTATCGGGGAATGAGCC

CG17544-F: *GTACTATGCTCTGACGCGCT*

CG17544-R: *GGTGGTCATGTGCTTCTCCA*

CG9527-F*: CTGCTTCCCCTGCTGAAGAA*

CG9527-R*: GACGTTTCCATCGCTCATGC*

FASN-F: *CGCTCCACTCCAAGAACTCG*

FASN-R*: CAGGTTTAGTTGTAGGGGCTAGA*

FACS sorting of adult fly glia

Approximately 100 *repo>mCD8.GFP* adult fly brains were dissected in fresh

Schneider's Insect Medium (Invitrogen, Carlsbad, CA, USA, Cat# 21720024) and

transferred them to the phosphate-buffered saline (PBS) dissociation buffer which

contains papain (0.18 U/mL; Worthington, Freehold, NJ, USA, Cat# LK003178) and

liberase TM (2.5 mg/mL; Roche, Basel, Switzerland, Cat# 5401119001) to dissociate.

Note that papain requires pre-activation at 37℃ for 15 minutes prior to use. Ice-cold

Schneider's Insect Medium was used to inactivate enzymes. Samples were filtrated by

40 μm cell strainers (BD Falcon, Franklin Lakes, NJ, USA, Cat# 352340) into flow

cytometry tubes (BD Falcon, Cat# 352003) then sorted by the flow cytometer

(FACSMelody, BD Biosciences, Bergen County, NJ, USA).

Immunohistochemistry

Adult fly brains were dissected and fixed in 4% formaldehyde for 45 minutes, then

washed with PBT (PBS + 0.3% TX-100) for 3 times and dissected further to remove

additional debris in PBS solution. Clean and fixed brains were blocked in PBT

solution with 5% Normal Donkey Serum (NDS) or 10% Bovine Serum Albumin

(BSA) and subsequently stained with primary antibodies at 4□ overnight and

secondary antibodies at room temperature for 2 hours. Primary antibodies used:

mouse anti-Atg8a (1:1000, gift from Chao Tong) and rabbit anti-TH (1:1000, Cat.

#NB300-109, Novus). For all samples, secondary antibodies used were from Jackson

ImmunoResearch (West Grove, PA, USA): donkey anti-rabbit Cy3 (1:1000, Cat.

#711-166-152), donkey anti-mouse Cy5 (1:500, Cat. #715-175-150). All samples

were mounted in Vectashield mounting media (Cat. #H-1500-10, Vector Laboratories,

Burlingame, CA, USA).

In-vivo **time lapse analysis**

Live adult fly brains were dissected and kept in PBS solution. All samples were

mounted in PBS, and immediately scanned anteriorly to posteriorly (top to bottom) in

a serial Z-stack of average 5 sections, each of 0.125 μm thickness using Nikon C2

confocal microscope (Tokyo, Japan) with a 63X oil objective in the interval of 10

seconds for 5 minutes. Samples in Figure 6A were scanned in a serial Z-stack of 5

sections, each of 0.395 µm thickness using Zeiss LSM 700 (Oberkochen, Germany)

with a 63X oil objective in the interval of 10 seconds for 5 minutes. The frequency of

Lamp1.GFP and mCherry.mitoOMM contacts was analyzed by Imaris using

autoregressive motion tracking mode. Duration of contacts were analyzed by ImageJ.

Live fluorescent dye imaging

Live adult fly brains were dissected and kept in PBS solution. For the analysis of LDs,

brains were incubated with 1 μg/mL BODIPY™ 493/503 (Cat. #D3922, Invitrogen)

for 5 minutes at 25°C and/or 10 μM α-Lipoic acid (Cat #D118666, Aladdin) for 3

days. For the analysis of mitochondria-derived ROS levels, brains were incubated

with 50 μM DHE (Cat. #D11347, Invitrogen) for 5 minutes at 25°C. For the analysis

of mitochondrial membrane potential, brains were incubated with 500 nM

MitoTracker RedCMXRos (Cat. #M7512, Invitrogen) solution in PBS for 15 minutes

at 25°C. Images were acquired with the Nikon C2 or A1 confocal microscope (Tokyo,

Japan) with a 20X or 60X oil objective.

Transmission electron microscopy

Adult fly brains were fixed, embedded, stained, and dehydrated according to previous
protocols[44]. Ultrathin sections (70 nm) of each brain were cut with a Diatome
diamond knife on a Leica EM UC7 ultra-microtome (Leica Microsystems, Germany)
and collected on copper grids. Images were acquired with Talos L120C transmission
electron microscope (Thermo Fisher Scientific, Waltham, MA, UAS) operating at an
acceleration voltage of 120 kV.
664

Biochemistry
<u>Western blot analysis</u>: Adult fly heads were homogenized using a motorized pestle
(Cat. #116005500, MP Biomedicals, Irvine, CA, USA) in lysis buffer (0.4% NP-40,
0.2 mM EDTA, 150 mM NaCl, 20% glycerol, 100 mM Tris-HCl pH7.5, 2% Tween
20, 0.5 mM phosphodiesterase inhibitors, and 1 mM PMSF). Proteins were separated
on SDS-PAGE gels and transferred to PVDF membranes (Cat. #IPFL00010,
Millipore, Billerica, MA, USA). The membranes were blocked by 5% fat-free milk
diluted in PBS with 0.1% Tween-20 for 30 minutes. Samples were incubated with the
primary antibodies at 4°C overnight and HRP-conjugated secondary antibodies at
room temperature for 2 hours. Primary antibodies used: mouse-anti-HA (1:1000, Cat.
#12C10, Roche), rabbit anti-Myc (1:2000, Cat. #0912-2, Hua An Biotechnology),
mouse anti-Rab7 (1:100, DSHB), rabbit-anti-α-Tubulin (1:5000, Cat. #SB-AB0049,
Share Biotechnology). Bands were visualized by Clarity Western ECL Substrate (Cat.
#WBKLS0500, Millipore). Results are representative of at least three biological
replicates.
<u>Co-immunoprecipitation</u>: The cell lysates were collected as described above for
centrifugation at 12,000 rpm, at 4°C for 30 minutes, the soluble supernatant was
incubated with prewashed anti-HA magnetic beads (Cat. # HY-K0201;
MedChemExpress), anti-c-Myc magnetic beads (Cat.# HY-K0206A,
MedChemExpress) at 4°C overnight. After pull-downs, gels were extensively washed
in lysis buffer for 30 minutes 3 times and then subjected to western blot analysis. The
following antibodies were used for blotting: rabbit-anti-Myc (1:2000, Cat. #0912-2,
Hua An Biotechnology), mouse-anti-Flag (1:1000, Cat. #F3165, Sigma),
mouse-anti-HA (1:1000, Cat. #3F10, Roche), mouse anti-Rab7 (1:100, DSHB) and
rabbit-anti-α-Tubulin (1:5000, Cat. #SB-AB0049, Share Biotechnology).
<u>BafilomycinA1 treatment</u>: BafA1 (Cat. #BML-CM110-0100, ENZO) was diluted in
DMSO (Cat. #D8371-50, solarbio) in a concentration of 1 mM, then added to 0.5 ml
of liquid yeast for a final concentration of 4 μM. Yeast containing BafA1 was added
to the top of the fly food in standard vials kept overnight to allow the evaporation of
residual DMSO. Flies of different genotypes (20 females and 20 males) were then put
in each vial and kept at 25°C for 5 days. After treatment, flies were collected for
imaging analysis. The TAG content in the LD was quantified by an triglyceride assay
kit (Cat #E1103, Applygen Technologies).
698

Confocal microscopy and statistical analysis
Images of adult fly brains were acquired by scanning a serial Z-stack of average 5-20
sections, each of 0.5, 0.8 or 0.125 μm thickness, using Zeiss LSM 700 (Oberkochen,

Germany), Nikon A1 and C2 confocal microscope (Tokyo, Japan) with the 20X or
63X oil objective, respectively. The whole brain was positioned so that they can be
scanned anteriorly to posteriorly (top to bottom). For statistical analyses of the
mitochondrial area, original and unmodified images were imported into ImageJ
(National Institutes of Health), and analyzed automatically using the Tubeness plugin.
Mitochondrial volume was analyzed by Imaris (the Surface module) for 3D
quantifications. For statistical analyses of LD number and intensity, original and
unmodified images were imported into ImageJ, and the intensity threshold for the
relevant channel was set so that maximum number of dots were selected without
miscounting two adjacent dots into one. For colocalization measurements, primary
images were imported into ImageJ, and analyzed automatically using the
colocalization plugin with the intensity correlation tool, showing as Pearson's
correlation coefficient (R) or colocalized trace. All data were imported into GraphPad
Prism 9 and shown in scatter plots or columnal bar graphs. For calculating the
statistical significance, two-tailed unpaired t-test or ordinary one-way ANOVA
followed by Tukey's multiple comparisons test was used. P value less than 0.05 is
considered significant. ns: no significance, $p \geq 0.05$; *: $p<0.05$; **: $p<0.01$; ***:
$p<0.001$; ****: $p<0.0001$.
720

Figure Legends
Figure 1. Mitochondrial elongation in VAMP7-deficient adult fly glia (**A-C**)
Representative TEM images (A) of mitochondria (double-membraned with cristae) in
control and *repo>vamp7*-RNAi[43543] adult fly brains. *UAS-HRP* was also expressed in
glia, attempting to label the glial processes in black. Note the presence of elongated
mitochondria (white dashed lines) upon glial VAMP7 reduction. Quantifications (B
and C) of mitochondrial morphology are designated as mitochondrial length/diameter
(L/D, B) and the ratio of mitochondria with a L/D number higher or lower than 2.5
over the total number (C). Note that the ratio of mitochondria with L/D number
greater than 2.5 is significantly higher upon glial VAMP7 reduction. (**D**) Illustration of
an adult fly brain with symmetrical mushroom body calyces located posteriorly
(green). Dashed square in black indicates the dorsal-medial region where images
throughout the study were acquired. (**E-H**) Representative images (E and H) and
quantifications (F and G) of mitochondrial morphology labeled by *UAS-mito.GFP* (E
and E') or *UAS-mCherry.mitoOMM* (H and H') in control and
vamp7-RNAi[43543]-expressing adult fly glia. Note that the average mitochondrial area
quantified by ImageJ in 2D is significantly larger than the control glia when
expressing *vamp7*-RNAi[43543]. (**I-L**) Representative images (I and L) and
quantifications (J and K) of mitochondrial morphology labeled by *UAS-mito.GFP* (I
and I') or *LexAOP-mCherry.mitoOMM* (L and L') in control and
vamp7-RNAi[43543]-expressing adult fly glia. Note that Co-expression of 6xMyc-*vamp7*
suppresses the elongation (I and J). Neuronal mitochondria were analyzed in adult fly
brains expressing *LexAOP-mCherry.mitoOMM* under the pan-neuronal driver
GMR57C10-LexA and *UAS-vamp7*-RNAi[43543] by *repo*-GAL4. Note that neuronal
mitochondria remain largely intact upon glial VAMP7 reduction. (**M-P**)

Representative images (M,M',O,O') and quantifications (N and P) of glial
mitochondria labeled by *UAS-mito.GFP* (*repo>mito.GFP*) upon expression of the
fission/fusion factors Drp1 (*drp1*-O/E), *marf*-RNAi, or *opa1*-RNAi with or without
VAMP7. Note that increasing expression of the fission factor Drp1 or downregulating
the expression of the fusion factor Marf or Opa1 causes increased fission, opposite to
the mitochondrial elongation phenotype when silencing *vamp7* expression in glia.
Co-expression of *drp1*-O/E, *marf*-RNAi, or *opa1*-RNAi with *vamp7*-RNAi[43543] in glia
suppresses mitochondrial elongation. The selected white squares in E,H,I,L,M,O are
enlarged and aligned in E',H',I',L',M',O'. All samples examined are brains of
3-day-old adult flies unless noted otherwise. For single transgene experiment,
Luciferase-RNAi (*luc*-RNAi) was used as a control. For double transgene experiment,
UAS-LacZ; *luc*-RNAi was used as a control. Scale bars of different sizes are indicated
on the images (TEM=500 nm). For TEM, all mitochondria found in more than 20
brain regions were analyzed per sample. Mitochondrial length, diameter, and area are
quantified by ImageJ as detailed in the Materials and Methods. Serial confocal
Z-stack sections were taken at similar planes across all genotypes, showing
representative single-layer images. Statistical graphs are shown with scatter dots and
the number of brains or mitochondria listed on top. For TEM, one scatter dot
represents the average of mitochondria L/D or ratio analyzed. Data are shown as mean
± SEM. P-values of significance (indicated with asterisks, ns no significance, *
$p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) are calculated by two-tailed
unpaired t-test between two sample groups or by one-way ANOVA among three
sample groups followed by Tukey's multiple comparisons test.
769
Figure 2. VAMP7 regulates the dynamics of mitochondria-lysosome contacts in
adult fly glia (**A-G**) Representative images (**A,B,E**) and quantifications (**C,D,F,G**) on
in-vivo time lapse analysis of mitochondria-lysosome contacts (MLCs) in adult fly
glia expressing two reporters: *UAS-mCherry.mitoOMM* (labels the mitochondrial
outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green). In control glia,
the contacts start untethered at t=120s (white arrow, A) or t=100s (white arrow in top
panel, B), whereas the untethering of contacts is delayed up to t=200s in
vamp7-RNAi[43543]-expressing glia (middle and bottom panels, B). Note that the
frequency and duration of the contacts increase (C and D) in glia expressing either
vamp7-RNAi. In a parallel experiment, 6xMyc-*vamp7* expression restores the contact
dynamics (untethering at t=170s, white arrows in bottom two panels, E). An
overexposed panel is shown for the rescue 6xMyc-*vamp7* genotype for clarity (bottom
panel, E). Note that the frequency and duration of the contacts (F and G) are restored
upon 6xMyc-*vamp7* co-expression in glia. All samples examined are brains of
3-day-old adult flies unless noted otherwise. For single transgene experiment,
luc-RNAi was used as a control. For double transgene experiment, *UAS-LacZ*;
luc-RNAi was used as a control. White arrows indicate the untethering. Scale bars of
different sizes are indicated on the images. Serial confocal Z-stack sections were
taken at similar planes across all genotypes, showing representative merged images.
Statistical graphs are shown with scatter dots and the number of brains (one MLC per

brain) listed on top. Images for the time lapse series are shown in 3x3 μm squares
centering on the target MLC selected from the movies (Movies S1-S7). Parameters for
the contact dynamics in C,D,F,G are quantified using Imaris and ImageJ. Data are
shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no
significance, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) are calculated by
one-way ANOVA followed by Tukey's multiple comparisons test.

Figure 3. GTP-bound Rab7 levels increase upon glial VAMP7 reduction (**A** and **B**)
Representative WB images (A) and quantifications (B) of Rab7-RILPL interaction.
GTP-bound Rab7 are present in the eluates of glial RILPL pull-downs by the
HA-conjugated beads (*repo>3xHA-RILPL*, A). Note that GTP-bound Rab7 levels
increases when silencing *vamp7* expression or overexpressing Rab7^{Q67L} in glia. (**C-H**)
Representative images (C and H) and quantifications (D-G) of transgenic flies
expressing PLEKHM1(CG6613)-GFP. Note that GFP signals exhibit aggregated
pattern in forms of multiple puncta and elevated intensities upon glial VAMP7
reduction. All samples examined are brains of 3-day-old adult flies unless noted
otherwise. For single transgene experiment, *luc*-RNAi was used as a control. For
double transgene experiment, *UAS-LacZ*; *luc*-RNAi was used as a control. Scale bars
of different sizes are indicated on the images. Serial confocal Z-stack sections were
taken at similar planes across all genotypes, showing representative merged images.
Statistical graphs are shown with scatter dots and the number of brains listed on top.
For Co-IP analysis, 200 independent fly brains for each genotype were analyzed. Data
are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no
significance, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) are calculated by
ordinary one-way ANOVA followed by Tukey's multiple comparisons test.

Figure 4. VAMP7 regulates mitochondria-lysosome contacts via Rab7 hydrolysis
in adult fly glia (**A-C**) Representative images (A) and quantifications (B and C) on
in-vivo time lapse analysis of mitochondria-lysosome contacts (MLCs) in adult fly
glia expressing two reporters: *UAS-mCherry.mitoOMM* (labels the mitochondrial
outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green). Note that
VAMP7-deficient contacts exhibit a higher frequency and prolonged duration (>180s),
whereas the control contacts start untethering at t=180s (white arrow in top panel, A).
Note that only the glial expression of the GDP-bound Rab7 (*3xFlag-Rab7^{T22N}*), but
not Rab7 (*3xFlag-Rab7^{wt}*) nor GTP-bound Rab7 (*3xFlag-Rab7^{Q67L}*), restores the
contact dynamics (white arrows in bottom two panels in A, B and C). An overexposed
panel is shown for the rescue *3xFlag-Rab7^{T22N}* genotype for clarity (bottom panel, A).
(**D-G**) Representative images (D,D',F,F') and quantifications (E and G) of
mitochondrial morphology labeled by *UAS-mito.GFP* (*repo>mito.GFP*) in control,
3xFlag-Rab7^{wt}-,   *3xFlag-Rab7^{Q67L}*-,   *3xFlag-Rab7^{T22N}*-,   *or*
vamp7-RNAi^{43543}-expressing glia. Note that the average mitochondrial area quantified
by ImageJ is significantly larger upon Rab7^{Q67L} expression in glia. The mitochondrial
area is also restored when co-expressing Rab7^{T22N} and *vamp7*-RNAi^{43543}. The selected
white squares in D and F are enlarged and aligned in D' and F'. All samples examined

are brains of 3-day-old adult flies unless noted otherwise. For single transgene
experiment, *luc*-RNAi was used as a control. For double transgene experiment,
UAS-LacZ; *luc*-RNAi was used as a control. White arrows indicate the untethering.
Scale bars of different sizes are indicated on the images. Serial confocal Z-stack
sections were taken at similar planes across all genotypes, showing representative
single layer or merged (MLCs) images. Statistical graphs are shown with scatter dots
and the number of brains (one MLC per brain) listed on top. Images for the time lapse
series are shown in 3x3 μm squares centering on the target MLC selected from the
movies (Movies S8-S12). Parameters for the contact dynamics in B-D are quantified
using Imaris and ImageJ. Data are shown as mean ± SEM. P-values of significance
(indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and
**** $p<0.0001$) are calculated by one-way ANOVA followed by Tukey's multiple
comparisons test.
847

Figure 5. VAMP7 interacts with TBC1D15-17 and expression of TBC1D15-17
lacking the GAP activity causes increased GTP-bound Rab7 levels, impaired
MLCs, and mitochondrial elongation in adult fly glia (**A** and **B**) Representative
WB images for Co-IP interaction between VAMP7 and Rab7 (A) or TBC1D15-17 (B).
Note that VAMP7 interacts with TBC1D15-17, not Rab7. (**C** and **D**) Representative
WB images (B) and quantifications (C) of TBC1D15-17-Rab7 interaction with or
without glial VAMP7. Cell lysates from the adult fly heads expressing
3xHA-TBC1D15-17 and *luc*-RNAi or *vamp7*-RNAi[43543] were collected and
immunoprecipitated. Note that TBC1D15-17 remains associated with Rab7 upon glial
VAMP7 reduction. (**E**) An illustration of the TBC1D15-17 protein structure. The
disordered domain (259-325aa) and GAP domain (372-582aa) are labeled. (**F-H**)
Representative images (F) and quantifications (G and H) of transgenic flies expressing
PLEKHM1(CG6613)-GFP and truncated TBC1D15-17 variants. Note that GFP
signals exhibit aggregated patterns in forms of multiple puncta and elevated intensities
when expressing *vamp7*-RNAi[43543], *3xFlag-TBC1D15-17^{wt}*, or
3xFlag-TBC1D15-17^{ΔGAP}. (**I-M**) Representative images (I and L) and quantifications
(J,K,M) on *in-vivo* time lapse analysis of mitochondria-lysosome contacts (MLCs, I)
and mitochondrial morphology (L) in adult fly glia. Two reporters:
UAS-mCherry.mitoOMM (labels the mitochondrial outer membrane, red) and
UAS-Lamp1.GFP (labels lysosome, green) were expressed in glia to label MLCs,
whereas mitochondrial morphology was labeled by *UAS-mito.GFP* (green). Note that
TBC1D15-17^{ΔGAP} expression in glia causes prolonged MLCs and elongated
mitochondria. The selected white squares in L are enlarged and aligned in L'. All
samples examined are brains of 3-day-old adult flies unless noted otherwise. For
single transgene experiment, *luc*-RNAi was used as a control. For double transgene
experiment, *UAS-LacZ*; *luc*-RNAi was used as a control. White arrows indicate the
untethering (I). Scale bars of different sizes are indicated on the images. Serial
confocal Z-stack sections were taken at similar planes across all genotypes, showing
representative single layer or merged (MLCs) images. Statistical graphs are shown
with scatter dots and the number of brain (one MLC per brain) listed on top. Images

for the time lapse series are shown in 3x3 μm squares centering on the target MLC
selected from the movies (Movies S13-S16). Parameters for the contact dynamics in J
and K are quantified using Imaris and ImageJ. For Co-IP analysis, 200 independent
fly brains for each genotype were analyzed. Data are shown as mean ± SEM. P-values
of significance (indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, ***
$p<0.001$ and **** $p<0.0001$) are calculated by one-way ANOVA followed by Tukey's
multiple comparisons test.
885
Figure 6. VAMP7 regulates mitochondria-lysosome contacts via Rab7 GAP
TBC1D15-17 in adult fly glia (**A-E**) Representative images (A,D) and
quantifications (B,C,E) on *in-vivo* time lapse analysis of mitochondria-lysosome
contacts (MLCs, A) and mitochondrial morphology (D) in adult fly glia. Two
reporters: *UAS-mCherry.mitoOMM* (labels the mitochondrial outer membrane, red)
and *UAS-Lamp1.GFP* (labels lysosome, green) were expressed in glia to label MLCs,
whereas mitochondrial morphology was labeled by *UAS-mito.GFP* (green). Whereas
the contacts persist up to t=130s upon glial VAMP7 reduction, co-expression of
TBC1D15-17^{WT} or TBC1D15-17$^{\Delta disordered}$ restores the prolonged contact dynamics
(t=90s and 50s, middle two panels in A). Note that TBC1D15-17$^{\Delta GAP}$ expression fails
to restore the frequency and the duration of the contacts (B and C). The average
mitochondrial area quantified by ImageJ is significantly restored when co-expressing
vamp7-RNAi^{43543} and TBC1D15-17^{WT} or TBC1D15-17$^{\Delta disordered}$, but not
TBC1D15-17$^{\Delta GAP}$ (D and E). The selected white squares in D are enlarged and
aligned in D'. All samples examined are brains of 3-day-old adult flies unless noted
otherwise. For single transgene experiment, *luc*-RNAi was used as a control. For
double transgene experiment, *UAS-LacZ*; *luc*-RNAi was used as a control. White
arrows indicate the untethering (A). Scale bars of different sizes are indicated on the
images. Serial confocal Z-stack sections were taken at similar planes across all
genotypes, showing representative single layer (mitochondrial morphology) or
merged (MLCs) images. Statistical graphs are shown with scatter dots and the number
of brains (one MLC per brain) listed on top. Images for the time lapse series are
shown in 1.82x1.82 μm squares centering on the target MLC selected from the movies
(Movies S17-S21). Parameters for the contact dynamics in B and C are quantified
using Imaris and ImageJ. Data are shown as mean ± SEM. P-values of significance
(indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and
**** $p<0.0001$) are calculated by one-way ANOVA followed by Tukey's multiple
comparisons test.
914
Figure 7. Silencing *vamp7* expression increases ROS levels and disrupts
mitochondrial function in adult fly glia (**A**) DHE fluorescent intensities indicating
cellular ROS levels are significantly increased upon glial VAMP7 or SOD1 reduction.
(**B**) RT-qPCR analyses reveal increased transcription levels of genes related to
mitochondrial function including the ones encoding acyl-coA dehydrogenases
(*CG17544* and *CG9527*) and fatty acid synthase (*FASN*). (**C-H**) The intensity and
number of LDs labeled by the BODIPY dye in both larval and adult fly brains (C-E)

and the TAG content in the adult fly brains (F) are significantly increased upon glial
VAMP7 reduction. Note that the intensities of LDs are quantified in selected regions
of 3^{rd} instar larval brains (D) vs. the whole brains in 3-day-old adult flies (E). Glial
VAMP7 regulates LD formation as the addition of α-lipoic acid suppresses ROS
levels in both control and *repo>vamp7*-RNAi[43543] adult fly brains (G and H). (**I-P**)
Representative images (I and M) and quantifications (K and O) of the mitochondrial
H_2O_2 reduction-oxidation activity monitored by the reporter *UAS-mito-roGFP-Orp1*
in 10-day-old *repo>mito-roGFP-Orp1* adult fly brains. Note that the mitochondrial
H_2O_2 reduction-oxidation activity is upregulated when expressing *vamp7*-RNAi[43543]
or Rab7[Q67L] (I and K). The increased activity upon glial VAMP7 reduction is
suppressed by co-expressing Rab7[T22N] or TBC1D15-17[WT] (M and O). Representative
images (J,J',N,N') and quantifications (L and P) of the MitoTracker Red CMXRos
fluorescent intensities detecting cellular mitochondrial membrane potential per
mito.GFP-positive mitochondria. Note that the mitochondrial membrane potential is
hyperpolarized when expressing *vamp7*-RNAi[43543] or Rab7[Q67L] (J,J',L). The
hyperpolarization in VAMP7-deficient glia is suppressed by co-expressing Rab7[T22N]
or TBC1D15-17[WT] (N,N',P). The selected white squares in J and N are enlarged and
aligned in J' and N'. All samples examined are brains of 3-day-old adult flies unless
noted otherwise. For single transgene experiment, *UAS-LacZ* (*LacZ*) or *luc*-RNAi was
used as a control. For double transgene experiment, *UAS-LacZ*; *luc*-RNAi was used as
a control. Scale bars of different sizes are indicated on the images. Serial confocal
Z-stack sections were taken at similar planes across all genotypes, showing
representative single layer or merged (LD) images. Statistical graphs are shown with
scatter dots and the number of brains listed on top. For TAG statistics, 20 independent
fly brains for each genotype were analyzed. Data are shown as mean ± SEM. P-values
of significance (indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, and
*** $p<0.001$) are calculated by two-tailed unpaired t-test compared two groups
experiments or by one-way ANOVA compared three groups followed by Tukey's
multiple comparisons test.
951
Figure 8. Glial VAMP7 contributes to DA neuron survival (**A**) An schematic
diagram illustrating the DA neuron clusters in adult fly brains. The black dashed box
encloses the area containing DA neuron clusters (PPM1/2, PPL1, and PPM3) and is
visualized in B,E,G,I,K,M. (**B-D**) The numbers of DA neurons in PPM1/2, PPM3, and
PPL1 are significantly decreased (white arrows in B) in the aged flies (10-day-old)
upon glial VAMP7 reduction. (**E-N**) Representative images (E,G,I,K,M) and
quantifications (F,H,J,L,N) of DA neuron number in the PPM1/2 cluster under
different genetic conditions. Note that DA neuron number is consistently decreased in
the 10-day-old flies expressing *vamp7*-RNAi[43543] in the presence of Elav-Gal80,
which blocks Gal4 expression in neurons (E and F). DA neuron number is also
significantly decreased in the 10-day-old flies expressing Rab7[Q67L] (G and H). The
DA neurodegeneration upon glial VAMP7 reduction is restored by co-expressing
Rab7[T22N] or TBC1D15-17[WT] (I and J). Expressing SOD1 in glia (K and L) or neurons
(M and N) restores the DA neuron loss in PPM1/2 upon glial VAMP7 reduction. DA

neurons are detected with anti-TH antibodies. All samples examined are brains of 10-day-old adult flies unless noted otherwise. For single transgene experiment, *luc*-RNAi was used as a control. For double transgene experiment, *UAS-LacZ*; *luc*-RNAi was used as a control. Scale bars of different sizes are indicated on the images. Serial confocal Z-stack sections were taken at similar planes across all genotypes, showing representative merged images. Statistical graphs are shown with scatter dots and the number of brains listed on top. Data are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no significance, * p<0.05, ** p<0.01, *** p<0.001 and **** p<0.0001) are calculated by ordinary one-way ANOVA followed by Tukey's multiple comparisons test.

Figure 9. VAMP7 regulates mitochondria-lysosome contacts in glia An illustrated mechanism of VAMP7-mediated mitochondrial dynamics in glia. Under physiological conditions, VAMP7 interacts with TBC1D15-17, facilitating Rab7 GTP hydrolysis for the untethering of MLCs in adult fly glia. The contacts determine the site for fission. Upon glial VAMP7 reduction, TBC1D15-17 remains associated with Rab7, yet is unable to promote Rab7 GTP hydrolysis, leading to persistent contact and mitochondrial elongation.

Figure S1. *vamp7*-RNAi expression reduces *vamp7* mRNA levels and causes mitochondrial elongation in adult fly glia (**A** and **B**) RT-qPCR analysis indicates that glial expression of *vamp7*-RNAi reduces the *vamp7* transcription levels significantly in both adult fly brains (RNAi[38300] and RNAi[43543], A) and FACS-sorted glia (RNAi[43543], B). (**C** and **D**) Representative images (C,C') and quantifications (D) of mitochondrial morphology in adult fly glia. Note that the average mitochondrial volume quantified by Imaris in 3D is significantly larger than the control glia when expressing *vamp7*-RNAi[43543]. All samples examined are 3-day-old adult fly brains unless noted otherwise. For RT-qPCR statistics, 20 independent fly brains for each genotype were analyzed. For FACS, GFP-positive glia cells from 200 independent fly brains for each genotype were collected. For single transgene experiment, *luc*-RNAi was used as a control. Scale bars of different sizes are indicated on the images. Serial confocal Z-stack sections were taken at similar planes across all genotypes, showing representative single-layer images. Statistical graphs are shown with scatter dots and the number of brains listed on top. Data are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no significance, * p<0.05, ** p<0.01, *** p<0.001 and **** p<0.0001) are calculated by ordinary one-way ANOVA followed by Tukey's multiple comparisons test.

Figure S2. VAMP7 regulates mitochondrial dynamics in glia independently of autophagosome-lysosome fusion and mitophagy (**A** and **B**) Autophagic flux monitored by the dual reporter *UAS-GFP-mCherry-Atg8a* is impaired upon glial VAMP7 reduction. Note that the Pearson's coefficient value analyzed by Imaris reveals the increased colocalization of GFP- and mCherry-positive signals when silencing *vamp7* expression in glia. (**C** and **D**) Representative quantifications (C) and

images (D and D') of mitochondrial morphology in glia with or without Bafilomycin A1 (BafA1). Note that no change in the mitochondrial area is observed when autophagosome-lysosome fusion is blocked by BafA1. (**E** and **F**) Glial STX17 inhibition fails to induce mitochondrial elongation. Note that the mitochondrial area remains unaffected when expressing *stx17*-RNAi in glia. (**G** and **H**) Representative images (H) and quantifications (G) of the DA neurons in 10-day-old flies expressing *luc*-RNAi, *vamp7*-RNAi[43543], *stx17*-RNAi, or *snap29*-RNAi. Note that only vamp7-RNAi[43543] expression causes DA neuron loss. (**I** and **J**) Representative quantifications (I) and images (J,J') of mitochondrial morphology in glia when downregulating the expression of autophagy-related genes. Note that glial mitochondrial area remains largely intact when expressing *atg5*-RNAi or *ref(2)p*-RNAi, and smaller when expressing *atg1*-RNAi and *atg8*-RNAi. (**K-O**) Representative images (K) and quantifications (L-O) on mitochondria-lysosome contacts in fly glia expressing two reporters: *UAS-mCherry.mitoOMM* (labels the mitochondrial outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green). Contacts were co-labeled with the anti-Atg8a antibodies (Atg8a, magenta). Note a significant increase in the colocalization of mCherry- and GFP-positive signals upon glial VAMP7 reduction. Atg8a is minimally present in these dual-labeled contact sites, suggesting the absence of mitophagy. The selected areas in white squares in D,E,I are enlarged and aligned in D',E',I'. DA neurons are detected with anti-TH antibodies. All samples examined are brains of 3- or 10-day-old adult flies unless noted otherwise. For single transgene experiment, *luc*-RNAi was used as a control. Scale bars of different sizes are indicated on the images. A serial confocal Z-stack sections were taken at the similar plane across all genotypes, with representative merged images shown in A and H and single layer images shown in other panels. Statistical graphs are shown with scatter dots and the number of brains listed on top. Parameters for the colocalization and the mitochondrial area are quantified using Imaris and ImageJ as detailed in the Materials and Methods. Data are shown as mean ± SEM. P-values of significance (indicated with asterisks, ns no significance, * $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) are calculated by two-tailed unpaired t-test between two sample groups and by one-way ANOVA compared among three or more sample groups followed by Tukey's multiple comparisons test.

Figure S3. Astrocytic VAMP7 reduction causes DA neuron loss (A-H) Representative images (A, C, E, G) and quantifications (B, D, F, H) of the DA neurons in 10-day-old flies expressing *luc*-RNAi or *vamp7*-RNAi[43543] using the different glial subtype drivers. Note that *vamp7*-RNAi[43543] expression by the astrocyte driver GMR86E01-GAL4 causes decreased DA neuron number in the PPM1/2 cluster. DA neurons are detected with anti-TH antibodies. All samples examined are brains of 10-day-old adult flies unless noted otherwise. For single transgene experiment, *luc*-RNAi was used as a control. Scale bars of different sizes are indicated on the images. Serial confocal Z-stack sections were taken at similar planes across all genotypes, showing representative merged images. Statistical graphs are shown with scatter dots and the number of brains listed on top. Data are shown as mean ± SEM.

1054 P-values of significance (indicated with asterisks, ns no significance, * p<0.05, **
1055 p<0.01, *** p<0.001 and **** p<0.0001) are calculated by ordinary one-way
1056 ANOVA followed by Tukey's multiple comparisons test.
1057

Movie S1. *In-vivo* time lapse analysis of mitochondria-lysosome contact in the
control flies. Two reporters *UAS-mCherry.mitoOMM* (labels the mitochondrial outer
membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green) are co-expressed to
visualize the mitochondria-lysosome contacts. Note that mitochondria and lysosomes
contact and dissociate over the imaging period. 5 images Z-stack sections were taken
with 0.125 μm each at an interval of 10 seconds for 5 minutes. Merged images are
shown. Video corresponds to Figure 2A. Scale bar, 1 μm.
1065

Movie S2-4. *In-vivo* time lapse analysis of mitochondria-lysosome contact in flies
lacking glial VAMP7. Two reporters *UAS-mCherry.mitoOMM* (labels the
mitochondrial outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green)
are co-expressed to visualize mitochondria-lysosome contact. A representative contact
is shown (Movie S2, *repo>Lamp1GFP,mCherry.mitoOMM; Luc*-RNAi). Note the
significant increase in colocalization of mitoOMM and Lamp1 upon glial VAMP7
reduction (Movie S3, *repo>Lamp1GFP,mCherry.mitoOMM; vamp7*-RNAi[38300] and
Movie S4, *repo>Lamp1GFP,mCherry.mitoOMM; vamp7*-RNAi[43543]). 5 images
Z-stack sections were taken with 0.125 μm each at an interval of 10 seconds for 5
minutes. Merged images are shown. Videos correspond to Figure 2B. Scale bar, 1 μm.
1076

Movie S5-7. Glial VAMP7 regulates mitochondria lysosome contacts. Two
reporters *UAS-mitoOMM.mCherry* (labels the mitochondrial outer membrane, red)
and *UAS-Lamp1.GFP* (labels lysosome, green) are co-expressed to visualize
mitochondria-lysosome contact *in vivo*. A representative contact is shown (Movie S5,
repo>Lamp1GFP, mCherry.mitoOMM, LacZ; Luc-RNAi). Note the significant
increase in colocalization of mitoOMM and Lamp1 upon glial VAMP7 depletion
(Movie S6, *repo>Lamp1GFP, mCherry.mitoOMM, LacZ; vamp7*-RNAi[43543]).
Re-introduced 6xMyc-VAMP7 after decreased glial VAMP7 expression,
mitochondria-lysosome contacts were rescued (Movie S7, *repo>Lamp1GFP,*
mCherry.mitoOMM, 6xMyc-vamp7; vamp7-RNAi[43543]). 10 images Z-stack sections
were taken with 0.125 μm each at an interval of 28 seconds for 5 minutes. Merged
images are shown. Videos correspond to Figure 2E. Scale bar, 1 μm.
1089

Movie S8-12. Glial VAMP7 regulates mitochondria-lysosome contacts via Rab7
hydrolysis Two reporters *UAS-mCherry.mitoOMM* (labels the mitochondrial outer
membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green) are co-expressed to
visualize mitochondria-lysosome contact. A representative contact is shown (Movie
S8, *repo>Lamp1GFP, mCherry.mitoOMM, LacZ; Luc*-RNAi). Note the significant
increase in colocalization of mitoOMM and Lamp1 upon glial VAMP7 depletion
(Movie S9, *repo>Lamp1GFP, mCherry.mitoOMM, LacZ; vamp7*-RNAi[43543]).
Co-expressing wild type Rab7 (*3xflag-rab7[wt]*) or GTP-bound Rab7 (*3xflag-rab7[Q67L]*)

and *vamp7*-RNAi^{435435} formed a stable mitochondria-lysosome contact (Movie S10,

repo>Lamp1GFP, mCherry.mitoOMM, 3xflag-rab7^{wt}; vamp7-RNAi^{43543} and Movie

S11, *repo>Lamp1GFP, mCherry.mitoOMM, 3xflag-rab7^{Q67L}; vamp7*-RNAi^{43543}). On

the contrary, co-expressing GDP-bound Rab7 (*3xflag-rab7^{T22N}*) and

vamp7-RNAi^{435435} recovered the mitochondria-lysosome untethering (Movie S12,

repo>Lamp1GFP, mCherry.mitoOMM, 3xflag-rab7^{T22N}; vamp7-RNAi^{43543}). 5 images

Z-stack sections were taken with 0.125 μm each at an interval of 10 seconds for 5

minutes. Merged images are shown. Videos correspond to Figure 4A. Scale bar, 1 μm.

1106

Movie S13-16. Glial expression of TBC1D15-17 lacking the GAP activity disrupts

mitochondria-lysosome contacts. Two reporters *UAS-mCherry.mitoOMM* (labels the

mitochondrial outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green)

are co-expressed to visualize mitochondria-lysosome contact. A representative contact

is shown (Movie S13, *repo>Lamp1GFP, mCherry.mitoOMM, 3xflag*). Note the

significant increase in colocalization of mitoOMM and Lamp1 upon glial

TBC1D15-17 lacking the GAP domain (Movie S16, *repo>Lamp1GFP,*

mCherry.mitoOMM, 3xFlag-TBC1D15-17$^{\Delta GAP}$). Whereas expressing wild type Rab7

GAP TBC1D15-17^{WT} (Movie S14, *repo>Lamp1GFP, mCherry.mitoOMM,*

3xFlag-TBC1D15-17^{WT}) or TBC1D15-17 lacking the disordered domain (Movie S15,

repo>Lamp1GFP, mCherry.mitoOMM, 3xFlag-TBC1D15-17$^{\Delta disordered}$) in glia causes

mitochondria and lysosome untethering successfully after contacts. 5 images Z-stack

sections were taken with 0.395 μm each at an interval of 10 seconds for 5 minutes.

Merged images are shown. Videos correspond to Figure 5I. Scale bar, 1 μm.

1121

Movie S17-21. VAMP7 regulates mitochondria-lysosome contacts in glia via the

Rab7 GAP TBC1D15-17. Two reporters *UAS-mCherry.mitoOMM* (labels the

mitochondrial outer membrane, red) and *UAS-Lamp1.GFP* (labels lysosome, green)

are co-expressed to visualize mitochondria-lysosome contact. A representative contact

is shown (Movie S17, *repo>Lamp1GFP, mCherry.mitoOMM, LacZ; Luc*-RNAi).

Note the significant increase in colocalization of mitoOMM and Lamp1 upon glial

VAMP7 depletion (Movie S18, *repo>Lamp1GFP, mCherry.mitoOMM, LacZ;*

vamp7-RNAi^{43543}). Co-expressing *vamp7*-RNAi^{43543} and wild type Rab7 GAP

3xflag-TBC1D15-17^{WT} (Movie S19, *repo>Lamp1GFP, mCherry.mitoOMM,*

3xFlag-TBC1D15-17^{WT}; vamp7-RNAi^{43543}) or TBC1D15-17 lacking the disordered

domain (Movie S20, *repo>Lamp1GFP, mCherry.mitoOMM,*

3xFlag-TBC1D15-17$^{\Delta disordered}$; vamp7-RNAi^{43543}) causes mitochondria and lysosome

untethering successfully after contacts. Whereas co-expressing glial *vamp7*-RNAi and

TBC1D15-17 lacking the GAP domain (Movie S16, *repo>Lamp1GFP,*

mCherry.mitoOMM, 3xFlag-TBC1D15-17$^{\Delta GAP}$; vamp7-RNAi) inhibits untethering

after mito.OMM and Lamp1.GFP contacts. 5 images Z-stack sections were taken with

0.395 μm each at an interval of 10 seconds for 5 minutes. Merged images are shown.

Videos correspond to Figure 6A. Scale bar, 1 μm.

1140

1141 **References**

1. Fenton AR, Jongens TA, Holzbaur ELF. Mitochondrial dynamics: Shaping and remodeling an organelle network. *Curr Opin Cell Biol* **68**, 28-36 (2020).
2. Gao S, Hu J. Mitochondrial Fusion: The Machineries In and Out. *Trends Cell Biol* **31**, 62-74 (2021).
3. Mattie S, Krols M, McBride HM. The enigma of an interconnected mitochondrial reticulum: new insights into mitochondrial fusion. *Curr Opin Cell Biol* **59**, 159-166 (2019).
4. Cho DH, Nakamura T, Lipton SA. Mitochondrial dynamics in cell death and neurodegeneration. *Cell Mol Life Sci* **67**, 3435-3447 (2010).
5. Itoh K, Nakamura K, Iijima M, Sesaki H. Mitochondrial dynamics in neurodegeneration. *Trends Cell Biol* **23**, 64-71 (2013).
6. Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. ER tubules mark sites of mitochondrial division. *Science* **334**, 358-362 (2011).
7. Kalia R*, et al.* Structural basis of mitochondrial receptor binding and constriction by DRP1. *Nature* **558**, 401-405 (2018).
8. Kamerkar SC, Kraus F, Sharpe AJ, Pucadyil TJ, Ryan MT. Dynamin-related protein 1 has membrane constricting and severing abilities sufficient for mitochondrial and peroxisomal fission. *Nat Commun* **9**, 5239 (2018).
9. Palmer CS, Elgass KD, Parton RG, Osellame LD, Stojanovski D, Ryan MT. Adaptor proteins MiD49 and MiD51 can act independently of Mff and Fis1 in Drp1 recruitment and are specific for mitochondrial fission. *J Biol Chem* **288**, 27584-27593 (2013).
10. Kageyama Y*, et al.* Mitochondrial division ensures the survival of postmitotic neurons by suppressing oxidative damage. *J Cell Biol* **197**, 535-551 (2012).
11. Imai Y. PINK1-Parkin signaling in Parkinson's disease: Lessons from Drosophila. *Neurosci Res* **159**, 40-46 (2020).
12. Pereira SL, Grossmann D, Delcambre S, Hermann A, Grünewald A. Novel insights into Parkin-mediated mitochondrial dysfunction and neuroinflammation in Parkinson's disease. *Curr Opin Neurobiol* **80**, 102720 (2023).
13. Uoselis L, Nguyen TN, Lazarou M. Mitochondrial degradation: Mitophagy and beyond. *Mol Cell* **83**, 3404-3420 (2023).
14. Wong YC, Peng W, Krainc D. Lysosomal Regulation of Inter-mitochondrial Contact Fate and Motility in Charcot-Marie-Tooth Type 2. *Dev Cell* **50**, 339-354 e334 (2019).
15. Wong YC, Ysselstein D, Krainc D. Mitochondria-lysosome contacts regulate mitochondrial fission via RAB7 GTP hydrolysis. *Nature* **554**, 382-386 (2018).
16. Stroupe C. This Is the End: Regulation of Rab7 Nucleotide Binding in Endolysosomal Trafficking and Autophagy. *Front Cell Dev Biol* **6**, 129 (2018).
17. Zhang Y*, et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929-11947 (2014).
18. Zhang Y*, et al.* Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37-53 (2016).
19. Hong W. SNAREs and traffic. *Biochim Biophys Acta* **1744**, 120-144 (2005).
20. Tian X, Teng J, Chen J. New insights regarding SNARE proteins in autophagosome-lysosome fusion. *Autophagy*, 1-9 (2020).
21. Knott AB, Bossy-Wetzel E. Impairing the mitochondrial fission and fusion balance: a new mechanism of neurodegeneration. *Ann N Y Acad Sci* **1147**, 283-292 (2008).
22. Ul Fatima N, Ananthanarayanan V. Mitochondrial movers and shapers: Recent insights into regulators of fission, fusion and transport. *Curr Opin Cell Biol* **80**, 102150 (2023).
23. Deus CM, Yambire KF, Oliveira PJ, Raimundo N. Mitochondria-Lysosome Crosstalk: From Physiology to Neurodegeneration. *Trends Mol Med* **26**, 71-88 (2020).
24. Kim S, Wong YC, Gao FD, Krainc D. Dysregulation of

mitochondria-lysosome contacts by dysfunction in dopaminergic neuronal models of Parkinson's disease. *Nature Communications* **12**, (2021).

25. Tabata K, Matsunaga K, Sakane A, Sasaki T, Noda T, Yoshimori T. Rubicon and PLEKHM1 negatively regulate the endocytic/autophagic pathway via a novel Rab7-binding domain. *Mol Biol Cell* **21**, 4162-4172 (2010).

26. Gillingham AK, Sinka R, Torres IL, Lilley KS, Munro S. Toward a comprehensive map of the effectors of rab GTPases. *Dev Cell* **31**, 358-373 (2014).

27. Lee MJ, Jang S, Nahm M, Yoon JH, Lee S. Tbc1d15-17 regulates synaptic development at the Drosophila neuromuscular junction. *Mol Cells* **36**, 163-168 (2013).

28. Liu Y, Bao H, Wang W, Lim HY. Cardiac Snail family of transcription factors directs systemic lipid metabolism in Drosophila. *PLoS Genet* **15**, e1008487 (2019).

29. Senos Demarco R, Uyemura BS, D'Alterio C, Jones DL. Mitochondrial fusion regulates lipid homeostasis and stem cell maintenance in the Drosophila testis. *Nat Cell Biol* **21**, 710-720 (2019).

30. Albrecht SC, Barata AG, Grosshans J, Teleman AA, Dick TP. In Vivo Mapping of Hydrogen Peroxide and Oxidized Glutathione Reveals Chemical and Regional Specificity of Redox Homeostasis. *Cell Metab* **14**, 819-829 (2011).

31. Budnik V, White K. Catecholamine-containing neurons in Drosophila melanogaster: distribution and development. *J Comp Neurol* **268**, 400-413 (1988).

32. Nassel DR, Elekes K. Aminergic neurons in the brain of blowflies and Drosophila: dopamine- and tyrosine hydroxylase-immunoreactive neurons and their relationship with putative histaminergic neurons. *Cell Tissue Res* **267**, 147-167 (1992).

33. Feany MB, Bender WW. A Drosophila model of Parkinson's disease. *Nature* **404**, 394-398 (2000).

34. Mauvezin C, Nagy P, Juhasz G, Neufeld TP. Autophagosome-lysosome fusion is independent of V-ATPase-mediated acidification. *Nat Commun* **6**, 7007 (2015).

35. Yoshimori T, Yamamoto A, Moriyama Y, Futai M, Tashiro Y. Bafilomycin A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification and protein degradation in lysosomes of cultured cells. *J Biol Chem* **266**, 17707-17712 (1991).

36. Hao T, *et al.* Hypoxia-reprogramed megamitochondrion contacts and engulfs lysosome to mediate mitochondrial self-digestion. *Nat Commun* **14**, 4105 (2023).

37. Gomes LC, Di Benedetto G, Scorrano L. During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. *Nature Cell Biology* **13**, 589-U207 (2011).

38. Purnell PR, Fox HS. Autophagy-mediated turnover of Dynamin-related Protein 1. *Bmc Neurosci* **14**, (2013).

39. Arasaki K, *et al.* A role for the ancient SNARE syntaxin 17 in regulating mitochondrial division. *Dev Cell* **32**, 304-317 (2015).

40. Sugo M, *et al.* Syntaxin 17 regulates the localization and function of PGAM5 in mitochondrial division and mitophagy. *Embo Journal* **37**, (2018).

41. McLelland GL, Lee SA, McBride HM, Fon EA. Syntaxin-17 delivers PINK1/parkin-dependent mitochondrial vesicles to the endolysosomal system. *J Cell Biol* **214**, 275-291 (2016).

42. Verkhratsky A, Matteoli M, Parpura V, Mothet JP, Zorec R. Astrocytes as secretory cells of the central nervous system: idiosyncrasies of vesicular secretion. *EMBO J* **35**, 239-257 (2016).

43. Pulipparacharuvil S, *et al.* Drosophila Vps16A is required for trafficking to lysosomes and biogenesis of pigment granules. *J Cell Sci* **118**, 3663-3673 (2005).

1258   44.   Li H, Li Y, Lei Z, Wang K, Guo A. Transformation of odor selectivity from
1259   projection neurons to single mushroom body neurons mapped with dual-color
1260   calcium imaging. *Proc Natl Acad Sci U S A* **110**, 12084-12089 (2013).
1261
1262

Fig 1

Fig 2

Fig 3

A *repo>3xHA-RILPL* Day 3

IP:HA
- HA-RILPL
- Rab7

Cell lysates
- HA-RILPL
- Rab7
- Flag-Rab7^{Q67L}
- α-Tubulin

B *repo>3xHA-RILPL* Day 3

Ratio of Rab7^{IP} (GTP-bound)/Total Rab7

C *repo>PLEKHM1-GFP* Day 10

luc-RNAi | *vamp7*-RNAi^{43543} | *3xFlag-rab7*^{Q67L}

PLEKHM1-GFP

D *repo>PLEKHM1-GFP* Day 10

Mean PLEKHM1-GFP intensity (A.U.)

E *repo>PLEKHM1-GFP* Day 10

Mean PLEKHM1-GFP іarea (µm²)

F *repo>PLEKHM1-GFP* Day 10

Mean PLEKHM1-GFP intensity (A.U.)

G *repo>PLEKHM1-GFP* Day 10

Mean PLEKHM1-GFP іarea (µm²)

H *repo>PLEKHM1-GFP* Day 10

LacZ; luc-RNAi | *LacZ; vamp7*-RNAi^{43543} | *3xFlag-rab7*^{wt}*; vamp7*-RNAi^{43543} | *3xFlag-rab7*^{Q67L}*; vamp7*-RNAi^{43543} | *3xFlag-rab7*^{T22N}*; vamp7*-RNAi^{43543}

PLEKHM1-GFP

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8

With VAMP7

Loss of VAMP7