Batch experiments demonstrating a two-stage bacterial process coupling

methanotrophic and heterotrophic bacteria for 1-alkene production from methane

Ramita Khanongnuch[a,*], Rahul Mangayil[a], Ville Santala[a], Anne Grethe Hestnes[b], Mette

Marianne Svenning[b], Antti J Rissanen[a]

[a]Faculty of Engineering and Natural Sciences, Tampere University, P.O. Box 541, 33014
Tampere, Finland
[b]Department of Arctic and Marine Biology, UiT, The Arctic University of Norway, 9037
Tromsø, Norway

*Corresponding author:
Ramita Khanongnuch, Faculty of Engineering and Natural Sciences, Tampere University,
P.O. Box 541, 33014 Tampere, Finland; email: ramita.khanongnuch@tuni.fi; Phone: +358
442415042

Abstract

Methane (CH$_4$) is a sustainable carbon feedstock source for aerobic CH$_4$-oxidizing

bacteria (methanotrophs) to produce value-added chemicals. Under substrate-limited (e.g.,

CH$_4$, oxygen and nitrogen) conditions, CH$_4$ oxidation results in the production of various

short-chain organic acids and platform chemicals. These CH$_4$-derived products could be

broadened by utilizing them as a feedstock for heterotrophic bacteria. As a proof of concept,

a two-stage system for CH$_4$ abatement and 1-alkene production was developed in this study.

Types I and II methanotrophs, i.e., *Methylobacter tundripaludum* SV96 and *Methylocystis*

rosea SV97, respectively, were investigated in batch tests under different CH$_4$ and air

supplementation schemes. CH$_4$ oxidation under either microaerobic or aerobic conditions

induced the production of formate, acetate, succinate, and malate in *M. tundripaludum* SV96,

accounting for 4.8-7.0% of consumed CH$_4$-carbon while *M. rosea* SV97 produced the same

compounds except for malate, and with lower efficiency than *M. tundripaludum* SV96,

accounting for 0.7-1.8% of consumed CH$_4$-carbon For the first time, the organic acids-rich

spent media of methanotrophs were successfully used for 1-alkene production using

engineered *Acinetobacter baylyi* ADP1 '*tesA-undA* cells. The highest yield of 1-undecene

was obtained from spent medium of *M. tundripaludum* SV96 at 68.9 ± 11.6 µmol mol

C$_{substrate}^{-1}$.

Keywords: methane, methanotrophs, organic acids production, 1-alkene production,

Acinetobacter baylyi ADP1

1. Introduction

Methane (CH_4) is the second most important greenhouse gas (GHG) after CO_2, having a global warming potential (GWP) of approximately 30 times higher than CO_2 on a 100-year time horizon (IPCC, 2021). Its emission from anthropogenic activities has been continuously increasing, being approximately as high as 60% of total CH_4 emission to the atmosphere (based on top-down estimates of Saunois et al., 2020). Hence, keeping on a stringent climate policy for the next several decades, particularly the use of CH_4 as an energy source, would significantly reduce CH_4 emissions (Harmsen et al., 2020). In environmental carbon flux, CH_4 oxidizing bacteria are the key regulators for CH_4 abatement (Hanson and Hanson, 1996; Saarela et al., 2020). Due to its abundance and potentiality as a sustainable carbon feedstock, the development of the biological processes for CH_4 conversion to bio-based chemicals/liquid fuel is promising and attractive research area (Liu et al., 2020; López et al., 2013; Sun et al., 2018).

Aerobic methane oxidizing bacteria (methanotrophs) containing methane monooxygenases (MMO) use CH_4 as the sole carbon and energy source and oxygen as an electron acceptor. Methanotrophs are classified into gammaproteobacterial (Types I and X) and alphaproteobacterial (Type II) methanotrophs based on their different pathways for formaldehyde assimilation into biomass, i.e., the ribulose monophosphate (RuMP) and the serine cycles, respectively (Kalyuzhnaya et al., 2015). Along with these pathways, CH_4 is potentially converted into various value-added products including methanol, single cell protein, ectoine and soluble metabolites (Cantera et al., 2018; Ge et al., 2014; Sheets et al., 2016; Strong et al., 2016). In various environmental processes and biological systems, methanotrophs have been reported to support other bacteria by producing organic carbon sources from CH_4, such as in wastewater treatment (Cao et al., 2019; Costa et al., 2000) and heavy metal bioremediation (Lai et al., 2016). The pathways for the excretion of organic

acids (e.g., formate, acetate, lactate and succinate) have been observed in the experiments and

deciphered from genomes of Type I methanotrophs during O_2 limiting conditions including

the Embden-Meyerhof-Parnas (EMP) and tricarboxylic acid (TCA) cycle (Fig. 1) (Gilman et

al., 2017; Kalyuzhnaya et al., 2013). Besides O_2 limiting conditions, acetate production was

also very recently detected in liquid culture of methanotrophs incubated under aerobic

conditions (Lee et al., 2021; Takeuchi and Yoshioka, 2021). These studies also showed the

potential of the methane-derived organic acids in biotechnological applications by cultivating

Type I methanotrophs and heterotrophs in co-culture systems, where the heterotrophs utilized

the organic acids produced by the methanotrophs (Lee et al., 2021; Takeuchi and Yoshioka,

2021). For example, acetate produced by *Methylocaldum marinum* S8 was successful used as

a growth medium for *Cupriavidus necator* cultivation (Takeuchi and Yoshioka, 2021) and

co-culture of *Methylococcus capsulatus* Bath and the engineered *Escherichia coli* SBA01

was carried out for mevalonate production (Lee et al., 2021).

Fig. 1.

Acinetobacter baylyi ADP1, naturally competent gram-negative gammaproteobacteria,

have been recently gained interest in bioengineering. The bacterium's capacity to metabolize

a wide range of carbon sources and genetic tractability has led to its application in long-chain

alkyl ester and 1-alkene compounds (e.g. 1-undecene) biosynthesis from aromatic lignin-

derived monomers and lignocellulose (Luo et al., 2019; Salmela et al., 2020). As a platform

chemical, 1-undecene can be used in the chemical synthesis of medium-chain poly-α-olefins

(C33) which are commonly used as lubricants (Salmela et al., 2020). Recently, the ability of

A. baylyi ADP1 to metabolize fermentation by-products has been exploited to broaden the

metabolic landscape in biological production processes (Mangayil et al., 2019; Salmela et al.,

2018). Hence, *A. baylyi* ADP1 would provide a promising, testable candidate for the

bioconversion of CH_4–derived organic acids. However, it is critical to find a suitable

methanotroph partner for such application. For instance, whether Type I or II methanotroph

would make a good candidate for converting CH_4 to organic acids for *A. baylyi* ADP1 is not

known, as to our knowledge, the comparison of organic acid production yields between

Types I and II methanotrophs has not been previously reported. Nevertheless, the CH_4-

derived compounds are limited to short carbon chain carbon compounds (C2-C6), which

could be further used as a feedstock for various heterotrophic bacteria, that can be easily

engineered in order to extend the range of CH_4-derived value-added (platform) chemicals.

With the aim to redirect the 'overflow carbon' from microaerobic methanotrophic

fermentation, organic acids obtained from CH_4 oxidation process could potentially provide as

important feedstocks towards high-value product formation. Thus, a two-stage bioprocess

setup was designed for successful integration of production of CH_4-derived organic acids

with aerobic synthesis. The growth and metabolite production profiles of Types I and II

methanotrophs, i.e., *Methylobacter tundripaludum* SV96 and *Methylocystis rosea* SV97,

respectively, were investigated under different gas supplementation schemes. Next, the

possibility of using the spent media of the methanotrophs as a growth substrate for the

cultivation of wild type *A. baylyi* ADP1 was tested. Finally, the synthesis of 1-alkenes from

CH_4 was demonstrated by a two-stage process with a methanotroph, i.e., *M. tundripaludum*

SV96 and *M. rosea* SV97, and an engineered *A. baylyi* ADP1 ('*tesA-undA*') strain (Luo et al.,

2019).

2. Materials and Methods

2.1. Strains and cultivation conditions

M. tundripaludum SV96 and *M. rosea* SV97, isolated from Arctic wetland soil, Norway

(Wartiainen et al., 2006a, 2006b), were used in this study. Nitrate mineral salts (NMS)

medium used for cultivating the methanotrophs was DSMZ 921 medium (German Collection

of Microorganisms and Cell Cultures, 2007) with initial pH ~6.80 and modification of adding

with 1 µM lanthanum chloride (LaCl$_3$). The pre-inoculum was grown in 120-mL serum

bottles containing 10 mL NMS medium and 20% CH$_4$ and 80% air in headspace and

incubated statically at 20 °C. All experiments were conducted under sterile conditions and the

serum bottles used for methanotrophs cultivation were sealed with butyl rubber stoppers and

capped with aluminum crimps.

For cocultivation studies, wild type and the engineered *A. baylyi* ADP1 strain (Luo et

al., 2019) carrying the plasmid pBAV1C-*'tesA-undA* (*A. baylyi* ADP1 *'tesA-undA*) were

used. For pre-inoculum, *A. baylyi* ADP1 cells were inoculated in 10 mL culture tubes

containing LB medium (5 g L^{-1} yeast extract, 10 g L^{-1} tryptone and 5 g L^{-1} NaCl)

supplemented with 0.5% glucose and 25 µg mL^{-1} chloramphenicol (for *A. baylyi* ADP1 *'tesA-*

undA). The inoculated tubes were aerobically grown overnight at 30 °C and 300 rpm. For 1-

undecene synthesis, *A. baylyi* ADP1 *'tesA-undA* was induced with 0.5 mM cyclohexanone

(Luo et al., 2019).

2.2. Evaluation of organic acid production by *M. tundripaludum* SV96 and *M. rosea* SV97

Batch tests were performed in 120-mL airtight serum bottles with a working volume of

15 mL NMS medium. The precultures of *M. tundripaludum* SV96 and *M. rosea* SV97 were

inoculated with an initial optical density at 600 nm (OD$_{600nm}$) of 0.02 and the growth, CH$_4$

utilization and organic acid profiles were monitored every two days for 14 days at 20°C and

static conditions. Both methanotrophs were tested under three different gas supplementation

schemes (Fig. 2). On day 0, the bottles were filled with 20% CH$_4$ and 80% air into headspace

accounting for the initial O$_2$/CH$_4$ molar ratio of ~1.2 and incubated for 7 days. After CH$_4$ and

O$_2$ concentrations were depleted on day 7, the batch bottles were supplemented with three

different gas compositions into headspace, i.e., test I: CH$_4$ + air (20% CH$_4$ and 80% air), test

II: only CH$_4$ (20% CH$_4$ and 80% N$_2$), and test III: only air (20% air and 80% N$_2$). The bottles

containing only NMS without bacterial cells were used as the control. The bottles were

incubated further for 7 days (days 8-14). The biomass growth (OD$_{600}$), gas compositions in

headspace and organic acid accumulation in liquid medium were monitored every two days.

Fig. 2.

2.3. Cultivation of wild type *A. baylyi* ADP1 in spent media of different methanotrophs

Prior to the study of 1-alkene production, the capacity of *A. baylyi* ADP1 cells to utilize

the spent media of methanotrophs were tested. In this test, *M. tundripaludum* SV96 and *M.*

rosea SV97 were cultivated in 500-mL airtight bottles with 60 mL NMS medium in

conditions as described previously (Section 2.2). To maximize the organic acid titer, the

bottle headspace was initially filled with 20% CH$_4$ and 80% air and cultivated for 28 days at

optimal conditions for *M. tundripaludum* SV96 and *M. rosea* SV97, individually. Gaseous

composition in headspace and organic acid concentrations in liquid medium were monitored

twice a week.

After 28 days, the cultivation of methanotrophs was stopped and their spent media were

used for *A. baylyi* ADP1 cultivation. The spent media were collected by centrifugation at

12000 rpm for 10 min. The 50-mL supernatant of both methanotrophs were transferred to 250

mL sterile flasks and used for *A. baylyi* ADP1 cultivation. The tests were conducted in

duplicate. After 4 h cultivation, the liquid culture was collected to monitor cell growth,

organic acid concentration and wax ester production. The spent media of methanotrophs

without *A. baylyi* ADP1 were used as contamination control (control 1) and *A. baylyi* ADP1

cells in NMS (fresh medium) were used to determine their background growth (control 2).

2.4. Cultivation of *A. baylyi* ADP1 *'tesA-undA* strain and 1-alkene synthesis in spent media of

different methanotrophs

In this test, the liquid culture samples from *M. tundripaludum* SV96 and *M. rosea*

SV97 microaerobic fermentation were collected and directly used to cultivate *A. baylyi* ADP1

'tesA-undA strain. The tests were conducted in quadruplicate in a sealed 20 mL glass tube

containing 5 mL of the spent medium. Subsequently, *A. baylyi* ADP1 '*tesA-undA* cells (initial

OD_{600nm}, 0.02) were added into the vials. After 1 h of incubation at 30 °C and 300 rpm, 0.5

mM cyclohexanone was added to the vials to induce 1-undecene synthesis, and the vials were

further incubated for 23 hours at 30 °C and 300 rpm. The liquid culture samples from *M.*

tundripaludum SV96 and *M. rosea* SV97 cultivation without *A. baylyi* ADP1 '*tesA-undA*

were included as the experimental control. The growth of *A. baylyi* ADP1 '*tesA-undA* was

estimated based on difference of OD_{600nm} between the tests and controls at 24-h incubation

(ΔOD_{600nm}).

2.5. Analytical methods

Liquid samples were filtered through a 0.2 µm membrane (Chromafil® Xtra PET-

20/25, Macherey-Nagel, Germany) prior to the analysis of liquid metabolites using a

Shimadzu high-performance liquid chromatograph (HPLC) equipped with Rezex RHM-

Monosaccharide H^+ column (Phenomenex, USA) as described in Okonkwo et al. (2018). The

gas samples in the headspace (CH_4, CO_2, O_2 and N_2) were measured using a Shimadzu gas

chromatograph GC-2014 equipped with a thermal conductivity detector (TCD) and a

Carboxen-1000 60/80 column (Agilent Technologies, USA). The oven temperature was held

at 35 °C for 3.75 min, increased 30 °C min^{-1} to 150 °C for 3 min. The injector and detector

were 155 °C and 160 °C, respectively. Helium was used as a carrier gas at 30 mL min^{-1}.

The production of 1-undecene in headspace was detected using gas chromatography-

mass spectrometry (GC-MS) as described by Luo et al. (2019). Compounds were identified

using the NIST/EPA/NIH Mass Spectral Library (NIST 05). The bacterial growth was

measured as OD_{600nm} using an Ultrospec 500 pro spectrophotometer (Amersham Biosciences,

UK) and as cell dry weight (CDW) using gravimetric method. The conversion factor between

CDW and OD_{600nm} obtained from the experimental measurement was 0.2914 g L^{-1} OD^{-1} (R^2 =

0.9948) and 0.3015 g L^{-1} OD^{-1} (R^2 = 0.9892) for *M. tundripaludum* SV96 and *M. rosea*

SV97, respectively.

Wax ester production was detected by using lipid extraction and thin layer

chromatography (TLC) analysis. Lipid extraction was done by using methanol-chloroform

extraction described by Santala et al. (2011) and the obtained lipid phase layer was used for

TLC analysis to visualize the total lipid composition. TLC analysis was done as described by

Santala et al. (2011). Briefly, 60 µl of lipid phase of a sample was applied to TLC plate (10 ×

20 cm HPTLC Silica Gel 60 F_{254} glass plates with 2.5 × 10 cm concentrating zone, Merck,

USA) using n-hexane:diethyl ether:acetic acid of 90:15:1 as a mobile phase. The

visualization was done by dyeing with iodine and Jojoba oil was used as a standard.

2.6. Statistical analysis

The statistical analysis was performed using Minitab16.0 (USA). The significant

differences of the obtained data sets (e.g., growth of the tested strains, gas concentrations and

utilization, concentrations and yields of the products produced by each strain) within the

varied treatment were analyzed using one-way of variance (ANOVA) with Tukey's multiple

comparison tests at the 95% confidence interval, where *P-value* ≤ 0.05 was considered

statistically significant.

3. Results and discussion

3.1. CH_4-derived organic acid production of *M. tundripaludum* SV96 and *M. rosea* SV97 in

different gas supplementation tests

Both *M. tundripaludum* SV96 and *M. rosea* SV97 used CH_4 as the sole carbon and

energy sources and O_2 as an electron acceptor. In all gas supplementation tests, *M.*

tundripaludum SV96 showed higher in biomass production than *M. rosea* SV97 (Fig. 3).

Highest biomass production of both methanotrophs was observed in the supplementation of

both CH_4+air (test I) on day 14 at concentrations of 0.60 ± 0.03 and 0.48 ± 0.05 g CDW L^{-1}

for *M. tundripaludum* SV96 and *M. rosea* SV97, respectively. In all tests, the growth yield of

M. rosea SV97 was lower than *M. tundripaludum* SV96 ($P < 0.05$) due to typical carbon

assimilation pathway of Type I methanotrophs (RuMP) showing more efficient channeling of

CH_4-carbon to biomass than in Type II methanotrophs (serine cycle) (Kalyuzhnaya et al.,

2015).

Fig. 3.

Spent media of both methanotrophs contained similar organic acid compounds

including formate, acetate, and succinate, while malate was only present in the spent medium

of *M. tundripaludum* SV96. However, the results suggested that *M. tundripaludum* SV96 was

more efficient and promising in CH_4 conversion into organic acids than *M. rosea* SV97. The

spent medium of *M. tundripaludum* SV96 contained higher concentrations and yields of

organic acids than those in *M. rosea* SV97 in all gas supplementation tests ($P < 0.05$) (Fig.

3). The efficiency of conversion of CH_4-C into organic acids was 4.8-7.0% and 0.7-1.8% (of

consumed CH_4-C) for *M. tundripaludum* SV96 and *M. rosea* SV97, respectively (Table S1).

This difference could be due to their different carbon assimilation pathways, i.e. that RuMP

pathway of Type I methanotrophs efficiently links to glycolytic pathway where pyruvate is

converted to organic acids, while the serine cycle of Type II methanotrophs has high flux

through acetyl-CoA (Kalyuzhnaya et al., 2015) which is derived into an intracellular storage

compound, like polyhydroxybutyrate (PHB), as previously observed in *M. rosea* SV97 during

isolation characterization (Wartiainen et al., 2006b). Regarding the production of organic

acids in Type II methanotrophs, which has not been widely reported, Costa et al. (2000)

observed methanotroph-driven CH_4 conversion into acetate which was subsequently

consumed by heterotrophs in the denitrification bioreactor using CH_4 as an electron donor.

Vecherskaya et al. (2009) also observed succinate, acetate and 2,3- butanediol excreted in

culture medium of Type II methanotroph, *Methylocystis parvus*, during CH_4 oxidation under

microaerobic and anaerobic conditions. Based on ^{13}C analysis in their study, those organic

acids were likely from PHB degradation during microaerobic conditions (5-10% O_2)

(Vecherskaya et al. 2009).

During 14-day incubation in all gas supplementation tests, organic acid concentrations

gradually accumulated corresponding with the increase of biomass concentration in both

methanotroph liquid cultures (Fig. 3). On days 10-14 onwards, different gas supplementation

tests likely induced different conditions to headspace including microaerobic (O_2-limited),

anaerobic and aerobic conditions. These conditions resulted in O_2/CH_4 molar ratios in

headspace of 0.2-0.3, < 0.01 and 2-10 for the addition of CH_4 + air, only CH_4 and only air,

respectively (Fig. S1). Interestingly, the three different gas supplementation tests in our study

showed similar total organic acid production yields (Fig. 4a and b). Lee et al. (2021)

compared the production of organic acids in *M. capsulatus* Bath culture under various

conditions including aerobic, oxygen-limited, sulfur-limited and nitrogen-limited conditions.

The authors observed acetate production in all studied conditions, and nitrate-nitrogen

limitation induced the highest acetate production (approximately 1.9 mmol-acetate g^{-1} CDW).

Furthermore, the previous studies on organic acid production by type I methanotrophs (Lee et

al., 2021; Takeuchi and Yoshioka, 2021) reported that acetate and formate were also

produced under aerobic conditions but in lower concentrations than in O_2-limited conditions.

In our study, however, the three different gas supplementation tests likely varied in the

distribution of organic acids contained in the spent medium, particularly, of *M.*

tundripaludum SV96, type I methanotrophs. Regarding organic acid production yields per

consumed CH_4-carbon, both formate (3.5%) and acetate (2.5%) were dominant in the test

with CH_4+air supplementation (test I), while only acetate (5.1%) was dominant in the test

with only air supplementation (test III) (Fig. 4c and d). These results suggest that it would be

possible to target dominant organic acid compounds by controlled feeding CH_4 and O_2. The

excretion of high formate concentration of *M. tundripaludum* SV96 at CH_4+air

supplementation (test I) might be due to imbalanced growth during O_2-limited conditions,

which was previously reported in type I methanotrophs by Gilman et al. (2017). This

circumstance also corresponded with pH reduction observed only in the test with CH_4+air

supplementation (test I) (Fig. S2).

Fig. 4.

Comparing to organic acid production in different Type I methanotroph species under

O_2-limited conditions, *M. tundripaludum* SV96 used in our study showed comparable and

higher organic acid yields per biomass production (CDW), i.e., for formate, acetate, and

succinate (Table 1). Those previous studies of organic acid production from microaerobic

CH_4 oxidation cultivated the methanotrophs with the initial headspace CH_4 and O_2

concentrations of 20% and 5%, respectively (Gilman et al., 2017, 2015; Kalyuzhnaya et al.,

2013). However, in this study, *M. tundripaludum* SV96 was prior cultivated aerobically

before allowing O_2 limiting condition to occur. Whether the prior cultivation in aerobic

conditions enhanced organic acid production in the subsequent microaerobic conditions

requires further studies. Additionally, the organic acid yields of *M. tundripaludum* SV96

obtained from our study (4.8-7.0% of consumed CH_4-carbon) were higher than those of *M.*

capsulatus Bath (<5% of consumed CH_4) (Lee et al., 2021), while *M. alcaliphilum* 20Z

enabled to convert 40-50% of the consumed CH_4 into mostly acetate and formate under O_2-

limited conditions (20% CH_4:5% O_2) (Kalyuzhnaya et al., 2013).

Table 1.

3.2. Utilization of organic acids rich spent medium of methanotrophs by *A. baylyi* ADP1

Wild type *A. baylyi* ADP1 cells enabled to grow in the organic acid-rich spent media of

both methanotrophs, containing formate, acetate, and succinate (and malate for *M.*

tundripaludum SV96). During 4-h incubation, *A. baylyi* ADP1 cells incubated in the spent

media from *M. tundripaludum* SV96 and *M. rosea* SV97 grew to an OD_{600nm} of 0.14 ± 0.01

and 0.12 ± 0.03, respectively (Fig. S3). Except for formate, acetate, succinate, and malate

were utilized as carbon sources for biomass assimilation (Fig. S3). *A. baylyi* ADP1 cells do

not utilize formate as a sole carbon source, but it is rather used for maintaining the cellular

redox balance (Kannisto et al., 2015). The growth of *A. baylyi* ADP1 was also confirmed by

the present of fatty acid fractions derived from *A. baylyi* ADP1 biomass on TLC plates.

Typically, *A. baylyi* ADP1 produce wax esters as carbon storage compounds which often

associated with growth (Mangayil et al., 2019). However, they were not detected in this study

(Fig. S4). This could be due to rapid degradation of the accumulated wax ester during carbon

limiting conditions. Under carbon limiting conditions *A. baylyi* ADP1 cells utilize the stored

carbon for maintaining cellular activities (Fixter et al., 1986; Salmela et al., 2018). The

application of using the spent media of methanotrophs for wax production would be possible

for further study but likely require higher quantity of organic acids.

3.3. 1-alkene synthesis from organic acids rich spent medium of methanotrophs

After confirming the growth of wild type *A. baylyi* ADP1 on the methanotroph spent

media, the organic acid-rich spent media were hereby used as carbon sources for 1-alkene

production by an engineered *A. baylyi* ADP1 '*tesA-undA*. The results showed that *A. baylyi*

ADP1 '*tesA-undA* completely utilized acetate (0.3 mM), succinate (0.05 mM), and malate

(0.2 mM) present in the spent media of methanotrophs, except formate, similar to wild type

ADP1 cells (Fig. 5a). The production of 1-undecene from *M. tundripaludum* SV96 spent

medium ($14.1 \pm 2.7\ \mu g\ L^{-1}$) was higher than that from *M. rosea* SV97 cultivations (1.0 ± 0.5

$\mu g\ L^{-1}$) (Fig. 5c), corroborating with the organic acid concentrations (Fig. 5a and 5b).

Likewise, the growth of *A. baylyi* ADP1 '*tesA-undA* was higher in spent medium from *M.*

tundripaludum SV96 ($0.115 \pm 0.06\ OD_{600nm}$) than from *M. rosea* SV97 (0.070 ± 0.054

OD_{600nm}) (Fig. 5c). The production of 1-undecene was not detected in the control cultivations

(Fig. S5). In addition, the growth occurred during the co-cultivation was from solely *A. baylyi*

ADP1 '*tesA-undA* as both methanotrophs could not grow without CH_4 as a carbon source.

Fig. 5.

The obtained 1-undecene concentrations in this study (1.0 and 14.1 µg L^{-1}) were lower

than that previously reported for *A. baylyi* ADP1 '*tesA-undA* (Luo et al., 2019; Salmela et al.,

2020). This phenomenon can be attributed towards the higher substrate concentrations used

in those studies. For example, Luo et al. (2019) developed a high ferulate-tolerant *A. baylyi*

ADP1 strain for 1-undecene production using adaptive laboratory evolution. The authors

tested the use of high concentration of ferulate (100 mM) as a sole carbon source and

obtained the 1-undecene production titer of 72 ± 7.5 µg L^{-1} (yield of 1.0 µmol mol $C_{substrate}^{-1}$).

In another study, Salmela et al. (2020) obtained 1-undecene concentration of up to $\sim107 \pm 8$

µg L^{-1} from a two-stage system for 1-undecene production from cellulose which was

converted into organic metabolites by *Clostridium cellulolyticum* (containing 5.2 mM

glucose, 4.9 mM acetate and 6.8 mM lactate). The authors reported the highest 1-undecene

production yield of ~35 µmol mol $C_{substrate}^{-1}$ using lactate as a substrate. In our study, 1-

undecene production from the spent media of methanotrophs was promising and comparable

to those in previous studies, resulting in 1-undecene yields of 68.9 ± 11.6 and 40.6 ± 19.8

µmol mol $C_{substrate}^{-1}$ for *M. tundripaludum* SV96 and *M. rosea* SV97, respectively. The

obtained carbon recovery for 1-undecene production accounted for 0.065% and 0.045% of

consumed total organic acids-carbon for *M. tundripaludum* SV96 and *M. rosea* SV97,

respectively (Table S2). The spent media of methanotrophs likely did not contain

intermediate compounds being toxic to growth and 1-undecene production of *A. baylyi* ADP1

'*tesA-undA*. Furthermore, the media could be directly used for the cultivation without

purification or additional downstream processing.

The results indicate that the spent media from microaerobic fermentation by

methanotrophs is an excellent carbon source for heterotrophs. This two-stage bacterial

process extended the range of CH_4-derived products to C11 compound (1-undecene). In

further studies, the process optimization, scale-up as well as statistical optimization of the

system for methanotroph cultivation will be developed to enhance organic acid concentration,

which will consequently improve the production efficiency of *A. baylyi* ADP1. As produced

organic acids, particularly acetate and succinate, by methanotrophs were also observed under

aerobic conditions, this study can also lead to develop a single system for CH_4 conversion

into high value-added products using coculture of methanotrophs and *A. baylyi* ADP1.

4. Conclusion

A two-step bioprocess setup was designed for successful integration of microaerobic

methane fermentation with aerobic synthesis. This study shows a proof of concept for

integrating greenhouse gas (GHG) utilization and platform chemical production: the

application of organic acids produced by methanotrophs for 1-undecene (C11) production. A

Type I methanotroph, *M. tundripaludum* SV96, showed higher potential for organic acid

production than a Type II methanotroph, *M. rosea* SV97 under aerobic and microaerobic

conditions. The organic acids-rich spent media of methanotrophs could be directly used as a

medium for the cultivation of the wild type *A. baylyi* ADP1 and the engineered *A. baylyi*

ADP1 '*tesA-undA* without additional downstream processes or purification. Acetate,

succinate, and malate contained in the spent media were completely utilized by *A. baylyi*

ADP1 '*tesA-undA* for their 1-undecene production. The highest yield of 1-undecene was

obtained from spent medium of *M. tundripaludum* SV96 at $68.9 \pm 11.6\ \mu mol\ mol\ C_{substrate}^{-1}$.

Acknowledgement

The research work was supported by Kone foundation [grant number 201803224] and

Academy of Finland [grant number 323214]. The authors would like to acknowledge Jin Luo

for helping with using GC-MS for 1-undecene measurement.

Appendices

Supplementary data is provided as a separate file attached with the manuscript.

References

Cantera, S., Muñoz, R., Lebrero, R., López, J. C., Rodríguez, Y., & García-Encina, P. A.

(2018). Technologies for the bioconversion of methane into more valuable products.

Current Opinion in Biotechnology, *50*, 128–135.

https://doi.org/10.1016/j.copbio.2017.12.021

Cao, Q., Liu, X., Ran, Y., Li, Z., & Li, D. (2019). Methane oxidation coupled to

denitrification under microaerobic and hypoxic conditions in leach bed bioreactors.

Science of the Total Environment, *649*, 1–11.

https://doi.org/10.1016/j.scitotenv.2018.08.289

Chen, S., & Smith, A. L. (2018). Methane-driven microbial fuel cells recover energy and

mitigate dissolved methane emissions from anaerobic effluents. *Environmental Science:*

Water Research and Technology, *4*(1), 67–79. https://doi.org/10.1039/c7ew00293a

Costa, C., Dijkema, C., Friedrich, M., Garcõâ A-Encina, P., Fernaâ Ndez-Polanco, F., &

Stams, A. J. M. (2000). Denitrification with methane as electron donor in oxygen-

limited bioreactors. *Applied Microbiology and Biotechnology*, *53*, 754–762.

Fixter, L. M., Nagi, M. N., Mccormacks, J. G., & Fewson, C. A. (1986). Structure,

distribution and function of wax esters in *Acinetobacter calcoaceticus*. *Journal of*

General Microbiology, *132*, 3147–3157.

Ge, X., Yang, L., Sheets, J. P., Yu, Z., & Li, Y. (2014). Biological conversion of methane to

liquid fuels: Status and opportunities. *Biotechnology Advances*, *32*(8), 1460–1475.

https://doi.org/10.1016/j.biotechadv.2014.09.004

Gilman, A., Fu, Y., Hendershott, M., Chu, F., Puri, A. W., Smith, A. L., Pesesky, M.,

Lieberman, R., Beck, D. A. C., & Lidstrom, M. E. (2017). Oxygen-limited metabolism

in the methanotroph *Methylomicrobium buryatense* 5GB1C. *PeerJ*, *5*, e3945.

https://doi.org/10.7717/peerj.3945

Gilman, A., Laurens, L. M., Puri, A. W., Chu, F., Pienkos, P. T., & Lidstrom, M. E. (2015).

Bioreactor performance parameters for an industrially-promising methanotroph

Methylomicrobium buryatense 5GB1. *Microbial Cell Factories*, *14*(1), 182.

https://doi.org/10.1186/s12934-015-0372-8

Hanson, R. S., & Hanson, T. E. (1996). Methanotrophic Bacteria. *Microbiological Reviews*,

60(2), 439–471. http://mmbr.asm.org/

Harmsen, M., van Vuuren, D. P., Bodirsky, B. L., Chateau, J., Durand-Lasserve, O., Drouet,

L., Fricko, O., Fujimori, S., Gernaat, D. E. H. J., Hanaoka, T., Hilaire, J., Keramidas, K.,

Luderer, G., Moura, M. C. P., Sano, F., Smith, S. J., & Wada, K. (2020). The role of

methane in future climate strategies: mitigation potentials and climate impacts. *Climatic*

Change, *163*, 1409–1425. https://doi.org/10.1007/s10584-019-02437-2

IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working

Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate

Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N.

Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R.

Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)].

Cambridge University Press, Cambridge, UK. *In Press*.

https://www.ipcc.ch/report/ar6/wg1/ (accessed 16.11.2021)

Kalyuzhnaya, M. G., Yang, S., Rozova, O. N., Smalley, N. E., Clubb, J., Lamb, A., Gowda,

G. A. N., Raftery, D., Fu, Y., Bringel, F., Vuilleumier, S., Beck, D. A. C., Trotsenko, Y.

A., Khmelenina, V. N., & Lidstrom, M. E. (2013). Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nature Communications*, *4*, 2785. https://doi.org/10.1038/ncomms3785

Kalyuzhnaya, Marina G., Puri, A. W., & Lidstrom, M. E. (2015). Metabolic engineering in methanotrophic bacteria. *Metabolic Engineering*, *29*, 142–152. https://doi.org/10.1016/j.ymben.2015.03.010

Kannisto, M. S., Mangayil, R. K., Shrivastava-Bhattacharya, A., Pletschke, B. I., Karp, M. T., & Santala, V. P. (2015). Metabolic engineering of *Acinetobacter baylyi* ADP1 for removal of *Clostridium butyricum* growth inhibitors produced from lignocellulosic hydrolysates. *Biotechnology for Biofuels*, *8*, 198. https://doi.org/10.1186/s13068-015-0389-6

Lai, C. Y., Zhong, L., Zhang, Y., Chen, J. X., Wen, L. L., Shi, L. D., Sun, Y. P., Ma, F., Rittmann, B. E., Zhou, C., Tang, Y., Zheng, P., & Zhao, H. P. (2016). Bioreduction of chromate in a methane-based membrane biofilm reactor. *Environmental Science and Technology*, *50*(11), 5832–5839. https://doi.org/10.1021/acs.est.5b06177

Lee, H., Baek, J. I., Lee, J. Y., Jeong, J., Kim, H., Lee, D. H., Kim, D. M., & Lee, S. G. (2021). Syntrophic co-culture of a methanotroph and heterotroph for the efficient conversion of methane to mevalonate. *Metabolic Engineering*, *67*, 285–292. https://doi.org/10.1016/j.ymben.2021.07.008

Liu, T., Li, J., Khai Lim, Z., Chen, H., Hu, S., Yuan, Z., & Guo, J. (2020). Simultaneous removal of dissolved methane and nitrogen from synthetic mainstream anaerobic effluent. *Environmental Science and Technology*, *54*(12), 7629–7638. https://doi.org/10.1021/acs.est.0c00912

López, J. C., Quijano, G., Souza, T. S. O., Estrada, J. M., Lebrero, R., & Muñoz, R. (2013). Biotechnologies for greenhouse gases (CH_4 , N_2O, and CO_2) abatement: State of the art

and challenges. *Applied Microbiology and Biotechnology*, *97*, 2277–2303.

https://doi.org/10.1007/s00253-013-4734-z

Luo, J., Lehtinen, T., Efimova, E., Santala, V., & Santala, S. (2019). Synthetic metabolic

pathway for the production of 1-alkenes from lignin-derived molecules. *Microbial Cell*

Factories, *18*, 48. https://doi.org/10.1186/s12934-019-1097-x

Mangayil, R., Efimova, E., Konttinen, J., & Santala, V. (2019). Co-production of 1,3

propanediol and long-chain alkyl esters from crude glycerol. *New Biotechnology*, *53*,

81–89. https://doi.org/10.1016/j.nbt.2019.07.003

Okonkwo, O., Lakaniemi, A. M., Santala, V., Karp, M., & Mangayil, R. (2018). Quantitative

real-time PCR monitoring dynamics of *Thermotoga neapolitana* in synthetic co-culture

for biohydrogen production. *International Journal of Hydrogen Energy*, *43*(6), 3133–

3141. https://doi.org/10.1016/j.ijhydene.2017.12.002

Saarela, T., Rissanen, A. J., Ojala, A., Pumpanen, J., Aalto, S. L., Tiirola, M., Vesala, T., &

Jäntti, H. (2020). CH_4 oxidation in a boreal lake during the development of hypolimnetic

hypoxia. *Aquatic Sciences*, *82*(2), 19. https://doi.org/10.1007/s00027-019-0690-8

Salmela, M., Lehtinen, T., Efimova, E., Santala, S., & Mangayil, R. (2018). Metabolic

pairing of aerobic and anaerobic production in a one-pot batch cultivation.

Biotechnology for Biofuels, *11*(1), 187. https://doi.org/10.1186/s13068-018-1186-9

Salmela, M., Lehtinen, T., Efimova, E., Santala, S., & Santala, V. (2020). Towards

bioproduction of poly-α-olefins from lignocellulose. *Green Chemistry*, *22*(15), 5067–

5076. https://doi.org/10.1039/d0gc01617a

Santala, S., Efimova, E., Kivinen, V., Larjo, A., Aho, T., Karp, M., & Santala, V. (2011).

Improved triacylglycerol production in *Acinetobacter baylyi* ADP1 by metabolic

Engineering. *Microbial Cell Factories*, *10*, 36.

http://www.microbialcellfactories.com/content/10/1/36

Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B.,

Raymond, P. A., Dlugokencky, E. J., Houweling, S., K. Patra, P., Ciais, P., K. Arora, V.,

Bastviken, D., Bergamaschi, P., R. Blake, D., Brailsford, G., Bruhwiler, L., M. Carlson,

K., Carrol, M., … Zhuang, Q. (2020). The global methane budget 2000-2017. *Earth*

System Science Data, *12*(3), 1561–1623. https://doi.org/10.5194/essd-12-1561-2020

Sheets, J. P., Ge, X., Li, Y. F., Yu, Z., & Li, Y. (2016). Biological conversion of biogas to

methanol using methanotrophs isolated from solid-state anaerobic digestate. *Bioresource*

Technology, *201*, 50–57. https://doi.org/10.1016/j.biortech.2015.11.035

Strong, P. J., Kalyuzhnaya, M., Silverman, J., & Clarke, W. P. (2016). A methanotroph-based

biorefinery: Potential scenarios for generating multiple products from a single

fermentation. In *Bioresource Technology* (Vol. 215, pp. 314–323). Elsevier Ltd.

https://doi.org/10.1016/j.biortech.2016.04.099

Sun, M. T., Yang, Z. M., Fu, S. F., Fan, X. L., & Guo, R. B. (2018). Improved methane

removal in exhaust gas from biogas upgrading process using immobilized methane-

oxidizing bacteria. *Bioresource Technology*, *256*, 201–207.

https://doi.org/10.1016/j.biortech.2018.02.020

Takeuchi, M., & Yoshioka, H. (2021). Acetate excretion by a methanotroph, *Methylocaldum*

marinum S8, under aerobic conditions. *Bioscience, Biotechnology, and Biochemistry*.

https://doi.org/10.1093/bbb/zbab150

Vecherskaya, M., Dijkema, C., Saad, H. R., & Stams, A. J. M. (2009). Microaerobic and

anaerobic metabolism of a *Methylocystis parvus* strain isolated from a denitrifying

bioreactor. *Environmental Microbiology Reports*, *1*(5), 442–449.

https://doi.org/10.1111/j.1758-2229.2009.00069.x

Wartiainen, I., Hestnes, A. G., McDonald, I. R., & Svenning, M. M. (2006a). *Methylobacter*

tundripaludum sp. nov., a methane-oxidizing bacterium from Arctic wetland soil on the

Svalbard islands, Norway (78° N). *International Journal of Systematic and Evolutionary*

Microbiology, *56*(1), 109–113. https://doi.org/10.1099/ijs.0.63728-0

Wartiainen, I., Hestnes, A. G., McDonald, I. R., & Svenning, M. M. (2006b). *Methylocystis*

rosea sp. nov., a novel methanotrophic bacterium from Arctic wetland soil, Svalbard,

Norway (78° N). *International Journal of Systematic and Evolutionary Microbiology*,

56(3), 541–547. https://doi.org/10.1099/ijs.0.63912-0

Table 1. The production of organic acids and other metabolites obtained from different Type I methanotrophs cultivated under O_2 limiting conditions.

Methanotrophs	Test mode	Volume	OD (CDW, g L^{-1})	Initial $CH_4:O_2$ (%)	Yield (mmol g CDW^{-1})				Note	Reference
					Formate	Acetate	Succinate	Malate		
M. tundripaludum SV96	Batch	15 mL in 120 mL vial	~2 (0.6)	20:15 (10:3)[†]	3.20 ± 0.9	1.15 ± 0.17	0.16 ± 0.01	0.07 ± 0.01		Our study
Methylomicrobium alcaliphilum 20Z	fed-batch/chemostat	1 L in 2 L chemostat	~2 ± 0.2	20:5	1.53 ± 0.27	0.020	0.0025	ND	H_2, lactate detected	Kalyuzhnaya et al. (2013)
Methylomicrobium buryatense 5GB1	continuous	1 L in 2.5 L vessel	(0.45-0.46)	20:5	0.29-0.30	0.036-0.085	ND	ND	lactate detected	Gilman et al. (2015)
	fed-batch	1 L in 2.5 L vessel	(0.31-0.34)	10:5	0.40-0.67	0.096-0.123	ND	ND	lactate detected	
Methylomicrobium buryatense 5GB1C (the aa3 cytochrome oxidase mutant strain)	Batch	1 L in 2.5 L vessel	(0.21)	20:5	1.22-3.00	0.081-0.086	ND	ND		Gilman et al. (2017)

Note: CDW, cell dry weight; ND, no data; [†]The $CH_4:O_2$ in headspace when O_2 limiting conditions occurring in the system

Figure Captions

Fig. 1. The principle of a two-stage bacterial process. The estimated pathway of the production of various organic acids could occur in methanotrophs (a). Organic acid rich spent medium was fed to the engineered *Acinetobacter baylyi* ADP1 for alkene production (b). Aerobic CH_4 oxidation pathway in Type I (RuMP cycle) and Type II (Serine cycle) methanotrophs (black arrows) and the possible pathway of CH_4 oxidation under O_2-limiting conditions via glycolysis-based CH_4 fermentation mode (brown line). Abbreviations: **RuMP**, ribulose monophosphate; **EMP**, Embden–Meyerhof–Parnas; **TCA**, Tricarboxylic acid; **MMO**, methane monooxygenase; **MDH**, methanol dehydrogenase; **FaDH**, formaldehyde dehydrogenase; **H4F**, tetrahydrofolate pathway; **H4MPT**, tetrahydromethanopterin pathway; **FDH**, formate dehydrogenase; **PTa**, phosphate acetyl-transferase; **AcK**, acetate kinase; **XFP**, phophoketolase xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Fig. 2. Experimental set up for methanotrophs under three different gas supplementation schemes applied on day 7 including test I: both CH_4 and air added, test II only CH_4 added and test III only air added.

Fig. 3. The accumulation of organic acids production and biomass production during the 14-day incubation of *M. tundripaludum* SV96 (a-c) and *M. rosea* SV97 (d-f) under three different gas supplementation schemes applied on day 7: (i) both CH_4 and air added (a and d), (ii) only CH_4 added (b and e), and (iii) only air added (c and f). Error bars indicate the standard deviation of triplicate samples.

Fig. 4. Carbon mass balance applied to CH_4 oxidation to CO_2, biomass, and organic acids (a and b) and distribution of each organic acid compound (c, d) of *M. tundripaludum* SV96 (left) and *M. rosea* SV97 (right) in three different gas supplementation tests for the 14-day incubation. Error bars indicate the standard deviation of sum of total yields in triplicate.

Fig. 5. Organic acids utilization by *A. baylyi* ADP1 *'tesA-undA* in spent media obtained from the cultivation of *M. tundripaludum* SV96 (MT) (a) and *M. rosea* SV97 (MR) (b) under O_2 limiting conditions. Growth and 1-undecene production from *A. baylyi* ADP1 '*tesA-undA* after 24 h incubation (c). ΔOD indicates the difference in OD_{600nm} of *A. baylyi* ADP1 '*tesA-undA* between 0 and 24 h. Error bars indicate the standard deviation of four replicate samples. The spent media from methanotrophs fermentation without *A. baylyi* ADP1 were used as control (Control-24 h).

Fig. 1. The principle of a two-stage bacterial process. The estimated pathway of the production of various organic acids could occur in methanotrophs (a). Organic acid rich spent medium was fed to the engineered *Acinetobacter baylyi* ADP1 for alkene production (b). Aerobic CH_4 oxidation pathway in Type I (RuMP cycle) and Type II (Serine cycle) methanotrophs (black arrows) and the possible pathway of CH_4 oxidation under O_2-limiting conditions via glycolysis-based CH_4 fermentation mode (brown line). Abbreviations: **RuMP**, ribulose monophosphate; **EMP**, Embden–Meyerhof–Parnas; **TCA**, Tricarboxylic acid; **MMO**, methane monooxygenase; **MDH**, methanol dehydrogenase; **FaDH**, formaldehyde dehydrogenase; **H₄F**, tetrahydrofolate pathway; **H₄MPT**, tetrahydromethanopterin pathway; **FDH**, formate dehydrogenase; **PTa**, phosphate acetyl-transferase; **AcK**, acetate kinase; **XFP**, phophoketolase xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

M. tundripaludum SV96/*M. rosea* SV97
Initial gas in headspace: $20\%CH_4 + 80\%$ air

Headspace supplementation after 7-day incubation

(i)   (ii)   (iii)

Test I: CH_4 + air   Test II: only CH_4 Test III: only air
$(20\%CH_4 + 80\%air)$ $(20\%CH_4 + 80\%N_2)$ $(20\%air + 80\%N_2)$

Fig. 2. Experimental set up for methanotrophs under three different gas supplementation schemes applied on day 7 including test I: both CH_4 and air added, test II only CH_4 added and test III only air added.

Fig. 3. The accumulation of organic acids production and biomass production during the 14-day incubation of *M. tundripaludum* SV96 (a-c) and *M. rosea* SV97 (d-f) under three different gas supplementation schemes applied on day 7: (i) both CH_4 and air added (a and d), (ii) only CH_4 added (b and e), and (iii) only air added (c and f). Error bars indicate the standard deviation of triplicate samples.

Fig. 4. Carbon mass balance applied to CH_4 oxidation to CO_2, biomass, and organic acids (a and b) and distribution of each organic acid compound (c, d) of *M. tundripaludum* SV96 (left) and *M. rosea* SV97 (right) in three different gas supplementation tests for the 14-day incubation. Error bars indicate the standard deviation of sum of total yields in triplicate.

Fig. 5. Organic acids utilization by *A. baylyi* ADP1 *'tesA-undA* in spent media obtained from the cultivation of *M. tundripaludum* SV96 (MT) (a) and *M. rosea* SV97 (MR) (b) under O_2 limiting conditions. Growth and 1-undecene production from *A. baylyi* ADP1 *'tesA-undA* after 24 h incubation (c). ΔOD indicates the difference in OD_{600nm} of *A. baylyi* ADP1 *'tesA-undA* between 0 and 24 h. Error bars indicate the standard deviation of four replicate samples. The spent media from methanotrophs fermentation without *A. baylyi* ADP1 were used as control (Control-24 h).