1

In Vivo Multi-Day Calcium Imaging of CA1 Hippocampus in Freely Moving Rats Reveals a High Preponderance of Place Cells and No Detectable Photobleaching

Hannah S Wirtshafter[1][*], John F Disterhoft[1]

*Lead contact: hsw@northwestern.edu (HSW)

1. Department of Neuroscience, Northwestern University Feinberg School of Medicine, Chicago, IL, USA.

Abstract

Calcium imaging using GCaMP calcium indicators and miniature microscopes has been used to image cellular populations during long timescales and in different task phases, as well as to determine neuronal circuit topology and organization. Because the hippocampus (HPC) is essential for many tasks of memory, spatial navigation, and learning, calcium imaging of large populations of HPC neurons can provide new insight on cell changes and organization over time during these tasks. To our knowledge, all reported HPC *in vivo* calcium imaging experiments have been done in mouse. However, rats have many behavioral and physiological experimental advantages over mice, and, due to their larger size, rats are able to support larger implants, thereby enabling the recording of a greater number of cells. In this paper, we present the first *in vivo* calcium imaging from CA1 hippocampus in freely moving rats. Using GCaMP7c and the UCLA Miniscope, we demonstrate that hundreds of cells (mean 240+-90 cells per session, maximum 428 cells) can reliably be visualized and held across weeks, and that calcium events in these cells are correlated with periods of movement. We additionally show proof of method by showing that an extremely high percent of place cells (82.3%+-8.1%, far surpassing the percent seen during mouse calcium imaging) can be recorded on a navigational task, and that these place cells enable accurately decoding of animal position. Finally, we show that calcium imaging in rats is not prone to photobleaching during hour-long recordings and that cells can be reliably recorded for an hour or more per session. A detailed protocol for this technique, including notes on the numerous parameter changes needed to use Ca2+ imaging in rats, is included in the Materials and Methods section, and implications of these advancements are discussed.

Introduction

The explosion in the use of *in vivo* calcium imaging has allowed the visualization of hundreds of identified neurons over long time periods [1-8]. Calcium (Ca2+) imaging using GCaMP calcium indicators and miniature microscopes has been used to image cellular populations during long timescales [9-15] and in different task phases [16-18], as well as to determine neuronal circuit topogrophy and organization [19-22].

The hippocampus (HPC) has been studied with cellular resolution using *in vivo* electrophysiology, but this technique does not allow the same cells to be definitively followed over days, nor does it allow the visualization of cellular organization within the structure [23-27]. *In vivo* calcium imaging is particularly well suited for imaging the rodent HPC, as the orientation of the horizontal cell layer permits imaging of large numers of neurons with insertion of a 1mm diameter or smaller lens [12, 13, 22, 28]. Because the hippocampus is essential for many tasks involving memory [15, 29-37], spatial navigation [38-43], and learning [44-47], Ca2+ imaging of large populations of HPC neurons can shed new insight on cell changes and organization over time during these tasks.

To our knowledge, all reports of hippocampal *in vivo* calcium imaging have been done in mouse (although calcium imaging has been used to visualize other structures in rat [48-51]). Using calcium imaging in rat hippocampus requires optimizing a number of methodological parameters, including aspiration and lens implantation techniques, to account for the increased thickness of the rat alveus compared to mouse which makes focusing on the CA1 cell layer difficult [52-54]. However, rats have many basic experimental advantages over mice: First, rats are able to learn and perform more complex tasks than mice, and employ more advanced strategies, and are thus suited for a wider variety of behavioral experiments [55-60]. Second, compared to mouse physiology, rat physiology for a number of conditions and disorders (including addiction, aging, and schizophrenia, among others) more closely mirrors human physiology [61-64]. Third, rats are less stressed by handling than mice, thus introducing less variability in experimental procedures [65].

Rats are also an advantageous organism to use for calcium imaging: Due to their

larger size, rats are able to support larger implants, including a wider diameter imaging

lens and a larger camera, thereby enabling the recording of a greater number of cells [2,

66, 67]. Additionally, rat hippocampal neurons are less excitable than those in mice [54,

59], and rats are less prone to post-operative seizures than mice [68-70]. Rats may

therefore, compared to mice, be at a decreased risk of the seizures (and potentially to

cell death) seen in mice after prolonged GCaMP expression [5, 6, 71-73].

Although *in vivo* Ca2+ imaging would appear to hold substantial promise for the

study of hippocampus in freely moving rats, there is currently no direct evidence that it

will work in this system. To our knowledge, there are no studies employing Ca2+

imaging in rats, and the technique appears to have obtained a reputation for being

difficult to use successfully to image rat CA1. In this paper, we present the first (to our

knowledge) *in vivo* Ca2+ recording from CA1 hippocampus in freely moving rats. We

demonstrate that hundreds of cells are reliably visualized and held across weeks using

this technique, and that there is little to no loss of fluorescence even in long (30-60m)

recording sessions. We additionally show proof of method by showing that a uniquely

high percentage (>80%) of place cells were recorded on a navigational task involving

retrieving food reward on a linear track. A detailed protocol for this technique, including

notes on the numerous parameter changes needed to use Ca2+ in rats, is included in

the Materials and Methods section, and implications of these advancements are

discussed.

Results

To initiate Ca2+ imaging, we injected AAV9-Syn-GCaMP7c below the CA1 cell

layer in five male Fischer 344 x Brown Norway rats (Figure 1a, see Materials and

Methods) [74]. After injection, in the same surgery, cortex and corpus collosum were

aspirated through a craniotomy until the alveus was clearly visible (Figure 1b, see

Materials and Methods). A 2mm diameter GRIN lens was inserted into the craniotomy

and cemented in place. Three animals were checked for GCaMP expression four weeks

after surgery, and all three animals showed expression and were subsequently affixed with a base plate to secure the position of the miniscope. The other two animals were checked six weeks after surgery and showed expression and were base plated. After the completion of the experiment, animals were sacrificed and expression of GCaMP in CA1 pyramidal cells was verified with microscopy (Figure 1c).

Animals were then food deprived to 85% body weight and trained on a 2.64m linear track (Figure S1a). Animals were rewarded with grain pellets on alternating sides of the linear track and quickly learned to run back and forth on the track (Figure S1b). A visit to one reward well signified a trial. Animals were run for a minimum of 30 minutes or 50 trials, whichever came later, with total track time not to exceed one hour per session. Cells were recorded for 7 days for one animal, 11 days for one animal, and 16 days for two animals (n=4 rats, one of the five animals was eliminated when the implant was lost).

Six sessions were analyzed from each animal, with first and last recorded sessions automatically included. These sessions were analyzed for cellular activity using the CNMF-E method implemented by the CIATAH software package [75-77] (See Materials and Methods). A total of 5761 neurons were recorded over the 24 sessions, with a maximum of 428 cells from a single session (mean of 239.6+-90.0 cells) (Figure 2a-d). Calcium activity appeared to be heavily modulated by the animal's activity, with periods of higher activity when the animal was running rather than stationary on the track (Figure 2c). The number of cells recorded from each animal across sessions was relatively constant (Figure 2d).

We then determined if there were place cells on any of the trials. A cell was considered to be a place cell if it met two criteria: First, the cell must have a mean firing rate greater than 0.01Hz. Second, the mutual information (MI) computed in either direction of travel must be greater than 95% of MI scores computed 500 times from shuffled cells [78]. Based on these criteria, all trials contained place cells (Figure 3a, Figure S2). An average of 82.3%+-8.1% of cells with firing rates >0.01Hz were place cells (Figure 3b-c).

We then aimed to determine whether place cell firing was sufficient to decode the

position of the animal. We split the track in 10cm segments and used Bayesian

decoding to determine the position of the animal in non-overlapping 0.5 second

increments [79]. We found that the best positional decoding was accurate to a median

error of 4.3cm (1.6% of the 2.64m track). Across all animals, accuracy ranged from a

median of 4.3cm to 13.6cm (decoding errors were below 10cm in 23 sessions and

above 10cm in one session) with a median error of 5.9cm (Figure 4a). To confirm that

decoding accuracy was better than chance we shuffled firing rate per position bins and,

using the shuffled data, decoded position. Median error for shuffled data ranged from

59.1cm to 77.6cm, which was significantly worse than non-shuffled data (Kruskal-Wallis

test, $x^2(47)= 35.3$, $p=2.9*10^{-9}$) (Figure 4b).

We then aimed to determine if the same cells could be held over the duration of

our recordings (six recording sessions, spanning 7, 11, or 16 days). 40% of the cells

imaged on day one of the 7 day session were also present on day seven. Of the rat run

for 11 days, 31% of the cells present on day one were also present on day eleven. For

the two rats run 16 days, 28% and 50% of cells, respectively, that were present on day

one were also present on day sixteen. Across all four animals, cells that were imaged

on day one of recording had, on average, a 72+-6% chance of being imaged on at least

one subsequent day of recording (Figure 5a). (Of note, recordings each day were made

to optimize total cell number in each day as proof of method, rather than to maintain the

same landmarks or focal planes.)

Finally, our recording times lasted up to an hour per day, but we observed

obvious no photobleaching within our sessions. We thus aimed to quantify if GCaMP

photobleaching occurred within each session. To determine this, we compared the

mean fluorescence of each cell for five minutes at the start of each session to the mean

fluorescence of each cell for five minutes at the end of the session [80]. We found there

was no consistent loss of fluorescence (mean change = .008, median change = -

$2.01*10^{-4}$) (Figure 5b). In 22 out of the 24 sessions, the end mean fluorescence of each

cell was not less than the starting mean fluorescence (one-tailed paired t-test, p>0.05),

and for 6 of these sessions, the end mean fluorescence was actually greater than the

starting fluorescence (one-tailed paired t-test, $p < 0.05$). Two sessions (8% of sessions)

showed a decrease in fluorescence (one-tailed paired t-test, $t(368) = 2.47$, $p < 0.01$, and

$t(231) = 1.98$, $p < 0.05$). We questioned whether the lack of decrease in fluorescence

might have been due to an increase in calcium events in the latter part of the session,

so we compared the number of events for each cell in the early five-minute period with

the late five-minute period. In every session, there were not significantly more calcium

events in the latter five-minute period than in the former (one-tailed paired t-test,

$p > 0.05$). To additionally control for the number of calcium events, we normalized the

fluorescence in each five-minute period by the number of events in the period, and then

found the change in fluorescence from beginning to end of session (mean change = -

$1.77 \ast 10^{-4}$, median change = $-5.7 \ast 10^{-8}$)(Figure 5c). Even after normalization, still only two

sessions (although a different two) showed a decrease in fluorescence from the start

period to the end period (one-tailed paired t-test, $t(226) = 1.7$, $p = 0.04$, and $t(349) = 2.1$,

$p = 0.02$), while three sessions showed an increase (one-tailed paired t-test, $t(123) = -1.9$;

$p = 0.03$, $t(173) = -3.0$, $p < 0.005$; $t(166) = -4.0$, $p < 0.00005$). There was therefore no

consistent loss of fluorescence within each session.

Discussion

In this manuscript, we have demonstrated calcium imaging recordings can be

performed in the CA1 region of the hippocampus in freely moving rats. These

recordings can capture hundreds of cells in a single recording session, including a high

number and percentage of place cells sufficient to decode the animal's position to within

centimeters of accuracy. We additionally showed that recordings are stable across time,

and that there is no evidence of significant photobleaching during recording. Calcium

imaging in rat will allow for more behaviorally complex and translational experiments

than have been done in mouse.

Significantly, we found a higher percentage of place cells in our calcium imaging

experiments in rats than those reported in mouse CA1 calcium imaging experiments,

even after the mice experiments were corrected for cells with low spiking rates. (The

studies with mice reported 20-30% of imaged CA1 cells were place cells, which is also

substantially below most electrophysiology experiments [14, 81]). Previous work has

reported that mouse place cells are less spatially tuned, contain less spatial information,

and have less stable place fields than in rat [59, 82], so some of the difference in place

cell percent may be due to species differences in spatial representations. However,

there are also likely physiological differences between rats and mice that affect the type

of activity captured in calcium imaging recordings. CA1 cells in mice are more excitable

than those in rat [54, 59], and prolonged GCaMP expression in mice has been shown to

lead to cell hyperexcitability, seizures, and even cell death [5, 6, 71-73]. The higher

baseline excitably of mouse cells, combined with a further increase in cellular excitability

after GCaMP expression may result in a lower threshold for cell firing and less defined

place fields, less stable fields, and, therefore, a seemingly lower number of place cells

compared to electrophysiology recordings [59]. Because rats have a lower baseline

excitability, they may be less prone to these GCaMP effects, although further slice

electrophysiology work will need to be done to directly compare the effect of GCaMP on

mouse and rat neurons.

The percentage of hippocampal CA1 cells found to be place cells using calcium

recordings (82.3%) was also slightly above findings from electrophysiological data,

which report that 35-75% of CA1 pyramidal cells recorded on a navigational task are

place cells (the number varies widely depending on task parameters) [83-85]. There are

multiple possible reasons related to recording technology as to why our calcium imaging

experiments detected a higher percentage of place cells than traditional

electrophysiology experiments: First, the GCaMP we used (AAV-syn-jGCaMP7c)

appeared to be preferentially expressed in pyramidal neurons (Figure 1c), which has

previously been reported at specific titers [74]. We are therefore preferentially imaging

cells that are expected to have place fields; while place information can be extracted

from CA1 inhibitory interneurons, they do not have place fields [86]. (Although

interneurons are typically removed from place field electrophysiological place cell

analysis, this can be difficult to do in practice.) Second, in electrophysiology

experiments with moveable electrodes (such as tetrodes), electrodes are typically

positioned while rats are inside a "sleep box" or other non-training apparatus until cells

are detected. The animal is then placed on a track for experimentation. This practice

optimizes cell detection for place cells that are active in the "sleep box" rather than on

the track. Because calcium imaging uses a fixed lens location and images a large

number of cells at once, recording is not optimized for a particular environment. This

may result in the recording of a higher percentage of active place cells on the recording

apparatus. Finally, compared to electrophysiology, calcium imaging provides additional

characteristics (cell shape, size, and brightness) by which to separate adjacent or

nearby cells. Because conflation of multiple cells may disguise place fields and disrupt

place field measurements (such as mutual information or bits per spike), the additional

resolution afforded by calcium imaging may allow for more accurate place cell

differentiation and identification.

In addition to potentially allowing the recording of a higher number of place cells,

the lower excitability of rat CA1 neurons, compared to mouse, also has other

advantages. We saw no behavioral evidence of post-operative seizures in any of our

implanted animals, and no behavioral or imaging evidence of seizures several months

post viral injection and lens implementation. This is in contrast to calcium imaging

experiments in mice, which frequently report that animals have to be removed from

behavior or analysis due to the development of seizure activity [71, 87].

Rats' larger weight and body size also allows the implantation of a larger lens

(we used a 2mm diameter lens, as opposed to the 0.5mm or 1.0mm lenses typically

used in mouse CA1 imaging, and believe larger lenses could also be used), which in

turn allows for the imaging of larger areas and more cells. It also appears that, unlike in

mouse calcium imaging experiment, where cells are often not imaged for longer than

10min periods to avoid photobleaching [8, 13, 78], cells in rat hippoocampus can be

reliably recorded for an hour or more without any adverse effects on brightness (Figure

5b-c). Although the reason for this remains unclear, there is evidence that larger cells

may show less photobleaching [88]. The larger pyramidal neurons in the rat than mouse

[54] may therefore contribute to the reduction of photobleaching. Rats' lower baseline

excitability may also cause less baseline calcium expression, resulting in fewer chances

for photobleaching. Regardless, the ability to do long (60+ min) recording experiments

provide another compelling reason to use rats in calcium imaging experiments.

During our recording sessions, we saw clear evidence that calcium events were

strongly correlated with periods of movement (Figure 2c). This result is similar to

calcium imaging experiments done in mice by Zhou et al., in which more calcium events

were seen during movement in an environment where food reward could be obtained

than during quiet wakefulness in a neutral box [8]. In our experiment, periods of quiet

wake were in the same environment (the track) as periods of movement, which shows

that changes in calcium event frequency are related to movement and not to contextual

associations. Zhou et al. posit that the high amount of calcium activity state-dependent

during movement, but not during sharp wave ripples in mouse, implicate calcium events

in theta dependent processes, such as memory encoding and planning [89-95]. Further

work will be needed to determine if, in rat, frequent calcium events are also unique to

movement, as opposed to sharp wave ripples.

As previously mentioned, similarities between rat and human physiology, as well

as the increased capacity for complex tasks of learning and memory [55-60], position

rats as a well-suited model organism for studying hippocampal function, with several

advantages over the mouse. Rats are advantageous over mice in that, for many

diseases and disorders, including those with substantial hippocampal involvement, their

physiology and behavior more closely mirrors that of humans [63]. For instance, 5-HT

serotonin receptors, which are prominent in the hippocampus and are implicated in a

variety of disorders (including mood and anxiety disorders, psychiatric disorders such as

schizophrenia, addiction and ADHD), are distributed in rats, but not in mice, in a similar

pattern as seen in humans [63, 96]. Hippocampal neurogenesis, which is believed to

play an important role in learning, memory, and the treatment of depression, is believed

to follow a similar pattern in rats and humans, and is less pronounced in mice [97]. In

addition, both pharmacologic and behavioral responses to addiction are more similar

between rats and humans than mice and humans [63, 98, 99].

Having the ability to do calcium imaging in rat CA1 will allow increased research

into these disorders, as well as basic research in learning and memory. Because rats

are substantially larger than mice, they can support larger lens implants and cameras,

which may lead to the ability to record from thousands of hippocampal cells in a single

session during complex tasks. The ability to record a large number of cells across days

will allow research into cell firing changes in the hippocampus, which will provide insight

into more advanced learning and disease mechanisms that are difficult to understand

when only investigating mice. Additionally, the uniquely high percentage of recorded

place cells (>80% of recorded cells) makes CA1 calcium imaging in rats an ideal

method by which to study spatial navigation and learning over days, including

mechanisms behind memory consolidation, spatial remapping, and planning over time.

Materials and Methods

All procedures were performed within Northwestern Institutional Animal Care and Use Committee and NIH guidelines. Five male Fischer 344 x Brown Norway rats (275g to 325g) were sourced from National Institute on Aging colony at Charles River Laboratories, injected with AAV9-GCaMP7c, implanted with a 2mm GRIN lens and run on linear maze (Figure S2). Animals were individually housed in an animal facility with a 12h light dark cycle.

METHOD DETAILS

GCaMP7s Injection and Lens Implantation

Rats were anesthetized with isoflurane (induction 4%, maintenance 1-2%). Following anesthesia, ear bars, and skull leveling, the skull was cleaned and a craniotomy site was marked above CA1 (stereotaxic coordinates Bregma AP -4.00mm, ML 3.00mm). The skull was scored around the craniotomy site using a razor blade to provide texture for future application of epoxy. A craniotomy was made with a 2mm trephine around the marked site and dura was removed with a bent 30 gage needle. A Hamilton syringe (Gastight® 1700 Series Syringes, Hamilton, Model 1705 Small RN Syringe, Volume=50 μL, Point Style=3, Gauge=22s, Needle Length=50.8 mm) loaded with AAV9-GCaMP7c (obtained from Viogene Biosciences, packaged AAV9 of pGP-AAV-syn-jGCaMP7c-WPRE, titer $1.73*10^{13}$GC/mL) was lowered straight down to about 0.2mm past the CA1 cell layer (approximate coordinates Bregma AP -4.00mm, ML 3mm, DV 2.95mm relative to skull)—this was often slightly off-center of the craniotomy. 0.6uL of virus were injected over 12 minutes using an automated stereotaxic Injector.

The syringe was then raised 0.2mm (Bregma AP -4.00mm, ML 3mm, DV 2.75mm) and another 0.6uL of virus were injected over 12 minutes (Figure 1a). The syringe was then left for 10 minutes before slowly raising to the skull. Because the previous injection was often slightly off center of the craniotomy, the process of lowering the syringe and doing two injections was repeated so that the two syringe "lowering points" were evenly spread throughout the craniotomy site. There were thus a total of

four injections of 0.6uL each (2.4uL total): two injections per each brain entrance site,

each at different depths (Figure 1a).

Following injection, four skull screws were screwed into the skull: one rostral to

bregma above the same hemisphere as the craniotomy, two between bregma and

lambda on the hemisphere opposite the craniotomy, and one behind lambda on the

hemisphere with the craniotomy. Under a dissecting microscope, tissue from the

craniotomy was then aspirated using a vacuum pump and 25 gage needle that had

been previously blunted using a Dremel drill. Tissue was quickly aspirated until the start

of horizontal striations could be seen in the craniotomy hole. Care was then taken to

"level" the hole and make sure aspiration was at an even depth. The horizontal

striations of the corpus callosum were then carefully and slowly aspirated until the

vertical striations of the alveus were visible in the entirety of the craniotomy hole (Figure

1b). If the aspiration hole was bleeding, a small twisted piece of gel foam was inserted

into the hole for approximately five minutes and then removed.

Notes differentiating this method from mouse Ca2+ imaging: We found the best

AAV for expression in rats was AAV9. Because of rat's lower seizure threshold and,

anecdotally noted, lower GCaMP expression than in mice, four injection sites appear

both necessary and sufficient for ideal GCaMP expression. The injection volumes are

much larger than those used in mice [14, 15], which is made possible due to the larger

hippocampal area and the technique of injecting below the CA1 cell layer, instead of on

it. Unlike in mice, where just the cortex or most dorsal part of the corpus callosum is

aspirated [15, 100], it is important that the entire corpus callosum be completely

aspirated in order to get very close to the alveus.

A 2mm GRIN lens (obtained from Go!Foton, CLH lens, 2.00mm diameter, 0.448

pitch, working distance 0.30mm, 550nm wavelength) was then held on the end of a

pipette using suction from the vacuum pump. The pipette was placed in the stereotaxic

instrument which was zero-ed with the bottom of the lens on bregma. The lens was then

moved to the center of the craniotomy and lowered to a depth of approximately 3mm

from skull surface. The lens was then fixed in place with UV curing epoxy and the

vacuum pump was turned off and the pipette holding the lens removed. The epoxied

lens was then secured to the skull screws using dental acrylic. Animals were then given

buprenorphine (0.05mg/kg) and 20mL saline, taken off anesthesia, and allowed to

recover in a clean cage place upon a heat pad.

Notes differentiating this method from mouse Ca2+ imaging: A much larger lens

(2mm diameter as opposed to 0.5mm or 1.0mm diameter) lens can be used in rats due

to their larger size. Of high importance: in order to be as close to the CA1 layer as

possible, the lens is inserted further down (about .5mm) lower than the cell layer. This

serves, presumably, to compress the alveus to allow for in focus imaging of the CA1.

We had zero visualization of cells unless the lens was deeply inserted. Additionally,

unlike in mouse implantation surgeries, did not have success using meloxicam after

surgery as it appeared to cause excessive bleeding around the lens.

Four to six weeks after surgery, animals were again anesthetized with isoflurane

and checked for cells expressing GCaMP. If expression was seen, the animal's lens

was base-plated. The baseplate was attached using UV-curing epoxy and dental

acrylic.

Notes differentiating this method from mouse Ca2+ imaging: During base plating,

is gets progressively harder to see cells the longer the animal is under anesthesia.

Additionally, while a few cells will appear to be bright circles in the captured image, cells

do not appear to be flashing at all while the rat is under isoflorane, even after a tail pinch

or loud noise (techniques often used to incite a calcium during mouse base plating). We

recommend focusing on the bright, not flashing cells while base plating. The entire field

of view should become progressively brighter as the animal wakes up. We recommend

not imaging for experiments within 24 hours of isoflurane exposure and base plating.

Behavioral Training

Prior to training, animals were food deprived to 85% body weight and maintained

at this weight throughout the experiment. Implanted animals were trained on a 2.64m

Med Associates linear track with pellet dispensers at either end. Animals learned to run

back and forth between the sides of the track and received a grain pellet at alternate

ends. Each pellet received was considered a trial. The fasted animals were able to

achieve over 120 trials in 30 minutes. (Animal were fed after daily training to maintain

weight.) Six sessions were selected for analysis for each animal, with first and last

recorded sessions automatically included. When not being run, animals were housed in

individual cages with a 12hr light- 12hr dark light cycle.

398   Calcium Imaging

One photon calcium imaging was done using UCLA V4 Miniscopes [101, 102],

assembled with standard lens configurations (two 3mm diameter, 6mm FL achromat

lens used in the objective module and one 4 mm diameter, 10mm FL achromat lens

used in the emission module). We tested Ca2+ recording when one of the objective

module lenses was replaced by a 3mm diameter, 9mm FL achromat lens, but no

improvement in cell visualization was noted. Gain was set at high and Miniscope LED

intensity varied between 10 and 38.

Notes differentiating this method from mouse Ca2+ imaging: Base fluorescence

appears lower in rats as compared to mice. While we have been advised that, using

Miniscopes, gain can often be on "low" for mouse imaging, we found it necessary to

have gain on "high" to visualize the highest number of cells.

411   GCaMP Visualization in Slice

412   Following conclusion of the study, animals were anesthetized and perfused with

413   PBS then 10% formalin. The brain was then sliced into 60 micron coronal sections,

414   mounted, let dry, and coverslipped with Vectashield. Sections were then imaged with a

415   confocal microscope.

418   *QUANTIFICATION AND STATISTICAL ANALYSIS*

419   Means are expressed as mean +- the standard deviation. All analysis code is available

420   at https://github.com/hsw28/ca_imaging.

422   Position and speed analysis

Position was sampled by an overhead camera at 30Hz. Position tracking was

done post-recording using Bonsai [103]. Speed was determined by taking the

hypotenuse of the coordinates of the point one before and one after the time point of

interest. Speed was then smoothed using a Gaussian kernel of 1s and speed was

converted from pixels/s to cm/s.

Video pre-processing

Videos were recorded with Miniscope software at 15 frames/second and 752 by

480 pixels. All video processing was done using the Matlab-based CIATAH software

[76, 77] and processing files are available at https://github.com/hsw28/ca_imaging/ .

Videos were down sampled in space (3x using bi-linear interpolation) and time (2x). The

video was normalized by subtracting the mean value of each frame from the frame.

Each frame was then spatially filtered (normalized) using a bandpass FFT filter (70-

100cycles/pixel). The video was then motion corrected to a chosen reference frame

using TurboReg [104]. Videos were then converted to relative florescence (dF/F_0),

where F_0 was the mean over the entire video.

Cell identification

Cells were first automatically identified using the Matlab-based CIATAH software

[76, 77], using the CNMF-E method [75]. Processing files are available at

https://github.com/hsw28/ca_imaging/ . Images were filtered with a gaussian kernel of

width 3 pixels. Neuron diameter was set at a pixel size of 13. The threshold for merging

neurons was set at a calcium trace correlation of 0.65, and neurons were merged if their

distances were smaller than 2 pixels and they had highly correlated spatial shapes

(correlation>0.8) and small temporal correlations (correlation <0.4).

All cells identified using CNMF-E were then scored as neurons or not by a

human scorer. Scoring was also done within the Matlab-based CIATAH software [76,

77] in a Matlab GUI. Human scoring was done while visualizing an activity trace,

average Ca2+ waveform, a cropped movie montage of the candidate cell's Ca2+

events, and a maximum projection of all cells on which the candidate cell was

highlighted. The relative fluorescence ($\Delta F/F_0$) local maxima of each identified cell were

considered calcium event times.

Place cell identification

Place cells were identified using two criteria. First, cells must have a mean firing

rate greater than 0.01Hz during periods of movement (>12cm/s) [83, 105]. Second, the

mutual information (MI) computed in either direction of travel must be greater than 95%

of MI scores computed 500 times from shuffled cells [78]. To compute MI, the track was

divided lengthwise into 4cm bins the firing rate of each cell and the occupancy of the

animal were found for each bin. Rate and occupancy were smoothed with a 2cm

Gaussian kernel. Mutual information was computed during periods of movement in each

direction of travel as follows [78, 83, 106]:

$P_s = calcium\ event\ probability\ in\ each\ bin$

$P_o = occupancy\ probability\ at\ each\ bin$

$p = \dfrac{P_s}{P_o}$

$M_s = \sum P_s$

$M_o = \sum P_o$

$$MI = \sum p * \log_2\left(\frac{p}{M_s * M_o}\right)$$

Position decoding

Position was decoded using all recorded cells in a session and was decoded for

times when the rat was moving (velocity>12cm/s). The track was split in 10cm

segments and the time window of decoding was 0.5 seconds with non-overlapping time

bins. Decoding was done using Bayesian decoding [79].

$x = positions, f_i(x) = average\ firing\ rate\ per\ second\ for\ cell\ N\ at\ position\ X$

$n = calcium\ events\ per\ cell, n_i = number\ of\ events\ in\ time\ window$

$N = total\ number\ of\ cells$

$t = time\ window\ of\ decoding$

$C(t,n) = normalization\ factor\ for\ probabilities\ to\ equal\ 1$

$P(x|n) = odds\ of\ being\ at\ position\ x\ giving\ calcium\ events\ n$

484
$$P(x|n) = C(t,n)\left(\prod_{i=1}^{N} f_i(x)^{n_i}\right)\exp\left(-t\sum_{i=1}^{N} f_i(x)\right)$$

Accuracy decoding was determined by randomly shuffling firing per position bins

and re-decoding. The difference between actual and decoded position was then

determined per time bin and compared to decoded shuffled data.

Validation across days and within session

Validation was done using the Matlab-based CIATAH software [76, 77]. Videos

underwent 3 rounds of registration comprised of Turboreg image rotation [104]. An

image binarization threshold of 40% of the images' maximum value was used to remove

background noise and axons and dendrites. A distance threshold of a maximum of 5

pixels was used to match cells across sessions, with a minimum 2-D correlation

coefficient of 0.5. All sessions were aligned to the last recording session.

Photobleaching analysis was performed by computing the mean brightness for

each identified cell for a five-minute period towards the beginning of the session and

comparing it to the brightness for a five-minute period towards the end of the session.

(Mean brightness was the mean $\Delta F/F_0$ for each cell over the time period). The sampling

periods were separated from session start and end by five minutes to avoid behavioral

correlates related to starting and finishing the task period (Figure 5b). Results from

beginning to end of session were compared with a paired T-test. To normalize

brightness by the number of calcium events that occurred in each five-minute time

period, the number of events per cell were counted for the period. The mean

fluorescence for the period for that cell was then divided by the mean number of events.

Figure captions

Figure 1: GCaMP is expressed in the CA1 cell layer of rat hippocampus

  a. Diagram showing injection locations of AAV9-Syn-GCaMP7c, right below the CA1 cell layer. There are four total injection sites, with 0.6uL injected at each site. (CC = corpus callosum, DG = dentate gyrus)

  b. Diagram showing level of aspiration in brain tissue. The entire corpus callosum is aspirated until the vertical striations of the alveus are visible.

  c. GCaMP expression in the hippocampus, taken with a confocal microscope. The CA1 cell layer is labeled.

Figure 2: Hundreds of cells can be captured in a single Ca2+ imaging session

  a. Examples of data from two (top and bottom rows) recording sessions. First column: example image from pre-processing. Second column: image post-processing (see materials and methods). Third column: all cell extractions, prior to manual cell sorting. Fourth column: post processing image overlayed with cell outlines identified after manual sorting.

  b. Example traces from 15 cells taken during 124 seconds of calcium imaging.

  c. Raster plot of cell firing during linear track running. Top: the animal's speed. Bottom: raster of peak calcium transients of 352 identified cells.

  d. Number of cells imaged per session for each rat. Dotted red line indicates the average for each animal.

Figure 3: A high percentage of place cells are recorded on the linear track using Ca2+ imaging

  a. Graph showing the percentage of place cells (out of total recorded cells with a firing rate >0.1Hz during movement) per session for each rat. Dotted red line indicates the average for each animal.

  b. Nine example place cells. A firing rate map in both directions of travel is provided for each cell, with the top map corresponding with running to the

right, and the bottom map corresponding with running to the left. Note the

colorbar scales may be different for each direction of travel. The difference in

mutual information scores between the cell and the top 95% of shuffled data

(actual MI-shuffled MI) is indicated in the left corner. If the actual MI was

greater than 95% of shuffled data, the number is printed in bold white.

c.   Distribution of directional mutual information scores after shuffling cells seen

in figure 3b. Event data was shuffled 500 times and the top 95% of mutual

information scores were determined from shuffled data (red dotted line). If the

actual mutual information score (green line) was greater than the upper 95%

MI score of shuffled data, the cell was considered to be a place cell.

Figure 4: Place cells are sufficient to decode the animal's position

a.   Four decoding examples as the animal traverses the track. Black line

indicates the animal's actual position, with the stars indicated sample point.

The colored circles connected by the red line indicate decoded position. The

color of the circles indicates decoding certainty.

b.   Same as A but decoding is using shuffled units. Decoding is much less

accurate using shuffled data.

Figure 5: Individual cells can be maintained across session and within session.

a.   Cells can be maintained across sessions weeks apart. Examples from two

animals of data taken on the first and final days of recording. Cells are color-

matched from the first to last day. Cells in gray represent cells with no

matches between the two days. Of note, these recordings were prioritized for

cell number as proof of concept rather than maintaining the same landmarks

(such as blood vessels) across days.

b.   Top: method for comparing calcium signal brightness across session. The

mean brightness for each identified cell was computed for a five minute

period towards the beginning of the session and compared to the mean

brightness for a five minute period towards the end of the session. The

sampling periods were separated from session start and end by five minutes

to avoid psychological or behavioral correlates related to starting and finishing

the task period. Bottom: Histogram of change in fluorescence from starting to

ending, for all 5751 recorded cells (mean change = .008, median change = -

$2.01*10^{-4}$), There was no consistent change in fluorescence when examining

at the cell level. Examining by groups of cells in each session, in 22 out of the

24 sessions, the end mean fluorescence of each cell was not less than the

starting mean fluorescence (one-tailed paired t-test, $p>0.05$), and for 6 of

these sessions, the end mean fluorescence was actually greater than the

starting fluorescence (one-tailed paired t-test, $p<0.05$). Two sessions (8% of

sessions) showed a decrease in fluorescence (one-tailed paired t-test,

$t(368)=2.47$, $p<0.01$, and $t(231)=1.98$, $p<0.05$). For ease of viewing, outliers

were removed from visualization (but not analysis).

c. To control for the number of calcium events, we normalized the fluorescence

in each five-minute period by the number of events in the period and then

found the change in fluorescence from beginning to end of session (mean

change = $-1.77*10^{-4}$, median change = $-5.7*10^{-8}$). Even after normalization,

still only two sessions (although a different two) showed a decrease in

fluorescence from the start period to the end period (one-tailed paired t-test,

$t(226)=1.7$, $p=0.04$, and $t(349)=2.1$, $p=0.02$), while three sessions showed an

increase (one-tailed paired t-test, $t(123)=-1.9$; $p=0.03$, $t(173)=-3.0$, $p<0.005$;

$t(166)=-4.0$, $p<0.00005$). There was therefore no consistent loss of

fluorescence within each session. For ease of viewing, outliers were removed

from visualization (but not analysis).

Figure S1: Animals ran on a linear track.

a. Diagram of linear track. The animal would run back and forth between the two

sides to receive a pellet reward

b. Trace of animal position over one session

Figure S2: Percentage of cells that are place cells without using a firing rate cutoff. Percent is indicated for each day of recording, by animal. Red dotted line indicates the average for the animal.

Acknowledgements

We would like to thank all members of the Disterhoft lab, especially Matthew Oh, Craig Weiss, and Kent Park. We would also like to thank the Miniscope team, in particular Daniel Aharoni and Federico Sangiuliano Jimka. We additionally thank Biafra Ahanonu for assistance with CIATAH, Amy Christensen for providing advice during imaging setup, David Wirtshafter for discussion and editing help, and Drew Ames for consultation on figures.

Funding Disclosure

This work was supported by an NIA T32 (T32-AG020506/AG/NIA), an NIA R37 (R37-AG008796/AG/NIA) and an NINDS R01 (R01 NS113804/NS/NINDS).

Competing Interests

The authors declare no competing interests.

Related Manuscripts

Authors do not have a related or duplicate manuscript under consideration (or accepted) for publication elsewhere.

Citations

1.   Grienberger C, Konnerth A. Imaging calcium in neurons. Neuron. 2012;73(5):862-85. doi: 10.1016/j.neuron.2012.02.011. PubMed PMID: 22405199.

2.   Aharoni D, Hoogland TM. Circuit Investigations With Open-Source Miniaturized Microscopes: Past, Present and Future. Frontiers in cellular neuroscience. 2019;13:141. doi: 10.3389/fncel.2019.00141. PubMed PMID: 31024265; PubMed Central PMCID: PMCPMC6461004.

3.   Aharoni D, Khakh BS, Silva AJ, Golshani P. All the light that we can see: a new era in miniaturized microscopy. Nat Methods. 2019;16(1):11-3. doi: 10.1038/s41592-018-0266-x. PubMed PMID: 30573833.

4.   Rickgauer JP, Deisseroth K, Tank DW. Simultaneous cellular-resolution optical perturbation and imaging of place cell firing fields. Nat Neurosci. 2014;17(12):1816-24. doi: 10.1038/nn.3866. PubMed PMID: 25402854; PubMed Central PMCID: PMCPMC4459599.

5.   Resendez SL, Jennings JH, Ung RL, Namboodiri VM, Zhou ZC, Otis JM, et al. Visualization of cortical, subcortical and deep brain neural circuit dynamics during naturalistic mammalian behavior with head-mounted microscopes and chronically implanted lenses. Nature protocols. 2016;11(3):566-97. Epub 2016/02/26. doi: 10.1038/nprot.2016.021. PubMed PMID: 26914316; PubMed Central PMCID: PMCPMC5239057.

6.   Grienberger C, Konnerth A. Imaging calcium in neurons. Neuron. 2012;73(5):862-85.

7.   Hamel EJ, Grewe BF, Parker JG, Schnitzer MJ. Cellular level brain imaging in behaving mammals: an engineering approach. Neuron. 2015;86(1):140-59. Epub 2015/04/10. doi: 10.1016/j.neuron.2015.03.055. PubMed PMID: 25856491; PubMed Central PMCID: PMCPMC5758309.

8.   Zhou H, Neville KR, Goldstein N, Kabu S, Kausar N, Ye R, et al. Cholinergic modulation of hippocampal calcium activity across the sleep-wake cycle. eLife. 2019;8. Epub 2019/03/08. doi: 10.7554/eLife.39777. PubMed PMID: 30843520; PubMed Central PMCID: PMCPMC6435325.

9.   Sheintuch L, Rubin A, Brande-Eilat N, Geva N, Sadeh N, Pinchasof O, et al. Tracking the Same Neurons across Multiple Days in Ca(2+) Imaging Data. Cell reports. 2017;21(4):1102-15. doi: 10.1016/j.celrep.2017.10.013. PubMed PMID: 29069591; PubMed Central PMCID: PMCPMC5670033.

10.   Hainmueller T, Bartos M. Parallel emergence of stable and dynamic memory engrams in the hippocampus. Nature. 2018;558(7709):292-6. doi: 10.1038/s41586-018-0191-2. PubMed PMID: 29875406.

11.   Dombeck DA, Harvey CD, Tian L, Looger LL, Tank DW. Functional imaging of hippocampal place cells at cellular resolution during virtual navigation. Nat Neurosci. 2010;13(11):1433-40. doi: 10.1038/nn.2648. PubMed PMID: 20890294; PubMed Central PMCID: PMCPMC2967725.

12.   Kinsky NR, Mau W, Sullivan DW, Levy SJ, Ruesch EA, Hasselmo ME. Trajectory-modulated hippocampal neurons persist throughout memory-guided navigation. Nature communications. 2020;11(1):2443. Epub 2020/05/18. doi: 10.1038/s41467-020-16226-4. PubMed PMID: 32415083; PubMed Central PMCID: PMCPMC7229120.

13.   Sheintuch L, Geva N, Baumer H, Rechavi Y, Rubin A, Ziv Y. Multiple Maps of the Same
Spatial Context Can Stably Coexist in the Mouse Hippocampus. Current biology : CB.
2020;30(8):1467-76 e6. Epub 2020/03/30. doi: 10.1016/j.cub.2020.02.018. PubMed PMID:
32220328.
14.   Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, et al. Long-term dynamics of
CA1 hippocampal place codes. Nat Neurosci. 2013;16(3):264-6. Epub 2013/02/12. doi:
10.1038/nn.3329. PubMed PMID: 23396101; PubMed Central PMCID: PMCPMC3784308.
15.   Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, et al. A shared neural ensemble
links distinct contextual memories encoded close in time. Nature. 2016;534(7605):115-8. Epub
2016/06/03. doi: 10.1038/nature17955. PubMed PMID: 27251287; PubMed Central PMCID:
PMCPMC5063500.
16.   Jimenez JC, Berry JE, Lim SC, Ong SK, Kheirbek MA, Hen R. Contextual fear memory
retrieval by correlated ensembles of ventral CA1 neurons. Nature communications.
2020;11(1):3492. Epub 2020/07/15. doi: 10.1038/s41467-020-17270-w. PubMed PMID:
32661319; PubMed Central PMCID: PMCPMC7359370.
17.   Zhang X, Li B. Population coding of valence in the basolateral amygdala. Nature
communications. 2018;9(1):5195. Epub 2018/12/07. doi: 10.1038/s41467-018-07679-9.
PubMed PMID: 30518754; PubMed Central PMCID: PMCPMC6281657.
18.   Shuman T, Aharoni D, Cai DJ, Lee CR, Chavlis S, Page-Harley L, et al. Breakdown of spatial
coding and interneuron synchronization in epileptic mice. Nat Neurosci. 2020;23(2):229-38.
Epub 2020/01/08. doi: 10.1038/s41593-019-0559-0. PubMed PMID: 31907437; PubMed
Central PMCID: PMCPMC7259114.
19.   Shin JH, Song M, Paik SB, Jung MW. Spatial organization of functional clusters
representing reward and movement information in the striatal direct and indirect pathways.
Proc Natl Acad Sci U S A. 2020;117(43):27004-15. Epub 2020/10/16. doi:
10.1073/pnas.2010361117. PubMed PMID: 33055217; PubMed Central PMCID:
PMCPMC7604453.
20.   Wang X, Liu Y, Li X, Zhang Z, Yang H, Zhang Y, et al. Deconstruction of Corticospinal
Circuits for Goal-Directed Motor Skills. Cell. 2017;171(2):440-55 e14. Epub 2017/09/26. doi:
10.1016/j.cell.2017.08.014. PubMed PMID: 28942925; PubMed Central PMCID:
PMCPMC5679421.
21.   Klaus A, Martins GJ, Paixao VB, Zhou P, Paninski L, Costa RM. The Spatiotemporal
Organization of the Striatum Encodes Action Space. Neuron. 2017;95(5):1171-80 e7. Epub
2017/09/01. doi: 10.1016/j.neuron.2017.08.015. PubMed PMID: 28858619; PubMed Central
PMCID: PMCPMC5584673.
22.   Guo W, Zhang JJ, Newman JP, Wilson MA. Latent learning drives sleep-dependent
plasticity in distinct CA1 subpopulations. bioRxiv. 2020:2020.02.27.967794. doi:
10.1101/2020.02.27.967794.
23.   O'Keefe J. Place units in the hippocampus of the freely moving rat. Exp Neurol.
1976;51(1):78-109. Epub 1976/04/01. PubMed PMID: 1261644.
24.   Wilson MA, McNaughton BL. Dynamics of the hippocampal ensemble code for space.
Science. 1993;261(5124):1055-8. Epub 1993/08/20. PubMed PMID: 8351520.

25.   Leutgeb S, Leutgeb JK, Treves A, Moser MB, Moser EI. Distinct ensemble codes in
hippocampal areas CA3 and CA1. Science. 2004;305(5688):1295-8. Epub 2004/07/24. doi:
10.1126/science.1100265
1100265 [pii]. PubMed PMID: 15272123.
26.   McEchron MD, Disterhoft JF. Hippocampal encoding of non-spatial trace conditioning.
Hippocampus. 1999;9(4):385-96. doi: 10.1002/(SICI)1098-1063(1999)9:4<385::AID-
HIPO5>3.0.CO;2-K. PubMed PMID: 10495020.
27.   Schimanski LA, Lipa P, Barnes CA. Tracking the course of hippocampal representations
during learning: when is the map required? J Neurosci. 2013;33(7):3094-106. doi:
10.1523/JNEUROSCI.1348-12.2013. PubMed PMID: 23407964; PubMed Central PMCID:
PMCPMC3700421.
28.   Stefanini F, Kushnir L, Jimenez JC, Jennings JH, Woods NI, Stuber GD, et al. A Distributed
Neural Code in the Dentate Gyrus and in CA1. Neuron. 2020;107(4):703-16 e4. Epub
2020/06/11. doi: 10.1016/j.neuron.2020.05.022. PubMed PMID: 32521223; PubMed Central
PMCID: PMCPMC7442694.
29.   Scoville WB, Milner B. Loss of recent memory after bilateral hippocampal lesions.
Journal of neurology, neurosurgery, and psychiatry. 1957;20(1):11-21. Epub 1957/02/01.
PubMed PMID: 13406589; PubMed Central PMCID: PMC497229.
30.   Squire LR. Memory and the hippocampus: a synthesis from findings with rats, monkeys,
and humans. Psychol Rev. 1992;99(2):195-231. Epub 1992/04/01. PubMed PMID: 1594723.
31.   Hasselmo ME, McClelland JL. Neural models of memory. Curr Opin Neurobiol.
1999;9(2):184-8. Epub 1999/05/14. doi: S0959-4388(99)80025-7 [pii]. PubMed PMID:
10322183.
32.   Ferbinteanu J, Shapiro ML. Prospective and retrospective memory coding in the
hippocampus. Neuron. 2003;40(6):1227-39. Epub 2003/12/23. doi: S0896627303007529 [pii].
PubMed PMID: 14687555.
33.   Smith DM, Mizumori SJ. Hippocampal place cells, context, and episodic memory.
Hippocampus. 2006;16(9):716-29. Epub 2006/08/10. doi: 10.1002/hipo.20208. PubMed PMID:
16897724.
34.   Vann SD. Dismantling the Papez circuit for memory in rats. eLife. 2013;2:e00736. Epub
2013/06/28. doi: 10.7554/eLife.00736. PubMed PMID: 23805381; PubMed Central PMCID:
PMC3691571.
35.   Josselyn SA, Tonegawa S. Memory engrams: Recalling the past and imagining the future.
Science. 2020;367(6473). Epub 2020/01/04. doi: 10.1126/science.aaw4325. PubMed PMID:
31896692.
36.   Eichenbaum H, Dudchenko P, Wood E, Shapiro M, Tanila H. The hippocampus, memory,
and place cells: is it spatial memory or a memory space? Neuron. 1999;23(2):209-26. Epub
1999/07/10. doi: S0896-6273(00)80773-4 [pii]. PubMed PMID: 10399928.
37.   Ferbinteanu J, Kennedy PJ, Shapiro ML. Episodic memory—from brain to mind.
Hippocampus. 2006;16(9):691-703.
38.   McNaughton BL, Battaglia FP, Jensen O, Moser EI, Moser MB. Path integration and the
neural basis of the 'cognitive map'. Nat Rev Neurosci. 2006;7(8):663-78. Epub 2006/07/22. doi:
nrn1932 [pii]
10.1038/nrn1932. PubMed PMID: 16858394.

39.   Foster DJ, Knierim JJ. Sequence learning and the role of the hippocampus in rodent
navigation. Curr Opin Neurobiol. 2012;22(2):294-300. Epub 2012/01/10. doi:
10.1016/j.conb.2011.12.005
S0959-4388(11)00223-6 [pii]. PubMed PMID: 22226994.
40.   Foster DJ, Morris RG, Dayan P. A model of hippocampally dependent navigation, using
the temporal difference learning rule. Hippocampus. 2000;10(1):1-16. Epub 2000/03/08. doi:
10.1002/(SICI)1098-1063(2000)10:1<1::AID-HIPO1>3.0.CO;2-1. PubMed PMID: 10706212.
41.   Avigan PD, Cammack K, Shapiro ML. Flexible spatial learning requires both the dorsal
and ventral hippocampus and their functional interactions with the prefrontal cortex.
Hippocampus. 2020;30(7):733-44.
42.   Moser EI, Moser M-B, McNaughton BL. Spatial representation in the hippocampal
formation: a history. Nature neuroscience. 2017;20(11):1448-64.
43.   Rosenzweig ES, Redish AD, McNaughton BL, Barnes CA. Hippocampal map realignment
and spatial learning. Nature neuroscience. 2003;6(6):609-15.
44.   Moyer JR, Jr., Deyo RA, Disterhoft JF. Hippocampectomy disrupts trace eye-blink
conditioning in rabbits. Behav Neurosci. 1990;104(2):243-52. PubMed PMID: 2346619.
45.   Andrzejewski ME, Ryals C. Dissociable hippocampal and amygdalar D1-like receptor
contribution to discriminated Pavlovian conditioned approach learning. Behav Brain Res.
2016;299:111-21. doi: 10.1016/j.bbr.2015.11.034. PubMed PMID: 26632336; PubMed Central
PMCID: PMCPMC4866504.
46.   Ito R, Everitt BJ, Robbins TW. The hippocampus and appetitive Pavlovian conditioning:
effects of excitotoxic hippocampal lesions on conditioned locomotor activity and autoshaping.
Hippocampus. 2005;15(6):713-21. doi: 10.1002/hipo.20094. PubMed PMID: 15906393.
47.   Chan KH, Morell JR, Jarrard LE, Davidson TL. Reconsideration of the role of the
hippocampus in learned inhibition. Behav Brain Res. 2001;119(2):111-30. PubMed PMID:
11165328.
48.   Pritchard A, Hornung R, Kramer PR. Measuring calcium activity within individual neurons
within the rat thalamus. MethodsX. 2021;8:101273.
49.   Anner P, Passecker J, Klausberger T, Dorffner G. Ca2+ imaging of neurons in freely
moving rats with automatic post hoc histological identification. Journal of neuroscience
methods. 2020;341:108765.
50.   Scott BB, Thiberge SY, Guo C, Tervo DGR, Brody CD, Karpova AY, et al. Imaging cortical
dynamics in GCaMP transgenic rats with a head-mounted widefield macroscope. Neuron.
2018;100(5):1045-58. e5.
51.   Hart EE, Blair GJ, O'Dell TJ, Blair HT, Izquierdo A. Chemogenetic modulation and single-
photon calcium imaging in anterior cingulate cortex reveal a mechanism for effort-based
decisions. Journal of Neuroscience. 2020;40(29):5628-43.
52.   Swanson L. Brain maps: structure of the rat brain: Gulf Professional Publishing; 2004.
53.   Franklin KBJ, Paxinos G. Paxinos and Franklin's The mouse brain in stereotaxic
coordinates2013.
54.   Routh BN, Johnston D, Harris K, Chitwood RA. Anatomical and electrophysiological
comparison of CA1 pyramidal neurons of the rat and mouse. Journal of neurophysiology.
2009;102(4):2288-302.

55.　　Whishaw IQ. A comparison of rats and mice in a swimming pool place task and matching
to place task: some surprising differences. Physiol Behav. 1995;58(4):687-93. Epub 1995/10/01.
doi: 10.1016/0031-9384(95)00110-5. PubMed PMID: 8559777.

56.　　Frick KM, Stillner ET, Berger-Sweeney J. Mice are not little rats: species differences in a
one-day water maze task. Neuroreport. 2000;11(16):3461-5. Epub 2000/11/30. doi:
10.1097/00001756-200011090-00013. PubMed PMID: 11095500.

57.　　Rosenfeld CS, Ferguson SA. Barnes maze testing strategies with small and large rodent
models. J Vis Exp. 2014;(84):e51194. Epub 2014/03/19. doi: 10.3791/51194. PubMed PMID:
24637673; PubMed Central PMCID: PMCPMC4140524.

58.　　Cressant A, Besson M, Suarez S, Cormier A, Granon S. Spatial learning in Long-Evans
Hooded rats and C57BL/6J mice: different strategies for different performance. Behav Brain
Res. 2007;177(1):22-9. Epub 2006/12/13. doi: 10.1016/j.bbr.2006.11.010. PubMed PMID:
17157932.

59.　　Hok V, Poucet B, Duvelle É, Save É, Sargolini F. Spatial cognition in mice and rats:
similarities and differences in brain and behavior. Wiley Interdisciplinary Reviews: Cognitive
Science. 2016;7(6):406-21.

60.　　Whishaw IQ, Metz GA, Kolb B, Pellis SM. Accelerated nervous system development
contributes to behavioral efficiency in the laboratory mouse: a behavioral review and
theoretical proposal. Developmental Psychobiology: The Journal of the International Society for
Developmental Psychobiology. 2001;39(3):151-70.

61.　　Vengeliene V, Bilbao A, Spanagel R. The alcohol deprivation effect model for studying
relapse behavior: a comparison between rats and mice. Alcohol. 2014;48(3):313-20. Epub
2014/05/09. doi: 10.1016/j.alcohol.2014.03.002. PubMed PMID: 24811155.

62.　　Carter CS, Richardson A, Huffman DM, Austad S. Bring Back the Rat! J Gerontol A Biol Sci
Med Sci. 2020;75(3):405-15. Epub 2020/01/03. doi: 10.1093/gerona/glz298. PubMed PMID:
31894235; PubMed Central PMCID: PMCPMC7021637.

63.　　Ellenbroek B, Youn J. Rodent models in neuroscience research: is it a rat race? Dis Model
Mech. 2016;9(10):1079-87. Epub 2016/10/14. doi: 10.1242/dmm.026120. PubMed PMID:
27736744; PubMed Central PMCID: PMCPMC5087838.

64.　　Iannaccone PM, Jacob HJ. Rats! Dis Model Mech. 2009;2(5-6):206-10. Epub 2009/05/02.
doi: 10.1242/dmm.002733. PubMed PMID: 19407324; PubMed Central PMCID:
PMCPMC2675817.

65.　　Buerge T, Weiss T. Handling and Restraint. In: Hedrich HJ, Bullock G, editors. The
Laboratory Mouse (Handbook of Experimental Animals): Elsevier; 2004. p. 517-26.

66.　　Voigts J, Newman JP, Wilson MA, Harnett MT. An easy-to-assemble, robust, and
lightweight drive implant for chronic tetrode recordings in freely moving animals. J Neural Eng.
2020;17(2):026044. Epub 2020/02/20. doi: 10.1088/1741-2552/ab77f9. PubMed PMID:
32074511.

67.　　Voigts J, Siegle JH, Pritchett DL, Moore CI. The flexDrive: an ultra-light implant for optical
control and highly parallel chronic recording of neuronal ensembles in freely moving mice.
Front Syst Neurosci. 2013;7:8. Epub 2013/05/30. doi: 10.3389/fnsys.2013.00008. PubMed
PMID: 23717267; PubMed Central PMCID: PMCPMC3652307.

68.   Bolkvadze T, Pitkanen A. Development of post-traumatic epilepsy after controlled
cortical impact and lateral fluid-percussion-induced brain injury in the mouse. J Neurotrauma.
2012;29(5):789-812. Epub 2011/10/26. doi: 10.1089/neu.2011.1954. PubMed PMID: 22023672.
69.   Hunt RF, Scheff SW, Smith BN. Posttraumatic epilepsy after controlled cortical impact
injury in mice. Exp Neurol. 2009;215(2):243-52. Epub 2008/11/18. doi:
10.1016/j.expneurol.2008.10.005. PubMed PMID: 19013458; PubMed Central PMCID:
PMCPMC4694635.
70.   Hunt RF, Scheff SW, Smith BN. Regionally localized recurrent excitation in the dentate
gyrus of a cortical contusion model of posttraumatic epilepsy. J Neurophysiol.
2010;103(3):1490-500. Epub 2010/01/22. doi: 10.1152/jn.00957.2009. PubMed PMID:
20089815; PubMed Central PMCID: PMCPMC2887624.
71.   Steinmetz NA, Buetfering C, Lecoq J, Lee CR, Peters AJ, Jacobs EAK, et al. Aberrant
Cortical Activity in Multiple GCaMP6-Expressing Transgenic Mouse Lines. eNeuro. 2017;4(5).
Epub 2017/09/22. doi: 10.1523/ENEURO.0207-17.2017. PubMed PMID: 28932809; PubMed
Central PMCID: PMCPMC5604087.
72.   Yang Y, Liu N, He Y, Liu Y, Ge L, Zou L, et al. Improved calcium sensor GCaMP-X
overcomes the calcium channel perturbations induced by the calmodulin in GCaMP. Nature
communications. 2018;9(1):1504. Epub 2018/04/19. doi: 10.1038/s41467-018-03719-6.
PubMed PMID: 29666364; PubMed Central PMCID: PMCPMC5904127.
73.   Tian L, Hires SA, Mao T, Huber D, Chiappe ME, Chalasani SH, et al. Imaging neural
activity in worms, flies and mice with improved GCaMP calcium indicators. Nat Methods.
2009;6(12):875-81. Epub 2009/11/10. doi: 10.1038/nmeth.1398. PubMed PMID: 19898485;
PubMed Central PMCID: PMCPMC2858873.
74.   Nathanson JL, Yanagawa Y, Obata K, Callaway EM. Preferential labeling of inhibitory and
excitatory cortical neurons by endogenous tropism of adeno-associated virus and lentivirus
vectors. Neuroscience. 2009;161(2):441-50. Epub 2009/03/26. doi:
10.1016/j.neuroscience.2009.03.032. PubMed PMID: 19318117; PubMed Central PMCID:
PMCPMC2728494.
75.   Zhou P, Resendez SL, Rodriguez-Romaguera J, Jimenez JC, Neufeld SQ, Giovannucci A, et
al. Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data.
eLife. 2018;7. Epub 2018/02/23. doi: 10.7554/eLife.28728. PubMed PMID: 29469809; PubMed
Central PMCID: PMCPMC5871355.
76.   Ahanonu B. CIAtah: a software package for analyzing one- and two-photon calcium
imaging datasets. v1.0.0 ed: Zenodo; 2018.
77.   Corder G, Ahanonu B, Grewe BF, Wang D, Schnitzer MJ, Scherrer G. An amygdalar neural
ensemble that encodes the unpleasantness of pain. Science. 2019;363(6424):276-81. Epub
2019/01/19. doi: 10.1126/science.aap8586. PubMed PMID: 30655440; PubMed Central PMCID:
PMCPMC6450685.
78.   Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB. Hippocampal Place
Fields Maintain a Coherent and Flexible Map across Long Timescales. Current biology : CB.
2018;28(22):3578-88 e6. Epub 2018/11/06. doi: 10.1016/j.cub.2018.09.037. PubMed PMID:
30393037; PubMed Central PMCID: PMCPMC6331214.

79.   Zhang K, Ginzburg I, McNaughton BL, Sejnowski TJ. Interpreting neuronal population
activity by reconstruction: unified framework with application to hippocampal place cells. J
Neurophysiol. 1998;79(2):1017-44. doi: 10.1152/jn.1998.79.2.1017. PubMed PMID: 9463459.
80.   Li P, Geng X, Jiang H, Caccavano A, Vicini S, Wu JY. Measuring Sharp Waves and
Oscillatory Population Activity With the Genetically Encoded Calcium Indicator GCaMP6f.
Frontiers in cellular neuroscience. 2019;13:274. Epub 2019/07/06. doi:
10.3389/fncel.2019.00274. PubMed PMID: 31275115; PubMed Central PMCID:
PMCPMC6593119.
81.   Stefanini F, Kushnir L, Jimenez JC, Jennings JH, Woods NI, Stuber GD, et al. A distributed
neural code in the dentate gyrus and in CA1. Neuron. 2020;107(4):703-16. e4.
82.   Mou X, Cheng J, Yu YS, Kee SE, Ji D. Comparing mouse and rat hippocampal place cell
activities and firing sequences in the same environments. Frontiers in cellular neuroscience.
2018;12:332.
83.   Wirtshafter HS, Wilson MA. Differences in reward biased spatial representations in the
lateral septum and hippocampus. eLife. 2020;9. Epub 2020/05/27. doi: 10.7554/eLife.55252.
PubMed Central PMCID: PMCPMC7274787.
84.   Fenton AA, Kao H-Y, Neymotin SA, Olypher A, Vayntrub Y, Lytton WW, et al. Unmasking
the CA1 ensemble place code by exposures to small and large environments: more place cells
and multiple, irregularly arranged, and expanded place fields in the larger space. Journal of
Neuroscience. 2008;28(44):11250-62.
85.   Markus EJ, Barnes CA, McNaughton BL, Gladden VL, Skaggs WE. Spatial information
content and reliability of hippocampal CA1 neurons: effects of visual input. Hippocampus.
1994;4(4):410-21.
86.   Ego-Stengel V, Wilson MA. Spatial selectivity and theta phase precession in CA1
interneurons. Hippocampus. 2007;17(2):161-74. doi: 10.1002/hipo.20253. PubMed PMID:
17183531.
87.   Huang L, Ledochowitsch P, Knoblich U, Lecoq J, Murphy GJ, Reid RC, et al. Relationship
between simultaneously recorded spiking activity and fluorescence signal in GCaMP6 transgenic
mice. eLife. 2021;10:e51675.
88.   Delon A, Usson Y, Derouard J, Biben T, Souchier C. Continuous photobleaching in
vesicles and living cells: a measure of diffusion and compartmentation. Biophys J.
2006;90(7):2548-62. Epub 2006/01/24. doi: 10.1529/biophysj.105.069815. PubMed PMID:
16428281; PubMed Central PMCID: PMCPMC1403194.
89.   Buzsaki G. Two-stage model of memory trace formation: a role for "noisy" brain states.
Neuroscience. 1989;31(3):551-70. Epub 1989/01/01. doi: 0306-4522(89)90423-5 [pii]. PubMed
PMID: 2687720.
90.   Colgin LL. Mechanisms and functions of theta rhythms. Annu Rev Neurosci.
2013;36:295-312. doi: 10.1146/annurev-neuro-062012-170330. PubMed PMID: 23724998.
91.   Hasselmo ME. What is the function of hippocampal theta rhythm?--Linking behavioral
data to phasic properties of field potential and unit recording data. Hippocampus.
2005;15(7):936-49. doi: 10.1002/hipo.20116. PubMed PMID: 16158423.
92.   Hasselmo ME, Hay J, Ilyn M, Gorchetchnikov A. Neuromodulation, theta rhythm and rat
spatial navigation. Neural Netw. 2002;15(4-6):689-707. Epub 2002/10/10. doi: S0893-
6080(02)00057-6 [pii]. PubMed PMID: 12371520.

93.   Wikenheiser AM, Redish AD. Hippocampal theta sequences reflect current goals. Nat
Neurosci. 2015;18(2):289-94. Epub 2015/01/07. doi: 10.1038/nn.3909. PubMed PMID:
25559082; PubMed Central PMCID: PMC4428659.

94.   Wirtshafter HS, Wilson MA. Lateral septum as a nexus for mood, motivation, and
movement. Neurosci Biobehav Rev. 2021;126:544-59. Epub 2021/04/14. doi:
10.1016/j.neubiorev.2021.03.029. PubMed PMID: 33848512.

95.   Shirvalkar PR, Rapp PR, Shapiro ML. Bidirectional changes to hippocampal theta–gamma
comodulation predict memory for recent spatial episodes. Proceedings of the National
Academy of Sciences. 2010;107(15):7054-9.

96.   Hirst WD, Abrahamsen B, Blaney FE, Calver AR, Aloj L, Price GW, et al. Differences in the
central nervous system distribution and pharmacology of the mouse 5-hydroxytryptamine-6
receptor compared with rat and human receptors investigated by radioligand binding, site-
directed mutagenesis, and molecular modeling. Molecular pharmacology. 2003;64(6):1295-308.

97.   Snyder JS, Choe JS, Clifford MA, Jeurling SI, Hurley P, Brown A, et al. Adult-born
hippocampal neurons are more numerous, faster maturing, and more involved in behavior in
rats than in mice. Journal of Neuroscience. 2009;29(46):14484-95.

98.   Parker CC, Chen H, Flagel SB, Geurts AM, Richards JB, Robinson TE, et al. Rats are the
smart choice: rationale for a renewed focus on rats in behavioral genetics. Neuropharmacology.
2014;76:250-8.

99.   Jupp B, Caprioli D, Dalley JW. Highly impulsive rats: modelling an endophenotype to
determine the neurobiological, genetic and environmental mechanisms of addiction. Dis Model
Mech. 2013;6(2):302-11. Epub 2013/01/29. doi: 10.1242/dmm.010934. PubMed PMID:
23355644; PubMed Central PMCID: PMCPMC3597013.

100.   Schoenfeld G, Carta S, Rupprecht P, Ayaz A, Helmchen F. In vivo calcium imaging of CA3
pyramidal neuron populations in adult mouse hippocampus. eNeuro. 2021. Epub 2021/08/01.
doi: 10.1523/ENEURO.0023-21.2021. PubMed PMID: 34330817.

101.   Aharoni D, Hoogland TM. Circuit investigations with open-source miniaturized
microscopes: past, present and future. Frontiers in cellular neuroscience. 2019;13:141.

102.   Silva AJ. Miniaturized two-photon microscope: seeing clearer and deeper into the brain.
Light: Science & Applications. 2017;6(8):e17104-e.

103.   Lopes G, Bonacchi N, Frazao J, Neto JP, Atallah BV, Soares S, et al. Bonsai: an event-
based framework for processing and controlling data streams. Front Neuroinform. 2015;9:7.
Epub 2015/04/24. doi: 10.3389/fninf.2015.00007. PubMed PMID: 25904861; PubMed Central
PMCID: PMCPMC4389726.

104.   Thevenaz P, Ruttimann UE, Unser M. A pyramid approach to subpixel registration based
on intensity. IEEE Trans Image Process. 1998;7(1):27-41. Epub 2008/02/13. doi:
10.1109/83.650848. PubMed PMID: 18267377.

105.   Wirtshafter HS, Wilson MA. Locomotor and Hippocampal Processing Converge in the
Lateral Septum. Current biology : CB. 2019;29(19):3177-92. doi: 10.1016/j.cub.2019.07.089.
PubMed PMID: 31543450.

106.   Olypher AV, Lansky P, Muller RU, Fenton AA. Quantifying location-specific information in
the discharge of rat hippocampal place cells. J Neurosci Methods. 2003;127(2):123-35. Epub
2003/08/09. doi: 10.1016/s0165-0270(03)00123-7. PubMed PMID: 12906942.

Figure 1

A

CC

CA1

x x
x x

Alveus

Cell layer

DG

CA3

B

Aspiration

CC

CA1

Alveus

Cell layer

DG

CA3

C

100 μm

CA1
cell layer

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

A

Sites of pellet
dispensing

2.64m

B

| 0 | 50 | 100 | 150 | 200 | 250 |

cm

Figure S2