1 Similarities and differences of face and body perception in

2 the dog (*Canis familiaris*) and human brain

Magdalena Boch[1,2*], Isabella C. Wagner[1], Sabrina Karl[3], Ludwig Huber[3+] and Claus

Lamm[1,4+]

[1]Social, Cognitive and Affective Neuroscience (SCAN) Unit, Department of Cognition, Emotion, and

Methods in Psychology, Faculty of Psychology, University of Vienna, Vienna, Austria

[2]Department of Cognitive Biology, Faculty of Life Sciences, University of Vienna, Vienna, Austria

[3]Comparative Cognition, Messerli Research Institute, University of Veterinary Medicine Vienna,

Medical University of Vienna and University of Vienna, Vienna, Austria

[4]Vienna Cognitive Science Hub, University of Vienna, Austria

[+]These authors share senior authorship.

*Corresponding author:

Email: magdalena.boch@univie.ac.at

Abstract

Accurately recognizing other individuals is fundamental for successful social interactions. While the neural underpinnings of this skill have been studied extensively in humans, less is known about the evolutionary origins of the brain areas specialized for recognising faces or bodies. Studying dogs (*Canis familiaris*), a non-primate species with the ability to perceive faces and bodies similarly to humans, promises insights into how visuo-social perception has evolved in mammals. We investigated the neural correlates of face and body perception in dogs (*N* = 15) and humans (*N* = 40) using functional MRI. Combining uni- and multivariate analysis approaches, we identified activation levels and patterns that suggested potentially homologous occipito-temporal brain regions in both species responding to faces and bodies compared to inanimate objects. Crucially, only human brain regions showed activation differences between faces and bodies and partly responded more strongly to humans compared to dogs. Moreover, only dogs represented both faces and dog bodies in olfactory regions. Overall, these findings advance our understanding of the similarities and differences in brain function between two phylogenetically distant mammal species. This likely reflects differences in the perceptual systems these species rely on to recognize others, and provides important insights into the foundations of social cognition and behaviour.

1 Introduction

The ability to detect and recognize another individual is a socio-cognitive skill

crucial for successful social interactions and survival. The face and body of another

human or non-human animal convey a wealth of social information beyond their

identity, such as when facilitating the correct categorization of another individual's

emotional state [1–7]. Consequently, accurate perception of faces and bodies is an

important building block for social perception.

Therefore, the neural underpinnings of face and body perception have been

studied extensively in humans. Decades of research revealed a predominant role of

the occipito-temporal cortex for the perception of others with distinct, but adjacent

regions specialized for face and body perception as part of the so-called ventral visual

pathway [8–17]. While the ventrolateral visual cortex responds to animate compared

to inanimate stimuli more generally [18–20], the sensitivity in the occipital face and

extrastriate body area (lateral occipital cortex), as well as the fusiform face and fusiform

body area (inferior temporal cortex) appear increased for faces or bodies [10–13,15–

17,21–28]. Regarding the recognition of another individual's species-identity, previous

research suggests a preference for human faces compared to faces of other species

(such as dogs or horses) [29,30], potentially reflecting increased salience of

conspecifics over heterospecifics.

The evolutionary history of the neural underpinnings of face and body perception

has been mainly investigated by comparative research in non-human primates [21–

24,28,31], and this has revealed a homologous visual pathway [e.g. 8,32 for reviews].

However, apart from primate species, the neural bases of this socio-cognitive ability

and its potential origin in the mammalian brain remain largely unstudied. Dogs' (*Canis*

familiaris) ability to perceive and understand humans and conspecifics has already

been well-investigated using behavioural and eye-tracking measures and yielded

numerous comparable skills between dogs and humans. For example, dogs can not

only differentiate between faces [33,34], but even discriminate facial emotional cues of

humans and conspecifics [7,35,36]; and they demonstrate complex behaviours such

as visual perspective taking [37,38]. Thus, dogs constitute an excellent model to study

the evolution of visuo-social perception in mammals [39].

While first non-invasive canine neuroimaging studies [29,40–43] suggested an

involvement of the temporal cortex in face perception, they triggered a debate on

whether the occipito-temporal specialization for face perception in dogs matches that

of humans [29,40–43]. Previous work reported increased activation for faces compared

to objects [41,43], scenes [41], or a mono-coloured surface [40], but not compared to

other low-level visual controls (i.e., scrambled images [40,41] or the back of the head

[29]). Some studies lacked non-facial controls entirely [42]. Studies additionally

investigating responses to con- and heterospecifics were inconclusive ranging from

reports of increased responsiveness to dog stimuli in general (i.e., face and back of

the head) but no specialization for faces [29], over separate brain regions specialized

for human or dog faces [42], to no species- but general face-preference [41]. Further,

no previous work investigated the neural bases of body perception. This is a major

limitation, as bodies play an important role for social perception in general, but in

particular for dogs - as indicated e.g., by dogs' high responsiveness to emotional or

ostensive-referential bodily cues, such as tail-wagging [44] or human gestures [44–

46]. It is particularly noteworthy in this respect, that dogs even outperform humans'

closest living relatives, chimpanzees, in utilizing human ostensive cues [47].

Thus, the findings so far do not allow for a distinction between face and body or

general animacy perception and are inconclusive regarding the influence of the other

individual's species-identity. The present comparative neuroimaging study aimed to

resolve this debate. We performed the, to our knowledge, first systematic comparison

of the neural bases of face, body and inanimate object perception in dogs and humans,

in order to gain insights into how social perception has evolved in two phylogenetically

distant mammalian species. [39,48].

2   Results

Fifteen awake and unrestrained pet dogs (**Figure 1A**) and forty human

participants underwent functional magnetic resonance imaging (MRI) while viewing

images of human and dog bodies, faces, inanimate objects (e.g., different toys), and

grid-scrambled versions of these images serving as visual controls (**Figure 1B**). Over

the course of two 5 min runs, participants saw 180 different images presented in a

block design interspersed with an implicit visual baseline. We analysed the dog fMRI

data with a newly developed analysis protocol shown to significantly improve neural

signal detection power [49]. All statistical tests were corrected for multiple comparisons

(see **fMRI data analysis, Figures 1-3**, **Supplementary file 1: Tables S1-S5**).

Figure 1. *Overview of the comparative experimental approach to study the neural bases of face and body perception in dogs and humans.* (**A**) We obtained all imaging data using a 3T Siemens Skyra MR-system, equipped with a 15-channel human knee coil to scan the dog participants, and using a 32-channel human head coil for the human participants (not depicted). Dogs received extensive training [50] to stay motionless without restraints or sedation and could leave the MR scanner at any time via a custom-made ramp positioned at the scanner bed. All (dog and human) participants wore earplugs for noise protection and dog participants had an additional head bandage to secure optimal positioning of the earplugs throughout the scan session. (**B**) All participants viewed images of human and dog bodies, human and dog faces, inanimate objects (e.g., toys) or grid-scrambled versions of these images presented in a block design. (**C**) We split the data sets and used data from the first task run to compare activation for visual stimulation (all stimuli) to the visual baseline (i.e., the localizer data set). In both participant samples, we observed activation in the occipito-temporal cortices. These task-responsive voxels then served as a search space to determine individual functional regions-of-interest (fROIs), additionally constrained by bilateral anatomical regions (see **Figure 2A**). Results are displayed at *p* < 0.05, *FWE*-corrected at cluster-level using a cluster-defining threshold of *p* < .001 (dogs, **Supplementary File 1: Table S1**), and *p* < 0.05 whole-brain *FWE*-corrected (humans; **Supplementary File 1: Table S1**) Results for the dogs are overlaid onto a canine breed-averaged surface render [51], and onto the FreeSurfer "FsAverage" surface for humans (https://surfer.nmr.mgh.harvard.edu/fswiki). Anatomical nomenclature for all figures refers to the dog brain atlas from Czeibert and colleagues [52] and to the human brain atlas from the Laboratory for Neuro Imaging (LONI) Brain Atlas [53] (LPBA40; http://www.loni.usc.edu/atlases/). R, right; L, left; A, anterior; P, posterior; STS, superior temporal sulcus; MT, middle temporal visual area (V5); LOC, lateral occipital cortex.

2.1 Mass-univariate analysis: Face- or body-sensitive areas in the occipito-temporal cortices of dogs and humans

To start out, we investigated whether dogs and humans have specialized cortical regions for face and body perception, and whether these regions responded differently to images of con- or heterospecifics. To this end, we used a standard functional region-

of-interest analysis approach (fROI,[54–56]). We split the data into two orthogonal data

sets, (1) a localizer data set (first task run) to define potential face- or body-sensitive

areas in visual-responsive brain regions (**Figure 1C**, **2A**) and (2) a test data set

(second task run) to extract activation levels from these regions and to test potential

category sensitivity. We chose this approach for two reasons. First, we aimed at

directly testing the potential category-sensitivity, quantified by increased activation

levels. Second, our goal was to account for differences in the location of activation

peaks between individuals as reported in previous dog imaging studies [41,43].

2.1.1 Functional regions-of-interest (fROIs)

For each participant, we defined bilateral fROIs within constrained search spaces

to preserve spatial information (i.e., the rough anatomical location of activation peaks).

For the human participants, we used previously reported anatomical regions known to

be engaged in face and body perception [57]: the extrastriate body area, fusiform body

area, occipital face area and fusiform face area (**Figure 2A**). For the dog participants,

we could not build on previous research due to lack of shared template space and

availability of data. Therefore, we first compared activation levels associated with the

visual presentation of all stimuli compared to implicit baseline (i.e., white cross

presented on grey background) using the localizer data set. This revealed task-

responsive activation within the occipital-, splenial-, ectomarginal-, caudal-, medial

suprasylvian- and marginal gyri, partially overlapping with results from previous studies

investigating face perception [29,40–43,58,59] (**Figure 1C**, **Supplementary File 1:**

Table S1). We then used anatomical masks [52] of these regions as search spaces for

the dog participants (**Figure 2A**). For comparison, we also report task-responsive

voxels for the human participants, confirming involvement of the occipito-temporal

cortex including the lateral occipital cortex and fusiform gyrus (**Figure 1C**,

Supplementary File 1: Table S1; see also **Figure 4** for schematic figures of the dog

and human brain).

We then defined individual category-sensitive regions within each search space

by selecting the top-10% voxels with the strongest activation for bodies or faces

compared to inanimate objects based on the individual data from each participant.

2.1.2 Sensitivity for animate stimuli but sub-division in face- and body-sensitive regions exclusive to humans

Next, we extracted the mean activation signal during viewing of faces, bodies,

inanimate objects and scrambled images (all compared to the implicit baseline) from

the individual fROIs using the test data set. In dogs, we observed significantly

increased activation levels for faces and bodies compared to inanimate objects or

scrambled images in both face and body fROIs, located in the medial and caudal

suprasylvian- and ectomarginal gyrus. In the caudal suprasylvian face- and

ectomarginal body fROI, the increased activation for faces showed the same trend but

did not always reach significance. We did not find consistent evidence for a further

sub-division into face- and body-selective areas in the occipito-temporal cortex, as

there was no significant activation difference between faces and bodies in seven out

of eight fROIs. We only observed significantly increased activation for bodies

compared to faces in the caudal suprasylvian face but not body fROI. The occipital and

splenial face and body fROIs did not result in differential activation changes in

response to any of the stimulus categories (**Supplementary File 1: Table S2**).

Regarding the parietal cortex, the marginal gyrus appeared solely responsive to bodies

compared to all other conditions with no activation differences for faces compared to

inanimate objects or scrambled controls (**Figure 2B**).

A Bilateral anatomical search spaces to create individual functional regions-of-interest (fROIs)

B fROI analysis: mean activation extracted from independent test data set

Figure 2. Strongest activation levels for animate stimuli in both species, but differences between faces and bodies exclusive in humans. (**A**) Based on the localizer data set, we defined individual category-specific regions-of-interest (functional region-of-interest approach, fROI) within multiple anatomically constrained search spaces using the contrasts bodies > inanimate objects (body fROIs) and faces > inanimate objects (face fROIs). (**B**) From these individual fROIs, we then extracted activation estimates during the viewing of faces, bodies, inanimate objects and scrambled images (all compared to the implicit baseline) using the data from the independent test data set to test potential face- or body-sensitivity. (**B**) In dogs, we observed increased activation levels for faces and bodies compared to inanimate objects or scrambled images in three occipito-temporal regions (medial and caudal suprasylvian, ectomarginal gyrus) in face and body fROIs. However, no difference between faces and bodies except in the caudal suprasylvian face fROI. The face and body fROIs located in the occipito-parietal marginal gyri resulted in increased activation levels primarily for bodies. In humans, similar to dogs, we observed the strongest activation levels for faces and bodies in all fROIs. In contrast to the dog data, the human data analysis consistently resulted in a significant activation change for bodies compared to faces in body fROIs and vice versa (**Supplementary File 1: Table S2**). Planned comparisons were corrected for multiple comparisons using Tukey method. *$p < .05$, **$p < .01$, ***$p < .001$, ***$p < .0001$, error bars represent the standard error of the mean; a.u., arbitrary units. Atlas maps and search spaces refer to the dog brain atlas from Czeibert and colleagues [52] and human parcels were retrieved from previous work [57]. A, anterior; P, posterior; S, superior; L, left; R, right.

Results for the human participants confirmed earlier research [10–13,15–17,21–

28] with the strongest overall activation levels for bodies in the extrastriate and fusiform

body areas and for faces in the occipital and fusiform face areas (**Figure 2B**,

Supplementary File 1: Table S2). In summary, we found multiple occipito-temporal

regions in the dog and human cortices responsive to animate stimuli, but detected a

further sub-division into face- and body-sensitive regions exclusively in the human

brain (see also **Figure 4** for a summary of the results).

163 2.1.3 Conspecifics preference in human extrastriate body and fusiform
164   face area

Next, we investigated whether processing of faces or bodies differed when

participants saw images of con- compared to heterospecifics (**Supplementary File 1:**

Table S3). In humans, this revealed increased activation for human (mean: 4.46, *SD*

= 1.83) compared to dog bodies (mean: 3.76, *SD* = 1.66) in the extrastriate body area

($t(39)$ = 3.14, *p* = 0.003, mean$_{Diff}$: 0.7, 95% CI [0.25, 1.15]), as well as a general

activation increase for human (mean: 1.89, *SD* = 0.87) compared to dog (mean: 1.64,

SD = 0.81) stimuli in the fusiform face area ($t(79)$ = 3.71, *p* < 0.0001, mean$_{Diff}$: 0.25 ,

95% CI$_{meandiff}$ [0.16, 0.38]). For the dogs, we did not find a difference between the

perception of dog and human stimuli.

174 2.2 Multivariate pattern analysis of animacy perception

Despite the overall activation (measured using mass-univariate analyses) to

stimulus categories may not differ, we reasoned that more fine-grained analyses of

multivariate activation patterns could reveal distinctions [60]. Therefore, we

investigated the neural representations of faces and bodies in dogs and humans and

their potential correspondence using whole-brain representational similarity analyses

(RSA). We moved a 4 and 8 mm searchlight across the whole dog and human brain,

respectively, in order to determine individual pattern similarity maps between all trials

of each stimulus category [61–63] (**Figure 3A**). On the group-level, we then conducted

paired *t*-tests to compare the pattern similarities maps of (1) animate vs. inanimate

stimuli, (2) bodies vs. inanimate objects, (3) faces vs. inanimate objects, (4) faces vs.

bodies, and (5,6) within the face and body categories: images of conspecifics vs.

heterospecifics (i.e., dog faces vs. human faces; dog bodies vs. human bodies).

Results revealed increased (within-category) pattern similarity for animate

compared to inanimate stimuli in the occipito-temporal cortex of dogs (i.e., caudal

suprasylvian and splenial gyrus) and humans (i.e., middle occipital gyrus) overlapping

with the identified fROIs. In human participants, we additionally observed increased

similarity in the cerebellum and fronto-parietal regions (**Supplementary File 1: Tables**

S4-S5). In brief, this indicates distinct neural representations of animate compared to

inanimate stimuli in higher-order visual regions of both species.

2.2.1 Similar and divergent neural representations of bodies and faces

Moving on to neural representations for bodies and faces, for bodies we observed

increased pattern similarity compared to inanimate objects in higher-order visual areas

in the occipito-temporal cortex, partially overlapping with the identified fROIs, and the

cerebellum in both species (**Figure 3B**). Within the same regions (and again, in both

species), we also observed distinct activation patterns for conspecific compared to

heterospecific bodies. Furthermore, when the dogs viewed dog bodies, results

revealed increased similarity in limbic structures and regions associated with olfaction

(**Figure 3C**). In humans, we observed increased similarity for human compared to dog

bodies in fronto-parietal regions associated with action perception [64–66] (**Figure 3C**).

Observing faces compared to inanimate objects resulted in significantly

increased pattern similarity in occipito-temporal cortices of both species and again in

dog olfactory structures (**Figure 3D**). Comparing similarity patterns for faces vs.

bodies, we only found significant differences for the human participants (see

Supplemental file 1: Tables S4-5 for detailed results and **Figure 4** for a schematic

summary of the main results).

Figure 3. *Whole-brain representational similarity analysis (RSA) revealed similar and divergent neural patterns of activation for animate vs. inanimate stimuli but differences between faces and bodies again exclusive in human.* (**A**) We moved a 4/8 mm (dog/human) spherical searchlight across the whole brain to quantify the neural patterns underlying stimuli categories. Pattern similarity matrices were computed across the respective quadrants marked in black. (**B,D**) In both species, we observed increased pattern similarity for bodies or faces compared to inanimate stimuli in higher-order visual areas in the occipito-temporal cortex and the cerebellum. (**C,D**) In dogs we additionally detected distinct neural patterns of activation for faces and conspecific (= dog) bodies in olfactory and limbic regions. In humans, we observed an increased pattern similarity for (conspecific) bodies in fronto-parietal regions associated with action perception [64–66]. All results are displayed at *p* < 0.05, *FWE*-corrected at cluster-level, and using a cluster-defining threshold of *p* < .005 for the dog and *p* < .001 for the human data (**Supplementary file 1: Tables S4-5**). Anatomical locations are shown in panel B for the dog and in D for the human data: superior (S), anterior (A); all sagittal, coronal and axial planes displayed have the same orientation. Coordinates refer to a canine-breed averaged template [51] or to MNI space for the human data. Person's *r*, correlation coefficient; t, *t*-value; g., gyrus; PFC, prefrontal cortex LOC, lateral occipital cortex; MT, middle temporal visual area (V5); pSTS, posterior superior temporal sulcus; R, right; L, left.

3 Discussion

Our findings revealed that the occipito-temporal cortex has a prominent role in the perception of animate entities in both dogs and humans. This suggests that the neural processing of animate vs. inanimate stimuli across mammalian brains may either have an old evolutionary origin in the common ancestors of canines and primates or is the result of convergent evolution [48]. However, within these regions, only humans displayed an additional sub-division into distinct face- and body-sensitive brain regions. Moreover, only dogs showed distinct responses to human and dog faces as well as dog bodies in olfactory areas (see **Figure 4** for schematic summary of the results). These findings suggest a divergent evolution of the neural bases of face and body perception (or processing) in dogs and humans. The differential engagement of visual and olfactory brain functions would fit particularly well with the differential sensitivity and preferential use of these perceptual systems to infer social and contextual information.

Detecting biological agents is crucial for survival and social relationships. Hence, the visual differentiation between animate vs. inanimate entities is an evolutionary important category representation. Studies show that dogs target animate agents when, for example, presented with images of wild animals, humans or dogs embedded in natural landscapes [67] or of social interactions between humans and dogs [68]. Further, dividing stimuli into an animate vs. inanimate dimension is one of the first visually salient categorizations formed by human infants [69]. In this way, animacy representation provides the first building block for more complex visual categorizations such as faces vs. bodies. Thus, considering its biological significance and the observed cross-species similarities in our study, animacy might constitute a general organizing principle across mammalian brains.

Figure 4. *Summary of the main study findings.* Results from the functional regions of interest (univariate activation strength) and representational similarity analysis (multivariate activation pattern) illustrating the results on schematic brain figures. For simplification, observed results are always summarized on one hemisphere; they do not mark the exact but the approximate anatomical location.

238  Behavioural [7,34,35,44,70–73] and imaging studies [58,59] have also

239  demonstrated that dogs are able to perceive facial and bodily cues of dogs and humans

240  and display high responsiveness to ostensive-referential bodily cues [45,46,74,75],

241  already as puppies [76]. Our results do not contradict these findings, but suggest that

242  visual regions involved in the perception of faces are also involved in the perception of

243  other body parts, and may thus encode animate stimuli. A recent dog neuroimaging

244  study [77] also showed that the visual regions observed in our study play a crucial part

in the perception of complex social interactions, which would confirm their important

role for social perception.

While previous investigations of how dogs perceive bodies have mainly focused

on the decoding of human referential cues, we hope that our work will inspire more

research on how dogs perceive bodily social as well as emotional cues of con- as well

as heterospecifics. The absence of a further sub-specialization for face vs. body

perception in the occipito-temporal cortex of dogs might indeed indicate that dogs in

comparison to humans focus more on whole body social cues rather than on specific

sub-parts. This interpretation is notably well in line with a recent comparative eye-

tracking study [78].

Regarding species perception, our results indicate an increased salience for

human compared to dog stimuli in humans, but no evidence for a conspecific-

preference in the occipito-temporal cortex of dogs. Thus, human and dog stimuli

appear to be equally salient for dogs. This is in line with previous behavioural studies

suggesting, for example, no significant difference in the perception of human or dog

emotional facial [35], whole-body cues [78], or images of social interactions [68].

Further, two studies demonstrated that dogs, already as puppies, follow human

gestural-communication and show an interest in human faces [76], which has not been

observed in wolf puppies [79]. Additionally, half of the variation in these socio-cognitive

skills could be accounted for by genetic factors, suggesting that dogs' attention to

humans might have been enhanced during domestication [75 for review of both

studies]. Future studies investigating species-preferences in dogs might thus consider

adding further heterospecific stimuli of other familiar species, such as cats.

In accordance with a potential divergent evolution of face and body

representations in dogs and humans, we also observed distinct neural representations

for faces (regardless of species) and conspecific (dog) bodies in dog olfactory regions.

Although unexpected, these results might reflect dogs' significantly higher olfactory

sensitivity and it's use to infer social and contextual information [54 for review].

However, the link between cognition and olfaction remains speculative since

behavioural studies using olfactory cues so far mainly explored dogs' odour and

disease detection abilities (e.g., [82–86] while dog imaging studies on olfaction focused

on reward-processing [87,88] or general sensory abilities [89–91]. However, a recent

behavioural study investigating dog cognition using olfactory cues showed first

evidence that dogs form representations of their owners when smelling their tracks

[92].

In humans, results revealed distinct activation patterns for bodies in fronto-

parietal regions that have been consistently associated with the perception of actions

[64–66]. In dogs, we did not find activation in (pre-)motor cortices, but the increased

neural sensitivity for bodies in the occipito-parietal marginal gyrus might be initial

evidence for an analogous visual pathway to encode actions. However, this is beyond

our study's scope, and the precise neural underpinnings of action observation in dogs

have not yet been investigated.

We have taken several steps to maximize neural signal sensitivity already when

designing our study (e.g., by using a block design) and at the analysis stage (e.g., by

using dog-tailored hemodynamic models increasing response detection power [49]).

Nevertheless, the extensive training required for dogs [50] resulted in different sample

sizes for the two species, and the more extensive prior work in humans [57] resulted

in more constrained search spaces, and likely higher spatial sensitivity for the human

data. Further, the majority (80%) of our pet dogs were pure-bred herding dogs, which

prevented us from testing potential breed-specific differences [as in e.g. 90] and may

limit generalizability.

In conclusion, our study reveals novel evidence for similarities and differences of

animacy perception between two phylogenetically distant mammal species, advancing

our understanding of the foundations of social cognition and behaviour. Finally, we

provide insights into the differentially evolved sensory systems of dogs and humans

for the perception of faces and bodies.

4 Materials and Methods

4.1 Participants

Fifteen pet dogs (*Canis familiaris*; 11 females, age range: 4-11 years, mean age:

7.8 years) consisting of 10 Border collies, 2 Australian shepherds, 1 Labrador retriever

and 2 mixed-breed dogs and forty human participants (22 females, age range: 19-28

years, mean age: 23 years) participated in the present study.

For the dog sample size we aimed to collect data from a minimum of $N = 12$ dogs

during the data collection period from April 2019 to July 2020 which was the median

sample size of task-based dog fMRI studies at the time of planning the study (the

median sample size in 2021 is also $N = 12$). For the human participants, we aimed for

a sample size of $N = 40$ participants, based on previous studies in our lab with similar

task designs, and also being in line with a previous study [57] investigating the neural

bases of face and body perception using a functional region-of-interest approach with

a sample size of $N = 35$ participants (reporting *F*-values above 40 for all repeated

measures analyses).

Prior to MRI scanning, all dogs underwent initial medical examination concerning

their eyesight and general health and received extensive training to habituate to the

MRI environment [50]. They were fully awake, able to exit the MRI scanner at any time

and equipped with noise protection (**Figure 1A**). All caregivers gave informed written

consent to their dogs' participation. Human participants were right handed, had normal

or corrected-to-normal vision, they reported no history of neurological or psychiatric

disease, nor phobia of dogs, fulfilled the standard inclusion criteria for functional MRI,

and gave informed written consent to participate. The study complies with the ARRIVE

guidelines [94]. Dog data collection was approved by the institutional ethics and animal

welfare commission in accordance with Good Scientific Practice (GSP) guidelines and

national legislation at the University of Veterinary Medicine Vienna (ETK-06/06/2017),

based on a pilot study conducted at the University of Vienna. Human data collection

was approved by the ethics committee of the University of Vienna (reference number:

00565) and performed in line with the latest revision of the Declaration of Helsinki

(2013).

331   4.2 Task and procedure

We employed a block design (duration: 12 s) split in two 5 min runs where

participants saw images of faces and bodies of dogs or humans, objects and

scrambled versions of these images (5 images per block; **Figure 1B**, see below) on

an MR-compatible screen (32 inch) positioned at the end of the scanner bore. Crucially,

we used the same task for dogs and humans. Human participants were instructed to

watch the images presented on the MR screen and dogs were trained to attend to the

MR screen (passive viewing paradigm). Each run contained three blocks per condition,

block order was randomized but the same condition was never presented twice in a

row. Between blocks, participants saw a visual baseline jittered between 3-7 s with a

white cross presented on grey background. Image composition for each block and

order within each block was randomized across participants to ensure effects are not

driven by specific blocks; each image was presented once. The task was implemented

using PsychoPy [95]. Overall motion and wakefulness were live monitored via the

camera of an eye-tracker (Eyelink 1000 Plus, SR Research, Ontario, Canada)

positioned below the MR-compatible screen. For the dogs, the trainer stayed within the

scanner room but out-of-sight throughout the scan session to monitor and handle the

dogs. Human participants saw both runs within a single scanner session with a short

break in-between. For the dogs, the number of attempted sessions varied depending

on how many repetitions they needed to complete one run without substantive motion

and with sufficient attentiveness (i.e., eyes open and gazing towards the centre of the

screen); in-between task runs the dogs always were given a short break outside the

MR scanner. Realignment parameters were evaluated after each scan session. If

overall motion exceeded 4 mm in any of the three translation directions, the dog was

re-invited to repeat the run in a subsequent session and sessions were scheduled at

least one week apart. On average, dogs needed three sessions to complete both runs.

No data of the non-successful sessions were used for analysis.

4.2.1 Stimulus material

The stimulus set comprised 180 coloured images of faces and bodies of dogs

and humans, everyday inanimate objects (e.g., a toy, a chair), and phase-grid

scrambled versions of each category (30 images per condition) derived from Wagner

and colleagues [96], the Hemera Photo-Object data base (Hemera Technologies) and

the internet (see **Figure 1B** for examples). The images were resized to 600 × 600

pixels and presented in the centre of the MR screen on grey background. We cut out

the heads off from the body images, as well as objects from body images (e.g., a coffee

cup, a soccer ball). The images showed a variety of postures (e.g., jumping, looking

up), neutral and positive emotional displays (e.g., sleeping, smiling) and viewing

perspectives (e.g., from above, from a side angle). Luminance was set homogeneously

across stimuli and we controlled for spatial extent (see **Supplementary file 2A** for

details).

4.3 MRI data acquisition

We acquired all MRI data with a 3T Siemens Skyra MR-system (Siemens

Medical, Erlangen, Germany) and a 15-channel coil (initially designed for

measurements of the human knee) for data acquisition in dogs and a 32-channel

human head coil for data acquisition in humans. Functional scans of dogs used a 2-

fold multiband (MB) accelerated echo planar imaging (EPI) sequence including the

following parameters: voxel size = 1.5 x 1.5 x 2 mm^3, repetition time (TR) / echo time

(TE) = 1000/38 ms, field of view (FoV) = 144 × 144 × 58 mm^3, flip angle = 61°, 20%

gap and 24 axial slices covering the whole brain (interleaved acquisition, descending

order). Structural scans had a voxel size of 0.7 mm isotropic (TR/TE = 2100/3.13 ms,

FoV = 230 × 230 × 165 mm^3) and were acquired in a separate scan session prior to

functional data collection. Human functional scans were acquired using a 4-fold MB

accelerated EPI sequence including the following parameters: voxel size = 2 mm

isotropic, TR/TE = 1200/34 ms, FoV = 192 × 192 × 124.8 mm^3, flip angle = 66°, 20%

gap and 52 axial slices coplanar to the connecting line between anterior and posterior

commissure (interleaved acquisition, ascending order). Additionally, we obtained field

map scans to correct functional scans for magnetic field inhomogeneities using a

double echo gradient echo sequence with the following parameters: voxel size = 1.72

x 1.72 x 3.85 mm^3, TR/TE1/TE2 = 400/4.92/7.38 ms, FoV = 220 × 220 × 138 mm^3, flip

angle = 60° and 36 axial slices (same orientation as functional scans). Structural scans

had a voxel size of 0.8 mm isotropic (TR/TE = 2300/2.43 ms,

FoV = 256 × 256 × 166 mm^3) and were acquired after functional data acquisition.

4.4 Data processing and statistical analysis

Imaging data was preprocessed and analyzed using SPM12

(https://www.fil.ion.ucl.ac.uk/spm/software/spm12/), Matlab 2018b (MathWorks) and R

3.6.3 [97].

4.4.1 MRI data preprocessing

In both samples, we slice-time corrected (reference: middle slice) and realigned functional images to the mean image. Human imaging data was also unwarped using the acquired field map. Dog imaging data was manually reoriented with the rostral commissure set as a visual reference (SPM module: "*Reorient images / Set origin*") to match the template orientation [51] and structural images were skull-stripped using individual binary brain masks created using itk-SNAP [98]. Next, we co-registered the structural and functional images to the mean functional image, segmented the structural images in both samples and normalized the human data to the Montreal Neurological Institute (MNI) template space and the dog data to a breed-averaged stereotaxic template space [51]. Normalized images were resliced to 1.5 mm isotropic and smoothed with a 3-dimensional Gaussian kernel (full-width-at-half-maximum, FWHM; with twice the raw voxel resolution: dogs/humans = 3/4 mm; see [49] for an in-depth description of our dog data preprocessing pipeline). We then calculated individual scan-to-scan motion (frame wise displacement, FD) and added motion regressors to first-level GLMs for each scan exceeding the a priori set FD threshold of 0.5 mm (motion scrubbing [99,100]) to account for both translational and rotational displacements. For the dog participants, we removed on average 8% of the scans from each run (run 1: mean FD = 0.23 mm, 90th percentile = 0.36 mm; run 2: mean FD = 0.24 mm, 90th percentile = 0.38 mm) and 1% of the scans from each run of the human participants (run 1: mean FD = 0.17 mm, 90th percentile = 0.22 mm; run 2: mean FD = 0.18 mm, 90th percentile = 0.21 mm).

4.4.2 fMRI data analysis

Mass-univariate activation analysis. We analysed the functional data using a general linear model (GLM) approach implemented in SPM12. Individual GLM matrices included six regressors of interest for task (dog faces, dog bodies, human

faces, human bodies, inanimate objects, scrambled) and the six regressors from the

realignment procedure (see above) were added as nuisance regressors. All blocks

were estimated using a boxcar function time-locked to the onset of each block with a

duration of 12 s. For the dog data, the signal was convolved with a tailored dog

haemodynamic response function [49] (HRF), while the standard human canonical

HRF (i.e., the default HRF parameters provided by SPM12) was used for the human

data. The dog HRF reflects a previously observed 2-3 s earlier peak of the BOLD signal

than expected by the human HRF model [49]. Normalized, individual binary masks

served as explicit masks and we applied a high-pass filter with a cut-off at 128 s. We

then split the data in two sets (task run 1, task run 2). Based on the data from the first

task run, we estimated three subject-level contrast maps for the difference between

our conditions of interest (i.e., faces, bodies with equal weights for human and dog

images) and objects, and a visual stimulation contrast (all conditions > implicit visual

baseline). For the second task run, we computed eight subject-level contrasts, one for

each task regressor (i.e., dog bodies, human bodies, dog faces, human faces,

everyday inanimate objects, scrambled), averaged for faces (i.e., dog and human

faces) and averaged for bodies (i.e., dog and human bodies) compared to implicit

visual baseline.

Functional region of interest (fROI) approach. We implemented a standard

functional region-of-interest (fROI, [54–56]) approach to investigate potential category-

specificity of cortical regions. This approach allows us to directly test the potential

category-sensitivity quantified by increased activation level while accounting for slight

differences in the location of activation peaks between individuals as which have been

reported in previous dog imaging studies [41,43].

The participant-level contrast images from the first run served as localizer data to

define individual category-sensitive regions. Within anatomically constrained search

spaces (**Figure 2A**) we selected the top-10% voxel from each hemisphere with the

strongest signal for the condition-of-interest (i.e., face areas: faces > objects, body

areas: bodies > objects) to form bilateral individual fROIs. Selecting the top-10% voxel

was an *a priori* decision we made because the resulting fROI sizes (on average: 30

voxels) were similar to the average cluster size observed in a previous dog imaging

study that used two visual paradigms and the same smoothing kernel [49]. Further,

percentages instead of absolute numbers took also the varying search space sizes

into account (see **Supplementary file 2B** and **Figure 2 – figure supplements 1-2** for

more information). The data from the left out second run allowed then to directly test

potential category specificity in an independent data set. Thus, we extracted parameter

estimates from the conditions-of-interest contrasted against implicit visual baseline

from both body and faces areas (= individual fROIs) using the REX toolbox [101].

Group functional ROI analyses were performed running repeated measures

analyses of variance (ANOVAs). First, we tested our main research question, whether

the body areas resulted in increased sensitivity for bodies regardless of species and

vice versa for faces, and added everyday objects and scrambled images as further

conditions. Thus, we ran a one-way ANOVA with *stimuli category* (faces, bodies,

everyday objects, scrambled*)* as dependent variable. Next, to investigate if there is a

difference in activation between conspecific and heterospecific stimuli, we used 2 x 2

within-subjects ANOVAs (*species*: conspecific, heterospecific; *image category*: face,

body). *P*-values for group comparisons investigating the same research questions

were Bonferroni-adjusted, e.g., if the 6 potential body fROIs result in increased

sensitivity for bodies compared to all other categories results in an adjusted *p*-value of

0.008 (= 0.05/6). Adjusted *p*-values are reported along with the results

(**Supplementary File 1: Tables S2-3**). For whole-brain group analyses, we tested

significance applying a whole-brain familywise error (*FWE*) correction for the human

data and a cluster-level inference with a cluster defining threshold of $p < 0.001$ and a

cluster probability of $p < 0.05$ *FWE* corrected for multiple comparisons for the dog data.

Cluster extent (i.e., minimum spatial extend to be labelled significant) was calculated

using the SPM extension "CorrClusTh.m" (by Thomas Nichols, University of Warwick,

United   Kingdom,   and   Marko   Wilke,   University   of   Tübingen,   Germany;

https://warwick.ac.uk/fac/sci/statistics/staff/academic-research/nichols/scripts/spm/).

Representational similarity analysis. Next, we investigated the neural

representations for faces, bodies and objects and their potential convergence in dogs

and humans. To this end, we performed a whole-brain representational similarity

analysis (RSA; [61–63]) to determine neural pattern similarities within image

categories. GLMs were modelled identical to univariate GLMs (see above) but for each

block, we ran a separate GLM with the block as task regressor and remaining blocks

were combined in one regressor of no interest [102]; runs were modelled

independently. Thus, the analysis resulted in 36 beta images for each participant (6

conditions × 6 trials). RSA was performed using the smoothed functional data. For all

RSA analyses, we moved a spherical searchlight [103] (dogs: $r = 4$ mm, 81 voxel;

humans: $r = 8$ mm, 251 voxel) throughout individual whole-brain grey matter masks

computed based on the normalised segmentation output considering only searchlights

with a minimum of 15 grey matter voxel for the dog and 30 for the human data.

Within a given searchlight, we extracted single-trial beta estimates from all voxels

and sorted them according to their stimuli category (**Figure 3A**; dog / human bodies,

dog / human faces, inanimate objects) and reshaped the data to a trial × voxel matrix.

Next, we *z*-scored the data across trials and runs and computed a trial × trial similarity

matrix by correlating each beta estimate of each trial to all other trial estimates. Finally,

in order to retrieve overall similarity scores, we then Fisher's *z*-transformed the data

and calculated overall similarity matrices by averaging scores across the respective

quadrants. We were specifically interested in pattern similarities across animate vs.

inanimate (faces × bodies vs. inanimate objects), faces or bodies vs. inanimate objects,

faces vs. bodies and conspecific vs. heterospecific species dimensions within face and

body categories. We then assigned the overall similarity values to the centre voxel of

each searchlight resulting in individual whole-brain pattern similarity maps.

On the group-level, we then used paired *t*-tests to compare the pattern similarities

between trials of (1) faces vs. inanimate objects (i.e., [dog faces × human faces] vs.

inanimate objects), (2) bodies vs. inanimate objects (i.e., [dog bodies × human bodies]

vs. inanimate objects), (3) animate vs. inanimate images (i.e., [dog faces × human

faces x dog bodies × human bodies] vs. inanimate objects), (4) faces vs. bodies (i.e.,

[dog faces × human faces] vs. [dog bodies × human bodies]), and (5,6) within the face

and body categories: images of conspecifics vs. heterospecifics (i.e., dog faces vs.

human faces; dog bodies vs. human bodies). We computed permutation tests [104] to

determine group-level significance using the Statistical nonParametric Mapping

(SnPm13, http://www.nisox.org/Software/SnPM13/) toolbox running 5000

permutations for each paired *t*-test and applied cluster-level inference with a cluster

defining threshold of $p < .005/.001$ (dogs/humans) and a cluster probability of $p < 0.05$

familywise error corrected (*FWE*) for multiple comparisons.

Data visualization. Data was visualized using SPM12

(https://www.fil.ion.ucl.ac.uk/spm/software/spm12/), Matlab 2018b (MathWorks), itk-

SNAP [98], R 3.6.3 [97] mainly using the packages ggplot2 [105] and RainCloudPlots

[106] and the python project nilearn (http://nilearn.github.io).

5 Data availability statement

The code to create functional regions-of-interest (fROIs) within anatomical

search spaces and to extract activations levels, as well as the representational

similarity analysis (RSA) code are openly available on Github

(https://github.com/magdalenaboch/fROI-analysis,

https://github.com/isabellawagner/searchlight-rsa). The extracted fROI activation

levels along with the R code to visualize and analyse the data, the overlap maps of

individual fROIs, thresholded and unthresholded statistical maps of univariate (all >

implicit baseline) and RSA analysis and averaged structural images to serve as

background images to overlay the data on have been uploaded to the Open Science

Framework and are openly available at https://osf.io/kzcs2/.

6 Acknowledgments

535 We thank Prof. Nancy Kanwisher for her helpful comments on the preliminary

536 results and analysis plan. We also want to thank Morris Krainz, Anna Thallinger, Olaf

537 Borghi and Helena Manzenreiter for their help collecting the data, Boryana Todorova

538 for her support in preparing the stimuli set and all the dogs and their caregivers and all

539 human participants for taking part in this project.

7 CRediT authorship statement

Magdalena Boch: Conceptualization, Methodology, Software, Validation,

Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review &

editing, Visualization, Project administration. **Isabella C. Wagner**: Conceptualization,

Methodology, Resources, Writing - original draft, Writing - review & editing,

Supervision. **Sabrina Karl**: Investigation, Writing - review & editing. **Ludwig Huber**:

Conceptualization, Resources, Writing - review & editing, Supervision, Funding

acquisition. **Claus Lamm**: Conceptualization, Methodology, Resources, Writing -

original draft, Writing - review & editing, Supervision, Funding acquisition.

8 Declaration of Competing Interests

The authors declare no competing financial interests.

9 Supplementary Files

Supplementary file 1. Supplementary tables: **Table S1.** Task-related activation: localizer data set (task run 1), **Table S2.** Results from one-way[a] repeated measures analyses of variance (ANOVAs), **Table S3.** Results from 2 × 2[a] repeated measures analyses of variance (ANOVAs), **Table S4.** Dog data: Pattern similarity during face, body and object perception and **Table S5.** Human data: Pattern similarity during face, body and object perception.

Supplementary file 2. 2A: Stimulus material: luminance and spatial extent, **2B:** Anatomical search spaces and size individual fROIs

561 **10 Figure supplements**

Figure 2 – figure supplement 1. *Overlap of individual functional regions-of-interest (fROIs) from dog participants.* Individual fROIs were defined for faces (> inanimate objects) and bodies (> inanimate objects) within multiple anatomical search spaces (A-D) based on the data from localizer data set (i.e., first task run). The colour of the heat map indicates the amount of overlap between the fROIs of all dog participants. We observed slight individual differences in anatomical location between participants as indicated by a maximum overlap of *N* = 7 from overall 15 participants. The axial plane in panel A (top row) shows the anatomical locations posterior (P) and right hemisphere (R); all axial planes displayed have the same orientation. The sagittal plane displays the cut coordinates (z) and the anatomical locations superior (S) and anterior (A); z-coordinates refer to a canine-breed averaged template [51].

Figure 2 – figure supplement 2. *Overlap of individual functional regions-of-interest (fROIs) from human participants.* Individual fROIs were defined for faces (> inanimate objects) and bodies (> inanimate objects) within multiple anatomical search spaces (A-D) based on the data from the localizer data set (i.e. first task run). The colour of the heat map indicates the amount of overlap between the fROIs of all humans. We observed slight individual differences in anatomical location between participants as indicated by a maximum overlap of N = 30 from overall 40 participants. The axial plane in panel A shows the anatomical locations posterior (P) and right hemisphere (R); all axial planes displayed have the same orientation. The sagittal planes display the cut coordinates (z) and the anatomical locations superior (S) and anterior.

11 References

1. Rajhans P, Jessen S, Missana M, Grossmann T. Putting the face in context: Body expressions impact facial emotion processing in human infants. Dev Cogn Neurosci. 2016;19: 115–121. doi:10.1016/J.DCN.2016.01.004

2. Aviezer H, Trope Y, Todorov A. Body cues, not facial expressions, discriminate between intense positive and negative emotions. Science. 2012;338: 1225–1229. doi:10.1126/science.1224313

3. Barrett LF, Mesquita B, Gendron M. Context in emotion perception. Curr Dir Psychol Sci. 2011;20: 286–290. doi:10.1177/0963721411422522

4. Lecker M, Dotsch R, Bijlstra G, Aviezer H. Bidirectional contextual influence between faces and bodies in emotion perception. Emotion. 2020;20: 1154–1164. doi:10.1037/EMO0000619

5. Poyo Solanas M, Zhan M, Vaessen M, Hortensius R, Engelen T, de Gelder B. Looking at the face and seeing the whole body. Neural basis of combined face and body expressions. Soc Cogn Affect Neurosci. 2018;13: 135–144. doi:10.1093/scan/nsx130

6. de Gelder B, de Borst AW, Watson R. The perception of emotion in body expressions. Wiley Interdiscip Rev Cogn Sci. 2015;6: 149–158. doi:10.1002/wcs.1335

7. Müller CA, Schmitt K, Barber ALA, Huber L. Dogs can discriminate emotional expressions of human faces. Curr Biol. 2015;25: 601–605. doi:10.1016/j.cub.2014.12.055

8. Kravitz DJ, Saleem KS, Baker CI, Ungerleider LG, Mishkin M. The ventral visual pathway: An expanded neural framework for the processing of object quality. Trends Cogn Sci. 2013;17: 26–49. doi:10.1016/j.tics.2012.10.011

9. Ungerleider LG, Haxby J V. "What" and "where" in the human brain. Curr Opin Neurobiol. 1994;4: 157–165. doi:10.1016/0959-4388(94)90066-3

10. Schwarzlose RF, Baker CI, Kanwisher N. Separate face and body selectivity on the fusiform gyrus. J Neurosci. 2005;25: 11055–11059. doi:10.1523/JNEUROSCI.2621-05.2005

11. Peelen M V., Downing PE. Selectivity for the human body in the fusiform gyrus. J Neurophysiol. 2005;93: 603–608. doi:10.1152/jn.00513.2004

12. Peelen M V., Downing PE. The neural basis of visual body perception. Nat Rev Neurosci. 2007;8: 636–648. doi:10.1038/nrn2195

599   13.   Downing PE, Jiang Y, Shuman M, Kanwisher N. A cortical area selective for
600   visual processing of the human body. Science. 2001;293: 2470–2473.
601   doi:10.1126/science.1063414

602   14.   Spiridon M, Fischl B, Kanwisher N. Location and spatial profile of category-
603   specific regions in human extrastriate cortex. Hum Brain Mapp. 2006;27: 77–
604   89. doi:10.1002/hbm.20169

605   15.   Kanwisher N. Domain specificity in face perception. Nat Neurosci. 2000;3:
606   759–763. doi:10.1038/77664

607   16.   Yovel G, Kanwisher N. Face perception: domain specific, not process specific.
608   Neuron. 2004;44: 889–898. doi:10.1016/j.neuron.2004.11.018

609   17.   Wegrzyn M, Riehle M, Labudda K, Woermann F, Baumgartner F, Pollmann S,
610   et al. Investigating the brain basis of facial expression perception using multi-
611   voxel pattern analysis. Cortex. 2015;69: 131–140.
612   doi:10.1016/J.CORTEX.2015.05.003

613   18.   Wiggett AJ, Pritchard IC, Downing PE. Animate and inanimate objects in
614   human visual cortex: Evidence for task-independent category effects.
615   Neuropsychologia. 2009;47: 3111–3117.
616   doi:10.1016/J.NEUROPSYCHOLOGIA.2009.07.008

617   19.   Chao LL, Haxby J V., Martin A. Attribute-based neural substrates in temporal
618   cortex for perceiving and knowing about objects. Nat Neurosci. 1999;2: 913–
619   919. doi:10.1038/13217

620   20.   Downing PE, Chan AW-Y, Peelen M V., Dodds CM, Kanwisher N. Domain
621   specificity in visual cortex. Cereb Cortex. 2006;16: 1453–1461.
622   doi:10.1093/CERCOR/BHJ086

623   21.   Tsao DY, Freiwald WA, Knutsen TA, Mandeville JB, Tootell RBH. Faces and
624   objects in macaque cerebral cortex. Nat Neurosci. 2003;6: 989–995.
625   doi:10.1038/nn1111

626   22.   Desimone R, Albright TD, Gross CG, Bruce C. Stimulus-selective properties of
627   inferior temporal neurons in the macaque. J Neurosci. 1984;4: 2051–2062.
628   doi:10.1523/jneurosci.04-08-02051.1984

629   23.   Wachsmuth E, Oram MW, Perrett DI. Recognition of objects and their
630   component parts: Responses of single units in the temporal cortex of the
631   macaque. Cereb Cortex. 1994;4: 509–522. doi:10.1093/cercor/4.5.509

632   24.   Gross CG, Bender DB, Rocha-Miranda CE. Visual receptive fields of neurons
633   in inferotemporal cortex of the monkey. Science. 1969;166: 1303–1306.

634    doi:10.1126/science.166.3910.1303

635 25. Haxby J V., Horwitz B, Ungerleider LG, Maisog JM, Pietrini P, Grady CL. The

636    functional organization of human extrastriate cortex: a PET-rCBF study of

637    selective attention to faces and locations. J Neurosci. 1994;14: 6336–6353.

638    doi:10.1523/JNEUROSCI.14-11-06336.1994

639 26. Kanwisher N, Yovel G. The fusiform face area: a cortical region specialized for

640    the perception of faces. Philos Trans R Soc B Biol Sci. 2006;361: 2109–2128.

641    doi:10.1098/rstb.2006.1934

642 27. Tsao DY, Freiwald WA, Tootell RBH, Livingstone MS. A cortical region

643    consisting entirely of face-selective cells. Science. 2006;311: 670–674.

644    doi:10.1126/science.1119983

645 28. Tsao DY, Moeller S, Freiwald WA. Comparing face patch systems in macaques

646    and humans. Proc Natl Acad Sci U S A. 2008;105: 19514–19519.

647    doi:10.1073/pnas.0809662105

648 29. Bunford N, Hernández-Pérez R, Farkas EB, Cuaya L V., Szabó D, Szabó ÁG,

649    et al. Comparative Brain Imaging Reveals Analogous and Divergent Patterns of

650    Species and Face Sensitivity in Humans and Dogs. J Neurosci. 2020;40:

651    8396–8408. doi:10.1523/JNEUROSCI.2800-19.2020

652 30. Kanwisher N, Stanley D, Harris A. The fusiform face area is selective for faces

653    not animals. Neuroreport. 1999;10: 183–187. doi:10.1097/00001756-

654    199901180-00035

655 31. Pinsk MA, Arcaro M, Weiner KS, Kalkus JF, Inati SJ, Gross CG, et al. Neural

656    representations of faces and body parts in macaque and human cortex: a

657    comparative fMRI study. J Neurophysiol. 2009;101: 2581–2600.

658    doi:10.1152/jn.91198.2008

659 32. Hesse JK, Tsao DY. The macaque face patch system: a turtle's underbelly for

660    the brain. Nat Rev Neurosci 2020 2112. 2020;21: 695–716.

661    doi:10.1038/s41583-020-00393-w

662 33. Racca A, Amadei E, Ligout S, Guo K, Meints K, Mills D. Discrimination of

663    human and dog faces and inversion responses in domestic dogs (Canis

664    familiaris). Anim Cogn. 2010;13: 525–533. doi:10.1007/s10071-009-0303-3

665 34. Huber L, Racca A, Scaf B, Virányi Z, Range F. Discrimination of familiar human

666    faces in dogs (Canis familiaris). Learn Motiv. 2013;44: 258–269.

667    doi:10.1016/J.LMOT.2013.04.005

668 35. Albuquerque N, Guo K, Wilkinson A, Savalli C, Otta E, Mills D. Dogs recognize

dog and human emotions. Biol Lett. 2016;12: 20150883.

doi:10.1098/rsbl.2015.0883

36. Huber L. How Dogs Perceive and Understand Us. Curr Dir Psychol Sci.

2016;25: 339–344. doi:10.1177/0963721416656329

37. Maginnity ME, Grace RC. Visual perspective taking by dogs (Canis familiaris)

in a Guesser–Knower task: evidence for a canine theory of mind? Anim Cogn.

2014;17: 1375–1392. doi:10.1007/s10071-014-0773-9

38. Catala A, Mang B, Wallis L, Huber L. Dogs demonstrate perspective taking

based on geometrical gaze following in a Guesser–Knower task. Anim Cogn.

2017;20: 581–589. doi:10.1007/s10071-017-1082-x

39. Bunford N, Andics A, Kis A, Miklósi Á, Gácsi M. Canis familiaris As a Model for

Non-Invasive Comparative Neuroscience. Trends Neurosci. 2017;40: 438–452.

doi:10.1016/J.TINS.2017.05.003

40. Szabó D, Gábor A, Gácsi M, Faragó T, Kubinyi E, Miklósi Á, et al. On the Face

of It: No Differential Sensitivity to Internal Facial Features in the Dog Brain.

Front Behav Neurosci. 2020;14. doi:10.3389/fnbeh.2020.00025

41. Dilks DD, Cook PF, Weiller SK, Berns HP, Spivak M, Berns GS. Awake fMRI

reveals a specialized region in dog temporal cortex for face processing. PeerJ.

2015; 3:e1115. doi:10.7717/peerj.1115

42. Thompkins AM, Ramaiahgari B, Zhao S, Gotoor SSR, Waggoner P, Denney

TS, et al. Separate brain areas for processing human and dog faces as

revealed by awake fMRI in dogs (Canis familiaris). Learn Behav. 2018;46:

561–573. doi:10.3758/s13420-018-0352-z

43. Cuaya L V., Hernández-Pérez R, Concha L. Our faces in the dog's brain:

Functional imaging reveals temporal cortex activation during perception of

human faces. PLoS One. 2016;11: e0149431.

doi:10.1371/journal.pone.0149431

44. Siniscalchi M, Lusito R, Vallortigara G, Quaranta A. Seeing Left- or Right-

Asymmetric Tail Wagging Produces Different Emotional Responses in Dogs.

Curr Biol. 2013;23: 2279–2282. doi:10.1016/J.CUB.2013.09.027

45. Duranton C, Range F, Virányi Z. Do pet dogs (Canis familiaris) follow ostensive

and non-ostensive human gaze to distant space and to objects? R Soc Open

Sci. 2017;4: 170349. doi:10.1098/rsos.170349

46. Topál J, Kis A, Oláh K. Dogs' Sensitivity to Human Ostensive Cues: A Unique

Adaptation? The Social Dog: Behavior and Cognition. Elsevier Inc.; 2014. pp.

704   319–346. doi:10.1016/B978-0-12-407818-5.00011-5

705 47. Bräuer J, Kaminski J, Riedel J, Call J, Tomasello M. Making inferences about

706   the location of hidden food: Social dog, causal ape. J Comp Psychol.

707   2006;120: 38–47. doi:10.1037/0735-7036.120.1.38

708 48. Fitch WT, Huber L, Bugnyar T. Social Cognition and the Evolution of

709   Language: Constructing Cognitive Phylogenies. Neuron. 2010;65: 795–814.

710   doi:10.1016/J.NEURON.2010.03.011

711 49. Boch M, Karl S, Sladky R, Huber L, Lamm C, Wagner IC. Tailored

712   haemodynamic response function increases detection power of fMRI in awake

713   dogs (Canis familiaris). Neuroimage. 2021;224: 117414.

714   doi:10.1016/j.neuroimage.2020.117414

715 50. Karl S, Boch M, Virányi Z, Lamm C, Huber L. Training pet dogs for eye-tracking

716   and awake fMRI. Behav Res Methods. 2020;52. doi:10.3758/s13428-019-

717   01281-7

718 51. Nitzsche B, Boltze J, Ludewig E, Flegel T, Schmidt MJ, Seeger J, et al. A

719   stereotaxic breed-averaged, symmetric T2w canine brain atlas including

720   detailed morphological and volumetrical data sets. Neuroimage. 2019;187: 93–

721   103. doi:10.1016/j.neuroimage.2018.01.066

722 52. Czeibert K, Andics A, Petneházy Ö, Kubinyi E. A detailed canine brain label

723   map for neuroimaging analysis. Biol Futur. 2019;70: 112–120.

724   doi:10.1556/019.70.2019.14

725 53. Shattuck DW, Mirza M, Adisetiyo V, Hojatkashani C, Salamon G, Narr KL, et al.

726   Construction of a 3D probabilistic atlas of human cortical structures.

727   Neuroimage. 2008;39: 1064–1080. doi:10.1016/j.neuroimage.2007.09.031

728 54. Fedorenko E, Hsieh P-J, Nieto-Castañón A, Whitfield-Gabrieli S, Kanwisher N.

729   New method for fMRI investigations of language: Defining ROIs functionally in

730   individual subjects. J Neurophysiol. 2010;104: 1177–1194.

731   doi:10.1152/jn.00032.2010

732 55. Poldrack RA. Region of interest analysis for fMRI. Soc Cogn Affect Neurosci.

733   2007;2: 67–70. doi:10.1093/scan/nsm006

734 56. Saxe R, Brett M, Kanwisher N. Divide and conquer: A defense of functional

735   localizers. Neuroimage. 2006;30: 1088–1096.

736   doi:10.1016/J.NEUROIMAGE.2005.12.062

737 57. Julian JB, Fedorenko E, Webster J, Kanwisher N. An algorithmic method for

738   functionally defining regions of interest in the ventral visual pathway.

Neuroimage. 2012;60: 2357–2364. doi:10.1016/j.neuroimage.2012.02.055

58.   Karl S, Boch M, Zamansky A, van der Linden D, Wagner IC, Völter CJ, et al.

Exploring the dog–human relationship by combining fMRI, eye-tracking and

behavioural measures. Sci Rep. 2020;10: 22273. doi:10.1038/s41598-020-

79247-5

59.   Thompkins AM, Lazarowski L, Ramaiahgari B, Gotoor SSR, Waggoner P,

Denney TS, et al. Dog–human social relationship: representation of human

face familiarity and emotions in the dog brain. Anim Cogn. 2021;24: 251–266.

doi:10.1007/s10071-021-01475-7

60.   Cohen MA, Dilks DD, Koldewyn K, Weigelt S, Feather J, Kell AJE, et al.

Representational similarity precedes category selectivity in the developing

ventral visual pathway. Neuroimage. 2019;197: 565–574.

doi:10.1016/j.neuroimage.2019.05.010

61.   Kriegeskorte N, Kievit RA. Representational geometry: integrating cognition,

computation, and the brain. Trends Cogn Sci. 2013;17: 401–412.

doi:10.1016/J.TICS.2013.06.007

62.   Kriegeskorte N, Mur M, Bandettini P. Representational similarity analysis -

connecting the branches of systems neuroscience. Front Syst Neurosci.

2008;2. doi:10.3389/neuro.06.004.2008

63.   Kriegeskorte N, Mur M, Ruff DA, Kiani R, Bodurka J, Esteky H, et al. Matching

Categorical Object Representations in Inferior Temporal Cortex of Man and

Monkey. Neuron. 2008;60: 1126–1141. doi:10.1016/J.NEURON.2008.10.043

64.   Rizzolatti G, Fogassi L, Gallese V. Neurophysiological mechanisms underlying

the understanding and imitation of action. Nat Rev Neurosci. 2001;2: 661–670.

doi:10.1038/35090060

65.   Hamilton AF de C, Grafton ST. Goal Representation in Human Anterior

Intraparietal Sulcus. J Neurosci. 2006;26: 1133–1137.

doi:10.1523/JNEUROSCI.4551-05.2006

66.   Kilner JM, Neal A, Weiskopf N, Friston KJ, Frith CD. Evidence of Mirror

Neurons in Human Inferior Frontal Gyrus. J Neurosci. 2009;29: 10153–10159.

doi:10.1523/JNEUROSCI.2668-09.2009

67.   Törnqvist H, Somppi S, Kujala M V., Vainio O. Observing animals and humans:

Dogs target their gaze to the biological information in natural scenes. PeerJ.

2020;8: e10341. doi:10.7717/peerj.10341

68.   Törnqvist H, Somppi S, Koskela A, Krause CM, Vainio O, Kujala M V.

Comparison of dogs and humans in visual scanning of social interaction. R Soc
Open Sci. 2015;2: 150341. doi:10.1098/rsos.150341

69. Spriet C, Abassi E, Hochmann J-R, Papeo L. Visual object categorization in
infancy. bioRxiv. 2021; 2021.02.25.432436. doi:10.1101/2021.02.25.432436

70. Buttelmann D, Tomasello M. Can domestic dogs (Canis familiaris) use
referential emotional expressions to locate hidden food? Anim Cogn. 2013.
doi:10.1007/s10071-012-0560-4

71. Nagasawa M, Murai K, Mogi K, Kikusui T. Dogs can discriminate human
smiling faces from blank expressions. Anim Cogn. 2011;14: 525–533.
doi:10.1007/s10071-011-0386-5

72. Barber ALA, Randi D, Müller CA, Huber L. The Processing of Human
Emotional Faces by Pet and Lab Dogs: Evidence for Lateralization and
Experience Effects. Guo K, editor. PLoS One. 2016;11: e0152393.
doi:10.1371/journal.pone.0152393

73. Racca A, Guo K, Meints K, Mills DS. Reading Faces: Differential Lateral Gaze
Bias in Processing Canine and Human Facial Expressions in Dogs and 4-Year-
Old Children. PLoS One. 2012;7: e36076. doi:10.1371/journal.pone.0036076

74. Range F, Viranyi Z, Huber L. Selective Imitation in Domestic Dogs. Curr Biol.
2007;17: 868–872. doi:10.1016/J.CUB.2007.04.026

75. Soproni K, Miklósi Á, Topál J, Csányi V. Dogs' (Canis familiaris)
Responsiveness to Human Pointing Gestures. J Comp Psychol. 2002;116: 27–
34. doi:10.1037//0735-7036.116.1.27

76. Bray EE, Gnanadesikan GE, Horschler DJ, Levy KM, Kennedy BS, Famula TR,
et al. Early-emerging and highly heritable sensitivity to human communication
in dogs. Curr Biol. 2021;31: 1–5. doi:10.1016/j.cub.2021.04.055

77. Karl S, Sladky R, Lamm C, Huber L. Neural Responses of Pet Dogs
Witnessing their caregiver's Positive Interactions with a Conspecific: An fMRI
Study. Cereb Cortex Commun. 2021;tgab047. doi:10.1093/TEXCOM/TGAB047

78. Correia-Caeiro C, Guo K, Mills D. Bodily emotional expressions are a primary
source of information for dogs, but not for humans. Anim Cogn. 2021;1: 3.
doi:10.1007/s10071-021-01471-x

79. Salomons H, Smith KCM, Callahan-Beckel M, Callahan M, Levy K, Kennedy
BS, et al. Cooperative Communication with Humans Evolved to Emerge Early
in Domestic Dogs. Curr Biol. 2021;31: 3137-3144.e11.
doi:10.1016/J.CUB.2021.06.051

80.   Kaminski J. Domestic dogs: Born human whisperers. Curr Biol. 2021;31:
810   R891–R893. doi:10.1016/J.CUB.2021.05.063

811   81.   Siniscalchi M, D'Ingeo S, Minunno M, Quaranta A. Communication in Dogs.
812   Animals. MDPI AG; 2018. p. 131. doi:10.3390/ani8080131

813   82.   Long RA, Donovan TM, Mackay P, Zielinski WJ, Buzas JS. Effectiveness of
814   Scat Detection Dogs for Detecting Forest Carnivores. J Wildl Manage.
815   2007;71: 2007–2017. doi:10.2193/2006-230

816   83.   Jamieson LTJ, Baxter GS, Murray PJ. Identifying suitable detection dogs.
817   Applied Animal Behaviour Science. Elsevier B.V.; 2017. pp. 1–7.
818   doi:10.1016/j.applanim.2017.06.010

819   84.   Grimm-Seyfarth A, Harms W, Berger A. Detection dogs in nature conservation:
820   A database on their world-wide deployment with a review on breeds used and
821   their performance compared to other methods. Methods in Ecology and
822   Evolution. British Ecological Society; 2021. pp. 568–579. doi:10.1111/2041-
823   210X.13560

824   85.   Dunn M, Degenhardt L. The use of drug detection dogs in Sydney, Australia.
825   Drug Alcohol Rev. 2009;28: 658–662. doi:10.1111/j.1465-3362.2009.00065.x

826   86.   Guest C, Otto CM. Editorial: Canine Olfactory Detection. Frontiers in Veterinary
827   Science. Frontiers Media S.A.; 2020. p. 100. doi:10.3389/fvets.2020.00100

828   87.   Berns GS, Brooks AM, Spivak M. Scent of the familiar: An fMRI study of canine
829   brain responses to familiar and unfamiliar human and dog odors. Behav
830   Processes. 2015;110: 37–46. doi:10.1016/J.BEPROC.2014.02.011

831   88.   Prichard A, Chhibber R, Athanassiades K, Spivak M, Berns GS. Fast neural
832   learning in dogs: A multimodal sensory fMRI study. Sci Rep. 2018;8: 14614.
833   doi:10.1038/s41598-018-32990-2

834   89.   Jia H, Pustovyy OM, Wang Y, Waggoner P, Beyers RJ, Schumacher J, et al.
835   Enhancement of Odor-Induced Activity in the Canine Brain by Zinc
836   Nanoparticles: A Functional MRI Study in Fully Unrestrained Conscious Dogs.
837   Chem Senses. 2016;41: 53–67. doi:10.1093/chemse/bjv054

838   90.   Jia H, Pustovyy OM, Waggoner P, Beyers RJ, Schumacher J, Wildey C, et al.
839   Functional MRI of the Olfactory System in Conscious Dogs. Dhenain M, editor.
840   PLoS One. 2014;9: e86362. doi:10.1371/journal.pone.0086362

841   91.   Ramaihgari B, Pustovyy OM, Waggoner P, Beyers RJ, Wildey C, Morrison E,
842   et al. Zinc nanoparticles enhance brain connectivity in the canine olfactory
843   network: Evidence from an fMRI study in unrestrained awake dogs. Front Vet

Sci. 2018;5: 127. doi:10.3389/fvets.2018.00127

92.   Bräuer J, Blasi D. Dogs display owner-specific expectations based on olfaction.

Sci Reports 2021 111. 2021;11: 1–10. doi:10.1038/s41598-021-82952-4

93.   Hecht EE, Smaers JB, Dunn WJ, Kent M, Preuss TM, Gutman DA. Significant

Neuroanatomical Variation Among Domestic Dog Breeds. J Neurosci. 2019;39:

0303–19. doi:10.1523/JNEUROSCI.0303-19.2019

94.   Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving

Bioscience Research Reporting: The ARRIVE Guidelines for Reporting Animal

Research. PLoS Biol. 2010;8: e1000412. doi:10.1371/journal.pbio.1000412

95.   Peirce JW. PsychoPy-Psychophysics software in Python. J Neurosci Methods.

2007;162: 8–13. doi:10.1016/j.jneumeth.2006.11.017

96.   Wagner IC, van Buuren M, Bovy L, Fernández G. Parallel engagement of

regions associated with encoding and later retrieval forms durable memories. J

Neurosci. 2016;36: 7985–7995. doi:10.1523/JNEUROSCI.0830-16.2016

97.   R Core Team. R: A Language and Environment for Statistical Computing.

Vienna, Austria: R Foundation for Statistical Computing; 2020. Available:

https://www.r-project.org/

98.   Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, et al. User-

guided 3D active contour segmentation of anatomical structures: Significantly

improved efficiency and reliability. Neuroimage. 2006;31: 1116–1128.

doi:10.1016/J.NEUROIMAGE.2006.01.015

99.   Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but

systematic correlations in functional connectivity MRI networks arise from

subject motion. Neuroimage. 2012;59: 2142–2154.

doi:10.1016/J.NEUROIMAGE.2011.10.018

100. Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE.

Methods to detect, characterize, and remove motion artifact in resting state

fMRI. Neuroimage. 2014;84: 320–341. doi:10.1016/j.neuroimage.2013.08.048

101. Duff EP, Cunnington R, Egan GF. REX: Response Exploration for

Neuroimaging Datasets. Neuroinformatics 2007 54. 2007;5: 223–234.

doi:10.1007/S12021-007-9001-Y

102. Mumford JA, Turner BO, Ashby FG, Poldrack RA. Deconvolving BOLD

activation in event-related designs for multivoxel pattern classification

analyses. Neuroimage. 2012;59: 2636–2643.

doi:10.1016/j.neuroimage.2011.08.076

103. Kriegeskorte N, Goebel R, Bandettini P. Information-based functional brain
mapping. Proc Natl Acad Sci U S A. 2006;103: 3863–3868.
doi:10.1073/pnas.0600244103

104. Nichols TE, Holmes AP. Non-parametric procedures. Statistical Parametric
Mapping: The analysis of Functional Brain Images. Amsterdam: Elsevier; 2007.
pp. 253–272.

105. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York; 2016. Available: https://ggplot2.tidyverse.org

106. Allen M, Poggiali D, Whitaker K, Marshall TR, Kievit RA. Raincloud plots: A
multi-platform tool for robust data visualization [version 1; peer review: 2
approved]. Wellcome Open Res. 2019;4.
doi:10.12688/wellcomeopenres.15191.1

107. Notter MP, Gale D, Herholz P, Markello R, Notter-Bielser M-L, Whitaker K.
AtlasReader: A Python package to generate coordinate tables, region labels,
and informative figures from statistical MRI images. J Open Source Softw.
2019;4: 1257. doi:10.21105/JOSS.01257
895