Effects of climate warming on the pine processionary moth at the

southern edge of its range: a retrospective analysis on egg survival

in Tunisia

Asma BOUROUGAAOUI[1,2,3], Christelle ROBINET[3], Mohamed L. BEN JAMAA[1], Mathieu LAPARIE[3]

[1] Université de Carthage, Institut National de Recherches en Génie Rural, Eaux et Forêts-Laboratoire

de Gestion et de Valorisation des Ressources Forestières, Ariana, Tunisia

[2] Université de Carthage, Institut National Agronomique de Tunis, Tunis, Tunisia

[3] INRAE, URZF, 45075, Orléans, France

* Corresponding author: **Mathieu Laparie, mathieu.laparie@inrae.fr**

Journal: PCI Ecology

Type of paper: Research Article

Number of figures: 6

Number of Tables: 3

Supplementary Material: ~~6~~9

Running title: **Effects of climate warming on eggs of the pine processionary moth in Tunisia**

Abstract

In recent years, ectotherm species have largely been impacted by extreme climate events, essentially heatwaves. In Tunisia, the pine processionary moth (PPM), *Thaumetopoea pityocampa*, is a highly damaging pine defoliator, which typically lays eggs in summer. Its geographical range is expanding northwards in Europe while retracting from South Tunisia where summer temperatures can reach extremely high values. In this study, we aimed at exploring the effects of climate change on this species at its southern range edge. We investigated variations of fecundity and causes of egg mortality over time using historical and contemporary collections of egg masses from different Tunisian sites to seek relationships with regional climate change over three decades (1990-2019). Our results suggest negative effects of summer heat on egg survival, reflected in a decrease of hatching rate down to 0% in one site during a heatwave. Such a high hatching failure was found to result from both high egg sterility (our results did not allow distinguishing impeded mating success from failed egg maturation or early death of the embryo) and increased abortion of more developed embryos, but little effects of parasitism rate, thereby suggesting vulnerability to heat during embryonic development. We also observed decreasing female fecundity (*i.e.*, number of eggs laid per female) in regions where data were available both in the 1990s and the 2010s, which was associated with a decrease in parasitism rate, while the climatic variability increased. This study investigated direct hatching failure in nature that may be related to the magnitude of warming in summer. Previous studies have confirmed the thermal sensitivity of early instars of the PPM to temperatures observed in the present work, including one population from South Tunisia. However, further work is required to evaluate the relative importance of warming summers among populations because the risk of heat stress depends on the phenology of sensitive instars, and populations from the warmest areas may not necessarily be the most vulnerable to climate change if they already evolved phenological heat avoidance. In addition to heat-induced mortality, the ultimate fitness of individuals that survive challenging heat stresses during early developmental stages should also be explored to determine potential carry-over effects on subsequent life stages.

Keywords

Egg mass, heatwave, pine processionary moth, pinus, *Thaumetopoea pityocampa*, Tunisia

Introduction

During the period 1901-2010, land temperature has risen by 1.12 and 0.84°C in the Northern and Southern hemispheres, respectively (Jones et al., 2012). Climate warming~~This average climate warming already impacted phenology and distribution in many plant and ectotherm species~~ (Parmesan & Yohe, 2003) may induce heritable as well as plastic changes in a wide range of traits at multiple phenotypic levels (Pigliucci, 2001, 2005), such as (i) the timing of life cycle events such as flowering, migration and egg laying (Charmantier & Gienapp, 2014; Chuine, 2010; Chuine et al., 2013; Liu et al., 1995), (ii) the behaviour such as locomotor activity or movements to seek suitable microclimates (Pincebourde et al., 2021; Pincebourde & Woods, 2020; Woods et al., 2015), the morphology such as body size (Gardner et al., 2011; Ghosh et al., 2013; Sheridan & Bickford, 2011; Wu et al., 2019), or (iii) the physiology such as thermal acclimation or adaptation to thermal stressors. Those changes may ultimately translate into survival and in turn alter genetic frequencies, population density in given habitats, as well as distribution (Root et al., 2003). In the twentieth century, a wide range of taxa ranging from invertebrates to mammals and from grasses to trees have shifted their ranges poleward, upslope or both (Crozier, 2004; Hickling et al., 2005; Karban & Strauss, 2004; Parmesan et al., 1999; Parmesan & Yohe, 2003; Root et al., 2003; Walther et al., 2002). Climate change is also expected to increase temperature variability and, in turn, increase the magnitude and frequency of stochastic extreme thermal events, which is already increasingly observed over most parts of the world (Allen et al., 2012; Coumou & Rahmstorf, 2012; Fischer & Schär, 2010), particularly northern Africa (Fontaine et al., 2013; Nangombe et al., 2019; Zittis et al., 2021). In Tunisia, the mean average temperature has risen by about 1.4°C since 1901, with a remarkable increase in the last 30 years with an average warming of 0.4°C per decade, and most substantial warming has been observed during summer in southern regions where average maximum temperatures can often exceed 40°C (Verner et al., 2013). Together with average warming, increasing thermal fluctuations and extreme events may impact organismal fitness (Kingsolver et al., 2013; Thompson et al., 2013), and the~~their~~ combination of these is expected to dramatically impact survival, distribution and life cycles of many organisms.

The pine processionary moth (hereafter referred to as PPM), *Thaumetopoea pityocampa* (Denis & Schiffermüller, 1776) (Lepidoptera, Notodontidae), is a highly damaging pest of pine forests across the circum-Mediterranean region (Carus, 2009; Démolin, 1969; Jacquet et al., 2013; Sbay & Zas, 2018). The geographic range of the PPM extends from northern Africa to southern Europe, from the Atlantic coast to the western part of Turkey (EPPO, 2004; Roques, 2015). The PPM is a well-documented ~~Lepidopteran~~ insect that has been acknowledged by the Intergovernmental Panel on Climate Change (IPCC) as one of the few species for which the causal relationship between climate

warming and range expansion has been thoroughly proven (Battisti et al., 2005; Rosenzweig et al., 2007). The distribution range remained relatively steady until the late 1990s but then expanded towards higher latitudes and elevations in southern Europe. Indeed, warming winter temperatures have facilitated feeding in this winter-developing species and thus indirectly contributed to improving survival rate and growth rate in newly colonized areas (Battisti et al., 2005, Robinet et al., 2007).

Contrary to the beneficial effects of climate change demonstrated near the northern distribution edge of the PPM, adverse effects of climate change have been observed on the southern range edge (North Africa). Range retraction has been described in southern Tunisia and was found to result from increasing mortality rates of early life stages in a translocation experiment along a natural thermal gradient, which could be ascribed to local effects of climate warming (Bourougaaoui et al., 2021). Fecundity, hatching rate and predation at the egg stage (mostly from parasitoids) presumably play an important role in the PPM because this species is gregarious. Several studies have emphasized how larval performance depends on the realised group size, i.e. the number of neonates, and ultimately the survival of the whole colony until the end of larval growth (Clark & Faeth, 1997; Denno & Benrey, 1997; Ronnås et al., 2010). Colony density has been suggested to influence feeding activity and feeding efficiency of individual larvae, which is particularly critical in early stages when individuals have little desiccation and starvation resistances. The number of larvae was also shown to impact silk weaving activity to build and maintain the nests that shelter larvae during the day until their pupation in spring (Démolin, 1965; Martin, 2005). As a result, ~~Authors found that~~ the number of surviving tents and the average proportion of living larvae per tent were positively correlated to colony size (Pérez-Contreras et al., 2003; Roques et al., 2015). Focusing on the main drivers of colony size in early development is therefore of key importance to understand distribution changes and responses to climate change.

In Tunisia, the life cycle of PPM is generally univoltine, however it can extend over two years at high altitudes due to prolonged diapause in a fraction of the pupae (Roques, 2015). Flight periods are poorly documented, nonetheless a study conducted by Démolin and Rive in 1968 in high and medium latitudes, revealed that most individuals fly in the second half of July at high elevations and August to September at mid elevations (Ben Jamâa & Jerraya, 1999; Démolin & Rive, 1968). Due to the short lifespan of adults, egg laying occurs immediately after adult flights, and eggs and neonate larvae are presumably the instars that are most likely exposed to acute heat during the whole life cycle. Understanding the effects of warming on female fecundity, egg survival and egg parasitoids is crucial to explore the overall effects of climate change of this species at its southern range edge where warming is known to be of great magnitude.

In this study, we explored how climate warming over the last three decades may have impacted
egg survival and hatching rate in Tunisia. To address this question, we combined historical and
contemporary collections of egg masses originating from different Tunisian localities in the 1990s
(1992, 1993, and 1995) and in the 2010s (2010, 2014, 2017, 2018, and 2019). Egg phenotypes and
survival rate were investigated with regard to regional climatic features and contrasts analyzed from
30-year climatic data series across Tunisia. A cornerstone of this study is the identification of climate
regions computed from multiple meteorological series, which allows comparing eggs from multiple
sites within statistically consistent climates, instead of using arbitrary groups such as administrative
regions. A grouping method was mandatory to analyze the long term data available on PPM eggs
because exact sampling sites have changed over the years.

Materials and Methods

1   Historical data (1992-2014) and egg sampling done for this study (2017-2019)

A total of 755 egg masses from historical datasets and recent collections were analyzed in this
study. Egg masses originated from 22 sites distributed across the PPM distribution in Tunisia (Table 1;
Figure 1; Table SM1). Historical datasets on egg masses collected in 1992, 1993, 1995, 2010 and 2014
on Aleppo pine stands, *Pinus halepensis* Miller, were retrieved from institutional reports
(unpublished data, INRGREF). These datasets report the length of egg masses, the number of eggs
per egg mass, and the phenotype of individual eggs (parasitized, aborted, sterile, hatched). In
addition, we collected egg masses in various locations in 2017, 2018, and 2019. All these egg masses
were also collected on Aleppo pine stands, before hatching but as late as possible in each region to
ensure eggs were exposed to natural conditions, and then and kept at ambient temperature (25 ±
2°C) at the INRGREF laboratory near Tunis where we followed a protocol similar to that used for
historical collections of egg masses. Egg masses were kept individually in test tubes capped with
cotton to allow ventilation. Egg hatching was checked daily. After a period of at least 40 days with no
additional hatching, the protective scales that cover PPM egg masses were removed to observe
individual eggs under a binocular magnifier and collect data similar to that available in historical
datasets: length of egg masses, number of eggs per egg mass (fecundity), and egg phenotype. First,
hatched eggs were distinguished from unhatched eggs based on the presence of the characteristic
large jagged exit hole from which the neonate left the egg, and an empty transparent shell. Then,
unhatched eggs were dissected to assess the cause of mortality (parasitized, *i.e.*, eggs with a small
parasitoid exit hole and/or containing a dead parasitoid and/or containing parasitoid meconium;
aborted, *i.e.*, dead embryo or dead PPM larva; and sterile, *i.e.*, undeveloped egg with dried-up yolk)

(Imbert, 2012). Parasitism rate was calculated taking into account both emerged parasitoids found in

the test tubes and dead ones found inside unhatched eggs.

Table 1 . Collection of PPM egg masses in Tunisia (see Table SM1 for coordinates of the sites).
Calculated climate clusters are indicated to represent the amount of data available per cluster.

Site	Year of collection	Number of egg masses	Cluster	Nearest meteorological station (within 100 km and 350 m in elevation); distance
Sejnane	1995	20	1	Béja; 51.0 km
Testour	2014	12	1	Béja; 26.3 km
Ain Jamala	2010	15	1	Béja; 26.5 km
El Krib	2010	18	1	Kef; 38.7 km
Henchir Naam	1992	53	1	Kef; 41.5 km
Jebel Kbouch	1993	56	1	Kef; 21.3 km
El Ayoun	1993	30	1	Kasserine; 73.2 km
Chaambi	1995	27	1	Kasserine; 30.3 km
	2014	11		
Thélepte	2017	43	1	Kasserine; 1.9 km
	2019	30		
Jebel Sidi Aich	2014	31	1	Kasserine; 32.7 km
Kesra	2010	18	1	Siliana; 20.8 km
Mghila	2014	51	1	Kasserine ; 71.6 km
Bir Lahfay	2014	12	2	Sidi Bouzid; 32.5 km
Jebel Motlag	2017	38	2	Sidi Bouzid; 20.6 km
	2018	29		
Jebel Rihane	2017	25	2	Sidi Bouzid; 25.7 km
El Fej	2017	18	2	Gafsa; 36.6 km
El Menzeh	1992	19	3	Carthage; 3.2 km
	1993	57		
	2014	10		
Cité Ettahrir	2014	21	3	Carthage; 7.9 km
Manouba	2010	15	3	Carthage; 10.5 km
Korbous	1992	30	3	Carthage; 32.5 km
	2010	15		
Dar Chichou	1995	20	4	Kelibia; 17.1 km
Oued Laabid	1995	31	4	Kelibia; 33.3 km

Figure 1. Location of egg sampling sites and meteorological stations, with associated calculated climate cluster.

2   Climate data

We used series of daily temperatures recorded (by the Institut National de Météorologie, INM, Tunis, Tunisia) in eight meteorological stations distributed within the PPM range in Tunisia (Fig. 1; Table SM2). To fill missing data in INM time series, satellite measurement of daily temperatures were also retrieved from the NASA Prediction of Worldwide Energy Resources website (https://power.larc.nasa.gov/data-access-viewer/) on the grid cells of 0.5 degree × 0.625 degree (~ 50.8 km × 60.8 km) matching the location of INM weather stations (Table SM2). The similarity of both sources of data was evaluated using Pearson correlations tests for daily maximal and daily minimal temperatures in Tunis, where the data series from INM since 1990 was the most comprehensive. Daily maximal and minimal temperatures from both data sources were found to be strongly correlated (Pearson tests, r= 0.95, *p* < 0.001 and r = 0.94, *p* < 0.001, respectively). The two types of datasets where therefore combined in case of missing data in other INM series to reconstruct uninterrupted series for the period 1990-2019 (Table SM2).

Each site of egg sampling was assigned to the nearest meteorological station (< 100 km in all

cases) among those situated at an elevation within 350 meters of the egg site, an arbitrary threshold

we chose to mitigate potential climatic differences along elevation gradients (Table 1, Figure SM3).

To better understand climatic features in each of the eight meteorological series (Table SM2), (i)

the normal daily temperatures with seasonal contrasts over the period, as well as (ii) the overall

trend since 1990, were calculated. For (i), we averaged 30 years of daily maxima (TX) and minima

(TN) by day of the year, and calculated the likelihood for each day of temperatures below 0 or above

32 and 40°C, which have been suggested by (Démolin, 1969) and Huchon & Démolin (1970) as pivotal

thresholds for phenological strategies and survival in the PPM (see also discussion in Robinet et

al.Huchon & Démolin (1969) as pivotal thresholds for phenological strategies and survival in the PPM

(see also discussion in Robinet et al. 2015). For (ii), daily TX and TN were averaged per year and

represented along the 30 years of data, together with the total number of days below 0 or above 32

and 40°C. Those per-station climate summaries are provided in SM4.

3 Statistical analyses

Climate clusters. The unbalanced egg sampling design throughout historical data and recent

collections prevents allochronic comparisons of egg phenotypes within individual sampling sites.

Therefore, we investigated climatic similarities and dissimilarities among meteorological series in

order to identify regional climate clusters within which multiple meteorological series and associated

egg sampling sites could be statistically grouped together. Climate-based grouping appeared more

relevant and less arbitrary than using administrative regions because of the heterogeneous

landscape and overall size of some regions. To do so, the monthly averages of TN and TX were

calculated in each meteorological series over the period 1990-2019, resulting in a set of 24 variables

(2 × 12 months) and 30 values per series (30 years). A Principal Component Analysis (PCA) was used

on the covariance matrix of those variables to project the 30 years of data from each of the eight

meteorological locations and better visualize their intra- and inter-group variance on reduced

dimensionality. The resulting multivariate object then fed a K-medoid clustering analysis using the

PAM method (Partitioning Around Medoids, see Reynolds et al. 2006, Schubert and Rousseeuw 2019)

to identify relevant climate clusters (listed in Table1). The PCA could be performed on unscaled

temperature variables since they were all measured in the same unit (covariance PCA), thereby

giving most weight to summer months due to generally higher temperature values, without

neglecting other months in the overall variance structure. As a consequence, the climate clusters

identified using all four seasons are mostly influenced by the season eggs are exposed to (roughly

June to September). Details of cluster assignation to individual points in each meteorological series

are detailed in SM5. Monthly means of TN and TX of the medoid of each cluster, *i.e.,* the individual
point that best represents its cluster due to low average dissimilarity to all other points, are
represented in SM6.

Interannual fluctuation of maximal summer temperature within clusters. To explore regional
warming trends to which eggs are subjected within clusters over 1990-2019, the monthly means of
daily maximal temperatures from meteorological series within each cluster were calculated from
June to September. A linear model was then built for each cluster and each month to plot regressions
over time and determine the slope for each cluster. The adequacy of residuals to Normality was
checked using QQ plots. ~~Finally, the four months were averaged together per cluster and period to~~
~~obtain the mean maximal summer temperature in each cluster and period. Then, a two-way ANOVA~~
~~followed by a Tukey *post-hoc* procedure was performed to test the effects of cluster and period on~~
~~the maximal summer temperature.~~

Egg phenotype comparisons. Egg phenotype variables did not meet assumptions of
homoscedasticity and normality for parametric tests, therefore we used the non-parametric test
(RANCOVA) proposed by Quade (1967) to compare eggs sampled within each of the four clusters in
the 1990s or in the 2010s. First, the response variables (Fecundity, Hatching, Sterility, Abortion and
Parasitism rates<u>, Clutch length was discarded due to its high correlation and redundancy with</u>
<u>Fecundity</u>) and the covariate (monthly means of TX averaged from June to September per year per
cluster) were ranked separately. Second, residuals from the respective linear regression of each
ranked response variable on the ranked covariate were calculated. Third, the effects of grouping
<u>factor(s</u>~~factors period (1990s vs 2010s) and Cluster (1, 2, 3, 4)~~) on residuals were investigated for each
response variable using the Quade's RANCOVA. <u>Since</u>~~When a significant interaction was found, we~~
~~reported~~ the <u>dataset is unbalanced, with both past and recent samples</u>~~pairwise comparisons~~ in <u>only</u>
<u>two of the four climate</u>~~dependent variable between~~ clusters<u>, we conducted two separate Quade's</u>
<u>RANCOVA analyses:</u> ~~at each period level, as well as between period levels for each cluster using~~
~~pairwise t-tests with Bonferroni corrections for multiple testing. When no significant interaction was~~
~~found between period and cluster, a Tukey post hoc procedure was used to investigate pairwise~~
~~differences between clusters.~~

- <u>Analysis (1) was conducted on a subset of the data containing only clusters 1 and 3 (where</u>
<u>past and recent samples exist), in which we included both Cluster and Period factors as well</u>
<u>as their interaction. When the interaction was significant, pairwise comparisons were</u>
<u>performed using pairwise t-tests and Bonferroni correction.</u>
- <u>Analysis (2) synchronically compared clusters within each period subset, i.e., all clusters were</u>
<u>considered but compared only to other clusters during the period(s) they shared. A Tukey</u>

post hoc procedure was used to investigate pairwise differences when a main effect was

found.

Results

1   Climate clusters

The first plane (PC1 × PC2) of the PCA performed on climatic data from all eight meteorological

series based on monthly averages of TN and TX each year (n = 8 × 30 = 240 data points) accounted

for 71.28% of the total inertia (Fig. 2). PAM clustering on the PCA scores indicated four relevant

groups with little overlapping (Fig. 2). Cluster 1 grouped Kef, Kasserine, Siliana and Béja together,

cluster 2 grouped Sidi Bouzid and Gafsa together, while cluster 3 and cluster 4 corresponded to single

meteorological series, Carthage and Kélibia, respectively. Depending on the meteorological series,

between 76.67 and 100% of data points (years) were correctly assigned to their cluster (SM5).

Figure 2. PCA scores for each year of data from the eight meteorological series (INM and NASA series, see table SM2) and 24 TN monthly average and TX monthly average variables. Each of the eight series are identified with different symbol and colour combinations. PAM clustering results are overlaid on the PCA scores with 95% confidence ellipses and different symbols for different clusters. Open points correspond to the centroids of each meteorological series, while large thin crosses mark the medoid point of each cluster.

Per-cluster climate reconstructions averaged from daily~~monthly~~ means of TX over years ~~in each~~

~~cluster~~ (Fig. 3) indicated comparatively cold winters and hot summers with a high interseasonal

variability in cluster 1, ~~a shift to~~ warmer winters and summers in cluster 2 with similar interseasonal

variability, no extreme winters or summers and lower interseasonal variability in cluster 3, and the

lowest interseasonal variability with comparatively mild summers in cluster 4. Within-month

variability also appeared to be the highest over the last 30 years in clusters 1 and 2. The probability

to overreach 40°C in summer was found to be the highest in cluster 2, while cluster 4 showed the

lowest probability of overreaching 32°C, with clusters 1 and 3 sitting in between those extremes. July

and August are the warmest months in all clusters (Fig 3, Fig 4). – TN monthly average within each

medoid appeared to roughly reflect TX monthly average across each months of the year (SM6, SM9),

indicating that similar trends can be inferred for per-cluster TN monthly average.

Cluster 1

Cluster 2

Cluster 3

Cluster 4

Mean daily TX over 1990–2019
White quantile lines mark (0, 0.1], (0.1, 0.9]
and (0.9, 1] probability ranges over the period

Figure 3. Ridge lines showing the distribution (kernel density) of daily~~monthly~~ means of TX over 1990-2019 for each cluster. The area of each distribution equals 1 and white vertical lines mark 0.1 and 0.9 quantiles.

~~Averaging TX monthly means (from June to September each year) within clusters and periods (Fig. 4) confirmed significant differences among clusters (ANOVA, Fdf = 275.2453, p < 0.001; significant differences in all pairs of clusters but 1 and 3 (Tukey post hoc, p < 0.0001)). Period did not influence this summary summer variable significantly despite an increasing trend in clusters 2, 3 and 4 (ANOVA, Fdf = 2.5111, Fig. 4), and the interaction term between cluster and period was not significant (ANOVA, Fdf = 2.5111, p = 0.116).~~

Fluctuations of monthly averaged TX during summer over the last three decades (Fig. ~~4~~5) in each cluster indicated that September has not warmed significantly over time (no apparent slopes, non-significant linear regressions). Contrariwise, July got significantly warmer over time, particularly in cluster 2 and 3 (positive slopes, significant linear regressions). Cluster 2 appeared as the cluster with the most striking summer warming, with positive increases of TX monthly average spanning not only July, but also June (non-significant trend (p ~~=~~< 0.11~~1~~), similar to cluster 3) and August (significant increase). Contrariwise, summers in clusters 1 and 4 appeared as the least subject to warming since 1990.

Figure 45. Fluctuations over time of monthly maximal temperature within each cluster, from June to September. The upper and lower lines of each ribbon correspond to 95% confidence intervals.

***Clutch size.*~~Fecundity and hatching rate.~~** The length of egg masses was <u>measured in case the</u> <u>distance between eggs of a clutch would differ among areas or periods, but this variable was found</u> <u>to be</u>~~significantly and~~ highly positively correlated to fecundity (Spearman correlation test, rs = 0.72, *p* < 0.001), <u>thereby leaving little room for variations in the fecundity/length ratio. Therefore,</u>~~therefore~~ we focused further analyses on fecundity only<u>, which</u>~~. Fecundity across clusters and years~~ ranged from 121 ± 8.2 SE to 174.6 ± 5.9 SE <u>among clusters and periods</u> (Figure 5). Analysis (1) conducted on

clusters where both past and present samples have been collected (clusters 1 and 3) showed no significant difference in fecundity between those two 6). Fecundity did not differ significantly between clusters (Quade's RANCOVA, F_{df} = 2.88$_{1363}$, p = 0.09) or 071) and between periods (F_{df} = 3.01$_{1521}$, p = 0.08361). However, there was a significant crossover interaction suggested non-parallel trends over time between them(i.e., the effect of periods on the fecundity is opposite, depending on the value in each cluster) (Quade's RANCOVA, F_{df} = 6.42$_{1411}$, p = 0.012). Indeed, cluster 3 was the only cluster where fecundity changed over time (pairwise t-test, t_{df} = 3.1$_1$, p = 0.002 adjusted by Bonferroni correction), with an average decrease of 16 %. Analysis (2) performed on all clusters but synchronic periods revealed no) suggested that significant differences among clusters 1, 3 and 4 in the 1990s (Quade's RANCOVA, F_{dt} = 1.58$_2$, p = 0.208), whereas it confirmed significant differences among clusters 1, 2can be found in at least one of the two periods, and 3 in the 2010s (F_{df} = 4.40$_2$, p = 0.013).evolved differentially over time. Specifically, no differences were found among clusters in the 1990s, whereas fecundity was significantly higher in cluster 1 than cluster 3 (Tukey post hoc, p = 0.009).in the 2010s (pairwise t test, t_{df} = 3.2$_2$, p = 0.005 adjusted by Bonferroni correction). Cluster 3 was the only cluster where fecundity significantly changed over time (t_{df} = -3.1$_1$, p = 0.002), with a decrease of 16 %.

Figure 56. Fecundity (raw count of eggs), clutch length, hatching rate, sterility rate, abortion rate and parasitism rate per cluster and sampling year. Black markers at the bottom show years when samples have been collected, to distinguish true zeros from missing values. in each egg sampling site and sampling year. Error bars: SEM.

Hatching rate significantly differed between clusters (Quade's RANCOVA, F_{df} = 3.8$_3$, p=0.01) and between periods (F_{df} = 24.9$_1$, p <0.001). As found for fecundity, differences in hatching rate among clusters varied between the 1990s and the 2010s (Quade's RANCOVA, F_{df} = 21.7$_1$, *p* < 0.001). Hatching rate was significantly lower in cluster 3 than cluster 1 (pairwise t-test, t_{df} = -4.7$_2$, p < 0.001 adjusted by Bonferroni correction) and cluster 4 (t_{df} = -3.6$_2$, p = 0.001) in the 1990s, and higher in cluster 2 than cluster 1 in the 2010s (t_{df} = 2.4$_2$, p = 0.05) (Table 2). The mean difference between periods showed that hatching rate in cluster 1 was significantly higher in the past (t_{df} = 6.7$_1$, p < 0.001) (Table 2). Particularly, in 2017, hatching rate dropped to zero in cluster 1, and largely decreased in cluster 2 compared to other years (Figure 6).

Table 2. Descriptive statistics: observed mean (M), Quade's adjusted mean (Madj) and associated standard error (SE) for the different response variables.

Egg mortality factors. Base abortion and parasitism rates differed significantly between clusters 1 and 3 (Analysis (1), Quade's RANCOVA, abortion rate: F_{df} = 17.13$_1$, p < 0.001; parasitism rate: F_{df} = 4.53$_1$, p = 0.034) and periods (abortion rate: F_{df} = 50.92$_1$, p <0.001; parasitism rate: F_{df} = 20.36$_1$, p < 0.001). However, the temporal increase in abortion and decrease in parasitism were similar in the two clusters since no significant interaction was observed (abortion rate: Fdf = 0.08$_1$, p = 0.776;

parasitism rate: $F_{df}= 1.34_1$, p = 0.248). Yet, in 2017, hatching rate dropped to zero in cluster 1 with all

eggs visually scored as sterile. Consequently, abortion and parasitism rates were also the lowest (0 %)

in this cluster that year since those mortality factors can only occur at a later development stage

(Figure 5, Table SM7). Cluster 1 was the only cluster where such a dramatic change was observed,

and where hatching rate changed significantly between the past and recent periods (Analysis (1),

pairwise t-test, $t_{df} = 7.5_1$, p < 0.001 adjusted by Bonferroni correction). This average decrease over

time can be attributed not only to the striking drop in 2017, but also to decreases of lower magnitude

in other recent years (Figure 5). Those changes correlate negatively with the aforementioned trends

in abortion rate and sterility rate. Of the two cluster, the increase in abortion rate was most

prominent in cluster 1 (Figure 5), although it may be underestimated in recent years due to its null

value in 2017 when eggs did not develop enough to score abortion rate. Analysis (2) showed

synchronic differences among clusters in the rates of hatched, sterile, aborted and parasitized eggs

(see Tables 2 and 3 for details).

Finally, to evaluate the influence of the 2017 peculiarity on overall trends in cluster 1 and investigate

other changes that may have been concealed by this heatwave, similar analyses (1) and (2) have

been performed on a subset of the data without cluster 1 in 2017 (see descriptive statistics in SM8).

Analysis (1) revealed that hatching and abortion rates still changed over time in the two clusters, to a

lower extent than when considering data from 2017 in cluster 1 (hatching rate: $F_{df} = 8.2_1$, p = 0.004;

abortion rate: $F_{df} = 7.6_1$, p = 0.006), whereas sterility and parasitism did not change in any of the two

clusters (sterility: $F_{df} = 1_1$, p = 0.315; parasitism: $F_{df} = 0.3_1$, p = 0.563). Analysis (2) revealed different

patterns than those found when considering data from cluster 1 in 2017, since no difference among

clusters appeared in the 2010s in any of the variables (hatching rate: $F_{df} = 0.9_2$, p = 0.412; sterility

rate: $F_{df} = 2.8_2$, p = 0.063; abortion rate: $F_{df} = 0.01_2$, p = 0.993; parasitism rate: $F_{df} = 1.5_2$, p = 0.218).

Table 2: Results of Tukey's post hoc tests in the 1990s.

Variable	Clusters~~Period~~	Mean difference (I-J)~~Cluster~~	Standard error~~M (SE)~~	p value~~Madj (SE)~~
~~Fecundity~~	~~1990s~~	~~1~~	~~156.6 (3)~~	~~19.9 (1.1)~~
		~~2~~	~~NA~~	~~NA~~
		~~3~~	~~159.7 (4.3)~~	~~24.6 (2.1)~~
		~~4~~	~~145.4 (6.3)~~	~~-55.3 (4.3)~~
	~~2010s~~	~~1~~	~~153 (3)~~	~~14 (1)~~
		~~2~~	~~142.1 (4.5)~~	~~-12.8 (1.8)~~
		~~3~~	~~134.5 (5.6)~~	~~-84.2 (3.5)~~
		~~4~~	~~NA~~	~~NA~~
Hatching rate	1 vs 3~~1990s~~	43.5[*1]	11.6~~73.5 (1.7)~~	0.001~~76.8 (1)~~
		~~2~~	~~NA~~	~~NA~~

		~~3~~	~~63.3 (2.5)~~	~~-42.5 (1.7)~~
		~~4~~	~~79.3 (3.1)~~	~~82.2 (3.7)~~
	~~2010s~~	~~1~~	~~46.8 (2.5)~~	~~-59.8 (1)~~
	1 vs 4	-43.9[*2]	15~~48.9 (3.4)~~	0.01~~-4.9 (1.8)~~
	3 vs 4	-87.4[*3]	16~~63.2 (3.6)~~	<0.001~~2.8 (3.3)~~
		~~4~~	~~NA~~	~~NA~~
Sterility rate	1 vs 3~~1990s~~	9.8~~1~~	11~~6.8 (0.6)~~	0.68~~-30.8 (1)~~
		~~2~~	~~NA~~	~~NA~~
	1 vs 4	32.13	15.3~~6.6 (0.7)~~	0.09~~-22.5 (1.7)~~
	3 vs 4	22.34	16.5~~8 (0.7)~~	0.37~~-21.8 (3.2)~~
	~~2010s~~	~~1~~	~~27.1 (2.5)~~	~~46.7 (1.1)~~
		~~2~~	~~17.3 (2.6)~~	~~21.8 (1.9)~~
		~~3~~	~~6.6 (1.7)~~	~~-67.6 (3.2)~~
		~~4~~	~~NA~~	~~NA~~
Abortion rate	1 vs 3~~1990s~~	-42.8[*1]	11.8~~2.4 (0.2)~~	-96.2 ~~(0.~~001~~8)~~
		~~2~~	~~NA~~	~~NA~~
	1 vs 4	-13.93	15~~9.3 (1.9)~~	0.64~~22.2 (1.8)~~
	3 vs 4	29~~4~~	16.5~~8.1 (2.8)~~	0.19~~-2.6 (4.1)~~
	~~2010s~~	~~1~~	~~17.2 (1.8)~~	~~18.1 (1)~~
		~~2~~	~~24.4 (2.9)~~	~~36.2 (1.8)~~
		~~3~~	~~17.4 (2.9)~~	~~116.6 (3.3)~~
		~~4~~	~~NA~~	~~NA~~
<u>Parasitism rate</u>	1 vs 3~~1990s~~	-29.8[*1]	11.4~~17.2 (1.3)~~	0.036~~0.2 (1)~~
		~~2~~	~~NA~~	~~NA~~
	1 vs 4	72[*3]	14~~20.8 (1.7)~~	<0.001~~89.5 (1.8)~~
	3 vs 4	101.4[*]	16~~6.8 (1.5)~~	<0.001~~-173.4 (4.2)~~
	~~2010s~~	~~1~~	~~8.9 (0.7)~~	~~-55.3 (1)~~
		~~2~~	~~9.4 (1.1)~~	~~-1 (1.6)~~
		~~3~~	~~12.8 (1.8)~~	~~15.4 (3.3)~~
		~~4~~	~~NA~~	~~NA~~

~~***Egg mortality factors.***~~ <u>*. The mean difference is significant at the .05 level.</u>

~~Abortion and parasitism rates were the lowest (0%) in cluster 1 in 2017, while sterility was the~~

~~highest, reaching 100% that year (Figure 6). The latter varied significantly between clusters (Quade's~~

~~RANCOVA, $F_{df} = 2.8_3$, p = 0.037) and periods ($F_{df} = 5.9_1$, p = 0.016). Sterility differences among clusters~~

~~varied between the 1990s and 2010s (Quade's RANCOVA, $F_{df} = 9.1_1$, *p* = 0.003). No differences were~~

~~found between clusters in the 1990s, however, sterility was significantly lower in cluster 3 than~~

~~cluster 1 (Pairwise t-test, $t_{df} = -3.7_2$, p = 0.001 adjusted by Bonferroni correction) and cluster 2 ($t_{df} = -~~

~~2.6_2$, p = 0.025) in the 2010s. Cluster 1 was the only cluster where sterility significantly increased over~~

~~time ($t_{df} = 3.6_1$, p <0.001) (Table 2).~~

~~Abortion and parasitism rates significantly differed between clusters (abortion rate: F~~$_{df}$~~ = 9.2~~$_3$~~, p < 0.001; parasitism rate: F~~$_{df}$~~ = 17.6~~$_3$~~, p < 0.001) and periods (abortion rate: F~~$_{df}$~~ = 40.2~~$_1$~~, p <0.001; parasitism rate: F~~$_{df}$~~ = 36.8~~$_1$~~, p < 0.001). However, unlike the other variables, no significant interactions were found between the main effects (abortion rate: F~~$_{df}$~~ = 0.3~~$_1$~~, p = 0.607; parasitism rate: F~~$_{df}$~~= 1.2~~$_1$~~, p = 0.283). Pairwise differences for these variables between clusters are summarized in Table 3.~~

~~Table 3.~~

Table 3: Results of Tukey's ~~Turkey's~~ post hoc tests in the 2010s ~~at a threshold of 0.05 for abortion and parasitism rates~~.

Variable	Clusters ~~Cluster~~	Mean difference (I-J) ~~Observed mean (M)~~	Standard error ~~Quade's adjusted mean (Madj)~~	p value ~~Statistical significance for alpha = 0.05~~
Abortion Hatching rate	1 vs 2	-18.7 ~~10.6~~	13 ~~33~~.1	0.33 ~~a~~
	1 vs 3	-46.7[*]	16.8	0.02
	2 vs 3	-28	18.3	0.28
Sterility rate	1 vs 2	16.5 ~~24.4~~	13 ~~36~~.2	0.42 ~~bc~~
	1 vs 3	59.2[*12.3]	16.9 ~~56.7~~	<0.001 ~~bd~~
	2 vs 3 ~~4~~	42.6 ~~8.1~~	18.4 ~~2.6~~	0.054 ~~ab~~
Abortion rate	1 vs 2	-34.3[*]	13.2	0.03
	1 vs 3	-35.2	17	0.10
	2 vs 3	-0.9	18.5	0.99
Parasitism rate	1 vs 2	-21.3 ~~13~~	13.1 ~~3.5~~	0.24 ~~b~~
	1 vs 3 ~~2~~	-48.1[*9]	16.9 ~~1~~	0.01 ~~bc~~
	2 vs 3	-26.8 ~~18~~	18.4 ~~62.5~~	0.31 ~~d~~
	~~4~~	~~7~~	~~-173.4~~	~~a~~

[*]. The mean difference is significant at the .05 level.

Discussion

Climate change has been recognized to be one of the major phenomena that may affect forest insect populations (Ramsfield et al., 2016). While many studies reported ~~report~~ the occurrence of more frequent and larger insect outbreaks (Raffa et al., 2008; Robinet & Roques, 2010), the opposite ~~diminishments in outbreak severity~~ have also been observed (Pureswaran et al., 2018;

Rozenberg et al., 2020). Ongoing ~~The ongoing~~ global warming may exert mixed effects on population dynamics (Dreyer & Baumgärtner, 1996; Huang et al., 2008), and ultimately have an impact on species distributions, as has been observed with the PPM northern range expansion ~~-~~in Europe (Battisti et al. 2005) and the southern range retraction in Tunisia (Bourougaaoui et al. 2021).~~-~~ A report by the German Technical Cooperation Agency (GTZ et al. (2007)) has predicted more intense and longer heatwaves in Tunisia, with temperatures tending to rise even further in the coming century. To better understand potential adverse effects of climate change at the southern edge of the PPM range, the present study sought to explore variations in egg survival and hatching and their potential relationship with climate variations, based on a set of historical and recent field samplings across Tunisia.

~~Fluctuations of monthly averaged TX during summer months over the period 1990-2019 showed that July and August are the warmest months overall (Fig 3, 5).~~ The ~~fastest warming was observed in June and July with an increasing trend (although not always significant) in clusters 2, 3 and 4 for June, and in all the clusters for July. In August, positive increase trends were noted in clusters 2 and 3, but the trend was significant in cluster 2 only (Fig. 5).~~

~~Hatching rate tended to~~ decrease ~~between the 1990s and the 2010s in cluster 1 (for which the number of egg samples were the highest). Particularly, in 2017, a sharp decrease to 0%~~ in hatching rate ~~was~~ observed ~~(and reflected~~ in ~~an equally steep increase in sterility rate), with no PPM larva or parasitoids (Fig. 6). The decrease in hatching rate observed during other years of~~ the 2010s period was ~~less critical and~~ caused by a mixture of increases in the rate of sterile eggs, parasitized eggs, and ~~most importantly,~~ a clear increase in ~~the~~ abortion rate (i.e., fertile eggs with failed embryo development), the latter being possibly related to warmer conditions during embryonic development. The strikingly high sterility rate observed in 2017 could be related to an unusually long series of 10 consecutive days above 40 °C recorded that year (see Fig. 6 and Fig SM~~8-9~~for meteorological data from the closest station of the site sampled that year). While extreme compared to the last 30 years, this anomaly reflects the global increase in the total number of acute heat days recorded in August in this station (Fig. 6). This overall trend in turn corroborates the assumption that the likelihood of such stochastic events should increase with future climate change and cannot be neglected since they might represent a prime cause of mortality in the PPM, before the average warming. We found that July is the most rapidly warming month in Tunisia, but egg masses in sites within cluster 1 are mostly laid after July and occur in August. Since all 43 egg masses from cluster 1 in 2017 were collected in late August, after this long heatwave, egg development may have been directly impacted before any sign of embryogenesis could be detected (noted as "sterile" from visual inspections). Such acute heat may also have accelerated pheromone decay due to higher

evaporation rate, hence affecting mating success and egg fecundation in the first place (Groot & Zizzari, 2019; Linn et al., 1988), or adult gametes~~The unusually high sterility rate observed in 2017 could be explained by extreme temperatures during that particular summer, suggesting that the trend is more than a simple correlation and might be causal. Maximal temperature overreached 40°C during approximately 10 consecutive days in August, which matches the usual timing of egg occurrence in sites within cluster 1. Such acute heat may have directly damaged newly laid eggs during that period before any embryo could be visible (*i.e.*, not sterile per se, but aborted very early), or alternatively, the heat stress may have impacted adult gametes and fertility~~ (Sales et al., 2018). These results bear a close resemblance to those shown by Rocha et al. (2017), which revealed that negative effects appeared on Tunisian egg masses at 42°C after only 3 days of heatwaves, and no survivorship was noted at 44°C. ~~Caution must be taken when interpreting the hatching rate drop in 2017, however, because only one site (Thélepte) was sampled in cluster 1 during that peculiar year, and it remains unknown whether populations in the vicinity and similarly exposed to the heatwave have also incurred a severe drop in hatching rate.~~

Figure 6. Frequency of consecutive days in August overreaching 32 (orange) and 40°C (red) over 1990-1999 and 2010-2019 in the Kasserine station. This station is situated less than two kilometers away from Thélepte, the egg sampling site where 43 egg clutches (100 %) were sterile in 2017. Grey histograms correspond to the total distribution across both periods.

Hatching rate was slightly higher in cluster 2 than cluster 1 in the 2010s (Table SM7), which can be attributed to the drop in 2017 in cluster 1 (Figure 5). Despite being true for all egg masses in cluster 1 that year, caution must be taken before generalizing the low egg survival recorded in 2017 to a temporal trend in the whole cluster 1. First, all eggs of cluster 1 in 2017 originated from the single site of Thélepte, however egg masses were collected on multiple trees scattered across the area, thereby reducing the risk of biased sampling, and the sample size was one of the largest of the whole data series (43 eggs masses; see Table 1). Second, inter-annual variability in climate and other factors not measured in this study cannot be neglected, meaning that the recent years cannot be

summarized to the year of 2017 which has been shown to be extreme. However, global climatic models as well as the data presented in this manuscript suggest that these types of events are likely to increase in frequency with climate change, together with average warming, and therefore should be explicitly taken into account when analyzing PPM dynamics because they represent prime causes of lethality. By putting together long data series, the present study provides a broader view on the spatial and temporal variations in the fecundity penalty that may result from these extremely high temperatures, as well as how the timing of these heatwaves may differentially impact populations from different areas. Beyond the dramatic drop in egg survival observed in 2017 in cluster 1, smaller penalties on hatching rate have also been observed during other years of the 2010s. Those smaller decreases of egg survival may have a snowball effect on whole colony success, as shown in Spain by Pérez-Contreras et al. (2003) who found that 32 individuals is a threshold above which larval growth reaches its maximum and mortality drops substantially. A similar result was observed in an exploratory experiment in a French population during winter, where survival was null for colonies of less than 50 individuals (Roques et al., 2015).

The probability to overreach biological thresholds was found to be the highest in cluster 2 and to significantly increase over time, but no historical egg data from this cluster are available to confirm whether egg survival used to be higher in the 1990s. Our finding showed that August conditions in 2017 have likely been lethal in Thélepte (and possibly to a largest extent in cluster 1) may indicate that similar dramatic effects associated with stochastic meteorological events are to be expected in cluster 2 where summers are already both the hottest and the most rapidly warming. However, eggs occur later in cluster 2 (hatching starting in mid-October for recent samples) than in cluster 1 (hatching starting in early September), and may therefore mitigate summer heatwave threats thanks to phenological avoidance. This contrasting situation warrants the high relevance of future research in populations from cluster 1 and 2 to observe in real-time and test how climate change may cause the retraction of the PPM at its southern edge (see Bourougaaoui et al. 2021).

Temperature thresholds of 32°C and 40°C have been frequently used in the literature as pivotal for range limits of the PPM. They have been inferred from observations of survival in nature, but also appear consistent with more recent experiments in multiple populations of the PPM. Eggs from a French population were able to withstand a short transient exposure to a daily maximal temperature of 40°C during several consecutive days with no mortality impact (Robinet et al., 2013), while eggs from a nearby population could survive a single 6-hour-long exposure to up to 44°C (Poitou, 2021). However, mortality appeared on egg masses from a Tunisian population at 42°C after only three days of 4-hour daily exposures (Rocha et al., 2017). While slightly different methods have been used and make it difficult to compare populations, those results help narrowing down the tipping point at

which PPM egg survival is impacted, depending on the duration of exposure. Regarding larvae, the survivorship of L1 and L2 from Portuguese populations started to drop after 4-hour exposures to 36°C and 40°C, respectively (Santos et al., 2011), showing higher susceptibility in early larvae compared to eggs. Recently, Poitou et al. (2022) determined experimentally the thermal performance curves in development rate in the first four larval instars in a French population, highlighting that the 32°C threshold is above the optimal development temperature, within the decreasing performance phase. The 32°C and 40°C thresholds proposed by Démolin (1969) and (Huchon & Démolin, 1970) thus appear as conservative but consistent integrators of whether a population is facing stressful conditions regardless of the exact duration of exposure, which our results corroborate with full mortality after the 2017 heatwave but not after the intense yet less stringent 2019 heatwave (Figure SM89).

Hatching rate was slightly higher in cluster 2 than cluster 1 in the 2010s (Table 2), which can be explained by the lowest hatching rate that dropped to 0% in 2017 in cluster 1 (Figure 6). This dramatic decline contributed the most to the greater decrease in hatching observed in cluster 1 than cluster 2 over the entire 2010s. Although the probability to overreach 32°C (proposed by Démolin (1969) as a biological threshold above which physiological processes in eggs and/or larvae may be altered and epizootic may be favoured, although it remains unclear whether this threshold was experimentally verified and compared among populations) and 40°C in summer since the 1990s was found to be the highest in cluster 2, no historical egg data from cluster 2 are available to confirm whether hatching rate was lower in cluster 2 than cluster 1 in the 1990s. During uneven years with unusually hot summers, the biological thresholds of the PPM summer instars may be overreached in an unpredictable way and in turn negatively affect egg survival, as already demonstrated in southern Tunisia (Bourougaaoui et al. 2021). According to this study, temperatures exceeded 35°C on more than half of July and August months (nearly each year during the period 1980-2019), even 40°C for few days in the southern regions. In September, temperatures above 35°C and 40°C were found to be frequent in Tataouine over 1980-2019, which suggested the contribution of these high temperatures to population declines and to the disappearance of the PPM from southernmost areas (Bourougaaoui et al., 2021). The present study provides a broader view on the spatial and temporal variations in the fecundity penalty that may result from these extremely high temperatures. Beyond striking drops in fecundity observed in 2017 in cluster 1, lower fecundity penalties may have a snowball effect on whole colony survival. In Spain, Pérez-Contreras et al. (2003) found that 32 individuals is a threshold above which larval growth reaches its maximum and mortality drops substantially. A similar result was observed in an exploratory experiment in a French population during winter, where survival was null for colonies of 25 or 50 individuals (Roques et al., 2015).

Temperature is a crucial abiotic factor that can lead to local adaptations in insects and shape their

geographic range (Bush et al., 2016; Hoffmann et al., 2002; Kellermann et al., 2012; Sinclair et al.,

2012). Various thermal environments may therefore be associated with differences in behaviours

and even physiological tolerance (Calosi et al., 2010). Numerous studies have focused on

geographical variation in thermal tolerance in the fruit fly *Drosophila melanogaster*, considered as

one of the most widely distributed insect species, highlighting local variation in the thermal tolerance

and performance of distinct populations (see e.g., Sinclair et al., 2012). In this species, Hoffmann et

al. (2002) found opposing clines in resistance to temperature extremes when comparing numerous

Australian populations along a gradient from tropical to temperate latitudes, which suggests that

thermal tolerance traits are under direct climatic selection.

Likewise, summer temperature has been hypothesized as being a strong selection pressure in the

PPM due to the mortality observed in eggs (Rocha et al. 2017) and neonate larvae (Santos et al.

2011) after experimental heat exposure, combined with the wide range of climates under which the

PPM occurs. Variations in environmental parameters may induce phenological shifts in all stages of

this species either as a consequence of altered developmental time or responses to stressors (Berardi

et al., 2015; Robinet et al., 2015). In areas where summers~~summer~~ are the warmest, early mortality

caused by heat stress may favour late-emerging individuals through natural selection, especially as

adult females are shortlived and lay eggs only once (Rocha et al. 2017). A striking example of

phenological differentiation among PPM populations was recently discovered in Portugal, where two

sympatric forms exhibit contrasted life cycles: one with the typical overwinter larval development,

and one with a summer larval development (Santos et al., 2011). While little is known on the causal

factors that promoted the emergence of both forms in the same area, Godefroid et al. (2016)

demonstrated that the range of the summer population is restricted to central-coastal Portugal due

to climatic constraints, while winter populations expands northward. This may be due to the

comparatively low thermal resistance found in eggs of the summer form, which develop into mature

larvae before the peak of summer, as opposed to eggs and early instar larvae of the winter-

developing form which usually occur in the middle of summer and are more likely to be exposed to

heats (Rocha et al. 2017). The increase in climatic variability is expected to exert effects on insect

species that differ from those caused by gradual global warming (Schreven et al., 2017). Large-scale

heatwaves are known to have contrasted effects on different populations of the PPM depending on

local climates and phenologies (Robinet et al., 2015). In Europe, the summer heatwave in 2003 led to

a collapse of PPM populations in northern areas in France (Robinet et al., 2013), while it benefited to

adult dispersal and altitudinal expansion in the Italian Alps, likely due to flight thermal thresholds

more easily met than usual near the elevation edge~~)~~ (Battisti et al., 2006)~~.~~ These findings are

congruent with the available data in Tunisia where phenology in cluster 2 is delayed compared to

that in cluster 1, most likely due to the climatic and altitudinal differences found between those

clusters. This fact could contribute to explain the lower hatching rate found in cluster 1 than in

cluster 2 in 2017 (Figure 56), since eggs and neonate larvae occur after most summer heats in cluster

2 as a result ofdue to later adult emergences. This suggests that populations from the warmest areas

of Tunisia may not be the most vulnerable to climate change thanks to phenological adaptation

favouring heat avoidance, as opposed to populations from areas where individuals are close to their

physiological limits but did not evolve differentiated phenology in response to heat lethality.

Despite the impacts of the PPM on Aleppo pine forests in Tunisia, little is known about how

climate change can alter the phenology of Tunisian population and whether it is is spatially

structured by climate heterogeneity. According to Robinet et al. (2015), predictable unfavourable

conditions to which PPM populations are exposed can be alleviated by differentiated phenologies

(phenological plasticity and/or adaptations), which is supported by later adult emergences in cluster

2 on average. By contrast, unpredictable adverse conditions may be mitigated by bet-hedging

strategies such as prolonged diapause, as it creates heterogeneity within populations and siblings by

expanding the life cycle of only a fraction of individuals that stay sheltered in the ground at the pupal

stage and evade climate stochasticity (Salman et al., 2019). Diapausing individuals may therefore

contribute to reconstituting local population collapses after extreme events, but the cost is that

survival decreases with the total duration of diapause due to other factors of mortality (fungi, pupal

predation, etc.) (Salman et al. 2019). A significant relationship was found between the rate of

prolonged diapause and both cold and warm winter temperatures, presumably because they serve

as cues of the likelihood of either cold or hot lethal temperatures caused by inter-annual climate

uncertainty (Salman et al. 2019). The acute heat stress exerted on PPM egg masses in Tunisia may

help explain the retraction of this pest from southernmost regions, and further investigations on

phenological differentiations and prolonged diapause strategies among populations incurring

different levels climate harshness are now necessary to predict further distribution changes.

A growing body of literature reveals that parasitoids are often more sensitive to climate warming

than other trophic levels because of their higher position in the food web (Jeffs & Lewis, 2013;

Rosenblatt & Schmitz, 2016). Climate change can lead to phenological asynchrony between

parasitoids and their hosts in cases where the phenology of the interacting species respond variously

to the same climatic cue (Visser & Holleman, 2001), or when the interacting species use different

cues to initiate emergence or development (Jeffs & Lewis, 2013; Walther, 2010). Phenological

asynchrony may also appear if one of the interacting partners rapidly develops or has a seasonal

diapause in response to warming (Forrest, 2016). Parasitism rates therefore tend to decrease with

increasing climatic variability that impedes parasitoids from tracking host populations (Chidawanyika et al., 2019). Alternatively, but not exclusively, eggs may escape parasitic attacks thanks to various counter-adaptations whose relative weight may differ among regions and populations. The identification of eggs by natural enemies may be hindered making egg masses inconspicuous, as PPM female covering eggs by greyish-brown scales similar to the colour of pine twigs (Battisti et al., 2015). These scales function not only as visual protection but also as factors impacting the microclimate of eggs and therefore their development rate (Milani, 1990), and as physical barriers limiting parasitoids from locating individual eggs in the clutch and greatly reducing their success (Pérez-Contreras & Soler, 2004). The chief defence against parasitoids, widely discussed in insects and in particular Drosophila spp., is encapsulation, an immune responses which sequesters and kills foreign body (Cavigliasso et al., 2021; Wertheim et al., 2005). Such immune system with specific cells (hemocytes) is well-developed in larval stages as already observed in PPM larvae (Boudjahem et al., 2019), however, a study conducted by Reed et al. (2007) showed that hosts in the egg stage can mount a cellular immune response against parasitoid eggs and larvae (Reed et al., 2007). Research on a wide range of species reveals that small changes in temperature can significantly shape insect immunity as well as parasitoid fitness (Murdock et al., 2012). Increases in temperature can consequently promote or repress the encapsulation process, which in turn may influence the outcome of parasitic success, (Cavigliasso et al., 2021). The outbreaks of phytophagous insects are expected to increase in the future as parasitism decreases~~As a result, outbreaks of phytophagous insects are expected to increase in the future~~ (Stireman et al., 2005). At the southern edge of PPM distribution (North Africa), some studies showed that extremely high temperatures could both disrupt population regulation mechanisms and decrease the severity of outbreaks (Bouzar.Essaidi et al., 2021; Pureswaran et al., 2018). The lower PPM fecundity in this area (when compared to that at the northern edge, in southern Europe) due to lower plant quality (Bouzar.Essaidi et al., 2021; Pimentel et al., 2010) may explain why egg parasitism is also lower with climate warming. Our results showed that the mean number of eggs per egg mass (155 ± 2.1 in cluster 1, 142 ± 4.5 in cluster 2, 150 ± 3.5 in cluster 3, 145 ± 6.3 in cluster 4; 151 ± 1.7 when merging all clusters) was considerably lower than that found in Bulgaria (226 ± 43.2) and France (194.3 ± 50.1) (Georgiev et al., 2020). The average rate of parasitism also showed a different pattern than what was found in northern parts of PPM distribution, with increases in coastal regions namely in cluster 1 in Tunisia, while it decreases in France and Spain from core to front populations and along an altitudinal gradient, respectively (Georgiev et al., 2020; Hódar et al., 2021). Although parasitoids account in egg mortality, their influence here was low compared to other factors (sterility and abortion) (Figure 5), and is therefore unlikely to be the main driver of PPM collapses at the southern edge of the distribution.~~Similarly, the average rate of parasitism in the four clusters was low (13 ± 0.7% in cluster 1, 9 ± 1.1% in cluster 2, 18 ± 1.3% in cluster 3, 7 ± 1.5% in~~

cluster 4; 13 ±0.5% when merging all clusters) compared to northern parts of PPM distribution (24.4 % in France and 29% in Bulgaria) (Georgiev et al., 2020), and is therefore unlikely to be the main driver of PPM collapses at the southern edge of the distribution. Our findings confirm Bouzar et al. (2021)'s results showing that fecundity and parasitism were lower in Algeria than in southern Europe . Significant local warming in Tunisia appears as a prime candidate factor contributing to the sharp decrease of PPM populations (Bourougaaoui et al., 2021).

Factors other than climate warming may putatively affect the survival and persistence of the PPM at its southern range edge. Embryonic mortality can be impacted by excessive exposure to intense solar radiation, particularly in southern parts of the distribution, as PPM females tend to lay their eggs exposed to the sun (Démolin, 1969). Another factor often modulating the spatial occurrence of insects is food availability. Nevertheless, it is rarely a limiting factor in the PPM because larvae feed on evergreen trees that are well distributed in the environment, from natural or semi-natural stands to urban areas where they often occur in relatively high numbers as ornamental trees (Martin, 2005). Natural enemies such as pathogens or predators (mostly insect parasitoids) at early larval stages have been suspected to cascade into increasing mortality during larval development because of the impact on the colony size and silk weaving effort to build a tent (Auger Rozenberg et al., 2015; Roques et al., 2015), however (1) there is no evidence that the enemy pressure would differ among areas investigated here, and (2) temperature, particularly summer heat waves or early autumnal cold snaps, are often put forward as a major cause of early mortality (Robinet et al., 2015).

Heat tolerance has received close attention in insects, however its fluctuation throughout ontogeny and effects persisting from one developmental stage to another are still poorly documented. Besides the PPM, few case studies showed that the effects of acute heat stress received early in life cycle may be carried over to later instars. This was demonstrated in holometabolous insects such as the tropical butterfly, *Bicyclus anynana* (Klockmann et al., 2017). Beside consequences of heat on immediate mortality investigated in experimental work (e.g., Rocha et al 2017) or inferred in the present study by putting together long time series, the ultimate fitness of individuals that survive challenging heats at the egg stage or first larval instar would therefore be of great interest to understand the impacts of climate warming at the southern edge of the PPM. This insect remains one of the ideal models to study these questions owing to (i) the availability of historical data, (ii) and its already demonstrated spatial and phenotypic causal response to climate change (Battisti et al., 2005; Robinet et al., 2007; Poitou et al., 2022), and (iii) ongoing processes at play in its southernmost distribution affecting population persistence (Bourougaaoui et al. 2021; this study). (Battisti et al., 2005; Robinet et al., 2007).

Acknowledgements

We are grateful to Adel Ben Abada (INRGREF, Tunis, Tunisia) for his valuable help in the field. We
acknowledge the National Institute of Meteorology (INM) in Tunis for providing temperature
datasets.

Funding

This work was supported by the Tunisian Ministry of Higher Education of Scientific Research and
Technology and the University of Carthage: the university provided two grants to AB for internships
at INRAE URZF in France during her PhD.

Conflict of interest disclosure

The authors declare no conflict of interest.

Author contributions

Conceptualization: AB, CR, MLBJ, ML; Data curation: AB; Formal analysis: AB, ML; Funding
acquisition: CR, MLBJ; Investigation: AB; Methodology: AB, CR, ML; Project administration: CR, MLBJ;
Supervision: CR, MLBJ, ML; Writing – original draft: AB, CR, ML; Writing – review and editing: AB, CR,
MLBJ, ML.

Data, script and code availability

Data and R scripts are publicly available at https://doi.org/10.15454/RUEIOA.

https://doi.org/10.15454/RUEIOA.

References

Allen, S., Cardona, O., Cutter, S., Dube, O. P., Ebi, K., Handmer, J., Lavell, A., Mastrandrea, M.,
McBean, G., Mechler, R., & Nicholls, N. (2012). Managing the Risks of Extreme Events and
Disasters to Advance Climate Change Adaptation. Special Report of Working Groups I and II
of   the   Intergovernmental   Panel   on   Climate   Change.   In.
https://doi.org/10.13140/2.1.3117.9529
Auger Rozenberg, M. A., Barbaro, L., Battisti, A., Blache, S., Charbonnier, Y., Denux, O., Garcia, J.,
Goussard, F., Imbert, C.-E., Kerdelhué, C., Roques, A., Torres Leguizamon, M., & Vetillard, F.
(2015). Ecological Responses of Parasitoids, Predators and Associated Insect Communities to
the Climate-Driven Expansion of the Pine Processionary Moth. In A. Roques (Ed.),
Processionary Moths and Climate Change : An Update (pp. 311-357). Springer Netherlands.
https://doi.org/10.1007/978-94-017-9340-7_7
Battisti, A., Avcı, M., Avtzis, D. N., Jamaa, M. L. B., Berardi, L., Berretima, W., Branco, M., Chakali, G.,
El Alaoui El Fels, M. A., Frérot, B., Hódar, J. A., Ionescu-Mălăncuş, I., İpekdal, K., Larsson, S.,
Manole, T., Mendel, Z., Meurisse, N., Mirchev, P., Nemer, N., . . . Zamoum, M. (2015). Natural
History of the Processionary Moths (Thaumetopoea spp.): New Insights in Relation to Climate
Change. In A. Roques (Ed.), *Processionary Moths and Climate Change : An Update* (pp. 15-79).
Springer Netherlands. https://doi.org/https://doi.org/10.1007/978-94-017-9340-7_2
Battisti, A., Stastny, M., Buffo, E., & Larsson, S. (2006). A rapid altitudinal range expansion in the pine
processionary moth produced by the 2003 climatic anomaly. *Global Change Biology*, *12*(4),
662-671. https://doi.org/https://doi.org/10.1111/j.1365-2486.2006.01124.x
Battisti, A., Stastny, M., Netherer, S., Robinet, C., Schopf, A., Roques, A., & Larsson, S. (2005).
Expansion of geographic range in the pine processionary moth caused by increased winter
temperatures. *Ecol Appl*, *15*(6), 2084-2096. https://doi.org/https://doi.org/10.1890/04-1903
Ben Jamâa, M., & Jerraya, A. (1999). Essai de lutte contre la processionnaire du pin, Thaumetopoea
pityocampa Schiff.(Lep., Thaumetopoeïdae), à l'aide de Bacillus thuringiensis Kurstaki
(ECOTECH-PRO). Annales de l'INRGREF.
Berardi, L., Branco, M., Paiva, M., Santos, H., & Battisti, A. (2015). Development time plasticity of the
pine processionary moth (Thaumetopoea pityocampa) populations under laboratory
conditions. *Entomologia*, *3*, 19-24. https://doi.org/10.4081/entomologia.2015.273
Boudjahem, I., Brivio Fransisco, M., Berchii, S., Mastore, M., & Aoouati, A. (2019). Identification and
Quantification of the Most Abondant Hemocytes in the Pine Processionary Caterpillar;
ThaumetopoeaPityocampa   (Notodontidae).   *Energy   Procedia*,   *157*,   992-998.
https://doi.org/https://doi.org/10.1016/j.egypro.2018.11.266
Bourougaaoui, A., Jamâa, M. L. B., & Robinet, C. (2021). Has North Africa turned too warm for a
Mediterranean forest pest because of climate change? *Climatic Change*, *165*(3-4), 46.
https://doi.org/10.1007/s10584-021-03077-1
Bouzar.Essaidi, K., Branco, M., Battisti, A., Garcia, A., Fernandes, M. R., Chabane, Y., Bouzemarene,
M., & Benfekih, L. (2021). Response of the egg parasitoids of the pine processionary moth
to host density and forest cover at the southern edge of the range. *Agricultural and Forest
Entomology*, *23*(2), 212-221. https://doi.org/https://doi.org/10.1111/afe.12423
Bush, A., Mokany, K., Catullo, R., Hoffmann, A., Kellermann, V., Sgrò, C., McEvey, S., & Ferrier, S.
(2016). Incorporating evolutionary adaptation in species distribution modelling reduces
projected   vulnerability   to   climate   change.   *Ecology   Letters*,   *19*(12),   1468-1478.
https://doi.org/https://doi.org/10.1111/ele.12696
Calosi, P., Bilton, D. T., Spicer, J. I., Votier, S. C., & Atfield, A. (2010). What determines a species'
geographical range? Thermal biology and latitudinal range size relationships in European
diving beetles (Coleoptera: Dytiscidae). *Journal of Animal Ecology*, *79*(1), 194-204.
https://doi.org/10.1111/j.1365-2656.2009.01611.x

Carus, S. (2009). Effects of defoliation caused by the processionary moth on growth of Crimean pines in western Turkey. *Phytoparasitica*, *37*(2), 105-114. https://doi.org/10.1007/s12600-008-0018-z

Cavigliasso, F., Gatti, J. L., Colinet, D., & Poirié, M. (2021). Impact of Temperature on the Immune Interaction between a Parasitoid Wasp and Drosophila Host Species. *Insects*, *12*(7), 647. https://www.mdpi.com/2075-4450/12/7/647

Charmantier, A., & Gienapp, P. (2014). Climate change and timing of avian breeding and migration: evolutionary versus plastic changes. *Evolutionary applications*, *7*(1), 15-28. https://doi.org/10.1111/eva.12126

Chidawanyika, F., Mudavanhu, P., & Nyamukondiwa, C. (2019). Global Climate Change as a Driver of Bottom-Up and Top-Down Factors in Agricultural Landscapes and the Fate of Host-Parasitoid Interactions [Review]. *Frontiers in Ecology and Evolution*, *7*(80). https://doi.org/10.3389/fevo.2019.00080

Chuine, I. (2010). Why does phenology drive species distribution? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1555), 3149-3160. https://doi.org/doi:10.1098/rstb.2010.0142

Chuine, I., de Cortazar-Atauri, I. G., Kramer, K., & Hänninen, H. (2013). Plant Development Models. In M. D. Schwartz (Ed.), *Phenology: An Integrative Environmental Science* (pp. 275-293). Springer Netherlands. https://doi.org/10.1007/978-94-007-6925-0_15

Clark, B. R., & Faeth, S. H. (1997). The consequences of larval aggregation in the butterfly Chlosyne lacinia. *Ecological Entomology*, *22*(4), 408-415. https://doi.org/https://doi.org/10.1046/j.1365-2311.1997.00091.x

Coumou, D., & Rahmstorf, S. (2012). A decade of weather extremes. *Nature Climate Change*, *2*(7), 491-496. https://doi.org/10.1038/nclimate1452

Crozier, L. (2004). WARMER WINTERS DRIVE BUTTERFLY RANGE EXPANSION BY INCREASING SURVIVORSHIP. *Ecology*, *85*(1), 231-241. https://doi.org/https://doi.org/10.1890/02-0607

Démolin, G. (1965). Grégarisme et subsocialité chez Thaumetopoea pityocampa Schiff. Nids d'hiver– activité de tissage. Actes du Vᵉ Congress de L'Union Internationale pour L'étude des insectes Sociaux,

Démolin, G. (1969a). Bioécologie de la processionnaire du pin Thaumetopoea pityocampa Schiff. Incidences des facteurs climatiques. *Boletin del Servicio de Plagas Forestales*(23), 9-24. https://hal.inrae.fr/hal-02732616

Démolin, G. (1969b). Comportement des adultes de Thaumetopoea pityocampa Schiff. Dispersion spatiale, importance écologique. Annales des sciences forestières,

Démolin, G., & Rive, J. (1968). La processionnaire du pin en Tunisie. *Ann. I.N.R.F. Tunisie*, *1*(1), 1-19.

Denno, R., & Benrey, B. (1997). Aggregation facilitates larval growth in the neotropical nymphalid butterfly Chlosyne janais. *Ecological Entomology*, *22*(2), 133-141. https://doi.org/https://doi.org/10.1046/j.1365-2311.1997.t01-1-00063.x

Dreyer, H., & Baumgärtner, J. (1996). Temperature influence on cohort parameters and demographic characteristics of the two cowpea coreids Clavigralla tomentosicollis and C. shadabi. *Entomologia Experimentalis et Applicata*, *78*(2), 201-213. https://doi.org/https://doi.org/10.1111/j.1570-7458.1996.tb00783.x

EPPO. (2004). EPPO Standards: Thaumetopoea pityocampa- PM7/37. *Bulletin OEPP/EPPO Bulletin*, *34*, 295-298.

Fischer, E. M., & Schär, C. (2010). Consistent geographical patterns of changes in high-impact European heatwaves. *Nature Geoscience*, *3*(6), 398-403. https://doi.org/10.1038/ngeo866

Fontaine, B., Janicot, S., & Monerie, P.-A. (2013). Recent changes in air temperature, heat waves occurrences, and atmospheric circulation in Northern Africa. *Journal of Geophysical Research: Atmospheres*, *118*(15), 8536-8552. https://doi.org/https://doi.org/10.1002/jgrd.50667

Forrest, J. R. K. (2016). Complex responses of insect phenology to climate change. *Current Opinion in Insect Science*, *17*, 49-54. https://doi.org/https://doi.org/10.1016/j.cois.2016.07.002

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body size: a third
universal response to warming? *Trends in Ecology & Evolution*, *26*(6), 285-291.
https://doi.org/https://doi.org/10.1016/j.tree.2011.03.005

Georgiev, G., Rousselet, J., Laparie, M., Robinet, C., Georgieva, M., Zaemdzhikova, G., Roques, A.,
Bernard, A., Poitou, L., Buradino, M., Kerdelhue, C., Rossi, J. P., Matova, M., Boyadzhiev, P., &
Mirchev, P. (2020). Comparative studies of egg parasitoids of the pine processionary moth
(Thaumetopoea pityocampa, Den. & Schiff.) in historic and expansion areas in France
and Bulgaria. *Forestry: An International Journal of Forest Research*, *94*(2), 324-331.
https://doi.org/10.1093/forestry/cpaa022

Ghosh, S. M., Testa, N. D., & Shingleton, A. W. (2013). Temperature-size rule is mediated by thermal
plasticity of critical size in Drosophila melanogaster. *Proceedings. Biological sciences*,
280(1760), 20130174-20130174. https://doi.org/10.1098/rspb.2013.0174

Godefroid, M., Rocha, S., Santos, H., Paiva, M. R., Burban, C., Kerdelhué, C., Branco, M., Rasplus, J. Y.,
& Rossi, J. P. (2016). Climate constrains range expansion of an allochronic population of the
pine processionary moth. *Diversity and Distributions*, *22*(12), 1288-1300.
https://doi.org/https://doi.org/10.1111/ddi.12494

Groot, A. T., & Zizzari, Z. V. (2019). Does climate warming influence sexual chemical signaling? *Animal
Biology*, *69*(1), 83-93. https://doi.org/https://doi.org/10.1163/15707563-20191103

GTZ, MARH, & Exaconsult Gopa. (2007). *Stratégie nationale d'adaptation de l'agriculture tunisienne
et des écosystèmes aux changements climatiques*. Rapport d'étude dans le cadre de la
coopération Tuniso-allemande publié par Deutsche Gesellschaft für Internationale.

Hickling, R., Roy, D. B., Hill, J. K., & Thomas, C. D. (2005). A northward shift of range margins in British
Odonata. *Global Change Biology*, *11*(3), 502-506.
https://doi.org/https://doi.org/10.1111/j.1365-2486.2005.00904.x

Hódar, J. A., Cayuela, L., Heras, D., Pérez-Luque, A. J., & Torres-Muros, L. (2021). Expansion of
elevational range in a forest pest: Can parasitoids track their hosts? *Ecosphere*, *12*(4),
e03476. https://doi.org/https://doi.org/10.1002/ecs2.3476

Hoffmann, A. A., Anderson, A., & Hallas, R. (2002). Opposing clines for high and low temperature
resistance in Drosophila melanogaster. *Ecology Letters*, *5*(5), 614-618.
https://doi.org/https://doi.org/10.1046/j.1461-0248.2002.00367.x

Huang, Z., Ren, S., & Musa, P. D. (2008). Effects of temperature on development, survival, longevity,
and fecundity of the Bemisia tabaci Gennadius (Homoptera: Aleyrodidae) predator,
Axinoscymnus cardilobus (Coleoptera: Coccinellidae). *Biological Control*, *46*(2), 209-215.
https://doi.org/https://doi.org/10.1016/j.biocontrol.2008.04.004

Huchon, H., & Demolin, G. (1970). La bioécologie de la Processionnaire du pin : dispersion
potentielle, dispersion actuelle.

Imbert, C. E. (2012). *Expansion d'un ravageur forestier sous l'effet du réchauffement climatique : la
processionnaire du pin affecte-t-elle la biodiversité entomologique dans les zones
nouvellement colonisées ?* PhD dissertation, Université d'Orléans (France), pp. 198.

Jacquet, J.-S., Bosc, A., O'Grady, A. P., & Jactel, H. (2013). Pine growth response to processionary
moth defoliation across a 40-year chronosequence. *Forest Ecology and Management*, *293*,
29-38. https://doi.org/https://doi.org/10.1016/j.foreco.2012.12.003

Jeffs, C. T., & Lewis, O. T. (2013). Effects of climate warming on host–parasitoid interactions.
Ecological Entomology, *38*(3), 209-218. https://doi.org/https://doi.org/10.1111/een.12026

Jones, P. D., Lister, D. H., Osborn, T. J., Harpham, C., Salmon, M., & Morice, C. P. (2012). Hemispheric
and large-scale land-surface air temperature variations: An extensive revision and an update
to 2010. *Journal of Geophysical Research: Atmospheres*, *117*(D5).
https://doi.org/https://doi.org/10.1029/2011JD017139

Karban, R., & Strauss, S. Y. (2004). Physiological tolerance, climate change, and a northward range
shift in the spittlebug, Philaenus spumarius. *Ecological Entomology*, *29*(2), 251-254.
https://doi.org/https://doi.org/10.1111/j.1365-2311.2004.00576.x

Kellermann, V., Overgaard, J., Hoffmann, A. A., Fløjgaard, C., Svenning, J.-C., & Loeschcke, V. (2012).
Upper thermal limits of Drosophila are linked to species distributions and strongly
constrained phylogenetically. *Proceedings of the National Academy of Sciences*, *109*(40),
16228. https://doi.org/10.1073/pnas.1207553109

Kingsolver, J. G., Diamond, S. E., & Buckley, L. B. (2013). Heat stress and the fitness consequences of
climate change for terrestrial ectotherms. *Functional Ecology*, *27*(6), 1415-1423.
https://doi.org/https://doi.org/10.1111/1365-2435.12145

Klockmann, M., Kleinschmidt, F., & Fischer, K. (2017). Carried over: Heat stress in the egg stage
reduces subsequent performance in a butterfly. *PloS one*, *12*(7), e0180968-e0180968.
https://doi.org/10.1371/journal.pone.0180968

Linn, C. E., Campbell, M. G., & Roelofs, W. L. (1988). Temperature modulation of behavioural
thresholds controlling male moth sex pheromone response specificity. *Physiological
Entomology*, *13*(1), 59-67. https://doi.org/https://doi.org/10.1111/j.1365-
3032.1988.tb00909.x

Liu, S. S., Zhang, G. M., & Zhu, J. (1995). Influence of Temperature Variations on Rate of Development
in Insects: Analysis of Case Studies from Entomological Literature. *Annals of the
Entomological Society of America*, *88*(2), 107-119. https://doi.org/10.1093/aesa/88.2.107

Martin, J. (2005). La processionnaire du pin Thaumetopoea pityocampa (Denis et Schiffermüller).
Biologie et protection des forêts. Avignon: Avignon Editions, INRA, 1-62.
http://www.prodinra.inra.fr/prodinra /pinra/index.xsp

Milani, N. (1990). The temperature of the egg masses of Thaumetopoea pityocampa (Den. & Schiff.)
(Lepidoptera, Thaumetopoeidae). *Redia*, *73*(1), 149-161.

Murdock, C. C., Paaijmans, K. P., Cox-Foster, D., Read, A. F., & Thomas, M. B. (2012). Rethinking
vector immunology: the role of environmental temperature in shaping resistance. *Nature
Reviews Microbiology*, *10*(12), 869-876. https://doi.org/10.1038/nrmicro2900

Nangombe, S. S., Zhou, T., Zhang, W., Zou, L., & Li, D. (2019). High-Temperature Extreme Events Over
Africa Under 1.5 and 2 °C of Global Warming. *Journal of Geophysical Research: Atmospheres*,
124(8), 4413-4428. https://doi.org/https://doi.org/10.1029/2018JD029747

Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J. K., Thomas, C. D., Descimon, H., Huntley, B., Kaila, L.,
Kullberg, J., Tammaru, T., Tennent, W. J., Thomas, J. A., & Warren, M. (1999). Poleward shifts
in geographical ranges of butterfly species associated with regional warming. *Nature*,
399(6736), 579-583. https://doi.org/https://doi.org/10.1038/21181

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across
natural systems. *Nature*, *421*(6918), 37-42.
https://doi.org/https://doi.org/10.1038/nature01286

Pérez-Contreras, T., & Soler, J. J. (2004). Egg parasitoids select for large clutch sizes and covering
layers in pine processionary moths (Thaumetopoea pityocampa). *Annales Zoologici Fennici*,
41(4), 587-597. http://www.jstor.org/stable/23735942

Pérez-Contreras, T., Soler, J., & Soler, M. (2003). Why do pine processionary caterpillars
Thaumetopoea pityocampa (Lepidoptera, Thaumetopoeidae) live in large groups? An
experimental study. *Annales Zoologici Fennici*, *40*, 505-515.

Pigliucci, M. (2001). *Phenotypic plasticity: beyond nature and nurture*. The John Hopkins University
Press.

Pigliucci, M. (2005). Evolution of phenotypic plasticity: where are we going now? *Trends in Ecology &
Evolution*, *20*(9), 481-486. https://doi.org/https://doi.org/10.1016/j.tree.2005.06.001

Pimentel, C., Ferreira, C., & Nilsson, J. A. N. Å. (2010). Latitudinal gradients and the shaping of life-
history traits in a gregarious caterpillar. *Biological Journal of the Linnean Society*, *100*(1), 224-
236. https://doi.org/https://doi.org/10.1111/j.1095-8312.2010.01413.x

Pincebourde, S., Dillon, M. E., & Woods, H. A. (2021). Body size determines the thermal coupling
between insects and plant surfaces. *Functional Ecology*, *35*(7), 1424-1436.
https://doi.org/https://doi.org/10.1111/1365-2435.13801

Pincebourde, S., & Woods, H. A. (2020). There is plenty of room at the bottom: microclimates drive insect vulnerability to climate change. *Current Opinion in Insect Science*, *41*, 63-70. https://doi.org/https://doi.org/10.1016/j.cois.2020.07.001

Poitou, L. (2021). *Etude de l'impact du changement climatique sur la phénologie de la processionnaire du pin.* PhD dissertation, Université d'Orléans (France), pp. 300.

Poitou, L., Iaparie, M., pincebourde, S., Rousselet, J., Suppo, C., & robinet, C. (2022). Warming causes atypical phenology in a univoltine moth with differentially sensitive larval stages. *Frontiers in Ecology and Evolution (in press)*. https://doi.org/10.3389/fevo.2022.825875

Pureswaran, D. S., Roques, A., & Battisti, A. (2018). Forest Insects and Climate Change. *Current Forestry Reports*, *4*(2), 35-50. https://doi.org/10.1007/s40725-018-0075-6

Quade, D. (1967). Rank Analysis of Covariance. *Journal of the American Statistical Association*, *62*(320), 1187-1200. https://doi.org/10.2307/2283769

Raffa, K. F., Aukema, B. H., Bentz, B. J., Carroll, A. L., Hicke, J. A., Turner, M. G., & Romme, W. H. (2008). Cross-scale Drivers of Natural Disturbances Prone to Anthropogenic Amplification: The Dynamics of Bark Beetle Eruptions. *BioScience*, *58*(6), 501-517. https://doi.org/https://doi.org/10.1641/B580607

Ramsfield, T. D., Bentz, B. J., Faccoli, M., Jactel, H., & Brockerhoff, E. G. (2016). Forest health in a changing world: effects of globalization and climate change on forest insect and pathogen impacts. *Forestry: An International Journal of Forest Research*, *89*(3), 245-252. https://doi.org/10.1093/forestry/cpw018

Reed, D. A., Luhring, K. A., Stafford, C. A., Hansen, A. K., Millar, J. G., Hanks, L. M., & Paine, T. D. (2007). Host defensive response against an egg parasitoid involves cellular encapsulation and melanization. *Biological Control*, *41*(2), 214-222. https://doi.org/https://doi.org/10.1016/j.biocontrol.2007.01.010

Reynolds, A. P., Richards, G., de la Iglesia, B., & Rayward-Smith, V. J. (2006). Clustering Rules: A Comparison of Partitioning and Hierarchical Clustering Algorithms. *Journal of Mathematical Modelling and Algorithms*, *5*(4), 475-504. https://doi.org/10.1007/s10852-005-9022-1

Robinet, C., Baier, P., Pennerstorfer, J., Schopf, A., & Roques, A. (2007). Modelling the effects of climate change on the potential feeding activity of Thaumetopoea pityocampa (Den. & Schiff.) (Lep., Notodontidae) in France. *GLOBAL ECOL BIOGEOGR*, *16*(4), 460-471. https://doi.org/ https://doi.org/10.1111/j.1466-8238.2006.00302.x

Robinet, C., Laparie, M., & Rousselet, J. (2015). Looking Beyond the Large Scale Effects of Global Change: Local Phenologies Can Result in Critical Heterogeneity in the Pine Processionary Moth. *Frontiers in Physiology*, *6*(334). https://doi.org/10.3389/fphys.2015.00334

Robinet, C., & Roques, A. (2010). Direct impacts of recent climate warming on insect populations. *Integrative Zoology*, *5*(2), 132-142. https://doi.org/https://doi.org/10.1111/j.1749-4877.2010.00196.x

Robinet, C., Rousselet, J., Pineau, P., Miard, F., & Roques, A. (2013). Are heat waves susceptible to mitigate the expansion of a species progressing with global warming? *Ecol Evol*, *3*(9), 2947-2957. https://doi.org/https://doi.org/10.1002/ece3.690

Rocha, S., Kerdelhué, C., Ben Jamaa, M. L., Dhahri, S., Burban, C., & Branco, M. (2017). Effect of heat waves on embryo mortality in the pine processionary moth. *Bull Entomol Res*, *107*(5), 583-591. https://doi.org/https://doi.org/10.1017/S0007485317000104

Ronnås, C., Larsson, S., Pitacco, A., & Battisti, A. (2010). Effects of colony size on larval performance in a processionary moth. *Ecological Entomology*, *35*, 436-445. https://doi.org/10.1111/j.1365-2311.2010.01199.x

Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C., & Pounds, J. A. (2003). Fingerprints of global warming on wild animals and plants. *Nature*, *421*(6918), 57-60. https://doi.org/10.1038/nature01333

Roques A, R. J., Avci M et al. (2015). Climate warming and past and present distribution of the

processionary moths (Thaumetopoea spp.) in Europe, Asia Minor and North Africa. In R. A (Ed.),
Processionary moths and climate change : an update (pp. 81-161). Springer.
https://doi.org/https://doi.org/10.

1007/978-94-017-9340-7_3

Roques, L., Rossi, J.-P., Berestycki, H., Rousselet, J., Garnier, J., Roquejoffre, J.-M., Rossi, L.,
Soubeyrand, S., & Robinet, C. (2015). Modeling the Spatio-temporal Dynamics of the Pine
Processionary Moth. In (pp. 227-263). https://doi.org/10.1007/978-94-017-9340-7_5

Rosenblatt, A. E., & Schmitz, O. J. (2016). Climate Change, Nutrition, and Bottom-Up and Top-Down
Food Web Processes. *Trends in Ecology & Evolution*, *31*(12), 965-975.
https://doi.org/https://doi.org/10.1016/j.tree.2016.09.009

Rosenzweig, C., Casassa, G., Karoly, D., Imeson, A., Liu, C., Menzel, A., Rawlins, S., Root, T., Seguin, B.,
& Tryjanowski, P. (2007). Assessment of observed changes and responses in natural and
managed systems. . In O. F. C. M.L. Parry, J.P. Palutikof, and P.J. van der Linden (Ed.), *Climate*
Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the
Fourth Assessment Report of the Intergovernmental Panel on Climate Change (pp. 79-131).
Cambridge University Press.

Rozenberg, P., Pâques, L., Huard, F., & Roques, A. (2020). Direct and Indirect Analysis of the
Elevational Shift of Larch Budmoth Outbreaks Along an Elevation Gradient [Original
Research]. *Frontiers in Forests and Global Change*, *3*(86).
https://doi.org/https://doi.org/10.3389/ffgc.2020.00086

Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, Ł., Hebberecht, L.,
Thomas, P., Franco, A., & Gage, M. J. G. (2018). Experimental heatwaves compromise sperm
function and cause transgenerational damage in a model insect. *Nature Communications*,
9(1), 4771. https://doi.org/10.1038/s41467-018-07273-z

Salman, M. H. R., Bonsignore, C. P., El Alaoui El Fels, A., Giomi, F., Hodar, J. A., Laparie, M., Marini, L.,
Merel, C., Zalucki, M. P., Zamoum, M., & Battisti, A. (2019). Winter temperature predicts
prolonged diapause in pine processionary moth species across their geographic range. *PeerJ*,
7, e6530-e6530. https://doi.org/10.7717/peerj.6530

Santos, H., Paiva, M. R., Tavares, C., Kerdelhué, C., & Branco, M. (2011). Temperature niche shift
observed in a Lepidoptera population under allochronic divergence. *J. Evol. Biol*, *24*(9), 1897-
1905. https://doi.org/https://doi.org/10.1111/j.1420-9101.2011.02318.x

Sbay, H., & Zas, R. (2018). Geographic variation in growth, survival, and susceptibility to the
processionary moth (Thaumetopoea pityocampa Dennis & Schiff.) of Pinus halepensis Mill.
and P. brutia Ten.: results from common gardens in Morocco. *Annals of Forest Science*, *75*(3),
69. https://doi.org/10.1007/s13595-018-0746-2

Schreven, S. J. J., Frago, E., Stens, A., de Jong, P. W., & van Loon, J. J. A. (2017). Contrasting effects of
heat pulses on different trophic levels, an experiment with a herbivore-parasitoid model
system. *PloS one*, *12*(4), e0176704-e0176704. https://doi.org/10.1371/journal.pone.0176704

Schubert, E., & Rousseeuw, P. J. (2019). Faster k-Medoids Clustering: Improving the PAM, CLARA, and
CLARANS Algorithms. In G. Amato, C. Gennaro, V. Oria, & M. Radovanović, *Similarity Search*
and Applications Cham.

Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change.
Nature Climate Change, *1*(8), 401-406. https://doi.org/10.1038/nclimate1259

Sinclair, B. J., Williams, C. M., & Terblanche, J. S. (2012). Variation in Thermal Performance among
Insect Populations. *Physiological and Biochemical Zoology: Ecological and Evolutionary*
Approaches, *85*(6), 594-606. https://doi.org/10.1086/665388

Stireman, J. O., Dyer, L. A., Janzen, D. H., Singer, M. S., Lill, J. T., Marquis, R. J., Ricklefs, R. E., Gentry,
G. L., Hallwachs, W., Coley, P. D., Barone, J. A., Greeney, H. F., Connahs, H., Barbosa, P.,
Morais, H. C., & Diniz, I. R. (2005). Climatic unpredictability and parasitism of caterpillars:
Implications of global warming. *Proceedings of the National Academy of Sciences of the*
United States of America, *102*(48), 17384. https://doi.org/10.1073/pnas.0508839102

Thompson, R. M., Beardall, J., Beringer, J., Grace, M., & Sardina, P. (2013). Means and extremes:
building variability into community-level climate change experiments. *Ecology Letters*, *16*(6),
799-806. https://doi.org/https://doi.org/10.1111/ele.12095

Verner, D., Wilby, R., Breisinger, C., Al-Riffai, P., Robertson, R., Wiebelt, M., Kronik, J., Clement, V.,
Levine, T., Esen, F., & Roos, P. (2013). *Tunisia in a changing climate : assessment and actions*
for increased resilience and development. World Bank Publications.
https://doi.org/https://doi.org/10.1596/978-0-8213-9857-9

Visser, M. E., & Holleman, L. J. M. (2001). Warmer springs disrupt the synchrony of oak and winter
moth phenology. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
268(1464), 289-294. https://doi.org/doi:10.1098/rspb.2000.1363

Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M.,
Hoegh-Guldberg, O., & Bairlein, F. (2002). Ecological responses to recent climate change.
Nature, *416*(6879), 389-395. https://doi.org/https://doi.org/10.1038/416389a

Walther, G. R. (2010). Community and ecosystem responses to recent climate change. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *365*(1549), 2019-2024.
https://doi.org/doi:10.1098/rstb.2010.0021

Wertheim, B., Kraaijeveld, A. R., Schuster, E., Blanc, E., Hopkins, M., Pletcher, S. D., Strand, M. R.,
Partridge, L., & Godfray, H. C. J. (2005). Genome-wide gene expression in response to
parasitoid attack in Drosophila. *Genome Biology*, *6*(11), R94. https://doi.org/10.1186/gb-
2005-6-11-r94

Woods, H. A., Dillon, M. E., & Pincebourde, S. (2015). The roles of microclimatic diversity and of
behavior in mediating the responses of ectotherms to climate change. *Journal of Thermal*
Biology, *54*, 86-97. https://doi.org/https://doi.org/10.1016/j.jtherbio.2014.10.002

Wu, C. H., Holloway, J. D., Hill, J. K., Thomas, C. D., Chen, I. C., & Ho, C.-K. (2019). Reduced body sizes
in climate-impacted Borneo moth assemblages are primarily explained by range shifts.
Nature Communications, *10*(1), 4612. https://doi.org/10.1038/s41467-019-12655-y

Zittis, G., Hadjinicolaou, P., Almazroui, M., Bucchignani, E., Driouech, F., El Rhaz, K., Kurnaz, L.,
Nikulin, G., Ntoumos, A., Ozturk, T., Proestos, Y., Stenchikov, G., Zaaboul, R., & Lelieveld, J.
(2021). Business-as-usual will lead to super and ultra-extreme heatwaves in the Middle East
and North Africa. *npj Climate and Atmospheric Science*, *4*(1), 20.
https://doi.org/10.1038/s41612-021-00178-7

37