Universal protein misfolding intermediates can bypass the proteostasis network and remain soluble and less functional

Daniel Nissley[1,‡], Yang Jiang[1,‡], Fabio Trovato[1], Ian Sitarik[1], Karthik Narayan[1], Philip To[2], Yingzi Xia[2], Stephen D. Fried[2,3], Edward P. O'Brien[1,4,5*]

[1]Department of Chemistry, Pennsylvania State University, University Park, PA 16802, USA
[2]Department of Chemistry, Johns Hopkins University, Baltimore, MD 21218, USA
[3]Thomas C. Jenkins Department of Biophysics, Johns Hopkins University, Baltimore, MD 21218, USA
[4]Bioinformatics and Genomics Graduate Program, The Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA 16802, USA
[5]Institute for Computational and Data Sciences, Pennsylvania State University, University Park, PA 16802, USA
[‡]These authors contributed equally to this research project
* to whom correspondence should be addressed: epo2@psu.edu

ABSTRACT

Misfolded protein conformations with decreased functionality can bypass the proteostasis machinery and remain soluble *in vivo*. This is an unexpected phenomenon as several cellular quality control mechanisms have evolved to rid cells of misfolded proteins. Three questions, then, are: how is it structurally possible for long-lived, soluble, misfolded proteins to bypass the proteostasis machinery and processes? How widespread are these soluble, misfolded states across the proteome? And how long do they persist for? Here, we address these questions using coarse-grain molecular dynamics simulations of the synthesis, termination, and post-translational dynamics of a representative set of cytosolic *E. coli* proteins. We predict that half of all proteins exhibit subpopulations of misfolded conformations that are likely to bypass molecular chaperones, avoid aggregation, and not be rapidly degraded. These misfolded states may persist for months or longer for some proteins. Structurally characterizing these misfolded states, we observe they have a large amount of native structure, but also contain localized misfolded regions from non-native changes in entanglement, in which a protein segment threads through a loop formed by another portion of the protein that is not found in the native state. The surface properties of these misfolded states are native like, suggesting they may bypass the proteostasis machinery and its regulatory processes to remain soluble, while their entanglements make these states long-lived kinetic traps, as disentanglement requires unfolding of already folded portions of the protein. In terms of function, we predict that one-third of proteins have subpopulations that misfold into less-functional states that have structurally perturbed functional sites yet remain soluble. Data from limited-proteolysis mass spectrometry experiments, which interrogate the misfolded conformations populated by proteins upon unfolding and refolding, are consistent with the structural changes seen in the entangled states of glycerol-3-phosphate dehydrogenase upon misfolding. These results provide an explanation for how proteins can misfold into soluble conformations with reduced functionality that can bypass cellular quality controls, and indicate, unexpectedly, this may be a wide-spread phenomenon in proteomes. Such entanglements are observed in many native structures, suggesting the non-native entanglements we observe are plausible. More broadly, these near-native entangled structures suggest a hypothesis for how synonymous mutations can modulate downstream protein structure and function, with these mutations partitioning nascent proteins between these kinetically trapped states.

INTRODUCTION

How soluble, misfolded protein populations with reduced functionality[1–3] bypass cellular quality control mechanisms[4] for long time periods is poorly understood. Further, how common this phenomenon is across organismal proteomes has not been assessed. These are important gaps in our knowledge to fill as the answers will offer a more complete picture of protein structure[5] and function in cells, may lead to refinement of the protein homeostasis model[6] of proteome maintenance, are likely to be relevant to how synonymous mutations have long-term impacts on protein structure and function[7], and could reveal long-term misfolding on a scale greater than previously thought.

Protein homeostasis ("proteostasis") refers to the maintenance of proteins at their correct concentrations and in their correct conformational states through the action of a cohort of chaperones, degradation machineries, and protein quality control pathways[6,8]. It is typically posited that under normal (*i.e.*, not stressed) cellular growth conditions globular proteins *in vivo* attain one of three states: folded/functional, misfolded/aggregated, or degraded. Various molecules work together to maintain proteostasis by catalyzing the interconversion of proteins between these states[9–11]. For example, some chaperones in *E. coli*, such as GroEL/GroES[9] and DnaK[10], can promote the folding of misfolded or unfolded proteins. Others, such as the set of enzymes associated with the ubiquitin-proteasome system in eukaryotes, covalently tag misfolded proteins for degradation[12]. Yet others, such as *E. coli*'s ClpXP, have the potential to break apart aggregates, allowing released monomeric proteins to be degraded[13]. Many caveats and nuances exist in the proteostasis model. For example, some insoluble protein aggregates, such as carboxysomes, are biologically beneficial by spatially concentrating protein function[14]. Some non-native protein oligomers are soluble[15], and recently discovered biomolecular condensates[16] represent a form of phase separation in which proteins within a condensate remain soluble but preferentially interact with each other over other cellular components. Additionally, soluble proteins are not always functional because some require co- or post-translational modifications[17,18].

The timescales involved with many proteostasis processes are often quite short. Co-translationally acting chaperones bind ribosome nascent chain complexes on time scales of tens of ms[19], the ubiquitin-degradation machinery tags up to 30% of eukaryotic nascent chains for immediate degradation after synthesis[20,21], and post-translationally acting chaperones can generally refold misfolded proteins in seconds or minutes[22]. Indeed, the FoldEco kinetic model of *E. coli* proteostasis indicates that conversions between various states within the network occur with rate constants typically on the order of seconds[23]. Thus, according to the proteostasis model, misfolded proteins should either be converted in a matter of seconds or minutes into their folded state, be degraded, or form aggregates provided cells are not stressed and the proteostasis machinery is not overwhelmed[24].

Synonymous mutations change the triplet of nucleotides between degenerate mRNA codons encoding the same amino acid, leading to an altered mRNA sequence that encodes the same protein primary structure. Such mutations can alter the translation-elongation rate of ribosomes and have been found to alter the structure and function of proteins for long timescales[7]. For example, translation of a synonymous variant of the *frq* gene in the fungus *Neurospora* resulted in the synthesis of FRQ protein with altered conformations that bound 50% less to a partner protein, resulting in a significantly altered circadian rhythm that persisted for multiple days[2].

Many other proteins have been reported to exhibit altered structure or function upon the introduction of synonymous mutations[25–27]. The fact that these functional changes occur in the soluble fraction of the proteome indicates it is not insoluble aggregation driving this phenomenon. And importantly, such observations are inconsistent with aspects of the proteostasis model, which predicts that any protein with a misfolded (and less functional) structure should either refold, aggregate, or be degraded on faster timescales.

One hypothesis that could resolve this discrepancy is that proteins can populate an additional state. In this state, proteins are kinetically trapped over long timescales in misfolded conformations with reduced functionality, but they do not have a propensity to aggregate or interact with proteostasis machinery in excess of that of folded proteins. If correct, this hypothesis raises a number of questions, including: (*i*) what type of structures are adopted in this state? (*ii*) how do those conformations simultaneously avoid folding, aggregation, and degradation in excess of that observed for the native ensemble? (*iii*) how long do they persist? And (*iv*) what fraction of the proteome exhibits this behavior?

Answering these questions requires a computational method that can access the second to minute timescale of protein synthesis and maturation while providing sufficient structural resolution to identify misfolded conformations and their properties. We use a topology-based coarse-grain model that represents proteins with one interaction site per residue placed at the coordinates of the C_α atom. This model folds proteins 4-million times faster, on average, than in real systems[28], and was previously used to accurately reproduce the co-translational folding time course of HemK N-terminal domain[29]. Coarse-grain simulations of a zinc-finger protein folding in the ribosome exit tunnel were also found to agree with experimental cryo-EM structures[30], indicating this method can reproduce realistic scenarios of folding on the ribosome. Additionally, excellent agreement has been found between such topology-based models and experimental assays monitoring force generation due to the folding of titin I27 domain on and off the ribosome[31]. These examples highlight the utility of such coarse-grain models to protein misfolding on and off the ribosome.

Here, we use such coarse-grain methods to simulate protein synthesis, co-translational and post-translational folding, and estimate the fraction of molecules that fold, misfold, interact with chaperones, aggregate, are degraded, or attain a functional conformation. After first confirming that our model can reproduce post-translational misfolding in Luciferase, we simulate a representative subset of the cytosolic *E. coli* proteome, finding that a substantial proportion of newly synthesized proteins can adopt misfolded conformations that are near-native in structure and thus likely to interact with co- and post-translational chaperones in a manner similar to that of their native states. These misfolded conformations expose a similar amount of aggregation-prone surface area as the native ensemble, and therefore do not have an increased propensity to aggregate. For some proteins, misfolding is localized near their functional sites, indicating their functionality is reduced. We estimate that many of these near-native misfolded states are kinetically trapped, exhibiting lifetimes on the order of days to months. Our simulations predict that there is a universal structural feature of these proteome-wide, soluble misfolded states.

RESULTS

A coarse-grain model reproduces experimentally observed misfolding of Firefly Luciferase. The model we use for protein synthesis, folding, and function has been

shown to accurately predict experimentally measured changes in enzyme specific
activities[32], indicating it reasonably describes protein structure-function relationships.
As an additional test, here we examine if the model is able to identify if a protein will
exhibit misfolded subpopulations. Firefly Luciferase, a 550-residue protein with four
domains, folds co-translationally[33]. Specific activity experiments[25] have found that
some soluble, nascent Luciferase molecules misfold when translation speed is
increased. Even when synthesized from its wild-type mRNA in *E. coli* some Luciferase
molecules still fail to fold correctly[25]. We therefore selected Luciferase as a test
system, judging that if it partitions into long-lived misfolded states in our simulations
when translated from its wild-type mRNA that our model is able to capture realistic
scenarios of misfolding.

We simulated Luciferase's synthesis, ejection from the ribosome exit tunnel,
and post-translational dynamics using a coarse-grain representation of the protein and
ribosome (Figure 1a-c, Table S1, Methods, and Eq. 1 and 2). Fifty statistically
independent trajectories were run. To characterize Luciferase's native conformational
ensemble we also simulated ten trajectories initiated from Luciferase's crystal
structure, which we refer to as "native-state simulations". To assess whether or not a
given Luciferase trajectory is misfolded we utilize the time-dependent mode of the
fraction of native contacts (Q_{mode}) and the probability that a non-native entanglement
($P(G_k)$) has formed. We categorize a trajectory as misfolded if it has (*i*) a mean Q_{mode},
over the final 100 ns of the post-translational phase of the simulation, that is less than
the average from the native-state simulations, or (*ii*) a mean $P(G_k)$ for the different
possible changes in non-covalent lasso threading denoted $k = \{0, 1, 2, 3, 4\}$ of 0.1 or
greater over the final 100 ns of the trajectory, or (*iii*) both (*i*) and (*ii*) occur (see
Methods). Conditions (*i*) and (*ii*) correspond to perturbations of structure relative to the
native state as defined by the fraction of native contacts and entanglement,
respectively. Based on this definition, 46% (95% Confidence Interval [32%, 60%],
calculated from bootstrapping 10^6 times) of nascent Luciferase molecules misfold
(Figure 1d-e, Methods). We observe that when Luciferase misfolds, it misfolds 100%
of the time in the second domain (based on $\langle Q_{\mathrm{mode}} \rangle$), which is composed of residues
13-52 and 212-355. These misfolded Luciferase structures are near-native, with a
5.5% decrease in the overall fraction of native contacts ($\langle Q_{\mathrm{overall}} \rangle = 0.86$, computed
over the final 100 ns of misfolded trajectories) compared to the native ensemble
($\langle Q_{\mathrm{overall}} \rangle = 0.91$). Thus, a large proportion of nascent Luciferase misfolds into near-
native conformations that typically involve misfolding of the second domain.

The motivating experiments on Luciferase were carried out in the presence of
the endogenous *E. coli* proteostasis machinery[25]. To predict whether the Luciferase
misfolded states produced by our model are likely to display reduced specific activity
in vivo we therefore need to determine four things. They must (*i*) evade chaperones to
remain misfolded, (*ii*) not aggregate, (*iii*) not get degraded, and (*iv*) the residues
involved in function must be structurally perturbed. The chaperone trigger factor (TF)
binds nascent proteins co-translationally, DnaK interacts both co- and post-
translationally, while GroEL/GroES is primarily a post-translational chaperone.
Interactions with TF[34] or GroEL/GroES[9] are thought to occur by the non-specific
recognition of exposed hydrophobic patches on client proteins co- and post-
translationally, respectively. DnaK, however, is hypothesized to interact with specific
binding sites within a protein's sequence[35].

To estimate whether misfolded Luciferase is likely to interact with TF we
computed the average relative difference between the hydrophobic solvent accessible
surface area (SASA) of each misfolded trajectory to the folded population (denoted

$\langle \zeta^{\text{co-t}}_{\text{hydrophobic}} \rangle$, Eq. 9 and Methods) during synthesis. The value of $\langle \zeta^{\text{co-t}}_{\text{hydrophobic}} \rangle$ is ≤10%
for 16 of 23 misfolded trajectories, meaning that they display less than a 10% increase
in hydrophobic SASA during synthesis relative to the folded population of trajectories.
This indicates that a majority of misfolded Luciferase molecules will not interact with
TF much more than a properly folded Luciferase molecule (see Methods). Though TF
accelerates protein folding under force[36], under normal conditions it is also thought to
act as a holdase[34]. Thus, we conclude that our co-translational Luciferase misfolded
states can misfold into conformations that do not interact with TF in a manner that
accelerates folding, allowing these misfolded states to persist post-translationally.
Next, to determine whether misfolded conformations of Luciferase are likely to
interact with GroEL/GroES post-translationally, we computed the average relative
difference between the hydrophobic SASA of each misfolded conformation in the final
100 ns and its value in the native-state simulations ($\langle \zeta_{\text{hydrophobic}} \rangle$, Eq. 10, Methods).
The value of $\langle \zeta_{\text{hydrophobic}} \rangle$ for Luciferase is ≤10% for 9 of 23 misfolded trajectories,
indicating that these misfolded states expose only a small excess of hydrophobic
SASA relative to the native ensemble, and are therefore not likely to be engaged by
GroEL/GroES.
Finally, to estimate whether misfolded Luciferase structures are more likely to
interact with DnaK than the native state, we computed $\langle \zeta_{\text{DnaK}} \rangle$ (Eq. 11), the average
relative difference in SASA of residues predicted to be DnaK binding sites by the Limbo
algorithm[35] in the final 100 ns for all misfolded trajectories. We find that 22 out of 23
misfolded trajectories have $\langle \zeta_{\text{DnaK}} \rangle \leq 10\%$, indicating that DnaK is unlikely to
preferentially bind to these misfolded states any more than it is to the native state.
Thus, some of Luciferase's misfolded states are unlikely to interact with TF,
GroEL/GroES, or DnaK, and thus bypass the *E. coli* chaperone network (Figure S1a).
The next key question is whether or not these misfolded Luciferase structures,
having bypassed chaperone quality controls, are likely to remain soluble or to
aggregate or be degraded. In the original experiments by Barral and co-workers,
centrifugation was used to remove aggregates from the soluble fraction. To estimate
whether the misfolded Luciferase structures from our simulations will aggregate we
used the AMYLPRED2[37] webserver to identify residues in the Luciferase amino acid
sequence that lead to aggregation when exposed to solvent. We then computed $\langle \zeta_{\text{agg}} \rangle$
(Eq. 12), the average relative difference in SASA between these aggregation-prone
residues in the final 100 ns of each misfolded trajectory in comparison to the same
residues in the native-state simulations. We find that 12 of 23 misfolded trajectories
have $\langle \zeta_{\text{agg}} \rangle \leq 10\%$, indicating that these misfolded conformations display only a minor
increase in aggregation propensity and are likely to remain soluble.
Finally, we considered the likelihood that misfolded Luciferase will be targeted
for degradation. Degradation in *E. coli* is carried out primarily by proteases coupled to
AAA+ ATPase motor proteins[38], including ClpXP and Lon, that recognize and degrade
misfolded or aggregated proteins. Misfolded protein structure contributes to
degradation[39], and therefore, like our GroEL/ES assessments, we use $\langle \zeta_{\text{hydrophobic}} \rangle$ to
quantify how similar misfolded Luciferase conformations are to the native state. For 9
of 23 misfolded Luciferase trajectories $\langle \zeta_{\text{hydrophobic}} \rangle$ is $\leq 10\%$, indicating they are
unlikely to be degraded more quickly than native Luciferase.
Having determined that some misfolded conformations of Luciferase can evade
chaperones, aggregation, and degradation, the final question is whether their function
is decreased relative to native Luciferase. To answer this question, we identified the
residues that take part in Luciferase's bioluminescence, defined as those resides

within 4.5 Å of the 5′-O-[N-(dehydroluciferyl)-sulfamoyl]-adenosine ligand in PDB
structure 4G36, in addition to all residues identified in the UniProt database[40] to have
a role in its catalytic mechanism. To quantify the difference in structure of residues
involved in Luciferase's catalytic mechanism, we compute the average relative
difference between the structures sampled in the final 100 ns of each misfolded
trajectory and native Luciferase in terms of the structural overlap function ($\langle \chi_{\text{func}} \rangle$) over
residues implicated in its function (see Eq. 13-15 and Figure 1b, c, f, g, and h). Positive
values of χ_{func} indicate perturbed structure relative to the native state simulations. We
find that 15 of 23 misfolded Luciferase trajectories have $\langle \chi_{\text{func}} \rangle \geq 10\%$, indicating that
they have significantly perturbed structure at functionally important sites relative to the
native state (Figure 1f). For example, Figures 1g and 1h show the binding pocket at
the final frames of folded and misfolded, soluble, but non-functional Luciferase
trajectories. The binding pocket structure is perturbed such that it impinges on the
substrate location. Since structure equals function, this result indicates that the
efficiency of the enzymatic reaction carried out by misfolded Luciferase will be less
efficient than in its native fold.
Cross-referencing the lists of misfolded trajectories that are likely to avoid
chaperones, aggregation, degradation, and exhibit reduced function, we find that one
trajectory displays all of these characteristics and likely remains soluble but less
functional than native Luciferase (Figure S1b). Our simulation results are thus
qualitatively consistent with the experimental observation that some nascent
Luciferase molecules misfold when translated from its wild-type mRNA. While a
misfolded state that is only populated by 2% of protein molecules is unlikely to strongly
influence the cell, perturbations to Luciferase translation-elongation kinetics by
synonymous mutations might increase this population beyond 2%. In general, these
results indicate that our coarse-grain simulation protocol for nascent protein synthesis,
ejection, and post-translational dynamics is able to recapitulate nascent protein
misfolding.
320
Simulating a representative subset of the _E. coli_ cytosolic proteome. It is not
computationally feasible to simulate all 2,600 cytosolic _E. coli_ proteins. Therefore, to
investigate the extent of nascent protein misfolding within the _E. coli_ proteome we
constructed models for a representative subset of 122 proteins. This set of proteins
has the same distributions of protein length and structural class, and a similar ratio of
multi- to single-domain proteins as the entire _E. coli_ proteome[41] (Table S2). The
details of the parameterization of these models are described in Ref. 41. Each protein
was synthesized on the same coarse-grain ribosome representation as Luciferase and
their post-translational dynamics simulated for 30 CPU days per trajectory. Larger
proteins take longer to simulate. Therefore, this fixed post-translational simulation run
time resulted in trajectories of different durations due to different protein sizes. The
simulation time in the post-translational phase therefore ranged between 2.7 and
154.1 μs per trajectory. Because our coarse-grain model exhibits an approximately
four-million fold acceleration of folding dynamics[28], due to decreased solvent
viscosity[42] and a smoother free-energy landscape[43], these post-translational
simulation times correspond approximately to experimental times of 11.0 to 611
seconds, respectively. As with Luciferase, ten trajectories were also initiated from the
crystal structure of each protein and simulated for 30 CPU days to serve as reference
simulations representing the native-state structural ensemble.
340

Two thirds of nascent *E. coli* proteins populate misfolded states. A fundamental question our simulation data set can address is how common nascent protein misfolding is across *E. coli*'s cytosolic proteome. As with Luciferase, we use the fraction of native contacts and entanglement as measures of misfolding. We find that 66% of proteins (80 out of 122) remain misfolded in at least one trajectory, 40% of proteins are misfolded in at least 20% of trajectories (49 out of 122), and 7% are misfolded in 100% of trajectories (9 out of 122). The proteins in these various categories are summarized in Table S4; Figure 2a displays a histogram of the probability of misfolding over the 122 different *E. coli* proteins simulated. In total, 27% of the simulation trajectories (1,631 out of 6,100) of the *E. coli* cytosolic proteome remain in misfolded conformations after 30 CPU days of post-translational dynamics.

Many misfolded states are similar to the native state. Misfolded conformations that are very different from the native state will likely interact with the proteostasis machinery. To characterize the closeness of misfolded states to the native state across our set of misfolded conformations we calculated the absolute percent change in the mean overall fraction of native contacts $\langle Q_{overall} \rangle$ (in this case, computed over all residues in secondary structures within each protein, rather than for individual domains or interfaces) between each protein's native state simulations and the mean Q in the final 10 ns of each misfolded trajectory (Figure 2b). We observe that 76% of misfolded trajectories (1,242 out of 1,631) have ≤20% change in mean Q in comparison to the native state, while 58% of trajectories (939 out of 1,631) misfold and have a ≤10% change in Q. 9% of trajectories (144 out of 1,631) have a ≤1% change in Q. These calculations indicate that a large proportion of trajectories that misfold populate states that are native-like. Therefore, many *E. coli* proteins can populate kinetically trapped near-native conformations that are structurally similar to the native state.

Misfolded states can persist for days or longer after release from the ribosome. Misfolded conformations that persist for just a few minutes before properly folding are unlikely to have downstream consequences in a cell. To estimate the range of folding times for misfolded conformations we computed the survival probability of the unfolded state, $S_U(t)$, for each protein domain and interface and extracted their characteristic folding timescales using a three-state folding model, which reports folding timescales for the fast- and slow-folding phases (see Methods). For a protein to be considered folded all its component domains and interfaces must be folded. Furthermore, since folding pathways that pass through misfolded states take longer to reach the native state, the slow-folding phase reflects the time scale of these pathways. Therefore, for a given protein, we interpret the longest, slow-folding phase time as the time scale of the misfolded state reaching the native state. In total, we are able to reliably determine folding times for 73 out of 122 proteins, with fit equations for other domains having small Pearson R^2 values indicative of low-quality estimates. These extrapolated folding times for the slow phase were then mapped onto experimental times using the acceleration factor associated with the coarse-grained model[28] (see Methods). The 25th, 50th, 75th, and 95th percentile mean folding times for the slow phase are 1.41 s, 50.9 s, 1.19 x 10^7 d, and 3.83 x 10^{16} d, respectively, and the full range of times extend from 0.04 s to 1.08 x 10^{22} d (Figure 2c, Table S5). While values at very long times have larger uncertainties, as small differences in the fit parameters will lead to large variation in the extrapolated folding times, these results clearly indicate that many of these misfolded states can persist for many days or longer after synthesis.

Half of the proteome misfolds and bypasses the chaperone machinery. Misfolded proteins are engaged by various chaperones both co- and post-translationally that help direct their correct folding. So, we next determined how many of the trajectories that exhibit misfolding in our simulations are likely to evade chaperone-dependent quality control mechanisms. As was done for Luciferase, we considered the interactions of each of our 1,631 misfolded trajectories with TF, GroEL/GroES, and DnaK based on the relative difference between the SASA of specific subsets of residues in the misfolded ensemble versus the native state ensemble (see Methods, Eqs. 9-11). We find that 1,053 misfolded trajectories, representing 70 unique proteins, are not likely to interact with TF, as they display $\langle \zeta_{\text{hydrophobic}}^{\text{co-t}} \rangle \leq 10\%$ or are too short to engage with it co-translationally (see Methods, Table S6, and Figure S2). A total of 1,411 of misfolded trajectories representing 80 unique proteins are either not known GroEL/GroES[44–46] clients or have $\langle \zeta_{\text{hydrophobic}} \rangle \leq 10\%$, and are therefore not likely to interact excessively with GroEL/GroES (Table S6, Figure 2d). Finally, we find that 1,115 misfolded trajectories representing 74 unique proteins are either not confirmed DnaK clients or have $\langle \zeta_{\text{DnaK}} \rangle \leq 10\%$ and are therefore unlikely to interact with DnaK excessively (Figure S3). A total of 731 trajectories representing 64 different proteins are misfolded and unlikely to interact with TF, GroEL/GroES, or DnaK (Table S6, Figure S4). These results indicate that 52% of proteins in our representative sample (64 out of 122 unique proteins) exhibit misfolded subpopulations that can bypass chaperones.

Half of the proteome misfolds and remains soluble. We next assessed how many of the 1,631 trajectories in which the protein misfolds represent conformational states that are likely to remain soluble. For each protein we computed $\langle \zeta_{\text{agg}} \rangle$, the average relative difference in SASA of residues predicted to be aggregation prone computed over the final 100 ns for each misfolded trajectory, to quantify the difference in aggregation propensity for the misfolded population relative to the native state simulations (see Methods, Eq. 12). Of the 1,631 misfolded trajectories, 814 have $\langle \zeta_{\text{agg}} \rangle \leq 10\%$, indicating they are not likely to aggregate in excess of what is observed for the native state (Table S7, Figure S5). We conclude that these trajectories, representing 56% of the proteins in the sample (68 out of 122), are unlikely to aggregate.

Half of the proteome misfolds and does not exhibit excess degradation. Next, we examined how many misfolded proteins are likely to avoid rapid degradation. We did this by computing $\langle \zeta_{\text{hydrophobic}} \rangle$, which characterizes the percent difference between the total hydrophobic SASA of misfolded trajectories in comparison to the set of native-state simulations (Eq. 10). The values of $\langle \zeta_{\text{hydrophobic}} \rangle$ for 896 misfolded trajectories are ≤10%, indicating they are unlikely to be targeted for degradation. These 896 misfolded trajectories predicted to bypass degradation represent 57% (70 out of 122) unique proteins (Table S7, Figure 2d). Thus, a majority of proteins can populate, to varying degrees, misfolded states that are not expected to be degraded at rates much faster than their native fold.

Half of the proteome misfold into conformations that bypass all aspects of the proteostasis machinery in *E. coli*. Misfolded conformations that do not engage chaperones, do not aggregate, and are not degraded in excess of the native state will

remain soluble within the cell for a similar time scale as the native state. We cross-
referenced our lists of misfolded trajectories that fall into each of these categories,
finding that 8% of all trajectories simulated (517 out of 6,100) misfold into such soluble
conformations, and 47% of proteins (57 out of 122) have at least one such trajectory
(Table S8). Thus, nearly half of proteins in our sample have subpopulations of
misfolded states that will bypass all aspects of protein homeostasis and stay misfolded
for biologically long time periods.
447
Half of the proteome misfolds and will exhibit altered function. Next, we examined
what percentage of the proteome misfolds and is likely to exhibit reduced function. To
answer this question, we constructed a database identifying residues that take part in
the function of each of the 122 proteins in our data set based on information available
in PDB and UniProt database entries. These functional residues were identified based
on whether they were in contact with substrates (such as other biomolecules, small-
molecule compounds, or ions) in their PDB structures, as well as based on UniProt's
identification of functional residues (see Methods). We then computed the mean
relative difference in the structural overlap function of these functional residues in the
final 100 ns of misfolded trajectories relative to the native state reference simulations
($\langle \chi_{\text{func}} \rangle$, see Methods). We find that 62% of misfolded trajectories (1,019 out of 1,631)
have $\langle \chi_{\text{func}} \rangle \geq 10\%$, indicating that structure at their functional sites are significantly
perturbed as well as their function. These trajectories represent misfolded
conformations of 69 unique proteins, indicating that 57% of the proteome can populate
misfolded conformations likely to exhibit reduced function (Table S8, Figure 2e).
463
One-third of proteins exhibit soluble, misfolded, native-like states with reduced
functionality. We next determined which of our 122 proteins misfold, evade
chaperones, aggregation, degradation, and display reduced function. We find that
31% of proteins (38 out of 122) and 3% of all trajectories (186 out of 6,100) can bypass
proteostasis machinery and display decreased function (Table S8, Figure 2f). The
extrapolated folding times of these 38 soluble but non-functional proteins range from
2.13 s to 1.07 x 10^{22} days with a median predicted folding time of 1.05 x 10^{12} days,
indicating that their function is likely to be perturbed for long timescales.
472
Intra-molecular entanglement drives long-lived, soluble misfolded
conformations. To determine what characteristics, if any, the misfolded
conformations of different proteins have in common, we used the Gauss linking
number calculated from linking between a closed loop formed by a native contact
between residues i and j and the pseudo-closed loops formed by the flanking termini,
$g(i, j)$[47]. This quantity provides a useful measure of whether subsections of the protein
chain are entangled with each other[48]. Misfolding, or changes in the linkage between
the two closed loops, can then be identified by changes in the Gauss linking number
of specific native contacts between a reference structure and a target structure (Figure
3a, b).
To determine whether or not misfolded states tend to be entangled, we
generated a 2-by-2 contingency table (Figure 3c) tabulating the co-occurrence of
misfolding based on $\langle Q_{\text{mode}} \rangle$ and the presence of an entanglement. We find an odds
ratio of 48.1 ($p < 10^{-100}$, Fisher's Exact test), indicating that entanglement and
misfolding frequently co-occur, with 82% of misfolded states containing an
entanglement. Thus, misfolding is predominantly driven by entanglement of segments
of the nascent protein with each other.

We hypothesized that due to the large energetic barrier needed to disentangle
entanglements, the most long-lived misfolded states in the *E. coli* proteome would tend
to be entangled. To test this hypothesis, we generated a second 2-by-2 contingency
table and counted how frequently slow- and fast-folding proteins tend to be entangled
(Figure 3d). Proteins with an extrapolated folding time for the slow phase greater than
the median were considered to be slow folding; and a protein's misfolded state is
considered entangled if ≥50% of its misfolded trajectories display an entanglement.
We find an odds ratio of 15.0 ($p = 5.0 \times 10^{-7}$, Fisher's Exact test) indicating that the
presence of entangled misfolded structures are 15 times more likely to be associated
with slow folding. Thus, entanglement is the primary cause of long-lived misfolded
states.
Finally, we further hypothesized that, because entangled conformations can
represent local minima with only small structural perturbations relative to the native
state, the set of 186 trajectories predicted to bypass proteostasis machinery to remain
soluble but non-functional for long timescales should be enriched in entangled
structures. We find that 94% of these trajectories are entangled (174 out of 186), and
that there is a strong association between escaping proteostasis machinery and the
presence of an entanglement (odds ratio 59.2, $p = 3.4 \times 10^{-102}$, Fisher's Exact test;
Figure 4).
Taken together, these results demonstrate that the formation of entanglements
entangled misfolded states lead to long-lived kinetic traps that can bypass the
proteostasis machinery.
512
An in-depth case study. To illustrate our key findings, it is useful to consider the
structural basis of misfolding for one protein in-depth. We focus on glycerol-3-
phosphate dehydrogenase, which has the largest proportion of misfolded trajectories
that are predicted to bypass the proteostasis machinery (see Figure 4). It consists of
two domains composed of residues 1-387 and 388-501 (Figure 5a). As part of its
biological function, glycerol-3-phosophate dehydrogenase uses a flavin adenine
dinucleotide cofactor (Figure 5a, dark blue). In our post-translational simulations 74%
(=37/50) of trajectories misfold, yet they only exhibit a 4.7% decrease in the fraction of
native contacts relative to the native-state simulations (Figures 5b, c). Thus, these
misfolded states resemble the native ensemble. This protein also folds extremely
slowly, with Domain 1, Domain 2, and the interface between Domains 1 and 2
estimated to require, respectively, on the order of 10^{16}, 10^{15}, and 10^{21} seconds to fold
(Figure 5d). These misfolded states are, however, expected to evade chaperones,
aggregation, and degradation to remain soluble based on the similarity of their surface
properties to that of the native ensemble (Figure 5e). Twenty-five misfolded
trajectories (50%) also exhibit notably reduced structure at functional sites, including
around the cofactor, despite being well folded overall (Figure 5f, g, and h). In 92%
(=34/37) of these misfolded trajectories a non-native entanglement is present. These
results exemplify how entangled misfolded states can perturb portions of a protein
critical for function in ways that are structurally subtle compared to gross deformations
typically associated with misfolded proteins (Figure 5d and h).
534
An experimental test for structural changes associated with entanglement. To
test these predictions for glycerol-3-phosphate dehydrogenase we carried out
protease digestion mass spectrometry (see Methods) in which whole extracts from
cells were globally unfolded by incubation in 6 M guanidinium chloride, and refolded
by rapid dilution. The structures of the refolding proteins were then interrogated with

pulse proteolysis with proteinase K (PK), which specifically cuts at exposed or
unstructured sites. The resulting fragments were identified and quantified with mass
spectrometry and compared to those from native lysates that were never unfolded.
Protease digestion was carried out at 1-min, 5-min, and 120-min timepoints after
refolding conditions were established, and glycerol-3-phosphate dehydrogenase's
digestion pattern is observed to change over these time points (see Methods and
Supplementary Data File 1). We consider in our analysis only those peptides that show
a greater than 3.5-fold difference in abundance in the refolded sample versus native
sample ($\left| \log_2 \left(\frac{R}{N} \right) \right| > 1.8$, Column W in Supplementary Data File 1 sheet labeled
"GlpD") and whose difference is statistically significant ($p < 0.01, -\log_{10}(p) > 2$,
Column Y in same sheet, see Methods[49]). A total of ten unique peptides meet these
criteria at one or more experimental time points. At 1 min residue V203 and residues
[333-354] are significantly more exposed in the refolded sample than in the native
sample. At 5 min L293, F351, Q487, and P387 are more exposed in the refolded than
native sample, while V203 and [333-354] are no longer found to be different between
refolded and native. After 120 min, seven exposed peptides are found: [333-354],
F351, and L293 once again appear more exposed in refolded than native, while Y313,
[284-302], D437, and G422 emerge as more exposed in the refolded sample. No one
peptide is found to be more exposed in the refolded than native samples at all three
time points, though L293, F351, and [333-354] are more exposed at two time points.
These experimental data indicate that some glycerol-3-phosphate dehydrogenase
molecules that fail to arrive at the native structure rapidly populate misfolded
structures.
To test if the entanglements we observe in simulations of glycerol-3-phosphate
dehydrogenase can explain these digestion patterns, we structurally clustered the
coarse-grain conformations from the final 100 ns of our simulations (based on their G
and Q_{overall}) into eight metastable states denoted {S1, S2, …, S8}. We focus on
segment [333-354] and residues F351 and L293 because these peptides persist in the
experiments, being present at either the 1- or 5-min timepoints *and* the 120-min
timepoint. In seven out of eight states an entanglement is present, with states S5, S7,
and S8 the most native like (Table S11). The entangled loop or threading segments in
these states overlap with one or more peptide fragments in five out of eight states.
Structure near cleavage sites is still perturbed even when the entangled region does
not overlap with them. For example, the threading of residues 218-237 through the
loop formed by residues 271-288 in S2 does not contain residues L293, F351, or
segment [333-354]. However, this entanglement increases the solvent accessible
surface area of these segments (Figure 5i). This is most clearly seen for F351, which
in the native state forms part of a β-sheet buried beneath the threading segment
residues (Figure 5i, middle panel, "Folded" structure). When this set of residues
becomes entangled by threading through the loop ("Misfolded" structure in Figure 5i),
the thread is kinetically trapped in a position that exposes F351 much more than in the
native fold. Calculating the solvent accessible surface area change (Eq. 16) of these
fragments in each metastable state we find broad agreement with the experimental
data (Table S11). Each of the seven entangled metastable states displays increased
solvent accessibility at each of the three locations.
585
586
587
588

DISCUSSION

Previous work has established that soluble, long-lived, non-functional protein misfolded states can arise from alteration of translation-elongation kinetics. To the best of our knowledge, this study is the first to estimate the extent of this phenomenon across the nascent proteome of an organism and examine the structural and kinetic properties of these kinetically trapped states. We predict that a majority of cytosolic *E. coli* proteins exhibit subpopulations of misfolded, kinetically trapped states, and that many of these misfolded states are similar enough to the native state to evade the proteostasis machinery in *E. coli*. We estimate that one-third of cytosolic *E. coli* proteins have subpopulations that misfold into near-native conformations that have reduced function and bypass the proteostasis network to remain soluble and non-functional for days or longer.

To appreciate these results, it is useful to understand the types of misfolding that can and cannot occur in our simulation model. The coarse-grain forcefield is parameterized for each protein based on its crystal structure, with this native-state conformation encoded as the potential energy minimum in the form of a Gō-based energy function (Eq. 1). This means that the native state is the global free energy minimum at our simulation temperatures; any other state is metastable. Another consequence of this type of model is that misfolding involving non-native tertiary structure formation is not possible. Thus, the misfolded states our model can populate are topologically frustrated states that are kinetic traps[50]. A kinetic trap is a local minimum separated from other conformations in the ensemble by energy barriers much larger than thermal energy, making the attainment of the native state a slow process for some protein subpopulations. In our model, intra-molecular entanglements can occur. These entanglements consist of two parts: a contiguous segment of the protein that forms a 'closed' loop, where the loop closure is geometrically defined as a backbone segment that has a native contact between two residues at its ends, and another segment of the protein that threads through this loop (Figure 3).

Proteins can misfold by a variety of mechanisms. For example, Bitran and co-workers[51] suggest that non-native contacts appear to play an important role in kinetic trapping of some proteins. Misfolding has also been observed via domain swapping[52], in which highly similar portions of proteins swap with one another. Our results are not mutually exclusive with these other types of misfolding; indeed, one can imagine situations in which domain swapping involves the introduction of an entanglement, or in which non-native contacts form entanglements. Understanding the overlap and interplay of these various types of misfolding in real systems is an interesting open question.

It was previously hypothesized[53] that like protein topological knots[54] (which persist when pulling on both termini), this type of entanglement, which we refer to as a non-covalent lasso entanglement[55], would generate topological frustration and be a kinetic trap. Simulations of proteins with topological knots in their native state[50] observed that the wrong knot could form and that many of these states were long-lived kinetic traps as they required 'backtracking'[56] (*i.e.*, unfolding) to fix the knot. While only 3 of the 122 proteins (gene names rlmB, metK, and rsmE) in our study contain topological knots in the native state, the non-covalent lasso entanglement intermediates we observe are non-native pseudoknots[54] (which unravel when pulling on both termini), and require either reptation of the threaded protein segment out of the closed loop (Figure 3) or local unfolding of the loop surrounding the threaded segment to disentangle. Thus, the results of this study bridge the rich field of polymer

topology with biologically important consequences for in vivo protein structure and
function.
A potential criticism of this work is that the non-native entanglements we
observe may be an artifact of our coarse-grained modeling of proteins. Several lines
of evidence indicate this criticism is unfounded. A sufficiently long linear polymer
performing a random walk will always sample knotted structures. Thus, it is a
fundamental polymer property that knots and entanglements have the potential to
form[57]. In a recent study, four entangled structures produced from our coarse-grained
model were back-mapped and simulated using classical, all-atom molecular
dynamics[32]. The entanglements, and native-like structure of these states persisted for
the entire 1 μs simulation time. In another study, one-third of the protein crystal
structures in the CATH database were found to contain in the native state the same
types of entanglements we observe as intermediates[58]. Taken together, these results
indicate that the entanglements we observe are realistic non-native intermediates that
have the potential to be populated by many proteins.
Two differences between our simulations and the limited-proteolysis data we
compare to lie in the preparation of the proteins and limits of detection. In the
experiment proteins are prepared in a chemically denatured state and then allowed to
refold, compared to folding concomitant with or after translation. Protein's that are
prone to misfolding during translation are likely to be prone to misfolding during bulk
refolding. Thus, while it is not necessary that the same misfolded states be populated
under these two different situations, the consistency between the misfolded entangled
states of glycerol-3-phosphate dehydrogenase and the persistent and significant
protease fragments from the experiment indicates similar misfolded states do occur.
Secondly, our computational workflow can detect proteins that misfold as little as 2%
of the time (1 misfolded trajectory out of 50); on the other hand, to filter signal from
noise, protein regions are only considered more (or less) exposed in the refolded form
relative to native if the corresponding PK-fragment is >2-fold (*i.e.*, $\log_2\left(\frac{R}{N}\right) > 1$ with p
< 0.01) more (or less) abundant in the proteolysis reaction. Thus, the sensitivity of this
approach to misfolded states with low populations is lower, and needs to be
considered when comparing to the simulations results.
The majority of proteins simulated in this study misfold in some capacity. And
38 unique proteins, or about one-third of the proteins we simulated, have one or more
trajectories that remain soluble and non-functional due to misfolding. Projecting this
proportion across the entire set of 2,600 proteins that make up the cytosolic *E. coli*
proteome, we estimate that approximately 806 proteins may exhibit misfolding into
soluble states. Given that a reduction in the function of a protein has the potential to
influence multiple cellular processes, this result suggests that these misfolded states
could exert wide-spread influences on cell behavior and phenotype. We also note that
it is not only proteins that bypass all aspects of proteostasis and remain non-functional
that can negatively impact cells. For example, protein conformations that avoid
chaperones and degradation but then go on to aggregate may lead to the
accumulation of amyloid fibrils.
Changes to the speed of translation, such as those that may be introduced by
synonymous mutations in a protein's mRNA template, can strongly influence the ability
of proteins to fold[7]. The simulation results described here were generated using the
wild-type translation rate profile for each protein (see Methods). Experiments have
shown that changing translation speed can alter the subpopulation of soluble, less
functional states several fold. Thus, the population of proteins that misfold have the

potential to be significantly altered in our computer simulations what translation-
elongation rates are altered. These complexities make exploration of the influence of
translation kinetics on the propensity of the *E. coli* proteome to misfold an interesting
direction for future research.
One of the most fundamental timescales of a protein is its half-life, which gives
a measure of the lifetime of a typical copy of a protein between its synthesis and
degradation. If misfolded states persist on the same timescale as the protein half-life
then protein function will be perturbed for most of that protein's existence.
Unfortunately, we are unaware of any proteome-wide studies of protein half-lives in *E.*
coli. However, based on studies of protein lifetimes in budding yeast[59] and human
cells[60], which found median half-lives of 43 min (range 2 min to 81 days) and 36 h
(range: 8 h to 153 days), respectively, we estimate that typical half-lives in *E. coli* range
from minutes to hours. Many of our extrapolated folding times from our simulations are
on the same order of magnitude as these values or greater, indicating that misfolded
but states with reduced function can persist for the entire lifetime of a protein. This is
consistent with the experimental observation that misfolding can influence folding and
function for extended periods[1,2,61].
If the half-life is a fundamental time scale of a protein, then the cell-division time
is a fundamental time scale of a bacterium. In *E. coli*, doubling times during exponential
growth phase range from tens of minutes to hours depending on the growth medium[62].
A total of 31 of our 122 proteins have extrapolated folding times for the slow phase
longer than 40 min. And, of the 38 proteins that misfold into soluble but less-functional
states, 14 have extrapolated folding times longer than 40 min. Since these folding
times are on a similar time scale as the doubling time, soluble misfolded conformations
will be split between the daughter cells. This suggests that the memory of those events
can be encoded in these kinetically trapped states and transferred to the daughter
cells. It will be an interesting area of future study to determine whether inheritance of
soluble, misfolded proteins with potentially altered function can act as a mechanism
for epigenetic inheritance and influence daughter cell behavior.
A key question potentially addressed by our simulations is what allows these
misfolded states to remain misfolded in non-functional states for such long timescales?
Entanglements appear to allow these misfolded states to persist for long time scales,
and their largely native topologies mean they are not excessively acted upon by the
proteostasis machinery. In many instances, large-scale unfolding would need to take
place in order for the entangled protein to disentangle[50] to a state from which the native
fold is more readily accessible. One interesting avenue for future research is
comparison of our results concerning misfolding after ribosomal synthesis with
simulations of refolding from denatured chains. Such work would provide a clearer
comparison to LiP-MS experiments and enable us to test the hypothesis of whether
protein synthesis reduces protein misfolding.
In summary, we have found that the majority of *E. coli* proteins misfold in our
simulations, and that some proteins misfold into states that likely bypass cellular
proteostasis machinery to remain soluble but with reduced function. We find that these
misfolded conformations are able to remain soluble because they are, overall, very
similar to the native state, but with certain entanglements that lead to perturbed
structure and function. Given that self-entanglement is a fundamental polymer
property, the entanglements we have observed represent a universal type of
misfolding that has the potential to impact a range of proteins and functions.
Specifically, our simulation results suggest the hypothesis that entangled states may
be the source of reduced dimerization[2], enzymatic function[63], and small-molecular

transport[27] upon changes in translation kinetics induced by synonymous mutations.
Future theoretical and experimental efforts should focus on the structural
characterization of non-native entangled states and their influence on protein function.
741
742 **METHODS**
743
Selection of proteins and parameterization of their coarse-grain models. A data
set of 50 multi- and 72 single-domain proteins was selected at random from a
previously developed database of *E. coli* proteins with solved X-ray diffraction or NMR
structures[41,64]. This data set contains proteins with realistic distributions of protein size
and structural class (see Figure S1 and Table S1, respectively, of Ref. 41). Small
sections of missing residues (<10) were rebuilt and minimized in CHARMM, while
large missing sections for some multi-domain proteins were rebuilt based on
homologous protein structures (see Table S4 of Ref. 41). Each of the rebuilt all-atom
models was then converted to a Cα coarse-grain representation. The potential energy
forcefield of this coarse-grain model is given by the equation

$$E = \sum_i k_b (r_i - r_0)^2 + \sum_i \sum_j^4 k_{\varphi,ij} \left(1 + \cos[j\varphi_i - \delta_{ij}]\right) + \sum_i -\frac{1}{\gamma} \ln \left\{ \exp[-\gamma(k_\alpha(\theta_i - \theta_\alpha)^2 + \right.$$

$$\left. \varepsilon_\alpha)] + \exp\left[-\gamma k_\beta \left(\theta_i - \theta_\beta\right)^2\right] \right\} + \sum_{ij} \frac{q_i q_j e^2}{4\pi\varepsilon_0\varepsilon_r r_{ij}} \exp\left[-\frac{r_{ij}}{l_D}\right] + \sum_{ij \in \{NC\}} \epsilon_{ij}^{NC} \left[13\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - \right.$$

$$\left. 18\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{10} + 4\left(\frac{\sigma_{ij}}{r_{ij}}\right)^6\right] + \sum_{ij \notin \{NC\}} \epsilon_{ij}^{NN} \left[13\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - 18\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{10} + 4\left(\frac{\sigma_{ij}}{r_{ij}}\right)^6\right]. \quad [1]$$

These forcefield terms represent, from left to right, the contributions from Cα − Cα
bonds, torsion angles, bond angles, electrostatic interactions, Lennard-Jones-like
native interactions, and repulsive non-native interactions to the total potential energy.
Full details of the model parameters can be found in Ref. 41. The value of ϵ_{ij}^{NC}, which
determines the global energy minimum for a native contact, is calculated as

$$\epsilon_{ij}^{NC} = n_{ij}\epsilon_{HB} + \eta\epsilon_{ij}. \quad [2]$$

In Eq. 2 ϵ_{HB} and ϵ_{ij} represent energetic contributions from the hydrogen bonds and
van der Waals interactions between residues i and j found within the all-atom structure
of the protein, respectively. n_{ij} is the integer number of hydrogen bonds between
residues i and j and $\epsilon_{HB} = 0.75$ kcal/mol. The value of ϵ_{ij} is initially set based on the
Betancourt-Thirumalai potential[65] and the value of η for each individual domain and
interface set based on a previously published training set[28]. The values of η used for
all production simulations are listed in Ref. 41 Tables S2 and S3 alongside all protein
names and the chain identifiers used during model building. For simplicity, all proteins
are referred to using the PDB ID of the entry from which they were primarily derived.
The parameters for the coarse-grain model of firefly luciferase (PDB ID: 4G36) have
not previously been reported and are therefore provided in Table S1.
774
Simulations of nascent protein synthesis, ejection, and post-translational
dynamics. All simulations were performed using CHARMM and the coarse-grain
forcefield described in Eq. 1 with an integration timestep of 0.015 ps and a Langevin
integrator with friction coefficient of 0.050 ps^{-1} at a temperature of 310 K. The synthesis
and ejection of each protein was simulated using a previously published protocol and
a coarse-grain cutout of the ribosome exit tunnel and surface (for complete simulation
details see Ref. 41). In this model, ribosomal RNA is represented by one interaction
site each for each ribose sugar, phosphate group, and pyrimidine base and two
interaction sites for each purine bases. Ribosomal proteins are represented at the Cα

level. Post-translational dynamics simulations were initiated from the final protein
structure obtained after ejection with the ribosome deleted. Fifty statistically
independent trajectories were run for each of the 122 proteins in the *E. coli* cytosolic
proteome data set and for Luciferase. Post-translational dynamics was run for 30 CPU
days for each trajectory. Ten trajectories were also initiated from the native-state
coordinates for all proteins and run for 30 CPU days each to provide a realistic
reference ensemble for each protein's folded state.
791
Identification of misfolded trajectories. Two order parameters, Q_{mode} and
$P(G_k \mid PDB, traj)$, were used to determine whether or a not a given trajectory folds.
Detailed definitions of these order parameters are given in the following two Methods
sections. A given trajectory is considered to be misfolded if either its Q_{mode} or
$f_c(G_k \mid PDB, traj)$ values (or both, as described below) indicate that the trajectory is
significantly different from the native state reference simulations.
798
Calculation of Q_{mode} and its use as an order parameter for protein folding. The
fraction of native contacts, Q, was calculated for each domain and interface of all 122
proteins during their synthesis, ejection, and post-translational dynamics. Only
contacts between pairs of residues both within secondary structural elements as
identified by STRIDE[66] based on the final rebuilt all-atom structures were considered.
To determine when a given domain or interface within a protein folded, the mode of
the Q values over a 15-ns sliding window (Q_{mode}) was compared to the representative
value of the native state computed as the average Q_{mode} over all windows of the ten
native-state simulations denoted ($\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle$). A given trajectory is defined as misfolded
if its average Q_{mode} over the final 100 ns of the post-translational dynamics portion of
the simulation, denoted $\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle$, is less than $\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle - 3\sigma$, where σ is the standard
deviation of $\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle$.
811
Generation of entanglement metric distributions and use as an order parameter
for protein folding. To detect non-covalent lasso entanglements we use linking
numbers[67]. A link is defined as the entanglement of two closed curves; here, we use
(1) the closed curve composed of the backbone trace connecting residues i and j that
form a native contact and (2) the open curves formed by the terminal tails. The native
contact between i and j in (1) is considered to close this loop, even though there is no
covalent bond between these two residues. Outside this loop is an N-terminal
segment, composed of residues 1 through $i - 1$, and a C-terminal segment, composed
of residues $j + 1$ through N, whose entanglement through the closed loop we
characterize with partial linking numbers denoted g_{N} and g_{C}[47]. For a given structure
of an *N*-length protein, with a native contact present at residues (i, j), the coordinates
R_l and the gradient dR_l of the point l on the curves were calculated as

$$\begin{cases} R_l = \dfrac{1}{2}(r_l + r_{l+1}) \\ dR_l = r_{l+1} - r_l \end{cases}, \qquad [3]$$

825 where r_l is the coordinates of the Cα atom in residue l. The linking numbers $g_{\mathrm{N}}(i,j)$
826 and $g_{\mathrm{C}}(i,j)$ were calculated as

$$827 \quad \begin{cases} g_{\text{N}}(i,j) = \dfrac{1}{4\pi} \sum\limits_{m=6}^{i-5} \sum\limits_{n=i}^{j-1} \dfrac{\boldsymbol{R}_m - \boldsymbol{R}_n}{|\boldsymbol{R}_m - \boldsymbol{R}_n|^3} \cdot (\text{d}\boldsymbol{R}_m \times \text{d}\boldsymbol{R}_n) \\[2em] g_{\text{C}}(i,j) = \dfrac{1}{4\pi} \sum\limits_{m=i}^{j-1} \sum\limits_{n=j+4}^{N-6} \dfrac{\boldsymbol{R}_m - \boldsymbol{R}_n}{|\boldsymbol{R}_m - \boldsymbol{R}_n|^3} \cdot (\text{d}\boldsymbol{R}_m \times \text{d}\boldsymbol{R}_n) \end{cases}, \qquad [4]$$

where we excluded the first 5 residues on the N-terminal curve, last 5 residues on the
C-terminal curve and 4 residues before and after the native contact for the purpose of
eliminating the error introduced by both the high flexibility and contiguity of the termini
and trivial entanglements in local structure. The above summations yield two non-
integer values, and the total linking number for a native contact (i,j) was therefore
estimated as

$$834 \qquad g(i,j) = \text{round}\big(g_{\text{N}}(i,j)\big) + \text{round}\big(g_{\text{C}}(i,j)\big), \qquad\qquad [5]$$

Comparing the absolute value of the total linking number for a native contact (i,j) to
that of a reference state allows us to ascertain a gain or loss of linking between the
backbone trace loop and the terminal open curves as well as any switches in chirality.
Therefore, there are 6 change in linking cases we should consider (Table S3) when
using this approach to quantify entanglement.
To examine the distribution of change in linking (entanglement) detected for a
given protein model and statistically independent post-translational trajectory we can
generate a discrete probability distribution of the 6 cases in Table S3 as

$$843 \qquad P(G_k \mid PDB, traj) = \frac{N_k}{N_E} \qquad\qquad [6]$$

Where G_k for $k \in \{0,1,2,3,4,5\}$ is the change in entanglement case of interest from
Table S3, N_E is the total number of changes in entanglement instances detected in the
trajectory, and N_k is the total number of changes in entanglement in the trajectory of
type k. As the change in entanglement is held relative to the static crystal structure, it
is necessary to correct the post-translational distribution to remove transient changes
in entanglement present in the reference state dynamics. This was done by subtraction
of the reference distribution from the post-translational distribution considering
nontrivial changes in linkage:

$$852 \quad f_c(G_k \mid PDB, traj) = \big| P_{pt}(G_k \mid PDB, traj) - P_{ref}(G_k \mid PDB) \big|, \quad k \in \{0,1,2,3,4\}. \qquad [7]$$

For a given case of entanglement change k the magnitude of $f_c(G_k \mid PDB, traj)$
increases as the probability of that mode of entanglement change deviates from the
reference simulations. We define a given trajectory as misfolded if the average of any
of its corrected entanglement values from Eq. 7 over the final 100 ns of the trajectory
are ≥0.1.
This trajectory-level analysis is useful for classifying statistically independent
sample sets by the level and types of changes in entanglement they exhibit, but a time
series metric which conveys the same information was desired to allow for folding time
extrapolations. G is a time dependent order parameter that reflects the extent of the
topological entanglement changes in a given structure compared to the native
structure and is calculated as

$$864 \qquad G(t) = \frac{1}{N} \sum_{(i,j)} \Theta\left((i,j) \in nc \cap g(i,j,t) \neq g^{\text{native}}(i,j) \right), \qquad\qquad [8]$$

where (i,j) is one of the native contacts in the native crystal structure; nc is the set of
native contacts formed in the current structure at time t; $g(i,j,t)$ and $g^{\text{native}}(i,j)$ are,
respectively, the total entanglement number of the native contact (i,j) at time t, and
native structures estimated using Eq. 5; N is the total number of native contacts within

the native structure and the selection function θ equals 1 when the condition is true
and equals 0 when it is false. The larger G is, the greater the number of native contact
residues that have changed their entanglement status relative to the native state. The
utility of entanglement for detecting structural perturbations not apparent by fraction of
native contacts or root mean square deviation is visually described in Figure S6.
874
Calculation and extrapolation of folding times. Folding times were determined for
each domain and interface from their post-translational Q_{mode} and G time series. The
folding time for a domain or interface is taken as the first t at which Q_{mode} is greater
than or equal to $\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle - 3\sigma$ and with $G \leq G_{\mathrm{xs}}$. The survival probability of the unfolded
state was then computed based on these folding times and fit to the double-
exponential function $S_U(t) = f_1 \exp(k_1 * t) + f_2 \exp(k_2 * t)$ with $f_1 + f_2 \equiv 1$. This
double-exponential fit equation represents a kinetic scheme in which the unfolded and
misfolded states each proceed irreversibly to the folded state by parallel folding
pathways and there is no inter-transitions between unfolded and misfolded states.
Folding times for each kinetic phase were computed as $\tau_{\mathrm{F},1} = \frac{1}{k_1}$ and $\tau_{\mathrm{F},2} = \frac{1}{k_2}$, with the
overall folding time of the domain or interface taken as the longer of the two folding
times. The folding time reported for each protein is the longest folding time from any
of its constituent domains or interfaces. The assumption of double-exponential folding
kinetics is not a good assumption for all 122 proteins simulated. We therefore only
consider domains and interfaces whose $S_U(t)$ time series are fit with a Pearson $R^2 >$
0.90. We are also unable to compute folding times for the nine proteins for which no
trajectories folded. In total, we found reliable folding times for 73 of our 122 proteins.
Simulated folding times were extrapolated to experimental timescales using the
equation $\tau_{\mathrm{exp}} = \tau_{\mathrm{sim}} * \alpha$, where τ_{sim} is a given protein's simulated folding time and $\alpha =$
3,967,486 is the mean acceleration of folding in our coarse-grain simulations relative
to real timescales[28].
896
897 **Identifying misfolded proteins unlikely to interact with trigger factor.** To
898 determine whether or not a given protein in our data set is likely to interact with trigger
899 factor (TF) during synthesis we computed the relative difference in the hydrophobic
900 SASA between misfolded trajectories and folded trajectories on the ribosome using
901 the equation

$$\zeta_{\mathrm{hydrophobic}}^{\mathrm{co-t}}(t,l) = \left(\frac{A_{\mathrm{hydrophobic}}(t,l)}{\langle A_{\mathrm{hydrophobic}}(t,l) \rangle_{\mathrm{F}}} - 1 \right) * 100\%. \tag{9}$$

In Eq. 9, $A_{\mathrm{hydrophobic}}(t,l)$ is the total hydrophobic SASA of residues in the nascent
protein at time t and nascent chain length l exposed outside of the ribosome exit tunnel
(defined as having an x-coordinate ≥ 100 Å in the internal CHARMM coordinate system;
see Ref. 41 for details). The term $\langle A_{\mathrm{hydrophobic}}(t,l) \rangle_{\mathrm{F}}$ is the mean total hydrophobic
SASA of residues at time t and length l outside of the exit tunnel calculated over all
frames of synthesis trajectories identified to be folded by Q_{mode} and G analysis. Note
well, the reference states for Equations 10-12 are the native-state reference
simulations initiated in bulk solution, but due to the co-translational nature of TF
interactions we use the folded subpopulation of synthesis trajectories as the reference
state in Eq. 9. As TF is thought to only interact with nascent proteins of 100 residues
or longer[68], we compute Eq. 9 for $l = \{100, 101, \dots, N\}$ for each protein and trajectory,
where N is the total number of residues in the full-length protein. Proteins shorter than
100 amino acids are considered not to interact with TF. To quantify the overall
propensity of a misfolded trajectory of a given protein to interact with TF we compute

$\langle\zeta_{\text{hydrophobic}}^{\text{co-t}}\rangle$ as the mean of $\zeta_{\text{hydrophobic}}^{\text{co-t}}(t, l)$ for all t and allowed values of l. Based
on examination of $\zeta_{\text{hydrophobic}}^{\text{co-t}}(t, l)$ time series, we determined that a value of
$\langle\zeta_{\text{hydrophobic}}^{\text{co-t}}\rangle \leq 10\%$ corresponds to proteins that are unlikely to engage TF
significantly more than folded conformations on the ribosome. For the purposes of this
calculation and all others in this work that consider sets of hydrophobic residues,
coarse-grain interactions sites representing {Ile, Val, Leu, Phe, Cys, Met, Ala, Gly, Trp}
are considered to be hydrophobic.
924
Identifying misfolded proteins unlikely to interact with GroEL/GroES. Our data
set of 122 *E. coli* proteins was first cross-referenced with the list of 276 confirmed
GroEL/GroES substrates[44–46]. The 103 proteins that do not appear in this list of
confirmed clients are considered to not be GroEL/GroES client proteins. GroEL/GroES
is thought to identify and bind regions of exposed hydrophobic surface area on nascent
proteins. To determine whether the misfolded conformations of proteins that can
interact with GroEL/GroES are likely to do so, we compared the total hydrophobic
SASA of misfolded conformations with the native-state ensemble using the equation

933
$$\zeta_{\text{hydrophobic}}(t) = \left(\frac{A_{\text{hydrophobic}}(t)}{\langle A_{\text{hydrophobic}}(t)\rangle_{\text{NS}}} - 1\right) * 100\%. \qquad [10]$$

This equation provides the quantity $\zeta_{\text{hydrophobic}}(t)$, which measures the relative
difference between the SASA of hydrophobic residues within a misfolded conformation
at time t $(A_{\text{hydrophobic}}(t))$ in comparison to to the mean hydrophobic SASA calculated
over all frames of the native-state reference simulations $(\langle A_{\text{hydrophobic}}(t)\rangle_{\text{NS}})$. As
GroEL/GroES interacts with proteins post-translationally, Eq. 10 was applied to the
post-translational simulation data for all misfolded conformations of a given protein
and the average over the final 100 ns computed for each trajectory $(\langle\zeta_{\text{hydrophobic}}\rangle)$. The
full set of proteins with misfolded conformations that we predict will not interact with
GroEL/GroES is taken as the union of the sets of all trajectories for proteins not in the
list of experimentally confirmed clients and the list of trajectories with $\langle\zeta_{\text{hydrophobic}}\rangle \leq$
10%.
945
Identifying misfolded proteins unlikely to interact with DnaK. We first cross-
referenced our list of 122 simulated proteins with a list of DnaK client proteins. The 34
proteins that do not appear on this list are considered to not interact with DnaK. To
determine whether misfolded conformations of proteins that can interact with DnaK
are likely to do so, we predicted DnaK binding sites using the Limbo webserver[35] and
then compared the total SASA of residues in predicted binding sites between
misfolded conformations and the native state using the equation

953
$$\zeta_{\text{DnaK}}(t) = \left(\frac{A_{\text{DnaK}}(t)}{\langle A_{\text{DnaK}}(t)\rangle_{\text{NS}}} - 1\right) * 100\%. \qquad [11]$$

In this equation $A_{\text{DnaK}}(t)$ is the total SASA of residues in DnaK binding sites as a
function of time t and $\langle A_{\text{DnaK}}(t)\rangle_{\text{NS}}$ is the mean total SASA of residues in DnaK binding
sites averaged over all frames of the native-state reference simulations. Equation 11
was applied to the post-translational simulation data of all misfolded conformations for
each misfolded trajectory and the average over the final 100 ns of each trajectory
computed $(\langle\zeta_{\text{DnaK}}\rangle)$. The full set of trajectories with misfolded conformations that we
predict will not interact with DnaK is taken as the union of the sets of trajectories for
proteins not in the list of experimentally confirmed clients and the list of trajectories
with $\langle\zeta_{\text{DnaK}}\rangle \leq 10\%$. We note that three proteins, PDB IDs 2JRX, 2V81, and 2KFW,
are predicted by Limbo to have no DnaK binding sites. 2JRX and 2KFW appear on the

list of confirmed DnaK clients, and are therefore considered to be DnaK binders. 2V81,
however, is not a confirmed DnaK client and is thus counted as not likely to interact
with DnaK.

Identifying misfolded proteins unlikely to aggregate. We used the AMYLPRED2
webserver[37] to predict the sets of aggregation-prone residues within the primary
sequences of our 122 proteins. Whether or not a given trajectory for a protein is likely
to aggregate was determined by comparing the SASA of aggregation-prone regions
within the misfolded trajectory to the mean SASA of aggregation-protein regions in the
native-state reference ensemble using the equation

974
$$\zeta_{\mathrm{agg}}(t) = \left(\frac{A_{\mathrm{agg}}(t)}{\langle A_{\mathrm{agg}}(t) \rangle_{\mathrm{NS}}} - 1 \right) * 100\%. \qquad [12]$$

In this equation, $A_{\mathrm{agg}}(t)$ is the total SASA of residues in aggregation-prone regions as
a function of time, t, and $\langle A_{\mathrm{agg}}(t) \rangle_{\mathrm{NS}}$ is the mean total SASA of residues in aggregation-
prone regions averaged over all frames of the native-state reference simulations.
Equation 12 was applied to the post-translational simulation data for all misfolded
trajectories of a given protein and the average over the final 100 ns for each trajectory
($\langle \zeta_{\mathrm{agg}} \rangle$). Trajectories with $\langle \zeta_{\mathrm{agg}} \rangle \leq 10\%$ are considered to be unlikely to aggregate.
Note that our methods do not account for the presence of forms of aggregates other
than amyloid, as our calculation is based on the AMYLPRED2 algorithm.

Identifying misfolded proteins unlikely to be degraded. Whether or not a protein's
misfolded conformations are likely to be targeted for degradation was determined on
the basis of $\zeta_{\mathrm{hydrophobic}}$ (Eq. 10). Proteins with $\langle \zeta_{\mathrm{hydrophobic}} \rangle \leq 10\%$ are considered to
be unlikely to be degraded.

Creating a database of functional residues for *E. coli* proteins. Information from
UniProt and RCSB was unified and parsed to create a database of residues implicated
in function for each of our 122 proteins. Residues involved in interactions with small
molecules were identified as those residues with heavy atoms within 4.5 Å of any
heteroatom (identified by the HETATM keyword in PDB records) other than water and
non-native amino acids such as selenidomethionine (*i.e.*, MSE residues). Many
proteins must form multimeric complexes in order to exercise their function. To
consider these interactions, we also identified residues with heavy atoms within 4.5 Å
of heavy atoms in a different chain ID within the same PDB structure.
Our 122 coarse-grain models were built from single PDB structures or, in the
case of some multi-domain protein models, the merging of multiple structures. These
structures used for model building often lack ligands or protein binding partners that
may be required for function due to differences in crystallographic conditions and/or
the intention of the original crystallographers. To provide a broader view of functional
residues we therefore also considered all PDB structures identified by UniProt to
represent the same gene product. Functional residues were identified in these
alternative structures as described above for the initial structures. PDB entries
representing the same protein often have different residue numbering schemes and a
small number of mutations. We therefore used amino acid alignments in BLAST to
determine the mapping from alternative numbering schemes to the numbering scheme
within the structure used for model building. Only those domains with at least 97%
sequence identity were considered for this analysis to allow for small mutational
changes while excluding significantly different proteins. A summary of each of the
terms in the database and their meanings is provided in Table S10.

Determining which misfolded conformations are likely non-functional. The relative difference in function between misfolded trajectories and native-state reference trajectories was determined by calculating the relative difference of the structural overlap of residues identified to be involved in protein function using the equation

$$\chi_{\text{func}} = \left(1 - \frac{\chi(t)}{\langle\chi(t)\rangle_{\text{NS}}}\right) * 100\%. \qquad [13]$$

In this equation, $\chi(t)$ is the structural overlap between residues implicated in function at time t in a misfolded trajectory with the native-state reference structure and $\langle\chi(t)\rangle_{\text{NS}}$ is the mean of this same value computed over all simulation frames of the native-state reference simulations. The value of $\chi(t)$ is calculated as

$$\chi(t) = \frac{1}{N}\sum_{i \in \{\text{Func}\}} \sum_{j \geq i+2; \, j \in \{\text{Func}\}} \theta\left[\epsilon - \left|r_{ij}(t) - r_{ij}^0\right|\right] \qquad [14]$$

and gives the fraction of pairwise distances that are at native-like values at time t. The set $\{\text{Func}\}$ contains all residues implicated in protein function. The indices i and j correspond to residues in $\{\text{Func}\}$ for the protein being analyzed. N is the total number of pairwise contacts between residues i and j both in $\{\text{Func}\}$ that also satisfy the condition $j \geq i+2$. The parameters $r_{ij}(t)$ and r_{ij}^0 are the distances between residues i and j at time t and between i and j in the native state reference structure, respectively. $\theta(x)$ is the step function given by

$$\theta(x) = \begin{cases} 1, & x \geq 0 \\ 0, & x < 0 \end{cases}. \qquad [15]$$

The value of ϵ is taken as $0.2 \cdot r_{c_\alpha}$ where $r_{c_\alpha} = 3.81$ Å is the virtual bond length between coarse-grain interaction sites in the coarse-grain simulation model. A particular pair of residues i and j contribute 1 to $\chi(t)$ if $\left|r_{ij}(t) - r_{ij}^0\right|$ is less than ϵ, such that $\epsilon - \left|r_{ij}(t) - r_{ij}^0\right| \geq 0$, and in all other situations contribute 0 to $\chi(t)$. Similar forms of this equation have been used previously to observe structural transitions in simulations of proteins[69]. The value of χ_{func} was averaged over the final 100 ns for given protein trajectory to give $\langle\chi_{\text{func}}\rangle$, the mean relative difference in structure of functional residues for the misfolded conformation in comparison to the native state. Trajectories with $\langle\chi_{\text{func}}\rangle \geq 10\%$ are considered to be less functional than the native state.

Selection of 10% threshold for classifying misfolded trajectories as native-like. We selected a threshold of 10% for $\zeta_{\text{hydrophobic}}(t)$, $\zeta_{\text{DnaK}}(t)$, $\zeta_{\text{agg}}(t)$, and χ_{func} (see Eqs. 10, 11, 12, and 13) based on an analysis of how frequently our native state reference trajectories for each protein explore conformations with 10% or greater difference from their native state mean (Table S9). Virtually all proteins explore conformations with ≥10% for each metric in their native state, indicating that 10% is a parsimonious threshold for determining if misfolded conformations are native like. In the case of $\zeta_{\text{hydrophobic}}^{\text{co}-\text{t}}(t,l)$, similar calculations as those performed for the other metrics would require running prohibitively expensive arrested-ribosome-nascent chain complex simulations for each of our 122 proteins. We therefore use a threshold of 10% for $\zeta_{\text{hydrophobic}}^{\text{co}-\text{t}}(t,l)$ for the sake of consistency with our other thresholds.

Protease mass spectrometry experiments. See the Supplementary Methods for a full description. Briefly, *E. coli* K12 cells (NEB) were grown in 2 sets of 3 × 50 mL (biological triplicates) MOPS EZ rich media from saturated overnight cultures with a

starting OD_{600} of 0.05. As described in Ref. 71, one set was supplemented with 0.5 mM [$^{13}C_6$]L-Arginine and 0.4 mM [$^{13}C_6$]L-Lysine and the other with 0.5 mM L-Arginine and 0.4 mM L-Lysine. Cells were cultured at 37°C with agitation (220 rpm) to a final OD_{600} of 0.8. Each heavy/light pair was pooled together; cells were collected by centrifugation at 4000 g for 15 mins at 4°C, supernatants were removed, and cell pellets were stored at -20°C until further use.

Frozen cell pellets were resuspended in a lysis buffer consisting of 900 µL of Tris pH 8.2 (20 mM Tris pH 8.2, 100 mM NaCl, 2 mM $MgCl_2$ and supplemented with DNase I to a final concentration (f.c.) of 0.1 mg mL^{-1}). Resuspended cells were cryogenically pulverized with a freezer mill (SPEX Sample Prep). Lysates were then clarified at 16000 g for 15 min at 4 °C to remove insoluble cell debris. To deplete ribosome particles, clarified lysates were ultracentrifuged at 33,300 rpm at 4 °C for 90 min using a SW55 Ti rotor. Protein concentrations of clarified lysates were determined using the bicinchoninic acid assay (Rapid Gold BCA Assay, Pierce) and diluted to 3.3 mg mL^{-1} using lysis buffer.

To prepare native samples, 3.5 µL of normalized lysates were diluted with 96.5 µL of Tris native dilution buffer (20 mM Tris pH 8.2, 100 mM NaCl, 10.288 mM $MgCl_2$, 10.36 mM KCl, 2.07 mM ATP, 1.04 mM DTT, 62 mM GdmCl) to a final protein concentration of 0.115 mg mL^{-1}. Native samples were then equilibrated by incubating for 90 min at room temperature. To prepare unfolded samples, 600 µL of normalized lysates, 100 mg of solid GdmCl, and 2.4 µL of a freshly prepared 700 mM DTT stock solution were combined, and solvent was removed using a vacufuge plus to a final volume of 170 µL. Unfolded lysates were incubated overnight at room temperature. To refold, 99 µL of refolding dilution buffer (19.5 mM Tris pH 8.2, 97.5 mM NaCl, 10.03 mM $MgCl_2$, 10.1 mM KCl, 2.02 mM ATP and .909 mM DTT) were rapidly added to 1 µL of unfolded extract. Refolded samples were then incubated at room temperature for 1 min, 5 min or 2 h.

100 µL of the native or refolded lysates was added to Proteinase K (enzyme:substrate ratio of 1:100 w/w ratio[70]), incubated for 1 min at room temperature, and quenched by boiling in a mineral oil bath at 110°C for 5. Boiled samples were transferred to tubes containing 76 mg urea. To prepare samples for mass spectrometry, dithiothreitol was added to a final concentration of 10 mM and samples were incubated at 37°C for 30 minutes. Iodoacetamide was added to a final concentration of 40 mM and samples were incubated at room temperature in the dark for 45 minutes. LysC was added to a 1:100 enzyme:substrate (w/w) ratio and samples were incubated at 37°C for 2 h, urea was diluted to 2 M using 100 mM ammonium bicarbonate pH 8, then trypsin was added to a 1:50 enzyme:substrate (w/w) ratio and incubated overnight at 25°C.

Peptides were acidified, desalted with Sep-Pak C18 1 cc Vac Cartridges, dried down, and resuspend in 0.1% formic acid, as previously described[71]. LC-MS/MS acquisition was conducted on a Thermo Ultimate 3000 UHPLC system with an Acclaim Pepmap RSLC C18 column (75 µm × 25 cm, 2 µm, 100 Å) in line with a Thermo Q-Exctive HF-X Orbitrap, identically as previously described[71].

Proteome Discoverer (PD) Software Suite (v2.4, Thermo Fisher) and the Minora Algorithm were used to analyze mass spectra and perform Label Free Quantification (LFQ) of detected peptides. Default settings for all analysis nodes were used except where specified. The data were searched against Escherichia coli (UP000000625, Uniprot) reference proteome database. For peptide identification, the PD MSFragger node was used, using a semi-tryptic search allowing up to 2 missed

cleavages[72]. A precursor mass tolerance of 10 ppm was used for the MS1 level, and
a fragment ion tolerance was set to 0.02 Da at the MS2 level. Additionally, a maximum
charge state for theoretical fragments was set at 2. Oxidation of methionine and
acetylation of the N-terminus were allowed as dynamic modifications while
carbamidomethylation on cysteines was set as a static modification. Heavy isotope
labeling ($^{13}C_6$) of Arginine and Lysine were allowed as dynamic modifications. The
Philosopher PD node was used for FDR validation. Raw normalized extracted ion
intensity data for the identified peptides were exported from the .pdResult file using a
three-level hierarchy (protein > peptide group > consensus feature). These data were
further processed utilizing custom Python analyzer scripts (available on GitHub, and
described in depth previously[49,71]).

1120
Clustering long-lived misfolded states of glycerol-3-phosphate dehydrogenase.
The structural distribution from the last 100 ns of the post-translational simulations of
glycerol-3-phosphate dehydrogenase was assessed as the pseudo free energy
$-ln(P)$, where P is the probability density, along the order parameters G and $Q_{overall}$.
To further analyze the post-translational structures, 400 clusters (micro-states) were
grouped from the last 100 ns trajectories using the k- means algorithm[73,74]. A Markov
state model (MSM) was built and the clusters were coarse-grained into a small number
of metastable states using the PCCA+ algorithm[75]. The number of metastable states
was chosen based on the existence of a gap in the eigenvalue spectrum of the
transition probability matrix[76]. Five representative structures of each metastable state
were randomly sampled from all microstates according to the probability distribution
of the microstates within the given metastable state. All the clustering and MSM
building were performed by using PyEmma package[77].

1134
1135 **Determining which coarse-grain structures have increased exposure of**
1136 **peptides**. The relative change in solvent accessible surface area of experimentally
1137 identified peptides for glycerol-3-phosphate dehydrogenase was calculated as

1138
1139
$$\zeta_{peptide}(t) = \left(\frac{A_{peptide}(t)}{\langle A_{peptide}(t)\rangle_{NS}} - 1\right) * 100\%. \qquad [16]$$

1140
In this equation, $A_{peptide}(t)$ is the total SASA of the residues within the peptide under
consideration at time t and $\langle A_{peptide}(t)\rangle_{NS}$ is the mean value of $A_{peptide}(t)$ computed
over all frames of the native-state reference simulations. Eq. 16 was applied to all
frames in the final 100 ns of each glycerol-3-phosphate dehydrogenase trajectory and
then averaged separately for each of its eight misfolded metastable states. Values of
$\langle \zeta_{[333-354]} \rangle$, $\langle \zeta_{[F351]} \rangle$, $\langle \zeta_{[L293]} \rangle$ are summarized in Table S11 for metastable states {S1,
S2, …, S8}.

1148
1149 **DATA AVAILABILITY**

1150
Raw data for Figures 1, 2, 3, 4, 5, S1, S2, and S3 are available in the included software
directory. We cannot feasibly provide all ~30 TB of trajectory data, but we do provide
sample trajectory files and use them to demonstrate our analysis methods.
Supplementary Data File 1 contains the annotated experimental data for glycerol-3-
phosphate dehydrogenase. Supplementary Data Files 2, 3, and 4 contain all peptide-
level data from the 1-, 5-, and 120-min time points, respectively. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD031425.

CODE AVAILABILITY

ACKNOWLEDGMENTS

SDF acknowledges support from the NIH Director's New Innovator Award (DP2GM140926) and from the National Science Foundation Division of Molecular and Cellular Biology (MCB-2045844). EPO acknowledges funding from the National Institutes of Health (R35-GM124818) and the National Science Foundation (MCB-1553291).

REFERENCES

(1)   Komar, A. A.; Lesnik, T.; Reiss, C. *FEBS Lett.* **1999**, *462* (3), 387–391.

(2)   Zhou, M.; Guo, J.; Cha, J.; Chae, M.; Chen, S.; Barral, J. M.; Sachs, M. S.; Liu, Y. *Nature* **2013**, *494* (7439), 111–115.

(3)   Fu, J.; Murphy, K. A.; Zhou, M.; Li, Y. H.; Lam, V. H.; Tabuloc, C. A.; Chiu, J. C.; Liu, Y. *Genes Dev.* **2016**, *30* (15), 1761–1775.

(4)   Jayaraj, G. G.; Hipp, M. S.; Ulrich Hartl, F. *Cold Spring Harb. Perspect. Biol.* **2020**, *12* (1).

(5)   Kuhlman, B.; Bradley, P. *Nat. Rev. Mol. Cell Biol.* **2019**, *20* (11), 681–697.

(6)   Hipp, M. S.; Kasturi, P.; Hartl, F. U. *Nat. Rev. Mol. Cell Biol.* **2019**, *20* (7), 421–435.

(7)   Sharma, A. K.; O'Brien, E. P. *Curr. Opin. Struct. Biol.* **2018**, *49*, 94–103.

(8)   Balch, W. E.; Morimoto, R. I.; Dillin, A.; Kelly, J. W. *Science.* **2008**, *319* (5865), 916–919.

(9)   Hayer-Hartl, M.; Bracher, A.; Hartl, F. U. *Trends Biochem. Sci.* **2016**, *41* (1), 62–76.

(10)   Mayer, M. P.; Bukau, B. *Cell. Mol. Life Sci.* **2005**, *62* (6), 670–684.

(11)   Calloni, G.; Chen, T.; Schermann, S. M.; Chang, H. C.; Genevaux, P.; Agostini, F.; Tartaglia, G. G.; Hayer-Hartl, M.; Hartl, F. U. *Cell Rep.* **2012**, *1* (3), 251–264.

(12)   Gong, B.; Radulovic, M.; Figueiredo-Pereira, M. E.; Cardozo, C. *Front. Mol. Neurosci.* **2016**, *9* (JAN), 1–16.

(13)   LaBreck, C. J.; May, S.; Viola, M. G.; Conti, J.; Camberg, J. L. *Front. Mol. Biosci.* **2017**, *4* (MAY), 1–13.

(14)   Rae, B. D.; Long, B. M.; Badger, M. R.; Price, G. D. *Microbiol. Mol. Biol. Rev.* **2013**, *77* (3), 357–379.

(15)   Haass, C.; Selkoe, D. J. *Nat. Rev. Mol. Cell Biol.* **2007**, *8* (2), 101–112.

(16)   Alberti, S.; Hyman, A. A. *Nat. Rev. Mol. Cell Biol.* **2021**, *22* (March).

(17)   Kramer, G.; Shiber, A.; Bukau, B. *Annu. Rev. Biochem.* **2019**, *88*, 337–364.

(18)   Conibear, A. C. *Nat. Rev. Chem.* **2020**, *4* (December).

(19) Fluitt, A.; Pienaar, E.; Viljoen, H. *Comput. Biol. Chem.* **2007**, *31* (5–6), 335–346.

(20) Qian, S. B.; Princiotta, M. F.; Bennink, J. R.; Yewdell, J. W. *J. Biol. Chem.* **2006**, *281* (1), 392–400.

(21) Schubert, U.; Antón, L. C.; Gibbs, J.; Norbury, C. C.; Yewdell, J. W.; Bennink, J. R. *Nature* **2000**, *404* (6779), 770–774.

(22) Wu, K.; Stull, F.; Lee, C.; Bardwell, J. C. A. *Nat. Commun.* **2019**, *10* (1).

(23) Powers, E. T.; Powers, D. L.; Gierasch, L. M. *Cell Rep.* **2012**, *1* (3), 265–276.

(24) Santra, M.; Dill, K. A.; De Graff, A. M. R. *Proc. Natl. Acad. Sci. U. S. A.* **2019**, *116* (44), 22173–22178.

(25) Spencer, P. S.; Siller, E.; Anderson, J. F.; Barral, J. M. *J. Mol. Biol.* **2012**, *422* (3), 328–335.

(26) Zhang, G.; Hubalewska, M.; Ignatova, Z. *Nat. Struct. Mol. Biol.* **2009**, *16* (3), 274–280.

(27) Kimchi-Sarfaty, C.; Oh, J. M.; Kim, I.-W.; Sauna, Z.; Calcagno, A. M.; Ambudkar, S.; Gottesman, M. *Science (80-.).* **2007**, *315* (November), 525–529.

(28) Leininger, S. E.; Trovato, F.; Nissley, D. A.; O'Brien, E. P. *Proc. Natl. Acad. Sci. U.S.A.* **2019**, *116* (12), 5523–5532.

(29) Nissley, D. A.; O'Brien, E. P. *J. Phys. Chem. B* **2018**, *122* (43), 9927–9937.

(30) Nilsson, O. B.; Hedman, R.; Marino, J.; Wickles, S.; Bischoff, L.; Johansson, M.; Müller-Lucks, A.; Trovato, F.; Puglisi, J. D.; O'Brien, E. P.; Beckmann, R.; von Heijne, G. *Cell Rep.* **2015**, *12* (10), 1533–1540.

(31) Tian, P.; Steward, A.; Kudva, R.; Su, T.; Shilling, P. J.; Nickson, A. A.; Hollins, J. J.; Beckmann, R.; Von Heijne, G.; Clarke, J.; Best, R. B. *Proc. Natl. Acad. Sci. U. S. A.* **2018**, *115* (48), E11284–E11293.

(32) Jiang, Y.; Neti, S. S.; Pradhan, P.; Booker, S. J.; O'Brien, E. P. O. *Under Review. bioRxiv* **2021**. DOI: https://doi.org/10.1101/2021.08.18.456802.

(33) Frydman, J.; Erdjument-Bromage, H.; Tempst, P.; Ulrich Hartl, F. *Nat. Struct. Biol.* **1999**, *6* (7), 697–705.

(34) Saio, T.; Guan, X.; Rossi, P.; Economou Anastassios , Kalodimos, C. G. *Science.* **2014**, *344*.

(35) Van Durme, J.; Maurer-Stroh, S.; Gallardo, R.; Wilkinson, H.; Rousseau, F.; Schymkowitz, J. *PLoS Comput. Biol.* **2009**, *5* (8).

(36) Tapia-rojo, R.; Eckels, E. C.; Valle-orero, J.; Fernandez, J. M. *Nat. Commun.* **2017**, 1–7.

(37) Tsolis, A. C.; Papandreou, N. C.; Iconomidou, V. A.; Hamodrakas, S. J. *PLoS One* **2013**, *8* (1), 1–6.

(38) Yedidi, R. S.; Wendler, P.; Enenkel, C. *Front. Mol. Biosci.* **2017**, *4* (JUN), 1–14.

(39) Goldberg, A. L.; Dice, J. F. *Annu. Rev. Biochem.* **1974**, 835–864.

(40) Consortium, T. U. *Nucleic Acids Res.* **2017**, *45*, 158–169.

(41) Nissley, D. A.; Vu, Q. V.; Trovato, F.; Ahmed, N.; Jiang, Y.; Li, M. S.; O'Brien, E. P. *J. Am. Chem. Soc.* **2020**, jacs.9b12264.

(42) Klimov, D. K.; Thirumalai, D. *Phys. Rev. Lett.* **1997**, *79* (2), 317–320.

(43) Kmiecik, S.; Gront, D.; Kolinski, M.; Wieteska, L.; Dawid, A. E.; Kolinski, A. *Chem. Rev.* **2016**, *116* (14), 7898–7936.

(44) Niwa, T.; Fujiwara, K.; Taguchi, H. *FEBS Lett.* **2016**, *590* (2), 251–257.

(45) Fujiwara, K.; Ishihama, Y.; Nakahigashi, K.; Soga, T.; Taguchi, H. *EMBO J.* **2010**, *29* (9), 1552–1564.

(46) Kerner, M. J.; Naylor, D. J.; Ishihama, Y.; Maier, T.; Chang, H. C.; Stines, A.

P.; Georgopoulos, C.; Frishman, D.; Hayer-Hartl, M.; Mann, M.; Hartl, F. U. *Cell* **2005**, *122* (2), 209–220.

(47) Baiesi, M.; Orlandini, E.; Seno, F.; Trovato, A. *J. Phys. A Math. Theor.* **2017**.

(48) Rolfson, D. *Knots and Links*; 1976.

(49) To, P.; Whitehead, B.; Tarbox, H. E.; Fried, S. D. *J. Am. Chem. Soc.* **2021**, *143* (30), 11435–11448.

(50) Sułkowska, J. I.; Sułkowski, P.; Onuchic, J. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106* (9), 3119–3124.

(51) Bitran, A.; Jacobs, W. M.; Zhai, X.; Shakhnovich, E. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117* (3), 1485–1495.

(52) Lafita, A.; Tian, P.; Best, R. B.; Bateman, A. *Curr. Opin. Struct. Biol.* **2019**, *58*, 97–104.

(53) Norcross, T. S.; Yeates, T. O. *J. Mol. Biol.* **2006**, *362* (3), 605–621.

(54) Taylor, W. R.; Lin, K. *Nature* **2003**, *421* (6918), 25.

(55) Dabrowski-Tumanski, P.; Piejko, M.; Niewieczerzal, S.; Stasiak, A.; Sulkowska, J. I. *J. Phys. Chem. B* **2018**, *122* (49), 11616–11625.

(56) Gosavi, S.; Chavez, L. L.; Jennings, P. A.; Onuchic, J. N. *J. Mol. Biol.* **2006**, *357* (3), 986–996.

(57) Grosberg, A.; Nechaev, S. **1993**, *106*, 1–29.

(58) Baiesi, M.; Orlandini, E.; Seno, F.; Trovato, A. *Sci. Rep.* **2019**, *9* (1), 1–12.

(59) Belle, A.; Tanay, A.; Bitincka, L.; Shamir, R.; Shea, E. K. O. *Proc. Natl. Acad. Sci. USA* **2006**, *103* (35), 13004–13009.

(60) Cambridge, S. B.; Gnad, F.; Nguyen, C.; Bermejo, J. L.; Kruger, M.; Mann, M.; Atthias. *J. Proteome Res.* **2011**, *10*, 5275–5284.

(61) Buhr, F.; Jha, S.; Thommen, M.; Mittelstaet, J.; Kutz, F.; Schwalbe, H.; Rodnina, M. V.; Komar, A. A. *Mol. Cell* **2016**, *61* (3), 341–351.

(62) Neidhardt, F.; Ingraham, J.; Schaechter, M. *Physiology of the Bacterial Cell. A Molecular Approach*; 1990.

(63) Siller, E.; DeZwaan, D. C.; Anderson, J. F.; Freeman, B. C.; Barral, J. M. *J. Mol. Biol.* **2010**, *396* (5), 1310–1318.

(64) Ciryam, P.; Morimoto, R. I.; Vendruscolo, M.; Dobson, C. M.; O'Brien, E. P. *Proc. Natl. Acad. Sci.* **2013**, *110* (2), E132–E140.

(65) Betancourt, M. R.; Thirumalai, D. *Protein Sci.* **2008**, *8* (2), 361–369.

(66) Frishman, D.; Argos, P. *Proteins Struct. Funct. Genet.* **1995**, *23*, 566–579.

(67) Kauffman, L.; Balachandran, A. P. *Phys. Today* **1992**.

(68) Oh, E.; Becker, A. H.; Sandikci, A.; Huber, D.; Chaba, R.; Gloge, F.; Nichols, R. J.; Typas, A.; Gross, C. A.; Kramer, G.; Weissman, J. S.; Bukau, B. *Cell* **2011**, *147* (6), 1295–1308.

(69) Guo, Z.; Thirumalai, D. *J. Mol. Biol.* **1996**, *263* (2), 323–343.

(70) Feng, Y.; De Franceschi, G.; Kahraman, A.; Soste, M.; Melnik, A.; Boersema, P. J.; De Laureto, P. P.; Nikolaev, Y.; Oliveira, A. P.; Picotti, P. *Nat. Biotechnol.* **2014**, *32* (10), 1036–1044.

(71) To, P.; Lee, S. O.; Xia, Y.; Devlin, T.; Fleming, K. G.; Fried, S. D. *bioRxiv* **2021** DOI: 10.1101/2021.11.20.469408.

(72) Kong, A. T.; Leprevost, F. V.; Avtonomov, D. M.; Mellacheruvu, D.; Nesvizhskii, A. I. *Nat. Methods* **2017**, *14* (5), 513–520.

(73) Steinhaus, H. *Bull. L'Académie Pol. des Sci.* **1956**, *4*, 801–804.

(74) MacQueen, J. In *Proceedings of the fifth Berkeley symposium on mathematical statistics and probability, Oakland, CA, USA:*; 1967; pp 281–297.

(75) Röblitz, S.; Weber, M. *Adv. Data Anal. Classif.* **2013**, *7* (2), 147–179.

1307 (76) Buchete, N. V.; Hummer, G. *J. Phys. Chem. B* **2008**, *112* (19), 6057–6069.
1308 (77) Scherer, M. K.; Trendelkamp-Schroer, B.; Paul, F.; Pérez-Hernández, G.;
1309   Hoffmann, M.; Plattner, N.; Wehmeyer, C.; Prinz, J. H.; Noé, F. *J. Chem.*
1310   *Theory Comput.* **2015**, *11* (11), 5525–5542.

Figure 1. **Luciferase exhibits subpopulations that misfold into soluble but less-functional conformations**. (a) Simulations of translation elongation and ejection of nascent Luciferase were performed with a coarse-grain ribosome representation (ribosomal proteins and RNA are displayed in red and grey respectively). Domains 1, 2, 3, and 4 of Luciferase are displayed in silver, light green, magenta, and light purple, respectively. After ejection, the ribosome is removed and post-release dynamics is simulated for 30 CPU days per trajectory. (b) Primary structure diagram of Luciferase colored as described in (a); positions involved in the catalytic function of Luciferase as described in Methods are colored blue. (c) Cartoon diagram of Luciferase native state colored as described in (a) with the 5′-O-[N-(dehydroluciferyl)-sulfamoyl]-adenosine ligand colored dark blue. (d) Q_{mode} (see Methods) versus time for Domain 2 of a trajectory of Luciferase that folded correctly. Portions of the plot colored green, yellow, and blue correspond to the synthesis, ejection, and post-translation phases of the simulation. Note that the relatively short duration of ejection for this protein renders that section of the plot invisible at this resolution. The red line corresponds to $\langle Q_{\mathrm{mode}}^{\mathrm{NS}} \rangle$ minus three standard deviations and represents the threshold for defining this domain as folded (see Methods). (e) Same as (d) but for a trajectory that misfolds. Inset shows the final microsecond of the Q_{mode} time series. (f) Distributions of χ_{func} (Eq. 13) over the final 100 ns of the folded (magenta) and misfolded (orange) trajectories displayed in panels (d) and (e). The misfolded and folded distributions are different based on the Kolmogorov-Smirnov test with test statistic 0.33 and p-value of 1 x 10^{-66}. The misfolded distribution shows greater structural distortion (i.e., values of $\chi_{\mathrm{func}} > 0$) of the binding pocket. (g) Backmapped all-atom structure from the final frame of the folded simulation shown in (d) aligned based on the residues implicated in function to the native state. (h) Same as (g) except for the final structure from the misfolded trajectory in (e), showing a strand misfolding in the ligand binding pocket (indicated by black arrow). Steric conflict between where the substrate binds and the surrounding binding pocket of the misfolded structures indicates this misfolded state will have reduced function.

Figure 2. **One in three proteins exhibit subpopulations that misfold into soluble but less-functional conformations that evade proteostasis machinery.** (a) Histogram of the probability of misfolding being detected in the final 100 ns of the simulation, computed as the number of misfolded trajectories divided by 50, for each of the 122 proteins in the cytosolic *E. coli* proteome set. (b) Histogram of the percent change, computed as $\frac{|\langle Q_{overall}\rangle - \langle Q_{overall}^{NS}\rangle|}{\langle Q_{overall}^{NS}\rangle} * 100\%$, in fraction of native contacts within the final 100 ns of each of the 1,631 misfolded trajectories ($\langle Q_{overall}\rangle$) in the *E. coli* proteome data set relative to the average value from each protein's native state simulations ($\langle Q_{overall}^{NS}\rangle$). The majority of misfolded proteins are within 10% of the native value. (c) Histogram of extrapolated folding times for the slow-folding kinetic phase from survival probability curves for the 73 proteins in the cytosolic *E. coli* dataset with a reliable estimate (see Methods). (d) Cumulative distribution function (CDF) of $\langle \zeta_{hydrophobic}\rangle$ computed over the values of $\langle \zeta_{hydrophobic}\rangle$ (Eq. 10) for 1,631 misfolded (orange) and 4,469 folded (magenta) trajectories. The blue shaded region indicates the set of $\langle \zeta_{hydrophobic}\rangle$ values considered to have no significant increase in hydrophobic solvent-accessible surface area relative to the native-state ensemble. (e) Same as (d) but CDFs are computed over values of $\langle \chi_{func}\rangle$ for trajectories in the misfolded and folded populations. Blue shaded region indicates the set of values considered to result in perturbed function. (f) Venn diagram indicating the number of the 1,631 misfolded trajectories that evade chaperones (TF, DnaK, and GroEL/GroES), do not aggregate, are not degraded, and are non-functional. The 186 trajectories at the center of this diagram are misfolded states that are expected to evade the proteostasis machinery, remaining soluble but non-functional. All error bars are 95% confidence intervals computed from bootstrapping 10^6 times; the height of the CDF plots in (d) and (e) indicates the 95% confidence intervals.

		Entangled	
		Yes	No
Misfolded based on Q_{mode}	Yes	1,016 [959, 1,073]	294 [262, 327]
	No	321 [287, 356]	4,469 [4,400, 4,537]

Odds ratio = 48.1 [40.7, 57.5]
p-value < 1 x 10^{-100}

		Slow folding	
		Yes	No
Entangled	Yes	29 [21, 37]	8 [3, 14]
	No	7 [3, 12]	29 [21, 37]

Odds ratio = 15.0 [5.4, 70.0]
p-value = 5 x 10^{-7} [2 x 10^{-11}, 1 x 10^{-3}]

Figure 3. **Detecting non-native entanglements in a monomeric protein structure.** (a) Schematic of how self-entanglements can be detected by examining the change in the Gauss linking number g(i,j) (Eq. 5) between a closed loop (pink) formed by the backbone segment between residues i and j that form a native contact (gold dashed line) and another pseudo-closed loop formed by the C-terminal backbone segment (blue) and a pseudo-vector (dashed green line) connecting the C-terminal residue and the start of the C-terminal segment, which begins at residue j+1. Threading of the N-terminal segment (composed of residues 1 through i-1) is determined in a similar manner. Examples of different Gauss linking numbers and their corresponding structures are shown in this hypothetical illustration. The magnitude of g(i,j) is proportional to the number of threading events of the blue segment through the pink loop, while its sign is a function of the relative positioning of primary structure vectors at crossing points between the pink and blue segments. The structure with g(i,j) =0 exhibits no entanglement. (b) An example of a gain in entanglement of the protein YJGH (PDB: 1PF5), where the C-termini (cyan) threads a loop (pink) formed by the native contact between residues D72 & Y104 (gold). Black arrow indicates the location of the crossing point of the two entangled loops. (c) Contingency table indicating the number of trajectories that are misfolded/folded across our 122 proteins based on Q_{mode} analysis and entangled/not entangled. Indicated p-values and odds ratios were computed in SciPy using the Fisher Exact Test. (d) Same as (c) except contingency table displays the number of proteins that are entangled/not entangled and predicted to be slow folding/fast folding. For the purposes of this analysis, a protein is considered slow- or fast-folding if its computed folding time is above or below the median folding time from the set of 73 proteins with reliable estimates, respectively. A protein is considered entangled if ≥50% of its misfolded trajectories are entangled. All error bars are 95% confidence intervals computed from bootstrapping 10^6 times.

Figure 4. The vast majority of trajectories predicted to bypass cellular quality controls and exhibit reduced function are entangled. The percent of trajectories out of 50 for each of the 38 proteins that bypass quality controls and are predicted to have reduced function that are entangled (blue) or not entangled (red). A total of 174 out of 186 trajectories are entangled. Protein names were taken from UNIPROT; see Table S2 for the structures used and their corresponding gene names. Inset contingency table indicates the number of trajectories that are misfolded and escape proteostasis machinery while remaining non-functional and entangled/not entangled. Indicated p-value and odds ratio were computed in SciPy using the Fisher Exact Test. All error bars are 95% confidence intervals from bootstrapping 10^6 times.

Figure 5. The two-domain glycerol-3-phosphate dehydrogenase protein displays widespread misfolding into soluble but less-functional conformations. (a) Ribbon structure of glycerol-3-phosphate dehydrogenase (PDB ID: 2QCU). Domains 1 and 2 are composed of residues 1-387 and 388-501 and are colored cyan and green, respectively. The FAD cofactor is shown in a dark blue representation. (b) Number of folded and misfolded trajectories for this protein in our simulations. (c) Q_{mode} versus time for Domain 1 for the subpopulations of folded (magenta) and misfolded (orange) trajectories. Each line represents one independent trajectory. (d) Survival probability of the unfolded state versus time computed for Domain 1 (cyan), Domain 2 (green), and the Domain 1|2 interface as described in Methods. Dotted black lines are double-exponential fits used to extract rate constants. (e) Summary of key parameters for the folded and misfolded populations. (f) 50% [36%, 64%] of glycerol-3-phosphate dehydrogenase trajectories are predicted to remain soluble but non-functional. (g) Representative misfolded structure colored as in (a) aligned to the native-state reference structure. Despite a high χ_{func} value indicative of a less-functional conformation, the protein is largely native. (h) Representative folded and misfolded structures back-mapped to all-atom resolution and then aligned to the native state based on the FAD binding pocket residues. Steric conflict (indicated by black arrow) can be seen between the substrate binding location and the misfolded binding pocket, indicating reduced function of this conformation. (i) Three pairs of structures corresponding to the native state (left) and the first representative structure of metastable state S2 (right) with locations of [333-354], F351, and L293 indicated in magenta. The loop (residues 271-288) and threading (residues 218-237) segments of the entanglement present in S2 are shown in red and blue, respectively. The same regions are colored yellow and light purple in the native state for reference, though no entanglement is present. The CA atoms of residues 271 and 288 that form the contact closing the loop segment are represented by orange spheres. Values of $\langle\zeta_{\mathrm{peptide}}\rangle$ were calculated with Eq. 16; error bars are available in Table S11. All error bars are 95% confidence intervals computed from bootstrapping 10^6 times.

Supplementary Information

Universal protein misfolding intermediates can bypass the proteostasis network and remain soluble and less functional

Daniel Nissley[1,ǂ], Yang Jiang[1,ǂ], Fabio Trovato[1], Ian Sitarik[1], Karthik Narayan[1], Philip To[2], Yingzi Xia[2], Stephen Fried[2,3], Edward P. O'Brien[1,4,5*]

[1]Department of Chemistry, Pennsylvania State University, University Park, PA 16802, USA
[2]Department of Chemistry, Johns Hopkins University, Baltimore, MD 21218, USA
[3]Thomas C. Jenkins Department of Biophysics, Johns Hopkins University, Baltimore, MD 21218, USA
[4]Bioinformatics and Genomics Graduate Program, The Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, PA 16802, USA
[5]Institute for Computational and Data Sciences, Pennsylvania State University, University Park, PA 16802, USA
[ǂ]These authors contributed equally to this research project
* to whom correspondence should be addressed: epo2@psu.edu

Supplementary Methods

Preparation of K12 Cell Pellets. *E. coli* K12 cells (NEB) were grown in 2 sets of 3 × 50 mL (biological triplicates) of in-house prepared MOPS EZ rich media(-Arginine/-Lysine) from saturated overnight cultures with a starting OD_{600} of 0.05. Similar to reported elsewhere[71], one set was supplemented with 0.5 mM [$^{13}C_6$]L-Arginine and 0.4 mM [$^{13}C_6$]L-Lysine and the other with 0.5 mM L-Arginine and 0.4 mM L-Lysine. Cells were cultured at 37˚C with agitation (220 rpm) to a final OD_{600} of 0.8. Each heavy/light pair was pooled together and then transferred to 2 × 50 mL falcon tubes and collected by centrifugation at 4000 *g* for 15 mins at 4˚C. The supernatants were removed, and cell pellets were stored at -20˚C until further use.

Frozen cell pellets were resuspended in a lysis buffer consisting of 900 µL of Tris pH 8.2 (20 mM Tris pH 8.2, 100 mM NaCl, 2 mM $MgCl_2$ and supplemented with DNase I to a final concentration (f.c.) of 0.1 mg mL^{-1}). Resuspended cells were flash frozen by slow drip over liquid nitrogen and cryogenically pulverized with a freezer mill (SPEX Sample Prep) over 8 cycles consisting of 1 min of grinding (9 Hz), and 1 min of cooling. Pulverized lysates were transferred to 50 mL centrifuge tubes and thawed at room temperature for 20 min. Lysates were then transferred to fresh 1.5 mL microfuge tubes and clarified at 16000 *g* for 15 min at 4 °C to remove insoluble cell debris. To deplete ribosome particles, clarified lysates were transferred to 3 mL *konical* tubes and ultracentrifuged at 33,300 rpm at 4 °C for 90 min without sucrose cushions using a SW55 Ti rotor. Protein concentrations of clarified lysates were determined using the bicinchoninic acid assay (Rapid Gold BCA Assay, Pierce) in a microtiter format with a plate reader (Molecular Devices iD3) using BSA as a calibration standard. Protein concentrations were diluted to a standard concentration of 3.3 mg mL^{-1} using Tris lysis buffer. This generates the normalized lysates for all downstream workflows.

Preparation of Native and Refolded Lysates for Limited Proteolysis Mass Spectrometry. To prepare half-isotopically-labeled native samples, 3.5 µL of normalized lysates derived from pellets in which half of the cells were grown with [$^{13}C_6$]L-Arginine and [$^{13}C_6$]L-Lysine during cell culture and half of the cells were grown with natural abundance L-Arginine and L-Lysine during cell culture, were diluted with 96.5 µL of Tris native dilution buffer (20 mM Tris pH 8.2, 100 mM NaCl, 10.288 mM $MgCl_2$, 10.36 mM KCl, 2.07 mM ATP, 1.04 mM DTT, 62 mM GdmCl) to a final protein concentration of 0.115 mg mL^{-1}. Following dilution, the final concentrations are 20 mM Tris pH 8.2, 100 mM NaCl, 10 mM $MgCl_2$, 10 mM KCl, 2mM ATP, 1 mM DTT and 60 mM GdmCl. Native samples were then equilibrated by incubating for 90 min at room temperature prior to limited proteolysis.

The refolding samples were prepared as described previously[49]. Briefly: 600 µL of normalized lysates, 100 mg of solid GdmCl, and 2.4 µL of a freshly prepared 700 mM DTT stock solution were added to a fresh 1.5 mL microfuge tube, and solvent was removed using a vacufuge plus to a final volume of 170 µL, such that the final concentrations of all components were 11.6 mg mL^{-1} protein, 6 M GdmCl, 70 mM Tris pH 8.2, 350 mM NaCl, 7 mM $MgCl_2$, and 10 mM DTT. Unfolded lysates were incubated overnight at room temperature to complete unfolding prior to refolding.

To prepare refolding samples, 99 µL of refolding dilution buffer (19.5 mM Tris pH 8.2, 97.5 mM NaCl, 10.03 mM $MgCl_2$, 10.1 mM KCl, 2.02 mM ATP and .909 mM DTT) were added to a fresh 1.5 mL microfuge tube. 1 µL of unfolded extract was then added to the tube containing the refolding dilution buffer and quickly mixed by rapid

vortexing, diluting the sample by 100x, followed by flash centrifugation to collect liquids
to the bottom of the tube. The final concentrations were 20 mM Tris pH 8.2, 100 mM
NaCl, 10 mM MgCl$_2$, 10 mM KCl, 2mM ATP, 1 mM DTT and 60 mM GdmCl. Refolded
samples were then incubated at room temperature for 1min, 5 min or 2 h to allow for
proteins to refold prior to limited proteolysis.

To perform limited proteolysis, 2 µL of a PK stock (prepared as a 0.067 mg mL^{-1}
PK in a 1:1 mixture of Tris lysis buffer and 20% glycerol, stored at -20˚C and thawed
at most only once) were added to a fresh 1.5 mL microfuge tube. After refolded
proteins were allowed to refold for the specified amount of time (1 min, 5 min, or 2 h),
or native proteins were allowed their 90 min equilibration, 100 µL of the native/refolded
lysates were added to the PK-containing microfuge tube and quickly mixed by rapid
vortexing (enzyme:substrate ratio is a 1:100 w/w ratio[70]), followed by flash
centrifugation to collect liquids to the bottom of the tube. Samples were incubated for
exactly 1 min at room temperature before transferring them to a mineral oil bath
preequilibrated at 110°C for 5 min to quench PK activity. Boiled samples were then
flash centrifuged (to collect condensation on the sides of the tube), and transferred to
fresh 1.5 mL microfuge tube containing 76 mg urea such that the final urea
concentration was 8 M and the final volume was 158 µL. They are then vortexed to
dissolve the urea to unfold all proteins and quench any further enzyme activity
indefinitely, and flash centrifuged to collect liquids to the bottom of the tubes.

All protein samples were prepared for mass spectrometry as follows: 2.25 µL
of a freshly prepared 700 mM stock of DTT were added to each sample-containing
microfuge tube to a final concentration of 10 mM. Samples were incubated at 37°C for
30 minutes at 700 rpm on a thermomixer to reduce cysteine residues. 9 µL of a freshly
prepared 700 mM stock of iodoacetamide (IAA) were then added to a final
concentration of 40 mM, and samples were incubated at room temperature in the dark
for 45 minutes to alkylate reduced cysteine residues. 1 µL of a 0.1 µg µL^{-1} stock of
LysC (NEB) was added to the samples (to a final enzyme:substrate ratio of 1:100 w/w)
and incubated for 2 h at 37°C at 700 rpm. 471 µL of 100 mM ammonium bicarbonate
(pH 8) were added to the samples to dilute the urea to a final concentration of 2 M. 2
µL of a 0.1 µg µL^{-1} stock of Trypsin (NEB) were added to the samples (to a final
enzyme:substrate ratio of 1:50 w/w) and incubated overnight (15-16 h) at 25°C at 700
rpm (not 37˚C, so as to minimize decomposition of urea and carbamylation of lysines).
1596
Desalting of Mass Spectrometry Samples. Peptides were desalted with Sep-Pak
C18 1 cc Vac Cartridges (Waters) over a vacuum manifold. Tryptic digests were first
acidified by addition of 16.6 µL trifluoroacetic acid (TFA, Acros) to a final concentration
of 1% (vol/vol). Cartridges were first conditioned (1 mL 80% ACN, 0.5% TFA) and
equilibrated (4 x 1 mL 0.5% TFA) before loading the sample slowly under a diminished
vacuum (ca. 1 mL/min). The columns were then washed (4 x 1 mL 0.5% TFA), and
peptides were eluted by addition of 1 mL elution buffer (80% ACN, 0.5% TFA). During
elution, vacuum cartridges were suspended above 15 mL conical tubes, placed in a
swing-bucket rotor (Eppendorf 5910R), and spun for 3 min at 350 g. Eluted peptides
were transferred from Falcon tubes back into microfuge tubes and dried using a
vacuum centrifuge (Eppendorf Vacufuge). Dried peptides were stored at -80°C until
analysis. For analysis, samples were vigorously resuspended in 0.1% FA in Optima
water (ThermoFisher) to a final concentration of 0.5 mg mL^{-1}.
1610
LC-MS/MS Acquisition. Chromatographic separation of digests were carried out on
a Thermo UltiMate3000 UHPLC system with an Acclaim Pepmap RSLC, C18, 75 µm

× 25 cm, 2 µm, 100 Å column. Approximately, 1 µg of protein was injected onto the
column. The column temperature was maintained at 40 °C, and the flow rate was set
to 0.300 µL min^{-1} for the duration of the run. Solvent A (0.1% FA) and Solvent B (0.1%
FA in ACN) were used as the chromatography solvents. The samples were run
through the UHPLC System as follows: peptides were allowed to accumulate onto the
trap column (Acclaim PepMap 100, C18, 75 µm x 2 cm, 3 µm, 100 Å column) for 10
min (during which the column was held at 2% Solvent B). The peptides were resolved
by switching the trap column to be in-line with the separating column, quickly
increasing the gradient to 5% B over 5 min and then applying a 95 min linear gradient
from 5% B to 25% B. Subsequently, the gradient was increased from 35% B to 40%
B over 25 min and then increased again from 40% B to 90% B over 5 min. The column
was then cleaned with a sawtooth gradient to purge residual peptides between runs in
a sequence.
A Thermo Q-Exactive HF-X Orbitrap mass spectrometer was used to analyze
protein digests. A full MS scan in positive ion mode was followed by 20 data-dependent
MS scans. The full MS scan was collected using a resolution of 120000 (@ m/z 200),
an AGC target of 3E6, a maximum injection time of 64 ms, and a scan range from 350
to 1500 m/z. The data-dependent scans were collected with a resolution of 15000 (@
m/z 200), an AGC target of 1E5, a minimum AGC target of 8E3, a maximum injection
time of 55 ms, and an isolation window of 1.4 m/z units. To dissociate precursors prior
to their reanalysis by MS2, peptides were subjected to an HCD of 28% normalized
collision energies. Fragments with charges of 1, 6, 7, or higher and unassigned were
excluded from analysis, and a dynamic exclusion window of 30.0 s was used for the
data-dependent scans. Mass tags were enabled with Δm of 2.00671 Th, 3.01007 Th,
4.01342 Th, and 6.02013 Th.
1638
LC-MS/MS Data Analysis. Proteome Discoverer (PD) Software Suite (v2.4, Thermo
Fisher) and the Minora Algorithm were used to analyze mass spectra and perform
Label Free Quantification (LFQ) of detected peptides. Default settings for all analysis
nodes were used except where specified. The data were searched against Escherichia
coli (UP000000625, Uniprot) reference proteome database. For peptide identification,
the PD MSFragger node was used, using a semi-tryptic search allowing up to 2 missed
cleavages[72]. A precursor mass tolerance of 10 ppm was used for the MS1 level, and
a fragment ion tolerance was set to 0.02 Da at the MS2 level. Peptide lengths between
7 and 50 amino acid residues was allowed with a peptide mass between 500 and 5000
Da. Additionally, a maximum charge state for theoretical fragments was set at 2.
Oxidation of methionine and acetylation of the N-terminus were allowed as dynamic
modifications while carbamidomethylation on cysteines was set as a static
modification. Heavy isotope labeling ($^{13}C_6$) of Arginine and Lysine were allowed as
dynamic modifications. The Philosopher PD node was used for FDR validation. Raw
normalized extracted ion intensity data for the identified peptides were exported from
the .pdResult file using a three-level hierarchy (protein > peptide group > consensus
feature). These data were further processed utilizing custom Python analyzer scripts
(available on GitHub, and described in depth previously[49,71]). Briefly, normalized ion
counts were collected across the refolded replicates and the native replicates for each
successfully identified peptide group. Effect sizes are the ratio of averages (reported
in log$_2$) and P-values (reported as –log$_{10}$) were assessed using t tests with Welch's
correction for unequal population variances. Missing data are treated in a special
manner. If a feature is not detected in all three native (or refolded) injections and is
detected in all three refolded (or native) injections, we use those data, and fill the

missing values with 1000 (the ion limit of detection for this mass analyzer); this peptide becomes classified as an all-or-nothing peptide. If a feature is not detected in one out of six injections, the missing value is dropped. Any other permutation of missing data (e.g., missing in two injections) results in the quantification getting discarded. In many situations, our data provide multiple independent sets of quantifications for the same peptide group. This happens most frequently because the peptide is detected in multiple charge states or as a heavy isotopomer. In this case, we calculate effect size and P-value for all features that map to the same peptide group. If the features all agree with each other in sign, they are combined: the quantification associated with the median amongst available features is used and the P-values are combined with Fisher's method. If the features disagree with each other in sign, the P-value is set to 1. Coefficients of variation (CV) for the peptide abundance in the three replicate refolded samples are also calculated. Analyzer returns a file listing all the peptides that can be confidently quantified, and provides their effect-size, P-value, refolded CV, proteinase K site (if half-tryptic), and associated protein metadata.

1713
1714 **Figure S1**. Venn diagrams showing the overlap between Luciferase trajectories that are misfolded and
1715 (a) predicted to evade DnaK, GroEL, and TF or (b) predicted not to aggregate, not to be degraded, to
1716 evade all chaperones, and to remain non-functional. Error bars are 95% confidence intervals from
1717 bootstrapping 10^6 times.

1740
1741 **Figure S2**. Cumulative distribution function of $\langle \zeta^{co-t}_{hydrophobic} \rangle$ over the subset of 1,053 misfolded
1742 trajectories for which it was computed (see Methods). The blue shaded region indicates the subset of
1743 values taken to indicate no significant increase in trigger factor interactions relative to the folded
1744 population. The height of the CDF indicates the 95% confidence intervals.

Figure S3. Cumulative distribution function of $\langle \zeta_{DnaK} \rangle$ over the subset of 1,631 misfolded (orange) and 4,469 folded (magenta) trajectories. The blue shaded region indicates the subset of values taken to indicate no significant increase in DnaK interactions relative to the native-state population. The height of the CDFs indicates the 95% confidence intervals.

Figure S4. Venn diagram showing the overlap between those trajectories that are predicted not to engage TF, DnaK, or GroEL/GroES. A total of 1,533 trajectories evade at least on chaperone. Error bars are 95% confidence intervals from bootstrapping 10^6 times.

1811
1812 **Figure S5**. Cumulative distribution function of $\langle \zeta_{Agg} \rangle$ over the subset of 1,631 misfolded (orange) and
1813 4,469 folded (magenta) trajectories. The blue shaded region indicates the subset of values taken to
1814 indicate no significant increase in aggregation propensity relative to the native state simulations. The
1815 height of the CDFs indicates the 95% confidence intervals.

1842

Figure S6. The utility of the change in entanglement metric (G) for detecting structural perturbations not apparent by fraction of native contacts (Q) or root mean square deviation (RMSD) is exemplified by examining clusters of the trajectories for aminoimidazole ribonucleotide synthetase (PDB: 1CLI) relative to the cluster of trajectories started from the reference state. The 50 trajectories obtained from synthesis simulations can be clustered into 5 separate clusters characterized by the $\langle Q_{\mathrm{mode}} \rangle$ and the corrected discrete distribution of a give type of change in entanglement (Eq. 7). cID0 contains 8/50 trajectories all of which terminate to the native state with no changes in entanglement. cID1 contains 3/50 trajectories that are near native like (i.e. $\langle Q_{\mathrm{mode}} \rangle \approx \langle Q_{\mathrm{mode}}^{NS} \rangle$) that show appreciable gain and loss of entanglements as well as pure switch in chirality. cID2 contains 32/50 trajectories that show appreciable gain and loss of entanglements with less near native like conformations (i.e. $\langle Q_{\mathrm{mode}} \rangle \geq \langle Q_{\mathrm{mode}}^{NS} \rangle - \sigma \langle Q_{\mathrm{mode}}^{NS} \rangle$). cID3 contains 6/50 trajectories that show non-native like conformations (i.e. $\langle Q_{\mathrm{mode}} \rangle < \langle Q_{\mathrm{mode}}^{NS} \rangle - \sigma \langle Q_{\mathrm{mode}}^{NS} \rangle$) with appreciable gain and loss of entanglements. cID4 contains 1/50 trajectories that show non-native like conformations with appreciable gain and loss of entanglements and pure change in chirality. For the clusters which contain near native like conformations and changes in entanglement of any kind (cID1 light blue and cID2) Q and RMSD fail to distinguish different clusters while G does not. For those clusters that contain non-native like conformations and changes in entanglement of any kind (cID3 & cID4) Q and RMSD are much more reliable.

1869 **Table S1.** Coarse-grain model parameters for PDB ID 4G36

PDB ID	Protein name	Domain or interface	Structural class	η used in model building
4G36	Firefly Luciferase	Domain 1:1-12; 356-437	β	1.442
		Domain 2:13-52; 212-355	α/β	1.114
		Domain 3: 53-211	α/β	1.114
		Domain 4: 438-550	α/β	1.916
		Interface 1|2	-	1.235
		Interface 1|3	-	1.235
		Interface 1|4	-	1.507*
		Interface 2|3	-	1.235
		Interface 2|4	-	1.235
		Interface 3|4	-	1.235

1870 *indicates interface is unstable at all tested values of η and so median interface
1871 value from set of stable interfaces is used for all simulations

Table S2 *E. coli* proteome dataset information

Index	PDB ID	Chain ID	UniProt Entry name	Protein name	Gene names
1	1A69	A	DEOD_ECOLI	Purine nucleoside phosphorylase DeoD-type	deoD, pup, b4384, JW4347
2	1A6J	B	PTSN_ECOLI	Nitrogen regulatory protein	ptsN, rpoP, yhbl, b3204, JW3171
3	1A82	A	BIOD1_ECOLI	ATP-dependent dethiobiotin synthetase BioD 1	bioD1, b0778, JW0761
4	1AG9	A	FLAV_ECOLI	Flavodoxin 1	fldA, b0684, JW0671
5	1AH9	Model 1	IF1_ECOLI	Translation initiation factor IF-1	infA, b0884, JW0867
6	1AKE	A	KAD_ECOLI	Adenylate kinase	Adk, dnaW, plsA, b0474, JW0463
7	1B9L	A	FOLX_ECOLI	Dihydroneopterin triphosphate 2'-epimerase	folX, b2303, JW2300
8	1CLI	A	PUR5_ECOLI	Phosphoribosylformylglycinamidine cyclo-ligase	purM, purG, b2499, JW2484
9	1D2F	A	MALY_ECOLI	Protein MalY	malY, b1622, JW1614
10	1DCJ	Model 1	TUSA_ECOLI	Sulfur carrier protein TusA	tusA, sirA, yhhP, b3470, JW3435
11	1DFU	P	RL25_ECOLI	50S ribosomal protein L25	rplY, b2185, JW2173
12	1DUV	G	OTC1_ECOLI	Ornithine carbamoyltransferase subunit I	argI, b4254, JW4211
13	1DXE	A	GARL_ECOLI	5-keto-4-deoxy-D-glucarate aldolase	garL, yhaF, b3126, JW3095
14	1EF9	A	SCPB_ECOLI	Methylmalonyl-CoA decarboxylase	scpB, mmcD, ygfG, b2919, JW2886
15	1EIX	C	PYRF_ECOLI	Orotidine 5'-phosphate decarboxylase	pyrF, b1281, JW1273
16	1EM8	A	HOLC_ECOLI	DNA polymerase III subunit chi	holC, b4259, JW4216
17	1EUM	A	FTNA_ECOLI	Bacterial non-heme ferritin	ftnA, ftn, gen-165, rsgA, b1905, JW1893
18	1FJJ	A	YBHB_ECOLI	UPF0098 protein YbhB	ybhB, b0773, JW0756
19	1FM0	D	MOAD_ECOLI	Molybdopterin synthase sulfur carrier subunit	moaD, chlA4, chlM, b0784, JW0767
20	1FTS	A	FTSY_ECOLI	Signal recognition particle receptor FtsY	ftsY, b3464, JW3429
21	1FUI	A	FUCI_ECOLI	L-fucose isomerase, Fuclase	fucI, b2802, JW2773
22	1GER	B	GSHR_ECOLI	Glutathione reductase	gor, b3500, JW3467
23	1GLF	O	GLPK_ECOLI	Glycerol kinase	glpK, b3926, JW3897
24	1GQE	A	RF2_ECOLI	Peptide chain release factor RF2	prfB, supK, b2891, JW5847
25	1GQT	B	RBSK_ECOLI	Ribokinase	rbsK, b3752, JW3731
26	1GT7	A	RHAD_ECOLI	Rhamnulose-1-phosphate aldolase	rhaD, rhuA, b3902, JW3873
27	1GYT	L	AMPA_ECOLI	Cytosol aminopeptidase	pepA, carP, xerB, b4260, JW4217
28	1GZ0	C	RLMB_ECOLI	23S rRNA (guanosine-2'-O-)-methyltransferase RlmB	rlmB, yjfH, b4180, JW4138
29	1H16	A	PFLB_ECOLI	Formate acetyltransferase 1	pflB, pfl, b0903, JW0886
30	1H75	A	NRDH_ECOLI	Glutaredoxin-like protein NrdH	nrdH, ygaN, b2673, JW2648

Table S2 cont.

Index	PDB ID	Chain ID	UniProt Entry name	Protein name	Gene names
31	1I6O	B	CAN_ECOLI	Carbonic anhydrase 2	can, cynT2, yadF, b0126, JW0122
32	1JNS	Model 1	PPIC_ECOLI	Peptidyl-prolyl cis-trans isomerase C	ppiC, parVA, b3775, JW3748
33	1JW2	A	HHA_ECOLI	Hemolysin expression-modulating protein Hha	hha, b0460, JW0449
34	1JX7	A	YCHN_ECOLI	Protein YchN	ychN, b1219, JW1210
35	1K7J	A	YCIO_ECOLI	Uncharacterized protein YciO	yciO, b1267, JW5196
36	1KO5	A	GNTK_ECOLI	Thermoresistant gluconokinase	gntK, b3437, JW3400
37	1KSF	X	CLPA_ECOLI	ATP-dependent Clp protease ATP-binding subunit ClpA	clpA, lopD, b0882, JW0866
38	1L6W	A	FSAA_ECOLI	Fructose-6-phosphate aldolase 1	fsaA, fsa, mipB, ybiZ, b0825, JW5109
39	1M3U	A	PANB_ECOLI	3-methyl-2-oxobutanoate hydroxymethyltransferase	panB, b0134, JW0130
40	1MZG	B	SUFE_ECOLI	Cysteine desulfuration protein SufE	sufE, ynhA, b1679, JW1669
41	1NAQ	A	CUTA_ECOLI	Divalent-cation tolerance protein CutA	cutA, cutA1, cycY, b4137, JW4097
42	1NG9	A	MUTS_ECOLI	DNA mismatch repair protein MutS	mutS, fdv, b2733, JW2703
43	1ORO	A	PYRE_ECOLI	Orotate phosphoribosyltransferase	pyrE, b3642, JW3617
44	1P7L	A	METK_ECOLI	S-adenosylmethionine synthase	metK, metX, b2942, JW2909
45	1P91	A	RLMA_ECOLI	23S rRNA (guanine(745)-N(1))-methyltransferase	rlmA, rrmA, yebH, b1822, JW1811
46	1PF5	A	YJGH_ECOLI	RutC family protein YjgH	yjgH, b4248, JW4206
47	1PMO	B	DCEB_ECOLI	Glutamate decarboxylase beta	gadB, b1493, JW1488
48	1PSU	B	PAAI_ECOLI	Acyl-coenzyme A thioesterase PaaI	paaI, ydbV, b1396, JW1391
49	1Q5X	A	RRAA_ECOLI	Regulator of ribonuclease activity A	rraA, menG, yiiV, b3929, JW3900
50	1QF6	A	SYT_ECOLI	Threonine-tRNA ligase	thrS, b1719, JW1709
51	1QTW	A	END4_ECOLI	Endonuclease 4	Nfo, b2159, JW2146
52	1RQJ	A	ISPA_ECOLI	Farnesyl diphosphate synthase	ispA, b0421, JW0411
53	1SG5	Model 1	ROF_ECOLI	Protein rof	Rof, yaeO, b0189, JW0184
54	1SV6	A	MHPD_ECOLI	2-keto-4-pentenoate hydratase	mhpD, b0350, JW0341
55	1SVT	J	CH60_ECOLI	60 kDa chaperonin	groL, groEL, mopA, b4143, JW4103
56	1T4B	A	DHAS_ECOLI	Aspartate-semialdehyde dehydrogenase	asd, hom, b3433, JW3396
57	1T8K	A	ACP_ECOLI	Acyl carrier protein	acpP, b1094, JW1080
58	1U0B	B	SYC_ECOLI	Cysteine–tRNA ligase	cysS, b0526, JW0515
59	1U60	A	GLSA1_ECOLI	Glutaminase 1	glsA1, ybaS, b0485, JW0474
60	1UUF	A	YAHK_ECOLI	Aldehyde reductase YahK	yahK, b0325, JW0317
61	1W78	A	FOLC_ECOLI	Dihydrofolate synthase/folylpolyglutamate synthase	folC, dedC, b2315, JW2312

Table S2 cont.

Index	PDB ID	Chain ID	UniProt Entry name	Protein name	Gene names
62	1W8G	A	PLPHP_ECOLI	Pyridoxal phosphate homeostasis protein	yggS, b2951, JW2918
63	1WOC	C	PRIB_ECOLI	Primosomal replication protein N	priB, b4201, JW4159
64	1XN7	Model 1	FEOC_ECOLI	Probable [Fe-S]-dependent transcriptional repressor FeoC	feoC, yhgG, b3410, JW3373
65	1XRU	A	KDUI_ECOLI	4-deoxy-L-threo-5-hexosulose-uronate ketol-isomerase	kduI, yqeE, b2843, JW2811
66	1XVI	A	MPGP_ECOLI	Mannosyl-3-phosphoglycerate phosphatase	yedP, b1955, JW1938
67	1YQQ	A	XAPA_ECOLI	Purine nucleoside phosphorylase 2	xapA, pndA, b2407, JW2398
68	1ZYL	A	SRKA_ECOLI	Stress response kinase A	srkA, rdoA, yihE, b3859, JW3831
69	1ZZM	A	YJJV_ECOLI	Uncharacterized metal-dependent hydrolase YjjV	yjjV, b4378, JW4341
70	2A6Q	E	YOEB_ECOLI	Toxin YoeB	yoeB, b4539, JW5331
71	2AXD	Model 1	HOLE_ECOLI	DNA polymerase III subunit theta	holE, b1842, JW1831
72	2D1P	B	TUSC_ECOLI	Protein TusC	tusC, yheM, b3344, JW3306
73	2FEK	Model 1	WZB_ECOLI	Low molecular weight protein-tyrosine-phosphatase Wzb	wzb, b2061, JW2046
74	2FYM	A	ENO_ECOLI	Enolase	eno, b2779, JW2750
75	2GQR	A	PUR7_ECOLI	Phosphoribosylaminoimidazole-succinocarboxamide synthase	purC, b2476, JW2461
76	2H1F	A	A0A0H2VC26_ECOL6	Lipopolysaccharide heptosyltransferase-1	rfaC, c4447
77	2HD3	K	EUTN_ECOLI	Ethanolamine catabolic microcompartment shell protein EutN	eutN, cchB, yffY, b2456, JW2440
78	2HG2	A	ALDA_ECOLI	Lactaldehyde dehydrogenase	aldA, ald, b1415, JW1412
79	2HGK	Model 1	YQCC_ECOLI	Uncharacterized protein YqcC	yqcC, b2792, JW2763
80	2HNA	A	MIOC_ECOLI	Protein MioC	mioC, yieB, b3742, JW3720
81	2HNH	A	DPO3A_ECOLI	DNA polymerase III subunit alpha	dnaE, polC, b0184, JW0179
82	2HO9	Model 1	CHEW_ECOLI	Chemotaxis protein CheW	cheW, b1887, JW1876
83	2ID0	A	RNB_ECOLI	Exoribonuclease 2	rnb, b1286, JW1279
84	2JEE	C	ZAPB_ECOLI	Cell division protein ZapB	zapB, yiiU, b3928, JW3899
85	2JO6	Model 1	NIRD_ECOLI	Nitrite reductase (NADH) small subunit	nirD, b3366, JW3329
86	2JRX	Model 1	YEJL_ECOLI	UPF0352 protein YejL	yejL, b2187, JW2175
87	2KC5	Model 1	HYBE_ECOLI	Hydrogenase-2 operon protein HybE	hybE, b2992, JW2960
88	2KFW	Model 1	SLYD_ECOLI	FKBP-type peptidyl-prolyl cis-trans isomerase SlyD, PPIase	slyD, b3349, JW3311
89	2KX9	A	PT1_ECOLI	Phosphoenolpyruvate-protein phosphotransferase	ptsI, b2416, JW2409
90	2O1C	A	NUDB_ECOLI	Dihydroneopterin triphosphate diphosphatase	nudB, ntpA, b1865, JW1854
91	2OQ3	Model 1	PTMA_ECOLI	Mannitol-specific cryptic phosphotransferase enzyme IIA component	cmtB, b2934, JW2901
92	2PTH	A	PTH_ECOLI	Peptidyl-tRNA hydrolase	pth, b1204, JW1195
93	2PTQ	A	PUR8_ECOLI	Adenylosuccinate lyase	purB, b1131, JW1117
94	2QCU	B	GLPD_ECOLI	Aerobic glycerol-3-phosphate dehydrogenase	glpD, glyD, b3426, JW3389
95	2QVR	A	F16PA_ECOLI	Fructose-1,6-bisphosphatase class 1	fbp, fdp, b4232, JW4191

 Table S2 cont.

Index	PDB ID	Chain ID	UniProt Entry name	Protein name	Gene names
96	2R5N	A	TKT1_ECOLI	Transketolase 1	tktA, tkt, b2935, JW5478
97	2UYJ	A	TDCF_ECOLI	Putative reactive intermediate deaminase TdcF	tdcF, yhaR, b3113, JW5521
98	2V81	A	DGOA_ECOLI	2-dehydro-3-deoxy-6-phosphogalactonate aldolase	dgoA, yidU, b4477, JW5628
99	2WIU	A	HIPA_ECOLI	Ser/Thr-protein kinase HipA	hipA, b1507, JW1500
100	2WW4	A	ISPE_ECOLI	4-diphosphocytidyl-2-C-methyl-D-erythritol kinase, CMK	ispE, ipk, ychB, b1208, JW1199
101	2YVA	A	DIAA_ECOLI	DnaA initiator-associating protein DiaA	diaA, yraO, b3149, JW3118
102	3ASV	B	YDFG_ECOLI	NADP-dependent 3-hydroxy acid dehydrogenase YdfG	ydfG, b1539, JW1532
103	3BMB	B	RNK_ECOLI	Regulator of nucleoside diphosphate kinase	rnk, b0610, JW0602
104	3BRQ	B	ASCG_ECOLI	HTH-type transcriptional regulator AscG	ascG, b2714, JW5434
105	3GN5	B	MQSA_ECOLI	Antitoxin MqsA	mqsA, ygiT, b3021, JW2989
106	3HWO	A	ENTC_ECOLI	Isochorismate synthase EntC	entC, b0593, JW0585
107	3IV5	B	FIS_ECOLI	DNA-binding protein Fis	fis, b3261, JW3229
108	3M7M	X	HSLO_ECOLI	33 kDa chaperonin	hslO, yrfl, b3401, JW5692
109	3N1S	J	HINT_ECOLI	Purine nucleoside phosphoramidase	hinT, ycfF, b1103, JW1089
110	3NXC	A	SLMA_ECOLI	Nucleoid occlusion factor SlmA	slmA, ttk, yicB, b3641, JW5641
111	3OFO	D	RS4_ECOLI	30S ribosomal protein S4	rpsD, ramA, b3296, JW3258
112	3PCO	D	SYFB_ECOLI	Phenylalanine--tRNA ligase beta subunit	pheT, b1713, JW1703
113	3QOU	A	CNOX_ECOLI	Chaperedoxin	cnoX, ybbN, b0492, JW5067
114	4A2C	A	GATD_ECOLI	Galactitol 1-phosphate 5-dehydrogenase	gatD, b2091, JW2075
115	4DCM	A	RLMG_ECOLI	Ribosomal RNA large subunit methyltransferase G	rlmG, ygjO, b3084, JW5513
116	4DZD	A	CAS6_ECOLI	CRISPR system Cascade subunit CasE	casE, cas6e, ygcH, b2756, JW2726
117	4E8B	A	RSME_ECOLI	Ribosomal RNA small subunit methyltransferase E	rsmE, yggJ, b2946, JW2913
118	4FZW	A	PAAF_ECOLI	2,3-dehydroadipyl-CoA hydratase	paaF, ydbR, b1393, JW1388
119	4HR7	A	ACCC_ECOLI	Biotin carboxylase	accC, fabG, b3256, JW3224
120	4IM7	A	A0A0H2V7F2_ECOL6	Hypothetical oxidoreductase ydfI	c1968
121	4IWX	A	RIMK_ECOLI	Ribosomal protein S6--L-glutamate ligase	rimK, b0852, JW0836
122	4KN7	A	RPOB_ECOLI	DNA-directed RNA polymerase subunit beta, RNAP subunit beta	rpoB, groN, nitB, rif, ron, stl

1924 **Table S3**. Definitions of entanglement types

Type	Change in entanglement	Change in chirality	Conditional*
G0	Gain	No	$\left\|g^{\text{current}}(i,j)\right\| > \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) \geq 0$
G1	Gain	Yes	$\left\|g^{\text{current}}(i,j)\right\| > \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) < 0$
G2	Lose	No	$\left\|g^{\text{current}}(i,j)\right\| < \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) \geq 0$
G3	Lose	Yes	$\left\|g^{\text{current}}(i,j)\right\| < \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) < 0$
G4	None	Yes	$\left\|g^{\text{current}}(i,j)\right\| = \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) < 0$
G5	None	None	$\left\|g^{\text{current}}(i,j)\right\| = \left\|g^{\text{native}}(i,j)\right\|$ and $g^{\text{current}}(i,j) \times g^{\text{native}}(i,j) \geq 0$

1925 * conditional used to determine if a native contact between residues i and j has a
1926 change in a given type of entanglement in the current structure relative to the native
1927 state structure.

1965 **Table S4.** Proteins categorized by percent of trajectories misfolded based

Category	Number of proteins	PDB IDs
At least 2% misfolded	80	1A69, 1A6J, 1A82, 1AH9, 1DFU, 1DXE, 1FJJ, 1GQT, 1GT7, 1H16, 1I6O, 1K7J, 1L6W, 1M3U, 1NAQ, 1ORO, 1P91, 1PF5, 1PMO, 1Q5X, 1SV6, 1U60, 1W8G, 1WOC, 1YQQ, 1ZYL, 2HNA, 2JO6, 2O1C, 2PTH, 2V81, 2YVA, 3BMB, 3HWO, 4A2C, 1CLI, 1D2F, 1DUV, 1EF9, 1FTS, 1FUI, 1GER, 1GLF, 1GYT, 1GZ0, 1KSF, 1NG9, 1P7L, 1QF6, 1SVT, 1T4B, 1U0B, 1UUF, 1W78, 1XRU, 1XVI, 2FYM, 2H1F, 2HG2, 2HNH, 2ID0, 2KFW, 2KX9, 2PTQ, 2QCU, 2R5N, 2WIU, 2WW4, 3BRQ, 3GN5, 3M7M, 3PCO, 4DCM, 4DZD, 4E8B, 4FZW, 4HR7, 4IM7, 4IWX, 4KN7
At least 20% misfolded	49	1A69, 1DFU, 1DXE, 1GQT, 1H16, 1K7J, 1P91, 1SV6, 1U60, 1YQQ, 1ZYL, 2HNA, 2JO6, 2V81, 3HWO, 4A2C, 1CLI, 1DUV, 1FTS, 1FUI, 1GLF, 1GYT, 1GZ0, 1KSF, 1NG9, 1P7L, 1QF6, 1SVT, 1T4B, 1U0B, 1W78, 1XRU, 1XVI, 2FYM, 2HG2, 2HNH, 2ID0, 2KFW, 2KX9, 2QCU, 2R5N, 2WIU, 3GN5, 3PCO, 4E8B, 4HR7, 4IM7, 4IWX, 4KN7
At least 50% misfolded	27	1GQT, 1H16, 1K7J, 1P91, 1U60, 1YQQ, 2JO6, 2V81, 1CLI, 1FUI, 1GYT, 1GZ0, 1KSF, 1P7L, 1T4B, 1U0B, 1XVI, 2HNH, 2ID0, 2KX9, 2QCU, 2R5N, 3PCO, 4E8B, 4HR7, 4IM7, 4KN7
100% misfolded	9	1GQT, 1H16, 1FUI, 1GYT, 1GZ0, 1XVI, 4E8B, 4HR7, 4KN7

1978   **Table S5**. Extrapolated folding times for the 73 proteins with a reliable estimate

PDB ID	Simulated τ_F, ns	Estimated experimental τ_F, s	Estimated experimental τ_F, min	Estimated experimental τ_F, h	Estimated experimental τ_F, d
1SG5	1.08E+01	4.29E-02	7.15E-04	1.19E-05	4.97E-07
2AXD	1.08E+01	4.29E-02	7.15E-04	1.19E-05	4.97E-07
3IV5	1.15E+01	4.55E-02	7.58E-04	1.26E-05	5.27E-07
2JRX	1.57E+01	6.22E-02	1.04E-03	1.73E-05	7.20E-07
2KC5	1.68E+01	6.67E-02	1.11E-03	1.85E-05	7.72E-07
2HO9	2.10E+01	8.34E-02	1.39E-03	2.32E-05	9.65E-07
2JEE	2.28E+01	9.05E-02	1.51E-03	2.52E-05	1.05E-06
3QOU	3.77E+01	1.50E-01	2.49E-03	4.16E-05	1.73E-06
1H75	4.25E+01	1.69E-01	2.81E-03	4.68E-05	1.95E-06
1T8K	4.30E+01	1.70E-01	2.84E-03	4.73E-05	1.97E-06
2FEK	4.30E+01	1.70E-01	2.84E-03	4.74E-05	1.97E-06
1MZG	5.86E+01	2.32E-01	3.87E-03	6.45E-05	2.69E-06
3NXC	9.97E+01	3.96E-01	6.60E-03	1.10E-04	4.58E-06
1FM0	1.07E+02	4.26E-01	7.10E-03	1.18E-04	4.93E-06
2HGK	2.84E+02	1.13E+00	1.88E-02	3.13E-04	1.30E-05
4DCM	2.92E+02	1.16E+00	1.93E-02	3.21E-04	1.34E-05
1GQE	3.42E+02	1.36E+00	2.26E-02	3.76E-04	1.57E-05
1ZZM	3.51E+02	1.39E+00	2.32E-02	3.87E-04	1.61E-05
2KX9	3.55E+02	1.41E+00	2.35E-02	3.92E-04	1.63E-05
2OQ3	3.63E+02	1.44E+00	2.40E-02	4.00E-04	1.67E-05
1EIX	4.34E+02	1.72E+00	2.87E-02	4.78E-04	1.99E-05
1KO5	4.39E+02	1.74E+00	2.90E-02	4.84E-04	2.02E-05
1GT7	5.36E+02	2.13E+00	3.54E-02	5.91E-04	2.46E-05
3N1S	5.83E+02	2.31E+00	3.85E-02	6.42E-04	2.68E-05
2QVR	6.26E+02	2.48E+00	4.14E-02	6.90E-04	2.87E-05
1AG9	8.60E+02	3.41E+00	5.68E-02	9.47E-04	3.95E-05
1EUM	9.16E+02	3.64E+00	6.06E-02	1.01E-03	4.21E-05
3OFO	9.94E+02	3.94E+00	6.57E-02	1.10E-03	4.56E-05
2KFW	1.81E+03	7.20E+00	1.20E-01	2.00E-03	8.33E-05
3BMB	3.36E+03	1.33E+01	2.22E-01	3.70E-03	1.54E-04
2PTH	3.39E+03	1.35E+01	2.24E-01	3.74E-03	1.56E-04
2GQR	3.82E+03	1.51E+01	2.52E-01	4.21E-03	1.75E-04
1QTW	4.13E+03	1.64E+01	2.73E-01	4.55E-03	1.90E-04
1JX7	4.43E+03	1.76E+01	2.93E-01	4.88E-03	2.03E-04
2YVA	5.42E+03	2.15E+01	3.59E-01	5.98E-03	2.49E-04
1PSU	1.05E+04	4.16E+01	6.93E-01	1.15E-02	4.81E-04
1GLF	1.28E+04	5.09E+01	8.49E-01	1.41E-02	5.89E-04
1A6J	1.54E+04	6.12E+01	1.02E+00	1.70E-02	7.09E-04
1DXE	1.81E+04	7.18E+01	1.20E+00	2.00E-02	8.31E-04
2HD3	1.93E+04	7.65E+01	1.28E+00	2.13E-02	8.86E-04
4IM7	2.46E+04	9.77E+01	1.63E+00	2.71E-02	1.13E-03
1XRU	3.38E+04	1.34E+02	2.23E+00	3.72E-02	1.55E-03

 Table S5 cont.

PDB ID	Simulated τ_F, ns	Estimated experimental τ_F, s	Estimated experimental τ_F, min	Estimated experimental τ_F, h	Estimated experimental τ_F, d
1I6O	4.03E+04	1.60E+02	2.66E+00	4.44E-02	1.85E-03
1EF9	4.66E+04	1.85E+02	3.08E+00	5.13E-02	2.14E-03
1ORO	4.82E+04	1.91E+02	3.18E+00	5.31E-02	2.21E-03
1SV6	4.94E+04	1.96E+02	3.26E+00	5.44E-02	2.27E-03
1AH9	6.57E+04	2.61E+02	4.34E+00	7.24E-02	3.02E-03
3GN5	7.30E+04	2.90E+02	4.83E+00	8.04E-02	3.35E-03
1ZYL	8.02E+04	3.18E+02	5.30E+00	8.84E-02	3.68E-03
2PTQ	5.28E+05	2.09E+03	3.49E+01	5.82E-01	2.42E-02
2V81	5.58E+05	2.22E+03	3.69E+01	6.15E-01	2.56E-02
1FTS	4.60E+06	1.83E+04	3.04E+02	5.07E+00	2.11E-01
3M7M	7.33E+13	2.91E+11	4.85E+09	8.08E+07	3.36E+06
4IWX	7.61E+13	3.02E+11	5.03E+09	8.39E+07	3.50E+06
1KSF	2.59E+14	1.03E+12	1.71E+10	2.85E+08	1.19E+07
3BRQ	3.83E+18	1.52E+16	2.53E+14	4.22E+12	1.76E+11
2FYM	1.28E+19	5.06E+16	8.44E+14	1.41E+13	5.86E+11
1CLI	2.35E+19	9.31E+16	1.55E+15	2.59E+13	1.08E+12
2O1C	3.30E+19	1.31E+17	2.18E+15	3.64E+13	1.52E+12
1NG9	1.35E+20	5.37E+17	8.94E+15	1.49E+14	6.21E+12
1U60	1.65E+20	6.53E+17	1.09E+16	1.81E+14	7.56E+12
1K7J	6.72E+20	2.67E+18	4.45E+16	7.41E+14	3.09E+13
1W78	7.66E+20	3.04E+18	5.07E+16	8.45E+14	3.52E+13
2HNH	7.85E+20	3.11E+18	5.19E+16	8.65E+14	3.60E+13
1U0B	2.82E+21	1.12E+19	1.87E+17	3.11E+15	1.30E+14
1UUF	7.80E+21	3.09E+19	5.15E+17	8.59E+15	3.58E+14
2ID0	4.21E+22	1.67E+20	2.78E+18	4.64E+16	1.93E+15
1QF6	7.43E+22	2.95E+20	4.91E+18	8.19E+16	3.41E+15
2HG2	2.83E+23	1.12E+21	1.87E+19	3.12E+17	1.30E+16
2QCU	1.66E+24	6.60E+21	1.10E+20	1.83E+18	7.64E+16
2WIU	4.80E+27	1.91E+25	3.18E+23	5.29E+21	2.20E+20
1GER	3.73E+28	1.48E+26	2.47E+24	4.11E+22	1.71E+21
1YQQ	2.34E+29	9.30E+26	1.55E+25	2.58E+23	1.08E+22

Table S6. Proteins with at least one misfolded trajectory expected to bypass chaperones

Category	Number of unique proteins	PDB IDs
Not expected to interact with TF	70	2ID0, 4IWX, 1A69, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1ZYL, 3BRQ, 1XRU, 1EF9, 2PTQ, 2FYM, 1NG9, 1DXE, 2JO6, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 1GLF, 1GER, 2H1F, 1YQQ, 1DFU, 1SVT, 2V81, 1NAQ, 1U0B, 1PF5, 1FJJ, 2YVA, 1DUV, 1UUF, 3BMB, 1A6J, 1SV6, 1W78, 4IM7, 2KFW, 1Q5X, 1P7L, 1QF6, 1T4B, 4FZW, 1AH9, 1FTS, 2QCU, 1CLI, 2PTH, 2O1C, 2HNA, 1W8G, 3GN5, 1WOC, 4DCM, 1A82, 2HG2, 1L6W, 1D2F, 1I6O, 1M3U
Not expected to interact with GroEL/GroES	80	2ID0, 1XVI, 4IWX, 1A69, 1FUI, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1P91, 1ZYL, 3BRQ, 1XRU, 1EF9, 2PTQ, 2FYM, 1NG9, 1DXE, 4HR7, 2JO6, 2HG2, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 1GQT, 2WW4, 3M7M, 3PCO, 1GLF, 1GER, 2H1F, 1YQQ, 1DFU, 1SVT, 2V81, 1NAQ, 1U0B, 1PF5, 1FJJ, 2YVA, 1DUV, 1UUF, 3BMB, 1A6J, 1H16, 1SV6, 1W78, 4IM7, 2KFW, 1Q5X, 1P7L, 1QF6, 2PTH, 3GN5, 4FZW, 1AH9, 1FTS, 2QCU, 1GZ0, 1CLI, 2O1C, 2HNA, 4E8B, 1W8G, 1WOC, 4DCM, 1A82, 1L6W, 1T4B, 1D2F, 1I6O, 4KN7, 1GYT, 1M3U
Not expected to interact with DnaK	74	2ID0, 1XVI, 1A69, 1FUI, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1P91, 1ZYL, 3BRQ, 1XRU, 1EF9, 2PTQ, 2FYM, 4HR7, 1NG9, 1DXE, 2JO6, 2HG2, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 2H1F, 1YQQ, 1DFU, 1SVT, 2V81, 1NAQ, 1U0B, 1PF5, 1FJJ, 1DUV, 1UUF, 3BMB, 1H16, 1SV6, 1W78, 4IM7, 1Q5X, 1P7L, 1QF6, 2PTH, 3GN5, 4FZW, 1AH9, 2QCU, 1GZ0, 2O1C, 1A6J, 4E8B, 1W8G, 1FTS, 1GYT, 1WOC, 4DCM, 1A82, 2HNA, 1L6W, 1T4B, 1D2F, 1GER, 4KN7, 1CLI, 1M3U
Not expected to interact with TF, GroEL/GroES, or DnaK	64	2ID0, 1A69, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1ZYL, 3BRQ, 1XRU, 1EF9, 2PTQ, 2FYM, 1NG9, 1DXE, 2JO6, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 2H1F, 1YQQ, 1DFU, 1SVT, 2V81, 1NAQ, 1U0B, 1PF5, 1FJJ, 1DUV, 1UUF, 3BMB, 1SV6, 1W78, 4IM7, 1Q5X, 1P7L, 1QF6, 4FZW, 1AH9, 2QCU, 2PTH, 2O1C, 1A6J, 1W8G, 1FTS, 3GN5, 1WOC, 4DCM, 1A82, 2HNA, 1L6W, 1T4B, 1D2F, 1GER, 1CLI, 1M3U

2001 **Table S7**. Proteins with at least one misfolded trajectory expected not to aggregate or
2002 be degraded

Category	Number of unique proteins	PDB IDs
Not expected to aggregate	68	2ID0, 1XVI, 1A69, 1FUI, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1P91, 1ZYL, 3BRQ, 1XRU, 2PTQ, 2FYM, 4HR7, 1NG9, 2JO6, 2HG2, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 1DFU, 1SVT, 1U0B, 1PF5, 1FJJ, 1DUV, 1SV6, 1W78, 4IM7, 1YQQ, 2KFW, 1Q5X, 1P7L, 1QF6, 2PTH, 3GN5, 4FZW, 1AH9, 1FTS, 2QCU, 1GZ0, 1CLI, 2O1C, 2HNA, 1A6J, 1W8G, 1WOC, 4DCM, 1A82, 1L6W, 1T4B, 1D2F, 1NAQ, 1I6O, 1GER, 4KN7, 1GYT, 1M3U
Not expected to be degraded	70	2ID0, 1XVI, 1A69, 1FUI, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1P91, 1ZYL, 3BRQ, 1XRU, 2PTQ, 2FYM, 1NG9, 4HR7, 2JO6, 2HG2, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 1GER, 1YQQ, 1DFU, 1SVT, 1U0B, 1PF5, 1FJJ, 1DUV, 1SV6, 1W78, 4IM7, 2KFW, 1Q5X, 1NAQ, 1P7L, 1QF6, 2PTH, 3GN5, 4FZW, 1AH9, 1FTS, 2QCU, 1GZ0, 1CLI, 2O1C, 1A6J, 4E8B, 1W8G, 1WOC, 4DCM, 1A82, 2HNA, 1L6W, 1T4B, 1UUF, 1D2F, 1I6O, 4KN7, 1GYT, 1M3U

Table S8. Proteins with misfolded conformations expected to bypass proteostasis machinery and remain soluble but less functional

Category	Number of unique proteins	PDB IDs
Expected to bypass chaperones, not aggregate, and not be degraded	57	2ID0, 1A69, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 1KSF, 1ORO, 1ZYL, 3BRQ, 1XRU, 2PTQ, 2FYM, 1NG9, 2JO6, 4A2C, 1PMO, 2KX9, 2R5N, 1K7J, 1GT7, 2WW4, 3M7M, 3PCO, 1DFU, 1SVT, 1U0B, 1PF5, 1FJJ, 1DUV, 1SV6, 1W78, 4IM7, 1YQQ, 1Q5X, 1P7L, 1QF6, 4FZW, 1AH9, 2QCU, 2PTH, 2O1C, 1A6J, 1W8G, 1FTS, 3GN5, 1WOC, 4DCM, 1A82, 2HNA, 1L6W, 1T4B, 1D2F, 1GER, 1CLI, 1M3U
Expected to exhibit reduced function	69	1XVI, 4IWX, 1A69, 1FUI, 2WIU, 1U60, 4DZD, 2HNH, 3HWO, 2ID0, 1ZYL, 1XRU, 1P91, 1EF9, 2FYM, 4HR7, 1NG9, 1DXE, 2HG2, 1PMO, 2R5N, 1K7J, 1GQT, 3PCO, 1GLF, 2H1F, 1YQQ, 1DFU, 1SVT, 2V81, 1NAQ, 1U0B, 2YVA, 4KN7, 1DUV, 1UUF, 3BMB, 1A6J, 1H16, 1SV6, 1W78, 4IM7, 1Q5X, 2PTH, 1T4B, 4FZW, 1FTS, 2QCU, 1P7L, 1GZ0, 1KSF, 1CLI, 1GYT, 1WOC, 2PTQ, 1A82, 2WW4, 4A2C, 3GN5, 1L6W, 1GT7, 1QF6, 1D2F, 1I6O, 1GER, 2KX9, 2O1C, 1PF5, 3M7M
Expected to bypass proteostasis machinery and to exhibit reduced function	38	1A69, 2WIU, 1U60, 4DZD, 3HWO, 1ZYL, 2FYM, 1NG9, 1PMO, 2R5N, 1K7J, 1DFU, 1SVT, 1U0B, 1SV6, 1W78, 4IM7, 1YQQ, 1Q5X, 4FZW, 2QCU, 1KSF, 2PTH, 1FTS, 2HNH, 1WOC, 2PTQ, 1A82, 3GN5, 3PCO, 1L6W, 1T4B, 1A6J, 1GT7, 1D2F, 1GER, 1DUV, 2O1C

Table S9. Percent of native state simulation frames with metric ≥10% averaged over set of 122 *E. coli* proteins

Metric	Mean [95% CIs]*, %	Range (low-high), %
$\zeta_{\text{hydrophobic}}(t)$,	6.32 [5.44, 7.27]	0.08 – 32.31
$\zeta_{\text{DnaK}}(t)$	14.74 [13.03, 16.47]	0.17 – 38.35
$\zeta_{\text{agg}}(t)$	11.74 [10.32, 13.22]	0.60 – 39.75
χ_{func}	24.58 [22.97, 26.28]	9.52 – 55.47

*confidence intervals calculated from bootstrapping 10^6 times

2101 **Table S10**. Functional residues database terms and descriptions

Term in database	Description
PDB used in model	Uniprot: PDB ID used in the simulation
Gene name	Uniprot: Name of gene
Protein Uniprot ID	Uniprot: ID associated with gene in uniprot
Ordered Locus	Uniprot: Ordered locus associated with gene
Function	Uniprot: Function of the gene product
Catalytic activity	Uniprot: Catalytic activity of protein
Subunit	Uniprot: Information about n-mers structure
GO functions	Uniprot: Gene ontology functions
Binding site	Uniprot: Information about mapped and unmapped binding site
Active site	Uniprot: Information about mapped and unmapped active site
Metal binding site	Uniprot: Information about mapped and unmapped metal binding site
Site	Uniprot: Information about mapped and unmapped site
Absorption	Uniprot: Information about ligands absorbed
Calcium binding	Uniprot: Information about calcium binding
DNA binding	Uniprot: Information about DNA binding
EC number	Uniprot: Enzyme Commission number
Metal binding	Uniprot: Information about mapped and unmapped metal binding
Nucleotide binding	Uniprot: Information about mapped and unmapped about nucleotide binding
Pathway	Uniprot: Information about mechanism pathway
pH dependence	Uniprot: Information about the protein function's pH dependence
Temperature dependence	Uniprot: Information about the protein function's temperature dependence
Cofactor	Uniprot: Information about the protein's cofactors

 Table S10 continued

Term in database	Description
Kinetics	Uniprot: Information about the kinetics of the protein
mRNA sequence	mRNA sequence of the gene product
AA sequence	Amino acid sequence of the gene product
CIF sequence of # - X	PDB: Amino acid sequence of the chain X in CIF #
Unmapped active residues from CIF #	PDB: Chain ID, Residue # in CIF file, Residue Name, Ligand in contact with
Mapped active residues from CIF #	PDB: Chain ID, Mapped residue #, Residue Name, Ligand in contact with
Ligand abbreviation from CIF #	PDB: 3 letter abbreviation, ligand name
Unmapped interfacial residues from CIF #	PDB: Chain 1, residue name, unmapped residue # in CIF file, Chain 2, residue name, unmapped residue # in CIF file
Mapped interfacial residues from CIF #	PDB: Chain 1, residue name, mapped residue, Chain 2, residue name, unmapped residue # in CIF file
Small molecule active residues (mapped)	PDB: List of residues in contact with small molecules and ligands
Interfacial active residues (mapped)	PDB: List of interfacial residues in the crystal structure
Combined active residues (mapped)	Uniprot and PDB: List of functional residues from Uniprot, residues interacting with small molecules and interfacial residues

2108

2128 **Table S11**. Values of $\langle \zeta_{\text{peptide}} \rangle$ and locations of principle entanglements by
2129 metastable state
2130

Markov State	$\langle \zeta_{\text{peptide}} \rangle$, %			Principle entanglement*		
	[333-354]	**[F351]**	**[L293]**	**Loop segment**	**Threading segment**	**Overlapping with non-refoldable region?**
S1	1.56 [0.83, 2.30]	108.38 [84.49, 133.75]	50.45 [-12.64, 123.61]	**44-314**	**318-337**	Yes
S2	86.53 [84.74, 88.30]	6446.30 [6266.09, 6626.00]	28.83 [-27.88, 99.59]	271-288	218-237	No
S3	1.42 [1.11, 1.74]	51.22 [42.10, 60.67]	-12.30 [-42.09, 22.20]	-	-	-
S4	14.66 [13.73, 15.60]	1802.97 [1707.65, 1900.01]	358.21 [151.55, 608.55]	90-138	**318-337**	Yes
S5	22.42 [21.41, 23.43]	1644.79 [1567.43, 1723.38]	58.64 [-8.93, 154.10]	271-288	218-237	No
S6	1.03 [0.38, 1.69]	358.65 [312.01, 407.32]	87.72 [-2.56, 212.08]	41-207	**318-337**	Yes
S7	6.55 [6.41, 6.69]	506.27 [496.06, 516.54]	155.52 [125.11, 188.00]	**44-313**	**317-336**	Yes
S8	3.85 [3.61, 4.10]	302.66 [290.37, 315.22]	66.70 [25.23, 115.01]	**44-313**	**317-336**	Yes

2131 *principle entanglement refers to the entanglement present in a representative structure for each metastable
2132 state. Segments that overlap with [333-354], [F351], or [L293] are highlighted in bold.