Mutualists construct the ecological conditions that trigger the transition from parasitism

Léo Ledru[1], Jimmy Garnier[2], Matthias Rohr[1], Camille Noûs[1,2,3], Sébastien Ibanez[1]

[1]Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, 38000 Grenoble, France
[2]CNRS, Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, LAMA, 73000 Chambery, France
[3]Laboratory Cogitamus

Address for correspondence: ledru.leo@hotmail.fr

Abstract

The evolution of mutualism between hosts and initially parasitic symbionts represents a major transition in evolution. Although vertical transmission of symbionts during host reproduction and partner control both favour the stability of mutualism, these mechanisms require specifically evolved features that may be absent in the first place. Therefore, the first steps of the transition from parasitism to mutualism may suffer from the cost of mutualism at the organismic level. We hypothesize that spatial structure can lead to the formation of higher selection levels favouring mutualism. This resembles the evolution of altruism, with the additional requirement that the offspring of mutualistic hosts and symbionts must co-occur often enough. Using a spatially explicit agent-based model we demonstrate that, starting from a parasitic system with global dispersal, the joint evolution of mutualistic effort and local dispersal of hosts and symbionts leads to a stable coexistence between parasites and mutualists. The evolution of local dispersal mimics vertical transmission and triggers the formation of mutualistic clusters, counteracting the organismic selection level of parasites that maintain global dispersal. The transition occurs when mutualistic symbionts increase the density of hosts, which strengthens competition between hosts and disfavours hosts inhabiting areas dominated by parasitic symbionts: mutualists construct the ecological conditions that allow their own spread. Therefore, the transition to mutualism may come from an eco-evolutionary feedback loop involving spatially structured population dynamics.

Keywords: mutualism, major transition, spatial structure, dispersal evolution, eco-evolutionary feedbacks, niche construction

Introduction

In their bestseller, Szathmáry and Smith (1995) explored several major transitions in biology, from the origin of chromosomes to the evolution of social groups. Most of these transitions resulted from the formation of a larger entity from smaller entities, smaller entities specializing within the larger ones. Following Queller (1997), Szathmáry (2015) suggested a dichotomy between fraternal transitions arising from a division of labour among closely related units (such as multicellularity) and egalitarian transitions, where phylogenetically distant units come together to complement their functions in a larger unit. Egalitarian transitions are generally achieved through mutualistic symbiosis[1] between a relatively large host and its symbiont (Bronstein, 2015; Drew et al., 2021) and constitute one of the main sources of new lineages, underlying the origin of the eucaryotic cell and photosynthetic eucaryotes for instance (Margulis and Sagan, 2002). In many cases symbionts are unicellular microbes which are hosted by large eucaryotes, the whole corresponding to a holobiont (Gilbert et al., 2012; Bordenstein and Theis, 2015); in other cases symbionts are multicellular organisms physically associated with their host at various degrees (e.g. plant-fungi, plant-ant, plant-seed eating pollinator). While symbionts depend on their host from the start, hosts often become dependent on the symbionts during later stages (Roughgarden, 1975), e.g. for reproduction or resource acquisition, eventually making the transition irreversible.

For a transition to occur and persist, evolutionary conflicts between the subentities must not overtake the whole's fate. In the case of fraternal transitions, this is prevented by the strong relatedness between subentities (Hamilton, 1964b,a; Queller, 2000; Fisher et al., 2013). However, in the case of egalitarian transitions, the subentities generally belong to different species. Thus, it can be advantageous for them to remain autonomous and exploit the other subentities. This parasitic behaviour occurs at the expense of the whole, as for the tragedy

---

Author contributions: SI, JG and LL originally formulated the project; all authors developed the model; LL and MR performed the numerical analyses; LL, SI and JG wrote the manuscript.

[1]Symbiosis is used here in its etymological sense of "living together", encompassing parasitic and mutualistic symbiosis.

of the commons (Garrett, 1968; Hardin, 1998). For instance, a symbiont may remain parasitic rather than collaborate with its host (Drew et al., 2021). The resulting evolutionary conflict might be circumvented by vertical transmission of the symbionts, which ensures that all subentities share a common fate (Wilson and Sober, 1994). As a result, vertical transmission of symbionts indeed promotes the transition to mutualism (Smith, 1998; Herre et al., 1999; Wilkinson and Sherratt, 2001; Ferdy and Godelle, 2005; Kerr and Nahum, 2011; Akçay, 2015; Estrela et al., 2016; Queller and Strassmann, 2016; Doebeli and Knowlton, 1998), although symbionts vertically transmitted can persist without becoming mutualists (Saikkonen et al., 2002).

The importance of vertical transmission has been highlighted by experiments on microbial systems (Sachs et al., 2011; Shapiro et al., 2016; King et al., 2016; Shapiro and Turner, 2018) as well as *in natura* observations of a Wolbachia-insect system (Weeks et al., 2007). However, in many mutualistic systems, the symbiont is transmitted horizontally (Wilkinson and Sherratt, 2001), such as legume-rhizobium (Denison and Kiers, 2004), squid-vibrio (McFall-Ngai, 2014), mycorrhizae (Allen, 1991), endophytes (Saikkonen et al., 2004) or plant-ants (Bronstein et al., 2006; Rico-Gray and Oliveira, 2008). In such cases, several mechanisms such as partner choice, sanction or fidelity can counteract the selection for selfishness (Genkai-Kato and Yamamura, 1999; Wilkinson and Sherratt, 2001; Sachs et al., 2004; Foster and Wenseleers, 2006; Estrela et al., 2016; Akçay, 2017; Sachs et al., 2010b). For instance, in legume-rhizobium, mycorrhizal and plant-ant associations, the plants can sanction the less beneficial (or even detrimental) symbionts by allocating them fewer resources (West et al., 2002; Kiers et al., 2003; Denison and Kiers, 2004; Edwards et al., 2006; Bever et al., 2009; Akçay, 2015). However, it is unclear whether these mechanisms are present at the beginning of the transition to mutualism. Since they require the evolution of complex and specific traits, they may occur in later stages, providing additional stability to the system. In the absence of such traits, what mechanism could promote the transition in the first place? Using a theoretical model, the present work aims to show that the joint evolution between mutualistic effort and local dispersal of hosts and symbionts leads to a positive association between mutualistic hosts and symbionts and subsequently triggers the formation of mutualistic clusters .

A similar issue exists with respect to the evolution of altruism[2], since partner choice and control mechanisms, such as voluntary reciprocal altruism (Axelrod, 1981), may be restricted to higher animals or may appear during later evolutionary stages. In line with the intuition of Darwin (1871), spatial structure has been recognized as a general mechanism promoting the transition to altruism (Mitteldorf and Wilson, 2000; Lion and Van Baalen, 2007, 2008; Débarre et al., 2012). Spatial structure generally stems from local dispersal, which triggers the formation of clusters dominated by altruistic organisms, while organisms with similar phenotypes are positively assorted in space (Wilson and Dugatkin, 1997; Pepper, 2007). The balance between organismic-level selection favouring cheaters and cluster-level selection favouring altruists ultimately determines the evolutionary outcome (Van Baalen and Rand, 1998; Mitteldorf and Wilson, 2000). Moreover, the joint evolution of cooperation and dispersal can allow the emergence of altruism, with spatial clusters of altruistic organisms promoting the persistence and spread of altruistic phenotypes (Koella, 2000; Le Galliard et al., 2005; Hochberg et al., 2008; Purcell et al., 2012; Mullon et al., 2018). Empirical evidence on the evolution of reduced virulence (Boots and Mealor, 2007; Szilágyi et al., 2009), the evolution of altruism (Harcombe, 2010), and the evolution of restraint predation (Kerr et al., 2006) also supports the crucial role of the spatial structure.

Similarly, spatial structure can allow mutualists to overcome non-mutualists (Yamamura et al., 2004; Akçay, 2017; Doebeli and Knowlton, 1998; Frank, 1994), and this can come along with the evolution of dispersal (Mack, 2012). However, this may not be sufficient to account for the transition from parasitism to mutualism, since parasitic symbionts should discourage hosts from initiating the transition, whereas non-mutualists have a weaker impact (Yamamura et al., 2004; Mack, 2012). In the case of holobionts, starting from free living bacteria, Sachs et al. (2011) documented 27 transitions towards parasitism, 9 directly towards mutualism and 3 towards commensalism, whereas the transition from parasitism to mutualism occurred only 3 times. This highlights that the transition from parasitism to mutualism, although feasible, is relatively infrequent, and calls for a theoretical understanding of the mechanisms involved. Moreover, in previous attempts mutualistic efforts were initially polymorphic but were not subject to mutations (Mack, 2012). In that case, mutualistic clusters cannot be invaded from inside through parasitic mutations, which favours mutualism. The present work therefore constitutes, to our knowledge, the first spatially explicit eco-evolutionary model where the mutualistic efforts and dispersal abilities of hosts and symbionts coevolve, beginning from a parasitic interaction. If some hosts and symbionts simultaneously become mutualists and start dispersing locally, this may lead to the formation of mutualistic host-symbiont clusters producing more offspring than in areas where hosts are mainly associated with parasitic symbionts, thereby initiating the transition. Meanwhile, parasitic symbionts should continue dispersing globally and invade the mutualistic clusters, which could homogenize the spatial structure and compromise the transition. Also, densely populated mutualistic clusters might suffer from intraspecific competition between hosts, unless competition acts on a large spatial scale. In sum, it is unclear whether mutualists will invade, whether mutualists will replace parasites, or whether both strategies will coexist, as is often the case in nature (e.g. Després and Jaeger, 1999; Borges, 2015; Saikkonen et al., 2004).

---

[2]An altruistic trait benefits conspecifics, at a cost to its bearer. In contrast, a mutualistic trait benefits heterospecifics.

The concept of major transitions also implies that the host and the symbiont become dependent upon each other (Szathmáry and Smith, 1995; Szathmáry, 2015), with each partner needing the other to perform essential functions like nutrient provisioning (Fisher et al., 2017). Dependence is often accompanied by gene loss and gene exchange, rendering the transition irreversible (Estrela et al., 2016). Most symbionts cannot live freely and therefore completely depend on their host, but most hosts can complete their life cycle without their symbiont (e.g., in plant-ant, plant-fungi or legume-rhizobium mutualisms) and several reverse pathways are possible from mutualism to parasitism (Sachs and Simms, 2006; Werner et al., 2018; Week and Nuismer, 2021). However in some cases hosts depend on their symbiont, for instance the intracellular bacterial symbiont *Buchnera aphidicola* provides essential amino acids to its aphid host (Akman Gündüz and Douglas, 2009; Bennett and Moran, 2015). Since the present work focuses on the transition and not on later stages, we will not assume that hosts depend on their symbionts for their physiology or development, which would render the transition irreversible *by construction*. Instead, hosts will always be able to produce offspring when alone. Nevertheless, the number of offspring produced by the hosts will depend on the mutualistic efforts of both species as well as on the population densities, which are expected to change during the transition. Under these altered ecological conditions, isolated hosts may exhibit a negative population growth rate, although they are physiologically able to produce offspring. This mechanism is hereafter called *ecological dependence*.

To sum up, we will tackle the following issues:

- **Main hypothesis:** In the absence of vertical transmission and partner control, we expect that the transition from parasitism to mutualism can occur when the mutualistic efforts of both hosts and symbionts jointly evolve with local dispersal.

- **H1:** The formation of mutualistic clusters should be necessary for the initiation of the transition. The emergence of spatial structure should come along with the transition.

- **H2:** By maintaining global dispersal, non-mutualistic hosts and parasitic symbionts should be able to coexist with mutualists.

- **H3:** The transition to mutualism is due to the relatively higher fecundity of mutualistic clusters.

- **H4:** If competition between hosts is mostly local, this should hamper the formation of mutualistic clusters, thereby preventing the transition.

- **H5:** We expect that mutualistic hosts will become ecologically dependent on their symbiont.

To investigate these hypothesis, we built an agent-based model using a two-dimensional space lattice that supports an autonomous host and a host-dependent symbiont. Hosts compete for space and other resources, while symbionts compete for available hosts. This situation occurs in many biological systems, such as plant-fungi, plant-seed eating pollinator, plant-ant, and multicellular eucaryotes hosting bacteria. Less intimate associations like cleaning mutualisms or plant-pollinator interactions may also fit, provided that the animal is specialized and dependent on its host. To model the transition from parasitism, the symbiont is initially detrimental to the host, and the host provides it the minimal energy possible without any spontaneous mutualistic effort, as would be the case after an antagonistic evolutionary arms race. Moreover, the host-parasite system is ecologically viable even in the absence of any mutualistic agent in the landscape. At first, both species disperse globally; this situation corresponds to the most disadvantageous conditions for the emergence of mutualism. Through continuous mutations, mutualistic and locally dispersing symbionts and hosts can appear. The mutualistic effort encompass the provision of resources, shelter, immunity, anti-predator behaviours, digestive enzymes or any other type of benefit provided that this occurs at some cost. If mutualistic symbionts manage to persist for a while, they eventually change the population dynamics, triggering feedback on their own evolutionary dynamics. In addition to these general hypotheses, no assumptions specific to a particular biological system were required.

Model description

Main rules Our model considers two types of agents, hosts and symbionts, living on the same two-dimensional space lattice. The interaction between the two species occurs when they share the same cell. Each cell can assume three states: i) empty, ii) occupied by a solitary host, with only one host per cell), iii) occupied by a host-symbiont couple, with only one symbiont per host (but see Appendix A.7 for a relaxation of this assumption). Each organism bears two traits, an interaction trait α and a dispersal trait ε, which both influence fecundity. At every time point, agents undergo the following steps (see appendix A.1 and Figure A1 for more details):

- The host and symbiont die with fixed probability m.
- They produce offspring, possibly with mutations. The average offspring number of a parent depends on its traits and on its interactions with their cell-sharing partner, if any.
- The offspring are dispersed according to the parental trait ε.
- The host offspring may establish only in empty cells, while the symbiont offspring can only establish in cells already occupied by a solitary host. If several organisms come to implant in the same cell, a uniform lottery determines which one will implant, while the others die.

Fecundity and mutualism/parasitism Each agent produces offspring according to a Poisson distribution with parameter f, which corresponds to its fecundity. The fecundity defines the average number of offspring per agent. It results from an interaction fecundity positively dependent on the trait of its cell-sharing partner and a mutualistic cost negatively dependent on its trait.

Specifically, the fecundity of a symbiont f^s of trait α_s in interaction with a host of trait α_h is defined by:

$$f^s(\alpha_s, \alpha_h) = (1 - c_m \alpha_s)\big(f^s_{min} + (f_{max} - f^s_{min})\alpha_h\big) \tag{1}$$

where c_m is the maximal mutualistic cost and f^s_{min} and f_{max} are the minimal and maximal interaction fecundity of symbionts. Similarly, the fecundity of a host f^h of trait α_h in interaction with a symbiont of trait α_s is defined by:

$$f^h(\alpha_h, \alpha_s) = (1 - c_m \alpha_h)\big(f^h_{min} + (f_{max} - f^h_{min})\alpha_s^{\gamma_f}\big) \tag{2}$$

where f^h_{min} is the minimal interaction fecundity of hosts and γ_f describes the selection strength on the trait α_s – the selection strength on α_h is set to $\gamma_s = 1$.

Since hosts are autonomous, in absence of symbionts, their fecundity f^{ha} only depends on their trait α_h:

$$f^{ha}(\alpha_h) = (1 - c_m \alpha_h)f^a \tag{3}$$

where the fecundity alone f^a ranges between the minimal and maximal interaction fecundity: $f^h_{min} < f^a < f_{max}$. As a result, the establishment of a symbiont with a low interaction trait ($\alpha_s < \alpha_s^*$) reduces the fecundity of the host; the symbiont is parasitic. Instead, a symbiont with a large interaction trait ($\alpha_s > \alpha_s^*$) enhances the host's fecundity; the symbiont is mutualistic. The threshold α_s^* is defined by $f^h(\alpha_h, \alpha_s^*) = f^{ha}(\alpha_h)$ (see appendix A.1 for mathematical derivation of the threshold). In the simulations, $\alpha_s^* = 0.475$ (Figure 1).

Figure 1: Fecundity of hosts f^h and symbionts f^s according to the interaction trait of their partners (dashed black lines). Plain black line corresponds to the fecundity of a solitary host f^a. The dashed red line corresponds to the threshold $\alpha_s^* = 0.475$ separates symbionts, which either reduce or enhance the fecundity or their host (appendix A.1).

Mutation Offspring inherit traits from their parents with variability due to mutations. The effects of mutations on each trait are independent . However, the distribution of mutation effects does depend on the trait of the parents. We use a Beta distribution with shape parameters $(1, 3)$ to describe the amplitude of these effects, which could be either beneficial or detrimental. This mutation kernel allows for rare mutations with large effects. However, these effects can not exceed a maximal mutation size set to $\beta_{max} = 0.5$ (see Figure A3 in appendix A.1 for details).

Dispersal The parents do not disperse, while their descendants disperse either locally in one of the 8 cells around the parent or globally, with a uniform distribution across the entire space (see Figure A5 for a sketch of the process). The dispersal trait ε is defined as the proportion of offspring dispersed globally, as in Kéfi et al. (2007, 2008). These two modes of dispersal correspond to a mixture of short and long distance dispersal

events. For instance, fleshy fruits may be dispersed either by small birds having a short-distance behaviour, or by mammals and large birds which disperse the seeds at long distances (Jordano et al., 2007). Fruits may also remain unconsumed and fall locally. Depending on the fruit's traits, its propensity to be consumed by either type of frugivores may vary among organisms, which is captured by the dispersal trait ε. Since the investment in global dispersal may reduce fecundity (Harada, 1999; Bonte et al., 2012), we assumed a linear trade-off between fecundity and dispersal: $f_e = (1 - d\,\varepsilon)\,f$, with f_e the effective fecundity and d the dispersal cost intensity, which is the same for both hosts and symbionts.

Competition Hosts compete for empty cells, especially if they disperse locally. Beside space, hosts may also compete with each other for resources like water, light or food. In order to test hypothesis H3 we introduced intraspecific density-dependent competition, acting either at the local or the global scale. For instance, competition for light only involves the closest neighbors while competition for the water table might act at the entire space scale. The competition scale parameter w_h, ranging in $[0, 1]$, weights the effect of the local density ρ_h^{local} and the global density ρ_h^{global} of host on the competition. Competition reduces the establishment probability P_I of the offspring:

$$P_I = 1 - \left((1 - w_h)\rho_h^{local} + w_h\rho_h^{global} \right)^{\gamma_C} \tag{4}$$

The local host density ρ_h^{local} corresponds to the host density in the 8 neighbouring cells surrounding the offspring, while the global density ρ_h^{global} corresponds to the host density over the entire landscape (see Figure A2 for a schematic representation). The parameter γ_C corresponds to the competition strength. The competition is strong when $\gamma_C < 1$ (sub-linear function), while it is weak when $\gamma_C \geq 1$ (super-linear function).

Parameters		Reference values	Sensitivity analysis range
m	probability of mortality	0.06	$[0.005; 0.15]$
c_m	maximum mutualistic cost	0.3	$[0; 1]$
f_{max}	maximal host and symbiont interaction fecundity	8	fixed
f_{min}^h	minimal host interaction fecundity	0.1	fixed
f_{min}^s	minimal symbiont interaction fecundity	2.5	fixed
f^a	maximal solitary host fecundity	0.5	fixed
γ_f	selection strength on the symbiont interaction trait	4	fixed
β_{max}	maximum mutation size	0.5	$[0.1; 1]$
w_h	scale of host competition	1	0 or 1
γ_c	strength of host competition	0.2	$[0.1; 2]$
d	dispersal cost	0	$[0; 1]$

Table 1: List of parameters and their reference values used for the simulations. The parameters of host and symbiont fecundities are determined to ensure the viability of the antagonistic system, therefore they are fixed because they are constitutive of the model.

Parasitic system and transition To tackle the issue of transition to mutualism, we assume that the system is viable without mutualism (see appendix A.2 for details). More precisely, in the absence of mutation, the extinction probability of a population with parasitic global dispersers without dispersal cost (minimal interaction traits, $\alpha_s = \alpha_h = 0$, dispersal trait $\varepsilon = 1$ and dispersal cost $d = 0$) is given by $m/((1 - m)f^a)$ where m is the mortality rate and f^a the fecundity of a parasitic host without symbiont.. We choose parameters such that the extinction probability is less than 1. In this case, the population stabilizes around a demographic equilibrium called the "parasitic system" where host density is around 0.15 and the symbiont density is around 0.1 with parameters value set in Table 1 (see Figure 2b-c and appendix A.2). From our perspective, this situation is the worst-case scenario because interactions are parasitic and dispersal cost is minimal. Then, mutualistic symbionts can appear by mutation, which generates approximately 2% of mutualistic symbionts in the population (see dashed purple curve in Figure 2c and Figure A8d). Natural selection eventually leads to a significant increase of the percentage of mutualistic symbionts, far above the 2% generated by mutations (Figure 2). Using an approximation model, we show that the extinction probability of mutualistic symbionts falls below 1 when the percentage of symbiont rises above 10% (appendix A.2 for details). In the simulations, a high density of mutualistic symbionts indeed persists in the long term when the percentage of mutualistic symbionts stands above 10% (Figure A9), which therefore characterizes the transition to mutualism. The transition time was defined as the time at which the percentage of mutualistic symbionts rises above this threshold.

Assortment and aggregation indices To investigate the spatial structure, which comes along with the transition to mutualism, we compute assortment indices: intraspesific indices measuring the spatial autocorrelation among hosts and symbionts and an interspecific index quantifying the correlation between phenotypes of host and symbiont sharing the same location. More specifically, the intraspecific indices compute the similarity between the trait of an organism and the traits of its neighbors located in the 8 cells around it, and compare it with the similarity between the organismic trait and the mean trait over the landscape (details in appendix A.1). If the intraspecific index is positive (respectively negative), it means that on average the neighbors of any organism share similar (respectively dissimilar) traits. Similarly, the interspecific index is positive if hosts and symbionts sharing the same cell have similar interaction traits. Spatial aggregation indices for hosts, mutualistic symbionts and parasitic symbionts were also computed, measuring the formation of clusters (appendix A.1 for details).

Results

In the following, the maximum cost of mutualism c_m is 30%, and the other parameters are set to satisfy the viability of the parasitic system (Table 1 in appendix A.1 and appendix A.2 for a discussion of the effect of the cost of mutualism).

The transition from parasitism to mutualism Our main objective was to investigate whether the transition to mutualism is possible starting from a viable parasitic system, without dispersal cost, which constitutes the most stringent condition for the transition. In that case, the transition is more likely to occur under strong ($\gamma_C = 0.2$) intraspecific host competition (with frequency 0.95) than under weak ($\gamma_C = 1$) competition (0.086). Moreover, when the transition succeeds, it occurs more rapidly under strong competition (median transition time around $2.5.10^4$) than under weak competition (median transition time around 7.10^4, Figure 2a). When the cost of dispersal is large ($d = 0.45$) the transition occurs systematically (with frequency 1) and the median transition time is much lower (around 7.10^2), regardless of the strength of competition (Figure 2a). Dispersal cost was therefore used as an instrumental tool to speed up the transition when necessary.

The transition begins with weakly mutualistic symbionts, which rapidly increase their mutualistic effort toward 1 (Figure 2c). In contrast, the increase of the average host interaction trait is delayed in response to the symbionts' transition (Figure 2b). Moreover, the transition does not occur at the expense of parasitic symbionts; on the contrary their population density benefit from the increase in host density triggered by the mutualistic symbionts (Figure 2c).

Since the symbiont population is monomorphic at the beginning of every simulation, the two distinct phenotypic clusters visible in Figure 3a indicate that both traits diverged, resulting in two classes of symbionts: parasitic global dispersers ($\alpha_s << 1$ and $\varepsilon \sim 1$) and mutualistic local dispersers ($\alpha_s \sim 1$ and $\varepsilon << 1$). Furthermore, the mutualistic and dispersal traits of symbionts evolve at the same time, during the transition (details not shown). Conversely the host traits do not diverge; their joint evolution leads to a negative correlation between global dispersal and mutualism intensity (Figure 3b, $R^2=0.102$). After the transition, most hosts provide a non-zero mutualistic effort to the symbiont (most $\alpha_h > 0.2$).

The assortment indices indicate that after the transition to mutualism the organisms of both species are locally similar. Moreover, hosts and symbionts sharing the same location also tend to have the same interaction behaviour (Figure 4a). The intraspecific assortment is stronger than the interspecific assortment, which is not surprising since the formation of the intraspecific spatial structure simply requires a sufficient proportion of local dispersal. The aggregation indices (appendix A.1) behave similarly, after the transition the spatial aggregation of hosts, parasitic symbionts and mutualistic symbionts all increase, and the parasitic and the mutualistic symbionts reach the same level of aggregation (Figure A7). These results together indicate that the transition to mutualism comes along with the emergence of a spatial structure, with clusters of mutualistic hosts and symbionts (Figure 4c).

The effect of competition between hosts Figure 2a shows that the host competition promotes the transition to mutualism; we next investigate its quantitative effect on the percentage of mutualistic symbionts. The following results were obtained using a large dispersal cost ($d = 0.45$) to reduce the mean time of transition and thus save computational time.

The competition strength γ_C increases the percentage of mutualistic symbionts after the transition when competition is global, i.e. when hosts compete with all the hosts present in the landscape (Figure 5a). However, the transition can occur even in the absence of host competition, if the cost of mutualism is sufficiently low (e.g., a maximum cost of only 10% instead of 30% as in previous simulations, details not shown). When competition is more local the percentage of mutualistic symbionts decreases drastically, until it drops below the transition threshold (Figure 5b). In the absence of dispersal cost, when competition is reduced after the transition to mutualism, the system switches back to the parasitic state (Figure 5c, see Figure A10 for details).

Another way to investigate the effect of competition is to reduce host density, through the eradication of hosts in a region after a transition to mutualism. At first, the perturbed region is mainly recolonized by hosts

Figure 2: a) Histograms of the number of transitions over 1000 simulations as a function of time, with a maximum projection time of 10^5. Without dispersal cost there are a total of 86 transitions when the competition is weak and 951 when the competition is strong. With dispersal cost there are 1000 transitions whether the competition is weak or strong. Panels b) and c) represent the host and symbiont densities over time averaged over 100 simulations (coloured plain curves) under strong competition $\gamma_C = 0.2$ and no dispersal cost $d = 0$ and with a maximum projection time of 10^4 steps. The densities correspond to the proportion of occupied cells. The time series are adjusted so that all simulations have a transition time $t = 2000$. The colour gradient corresponds to the mean interaction trait α, and shaded regions correspond to the standard deviation for densities. In panel c), the purple dotted line and the right y-axis show the relative density of mutualistic symbionts, and the black line indicates the 10% transition threshold. For all panels, other parameters are $m = 0.06$, $c_m = 0.3$, $w_h = 1$, $f^h_{min} = 0.1$, $f^s_{min} = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$

272 and parasitic symbionts (Figure A11b), but mutualistic symbionts persist in the landscape. Due to the relaxation
273 of global competition, the probability of host establishment is better, and the mutualistic clusters outside the
274 perturbation zone gain in size, which explains why the proportion of mutualistic symbionts increases slightly de-
275 spite the recolonization of the centre by parasites (Figure A11a). In the end, once recolonization is complete, the
276 system returns to an equilibrium state whose trait distributions are close to distributions before the perturbation

Figure 3: Post transition traits distribution (ε, α) of symbionts (panel a) and hosts (panel b). The dashed red line in panel a indicates the threshold $\alpha_s^* = 0.475$ above which a symbiont benefits its host. The plain red line in panel b shows the linear regression between host traits (R^2=0.102). Distributions corresponds to 100 simulations with strong competition $\gamma_C = 0.2$, no dispersal cost $d = 0$ and with a maximum projection time of 10^4 steps. Other parameters are $m = 0.06$, $c_m = 0.3$, $w_h = 1$, $f_{min}^h = 0.1$, $f_{min}^s = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$.

(details not shown). A similar experiment with a perturbation causing the death of 50% of uniformly occupied cells leads to the same results.

Host dependency Under favourable conditions leading to the transition to mutualism, the population of mutualistic hosts always persists in the absence of mutualistic symbionts, which excludes any absolute dependency of hosts for symbionts. However, *ecological dependency* may occur, where isolated hosts may have a negative growth rate because of intraspecific competition, although they would be able to form stable populations at lower densities. The transition to mutualism co-occurs with an increase in host density and thus an increase in intraspecific competition. If this increase in competition is sustainable only in the presence of mutualistic symbionts, the hosts are *ecologically dependent* on the symbionts. In order to determine the occurrence of ecological dependency, the intensity of intraspecific competition between hosts was measured in a system at equilibrium after the transition to mutualism (e.g., at the end of Figure 2c), and subsequently used as a fixed parameter to test if the host population can now survive in the absence of mutualistic symbionts and mutation. We found that ecological dependency occurs in the hatched area of Figure 6, when the system evolves toward a mutualistic system in which the percentage of mutualistic symbionts is sufficiently large.

Figure 6 further shows that both dispersal cost and mortality promote mutualism. For the parameter pair in the area indicated by the grey star, where dispersal cost is zero, Figure 2a showed that the probability of transition during the 10^5 time steps is only 0.086, with a mean transition time of 7.10^4. This explains why no transition occurred in Figure 6, where 50 simulations per parameter combination were performed, with only 10^4 time steps. Finally, Figure 6 also shows that for some parameter combination, mutualism evolves even though the parasitic system is initially unviable. The viability of the parasitic system was assessed by simulations of 5000 time steps, without evolution. This implies that in a relatively short period of time in comparison to the transition times shown in Figure 2 for other parameter values, transitions can occur quickly enough and prevent the extinction of a parasitic system otherwise unviable. However this occurs rarely; Figure A14 shows that for some parameter combinations up to 90% of the simulations go extinct, the remaining being able to persist thanks to the evolution of mutualism. In those cases the mean percentage of mutualistic symbionts is much higher, ranging from 35 to 60%.

Discussion

The mechanisms underlying the transition to mutualism.

In line with our main hypothesis, our results indicate that the transition from parasitism to mutualism occurs when mutualistic efforts evolve together with dispersal, despite the absence of vertical transmission or partner control. The following paragraphs review the mechanisms which contribute to the transition, and related them with the hypothesis formulated earlier.

The formation of clusters. Before the transition, the host performs better when alone; therefore, it has no interest in increasing its mutualistic effort and natural selection keeps it as low as possible. In contrast the symbiont population is limited by the number of available hosts, which increases when the symbiont becomes

Figure 4: a) Spatial structures are described by the assortment index measuring the intraspecific assortment between hosts (plain line), the intraspecific assortment between symbionts (dashed line) and the interspecific assortment between hosts and symbionts (dash-dot line). Results are averaged over 100 simulations with strong competition $\gamma_C = 0.2$ and no dispersal cost $d = 0$. The time series are adjusted so that all simulations have a transition time $t = 2000$ (red dotted line). Grey areas show the standard deviation. The threshold separating mutualistic and antagonistic symbionts is as in Figures 1 and 3. b)-c) Snapshots of a region of 40x40 cells before (panel b) and after (panel c) the transition to mutualism. For the sake of the figure, a host is considered weakly mutualistic if its interaction trait is less than 0.5 and strongly mutualistic if it is greater than 0.5. Other parameters are $m = 0.06$, $c_m = 0.3$, $w_h = 1$, $f^h_{min} = 0.1$, $f^s_{min} = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$

mutualistic. Mutualistic symbionts, which help globally dispersing hosts, would be counter-selected. However, in spatially structured populations, rare mutants can interact with each other (Lion and Van Baalen, 2008), so if by chance mutations produce a mutualistic symbiont dispersing locally and interacting with a host dispersing locally as well, its offspring will benefit from the increased density of hosts in their neighbourhood and will form a mutualistic cluster (in line with hypothesis H1, Figures 4c) and A7). The cluster can then be invaded by parasitic symbionts dispersing globally, resulting in a dynamic equilibrium between mutualism and parasitism (in line with hypothesis H2, Figure 2c). Parasitic symbionts become themselves aggregated (Figure A7) since they develop around the mutualistic clusters, at their expense (Figure 4c). Joint evolution between mutualistic effort and dispersal results in a negative correlation between mutualism intensity and global dispersal (80% of mutualists disperses locally, Figure 3), which mirrors the link between altruistic behaviour and local dispersal (Koella, 2000; Le Galliard et al., 2005; Hochberg et al., 2008; Purcell et al., 2012; Mullon et al., 2018) as well as

Figure 5: The role of host competition in the transition to mutualism. Effects of competition strength γ_c (panel a) and spatial scale w_h (panel b) on the percentage of mutualistic symbionts with dispersal cost $d = 0.45$ and after 10^4 time steps and over 50 replicates. Black curves is the median and shaded regions are 95% confident intervals. Dashed red line is the transition threshold of 10%. In panel a) competition is global ($w_h = 1$) and in panel b) competition is strong $\gamma_C = 0.2$. Panel c) represents the effect of competition on the transition and on the maintenance of mutualism (simulation without dispersal cost $d = 0$). Panel d) presents the effect of a reduction in competition caused by a perturbation eradicating all organisms in a large square. The perturbation occurs around $t = 10^4$, which is 5000 time steps after the transition. Other parameters are $m = 0.06$, $c_m = 0.3$, $f_{min}^h = 0.1$, $f_{min}^s = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$.

the relationship between local interactions and avirulence evolution (Boots and Mealor, 2007).

The key role of intraspecific competition The invasion of a mutualistic cluster by parasites may cause its extinction and hinder the transition. We postulated that the higher fecundity of mutualistic clusters could compensate for their susceptibility to parasites (hypothesis H3). We instead found that, in the absence of dispersal cost, an eco-evolutionary feedback involving intraspecific competition between hosts was necessary for the transition. Indeed when competition between hosts is weak, the transition to mutualism rarely occurs (Figure 2a), and when it does, the percentage of mutualistic symbionts remains low (Figure 5a). Conversely, when hosts strongly compete for resources, the ecological conditions change dramatically. The formation of mutualistic clusters (Figure 4a) increases population densities (Figures 2b and 2c), which enhances competition between hosts. Areas dominated by hosts associated with parasitic symbionts were initially viable, but their population growth rate becomes negative following the increase in competition. This creates empty space that can be colonized by mutualists, which still disperse globally from time to time. By lowering the abundance of parasitic symbionts, this also reduces the frequency at which mutualistic clusters are invaded by parasites. The transition needs some time to occur (Figure 2a) because several obstacles must be overcome (simultaneity of the mutations, demographic stochasticity, possible invasions by parasites) before the mutualists are numerous enough to induce the shift in host competition. Therefore, contrary to hypothesis H3, the transition is not directly caused by the higher fecundity of mutualistic pairs (which would fit soft selection, Wallace, 1975) but only indirectly by the increase in host competition, which renders areas dominated by parasites unviable (hard

Figure 6: Percentage of mutualistic symbionts as a function of intrinsic mortality m and dispersal cost d. We run 50 simulations per parameter combination, with strong global competition $\gamma_C = 0.2$ and $w_h = 1$ with a maximum projection time of 10^4 steps. The percentages are averaged over the simulations leading to transition, if any occur. Above the black dotted line, the parasitic system is not viable, although the evolution of mutualism can occur above this line (evolutionary rescue). White cells correspond to the nonviability domain for the whole system, even with evolution. In the dark grey area, none of the simulations gave birth to transitions. The evolution of host ecological dependency occurs in the hatched area, where an average isolated host has a negative growth rate because of intraspecific competition. The grey star corresponds to the restrictive conditions of Figure 2. Other parameters are $c_m = 0.3$, $f_{min}^h = 0.1$, $f_{min}^s = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$.

selection). However, this only occurs when competition between hosts is partly global (Figure 5b); if it is purely local, mutualistic clusters cannot influence the viability of parasitic regions and will suffer from kin competition. In line with hypothesis H4, local competition between hosts for resources thereby prevents the transition to mutualism. Local competition between hosts for available space also occurs when hosts disperse locally, but this does not jeopardize the transition.

Empirical work has shown that the outcome of interactions between hosts and symbionts depends not only on the traits of the protagonists, but also on the surrounding ecological conditions (Bronstein, 1994). For instance, plants take advantage of seed-eating pollinators in the absence of alternative pollinators but not in their presence (Thompson and Cunningham, 2002). Mycorrhizae are beneficial for plants when soil resources are scarce while they are detrimental when resources are abundant (Johnson et al., 1997). In the above cases, the outcome of the interaction depends on both biotic and abiotic factors that are external to the host-symbiont system. Our model showed that the association with symbionts remains parasitic when host competition is low, while it evolves towards mutualism when host competition increases. In that case, the outcome of the interaction depends on intrinsic features of the interactions that are constructed by the eco-evolutionary dynamics of the system, as the emergence of mutualists increases host density.

The impact of dispersal cost and mortality As expected, dispersal cost speeds up the transition (Figure 2a and 6) because it induces a selection pressure at the organismic level in favour with local dispersal, which increases the likelihood of the formation of mutualistic clusters. Mortality also enhances the probability of transition (Figure 6), but with another mechanism. We have stressed that competition between hosts creates an eco-evolutionary feedback loop, where the evolution of mutualism increases global densities, which strengthens competition and therefore turns the growth rate of the parasitic system negative. Given that mortality pushes the parasitic system towards its viability boundary, high mortality enhances the ability of competition to launch the transition. Although the transition occurs in a wide range of parameters where the parasitic system is viable, it is more likely when the parasitic system is close to extinction (Figure 6). However, mortality cannot itself trigger the transition since the parasitic system is unviable from the start when mortality is too high. Finally, mortality may also facilitate the transition through the reduction of global densities, which decreases the threat

of parasites invading mutualistic clusters. The facilitation of mutualistic symbiosis in harsh environmental conditions has also been observed in previous empirical (Callaway et al., 2002; Maestre et al., 2003; Werner et al., 2015) and theoretical (Travis et al., 2006) works. However in the context of altruism the opposite relationship was found (Taylor and Irwin, 2000).

Evolutionary rescue As evidenced by Figure 6, the evolution of mutualism can prevent the extinction of the parasitic system for parameter combinations that are just above the upper limit of its viability domain. This echoes the concept of evolutionary rescue (Ferriere and Legendre, 2013; Gomulkiewicz and Holt, 1995), according to which the persistence time of a population is longer with than without evolution. In the present case, instead of a single population, the populations of two distinct species are rescued by evolution. More generally, the parasitic system benefits from the evolution of mutualism even when it is initially viable, through an increase in population densities (Figure 2).

The evolution of mutualistic hosts So far, only the mechanisms responsible for the evolution of mutualistic symbionts have been elucidated, but not those involved in the evolution of mutualistic hosts. Surprisingly, mutualistic hosts evolve after the transition (Figure 2c). Following the transition, the density of mutualistic symbionts is much higher, so that mutualistic hosts tend to be associated with mutualistic symbionts (Figure 4c), which disperse locally (Figure 3a). In that case, mutualistic hosts will increase the local density of mutualistic symbionts in the following generations, which will benefit their offspring provided that they disperse locally as well (Figure 3b). Symbionts may become less abundant for instance because of additional intraspecific competition between them, as in Appendix A.5. As a result, more hosts remain non-mutualistic because they are less often associated with a symbiont (Figure A13), which further highlights that the evolution of mutualistic hosts relies on high symbiont densities.

The role of quasi-vertical transmission Although mutualistic symbionts are environmentally acquired, when both hosts and symbionts disperse locally this produces a similar effect as vertical transmission (as for mycorrhizae, Wilkinson, 1997), which we term "quasi-vertical" transmission. However, local dispersal (even 100%) is not equivalent to vertical transmission because host and symbiont offspring can disperse to any of the 8 neighbouring cells, so vertical transmission due to specific reproductive and physiological adaptations would have produced transitions to mutualism more easily. Moreover, the colonization of empty space by a mutualistic pair requires that both species disperse to the same remote place by chance, whereas in the case of vertical transmission this always occurs. Nevertheless, since hosts need to colonize empty space a significant fraction of hosts with mutualistic phenotypes also dispersed globally ($\sim 40\%$, Figure 3), which partly counteracts the necessity of quasi-vertical transmission. As well as hosts, mutualistic symbionts may also suffer from limited dispersal when they need to percolate in a landscape of non-adjacent hosts, which explains why they maintain $\sim 20\%$ of global dispersal (Figure 3). On the other hand, parasitic symbionts also evolve towards an intermediate dispersal strategy, although they tend to disperse globally much more often ($\sim 80\%$, Figure 3). In purely parasitic systems it has been shown that some degree of vertical transmission, which is close to local dispersal in our case, is necessary for persistence in fragmented landscapes (Su et al., 2019; Schinazi, 2000). In those cases as well as here, the parasitic population needs some degree of local dispersal in order to exploit a patch of hosts, once it has been "found" by global dispersers. Intermediate dispersal strategies have been found to favor persistence of a variety of systems. For instance, frequent short-distance and rare long-distance dispersal together favor metacommunity persistence in fragmented habitats (Huth et al., 2015) and intermediate migration rate is required for the spread of cooperative strategies in spatial prisoner's dilemma games (Vainstein et al., 2007).

Colonization ability comes along with the avoidance of local overpopulation generated by mutualism. This is in line with both the evolution of altruism, which can be limited by kin competition (Wilson et al., 1992; Alizon and Taylor, 2008) , and the evolution of dispersal which is in part due to the reduction of kin competition (Hamilton and May, 1977; Poethke et al., 2007; Harada, 1999). A mixed strategy combining both dispersal modes takes advantage of kin selection and simultaneously maintains the opportunity to escape kin competition. Figure 5b shows that purely local competition between hosts prevents the transition to mutualism because kin competition overcomes kin selection. Similarly, the evolution of cooperation by group selection can be hindered if competition between groups is local (Akdeniz and Van Veelen, 2020). In nature, global competition between hosts may arise when plants compete for water present in the same groundwater (Lejeune et al., 1999; Rietkerk et al., 2002), while competition for light is more local. Thus, the evolution of mutualism may depend on the dominant form of competition for resources between hosts.

Assumptions, limitations and generality of the model

Our results rely on several hypothesis which have contrasting effects on the likelihood of the transition to mutualism.

No vertical transmission We excluded the possibility of vertical transmission because it is a complex feature involving many traits, which more likely evolve some time after the transition once the mutualistic relationship is well established. For this reason an alternative mechanism is needed, and our results demonstrate that the coevolution of mutualistic effort and limited dispersal in both species can mimic vertical transmission, as argued above. However, in some parasitic systems (e.g. birds displacing parasitic flatworms, ticks carried on large vertebrates) vertical transmission may be a passive feature, present from the start. In such cases the evolution of mutualism is theoretically possible even if hosts keep dispersing globally, provided that mutations turning the parasites into mutualists exist.

Benefits and costs Benefits only depend on the interaction trait of the partner. In turn, costs depend on the interaction trait of the focal organism as well as on the benefits provided by the partner (Appendix A.1). This would correspond for instance to the development of organs like plant domatia (Szilágyi et al., 2009): if the symbiotic ants are mutualistic, the plant can grow bigger, thereby producing more domatia, which is more costly in absolute terms. An alternative would be to assume that the costs do not increase with the mutualistic benefit; this would in any case be favourable to the evolution of mutualism. Moreover, in the model some cost is paid even if the partner is parasitic or if the host is free of symbiont. For instance, domatia or extrafloral nectaries are unconditionally produced (Bronstein, 1998), even though domatia size can be plastic (Kokolo et al., 2020). Also, plants produce costly floral displays even in the absence of pollinators. Finally, another alternative arises when partners interact repeatedly, for instance during their growth. Using an iterated prisoners' dilemma model, Doebeli and Knowlton (1998) assumed that large received benefits trigger higher investment in the relationship. The interaction traits therefore become subject to phenotypic plasticity, in function of the partner's trait. This assumption favors the transition to mutualism since mutualists benefit more from being associated with mutualists. In contrast, our set of assumptions is more conservative.

Antagonistic coevolution of the parasitic system The evolutionary dynamics of the parasitic system have been ignored here, although they might affect the probability of transition. In the model the hosts cannot become resistant against the parasitic symbiont, which fits with the "superpathogen" of the gene-for-gene model (Salathé et al., 2008). This can be interpreted as a monomorphic long-term result of Red Queen dynamics, some constrain preventing the appearance of new resistant and virulent alleles. However, if the host-parasite interaction is instead ruled by a matching allele model (Salathé et al., 2008), dispersal and the associated spatial structure is likely to maintain polymorphism (Sasaki et al., 2002). During the early stages of the transition, formerly parasitic symbionts turned mutualistic will inherit this matching genetic system and will need to find compatible hosts. This adds another requirement, rendering the transition less likely.

Asexual reproduction Many models of (co)evolutionary dynamics assume asexual reproduction (e.g. Kéfi et al., 2008; Loeuille and Loreau, 2005), especially within the framework of Adaptive Dynamics (e.g. Dieckmann et al., 1995; Loeuille and Loreau, 2005). In the case of sexual reproduction, recombination may soften the correlation between dispersal and interaction traits, which is nevertheless essential to the transition. However, the work of Dieckmann and Doebeli (1999) on the coevolution between a niche and a mating trait showed that linkage disequilibrium can itself evolve, thereby preserving the correlation between traits. In the present case, we speculate that sexual reproduction would lengthen the waiting time until a successful transition, without hindering the transition in the long term.

Symbiont competition within hosts We previously assumed that only a single symbiont could infect a host, however several strains may compete within the same (Bongrand and Ruby, 2019; Zytynska and Weisser, 2016; Alizon et al., 2013). The host may be able to prevent the proliferation of parasitic strains (Sachs et al., 2010b), but parasitic strain may also overcome the others, which could prevent the evolution of mutualism (Jones et al., 2012). An extension of the model, presented in Appendix A.7, includes superinfections where mutualistic symbionts can be dislodged by parasites reaching the same host. When superinfection probability rises above 50%, the transition is prevented, otherwise mutualistic symbionts can persist, although at lower densities than without superinfections (Figure A15). Thus, although superinfections are clearly detrimental to the transition, mechanisms favouring the evolution of mutualism in our present model can resist some degree of competitive exclusion by parasites.

The evolution of cheating Our main interest was to understand how mutualism can evolve from a parasitic relationship (Roughgarden, 1975; Drew et al., 2021) but mutualism may also have evolved in the first place, the classic evolutionary problem in this case being how can it resist to the invasion of "cheaters" (e.g. Sachs et al., 2010a; Jones et al., 2015; Ferriere et al., 2002). According to Jones et al. (2015), cheating "(1) increases the fitness of the actor above average fitness in the population and (2) decreases the fitness of the partner below average fitness in the partner population". The latter condition is always satisfied by parasitic symbionts, but the former remains to be checked. Simulations starting with mutualistic symbionts only are rapidly invaded by parasites, leading to an evolutionary equilibrium identical to the one reached by Figure 2c (details not shown). The population-level fitness (sensu Metz et al., 1992) of parasites is therefore positive when they

are rare, thereby satisfying condition (1), and it gradually decreases to zero until the evolutionary equilibrium is reached. Hence, our model also accounts for the invasion by cheaters of an initially mutualistic system, leading to a coexistence of both strategies. Mutualism may also evolve from a competitive interaction, if two competitors start exchanging resources, each being a better exploiter of the resource it provides, and limited by the resource it receives (De Mazancourt and Schwartz, 2010). However it is unknown to what extent this kind of mutualism is sensitive to cheating; spatial effects similar to those studied here might stabilize it.

The interplay between several levels of selection

Although the first models of group selection relied on well-defined groups (e.g. Wilson, 1975; Smith, 1964), multilevel selection theory has since been extended to fuzzy group boundaries and more complex landscapes (e.g. Lion and Van Baalen, 2008; Nunney, 1985; Tekwa et al., 2015) like in the present case. Earlier in the discussion, intermediate dispersal has been interpreted as the result of a balance between two components of inclusive fitness, kin selection and kin competition, which have been recognized as particular cases of multilevel selection (Goodnight, 2005; Lion and Van Baalen, 2008; Queller, 1992; Sober and Wilson, 1999). Although our model is too complex for an analytical derivation of inclusive fitness, this should be possible in principle, as it has been done for simpler models of the evolution of altruism (Hamilton and Fox, 1975; Lion et al., 2011; Lehmann et al., 2007; Marshall, 2011; Wade, 1980). However, the levels-of-selection problem is more a question about the level at which there is a *causal* link between character and fitness (Okasha, 2006, 2016; Sober, 1984; Sober and Lewontin, 1982), rather than the level at which a mathematical formulation of fitness can be derived ("bookkeeping" in the words of S. J. Gould 2002, p. 619). Following Sober (1984), we will consider that selection at a given level of organization occurs if the different entities belonging to this level are variable with respect to some property causally involved in the survival or reproduction of the organisms forming the entities. Since Sober's formulation has been originally framed in the context of group selection, we first discuss how the levels-of-selection problem for mutualism can be related to the group selection debate in the context of altruism. The mechanism by which parasitic symbionts and hosts can invade mutualistic clusters is a two-species version of the tragedy of the commons (Garrett, 1968; Feeny et al., 1990; Hardin, 1998). In the case of altruism, the tragedy of the commons can be bypassed by local dispersal which triggers the formation of cooperative clusters (Mitteldorf and Wilson, 2000; Le Galliard et al., 2003), as in the present case. The evolution of altruism results from the conflict between two levels of selection, the organismic-level favouring cheaters and the group-level favoring altruism (Van Baalen and Rand, 1998; Simon et al., 2013; Wilson and Sober, 1989). At a given time step, neighbouring altruistic organisms help each other, which favors their fecundity. Shortly after the local density of altruists increases, which is beneficial for their offspring's fecundity as well. Since the transition to mutualism is egalitarian whereas the transition to altruism is fraternal, it is unclear if the evolution of mutualism involves the same levels of selection as for altruism. Sure enough, mutualism is also counter-selected at the organismic level, since mutualism is costly to both hosts and symbionts. However, differences between altruism and mutualism may arise at higher organization levels because at a given time step mutualists help their heterospecific partners but not their neighbouring conspecifics. In the present model the evolution of mutualism involves selection at the level of the host-symbiont pair, since at a given time step the reproduction of each of its organisms depends on the properties of the pair (the interaction traits α_h and α_s). This resembles the tit-for-tat strategy where cooperators are selected at the pair level (Wilson, 2004; Sober and Wilson, 1999). The mutualistic host-symbiont holobiont therefore emerges as a new unit of selection (Roughgarden et al., 2018; Drew et al., 2021).

Considering several times steps in a row, another level of selection appears. Since mutualists also disperse locally (Figure 3), after some time a mutualistic pair may trigger the formation of a mutualistic cluster (Figure 4c). Neighbouring mutualistic pairs do not help each other directly at a given time step, but indirectly by increasing the likelihood that their offspring will encounter mutualistic partners in the subsequent time steps. Although only hosts and symbionts reproduce in the traditional sense of organismic reproduction, the association between mutualistic hosts and symbionts is also re-produced (Doolittle and Inkpen, 2018; Griesemer, 2001) via local dispersal and cluster formation. Selection at the cluster level therefore occurs, since clusters dominated by mutualistic pairs will favour the reproduction of organisms and the re-production of mutualistic pairs. The re-production of pairs constitutes a another mechanism of inheritance, different from the one occurring during organismic reproduction. It is therefore hard to match Hull's (1980) categorization of replicators (here, hosts and symbionts) and interactors (here, pairs), since during the transition mutualistic pairs also acquire a replicative power via the evolution of local dispersal. This also emphasizes that Sober's (1984) formulation of group selection needs to be generalized for the present context, since the properties of clusters favor not only the reproduction of organisms but also the transmission of higher-level properties. Mutualistic clusters are self-perpetuating systems (Lenton et al., 2021), some of their properties being homeostatic (Ibanez, 2020). However, we believe this is not enough to qualify to evolutionary individuality (sensu Godfrey-Smith et al., 2013) since conflicts are still vivid (Queller and Strassmann, 2016); mutualistic clusters being prone to the invasion by parasites (Figure 4c).

Lastly, in the absence of dispersal cost mutualism rarely invades when host competition is weak (Figure 2a), despite the occasional formation of mutualistic pairs. Without dispersal cost, competition between hosts at the global scale is necessary for the transition to mutualism (Figure 5b). The global scale therefore constitutes another level of organization involved in the transition to mutualism. Global competition between hosts acts as an

environmental factor mitigating selection at the different organization levels discussed above. This environmental factor is not fixed by a parameter but instead determined by the evolutionary dynamics of the whole system, it is at the same time subject and object of evolution (Lewontin, 1982, 1983).

Host dependency and irreversibility of the transition

Major transitions in evolution are characterized by their irreversibility and by the interdependence between the agents (Szathmáry and Smith, 1995; Estrela et al., 2016). The model does not include any physiological or developmental dependence of the host on its symbiont, or any loss of functions in the host due to gene transfers, because we assumed that this generally occurs during later stages of the evolution of mutualism. Instead, dependence has been defined from a population dynamics perspective: the host is *ecologically dependent* when its population growth rate is negative in the absence of the symbiont. In that case the host can produce offspring, although not enough to compensate for mortality. In line with hypothesis H5, we found that mutualistic hosts deprived of their symbiont exhibit a negative growth rate when the host density after the transition to mutualism becomes sufficiently large (Figure 6). This ecological dependency resulted from the density-dependent competition between hosts and the assumption that mutualism is costly for the host, even when its symbiont is absent (as discussed above). However, ecological dependency is not absolute: once the density of hosts becomes sufficiently low, the mutualistic hosts alone are viable. Dependency may become absolute for a sufficiently high cost of mutualism, but in these conditions the transition to mutualism will not occur.

If host competition strength decreases permanently, for instance following the continuous supply of extra resources, the reverse transition back to parasitism occurs (Figure 5c). This has been documented in nature as well (Pellmyr and Leebens-Mack, 2000; Kawakita et al., 2015; Sachs et al., 2011), although the mechanisms involved may well be different. Reversal towards parasitism occurs because ecological dependency relies on host competition, which change with host densities, highlighting that mutualistic symbiosis may be sensitive to environmental change (Drew et al., 2021). However, if host competition decreases punctually, e.g., following a perturbation of a fraction of the landscape, then mutualism persists (Figure A11) because mutualistic clusters take advantage of the reduction of global host competition to colonize free cells around them. This leads to an increase in host competition; in that case mutualism can restore the ecological conditions allowing its own persistence, as in a niche construction process (Lewontin, 1982, 1983; Odling-Smee et al., 2013; Laland et al., 2016). Niche construction is generally understood as the improvement of abiotic conditions (e.g. Arnoldi et al., 2020). In the context of mutualism, it is due to the improvement of host densities, which induces an increase in host competition. This also occurs at the beginning of the transition, when the first mutualistic clusters trigger an increase in global host density. Although this has not been tested formally, the reversion is also very likely to occur if host competition for resources shifts from global to local, since it is apparent from Figure 5b that local competition completely prevents mutualism, even in the presence of dispersal cost.

Conclusion

In the present paper, we aim to understand the mechanisms promoting the transition from parasitism to mutualism. To tackle this issue, we develop an agent based model on a lattice. In our general model, we only assume that the mutualistic interactions influence the fecundity of both partners and that hosts face density-dependent competition. In addition, we ensure that the antagonistic system is stable in absence of mutations. We show that in the absence of vertical transmission or partner control mechanisms, the joint evolution between mutualistic effort and local dispersal can trigger the transition from parasitism to mutualism, provided that intraspecific competition between host is sufficiently global and that either dispersal cost or competition strength is large enough.

Unexpectedly, we found that mutualistic clusters invade the antagonistic system thanks to their ability to increase the population densities of both partners, thereby triggering global competition between hosts and rendering regions where hosts are mainly associated with parasitic symbionts unsuitable. In contrast, the higher fecundity of mutualists is not advantageous enough to compensate for the ability of parasites to invade mutualistic clusters. Thus, our results suggest that the eco-evolutionary feedback involving competition between hosts might promote the transition from parasitism to mutualism in a wide range of biological systems, such as plant-fungi, plant-ant and plant-seed-eating pollinator interactions.

Acknowledgements

We are grateful to Sébastien Lion, Jean-François Arnoldi, Sonia Kéfi, Nicolas Loeuille, Cédric Gaucherel, François Massol and four anonymous reviewers for their thoughtful comments and discussion on this work.

 # Data and code accessibility

A   Appendix

A.1   Mathematical and numerical details of the model

We present here the mathematical underpinnings of the model as well as some details of the numerical compu-
tation.

Rules of the individual based model description   Our model follows the cycle presented in Figure A1:

- Host and symbiont die with fixed probability $m \in (0, 1)$.
- They produce offspring, possibly with different traits from them due to mutation. The fecundity of the
parents depends on their two traits $(\alpha, \varepsilon) \in [0, 1]^2$ and on their interactions with their possibly cell-sharing
partner.
- The offspring are dispersed according to the parents' dispersal traits ε.
- The offspring of the hosts may establish only in empty cells, while the offspring of the symbionts can only
establish in cells already occupied by a solitary host. If several organisms arrive in the same cell, a lottery
determines which one will establish, while the others die.

In our numerical computations, mutations occurred only after the descendant was successfully established in a
cell. This procedure saves computational time and did not influence our results because offspring dispersal and
establishment do not depend on their traits but only on their parent traits. Furthermore, the mortality process
was applied to both types of agents simultaneously, while the reproduction and dispersal processes were applied
consecutively to the hosts and then to the symbionts. We confirmed that the order of the algorithm did not
qualitatively affect our results.

Fecundity and the average offspring number   The fecundity f of an agent depends on its mutualistic
interaction trait α as well as the interaction trait of its cell-sharing partner. This continuous trait ranging between
0 and 1 determines the intensity of the agent investment in the mutualistic relationship.

We assumed a positive interaction trait dependence between agents. A mutualistic agent tends to increase
the fecundity of its cell-sharing partner. The interaction fecundity $f_I^i(\alpha_j)$ of an organism of type $i \in \{h, s\}$, (h =
host, s = symbiont) interacting with an organism of type $j \in \{s, h\}$ with trait α_j was defined by

$$f_I^h(\alpha_s) = f_{min}^h + \left(f_{max} - f_{min}^h\right)\alpha_s^{\gamma_f}$$
$$f_I^s(\alpha_h) = f_{min}^s + \left(f_{max} - f_{min}^s\right)\alpha_h \tag{5}$$

625 The coefficient γ_f corresponds to the selection strength on the interaction trait α_s. Using a coefficient $\gamma_f > 1$, we
626 create a convex function allowing a transition from parasitism to mutualism for a central value of the symbiont
627 interaction trait. However, note that modifying the shape of this fecundity curve (from concave to convex via
628 linear) does not qualitatively change our results.

On the other hand, a mutualistic agent has an intrinsic cost reducing its fecundity. The mutualism cost
$C_m(\alpha_i)$ of an organism of type $i \in \{h, s\}$ (h = host, s = symbiont) ranges between 0 and 1, and it increases with
interaction trait α_i of the agent. It is defined by

$$C_m^i(\alpha_i) = 1 - c_m\alpha_i \tag{6}$$

630 where c_m is the maximal cost of mutualism.

Thus, for the host as for the symbiont, the fecundity f_i of an organism i interacting with an organism j is
the product of the interaction fecundity $f_I(\alpha_j)$ defined by (5) and the cost of mutualism $C_m(\alpha_i)$ defined by (6).

$$f^h = C_m^h(\alpha_h)f_I^h(\alpha_s)$$
$$f^s = C_m^s(\alpha_s)f_I^s(\alpha_h) \tag{7}$$

When a host agent is alone in a cell, its fecundity is defined by its intrinsic host fecundity f^a weighted by its
mutualism cost $C_m(\alpha_h)$: Fecundity of the solitary host:

$$f^{ha} = C_m^h(\alpha_h)f^a. \tag{8}$$

Figure A1: Sketch representation of the individual based model. The host population (H) and the symbiont population (S) undergo intrinsic mortality, then reproduction, dispersal, establishment, and finally mutation. The mortality step is simultaneous for the host and the symbiont, while the other steps occur first for the host and then for the symbiont.

631 We assume that the cost of mutualism is paid regardless of whether the interaction is realized.

633 In general, the average offspring number is not integer, yet the number of offspring in our model can only
634 be represented by an integer. Thus, in the numerical algorithm, the fecundity was used as the λ parameter of
635 a Poisson distribution. If the value drawn from the distribution was greater than the maximum fecundity f_{max},
636 then it was set back to the maximum fecundity.

Mutualism/parasitism threshold In our model, the presence of a host always produces a net benefit for the symbiont. However, the presence of the symbiont might be detrimental for the host. Indeed, the fecundity of a host h interacting with a symbiont s is $f^h(\alpha_h, \alpha_s) = f_I^h(\alpha_s)C_m(\alpha_h)$, while the fecundity of the same host h without a symbiont is $f^{ha}(\alpha_h) = f^a C_m(\alpha_h)$. Thus, the host has net benefit only if its fecundity in association with a symbiont is larger than its fecundity alone. Therefore, mutualism only occurs when $f_I^h(\alpha_s) > f^a$. Otherwise, the interaction is parasitic. This criterion does not depend on the host mutualism trait α_h because hosts always pay the same mutualism cost. Thus, we can define the mutualism/parasitism threshold α_s^* such that $f_I^h(\alpha_s^*) = f^a$; thus, we obtain

$$\alpha_s^* = \left(\frac{f^a - f_{min}^h}{f_{max} - f_{min}^h} \right)^{1/\gamma_f} \approx 0.475 \tag{9}$$

Competition To test the effect of the spatial scale of the competition, we introduced a scale parameter $w_h \in [0, 1]$ that weighs the effect of local ρ_h^{local} and global ρ_h^{global} host density on the competition. The establishment probability thus satisfies

$$P_I = 1 - \left((1 - w_h)\rho_h^{local} + w_h \, \rho_h^{global} \right)^{\gamma_C} \tag{10}$$

The local host density ρ_h^{local} corresponds to the host density in the 8 neighbouring cells surrounding the implantation cell of the host, while the global density ρ_h^{global} corresponds to the host density all over the landscape (see Figure A2 for a schematic representation).

local host density = 3/8
global host density = 7/36

Figure A2: Local and global host densities influence the probability of establishment in the focus cell (pink filled square). The global density corresponds to the host density in the whole 36-cell landscape. The local density corresponds to the density in the eight cells vicinity around the focus cell.

These competition scales may have various ecological explanations. For instance, plants sharing the same water table face global competition for this resource. Conversely, the competition for light between plants is an example of local competition. Thus, our competition scale model allows us to describe the competition for several different resources that may appear at different scales. Following our previous examples, if the water supply represents 90% of the competition and light supply represents only 10%, then the competition scale w_h is $w_h = 0.9$ (90% global competition and 10% local competition).

Distribution of mutation effects During reproduction, organisms generate offspring with traits that can deviate from their traits due to mutation. The effects of mutation on each trait are independent. However, the mutation effect does depends on the trait of the parent. For instance, an organism with trait α will give birth to an organism of trait $\alpha + \beta$ where β is drawn from a distribution with probability distribution function given by $K(\beta\,|\alpha)$, which depends on the trait of the parent α (Figure A3). In our model, we use a modified Beta distribution with shape parameters $(1, 3)$ to describe the effects of the mutation. More precisely, for a parent of trait α, the effect of mutaion is a random variable β defined by

$$\beta = (\beta_{max}\, \xi\, B)\mathbf{1}_{0 \leq \alpha + \beta_{max}\, \xi\, B \leq 1}$$

where B is a random variable, which follows a Beta distribution, ξ is a random variable independednt of B, which follows a Bernouilli distribution ($\mathbf{P}(\xi = 1) = \mathbf{P}(\xi = -1) = 1/2$). In other word, the random variable β follows the probability distribution function $K(\beta\,|\alpha)$, with $\alpha \in [0, 1]$:

$$K(\beta\,|\alpha) = \frac{3}{2\beta_{max}^3}\left(\beta_{max} - |\beta|\right)^2 \mathbf{1}_{|\beta| \leq \beta_{max}} \mathbf{1}_{0 \leq \alpha + \beta \leq 1} + K_0(\alpha)\delta_{\beta=0} \tag{11}$$

where $\mathbf{1}$ is the indicators function, δ is the Dirac mass and the function $K_0(\alpha)$ is defined by

$$K_0(\alpha) = \begin{cases} \dfrac{1}{2\beta_{max}^3}(\beta_{max} - \alpha)^3 & \text{if } \alpha \leq \beta_{max} \\ \quad 0 & \text{if } \beta_{max} \leq \alpha \leq 1 - \beta_{max} \\ \dfrac{1}{2\beta_{max}^3}(\beta_{max} - 1 - \alpha)^3 & \text{if } \alpha \geq 1 - \beta_{max} \end{cases} \tag{12}$$

Figure A3: Distribution of mutational effects $K(\beta|\alpha)$. Each parent of trait, e.g. α_1 or α_2, produce offspring with trait $\alpha_i + \beta$ where β has the density $K(\beta|\alpha_i)$ depending on its parent traits (red and blue curves for α_1 and α_2, respectively).

Figure A4: Effect of the average mutation effect (parameter β_{max} of distribution kernel of mutation effect) on the proportion of mutualistic symbionts.

Moreover, we investigate the effect of the maximal effects of mutation β_{max} on the proportion of mutualistic symbionts. From our formula, we know that the mean effect of mutation depends on the trait of the parent α but it is proportional to β_{max}, and it ranges between $3\beta_{max}/8$ for parents with intermediate trait ($\alpha \sim 0.5$) and $3\beta_{max}/4$ for parents with trait either close to 1 or 0. We show in Figure A4 that increasing the mean effect of mutation increases the proportion of mutualistic symbionts in the population. Thus large effects of mutation favour the emergence of mutualism. In our simulations we fix the maximal effect of mutation to $\beta_{max} = 0.5$.

Dispersal At each time step, hosts and symbionts produce offspring which can disperse over the landscape either locally or globally. For each agent, the proportion of its offspring dispersing globally is given by the dispersal trait ε. The location of offspring dispersed locally is chosen randomly uniformly over the 8 neighbors of its parents, while the location of those dispersed globally is chosen uniformly over the entire landscape expected the location of the parent (Figure A5 for the description of the local and global scale). In particular, a globally dispersed organism can arrive in the local neighbor of parents as the locally dispersed one. Moreover, the offspring are dispersed independently from each other and their location is chosen independently of the current landscape. In particular, offspring can arrive at an already occupied location and symbionts' offspring are not only dispersed in location where there is already an host. For instance if a host disperse 2/3 of its offspring at large distance from it, its dispersal trait satisfies $\varepsilon = 2/3$. Then the 2/3 of its offspring are dispersed randomly uniformly in

665 the entire landscape (red stars in Figure A5) while the remaining 1/3 is dispersed locally around it (red circles
666 in Figure A5).

Figure A5: Local and global dispersal of offspring from the host in the red cell over the landscape. Local
dispersal (red circles) occurs only within the neighborhood of the host (red dashed square) while global
dispersal (red stars) occurs over the entire landscape (red plain square). The host has a dispersal trait
$\varepsilon = 2/3$ and it disperses 6 offspring: 4 globally (red stars) and 2 locally (red circles).

Assortment index To compute the assortment index, we measured the similarity between spatially neigh-
bouring phenotypes for the spacial repartition resulting from the transition to mutualism and for the same spacial
repartition but with phenotypes randomly redistributed among organisms. The assortment index corresponds to
the difference between the measurement made on the space resulting from the transition to mutualism and the
measurement on the randomly rearranged space. If the index shifts positively (resp. negatively) from zero, it
means that similar phenotypes are closer (resp. more distant) than different phenotypes compared to random
spatial distribution. This methodology is similar to that used in Pepper and Smuts (2002) and Pepper (2007).
674

Intraspecific assortment index. More precisely, for the intraspecific assortment index we use the following
similarity index for host and symbiont. For each simulation and time t, we compute the similarity indices S_h and
S_s respectively among hosts and symbionts, as follows

$$S_h(t) = 1 - \frac{1}{N_h} \sum_{h=1}^{N_h} |\alpha_h - \overline{\alpha}_h| \tag{13}$$

where α_h is the trait of the host h and N_h is the total number of host in the landscape at time t. The quantity
$\overline{\alpha}_h$ is the average trait in the neighborhood V_h of the host h. The neighborhood V_h of a host h is the 8 closest
cells surrounding it (figure A2). It is defined by

$$\overline{\alpha}_h = \frac{1}{|V_h|} \sum_{i \in V_h} \alpha_i.$$

675 The similarity index among symbionts S_s is computed similarly.
676 Then for each time, we reshuffle the traits among the location occupied by hosts and symbionts and we
677 compute the associated similarity indices using equation (13). We average those indices over 1000 replicates to
678 compute the similarity indice S_{rh} and S_{rs} corresponding to a random spatial distribution.
Finally, We build the assortment index A_h as the difference between the similarity index of host S_h observed
and the similarity index S_{rh} of host when we randomly assigned trait of the host over the landscape,

$$A_h(t) = S_h(t) - S_{rh}(t). \tag{14}$$

679 We also compare our assortment index with the spatial autocorrelation Moran index for the host and symbiont.
680 The two indices show the same pattern. A positive spatial autocorrelation is observed after the transition occurred
681 (Figure A6).
682

Interspecific assortment index. For the assortment index between host and symbiont, we also use a measure of similarity between the host and symbiont trait at each location of the couple. More precisely, we define for each simulation and each time t the similarity index S_{sh} between host and symbiont sharing the same location as follows

$$S_{sh}(t) = 1 - \frac{1}{N_s} \sum_{s=1}^{N_s} |\alpha_s - \alpha_h| \tag{15}$$

where N_s is the number of symbiont, which is also the number of host–symbiont couple. As before, we compare this observed index with the random index S_{rsh} defined by randomly rearranging pairs of symbiont and host and taking average over 1000 replicates. The assortment index A_{sh} is thus given by

$$A_{sh}(t) = S_{sh}(t) - S_{rsh}(t). \tag{16}$$

We also compare our index with the correlation coefficient between the interaction traits of hosts and symbionts. We find a positive correlation between trait in a same location (Figure A6).

Figure A6: Spatial autocorrelation among hosts (plain curve) and symbionts (dashed curve) are described by the Moran index. The spatial correlation between the host and symbionts are described by the correlation coefficient (dash-dotted curve). The shadow regions corresponds to the 95% confident interval and curves corresponds to the median over 100 replicates. The parameters are similar as Figure 4.

Aggregation index From the assortment index analysis, we show that the symbionts and hosts are spatially assorted according to their trait. Now we aim to investigate how they are aggregated in space. We use a relative aggregation index $\mathcal{A}$ based on a measure of the number of pair of neighbors. More precisely, we define for any spatial configuration the number of pairs of neighbors P where a neighbor of an organism is its 8 closest cells. For instance, Figure A2 provides a schematic representation of a host spatial configuration and the dashed square represents the neighborhood of the red organism. The number of pair of the red organism is 3 in this example. Then for any spatial configuration with n organisms, we can define the maximal number of possible pair of organism which is given by $P_{max} = 4n - \lceil 6\sqrt{n} \rceil$ (Harary and Harborth, 1976). Thus, we define the aggregation index $\mathcal{A}$ as the ratio between P and P_{max}:

$$\mathcal{A} = \frac{P}{P_{max}}.$$

We compute the aggregation index over time for the hosts, the parasitic symbionts ($\alpha_s < \alpha_s^*$) and the mutualistic symbionts ($\alpha_s \geq \alpha_s^*$) (Figure A7).

Hosts are always more aggregated than symbionts. Moreover, after the transition occurred, mutualistic and parasitic symbionts have the same spatial signature in terms of aggregation. This pattern was already observed in Figure 4 where we see mutualistic clusters surrounded by parasitic clusters.

A.2 Mathematical approximations

In order to provide some heuristics about our stochastic model, we develop some deterministic approximation. This mathematical analysis also provides some quantitative insights on our choice of parameters and the threshold used to describe our outcomes (mutualistic vs parasitic symbionts, emergence of mutualists). We first investigate a simple model with hosts and parasitic symbionts and then we investigate a model with hosts and both parasitic and mutualistic symbionts.

Figure A7: Aggregation index of the spatial distribution of hosts (plain curve), parasitic symbionts (dash-dotted curve) and mutualistic symbionts (dashed curve) over time. The shadow regions corresponds to the 95% confident interval and curves corresponds to the median over 100 replicates. The parameters are similar as Figure 4.

An approximation of the parasitic system. First, we aim to describe the expected proportion of sites occupied by hosts and parasitic symbionts at equilibrium. We assume no mutations of interaction or dispersal traits and hosts and symbionts disperse globally randomly over the landscape composed of N sites. According to our model, the dynamics of the proportion of sites occupied by the host alone ρ_{ha} or host with symbionts ρ_{hs} is given by

$$
\begin{aligned}
\rho_{ha}(t+1) &= (1-m)\rho_{ha}(t)\left[\left(1 - f^s(\alpha_h, \alpha_s)\rho_{hs}(t)\right) + f^{ha}(\alpha_h)\left(1 - \rho_h(t)^{\gamma_C}\right)\right] \\
&\quad + (1-m)\rho_{hs}(t)f^h(\alpha_h, \alpha_s)\left(1 - \rho_h(t)^{\gamma_C}\right) \\
\rho_{hs}(t+1) &= (1-m)\rho_{hs}(t)\left(1 + f^s(\alpha_h, \alpha_s)\rho_{ha}(t)\right)
\end{aligned}
\tag{17}
$$

where $\rho_h = \rho_{ha} + \rho_{hs}$ is the total proportion of hosts and α_h and α_s are interactions trait of host and symbionts respectively. In this model, the traits are fixed – if $\alpha_s < \alpha_s^*$ symbionts are parasitic while there are mutualistic if $\alpha_s \geq \alpha_s^*$. Since the symbionts need host to survive, the proportion of sites occupied by symbionts is ρ_{hs}. Even if hosts and symbiont does not share the same mortality rate m, the model holds true by multiplying the terms f^s by $(1 - m_s)$ the survival rate of symbionts instead of $(1 - m)$. We can check that the following qualitative properties holds true with a different mortality rate. However, it will modify the quantitative outcome of the model.

For any given pair of interaction traits, we can compute the equilibria of this dynamical system.

Extinction. The extinction equilibrium, which corresponds to $\rho_{ha} = \rho_{hs} = 0$, always exists but it is unstable if

$$
\frac{1}{f^{ha}(\alpha_h)}\frac{m}{(1-m)} = \frac{1}{(1 - c_m\alpha_h)f^a}\frac{m}{(1-m)} < 1
$$

We have picked parameters, which fulfill this criterion (Table 1). In particular, we can see from this formula that increasing the mutualism cost c_m can lead to non viability of more mutualistic host. In our simulations, we fix this value to $c_m = 0.3$.

Absence of symbionts. We first look at the case where the symbionts are absent, $\rho_{hs} = 0$. Then we have $\rho_{ha} = \rho_h$ which is given by

$$
\rho_h = \rho_{ha} = \left(1 - \frac{1}{f^{ha}(\alpha_h)}\frac{m}{(1-m)}\right)^{1/\gamma_C} = \left(1 - \frac{1}{(1 - c_m\alpha_h)f^a}\frac{m}{(1-m)}\right)^{1/\gamma_C}
\tag{18}
$$

This equilibrium should be positive for any host trait α_h. Our fixed parameters stated in Table 1 satisfy this assumption and the proportion of host ρ_h alone without symbionts ranges between 0.36 and 0.5.

Without symbiont, the proportion of host converges to $\rho_h = \rho_{ha} = 50, 5\%$ ($\gamma_C = 0.2$ and $\alpha_h = 0$). Thus in absence of any symbionts, the host survives. Its intrinsic growth rate is given by $(1 - m)(1 + f^{ha}(\alpha_h))$. It is decreasing with α_h and it ranges between 1.41 ($\alpha_h = 0$) and 1.27 ($\alpha_h = 1$). Thus in absence of symbionts, the hosts always survives.

However, this equilibrium is unstable in our parameters range – the eigenvalue of the jacobian around this equilibrium has an eigenvalue with modulus greater than 1. This suggests that a third equilibrium exists and may be stable.

Coexistence of symbionts and host.   We also have the coexistence equilibrium which is given by the following formulae

$$\rho_{ha} = \frac{1}{f^s(\alpha_h, \alpha_s)} \frac{m}{(1-m)}$$

$$0 = \rho_h \Big(f^h(\alpha_h, \alpha_s)\big(1 - \rho_h^{\gamma_C}\big) - m \Big) + (1-m)\rho_{ha}\big(f^{ha}(\alpha_h) - f_h(\alpha_h, \alpha_s)\big)\big(1 - \rho_h^{\gamma_C}\big)$$

(19)

In our parameters range, this equilibrium always exists and it is always stable and attractive for any values of the interactions traits (α_h, α_s).

So, in the presence of a parasitic symbiont ($\alpha_s = 0$), the proportion of hosts converges to $\rho_h = 12,5\%$ and the proportion of symbionts to $\rho_{hs} = 10\%$ which is in accordance with our simulation at initial time $t = 0$ (Figure 2 b and c).

In addition, when hosts ($\alpha_h = 0$) are associated with mutualistic symbionts ($\alpha_s = 1$), the proportion of hosts rises to $\rho_h = 96\%$ and the proportion of symbionts to $\rho_{hs} = 92,5\%$. Thus the gain of cohabiting with mutualistic symbionts is indeed huge.

An approximation of the mutualistic/parasitic system without mutations.   We now investigate the outcome of the competition between mutualistic symbionts and parasitic symbionts on a host. In this case, our previous model (17) extends to the following form

$$
\begin{aligned}
\rho_{ha}(t+1) &= (1-m)\rho_{ha}(t)\left[\Big(1 - \frac{f^s(\alpha_h, \alpha_{sp})\rho_{sp}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)} - \frac{f^s(\alpha_h, \alpha_{sm})\rho_{sm}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\Big)\right.\\
&\left.+f^{ha}(\alpha_h)\Big(1 - \rho_h(t)^{\gamma_C}\Big)\right]\\
&+(1-m)\rho_{hs}(t)\big(1 - \rho_h(t)^{\gamma_C}\big)\left(\frac{f^h(\alpha_h,\alpha_{sp})\rho_{sp}(t)}{\rho_{sp}(t)+\rho_{sm}(t)} + \frac{f^h(\alpha_h,\alpha_{sm})\rho_{sm}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)\\
\rho_{hs}(t+1) &= (1-m)\rho_{hs}(t) + (1-m)\rho_{ha}(t)\Big(\frac{f^s(\alpha_h,\alpha_{sp})\rho_{sp}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)} + \frac{f^s(\alpha_h,\alpha_{sm})\rho_{sm}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\Big)\\
\rho_{sp}(t+1) &= (1-m)\rho_{sp}(t)\left(1 + \frac{f^s(\alpha_h,\alpha_{sp})\rho_{ha}(t)\rho_{sp}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)\\
\rho_{sm}(t+1) &= (1-m)\rho_{sm}(t)\left(1 + \frac{f^s(\alpha_h,\alpha_{sm})\rho_{ha}(t)\rho_{sm}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)
\end{aligned}
$$

(20)

where ρ_{sp} and ρ_{sm} represent respectively, the proportion of parasitic and mutualistic symbionts. The nature of the symbiont only depends on the interaction trait α_s: parasitic symbiont ($\alpha_{sp} < \alpha_s^*$), mutualistic symbiont ($\alpha_{sm} > \alpha_s^*$) where α_s^* is the mutualistic/parasitic threshold defined by (9).

For this model, we still recover the three equilibria described in the previous section: extinction $(0,0,0,0)$, host alone $(\rho_h, 0, 0, 0)$ (equation (18)), host with one symbiont $(\rho_{ha}, \rho_{hs}, \rho_{sp}, 0)$ or $(\rho_{ha}, \rho_{hs}, 0, \rho_{sm})$ (equation (19)). However, another equilibrium may exists: host with two coexisting symbionts $(\rho_{ha}, \rho_{hs}, \rho_{sp}, \rho_{sm})$. In the following section, we will investigate the stability of this coexisting equilibrium.

Competitive exclusion or coexistence of symbionts?   The outcome of this model crucially depends on the initial proportion of parasitic and mutualistic symbionts.

The coexistence of parasitic and mutualistic symbionts with a host may occur when the initial relative proportion of mutualistic symbiont is exactly the proportion at coexistence equilibrium, which is given by

$$p^* = \frac{f^s(\alpha_h, \alpha_{sp})}{f^s(\alpha_h, \alpha_{sp}) + f^s(\alpha_h, \alpha_{sm})}$$

(21)

However, when the proportion of mutualistic symbionts is initially larger than the critical threshold p^*, the mutualistic symbionts can win the competition by excluding the parasitic symbiont. While if the parasitic symbionts are more present with a proportion larger than $1 - p^*$, they drive the mutualistic symbionts toward extinction.

As a consequence, the coexistence of mutualistic and parasitic symbiont seems unlikely to occur without mutations among symbionts because of competitive exclusion. Moreover, this mechanism should create segregation if we use a spatially explicit model.

An approximation of the mutualistic/parasitic system with mutations.   In our stochastic model, the interaction trait may change from one symbiont to its offspring. So now we assume that during reproduction, symbionts can mutate from one type to the other at a rate $U = 0.02$. In this situation, the previous model

extends to

$$
\begin{aligned}
\rho_{ha}(t+1) \quad &= (1-m)\rho_{ha}(t)\left[\left(1 - \frac{f^s(\alpha_h,\alpha_{sp})\rho_{sp}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)} - \frac{f^s(\alpha_h,\alpha_{sm})\rho_{sm}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)\right.\\
&\quad \left. + f^{ha}(\alpha_h)\left(1 - \rho_h(t)^{\gamma_C}\right)\right]\\
&\quad + (1-m)\rho_{hs}(t)\left(1 - \rho_h(t)^{\gamma_C}\right)\left(\frac{f^h(\alpha_h,\alpha_{sp})\rho_{sp}(t)}{\rho_{sp}(t)+\rho_{sm}(t)} + \frac{f^h(\alpha_h,\alpha_{sm})\rho_{sm}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)\\
\rho_{hs}(t+1) \quad &= (1-m)\rho_{hs}(t) + (1-m)\rho_{ha}(t)\left(\frac{f^s(\alpha_h,\alpha_{sp})\rho_{sp}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)} + \frac{f^s(\alpha_h,\alpha_{sm})\rho_{sm}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right)\\
\rho_{sp}(t+1) \quad &= (1-m)\rho_{sp}(t)\left(1 + (1-U)\frac{f^s(\alpha_h,\alpha_{sp})\rho_{ha}(t)\rho_{sp}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right) + U(1-m)\frac{f^s(\alpha_h,\alpha_{sm})\rho_{ha}(t)\rho_{sm}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\\
\rho_{sm}(t+1) \quad &= (1-m)\rho_{sm}(t)\left(1 + (1-U)\frac{f^s(\alpha_h,\alpha_{sm})\rho_{ha}(t)\rho_{sm}(t)}{\rho_{sp}(t)+\rho_{sm}(t)}\right) + U(1-m)\frac{f^s(\alpha_h,\alpha_{sp})\rho_{ha}(t)\rho_{sp}^2(t)}{\rho_{sp}(t)+\rho_{sm}(t)}
\end{aligned}
\tag{22}
$$

In this case, the competitive exclusion process is damped by the mutation process and the coexistence always occurs. However, the proportion of mutualistic truly varies and depends on its initial proportion. If the initial proportion of mutualistic symbiont is above p^* stated in (21), then its proportion stabilizes around an equilibrium value $\rho_{sm}^* \approx 0.02$ (this value coincides with the initial value of our simulation, Figure A8d and Figure 2b-c). Thus this last model provides an accurate approximation of the antagonistic system used as initial condition of our stochastic model.

Transition from parasitic system to mutualistic/parasitic system. Our model also allows us to quantify the transition between a parasitic system with few mutualistic symbiont ($\rho_{sm} \sim 0.02$) and a system with a larger proportion of mutualistic symbiont. From the previous analysis, we capture the threshold p^* such that the system becomes mainly mutualistic. However, in our stochastic simulations, we do not observe this huge transition (Figure A9). When the percentage of mutualistic symbiont evolves, it stabilizes around 12.5% and the host density converges around 0.35.

From our deterministic model with mutation, we know that the density of host alone in the parasitic system, is around $\rho_{ha}^* = 0.1$. Moreover, the intrinsic growth rate of the mutualistic symbiont R_s is given from (22) by

$$
R_s = (1-m)\left(1 + (1-U)f^s(\alpha_h,\alpha_{sm})\rho_{ha}p_{sm}\right) + U(1-m)f^s(\alpha_h,\alpha_{sp})\rho_{ha}\frac{(1-p_{sm})^2}{p_{sm}}
\tag{23}
$$

where p_{sm} represents the proportion of mutualistic symbionts in the population. The mutualistic symbiont population increases if its intrinsic growth rate R_s is larger than 1, $R_s > 1$. In the parasitic system with the host density alone $\rho_{ha} = \rho_{ha}^* \approx 0.1$, the growth rate is larger than 1 if the proportion of mutualistic symbiont p_{sm} rises above $p_{sm}^* \approx 0.1$, where p_{sm}^* is such that $R_s(p_{sm}^*) = 1$. So in the parasitic system, if the proportion of mutualistic symbiont rises above 10%, its intrinsic growth rate becomes greater than 1 and its extinction probability falls below 1. As consequences, we can assume that if the proportion of mutualistic symbiont rises above the threshold of 10%, their percentage will remain truly above the 2% observed in the parasitic system, which characterizes a transition from parasitism to mutualism.

A.3 Competition strength, perturbation and mutualism persistence

Competition strength determines mutualism persistence In the main text, we show that competition is essential for the transition to mutualism, but it is also important for its persistence, as shown here. In this section, we explored the effect of a sudden variation in competition strength γ_C on the persistence of mutualism. We started with a strong competition $\gamma_C = 0.2$. As expected from our previous results, a transition to mutualism occurred (Figure A10a), c) and d)). Then, around $t = 10^4$, we suddenly switched the competition strength to $\gamma_C = 2$, corresponding to negligible competition. We observed reversal of mutualism due to the proportion of mutualistic symbionts decreasing from 20% to less than 5% (Figure A10a) and e)). We observed that the reversal of mutualism due to a weakening of the competition corresponded with an increase in host and symbiont densities. This increase is due to the reduction of competition, which determines densities more than the presence or absence of mutualism does.

Density perturbation does not affect mutualism persistence Next, we tested how mutualism responds to a decrease in competition due to eradication of hosts and symbionts in a large homogeneous region of space (Figure A11). While previously we demonstrated that mutualism regresses when competition is set to be weak, we show here that mutualism persists in the face of decreased competition due to decreased host density.

(a) Without mutation and $p = 0.6$

(b) Without mutation and $p = p^*$

(c) Without mutation and $p = 0.15$

(d) With mutations and $p = p^*$

Figure A8: Evolution over time of the host-symbiont model (20) (panel a-c) and (22) (panel d) with two types of symbionts: parasitic symbionts ($\alpha_s = 0$) and mutualistic symbionts ($\alpha_s = 1$). We present the model (20) without mutation for various initial relative proportion of mutualistic symbiont p: (a) $p = 0.6$, (b) $p = p^* \approx 0.45$ and (c) $p = 0.15$. The model (22) with mutation is presented in (d) with initial proportion of mutualistic symbiont $p = p^*$.

Figure A9: Evolution of the percentage of mutualistic symbionts in the population over 129 simulations. Red curves corresponds to replicates such that the percentage of mutualistic symbionts remains greater than the threshold of 10% – transition to mutualism. Black curves corresponds to replicates where the percentage remains below the 10% threshold – no transition.

Figure A10: The transition to mutualism and collapse of mutualism depending on competition. a) Host and symbiont average interaction traits and the percentage of mutualistic symbionts over time. b) Host and symbiont abundance. There is strong competition from time $t = 0$ to $t = 10^4$ and negligible competition from $t = 10^4$ to the end. c) Dispersal and interaction traits distribution before the transition to mutualism. d) Dispersal and interaction traits distribution during mutualism persistence. e) Dispersal and interaction traits distribution after mutualism collapse. Parameters are $m = 0.06$, $c_m = 0.3$, $w_h = 1$, $f^h_{min} = 0.1$, $f^s_{min} = 2.5$, $f_{max} = 8$, $f^a = 0.5$, $\beta_{max} = 0.5$ and $d = 0$.

A.4 Mortality and dispersal cost can induce host dependency in emerging mutualistic systems.

In the main text, we focused on the effect of mortality m and dispersal cost d on the transition to mutualism and host dependency. Here, we present in more detail the effect of dispersal cost on the distribution of hosts and symbionts in trait space for three values of dispersal cost and fixed mortality rate (Figure A12). In addition, the table A1 shows the features of the clusters in the trait distribution.

We demonstrated that the dispersal cost favours the transition to mutualism. Moreover, even when the cost was high, the features of the clusters revealed that parasitic symbionts maintained a more global dispersal than

Figure A11: Maintenance of mutualism in the face of a reduction in competition caused by a perturbation creating a large square of free cells. Snapshot at several times: $t-1$ is the eco-evolutionary equilibrium with mutualism just before the perturbation, then at t_0 the perturbation, and then $t+5$, $t+10$, $t+100$ and $t+10000$ after the perturbation. In black, the free cells; in green, the hosts alone; in blue, the couples with parasitic symbionts; and in red, the couples with mutualistic symbionts. Parameters are $m=0.06$, $c_m=0.3$, $w_h=1$, $\gamma_C=0.2$, $f_{min}^h=0.1$, $f_{min}^s=2.5$, $f_{max}=8$, $f^a=0.5$, $\beta_{max}=0.5$ and $d=0$.

783 mutualistic symbionts.

Figure A12: Joint distribution of the host (green), the parasitic symbiont (blue) and the mutualistic symbiont (red) populations in the traits domain for mortality $m=0.03$ and an increasing dispersal cost $d \in \{0, 0.3, 0.75\}$. The ellipses correspond to the standard deviation. The 48 runs averaged in Figure 6 are plotted together. Other parameters are $c_m=0.3$, $w_h=1$, $\gamma_C=0.2$, $f_{min}^h=0.1$, $f_{min}^s=2.5$, $f_{max}=8$, $f^a=0.5$ and $\beta_{max}=0.5$.

dispersal cost	species	mean interaction trait	mean dispersal trait
$d = 0$	hosts	0.074	0.81
	parasitic symbionts	0.06	0.82
	mutualistic symbionts	** (density < 2%)*	** (density < 2%)*
$d = 0.3$	hosts	0.45	0.35
	parasitic symbionts	0.07	0.39
	mutualistic symbionts	0.88	0.17
$d = 0.75$	hosts	0.36	0.19
	parasitic symbionts	0.08	0.18
	mutualistic symbionts	0.84	0.10

Table A1: Features of the clusters in the traits domain

A.5 Density-dependent competition between symbionts

In the main text, symbionts compete for free hosts, which is a form of density-dependent competition. Other ecological factors may also lead to density-dependent competition between symbionts, for instance if symbionts compete for resources that are not provided by the hosts. Figure A13b shows that density-dependent competition between symbionts reduces symbiont density, as expected. Hosts are therefore free of symbionts more often, which selects for non-mutualistic hosts (Figure A13c, to be compared with Figure 3b).

Figure A13: Effect of density-dependent competition between symbionts on host evolution. a) Symbiont and host densities after and before the transition, without density-dependent competition between symbionts. b) Symbiont and host densities after and before the transition, with density-dependent competition between symbionts. c) Distribution of host traits after the transition, with density-dependent competition between symbionts. To be compared with Figure 3b.

A.6 Evolutionary rescue

Figure 6 provides evidence for evolutionary rescue, as discussed in the main text. Figure A14 shows that this occurs only in a fraction of the simulations, when mutualists arise soon enough to rescue the whole system.

Figure A14: Frequency at which evolutionary rescue occurs. This figure is identical to Figure 6, except that it maps the probability of extinction before the transition, instead of the mean percentage of mutualistic symbionts. The dotted black line indicates the upper boundary of viability for the parasitic system, without evolution. Above the dotted black line, in some cases the evolution of mutualism rescued the whole system, although the parasitic system is unviable alone. In the white region, the systems goes extinct, even with evolution

A.7   Symbionts competition within hosts

In our current model, a host can be colonised only by one symbiont and once the symbiont is established on a host, it cannot be replaced by another symbiont. Furthermore, when several symbionts arrive at the same time on an available host, the symbiont, which establishes, is chosen randomly uniformly among the contenders. Here, we relax these assumptions in order to model symbionts' competition within a host, or "superinfection". We assume that within a host, the most parasitic symbiont, with the lowest interaction trait, is the most competitive symbiont. Thus, it will be more efficient to establish in a host or dislodge a symbionts from the host.

Establishment of symbionts on a host.   Specifically, when N symbionts, with trait $\{\alpha_1, \ldots, \alpha_N\}$ arrive on a host, the establishment probability P_e^i of the symbiont i is given by :

$$P_e^i = \frac{p_i}{\sum\limits_{i=1}^{N} p_i} \quad \text{with} \quad p_i = \min\left(\max\left(\frac{1}{N} - (\alpha_i - \bar{\alpha})S_{max}, 0\right), 1\right), \quad \text{and} \quad \bar{\alpha} = \frac{1}{N}\sum_{j=1}^{N}\alpha_j \tag{24}$$

where S_{max} measures the superinfections' intensity, which corresponds to the maximal competitive advantage of a symbiont. For instance, when a truly parasitic symbiont $\alpha_1 = 0$ tries to establish with a truly mutualsistic symbiont $\alpha_2 = 1$, its establishment probability is $P_e^1 = (1 + S_{max})/2 \geq 1/2$. The establishment probability of the mutualsitic symbiont is $P_e^2 = (1 - S_{max})/2 \leq 1/2$. If $S_{max} = 0$, they have the same probability of establishment, while if $S_{max} = 1$, the parasitic symbiont always over-competes the mutualistic symbiont.

Replacement of a resident symbiont   When N symbionts with trait $\{\alpha_1, \ldots, \alpha_N\}$ arrive in a host already occupied by a resident symbiont with trait α_s, they may dislodge the resident. Specifically, the probability of the resident symbiont to persist P_p is given by

$$P_p = \min\left(1 - (\alpha_s - \bar{\alpha})S_{max}, 1\right) \quad \text{with} \quad \bar{\alpha} = \frac{1}{N}\sum_{i=1}^{N}\alpha_i. \tag{25}$$

In particular, if the resident has a trait α_s lower than the mean trait of the invaders $\bar{\alpha}$, then the resident always persists. Otherwise, the resident may be dislodged with a probability smaller than S_{max}. Then if the resident is dislodged, the establishment probability of the N invader symbionts is given by the previous formula (24).

Figure A15 shows the effect of the superinfection intensity S_{max} on the percentage of mutualistic symbionts. We show that despite the competitive advantage of parasitic symbionts when competing for a host, the transition to mutualism is possible when the superinfection intensity is not too large (if $S_{max} < 1/2$, transition occurs, that is the percentage of mutualistic symbionts stays above 10%). Moreover, when $S_{max} < 1/2$, the trait distribution of symbionts is bimodal.

Figure A15: a) Percentage of mutualistic symbionts in function of the maximum superinfection advantage S_{max} averaged over 20 simulations per parameter values. b) Distributions of symbionts population in traits domain according to five intensity of superinfection advantage S_{max}. Distributions corresponds to 20 simulations for each parameter values. These results are obtained with a maximum time projection of 5000 time steps, a strong and global competition ($\gamma_C = 0.2$) and a dispersal cost ($d = 0$). Others parameters are $m = 0.06$, $c_m = 0.3$, $f_{min}^h = 0.1$, $f_{min}^s = 2.5$, $f_{max} = 8$, $f^a = 0.5$ and $\beta_{max} = 0.5$.

References

Akçay, E. (2015). Evolutionary models of mutualism. In Bronstein, J. L., editor, *Mutualism*, pages 57–76. Oxford University Press.

Akdeniz, A. and Van Veelen, M. (2020). The cancellation effect at the group level. *Evolution*, 74(7):1246–1254.

Akman Gündüz, E. and Douglas, A. E. (2009). Symbiotic bacteria enable insect to use a nutritionally inadequate diet. *Proceedings of the Royal Society B: Biological Sciences*, 276(1658):987–991.

Akçay, E. (2017). Population structure reduces benefits from partner choice in mutualistic symbiosis. *Proceedings of the Royal Society B: Biological Sciences*, 284(1850):2016–2317.

Alizon, S., de Roode, J. C., and Michalakis, Y. (2013). Multiple infections and the evolution of virulence. *Ecology Letters*, 16(4):556–567.

Alizon, S. and Taylor, P. (2008). Empty sites can promote altruistic behavior. *Evolution: International Journal of Organic Evolution*, 62(6):1335–1344.

Allen, M. F. (1991). *The ecology of mycorrhizae*. Cambridge University Press.

Arnoldi, J.-F., Coq, S., Kéfi, S., and Ibanez, S. (2020). Positive plant–soil feedback trigger tannin evolution by niche construction: A spatial stoichiometric model. *Journal of Ecology*, 108(1):378–391.

Axelrod, R. (1981). The emergence of cooperation among egoists. *American political science review*, 75(2):306–318.

Bennett, G. M. and Moran, N. A. (2015). Heritable symbiosis: the advantages and perils of an evolutionary rabbit hole. *Proceedings of the National Academy of Sciences*, 112(33):10169–10176.

Bever, J. D., Richardson, S. C., Lawrence, B. M., Holmes, J., and Watson, M. (2009). Preferential allocation to beneficial symbiont with spatial structure maintains mycorrhizal mutualism. *Ecology letters*, 12(1):13–21.

Bongrand, C. and Ruby, E. G. (2019). Achieving a multi-strain symbiosis: strain behavior and infection dynamics. *The ISME journal*, 13(3):698–706. Publisher: Nature Publishing Group.

Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E., Mustin, K., and Saastamoinen, M. (2012). Costs of dispersal. *Biological reviews*, 87(2):290–312. Publisher: Wiley Online Library.

Boots, M. and Mealor, M. (2007). Local interactions select for lower pathogen infectivity. *Science*, 315(5816):1284–1286.

Bordenstein, S. R. and Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten Principles of Holobionts and Hologenomes. *PLOS Biology*, 13(8):e1002226.

Borges, R. M. (2015). How to be a fig wasp parasite on the fig–fig wasp mutualism. *Current Opinion in Insect Science*, 8:34–40.

Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in ecology & evolution*, 9(6):214–217.

Bronstein, J. L. (1998). The contribution of ant-plant protection studies to our understanding of mutualism 1. *Biotropica*, 30(2):150–161.

Bronstein, J. L. (2015). *Mutualism*. Oxford University Press, USA.

Bronstein, J. L., Alarcón, R., and Geber, M. (2006). The evolution of plant–insect mutualisms. *New Phytologist*, 172(3):412–428.

Callaway, R. M., Brooker, R. W., Choler, P., Kikvidze, Z., Lortie, C. J., Michalet, R., Paolini, L., Pugnaire, F. I., Newingham, B., Aschehoug, E. T., and Armas, C. (2002). Positive interactions among alpine plants increase with stress. *Nature*, 417(6891):844–848.

Darwin, C. (1871). *The descent of man, and selection in relation to sex*. John Murray, London.

De Mazancourt, C. and Schwartz, M. W. (2010). A resource ratio theory of cooperation. *Ecology letters*, 13(3):349–359. Publisher: Wiley Online Library.

Débarre, F., Lion, S., Van Baalen, M., and Gandon, S. (2012). Evolution of host life-history traits in a spatially structured host-parasite system. *The American Naturalist*, 179(1):52–63.

Denison, R. F. and Kiers, E. T. (2004). Why are most rhizobia beneficial to their plant hosts, rather than parasitic? *Microbes and Infection*, 6(13):1235–1239.

Després, L. and Jaeger, N. (1999). Evolution of oviposition strategies and speciation in the globeflower flies chiastocheta spp.(anthomyiidae). *Journal of Evolutionary Biology*, 12(4):822–831.

Dieckmann, U. and Doebeli, M. (1999). On the origin of species by sympatric speciation. *Nature*, 400(6742):354–357.

Dieckmann, U., Marrow, P., and Law, R. (1995). Evolutionary cycling in predator-prey interactions: population dynamics and the red queen. *Journal of Theoretical Biology*, 176(1):91–102.

Doebeli, M. and Knowlton, N. (1998). The evolution of interspecific mutualisms. *Proceedings of the National Academy of Sciences*, 95(15):8676–8680.

Doolittle, W. F. and Inkpen, S. A. (2018). Processes and patterns of interaction as units of selection: An introduction to ITSNTS thinking. *Proceedings of the National Academy of Sciences*, 115(16):4006–4014.

Drew, G. C., Stevens, E. J., and King, K. C. (2021). Microbial evolution and transitions along the parasite–mutualist continuum. *Nature Reviews Microbiology*, pages 1–16.

Edwards, D. P., Hassall, M., Sutherland, W. J., and Yu, D. W. (2006). Selection for protection in an ant–plant mutualism: host sanctions, host modularity, and the principal–agent game. *Proceedings of the Royal Society B: Biological Sciences*, 273(1586):595–602.

Estrela, S., Kerr, B., and Morris, J. J. (2016). Transitions in individuality through symbiosis. *Current opinion in microbiology*, 31:191–198.

Feeny, D., Berkes, F., McCay, B. J., and Acheson, J. M. (1990). The tragedy of the commons: twenty-two years later. *Human ecology*, 18(1):1–19.

Ferdy, J.-B. and Godelle, B. (2005). Diversification of transmission modes and the evolution of mutualism. *The American Naturalist*, 166(5):613–627.

Ferriere, R., Bronstein, J. L., Rinaldi, S., Law, R., and Gauduchon, M. (2002). Cheating and the evolutionary stability of mutualisms. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1493):773–780.

Ferriere, R. and Legendre, S. (2013). Eco-evolutionary feedbacks, adaptive dynamics and evolutionary rescue theory. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1610):20120081.

Fisher, R. M., Cornwallis, C. K., and West, S. A. (2013). Group formation, relatedness, and the evolution of multicellularity. *Current Biology*, 23(12):1120–1125.

Fisher, R. M., Henry, L. M., Cornwallis, C. K., Kiers, E. T., and West, S. A. (2017). The evolution of host-symbiont dependence. *Nature Communications*, 8(1):15973.

Foster, K. R. and Wenseleers, T. (2006). A general model for the evolution of mutualisms. *Journal of evolutionary biology*, 19(4):1283–1293.

Frank, S. A. (1994). Genetics of mutualism: the evolution of altruism between species. *Journal of Theoretical Biology*, 170(4):393–400.

Garrett, H. (1968). The tragedy of the commons. *Science*, 162(3859):1243–1248.

Genkai-Kato, M. and Yamamura, N. (1999). Evolution of mutualistic symbiosis without vertical transmission. *Theoretical population biology*, 55(3):309–323.

Gilbert, S. F., Sapp, J., and Tauber, A. I. (2012). A Symbiotic View of Life: We Have Never Been Individuals. *The Quarterly Review of Biology*, 87(4):325–341.

Godfrey-Smith, P., Goodnight, C. J., Clarke, E., Okasha, S., Pradeau, T., Gardner, A., van Baalen, M., Hamilton, A., Fewell, J., and Haber, M. (2013). *From Groups to Individuals: Evolution and Emerging Individuality*. MIT Press.

Gomulkiewicz, R. and Holt, R. D. (1995). When does evolution by natural selection prevent extinction? *Evolution*, pages 201–207. Publisher: JSTOR.

Goodnight, C. J. (2005). Multilevel selection: the evolution of cooperation in non-kin groups. *Population ecology*, 47(1):3–12.

Gould, S. J. (2002). *The structure of evolutionary theory*. Harvard University Press.

Griesemer, J. (2001). The units of evolutionary transition. *Selection*, 1(1):67–80.

Hamilton, W. D. (1964a). The genetical evolution of social behaviour. i. *Journal of Theoretical Biology*, 7(1):1–16.

Hamilton, W. D. (1964b). The genetical evolution of social behaviour. ii. *Journal of theoretical biology*, 7(1):17–52.

Hamilton, W. D. and Fox, R. (1975). Innate social aptitudes of man: an approach from evolutionary genetics. *Narrow roads of gene land. Vol. 1: Evolution of social behaviour*, pages 315–52.

Hamilton, W. D. and May, R. M. (1977). Dispersal in stable habitats. *Nature*, 269(5629):578–581.

Harada, Y. (1999). Short- vs. Long-range Disperser: the Evolutionarily Stable Allocation in a Lattice-Structured Habitat. *Journal of Theoretical Biology*, 201(3):171–187.

Harary, F. and Harborth, H. (1976). Extremal animals. *Journal of Combinatorics, Information & System Sciences*, 1:1–8.

Harcombe, W. (2010). Novel cooperation experimentally evolved between species. *Evolution: International Journal of Organic Evolution*, 64(7):2166–2172.

Hardin, G. (1998). Extensions of "the tragedy of the commons". *Science*, 280(5364):682–683.

Herre, E. A., Knowlton, N., Mueller, U. G., and Rehner, S. A. (1999). The evolution of mutualisms: exploring

the paths between conflict and cooperation. *Trends in ecology & evolution*, 14(2):49–53.

Hochberg, M. E., Rankin, D. J., and Taborsky, M. (2008). The coevolution of cooperation and dispersal in social groups and its implications for the emergence of multicellularity. *BMC Evolutionary Biology*, 8(1):1–14.

Hull, D. L. (1980). Individuality and selection. *Annual review of ecology and systematics*, pages 311–332.

Huth, G., Haegeman, B., Pitard, E., and Munoz, F. (2015). Long-distance rescue and slow extinction dynamics govern multiscale metapopulations. *The American Naturalist*, 186(4):460–469.

Ibanez, S. (2020). The Evolution of Ecosystem Phenotypes. *Biological Theory*, 15(2):91–106.

Johnson, N. C., Graham, J.-H., and Smith, F. A. (1997). Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New phytologist*, 135(4):575–585.

Jones, E. I., Afkhami, M. E., Akçay, E., Bronstein, J. L., Bshary, R., Frederickson, M. E., Heath, K. D., Hoeksema, J. D., Ness, J. H., and Pankey, M. S. (2015). Cheaters must prosper: reconciling theoretical and empirical perspectives on cheating in mutualism. *Ecology letters*, 18(11):1270–1284.

Jones, E. I., Bronstein, J. L., and Ferrière, R. (2012). The fundamental role of competition in the ecology and evolution of mutualisms. *Annals of the New York Academy of Sciences*, 1256(1):66–88.

Jordano, P., García, C., Godoy, J. A., and García-Castaño, J. L. (2007). Differential contribution of frugivores to complex seed dispersal patterns. *Proceedings of the National Academy of Sciences*, 104(9):3278–3282. Publisher: National Acad Sciences.

Kawakita, A., Mochizuki, K., and Kato, M. (2015). Reversal of mutualism in a leafflower–leafflower moth association: the possible driving role of a third-party partner. *Biological Journal of the Linnean Society*, 116(3):507–518. Publisher: Oxford University Press.

Kéfi, S., Rietkerk, M., Van Baalen, M., and Loreau, M. (2007). Local facilitation, bistability and transitions in arid ecosystems. *Theoretical population biology*, 71(3):367–379.

Kéfi, S., Van Baalen, M., Rietkerk, M., and Loreau, M. (2008). Evolution of local facilitation in arid ecosystems. *The American Naturalist*, 172(1):E1–E17.

Kerr, B. and Nahum, J. (2011). The evolution of restraint in structured populations: setting the stage for an egalitarian major transition. In *The Major Transitions in Evolution Revisited*, pages 127–140. MIT Press, Cambridge.

Kerr, B., Neuhauser, C., Bohannan, B. J. M., and Dean, A. M. (2006). Local migration promotes competitive restraint in a host–pathogen 'tragedy of the commons'. *Nature*, 442(7098):75–78.

Kiers, E. T., Rousseau, R. A., West, S. A., and Denison, R. F. (2003). Host sanctions and the legume–rhizobium mutualism. *Nature*, 425(6953):78–81.

King, K. C., Brockhurst, M. A., Vasieva, O., Paterson, S., Betts, A., Ford, S. A., Frost, C. L., Horsburgh, M. J., Haldenby, S., and Hurst, G. D. (2016). Rapid evolution of microbe-mediated protection against pathogens in a worm host. *The ISME journal*, 10(8):1915–1924.

Koella, J. C. (2000). The spatial spread of altruism versus the evolutionary response of egoists. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267(1456):1979–1985.

Kokolo, B., Attéké Nkoulémbéné, C., Ibrahim, B., M'Batchi, B., and Blatrix, R. (2020). Phenotypic plasticity in size of ant-domatia. *Scientific Reports*, 10(1):20948. Number: 1 Publisher: Nature Publishing Group.

Laland, K., Matthews, B., and Feldman, M. W. (2016). An introduction to niche construction theory. *Evolutionary ecology*, 30(2):191–202.

Le Galliard, J.-F., Ferriere, R., and Dieckmann, U. (2005). Adaptive evolution of social traits: origin, trajectories, and correlations of altruism and mobility. *The American Naturalist*, 165(2):206–224.

Le Galliard, J.-F., Ferriére, R., and Dieckmann, U. (2003). The Adaptive Dynamics of Altruism in Spatially Heterogeneous Populations. *Evolution*, 57(1):1–17.

Lehmann, L., Keller, L., West, S., and Roze, D. (2007). Group selection and kin selection: two concepts but one process. *Proceedings of the National Academy of Sciences*, 104(16):6736–6739.

Lejeune, O., Couteron, P., and Lefever, R. (1999). Short range co-operativity competing with long range inhibition explains vegetation patterns. *Acta Oecologica*, 20(3):171–183.

Lenton, T. M., Kohler, T. A., Marquet, P. A., Boyle, R. A., Crucifix, M., Wilkinson, D. M., and Scheffer, M. (2021). Survival of the Systems. *Trends in Ecology & Evolution*, 36(4):333–344.

Lewontin, R. C. (1982). Organism and environment. ic plotkin (ed.), learning, development and culture: Essays in evolutionary epistemology.

Lewontin, R. C. (1983). Gene, organism and environment. *Evolution from molecules to men*, 273:285.

Lion, S., Jansen, V. A., and Day, T. (2011). Evolution in structured populations: beyond the kin versus group debate. *Trends in ecology & evolution*, 26(4):193–201.

Lion, S. and Van Baalen, M. (2007). From infanticide to parental care: why spatial structure can help adults be

good parents. *The American Naturalist*, 170(2):E26–E46.

Lion, S. and Van Baalen, M. (2008). Self-structuring in spatial evolutionary ecology. *Ecology letters*, 11(3):277–295.

Loeuille, N. and Loreau, M. (2005). Evolutionary emergence of size-structured food webs. *Proceedings of the National Academy of Sciences*, 102(16):5761–5766.

Mack, K. M. (2012). Selective feedback between dispersal distance and the stability of mutualism. *Oikos*, 121(3):442–448.

Maestre, F. T., Bautista, S., and Cortina, J. (2003). Positive, negative, and net effects in grass–shrub interactions in mediterranean semiarid grasslands. *Ecology*, 84(12):3186–3197.

Margulis, L. and Sagan, D. (2002). *Acquiring genomes: a theory of the origins of species*. Basic Books, New York, NY, 1st ed edition.

Marshall, J. A. (2011). Group selection and kin selection: formally equivalent approaches. *Trends in Ecology & Evolution*, 26(7):325–332.

McFall-Ngai, M. J. (2014). The importance of microbes in animal development: lessons from the squid-vibrio symbiosis. *Annual review of microbiology*, 68:177–194.

Metz, J. A. J., Nisbet, R. M., and Geritz, S. A. H. (1992). How should we define 'fitness' for general ecological scenarios? *Trends in Ecology & Evolution*, 7(6):198–202.

Mitteldorf, J. and Wilson, D. S. (2000). Population viscosity and the evolution of altruism. *Journal of theoretical biology*, 204(4):481–496.

Mullon, C., Keller, L., and Lehmann, L. (2018). Social polymorphism is favoured by the co-evolution of dispersal with social behaviour. *Nature ecology & evolution*, 2(1):132–140.

Nunney, L. (1985). Group Selection, Altruism, and Structured-Deme Models. *The American Naturalist*, 126(2):212–230.

Odling-Smee, F. J., Laland, K. N., and Feldman, M. W. (2013). *Niche construction: the neglected process in evolution (MPB-37)*. Princeton university press.

Okasha, S. (2006). *Evolution and the Levels of Selection*. Oxford University Press.

Okasha, S. (2016). The relation between kin and multilevel selection: an approach using causal graphs. *The British Journal for the Philosophy of Science*, 67(2):435–470.

Pellmyr, O. and Leebens-Mack, J. (2000). Reversal of mutualism as a mechanism for adaptive radiation in yucca moths. *the american naturalist*, 156(S4):S62–S76. Publisher: The University of Chicago Press.

Pepper, J. W. (2007). Simple models of assortment through environmental feedback. *Artificial life*, 13(1):1–9.

Pepper, J. W. and Smuts, B. B. (2002). A mechanism for the evolution of altruism among nonkin: positive assortment through environmental feedback. *The American Naturalist*, 160(2):205–213.

Poethke, H. J., Pfenning, B., and Hovestadt, T. (2007). The relative contribution of individual and kin selection to the evolution of density-dependent dispersal rates. *Evolutionary Ecology Research*.

Purcell, J., Brelsford, A., and Avilés, L. (2012). Co-evolution between sociality and dispersal: the role of synergistic cooperative benefits. *Journal of Theoretical Biology*, 312:44–54.

Queller, D. C. (1992). Quantitative genetics, inclusive fitness, and group selection. *The American Naturalist*, 139(3):540–558.

Queller, D. C. (1997). Cooperators since life began. *The Quarterly Review of Biology*, 72(2).

Queller, D. C. (2000). Relatedness and the fraternal major transitions. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 355(1403):1647–1655.

Queller, D. C. and Strassmann, J. E. (2016). Problems of multi-species organisms: endosymbionts to holobionts. *Biology & Philosophy*, 31(6):855–873.

Rico-Gray, V. and Oliveira, P. S. (2008). *The ecology and evolution of ant-plant interactions*. University of Chicago Press.

Rietkerk, M., Boerlijst, M. C., Van Langevelde, F., HilleRisLambers, R., de Koppel, J. V., Kumar, L., Prins, H. H., and de Roos, A. M. (2002). Self-organization of vegetation in arid ecosystems. *The American Naturalist*, 160(4):524–530.

Roughgarden, J. (1975). Evolution of Marine Symbiosis–A Simple Cost-Benefit Model. *Ecology*, 56(5):1201–1208. Publisher: Wiley Online Library.

Roughgarden, J., Gilbert, S. F., Rosenberg, E., Zilber-Rosenberg, I., and Lloyd, E. A. (2018). Holobionts as units of selection and a model of their population dynamics and evolution. *Biological Theory*, 13(1):44–65.

Sachs, J. L., Ehinger, M. O., and Simms, E. L. (2010a). Origins of cheating and loss of symbiosis in wild bradyrhizobium. *Journal of evolutionary biology*, 23(5):1075–1089.

Sachs, J. L., Mueller, U. G., Wilcox, T. P., and Bull, J. J. (2004). The evolution of cooperation. *The Quarterly review of biology*, 79(2):135–160.

Sachs, J. L., Russell, J. E., Lii, Y. E., Black, K. C., Lopez, G., and Patil, A. S. (2010b). Host control over infection and proliferation of a cheater symbiont. *Journal of evolutionary biology*, 23(9):1919–1927. Publisher: Wiley Online Library.

Sachs, J. L. and Simms, E. L. (2006). Pathways to mutualism breakdown. *Trends in ecology & evolution*, 21(10):585–592.

Sachs, J. L., Skophammer, R. G., and Regus, J. U. (2011). Evolutionary transitions in bacterial symbiosis. *Proceedings of the National Academy of Sciences*, 108(Supplement 2):10800–10807.

Saikkonen, K., Ion, D., and Gyllenberg, M. (2002). The persistence of vertically transmitted fungi in grass metapopulations. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1498):1397–1403.

Saikkonen, K., Wäli, P., Helander, M., and Faeth, S. H. (2004). Evolution of endophyte–plant symbioses. *Trends in plant science*, 9(6):275–280.

Salathé, M., Kouyos, R. D., and Bonhoeffer, S. (2008). The state of affairs in the kingdom of the Red Queen. *Trends in ecology & evolution*, 23(8):439–445.

Sasaki, A., Hamilton, W. D., and Ubeda, F. (2002). Clone mixtures and a pacemaker: new facets of Red-Queen theory and ecology. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1493):761–772.

Schinazi, R. B. (2000). Horizontal versus vertical transmission of parasites in a stochastic spatial model. *Mathematical Biosciences*, 168(1):1–8.

Shapiro, J. W. and Turner, P. E. (2018). Evolution of mutualism from parasitism in experimental virus populations. *Evolution*, 72(3):707–712.

Shapiro, J. W., Williams, E. S., and Turner, P. E. (2016). Evolution of parasitism and mutualism between filamentous phage m13 and escherichia coli. *PeerJ*, 4:e2060.

Simon, B., Fletcher, J. A., and Doebeli, M. (2013). Towards a general theory of group selection. *Evolution*, 67(6):1561–1572.

Smith, J. M. (1964). Group selection and kin selection. *Nature*, 201(4924):1145–1147.

Smith, J. M. (1998). *Evolutionary genetics.* Oxford University Press.

Sober, E. (1984). *The nature of selection: evolutionary theory in philosophical focus.* University of Chicago Press.

Sober, E. and Lewontin, R. C. (1982). Artifact, cause and genic selection. *Philosophy of science*, 49(2):157–180.

Sober, E. and Wilson, D. S. (1999). *Unto others: The evolution and psychology of unselfish behavior.* Harvard University Press.

Su, M., Chen, G., and Yang, Y. (2019). Dynamics of host-parasite interactions with horizontal and vertical transmissions in spatially heterogeneous environment. *Physica A: Statistical Mechanics and its Applications*, 517:452–458.

Szathmáry, E. (2015). Toward major evolutionary transitions theory 2.0. *Proceedings of the National Academy of Sciences*, 112(33):10104–10111.

Szathmáry, E. and Smith, J. M. (1995). *The major transitions in evolution.* WH Freeman Spektrum Oxford, UK:.

Szilágyi, A., Scheuring, I., Edwards, D. P., Orivel, J., and Yu, D. W. (2009). The evolution of intermediate castration virulence and ant coexistence in a spatially structured environment. *Ecology Letters*, 12(12):1306–1316.

Taylor, P. D. and Irwin, A. J. (2000). Overlapping generations can promote altruistic behavior. *Evolution*, 54(4):1135–1141.

Tekwa, E. W., Gonzalez, A., and Loreau, M. (2015). Local densities connect spatial ecology to game, multilevel selection and inclusive fitness theories of cooperation. *Journal of theoretical biology*, 380:414–425.

Thompson, J. N. and Cunningham, B. M. (2002). Geographic structure and dynamics of coevolutionary selection. *Nature*, 417(6890):735–738.

Travis, J. M. J., Brooker, R. W., Clark, E. J., and Dytham, C. (2006). The distribution of positive and negative species interactions across environmental gradients on a dual-lattice model. *Journal of Theoretical Biology*, 241(4):896–902.

Vainstein, M. H., Silva, A. T., and Arenzon, J. J. (2007). Does mobility decrease cooperation? *Journal of theoretical biology*, 244(4):722–728.

Van Baalen, M. and Rand, D. A. (1998). The unit of selection in viscous populations and the evolution of altruism. *Journal of theoretical biology*, 193(4):631–648.

Wade, M. J. (1980). Kin selection: its components. *Science*, 210(4470):665–667.

Wallace, B. (1975). Hard and soft selection revisited. *Evolution*, pages 465–473. Publisher: JSTOR.

Week, B. and Nuismer, S. L. (2021). Coevolutionary arms races and the conditions for the maintenance of mutualism. *The American Naturalist*, 198(2):000–000.

Weeks, A. R., Turelli, M., Harcombe, W. R., Reynolds, K. T., and Hoffmann, A. A. (2007). From parasite to mutualist: rapid evolution of wolbachia in natural populations of drosophila. *PLoS Biol*, 5(5):e114.

Werner, G. D., Cornelissen, J. H., Cornwell, W. K., Soudzilovskaia, N. A., Kattge, J., West, S. A., and Kiers, E. T. (2018). Symbiont switching and alternative resource acquisition strategies drive mutualism breakdown. *Proceedings of the National Academy of Sciences*, 115(20):5229–5234.

Werner, G. D., Cornwell, W. K., Cornelissen, J. H., and Kiers, E. T. (2015). Evolutionary signals of symbiotic persistence in the legume–rhizobia mutualism. *Proceedings of the National Academy of Sciences*, 112(33):10262–10269.

West, S. A., Kiers, E. T., Simms, E. L., and Denison, R. F. (2002). Sanctions and mutualism stability: why do rhizobia fix nitrogen? *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1492):685–694.

Wilkinson, D. M. (1997). The role of seed dispersal in the evolution of mycorrhizae. *Oikos*, pages 394–396.

Wilkinson, D. M. and Sherratt, T. N. (2001). Horizontally acquired mutualisms, an unsolved problem in ecology? *Oikos*, 92(2):377–384.

Wilson, D. S. (1975). A theory of group selection. *Proceedings of the national academy of sciences*, 72(1):143–146.

Wilson, D. S. (2004). What is wrong with absolute individual fitness? *Trends in ecology & evolution*, 19(5):245–248.

Wilson, D. S. and Dugatkin, L. A. (1997). Group selection and assortative interactions. *The American Naturalist*, 149(2):336–351.

Wilson, D. S., Pollock, G. B., and Dugatkin, L. A. (1992). Can altruism evolve in purely viscous populations? *Evolutionary ecology*, 6(4):331–341.

Wilson, D. S. and Sober, E. (1989). Reviving the superorganism. *Journal of theoretical Biology*, 136(3):337–356.

Wilson, D. S. and Sober, E. (1994). Reintroducing group selection to the human behavioral sciences. *Behavioral and brain sciences*, 17(4):585–607.

Yamamura, N., Higashi, M., Behera, N., and Wakano, J. Y. (2004). Evolution of mutualism through spatial effects. *Journal of theoretical biology*, 226(4):421–428.

Zytynska, S. E. and Weisser, W. W. (2016). The natural occurrence of secondary bacterial symbionts in aphids. *Ecological Entomology*, 41(1):13–26. Publisher: Wiley Online Library.