The interferon-rich skin environment regulates Langerhans cell ADAM17 to promote photosensitivity in lupus

Authors: Thomas M. Li[1,†,§], Victoria Zyulina[1,2,§], Ethan S. Seltzer[1,‡,§], Marija Dacic[3,4,5], Yurii Chinenov[3], Andrea R. Daamen[6], Keila R. Veiga[1,7,8,††], Noa Schwartz[1,9,‡‡], David J. Oliver[3], Jose Lora[4,5], Ali Jabbari[10,§§], Yong Liu[11], William D. Shipman[1,12,13, ¶], William G. Ambler[1,7,8], Sarah F. Taber[8,9], Karen B. Onel[8,9], Jonathan H. Zippin[10], Mehdi Rashighi[14], James G. Krueger[10], Niroshana Anandasabapathy[11,12,13], Inez Rogatsky[2,3,4,13], Carl P. Blobel[4,5,15], Peter E. Lipsky[6], Theresa T. Lu[1,2,7,8,13]

Affiliations:
[1]Autoimmunity and Inflammation Program, Hospital for Special Surgery Research Institute, New York, NY 10021
[2]Department of Microbiology and Immunology, Weill Cornell Medical College; New York, USA
[3]David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery Research Institute; New York, USA
[4]Arthritis and Tissue Degeneration Program, Hospital for Special Surgery Research Institute, New York, USA
[5]Physiology, Biophysics, and Systems Biology Program, Weill Cornell Graduate School of Medical Sciences; New York, USA
[6]AMPEL BioSolutions, Charlottesville, VA, USA
[7]Pediatric Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, USA
[8]Department of Pediatrics, Weill Cornell Medical College; New York, USA
[9]Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, NY 10021
[10]Laboratory of Investigative Dermatology, Rockefeller University, New York, USA
[11]Department of Dermatology, Weill Cornell Medical College; New York, USA
[12]Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-PhD Program, Weill Cornell Medical College; New York, USA
[13]Immunology and Microbial Pathogenesis Program, Weill Cornell Graduate School of Medical Sciences; New York, USA
[14]Department of Dermatology, University of Massachusetts Medical School; Worcester, USA
[15]Department of Physiology, Biophysics, and Systems Biology, Weill Cornell Medical College; New York, USA

[§]These authors share first authorship

[†]Current address Icahn School of Medicine at Mount Sinai; New York, USA
[‡]Current address Rockefeller University; New York, USA

45   ††Current address Department of Rheumatology, Maria Fareri Children's Hospital; New York, USA
      ‡‡Current address Department of Medicine (Rheumatology), Montefiore Medical Center/Albert Einstein College of Medicine; New York, USA
      §§Current address Department of Dermatology, University of Iowa Hospital and Clinics; Iowa
50   City, USA
      ¶Current address Department of Dermatology, Yale School of Medicine; New Haven, USA

      Corresponding author
      Theresa T. Lu
55   Hospital for Special Surgery
      535 East 70th Street
      Tel: 212-774-2532
      Email: LuT@HSS.edu

ABSTRACT

60   The autoimmune disease lupus erythematosus (lupus) is characterized by photosensitivity, where

patients develop inflammatory skin lesions with even ambient ultraviolet radiation (UVR)

exposure. The beneficial effects of recently FDA-approved anifrolumab (anti-IFNAR) on lupus

skin disease along with other work support a pathogenic role for IFN-I , but mechanistic

understanding is limited. We have shown Langerhans cell (LC) dysfunction contributes to

65   photosensitivity. Healthy LCs limit photosensitivity via a disintegrin and metalloprotease 17

(ADAM17), a sheddase that releases soluble epidermal growth factor receptor (EGFR) ligands to

limit UVR-induced keratinocyte apoptosis. However, LCs from even non-lesional lupus model

skin show reduced ADAM17 activity and mRNA expression and data point to reduced LC-

mediated protection in human lupus. Here, we asked how the lupus skin environment contributes

70   to LC dysfunction. We show a shared IFN-rich environment in non-lesional skin across human

and murine model systems, that IFN-I inhibits LC ADAM17 activity, and that anti-IFNAR in

lupus models restores LC ADAM17 function and reduces photosensitivity in EGFR and LC

ADAM17-dependent manners. Reactive oxygen species (ROS) can mediate ADAM17 activity,

and we show reduced LC ROS expression in lupus models that is restored by anti-IFNAR. Our

75   findings suggest that IFN-I promotes photosensitivity by causing LC ADAM17 dysfunction and

that anifrolumab ameliorates lupus skin disease at least in part by restoring LC function. This

work provides insight into IFN-I-mediated disease mechanisms, LC regulation, and a mechanism

of action for anifrolumab in lupus.

INTRODUCTION

The autoimmune disease lupus erythematosus (lupus) manifests in part by photosensitivity, a sensitivity to ultraviolet radiation (UVR) whereby even ambient sunlight can trigger the development of inflammatory skin lesions. Patients with lupus skin disease, or cutaneous lupus erythematosus (CLE), can also have the systemic form of the disease, systemic lupus erythematosus (SLE), characterized by inflammation and damage to additional organs such as the kidneys, brain, and heart. In SLE patients, photosensitive skin responses can be associated with worsening systemic autoimmunity and end organ damage. A prominent interferon (IFN) signature indicative of exposure to type I interferon (IFN-I) is found in circulating cells and tissues in lupus (Baechler et al., 2003; Bennett et al., 2003; Catalina et al., 2019; Der et al., 2019; Jabbari et al., 2014; Kirou et al., 2004), and the development and recent FDA approval of anifrolumab (anti-IFNAR1) for SLE highlights the importance of IFN-I in disease pathogenesis. Anifrolumab is especially efficacious for skin disease in SLE patients, consistent with reduced skin lesions in IFNAR-deficient SLE model mice (Nickerson et al., 2013) and an IFN-I signature in both lesional and non-lesional skin (Billi et al., 2022; Der et al., 2017; Der et al., 2019; Reefman et al., 2008; Stannard et al., 2017). These findings suggest a prominent role for IFN-I in lupus skin disease, but mechanisms of IFN-I-mediated pathogenesis remain poorly understood.

We recently reported that Langerhans cells (LCs) can limit UVR-induced skin inflammation by expressing a disintegrin and metalloprotease 17 (ADAM17) which releases membrane-bound epidermal growth factor receptor (EGFR) ligands that then preserve epidermal integrity via EGFR stimulation of keratinocytes (Shipman et al., 2018). Two distinct photosensitive SLE mouse models showed reduced LC *Adam17* mRNA which corresponded to reduced LC ADAM17 sheddase activity, and photosensitivity was reduced by topical EGFR

ligand application. Together, these data suggested that the lack of LC ADAM17-dependent

105 EGFR-ligand sheddase activity contributes to photosensitivity. Non-lesional skin of SLE patients

showed reduced LC numbers along with reduced epidermal EGFR phosphorylation, further

supporting the potential for LC defects to contribute to photosensitivity in lupus. The factors that

lead to LC dysfunction, however, remain to be fully elucidated.

Here, we test the hypothesis that the high IFN-I environment present in even non-lesional

110 lupus skin promotes LC ADAM17 dysfunction to contribute to photosensitivity. We build upon

human data showing an IFN-I signature in non-lesional skin of CLE patients and establish that

the photosensitive MRL/lpr and B6.Sle1yaa models share IFN-I-related and other gene

expression modules with human skin. The inducible photosensitive imiquimod (IMQ) model

mice also upregulated IFN-associated genes in non-lesional skin. IFN-I is sufficient to

115 downregulate healthy murine and human LC ADAM17 activity, and anti-IFN-I receptor (anti-

IFNAR) treatment of lupus mouse models restores LC ADAM17 activity and reduces

photosensitivity in an EGFR and LC ADAM17-dependent manner. Lastly, reactive oxygen

species (ROS) are known to promote ADAM17 activity (Singh et al., 2009), and we show that

UVR-induced LC ROS are reduced in lupus model mice while anti-IFNAR restores ROS

120 expression. Together, our results establish that multiple photosensitive murine lupus models are

similar to human lupus patients in demonstrating IFN-I signatures in non-lesional skin and put

forth a mechanism whereby IFN-I contributes to photosensitivity at least in part by inhibiting LC

ADAM17 activity. These data provide insight into IFN-I-mediated contributions, delineate a

driver of LC dysfunction, and suggest that restoration of LC ADAM17 function is a mechanism

125 of action for anifrolumab in lupus skin disease.

RESULTS

Non-lesional skin in lupus model mice share an IFN-I signature and other functional gene

130   **expression modules with human lupus skin**

As we are interested in the mechanisms that predispose to photosensitivity and to the LC

defects seen in lupus mouse models and human SLE skin (Shipman et al., 2018), we sought to

better understand the environment in such non-lesional skin. Recent studies have suggested a

high IFN-I environment in non-lesional skin of lupus patients based on single cell RNAseq of

135   keratinocytes from SLE patients and skin from CLE patients (Billi et al., 2022; Der et al., 2017;

Der et al., 2019), CLE keratinocyte IFN-κ expression (Stannard et al., 2017), and upregulation of

IFN-stimulated genes (ISGs) such as *MX1* on tissue sections in SLE and incomplete SLE patients

(Lambers et al., 2019; Reefman et al., 2008). Consistent with these findings, our recent analysis

of publicly available bulk transcriptomic datasets from non-lesional DLE skin using Gene Set

140   Variation Analysis (GSVA), an unsupervised gene set analysis approach, showed a proportion of

samples with upregulated IFN gene set expression when compared to healthy controls, although

this effect was modest when compared to the upregulation in lesional skin (Martínez et al.,

2022). Here, we examined an additional non-lesional skin dataset from a CLE cohort with the

discoid form of CLE (DLE) and also non-lesional skin from multiple lupus models to understand

145   shared pathways.

The non-lesional skin has not been previously analyzed and comes from a DLE cohort

whose lesional skin had been shown to have a Th1 signature on microarray when compared to

healthy controls and psoriatic skin (Jabbari et al., 2014). This was among the lesional skin

cohorts that was shown by GSVA to express the IFN gene set at high levels (Martínez et al.,

150   2022). Normal (ie healthy subject) and non-lesional skin exhibited significantly different

Figure 1. **Analysis of a new cohort reinforces a model of an IFN-rich environment and LC and keratinocyte dysfunction in non-lesional DLE skin.** (A-H) Microarray analysis of gene expression from non-lesional skin of DLE patients (n=7) and healthy controls (n=13). In (E-F), samples were analyzed with lesional DLE and psoriasis (PSO) skin samples that were collected as part of the same study and previously published (Jabbari et al., 2014). (A,E) Principal component analysis (PCA) using top 250 genes. (B) Differentially expressed pathways were determined using QuSAGE pathway analysis against Molecular Signatures Database (MSigDB). (C) Volcano plot of differentially expressed genes. Genes from IFN α/β (red), IFNγ (blue) and IRF transcription factor (green) pathways are marked. CD207 (pink) is additionally marked. (D) Heatmap of gene expression in the IFN α/β signaling pathway. (F, G) Gene Set Variation Analysis (GSVA) of gene sets relevant to lupus (Martínez et al., 2022). (H) GSVA using gene sets comprising specific IFN sub-types. (F-H) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by unpaired t-test.

expression profiles (Figure 1A and Table S1). Differential pathway analysis via Quantitative Set

Analysis for Gene Expression (QuSAGE) showed that IFN-I and IFN-γ pathways were among

the most upregulated in the non-lesional skin compared to normal controls (Figure 1B and Table

S2). The IFN-I pathway was comprised of several upregulated IFN-stimulated transcription

170   factors, including IRF 1, 2, 7, 8, and 9 along with multiple IRF targets (ISG15, IFIT1, IFIT3,

MX1, MX2) (Figure 1C-D, S1A). Notably, *CD207*, the gene encoding Langerin and a lineage

marker of LCs, was downregulated (Figure 1C), consistent with accumulating data on LC

dysfunction in lupus (Billi et al., 2022; Martínez et al., 2022; Shipman et al., 2018). A supervised

approach to analyzing normal control versus non-lesional DLE skin gene expression data, then,

175   yielded results consistent with other cohorts showing upregulation of IFN pathways and LC

dysfunction.

We next performed GSVA using previously defined lupus-relevant gene sets (Martínez et

al., 2022) (Table S3). Analysis of the non-lesional samples along with the previously analyzed

lesional and psoriatic samples (Jabbari et al., 2014) reiterated the GSVA results of multiple

180   datasets. The gene expression profiles of the four conditions were significantly different from

each other (Figure 1E). Lesional DLE was characterized by enrichment of immune cell gene sets

including plasma cells, T cells, monocyte/myeloid cells, neutrophils, and plasmacytoid dendritic

cells (pDCs) in conjunction with de-enrichment of LCs when compared to normal controls

(Figure 1F). Lesional DLE samples were also enriched for inflammatory cytokine gene sets

185   including IL12, IL21, IL23, and TNF as well as a highly upregulated IFN gene signature as

previously seen in lesional DLE skin (Figure 1F) (Martínez et al., 2022). Non-lesional skin also

showed upregulation IFN when compared to healthy controls, although this change was dwarfed

when compared to enrichment in lesional skin (Figure 1F). GSVA analysis of just healthy

controls and non-lesional skin in the absence of the high IFN enrichment from lesional skin revealed clear upregulation of the IFN gene set in non-lesional skin as well as de-enrichment of LC and keratinocyte gene sets (Figure 1G). These results were consistent with our previous GSVA analyses of other non-lesional cohorts (Martínez et al., 2022), suggesting along with direct cellular analyses (Billi et al., 2022; Sarkar et al., 2018; Shipman et al., 2018) a loss of epidermal (LC and keratinocyte) function in the context of an IFN-rich proinflammatory environment. The IFN gene set upregulation in non-lesional skin was driven by IFNα, IFNκ, and IFNω gene set upregulation (Figure 1H). GSVA of these previously unanalyzed non-lesional DLE skin samples, then, reinforced previous gene expression analyses and revealed an IFN-I signature in this cohort, all pointing to perturbed epidermal and skin homeostasis even in non-lesional skin.

We examined non-lesional ear skin from MRL/lpr and B6.Sle1yaa SLE mouse models by RNAseq. The photosensitive MRL/lpr model has a *Fas* gene mutation and develops a lupus-like phenotype especially well on the MRL and but not the B6 genetic background. It has reduced LC ADAM17 sheddase function and mRNA, and topical EGFR ligand supplementation ameliorated photosensitivity (Shipman et al., 2018). Although the differences in gene expression between MRL/lpr mice and control MRL-MpJ (MRL/+) mice by RNAseq were less dramatic than those in the human cohort (Figures 1A-D and 2A-D) (Table S4), QuSAGE showed that the IFN-α/β and IFN-γ pathways were among the most highly expressed in MRL/lpr mice (Figure 2B and Table S5). Consistent with the activation of IFN pathways, several IRF transcription factors and their targets were expressed at higher levels in MRL/lpr mice (Figure 2C-D and Figure S1B).

The B6.Sle1yaa model is driven by the Sle1 lupus susceptibility locus derived from lupus-prone NZB2410 mice in combination with the Y chromosome autoimmune accelerator locus

Figure 2. Non-lesional skin of multiple photosensitive murine SLE models recapitulate gene expression patterns of human DLE non-lesional skin. (A-I) Non-lesional skin gene expression from MRL/lpr (LPR) and control MRL/+ (MRL) mice (A-D, F-G) and B6.Sle1yaa and control B6 mice (E, H-I) were analyzed by RNA sequencing. (A, E) PCA using top 250 genes. (B) Differentially expressed pathways determined by QuSAGE (continued on next page)

Figure 2 (continued), pathway analysis against MSigDB. (C) Volcano plot of differentially expressed genes. Genes from IFN α/β (red), IFNγ (blue) and IRF transcription factor (green) pathways are marked. (D) Heatmap showing gene expression in the IFN α/β signaling pathway. (F, H) GSVA of gene sets relevant to lupus, adapted for murine models (Kingsmore et al., 2021). (G, I) GSVA of gene signatures specific to distinct IFN subtypes. (J) Imiquimod (IMQ) model non-lesional and lesional ears and control mice ears were analyzed by qPCR of select IFN-related genes that are upregulated in non-lesional skin of human DLE and MRL/lpr and B6.Sle1yaa models. (A, E-J) Each symbol represents one mouse (A, E-I) or one ear, with non-lesional and lesional ears coming from the same mice (J). (F-J) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by unpaired t-test.

230   whose effects are attributed to *Tlr7* duplication (Pisitkun et al., 2006; Subramanian et al., 2006).

These mice are photosensitive and have reduced LC *Adam17* mRNA expression (Shipman et al.,

2018). As expected, compared to B6 controls, B6.Sle1yaa mice expressed elevated levels of

several lupus related genes from the Yaa locus including Tlr7 (Figure S1C). Although

differences between the B6 and B6.Sle1yaa transcriptomes were modest (Figure 2E)(Table S6),

235   several genes involved in the regulation of and response to IFN were expressed at higher levels,

including *Irf7, Irf9, Isg15, Xaf1, Selp,* and *Ube2l6* (Figure S1D). Although LC numbers were not

different at 10 months of age when this model is fully diseased (Shipman et al., 2018), *Cd207*

expression in younger (5 month old) diseased mice was downregulated when compared to B6

controls (Figure S1D), similar to the findings in CLE skin (Figure 1C). By supervised gene set

240   analysis, both the MLR/lpr and B6.Sle1yaa murine models showed similarities to human non-

lesional skin.

By GSVA of lupus-relevant murine gene sets (Table S7), parallels between the MLR/lpr

and B6.Sle1yaa murine models and human lupus were even more dramatic. MRL/lpr mice

showed increased enrichment of a number of inflammatory cell and cytokine gene sets including

245   T cells, myeloid cells, neutrophils, pDCs, IFN, IL1, IL21, and TNF (Figure 2F). However, in

contrast to human DLE, non-lesional skin of MRL/lpr mice was enriched for the keratinocyte

gene signature. LC numbers in the skin of MRL/lpr mice are not downregulated when compared

to MRL/+ controls (Shipman et al., 2018), which was reflected in the absence of a decreased

dendritic cell signature. The IFN gene set upregulation was broad, with significant enrichment of

250   gene signatures induced by stimulation from multiple IFNs (Figure 2G). B6.Sle1yaa mice

showed less dramatic differences than the MRL/lpr model, but also showed upregulation of the

IFN gene set (although there was not a significant contribution from any one IFN signature) and

downregulation of the DC gene set, potentially reflective of decreased LC function (Figure 2H-1). Together, our results showed a number of parallels to the human system, including an IFN signature in these lupus models.

We also tested for expression of ISGs in the imiquimod (IMQ) lupus model. This model is induced by 4-5 weeks of topical IMQ, a TLR 7/8 agonist, resulting in autoantibody production, enlarged spleen and lymph nodes, nephritis, and photosensitivity characteristic of SLE (Ambler et al., 2022; Goel et al., 2020; Yokogawa et al., 2014). The model can be induced in B6 mice, allowing for the use of transgenic models to assess for effects on lupus features without having to cross onto a different genetic background. We applied IMQ to the right ear to assess ISGs in both the painted right "lesional" and unpainted left "non-lesional" ears. We performed qPCR for a set of transcripts (*Gbp2, Ifi35, Ifit1, Ifit3, Ifit2, Irf7, Irf8, Isg15, Stat1,* and *Xaf1)* that were upregulated across human and murine non-lesional skin datasets using our datasets (Figures S1A, B, D) and that in (Billi et al., 2022) (Figure 2J). Most of the genes we examined showed upregulation in the "lesional" ear, including transcripts for IRF target genes *Irf7, Irf8* and *Stat1*, and the apoptosis marker *Xaf1* potentially reflecting tissue damage along with an IFN-rich environment. Several transcripts also showed trends toward upregulation in non-lesional skin when compared to mice that were not treated with IMQ (Figure 2J). Notably, *Isg15*, the only transcript that we examined that was upregulated across all the datasets, was also upregulated in the non-lesional ears of the IMQ mice. ISG15 has been shown to limit over-amplification of IFN-I effects, and its deficiency predisposes to skin inflammation among other findings (Martin-Fernandez et al., 2020; Zhang et al., 2015), suggesting that upregulation of *Isg15* may reflect an attempt to minimize IFN-I-mediated inflammation. These results showed that non-lesional skin

275   from human lupus and 3 murine models were characterized by upregulation of IFN-associated

transcripts, albeit at lower levels than in lesional skin.

Turning our attention to common pathways in human and murine lupus non-lesional skin,

we performed ranked Gene Set Enrichment Analysis (GSEA) separately for DLE and MRL/lpr

datasets using negative log-transformed p-values as a ranking variable, constructed pathway

280   similarity networks, and compared selected pathways (FDR<0.005) in humans and mice.

Consistent with QuSAGE analysis (Figure 1B and 2B), IFN-related pathways were enriched in

human DLE and MRL/lpr non-lesional skin (nodes 3, 4, 8), along with several other pathways

often associated with lupus (Figure 3A): T helper pathway (nodes 26); natural killer cell-

mediated cytotoxicity (nodes 28); Toll-like receptor pathways (nodes 36,46); JAK-STAT

285   signaling pathway (node 43); and multiple pathways associated with complement activation

(nodes 6, 23, 29, 34, 52, 53, 58, 61, 62, 64, 75, 88, 94, 95). For the full list of pathways identified

by ranked GSEA, see Table S8. Direct comparison of differentially expressed genes in DLE and

MRL/lpr skin (Figure 3B) revealed a gene signature typical of nucleic acid-driven

interferonopathies dominated by IFN-stimulated transcription factors (IRF1,3,9 and STAT1),

290   IFN-induced viral restriction factors (BST2, SAMHD1, DDX60, IFIH1, ZBP1), neutrophilic

markers (HMOX1, S100A8, S100A9, LCN2, RSAD2), and apoptosis-related genes (XAF1,

TNFSF10). Several keratins (KRT16, 6B and 34) that are rapidly induced upon skin damage and,

in turn, downregulate inflammatory responses (Lessard et al., 2013), were highly expressed in

both human CLE and MRL/lpr skin, likely reflecting disturbed epidermal homeostasis.

295   Comparisons of the GSVA analyses of non-lesional skin among the datasets (Figure 3C)

echoed the GSEA analysis in emphasizing the prominent enrichment of IFN in human CLE and

both MRL/lpr and B6.Sle1yaa lupus-prone mice. In addition, de-enrichment of the LC/DC gene

REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM
REACTOME_INNATE_IMMUNE_SYSTEM
REACTOME_INTERFERON_SIGNALING
REACTOME_INTERFERON_ALPHA_BETA_SIGNALING
REACTOME_ADAPTIVE_IMMUNE_SYSTEM
KEGG_CYTOKINE_CYTOKINE_RECEPTOR_INTERACTION
REACTOME_INTERFERON_GAMMA_SIGNALING
REACTOME_SIGNALING_BY_INTERLEUKINS
REACTOME_NEUTROPHIL_DEGRANULATION
KEGG_PRIMARY_IMMUNODEFICIENCY
REACTOME_CHEMOKINE_RECEPTORS_BIND_CHEMOKINES
KEGG_CHEMOKINE_SIGNALING_PATHWAY
KEGG_CELL_ADHESION_MOLECULES_CAMS
REACTOME_LEISHMANIA_INFECTION
KEGG_HEMATOPOIETIC_CELL_LINEAGE
BIOCARTA_THELPER_PATHWAY
KEGG_LEISHMANIA_INFECTION
KEGG_NATURAL_KILLER_CELL_MEDIATED_CYTOTOXICITY
REACTOME_HEMOSTASIS
KEGG_VIRAL_MYOCARDITIS
REACTOME_COSTIMULATION_BY_THE_CD28_FAMILY
REACTOME_INTERLEUKIN_10_SIGNALING
REACTOME_TOLL_LIKE_RECEPTOR_CASCADES
REACTOME_GPCR_LIGAND_BINDING
REACTOME_CLASS_A_1_RHODOPSIN_LIKE_RECEPTORS
BIOCARTA_GRANULOCYTES_PATHWAY
KEGG_JAK_STAT_SIGNALING_PATHWAY
REACTOME_INTERLEUKIN_2_FAMILY_SIGNALING
KEGG_TOLL_LIKE_RECEPTOR_SIGNALING_PATHWAY
REACTOME_ANTIVIRAL_MECHANISM_BY_IFN_STIMULATED_GENES
REACTOME_SIGNALING_BY_GPCR
56 REACTOME_PEPTIDE_LIGAND_BINDING_RECEPTORS
58 REACTOME_FCGAMMA_RECEPTOR_FCGR_DEPENDENT_
   PHAGOCYTOSIS
61 REACTOME_PLATELET_ACTIVATION_SIGNALING_AND_AGGREGATION
63 REACTOME_ANTI_INFLAMMATORY_RESPONSE_FAVOURING_
   LEISHMANIA_PARASITE_INFECTION
65 REACTOME_INTERLEUKIN_4_AND_INTERLEUKIN_13_SIGNALING
69 KEGG_APOPTOSIS
72 BIOCARTA_NEUTROPHIL_PATHWAY
REACTOME_METAL_SEQUESTRATION_BY_ANTIMICROBIAL_PROTEINS
BIOCARTA_MHC_PATHWAY
BIOCARTA_HIVNEF_PATHWAY
BIOCARTA_MONOCYTE_PATHWAY
REACTOME_CREATION_OF_C4_AND_C2_ACTIVATORS
KEGG_B_CELL_RECEPTOR_SIGNALING_PATHWAY
REACTOME_CASPASE_ACTIVATION_VIA_DEATH_RECEPTORS_IN_
   THE_PRESENCE_OF_LIGAND
94 REACTOME_INITIAL_TRIGGERING_OF_COMPLEMENT

Figure 3. Analysis of human and murine lupus models show shared gene expression pathways. (A) Pathway similarity network built for the most enriched pathways (adjusted p<0.005) in DLE non-lesional skin determined by ranked GSEA (complete list in Table S8). Colors indicate network community clusters identified by the walktrap algorithm. Red borders indicate pathways that were also enriched in LPR mice (listed). (B) Comparison of gene expression changes in DLE vs MRL/lpr mice. (C) Heatmap of GSVA scores for shared gene sets between non-lesional skin from DLE patients, LPR mice, and B6.Sle1yaa mice. Asterisks indicate significant differences in GSVA scores compared to controls for each dataset.

set was present in both human DLE and B6.Sle1yaa mouse non-lesional skin (Figure 3C).

Multiple approaches to analyze gene expression of non-lesional skin from human DLE and

murine lupus models, then, showed upregulation of IFN-related transcripts and disturbance of

LC and epidermal homeostasis. These data are consistent with the idea that non-lesional skin is

primed for photosensitive responses and that LCs sit within an IFN-rich environment in both

human lupus and multiple murine models that may cause LC dysfunction.

IFN-I inhibits LC ADAM17 function

We next examined the effects of IFN-I on LC ADAM17 function and expression. As

previous (Shipman et al., 2018), we assessed LC ADAM17 function by quantifying UVR-

induced cell surface TNFR receptor 1 (TNFR1) shedding. We confirmed that UVR treatment

performed on a mixture of digested epidermal cells induced LC loss of surface TNFR1 in an LC

ADAM17-dependent manner (Figure S2A-B), similar to our results with isolated LCs (Shipman

et al., 2018). IFN-κ, an IFN-I that is highly expressed by lupus keratinocytes (Tsoi et al., 2019),

reduced LC ADAM17 activity by nearly 40% (Figure 4A). Using a validated anti-murine

ADAM17 antibody (Lora et al., 2021) (Figure S2A), we observed that IFN-I did not reduce

ADAM17 cell surface levels by the same magnitude (Figure 4B) (Lora et al., 2021), and LC

viability was not affected (Figure 4C). The IFN-κ-driven inhibition of ADAM17 activity was

abrogated by the JAK1/3 inhibitor tofacitinib without altering ADAM17 protein levels (Figure

4D-E) or cell viability (Figure 4F), supporting an IFNAR-mediated inhibition of LC ADAM17

sheddase activity rather than loss of ADAM17 cell surface expression. Application of IFN-κ to

the skin of wild-type mice reduced LC ADAM17 activity by 57% (Figure 4G) while cell surface

ADAM17 was reduced by 9% (Figure 4H). Together, these data indicate that IFN-I is sufficient

Figure 4. IFN-I inhibits LC ADAM17 sheddase activity. (A-K) Epidermal cell suspensions were treated ex vivo or mice were treated in vivo with IFN-I prior to assaying for UVR-induced LC ADAM17 sheddase activity, ADAM17 levels, and numbers. Cells from WT mice were treated with IFNκ or vehicle (A-C) with and without tofacitinib (D-F); IFN κ or vehicle was applied topically to ears of WT mice 16-20 hours prior to generation of epidermal cell suspensions (G-H); suction blister epidermal cell suspensions from healthy human donors were treated with IFNβ or vehicle (I-K). (A, D, G, I) LC ADAM17 sheddase activity as indicated by the extent of UVR-induced cell surface TNFR1 loss. Percent change in cell surface TNFR1 mean fluorescence intensity (MFI) after UVR (left); relative LC ADAM17 activity calculated by normalizing TNFR1 loss to that of vehicle controls (right). In (I), lines connect samples from the same donor. (B, E, H, J) Cell surface ADAM17 levels as indicated by MFI relative to that of vehicle controls. (C, F, K) Relative numbers of LCs per sample after indicated treatments. (A-K) Each symbol represents cells from a single mouse or donor, bars represent average values, and error bars are SD. n=3-10 per condition over 3-5 independent experiments. *p<0.05, **p<0.01, ***p<0.001, n.s.=not significant by unpaired t-test.

to downregulate LC murine ADAM17 function and does so primarily at the level of sheddase activity ex vivo and in vivo.

350   We also asked how IFN-I affects human LCs. While LCs can be obtained from skin biopsies or discarded tissues associated with surgical procedures, these tissues often require lengthy enzymatic digestion to dissociate the cells (Shipman et al., 2018), which has the potential to disrupt cell function. We found that epidermal roofs yielded from suction blistering could be digested for a short amount of time to obtain an epidermal cell mixture containing LCs (Figure

355   S3) (Strassner et al., 2017). Addition of IFN-I reduced LC ADAM17 activity by 62% without altering ADAM17 expression or LC viability (Figure 4I-K). These results together showed that IFN-I can inhibit LC ADAM17 activity in both murine and human LCs and supported the idea that high levels of IFN-I could potentially contribute to LC ADAM17 dysfunction in disease.

360   **IFNAR is important for LC ADAM17 dysfunction in multiple lupus models**

     We next asked whether IFN-I was an important mediator of LC ADAM17 dysfunction in lupus models. Consistent with previous findings (Shipman et al., 2018), MRL/lpr mice showed a 42% reduction in LC ADAM17 sheddase activity when compared to control MRL/+ mice (Figure 5A). LC ADAM17 protein levels, however, were less affected, showing a 21% reduction

365   (Figure 5B). Intraperitoneal (IP) anti-IFNAR treatment at 500 μg/dose restored LC ADAM17 activity in MRL/lpr mice (Figure 5A) without increasing ADAM17 protein expression (Figure 5B). A lower 125 μg/dose of anti-IFNAR had similar effects in rescuing LC ADAM17 sheddase function (Figure 5C-D). These results together suggested that IFN-I is important for inhibiting LC ADAM17 sheddase activity in MRL/lpr mice.

Figure 5. Anti-IFNAR restores LC ADAM17 activity in multiple lupus models. (A-I) MRL/lpr (A-D), B6.Sle1yaa (E-F), and IMQ (G-I) lupus model mice and their controls were treated twice with anti-IFNAR or isotype control at indicated doses over 6 days prior to collection of non-lesional epidermal cells. (A, C, E, G) UVR-induced LC ADAM17 sheddase activity as in Figure 4. (B, D, F, H) Relative cell surface ADAM17 levels. (I) Relative LC numbers per sample. (A-I) Each symbol represents a mouse, bars represent average values, and error bars are SD. n=3-8 per condition over 3-4 independent experiments. *p<0.05, **p<0.01, ***p<0.001, n.s.=not significant by paired (A, C, E (left)) or unpaired ((A, C, E (right)), B, D, F-I) t-test.

LCs from the B6.Sle1yaa model showed a 63% reduction in ADAM17 activity when compared to control mice, without a corresponding change in ADAM17 protein levels (Figure 5E-F). Similar to MRL/lpr mice, anti-IFNAR restored LC ADAM17 activity in B6.Sle1yaa mice without increasing ADAM17 protein expression (Figure 5E-F).

385   IMQ model mice showed reduced LC ADAM17 sheddase activity without altering cell surface ADAM17 protein levels (Figure 5G-H), similar to data from MRL/lpr and B6.Sle1yaa models. Interestingly, unlike the MRL/lpr and B6.Sle1yaa models where LC numbers were unchanged from controls (Shipman et al., 2018), skin LC numbers in the IMQ model were reduced by 35% (Figure 5I), echoing findings in human skin (Shipman et al., 2018). Similar to

390 the MRL/lpr and B6.Sle1yaa models, anti-IFNAR restored LC ADAM17 activity in IMQ mice (Figure 5G). Together, the sufficiency of IFN-I in inhibiting LC ADAM17 activity and the restoration of LC ADAM17 activity by anti-IFNAR across 3 lupus models suggest that the IFN-I-rich environment drives LC ADAM17 dysfunction in non-lesional lupus skin.

395 **Anti-IFNAR reduces photosensitivity in an EGFR and LC ADAM17-dependent manner**

       We asked whether the anti-IFNAR-mediated restoration of LC ADAM17 function also reduced photosensitive skin responses. Anti-IFNAR treatment at 500 μg/dose reduced ear swelling in MRL/lpr mice (Figure 6A-B) but did not reduce epidermal permeability, a readout of skin barrier function (Figure 6C). Furthermore, in control MRL/+ mice, anti-IFNAR actually

400 increased UVR-induced ear swelling and epidermal permeability (Figure 6B-C), suggesting that anti-IFNAR increased photosensitivity in non-lupus mice. These results echoed findings by other groups that IFNAR deficiency in non-lupus B6 mice increased photosensitivity (Sontheimer et al., 2017) and reduced skin wound re-epithelialization (Gregorio et al., 2010), suggesting that the

Figure 6. Anti-IFNAR reduces photosensitivity in EGFR- and LC ADAM17- dependent manners. (A-I) MRL/lpr and (J-K) LCAd17 mice and controls were treated according to schematics in (A, G, J) and non-lesional ears were harvested. (B, D, H) Ear thickness. (C, E, I) Epidermal permeability as indicated by toluidine blue retention. (F, K) Neutrophils and monocytes per ear. (B-F, H-I, K). n=4-10 per condition over 4-6 independent experiments.

410 *p<0.05, **p<0.01, ***p<0.001, n.s.=not significant by paired (B, D, H) and unpaired (C, E, F, I, K) t-test.

500 μg anti-IFNAR dose was disrupting intrinsic skin functions and potentially obscuring the effects of LC ADAM17 activity restoration. Consistent with this idea, anti-IFNAR at 125 μg/dose still worsened MRL/+ ear swelling but no longer increased epidermal permeability

415 (Figure 6D-E), suggesting a decreased pathologic effect on homeostatic skin functions. This lower dose, which was sufficient to restore LC ADAM17 activity in MRL/lpr mice (Figure 5C), was sufficient to reduce both UVR-induced skin swelling and epidermal permeability (Figure 6D-E) and limit neutrophil and monocyte infiltration after UVR exposure (Figure 6F). At 125 μg/dose, then, anti-IFNAR rescued LC ADAM17 function and reduced UVR-induced skin

420 swelling, epidermal permeability, and inflammatory infiltrates in MRL/lpr mice.

We sought to understand the extent to which the anti-IFNAR-driven reduction in photosensitivity reflected the rescue of LC ADAM17 activity. As LC ADAM17 limits photosensitivity by stimulating EGFR signaling (Shipman et al., 2018), we blocked EGFR activity upon anti-IFNAR treatment of MRL/lpr mice (Figure 6G). EGFR blockade with the

425 small molecule inhibitor PD16 abrogated the anti-IFNAR-mediated reduction in ear swelling and epidermal permeability (Figure 6H-I), suggesting that anti-IFNAR requires EGFR signaling to ameliorate photosensitivity in MRL/lpr mice. This is consistent with the idea that anti-IFNAR reduces photosensitivity by restoring LC ADAM17 function.

In the IMQ model, anti-IFNAR also reduced inflammatory cell infiltrate in UVR-treated

430 skin (Figure 6J-K). In IMQ-treated LCAd17 mice lacking ADAM17 in LCs (Shipman et al., 2018), however, anti-IFNAR failed to reduce neutrophil and monocyte accumulation (Figure 6K), suggesting that the ameliorative effects of anti-IFNAR on UVR-induced inflammation are dependent on LC ADAM17. That anti-IFNAR reduced photosensitive responses across two

lupus models depended on EGFR signaling and LC ADAM17 provides strong support for the

idea that IFN-I contributes to photosensitivity at least in part by promoting LC ADAM17

dysfunction. These results also suggest that the beneficial effects of anifrolumab on SLE skin

may be attributable to the improvement of LC ADAM17 activity.

IFN-I inhibits UVR-induced LC ROS expression

UVR has been shown to trigger ADAM17 activity in fibroblasts via ROS generation

(Singh et al., 2009), and we asked if IFN-I modulation of ROS was associated with changes in

Figure 7. LC ROS is reduced in lupus model mice and restored by anti-IFNAR. (A-B) Epidermal cell suspensions from indicated mice were exposed to UVR or not and LC ROS content assayed by the ROS indicator CM-H2DCFDA. (A) WT mice; representative histograms (left) and LC ROS content relative to non-exposed condition (right). Dashed "control" line indicates signal without CM-H2DCFDA. (B) Control (-IMQ) and IMQ-treated mice (+IMQ) were treated as in Figure 5G with anti-IFNAR or isotype IgG prior to harvest of non-lesional skin; histograms (left) and fold change in LC ROS with UVR exposure (right). Each symbol

represents one mouse, bars represent average values, and error bars are SD. n=5-6 per condition over 4-5 experiments. *p<0.05, **p<0.01, n.s.=not significant by unpaired t-test.
LC ADAM17 activity. We observed the expected UVR-induced ROS generation in LCs from

WT skin (Figure 7A). Skin from IMQ-treated mice showed a reduction in UVR-induced

455   LC ROS generation when compared to control skin, and anti-IFNAR treatment of IMQ mice

restored UVR-driven ROS generation (Figure 7B). These results suggest that IFN-I could

potentially limit UVR-induced LC ROS to inhibit LC ADAM17 function.

       Together, our results suggested a model whereby the IFN-I-rich tissue microenvironment

in lupus skin causes dysfunction of UVR-induced LC ADAM17, leading to photosensitivity.

460   Anti-IFNAR restores this axis, suggesting that the therapeutic effects of anifrolumab on SLE

skin could, in part, be via restoration of LC function.

DISCUSSION

465   Here, we establish that multiple photosensitive lupus models resemble human lupus in showing upregulated IFN-I modules or IFN-regulated genes in non-lesional skin and that IFN-I in these lupus models causes LC ADAM17 dysfunction which contributes to photosensitive skin responses. These results suggest that the IFN-rich environment in lupus skin promotes photosensitivity by causing Langerhans cell ADAM17 dysfunction and that the beneficial effects

470   of anifrolumab on skin disease in SLE patients, in part, reflects restoration of LC function. Our results also suggest that photosensitivity in the MRL/lpr, B6.Sle1yaa, and IMQ lupus models shares pathogenic mechanisms with human lupus, underscoring the utility of these models for the study of lupus skin pathophysiology.

       How the IFN-I-rich environment in lupus skin contributes to photosensitive responses has

475   not been well understood. Kahlenberg and colleagues have shown that IFN-I can potentiate IL-6 expression and apoptosis of human keratinocytes (Sarkar et al., 2018; Stannard et al., 2017). Keratinocytes are likely a major local source of IFN-I to modulate LC function in non-lesional skin, as LCs are positioned in the suprabasal layer of the epidermis near basal and spinous keratinocytes that demonstrate high IFN signaling (Billi et al., 2022). PDCs may also regulate

480   LCs, as pDCs contribute to the IFN-rich environment in lupus skin (Furie et al., 2019; Karnell et al., 2021). Our results showing that anti-IFNAR restores LC ADAM17 function and reduces UVR-induced inflammation in an EGFR and LC ADAM17-dependent manner suggest that IFN-I contributes to photosensitivity by causing LC ADAM17 dysfunction. The accumulating evidence points to a pathogenic feed-forward loop between LCs and keratinocytes whereby

485   keratinocyte IFN contributes to LC ADAM17 dysfunction and LC ADAM17 dysfunction contributes to keratinocyte dysfunction and ultimate loss. It should be noted, however, that the

importance of IFN-I-mediated LC ADAM17 dysfunction does not rule out IFN-I-mediated keratinocyte-intrinsic mechanisms but points to multiple players in the ecosystem that maintain skin health.

490   Our findings showing that anti-IFNAR treatment could increase UVR-induced skin inflammation in control MRL/+ mice underscores the idea that excessive IFN-I contributes to skin pathogenesis in lupus. IFNAR is important for re-epithelialization during wound healing (Gregorio et al., 2010) and our data support the idea that our treatments should not compromise protective levels of IFNs. This notion is supported by the clinical findings that patients with high

495   IFN-I scores responded to anifrolumab (Morand et al., 2020) and, when comparable, with a greater effect size than patients with lower IFN-I scores (Furie et al., 2017). Understanding the regulation and sources of the "excess" IFN-I in lupus will help to further develop therapies that ameliorate disease while preserving skin-intrinsic functions.

Our results extend our understanding of ADAM17 regulation as a contributor to disease.

500   Increased ADAM17 activity can promote inflammation, such as by release of TNF or EGFR ligands within inflamed joints or lupus kidneys (Issuree et al., 2013; Qing et al., 2018) but loss of ADAM17 activity compromises barrier surfaces (Blaydon et al., 2011; Franzke et al., 2012). Here we identify IFN-I as a negative regulator of ADAM17 in LCs and our data suggest a scenario whereby IFN-I reduces LC ROS generation to inhibit ADAM17 function. Further

505   investigation will better delineate the locus and importance of LC ROS generation as well as contributions of other molecules such as the inactive Rhomboids (iRhoms) that regulate ADAM17 activity (Geesala et al., 2019).

High IFN-I states also include viral infections, genetic interferonopathies, and autoimmune diseases rheumatoid arthritis and dermatomyositis (Muskardin and Niewold, 2018), in which

510 photosensitivity is found in the former and is a disease hallmark in the latter (Werth et al., 2004; Wysenbeek et al., 1989). Autoimmune disease-like manifestations that include photosensitivity are among the immune-related adverse events in cancer patients treated with checkpoint inhibitors (Brahmer et al., 2018), a situation where distinct IFN-I-mediated processes may have specific roles in promoting anti-tumor immunity and treatment resistance (Benci et al., 2019).

515 Our findings will benefit patients with lupus and a wide spectrum of diseases.

MATERIALS AND METHODS

Mice

8-16 week old mice were used unless otherwise specified. Mice were sex and age-matched. For C57Bl/6 mice, LCAd17 (Langerin-cre; Adam17 f/f) mice, and wildtype littermate controls, males and females were used. Unless otherwise indicated, for MRL/+ and MRL/lpr mice, females were used at 8-12 weeks, while males were used at 12-16 weeks as females present with a disease phenotype earlier than males. For B6.Sle1yaa mice and their B6 controls, only males were used because the model depends on the autoimmune accelerator locus located on the Y chromosome. B6 mice were either bred in-house or purchased from Jackson Laboratory (Bar Harbor, ME). MRL/+, MRL/lpr mice, B6.Sle1yaa mice, and their B6 controls were either bred in-house or purchased from Jackson Laboratory. LCAd17 mice (Shipman et al., 2018) were bred in-house. All animal procedures were performed in accordance with the regulations of the Institutional Animal Care and Use Committee at Weill Cornell Medicine (New York, NY)

Mouse treatments

For IFN-κ painting, 37.5 ng IFN-κ (R&D Systems, Minneapolis, MN) was dissolved in DMSO (Sigma Aldrich, St. Louis, MO) and was applied topically onto each ear (18.75 ng on dorsal and ventral sides, each). For anti-IFNAR1 treatments, mice were injected IP with 500 µg/dose or 125 µg/dose of anti-IFNAR1 (MAR1-5A3) or isotype control (MOPC-21) (Bio X Cell, West Lebanon, NH). For UVR treatments, mice were exposed to 250 J/m^2 (25 mJ/cm^2)daily for four consecutive days using a bank of four FS40T12 sunlamps that emitted a combination of UVA and UVB radiation, similar to described (Shipman et al., 2018). Measurement using UVA and UVB meters revealed that these lamps emit 2.1mW/cm2 UVA and 0.8mW/cm2 UVB. To

measure ear swelling after UVR exposure, a caliper (Mitutoyo, Aurora, IL) was used to take three measurements throughout each ear, and the reported value is the average of both ears. Each measurement was taken 22-24 hours after the previous measurement. For the IMQ-induced lupus mouse model, mice were painted on the dorsal and ventral sides of both ears with 5% imiquimod cream (42 mg/ear/mouse) 3x/week for 5-6 weeks (Yokogawa et al., 2014) prior to, and continued through experiments. Indicated mice received 2.95 mg of the irreversible EGFR inhibitor PD168393 (0.74 mg on dorsal and ventral sides, each) (Selleck Chem, Pittsburgh, PA). Mice were randomized to treatments.

Human subjects

For microarray analysis, seven discoid CLE patients were examined. Both lesional and non-lesional skin were collected and the gene expression data from lesional tissue of these patients have been published, and patient characteristics are described (Jabbari et al., 2014). Healthy controls include the three that were previously described (Jabbari et al., 2014). All human tissue collection and research use adhered to protocols approved by the Institutional Review and Privacy Boards at the Rockefeller University (IRB# AJA-00740) and New York University (IRB# 10-02117), where participants signed written informed consents.

For suction blistering, 1 male and 3 female healthy participants between the ages of 23 and 55 participated. All human tissue collection and research use adhered to protocols approved by the Institutional Review and Privacy Board at the Hospital for Special Surgery (IRB# 2019-1998), where participants signed written informed consents.

Human skin cell collection and preparation

Suction blistering was performed as previously described (27). Eight 5 mm suction blisters were generated with a negative pressure instrument (NP-4 model, Electronic Diversities, Finksburg, MD) on the arm of healthy donors over 30-60 minutes. The blister fluid was then collected by aspiration, and epidermal roofs were collected. After collection, epidermal roofs from suction blisters were digested with dispase (2.42 U/mL; Thermo Fisher Scientific, Waltham, MA), collagenase type II (616 U/mL; Worthington Biochemical Corporation, Lakewood, NJ), and DNAseI (80 µg/mL; Sigma Aldrich, St. Louis, MO) for 40 minutes to generate single cell suspensions that were used for the ADAM17 sheddase assay.

Skin cell collection and flow cytometry staining and quantification

For murine epidermal single cell suspensions, skin on the trunk was excised and subcutaneous fat scraped off. Skin was incubated in dispase (2.42 U/mL) at 37°C. The epidermis was then gently peeled, finely minced, digested in type II collagenase (616 U/mL) and DNAse I (80 µg/mL).

For collection of ear skin, ears were cut along the cartilage ridge, and the dorsal and ventral sides were manually peeled. Ear skin was then finely minced and digested with a buffer containing dispase, type II collagenase, and DNAse I as described above. For murine LC TNFR1 staining for the ADAM17 sheddase activity assay (see below), epidermal cells were preincubated with Fc block and then stained so that we could gate on CD45+, CD3-, CD11c+, I-Ab+ for B6 mice and I-Ak+ for MRL mice, and EpCAM+ LCs and assess for TNFR1. LCs were sometimes stained with rabbit-anti-ADAM17 (Lora et al., 2021) followed by donkey anti-rabbit IgG (Jackson Immunoresearch, West Grove, PA). Monocytes and T cells were identified by gating for CD45+, CD11b+, Ly6C+, and CD45+, CD11c-, I-Ak/I-Ab-, and CD3+ cells, respectively.

4',6-diamidino-2-phenylindole dihydrochloride (DAPI) was used to exclude dead cells and debris (Sigma Aldrich, St. Louis, MO). All antibodies are from Biolegend, San Diego, CA unless otherwise indicated.

For human LC TNFR1 staining for the ADAM17 sheddase activity assay, single cell

590   suspensions of the epidermal roofs of suction blisters and blister fluid were preincubated with Fc block and subsequently stained for CD45+, CD11c+, CD14-, CD3- CD19-, CD56-, CD66b-, EpCAM+, and HLA-DR+ to identify LCs. DAPI was used to exclude dead cells and debris, and TNFR1 was also stained to assess ADAM17 activity. All antibodies are from Biolegend, San Diego, CA unless otherwise indicated. Cells were analyzed using a FACS Canto or FACS

595   Symphony A3 (BD Biosciences, San Jose, CA) and data were analyzed with FlowJo V10 Software (TreeStar, Ashland, OR).

For mean fluorescent intensity (MFI) calculations of TNFR1, each sample was assayed in triplicate, and the mean of the triplicates was recorded unless otherwise specified. The MFI of TNFR1 was divided by the MFI for the isotype control to generate a relative MFI. The percent

600   TNFR1 MFI change was calculated by setting the relative MFI of the control sample to 1 and normalizing the value from the experimental samples to that control. In experiments with more than one control sample, the control values were averaged, and the experimental samples were each normalized to the averaged value.

For antibodies used, please see Table S9.

Ex vivo ADAM17 sheddase activity assay

For murine skin, cell suspensions of epidermis were suspended at 200,000-500,000 cells/mL medium. For MRL/LPR and B6.Sle1yaa skin, medium was RPMI with 2% FBS, HEPES buffer,

L-glutamine, penicillin-streptomycin (all Thermo Fisher Scientific, Waltham, MA). For IMQ

610 mice skin, medium was phenol red-free RPMI (Corning, Edison, New Jersey), no serum, with

HEPES buffer, L-glutamine, penicillin-streptomycin. Cells were exposed to 500 (LPR and

B6.Sle1yaa model mice) or 250 (IMQ model mice) J/m^2 UVR, similar to previous (Shipman et

al., 2018), incubated for 45 minutes, and then stained for cell surface markers described above.

For IFN-κ treatment, 3.125 ng/mL IFN-κ (R&D Systems, Minneapolis, MN) was added to the

615 epidermal cell suspension that were incubated for 16-20 hours prior to UVR exposure. Indicated

cells were treated with 1nM tofacitinib (Selleck Chem, Pittsburgh, PA) during IFN-κ incubation.

For human skin, cells were incubated for 1 hour with recombinant human IFN-β (35.7

ng/mL; activity=2.8x10^8 UI/mg) and then exposed to 500 J/m^2 UVR as above. Cells were further

incubated for 45 minutes and collected for preincubation with Fc block. Cells were then stained

620 for cell surface markers described above.

Epidermal permeability assay

Epidermal barrier function was assessed as described (Shipman et al., 2018). Ear skin was

dehydrated and rehydrated in graded methanol and then incubated in 0.1% toluidine blue dye

625 (Sigma Aldrich, St. Louis, MO). The dye was extracted with a buffer consisting of 95%

methanol, 2.5% sulfuric acid, and 2.5% water and quantified with a colorimeter at 620 nm. Dye

concentration was calculated using a standard curve.

Ex vivo ROS assay

630 Epidermal cell suspensions in phenol red-free RPMI were exposed to 250 J/m2 UVR and

incubated with the ROS indicator 5-(and-6)-chloromethyl-2',7'-dichlorodihydrofluorescein

diacetate, acetyl ester (CM-H2DCFDA) (Invitrogen, Waltham, MA) at 10uM for 45 minutes
prior to cell surface marker staining for flow cytometry.

RNA extraction

Female MRL/+ and MRL/lpr 12-14 weeks old and 12 week old B6 and B6.Sle1yaa male mice
were used for RNA sequencing. RNA was extracted from ear skin using an RNEasy Kit (Qiagen,
Germantown, MD) and quality confirmed on a BioAnalyzer 2100 (Agilent Technologies, Santa
Clara, CA). For RT-qPCR of IMQ-treated mice, RNA was extracted with Trizol (Life

640   Technologies) following the manufacturer's protocol.

RNA sequencing

Libraries were prepared with non-stranded and poly-A selection TruSeq RNA Sample
Preparation kits. Sequencing libraries were sequenced by the Genomics Core Facility at Weill

645   Cornell Medicine using a HiSeq 2500 (MRL/lpr, MRL/+ samples) and NovaSeq (B6,
B6.Sle1yaa) at a depth of 15-30 million mappable paired-end reads per sample.

Computational analyses of gene expression data

Microarray analysis was performed using healthy controls from an existing GEO dataset

650   (GSE52471) obtained from GEO using GEOquery and combined with non-lesional skin sample
data kindly provided by the authors of the original study (Jabbari et al., 2014). Microarray data
were normalized with gcRMA normalization using the gcrma R package (Wu et al., 2021).
Differential gene expression analysis was performed with limma using the empirical bayes
method (Ritchie et al., 2015) and controlling for microarray hybridization kit and sex. Batch

655 correction was performed using ComBat (Leek et al., 2021). Visualizations were generated using

ComBat corrected data with plotly and ggplot2 in R.

For RNA sequencing analysis, read quality filtering and adapter trimming were performed

using *fastp* (Chen et al., 2018). Filtered reads were mapped to the mouse genome (mm10), and

exonic reads were counted against GENCODE release 27 annotation with the STAR aligner

660 (Dobin et al., 2013) using a customized pipeline available

athttps://gitlab.com/hssgenomics/pipelines. Differential gene expression analysis was performed

with edgeR 3.30.2 using quasi-likelihood framework. Genes with low expression levels (<3

counts per million in at least one group) were filtered from all downstream analyses. P-values

were adjusted to correct for multiple testing using the Benhamini-Hochberg FDR procedure.

665 Genes with adjusted p-values < 0.01 were considered differentially expressed. Downstream

analyses were performed in R using a Shiny-driven visualization platform (RNAseq DRaMA)

developed at the HSS David Z. Rosensweig Genomics Research Center.

Differentially regulated pathways were identified using R QuSAGE 2.22.0 package (Yaari

et al., 2013) with MSigDB C2 set (curated gene sets). All gene sets with less than 10 genes were

670 excluded from the analysis. Pathways with Benhamini-Hochberg adjusted p-values <0.005 (non-

lesional skin versus healthy control) or <0.05 (MRL/lpr versus MRL/+) were used for further

analyses.

Ranked GSEA analysis was performed to define gene sets from MSigDB C2 enriched in

human non-lessional skin and MRL/lpr mice. Log-transformed fold changes in relevant

675 comparisons (non-lesional skin versus healthy control and MRL/lpr versus MRL/+) were used as

ranking variables. Pathways with Benhamini-Hochberg adjusted p-values <0.05 were selected

and the pathways' similarity network was built to visualize gene overlap between pathways

using the R/Bioconductor package *hypeR* (1.7.0) (Federico and Monti, 2020) and compared to enriched pathways in human SLE patients and mouse models.

680   The R/Bioconductor package, GSVA (Hänzelmann et al., 2013)(v1.36.3), was utilized as a non-parametric, unsupervised method to estimate enrichment of pre-defined gene sets in gene expression data from human DLE skin and non-lesional skin from MRL/lpr and B6.Sle1yaa mice. The inputs for GSVA were a matrix of expression values for all samples and curated gene sets describing select immune/tissue cell types and inflammatory cytokines. Low-intensity

685   probes were filtered out if the interquartile range (IQR) of their expression values across all samples was not greater than 0. Enrichment scores were calculated using a Kolgomorov Smirnoff (KS)-like random walk statistic and represented the greatest deviation from zero for a particular sample and gene set. Scores across all samples were normalized to values between -1 (indicating no enrichment) and +1 (indicating enrichment). Gene sets used as input for GSVA

690   are listed in Tables S3 and S7. The human gene sets were previously published (Martínez et al., 2022). The mouse plasma cell, T cell, myeloid cell, neutrophil, pDC, dendritic cell, and keratinocyte gene sets were derived from Mouse CellScan, a tool for the identification of cellular origin of mouse gene expression datasets. The mouse IFN gene sets have been previously described (Kingsmore et al., 2021). The mouse cytokine signatures were generated by an

695   iterative process involving derivation through literature mining and GO term definitions provided by the Mouse Genome Informatics (MGI) GO Browser (Bult et al., 2019).

qPCR

Both "lesional" and "non-lesional" ears were collected from each IMQ-treated and control

700   mouse. Ears were flash-frozen with liquid nitrogen and stored at -80°C. Reverse transcription

was performed using random primers and gene expression assessed by qPCR. PowerSYBR Green PCR Master Mix (Applied Biosystems) was used and assays were run on a QuantStudio 6 Pro (Applied Biosystems). Gene expression was normalized to the control for each target gene. Primers used in the qPCR assay are listed in Table S10.

Sample size and statistics

Sample sizes were determined based on previous similar types of experiments. Two-tailed unpaired and paired Student's t-tests were used for comparisons between conditions to evaluate p-values as indicated. Significance was defined as $p < 0.05$. For figures showing normalized

710   values, the control sample was set to one, and the experimental samples were normalized relative to the control for that experiment. For experiments that contained more than one control sample, the mean was obtained for the control samples, and the individual control and experimental samples were calculated relative to this mean.

715   ### Data availability

All sequencing data will be submitted to the Gene Expression Omnibus database. All other data supporting the findings of this study are available within the paper and its Supplementary Materials.

REFERENCES

Ambler, W., M.J. Howlader, M. Latha Somaraju, E. Seltzer, J. Sim, J. Shin, N. Schwartz, W.D. Shipman, D. Dasoveanu, C. Carballo, E. Sevim, S. Rodeo, D. Erkan, R.P. Kataru, B. Mehrara, and T. Lu. 2022. Lymphatic dysfunction lupus contributes to cutaneous photosensitity and lymph node B cell responses. *bioRxiv : the preprint server for biology* 2022.2006.2013.495930.

Baechler, E.C., F.M. Batliwalla, G. Karypis, P.M. Gaffney, W.A. Ortmann, K.J. Espe, K.B. Shark, W.J. Grande, K.M. Hughes, V. Kapur, P.K. Gregersen, and T.W. Behrens. 2003. Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *PNAS* 100:2610-2615.

Benci, J.L., L.R. Johnson, R. Choa, Y. Xu, J. Qiu, Z. Zhou, B. Xu, D. Ye, K.L. Nathanson, C.H. June, E.J. Wherry, N.R. Zhang, H. Ishwaran, M.D. Hellmann, J.D. Wolchok, T. Kambayashi, and A.J. Minn. 2019. Opposing Functions of Interferon Coordinate Adaptive and Innate Immune Responses to Cancer Immune Checkpoint Blockade. *Cell* 178:933-948.e914.

Bennett, L., A.K. Palucka, E. Arce, V. Cantrell, J. Borvak, J. Banchereau, and V. Pascual. 2003. Interferon and Granulopoiesis Signatures in Systemic Lupus Erythematosus Blood. *J. Exp. Med.* 197:711-723.

Billi, A.C., F. Ma, O. Plazyo, M. Gharaee-Kermani, R. Wasikowski, G.A. Hile, X. Xing, C.M. Yee, S.M. Rizvi, M.P. Maz, C.C. Berthier, F. Wen, L.C. Tsoi, M. Pellegrini, R.L. Modlin, J.E. Gudjonsson, and J.M. Kahlenberg. 2022. Nonlesional lupus skin contributes to inflammatory education of myeloid cells and primes for cutaneous inflammation. *Science Translational Medicine* 14:eabn2263.

Blaydon, D.C., P. Biancheri, W. Di, V. Plagnol, R.M. Cabral, M.A. Brooke, D.A. van Heel, F. Ruschendorf, M. Toynbee, A. Walne, E.A. O'Toole, J.E. Martin, K. Lindley, T. Vulliamy, D.J. Abrams, T.T. MacDonald, J.I. Harper, and D.P. Kelsell. 2011. Inflammatory skin and bowel disease linked to ADAM17 deletion. *The New England Journal of Medicine* 365:1502-1508.

Brahmer, J.R., C. Lacchetti, B.J. Schneider, M.B. Atkins, K.J. Brassil, J.M. Caterino, I. Chau, M.S. Ernstoff, J.M. Gardner, P. Ginex, S. Hallmeyer, J. Holter Chakrabarty, N.B. Leighl, J.S. Mammen, D.F. McDermott, A. Naing, L.J. Nastoupil, T. Phillips, L.D. Porter, I. Puzanov, C.A. Reichner, B.D. Santomasso, C. Seigel, A. Spira, M.E. Suarez-Almazor, Y. Wang, J.S. Weber, J.D. Wolchok, J.A. Thompson, and N. National Comprehensive Cancer. 2018. Management of Immune-Related Adverse Events in Patients Treated With Immune Checkpoint Inhibitor Therapy: American Society of Clinical Oncology Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 36:1714-1768.

Bult, C.J., J.A. Blake, C.L. Smith, J.A. Kadin, and J.E. Richardson. 2019. Mouse Genome Database (MGD) 2019. *Nucleic Acids Res* 47:D801-d806.

Catalina, M.D., P. Bachali, N.S. Geraci, A.C. Grammer, and P.E. Lipsky. 2019. Gene expression analysis delineates the potential roles of multiple interferons in systemic lupus erythematosus. *Commun Biol* 2:140.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884-i890.

Der, E., S. Ranabothu, H. Suryawanshi, K.M. Akat, R. Clancy, P. Morozov, M. Kustagi, M. Czuppa, P. Izmirly, H.M. Belmont, T. Wang, N. Jordan, N. Bornkamp, J. Nwaukoni, J. Martinez, B. Goilav, J.P. Buyon, T. Tuschl, and C. Putterman. 2017. Single cell RNA sequencing to dissect the molecular heterogeneity in lupus nephritis. *JCI Insight* 2:

Der, E., H. Suryawanshi, P. Morozov, M. Kustagi, B. Goilav, S. Ranabothu, P. Izmirly, R. Clancy, H.M. Belmont, M. Koenigsberg, M. Mokrzycki, H. Rominieki, J.A. Graham, J.P. Rocca, N. Bornkamp, N. Jordan, E. Schulte, M. Wu, J. Pullman, K. Slowikowski, S. Raychaudhuri, J. Guthridge, J. James, J. Buyon, T. Tuschl, and C. Putterman. 2019. Tubular cell and keratinocyte single-cell

transcriptomics applied to lupus nephritis reveal type I IFN and fibrosis relevant pathways. *Nat Immunol* 20:915-927.

Dobin, A., C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T.R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15-21.

Federico, A., and S. Monti. 2020. hypeR: an R package for geneset enrichment workflows. *Bioinformatics* 36:1307-1308.

Franzke, C.-W., C. Cobzaru, A. Triantafyllopoulou, S. Löffek, K. Horiuchi, D.W. Threadgill, T. Kurz, N. van Rooijen, L. Bruckner-Tuderman, and C.P. Blobel. 2012. Epidermal ADAM17 maintains the skin barrier by regulating EGFR ligand-dependent terminal keratinocyte differentiation. *The Journal of Experimental Medicine* 209:1105-1119.

Furie, R., M. Khamashta, J.T. Merrill, V.P. Werth, K. Kalunian, P. Brohawn, G.G. Illei, J. Drappa, L. Wang, and S. Yoo. 2017. Anifrolumab, an Anti-Interferon-α Receptor Monoclonal Antibody, in Moderate-to-Severe Systemic Lupus Erythematosus. *Arthritis Rheumatol* 69:376-386.

Furie, R., V.P. Werth, J.F. Merola, L. Stevenson, T.L. Reynolds, H. Naik, W. Wang, R. Christmann, A. Gardet, A. Pellerin, S. Hamann, P. Auluck, C. Barbey, P. Gulati, D. Rabah, and N. Franchimont. 2019. Monoclonal antibody targeting BDCA2 ameliorates skin lesions in systemic lupus erythematosus. *J Clin Invest* 129:1359-1371.

Geesala, R., P.D. Issuree, and T. Maretzky. 2019. Novel functions of inactive rhomboid proteins in immunity and disease. *J Leukoc Biol* 106:823-835.

Goel, R.R., X. Wang, L.J. O'Neil, S. Nakabo, K. Hasneen, S. Gupta, G. Wigerblad, L.P. Blanco, J.B. Kopp, M.I. Morasso, S.V. Kotenko, Z.X. Yu, C. Carmona-Rivera, and M.J. Kaplan. 2020. Interferon lambda promotes immune dysregulation and tissue inflammation in TLR7-induced lupus. *Proc Natl Acad Sci U S A* 117:5409-5419.

Gregorio, J., S. Meller, C. Conrad, A. Di Nardo, B. Homey, A. Lauerma, N. Arai, R.L. Gallo, J. Digiovanni, and M. Gilliet. 2010. Plasmacytoid dendritic cells sense skin injury and promote wound healing through type I interferons. *J Exp Med* 207:2921-2930.

Hänzelmann, S., R. Castelo, and J. Guinney. 2013. GSVA: gene set variation analysis for microarray and RNA-Seq data. *BMC Bioinformatics* 14:7.

Issuree, P.D., T. Maretzky, D.R. McIlwain, S. Monette, X. Qing, P.A. Lang, S.L. Swendeman, K.H. Park-Min, N. Binder, G.D. Kalliolias, A. Yarilina, K. Horiuchi, L.B. Ivashkiv, T.W. Mak, J.E. Salmon, and C.P. Blobel. 2013. iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. *J Clin Invest* 123:928-932.

Jabbari, A., M. Suárez-Fariñas, J. Fuentes-Duculan, J. Gonzalez, I. Cueto, A.G. Franks, Jr., and J.G. Krueger. 2014. Dominant Th1 and minimal Th17 skewing in discoid lupus revealed by transcriptomic comparison with psoriasis. *J Invest Dermatol* 134:87-95.

Karnell, J.L., Y. Wu, N. Mittereder, M.A. Smith, M. Gunsior, L. Yan, K.A. Casey, J. Henault, J.M. Riggs, S.M. Nicholson, M.A. Sanjuan, K.A. Vousden, V.P. Werth, J. Drappa, G.G. Illei, W.A. Rees, and J.N. Ratchford. 2021. Depleting plasmacytoid dendritic cells reduces local type I interferon responses and disease activity in patients with cutaneous lupus. *Sci Transl Med* 13:

Kingsmore, K.M., P. Bachali, M.D. Catalina, A.R. Daamen, S.E. Heuer, R.D. Robl, A.C. Grammer, and P.E. Lipsky. 2021. Altered expression of genes controlling metabolism characterizes the tissue response to immune injury in lupus. *Sci Rep* 11:14789.

Kirou, K.A., C. Lee, S. George, K. Louca, I.G. Papagiannis, M.G. Peterson, N. Ly, R.N. Woodward, K.E. Fry, A.Y. Lau, J.G. Prentice, J.G. Wohlgemuth, and M.K. Crow. 2004. Coordinate overexpression of interferon-alpha-induced genes in systemic lupus erythematosus. *Arthritis Rheum* 50:3958-3967.

815 Lambers, W.M., K. de Leeuw, B. Doornbos-van der Meer, G.F.H. Diercks, H. Bootsma, and J. Westra. 2019. Interferon score is increased in incomplete systemic lupus erythematosus and correlates with myxovirus-resistance protein A in blood and skin. *Arthritis Research & Therapy* 21:260.

Leek, J.T., W.E. Johson, H.S. Parker, E.J. Fertig, A.E. Jaffe, Y. Zhang, J.D. Storey, and L. Callado Torres. 2021. sva: Surrogate Variable Analysis. . *R package version 3.40.0*

820 Lessard, J.C., S. Piña-Paz, J.D. Rotty, R.P. Hickerson, R.L. Kaspar, A. Balmain, and P.A. Coulombe. 2013. Keratin 16 regulates innate immunity in response to epidermal barrier breach. *Proc Natl Acad Sci U S A* 110:19537-19542.

Lora, J., G. Weskamp, T.M. Li, T. Maretzky, D.T.N. Shola, S. Monette, S.F. Lichtenthaler, T.T. Lu, C. Yang, and C.P. Blobel. 2021. Targeted truncation of the ADAM17 cytoplasmic domain in mice results in

825 protein destabilization and a hypomorphic phenotype. *J Biol Chem* 296:100733.

Martin-Fernandez, M., M. Bravo García-Morato, C. Gruber, S. Murias Loza, M.N.H. Malik, F. Alsohime, A. Alakeel, R. Valdez, S. Buta, G. Buda, M.A. Marti, M. Larralde, B. Boisson, M. Feito Rodriguez, X. Qiu, M. Chrabieh, M. Al Ayed, S. Al Muhsen, J.V. Desai, E.M.N. Ferre, S.D. Rosenzweig, B. Amador-Borrero, L.Y. Bravo-Gallego, R. Olmer, S. Merkert, M. Bret, A.K. Sood, A. Al-Rabiaah,

830 M.H. Temsah, R. Halwani, M. Hernandez, F. Pessler, J.L. Casanova, J. Bustamante, M.S. Lionakis, and D. Bogunovic. 2020. Systemic Type I IFN Inflammation in Human ISG15 Deficiency Leads to Necrotizing Skin Lesions. *Cell Rep* 31:107633.

Martínez, B.A., S. Shrotri, K.M. Kingsmore, P. Bachali, A.C. Grammer, and P.E. Lipsky. 2022. Machine learning reveals distinct gene signature profiles in lesional and nonlesional regions of

835 inflammatory skin diseases. *Sci Adv* 8:eabn4776.

Morand, E.F., R. Furie, Y. Tanaka, I.N. Bruce, A.D. Askanase, C. Richez, S.C. Bae, P.Z. Brohawn, L. Pineda, A. Berglind, and R. Tummala. 2020. Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *N Engl J Med* 382:211-221.

Muskardin, T.L.W., and T.B. Niewold. 2018. Type I interferon in rheumatic diseases. *Nature reviews.*

840 *Rheumatology* 14:214-228.

Nickerson, K.M., J.L. Cullen, M. Kashgarian, and M.J. Shlomchik. 2013. Exacerbated Autoimmunity in the Absence of TLR9 in MRL.Fas^lpr Mice Depends on Ifnar1. *The Journal of Immunology* 190:3889-3894.

Pisitkun, P., J.A. Deane, M.J. Difilippantonio, T. Tarasenko, A.B. Satterthwaite, and S. Bolland. 2006.

845 Autoreactive B cell responses to RNA-related antigens due to TLR7 gene duplication. *Science* 312:1669-1672.

Qing, X., Y. Chinenov, P. Redecha, M. Madaio, J.J. Roelofs, G. Farber, P.D. Issuree, L. Donlin, D.R. McLlwain, T.W. Mak, C.P. Blobel, and J.E. Salmon. 2018. iRhom2 promotes lupus nephritis through TNF-alpha and EGFR signaling. *J Clin Invest* 128:1397-1412.

850 Reefman, E., H. Kuiper, P.C. Limburg, C.G. Kallenberg, and M. Bijl. 2008. Type I interferons are involved in the development of ultraviolet B-induced inflammatory skin lesions in systemic lupus erythaematosus patients. *Ann Rheum Dis* 67:11-18.

Ritchie, M.E., B. Phipson, D. Wu, Y. Hu, C.W. Law, W. Shi, and G.K. Smyth. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*

855 43:e47.

Sarkar, M.K., G.A. Hile, L.C. Tsoi, X. Xing, J. Liu, Y. Liang, C.C. Berthier, W.R. Swindell, M.T. Patrick, S. Shao, P.-S. Tsou, R. Uppala, M.A. Beamer, A. Srivastava, S.L. Bielas, P.W. Harms, S. Getsios, J.T. Elder, J.J. Voorhees, J.E. Gudjonsson, and J.M. Kahlenberg. 2018. Photosensitivity and type I IFN responses in cutaneous lupus are driven by epidermal-derived interferon kappa. *Annals of the*

860 *Rheumatic Diseases* 77:1653-1664.

Shipman, W.D., S. Chyou, A. Ramanathan, P.M. Izmirly, S. Sharma, T. Pannellini, D.C. Dasoveanu, X. Qing, C.M. Magro, R.D. Granstein, M.A. Lowes, E.G. Pamer, D.H. Kaplan, J.E. Salmon, B.J. Mehrara,

J.W. Young, R.M. Clancy, C.P. Blobel, and T.T. Lu. 2018. A protective Langerhans cell–keratinocyte axis that is dysfunctional in photosensitivity. *Science Translational Medicine* 10:

865 Singh, B., M. Schneider, P. Knyazev, and A. Ullrich. 2009. UV-induced EGFR signal transactivation is dependent on proligand shedding by activated metalloproteases in skin cancer cell lines. *International Journal of Cancer* 124:531-539.

Sontheimer, C., D. Liggitt, and K.B. Elkon. 2017. Ultraviolet B Irradiation Causes Stimulator of Interferon Genes–Dependent Production of Protective Type I Interferon in Mouse Skin by Recruited

870 Inflammatory Monocytes. *Arthritis & Rheumatology* 69:826-836.

Stannard, J.N., T.J. Reed, E. Myers, L. Lowe, M.K. Sarkar, X. Xing, J.E. Gudjonsson, and J.M. Kahlenberg. 2017. Lupus Skin Is Primed for IL-6 Inflammatory Responses through a Keratinocyte-Mediated Autocrine Type I Interferon Loop. *J Invest Dermatol* 137:115-122.

Strassner, J.P., M. Rashighi, M. Ahmed Refat, J.M. Richmond, and J.E. Harris. 2017. Suction blistering the

875 lesional skin of vitiligo patients reveals useful biomarkers of disease activity. *J Am Acad Dermatol* 76:847-855.e845.

Subramanian, S., K. Tus, Q.Z. Li, A. Wang, X.H. Tian, J. Zhou, C. Liang, G. Bartov, L.D. McDaniel, X.J. Zhou, R.A. Schultz, and E.K. Wakeland. 2006. A Tlr7 translocation accelerates systemic autoimmunity in murine lupus. *Proc Natl Acad Sci U S A* 103:9970-9975.

880 Tsoi, L.C., G.A. Hile, C.C. Berthier, M.K. Sarkar, T.J. Reed, J. Liu, R. Uppala, M. Patrick, K. Raja, X. Xing, E. Xing, K. He, J.E. Gudjonsson, and J.M. Kahlenberg. 2019. Hypersensitive IFN Responses in Lupus Keratinocytes Reveal Key Mechanistic Determinants in Cutaneous Lupus. *The Journal of Immunology* 202:2121-2130.

Werth, V.P., M. Bashir, and W. Zhang. 2004. Photosensitivity in rheumatic diseases. *The journal of*

885 *investigative dermatology. Symposium proceedings* 9:57-63.

Wu, J., R. Irizarry, J. MadDonald, and J. Gentry. 2021. gcrma: Background Adjustment Using Sequence Information. . *R package version 2.64.0*

Wysenbeek, A.J., D.A. Block, and J.F. Fries. 1989. Prevalence and expression of photosensitivity in systemic lupus erythematosus. *Annals of the Rheumatic Diseases* 48:461-463.

890 Yaari, G., C.R. Bolen, J. Thakar, and S.H. Kleinstein. 2013. Quantitative set analysis for gene expression: a method to quantify gene set differential expression including gene-gene correlations. *Nucleic Acids Res* 41:e170.

Yokogawa, M., M. Takaishi, K. Nakajima, R. Kamijima, C. Fujimoto, S. Kataoka, Y. Terada, and S. Sano. 2014. Epicutaneous application of toll-like receptor 7 agonists leads to systemic autoimmunity in

895 wild-type mice: a new model of systemic Lupus erythematosus. *Arthritis Rheumatol* 66:694-706.

Zhang, X., D. Bogunovic, B. Payelle-Brogard, V. Francois-Newton, S.D. Speer, C. Yuan, S. Volpi, Z. Li, O. Sanal, D. Mansouri, I. Tezcan, G.I. Rice, C. Chen, N. Mansouri, S.A. Mahdaviani, Y. Itan, B. Boisson, S. Okada, L. Zeng, X. Wang, H. Jiang, W. Liu, T. Han, D. Liu, T. Ma, B. Wang, M. Liu, J.Y. Liu, Q.K. Wang, D. Yalnizoglu, L. Radoshevich, G. Uzé, P. Gros, F. Rozenberg, S.Y. Zhang, E.

900 Jouanguy, J. Bustamante, A. García-Sastre, L. Abel, P. Lebon, L.D. Notarangelo, Y.J. Crow, S. Boisson-Dupuis, J.L. Casanova, and S. Pellegrini. 2015. Human intracellular ISG15 prevents interferon-α/β over-amplification and auto-inflammation. *Nature* 517:89-93.

905   **ACKNOWLEDGEMENTS**

Acknowledgements: The authors thank the members of the Lu Lab for helpful discussions and critical reading of the manuscript.

910   **Funding:**

Alpha Omega Alpha Carolyn L. Kuckein Student Research Fellowship (TML)

Erwin Schrodinger Fellowship J 4638-B FWF (VZ)

National Institutes of Health grant T32AR071302-01 to the Hospital for Special Surgery Research Institute Rheumatology Training Program (NS and WDS)

National Institutes of Health grant MSTP T32GM007739 to the Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-PhD Program (WDS)

National Institutes of Health grant K08 AR069111 to the University of Iowa Department of Dermatology (AJ)

925   Veterans Administration VA Merit I01 BX004907 (AJ),

Physician Scientist Career Development Award from the Dermatology Foundation (AJ)

National Institutes of Health grant R01 DK099087 (IR)

National Institutes of Health grant R35GM134907 (CPB)

National Institutes of Health grant R01AI079178 (TTL)

935   National Institutes of Health grant R21 AR081493 (TTL)

Department of Defense grant W81XWH-21-LRP-IPA (TTL)

Lupus Research Alliance (TTL)

St. Giles Foundation (TTL)

National Institutes of Health Office of the Director grant S10OD019986 to Hospital for Special Surgery.

950   **AUTHOR CONTRIBUTIONS**

Conceptualization: TML, TTL

Performed experiments and acquired data: TML, VZ, ESS, MD, KRV, NS, AJ

Data analyses and interpretation: TML, VZ, ESS, MD, YC, ARD, DJO, IR, CPB, PEL, TTL

955   Methodology and reagents: JL, YL, WDS, WGA, JHZ, MR, NA, CPB

Supervision: SFT, KBO, JGK, IR, CPB, PEL, TTL

Writing—original draft: TML, VZ, MD, YC, AD, IR, PEL, TTL

Writing—review and editing: All authors

Competing interests: Authors declare that they have no competing interests.

SUPPLEMENTARY MATERIALS

Figure S1. Further analyses of gene expression in human DLE and multiple murine lupus models.

Figure S2. ADAM17 expression levels in LCAd17 mice, UVR-induced cell surface TNFR1 loss on LCs in mixed epidermal cell suspensions is dependent on LC ADAM17.

Figure S3. Langerhans cell yield and gating with suction blistering of human skin.

Table S1. Differentially expressed genes in DLE vs healthy control non-lesional skin microarray

Table S2. Differentially expressed pathways in DLE vs healthy control non-lesional skin using QuSAGE

Table S3. Human gene sets used for GVSA

Table S4. Differentially expressed genes in MRL/lpr vs control MRL/+ non-lesional skin RNAseq

Table S5. Differential expressed pathways in MRL/lpr vs control MRL/+ non-lesional skin using QuSAGE

Table S6. Differentially expressed genes in B6.Sle1yaa vs control B6 non-lesional skin RNAseq

Table S7. Murine gene sets used for GVSA

Table S8. Ranked GSEA enriched pathways in DLE vs normal control dataset

Table S9. Antibodies used

Table S10. Primer sequences for qPCR