The interferon-rich skin environment regulates Langerhans cell ADAM17 to promote photosensitivity in lupus

Authors: Thomas M. Li[1,†,§], Victoria Zyulina[1,2,§], Ethan S. Seltzer[1,‡,§], Marija Dacic[3,4,5], Yurii Chinenov[3], Andrea R. Daamen[6], Keila R. Veiga[1,7,8,††], Noa Schwartz[1,9,‡‡], David J. Oliver[3], Jose Lora[4,5], Ali Jabbari[10,§§], Yong Liu[11], William D. Shipman[1,12,13,¶], William G. Ambler[1,7,8], Sarah F. Taber[8,9], Karen B. Onel[8,9], Jonathan H. Zippin[10], Mehdi Rashighi[14], James G. Krueger[10], Niroshana Anandasabapathy[11,12,13], Inez Rogatsky[2,3,4,13], Carl P. Blobel[4,5,15], Peter E. Lipsky[6], Theresa T. Lu[1,2,7,8,13]

Affiliations:
[1]Autoimmunity and Inflammation Program, Hospital for Special Surgery Research Institute, New York, NY 10021
[2]Department of Microbiology and Immunology, Weill Cornell Medical College; New York, USA
[3]David Z. Rosensweig Genomics Research Center, Hospital for Special Surgery Research Institute; New York, USA
[4]Arthritis and Tissue Degeneration Program, Hospital for Special Surgery Research Institute, New York, USA
[5]Physiology, Biophysics, and Systems Biology Program, Weill Cornell Graduate School of Medical Sciences; New York, USA
[6]AMPEL BioSolutions, Charlottesville, VA, USA
[7]Pediatric Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, USA
[8]Department of Pediatrics, Weill Cornell Medical College; New York, USA
[9]Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, NY 10021
[10]Laboratory of Investigative Dermatology, Rockefeller University, New York, USA
[11]Department of Dermatology, Weill Cornell Medical College; New York, USA
[12]Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-PhD Program, Weill Cornell Medical College; New York, USA
[13]Immunology and Microbial Pathogenesis Program, Weill Cornell Graduate School of Medical Sciences; New York, USA
[14]Department of Dermatology, University of Massachusetts Medical School; Worcester, USA
[15]Department of Physiology, Biophysics, and Systems Biology, Weill Cornell Medical College; New York, USA

[§]These authors share first authorship

[†]Current address Icahn School of Medicine at Mount Sinai; New York, USA
[‡]Current address Rockefeller University; New York, USA

45 ††Current address Department of Rheumatology, Maria Fareri Children's Hospital; New York, USA

‡‡Current address Department of Medicine (Rheumatology), Montefiore Medical Center/Albert Einstein College of Medicine; New York, USA

§§Current address Department of Dermatology, University of Iowa Hospital and Clinics; Iowa

50 City, USA

¶Current address Department of Dermatology, Yale School of Medicine; New Haven, USA

Corresponding author
Theresa T. Lu

55 Hospital for Special Surgery
535 East 70th Street
Tel: 212-774-2532
Email: LuT@HSS.edu

ABSTRACT

Background: The autoimmune disease lupus erythematosus (lupus) is characterized by photosensitivity, where even ambient ultraviolet radiation (UVR) exposure inflames skin. Beneficial effects of anifrolumab (anti-interferon α/βreceptor (anti-IFNAR)) on lupus skin disease support a pathogenic role for IFN-I , but mechanistic understanding is limited. We have

65   shown that Langerhans cell (LC) dysfunction contributes to photosensitivity. Healthy LCs act via a disintegrin and metalloprotease 17 (ADAM17) to release epidermal growth factor receptor (EGFR) ligands that limit UVR-induced keratinocyte apoptosis and photosensitivity. However, LC ADAM17 activity is reduced in non-lesional lupus model skin, and data point to reduced LC-mediated protection in human lupus. Here, we asked about the role of the IFN-rich lupus skin

70   environment in LC dysfunction and the implications of this regulation for photosensitivity.

Methods: Gene expression patterns in non-lesional skin from human lupus and multiple murine models were examined. We used MRL/lpr, B6.Sle1yaa, and imiquimod models of lupus in *in vivo* studies to assess the role of IFN-I in LC ADAM17 dysfunction and photosensitivity.

Results: We show a shared IFN-rich environment in non-lesional skin across human and murine

75   model systems, that IFN-I inhibits LC ADAM17 activity, and that anti-IFNAR in lupus models restores LC ADAM17 function and reduces photosensitivity in EGFR and LC ADAM17-dependent manners. Reactive oxygen species (ROS) can mediate ADAM17 activity, and we show reduced LC ROS expression in lupus models that is restored by anti-IFNAR.

Conclusions: Our findings suggest that IFN-I promotes photosensitivity by causing LC

80   ADAM17 dysfunction and that anifrolumab ameliorates lupus skin disease at least in part by restoring LC function. This work provides insight into IFN-I-mediated disease mechanisms, LC regulation, and a mechanism of action for anifrolumab in lupus.

INTRODUCTION

The autoimmune disease lupus erythematosus (lupus) manifests in part by

85   photosensitivity, a sensitivity to ultraviolet radiation (UVR) whereby even ambient sunlight can

trigger the development of inflammatory skin lesions. Patients with lupus skin disease, or

cutaneous lupus erythematosus (CLE), can also have the systemic form of the disease, systemic

lupus erythematosus (SLE), characterized by inflammation and damage to additional organs such

as the kidneys, brain, and heart. In SLE patients, photosensitive skin responses can be associated

90   with worsening systemic autoimmunity and end organ damage. A prominent interferon (IFN)

signature indicative of exposure to type I interferon (IFN-I) is found in circulating cells and

tissues in lupus (Baechler et al., 2003; Bennett et al., 2003; Catalina et al., 2019; Der et al., 2019;

Jabbari et al., 2014; Kirou et al., 2004), and the development and recent FDA approval of

anifrolumab (anti-IFNAR1) for SLE highlights the importance of IFN-I in disease pathogenesis.

95   Anifrolumab is especially efficacious for skin disease in SLE patients, consistent with reduced

skin lesions in IFNAR-deficient SLE model mice (Nickerson et al., 2013) and an IFN-I signature

in both lesional and non-lesional skin (Billi et al., 2022; Der et al., 2017; Der et al., 2019;

Reefman et al., 2008; Stannard et al., 2017). These findings suggest a prominent role for IFN-I in

lupus skin disease, but mechanisms of IFN-I-mediated pathogenesis remain poorly understood.

100   We recently reported that Langerhans cells (LCs) can limit UVR-induced skin

inflammation by expressing a disintegrin and metalloprotease 17 (ADAM17) which releases

membrane-bound epidermal growth factor receptor (EGFR) ligands that then preserve epidermal

integrity via EGFR stimulation of keratinocytes (Shipman et al., 2018). Two distinct

photosensitive SLE mouse models showed reduced LC *Adam17* mRNA which corresponded to

105   reduced LC ADAM17 sheddase activity, and photosensitivity was reduced by topical EGFR

ligand application. Together, these data suggested that the lack of LC ADAM17-dependent EGFR-ligand sheddase activity contributes to photosensitivity. Non-lesional skin of SLE patients showed reduced LC numbers along with reduced epidermal EGFR phosphorylation, further supporting the potential for LC defects to contribute to photosensitivity in lupus. The factors that lead to LC dysfunction, however, remain to be fully elucidated.

Here, we test the hypothesis that the high IFN-I environment present in even non-lesional lupus skin promotes LC ADAM17 dysfunction to contribute to photosensitivity. We build upon human data showing an IFN-I signature in non-lesional skin of CLE patients and establish that the photosensitive MRL/lpr and B6.Sle1yaa models share IFN-I-related and other gene expression modules with human skin. The inducible photosensitive imiquimod (IMQ) model mice also upregulated IFN-associated genes in non-lesional skin. IFN-I is sufficient to downregulate healthy murine and human LC ADAM17 activity, and anti-IFN-I receptor (anti-IFNAR) treatment of lupus mouse models restores LC ADAM17 activity and reduces photosensitivity in an EGFR and LC ADAM17-dependent manner. Lastly, reactive oxygen species (ROS) are known to promote ADAM17 activity (Singh et al., 2009), and we show that UVR-induced LC ROS are reduced in lupus model mice while anti-IFNAR restores ROS expression. Together, our results establish that multiple photosensitive murine lupus models are similar to human lupus patients in demonstrating IFN-I signatures in non-lesional skin and put forth a mechanism whereby IFN-I contributes to photosensitivity at least in part by inhibiting LC ADAM17 activity. These data provide insight into IFN-I-mediated contributions, delineate a driver of LC dysfunction, and suggest that restoration of LC ADAM17 function is a mechanism of action for anifrolumab in lupus skin disease.

RESULTS

Non-lesional skin in lupus model mice share an IFN-I signature and other functional gene expression modules with human lupus skin

As we are interested in the mechanisms that predispose to photosensitivity and to the LC defects seen in lupus mouse models and human SLE skin (Shipman et al., 2018), we sought to

135 better understand the environment in such non-lesional skin. Recent studies have suggested a high IFN-I environment in non-lesional skin of lupus patients based on single cell RNAseq of keratinocytes from SLE patients and skin from CLE patients (Billi et al., 2022; Der et al., 2017; Der et al., 2019), CLE keratinocyte IFN-κ expression (Stannard et al., 2017), and upregulation of IFN-stimulated genes (ISGs) such as *MX1* on tissue sections in SLE and incomplete SLE patients

140 (Lambers et al., 2019; Reefman et al., 2008). Consistent with these findings, our recent analysis of publicly available bulk transcriptomic datasets from non-lesional DLE skin using Gene Set Variation Analysis (GSVA), an unsupervised gene set analysis approach, showed a proportion of samples with upregulated IFN gene set expression when compared to healthy controls, although this effect was modest when compared to the upregulation in lesional skin (Martínez et al.,

145 2022). Here, we examined an additional non-lesional skin dataset from a CLE cohort with the discoid form of CLE (DLE) and also non-lesional skin from multiple lupus models to understand shared pathways.

The non-lesional skin has not been previously analyzed and comes from a DLE cohort whose lesional skin had been shown to have a Th1 signature on microarray when compared to

150 healthy controls and psoriatic skin (Jabbari et al., 2014). This was among the lesional skin cohorts that was shown by GSVA to express the IFN gene set at high levels (Martínez et al., 2022). Normal (ie healthy subject) and non-lesional skin exhibited significantly different

155   **Figure 1**. **Analysis of a new cohort reinforces a model of an IFN-rich environment and LC and keratinocyte dysfunction in non-lesional DLE skin.** (A-H) Microarray analysis of gene expression from non-lesional skin of DLE patients (n=7) and healthy controls (n=13). In (E-F), samples were analyzed with lesional DLE and psoriasis (PSO) skin samples that were collected as part of the same study and previously published (Jabbari et al., 2014). (A,E) Principal
160   component analysis (PCA) using top 250 genes. (B) Differentially expressed pathways were determined using QuSAGE pathway analysis against Molecular Signatures Database (MSigDB). (C) Volcano plot of differentially expressed genes. Genes from IFN α/β (red), IFNγ (blue) and IRF transcription factor (green) pathways are marked. CD207 (pink) is additionally marked. (D) Heatmap of gene expression in the IFN α/β signaling pathway. (F, G) Gene Set Variation
165   Analysis (GSVA) of gene sets relevant to lupus (Martínez et al., 2022). (H) GSVA using gene sets comprising specific IFN sub-types. (F-H) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by unpaired t-test.

expression profiles (Figure 1A and Table S1). Differential pathway analysis via Quantitative Set

Analysis for Gene Expression (QuSAGE) showed that IFN-I and IFN-γ pathways were among

170   the most upregulated in the non-lesional skin compared to normal controls (Figure 1B and Table

S2). The IFN-I pathway was comprised of several upregulated IFN-stimulated transcription

factors, including IRF 1, 2, 7, 8, and 9 along with multiple IRF targets (ISG15, IFIT1, IFIT3,

MX1, MX2) (Figure 1C-D, S1A). Notably, *CD207*, the gene encoding Langerin and a lineage

marker of LCs, was downregulated (Figure 1C), consistent with accumulating data on LC

175   dysfunction in lupus (Billi et al., 2022; Martínez et al., 2022; Shipman et al., 2018). A supervised

approach to analyzing normal control versus non-lesional DLE skin gene expression data, then,

yielded results consistent with other cohorts showing upregulation of IFN pathways and LC

dysfunction.

    We next performed GSVA using previously defined lupus-relevant gene sets (Martínez et

180   al., 2022) (Table S3). Analysis of the non-lesional samples along with the previously analyzed

lesional and psoriatic samples (Jabbari et al., 2014) reiterated the GSVA results of multiple

datasets. The gene expression profiles of the four conditions were significantly different from

each other (Figure 1E). Lesional DLE was characterized by enrichment of immune cell gene sets

including plasma cells, T cells, monocyte/myeloid cells, neutrophils, and plasmacytoid dendritic

185   cells (pDCs) in conjunction with de-enrichment of LCs when compared to normal controls

(Figure 1F). Lesional DLE samples were also enriched for inflammatory cytokine gene sets

including IL12, IL21, IL23, and TNF as well as a highly upregulated IFN gene signature as

previously seen in lesional DLE skin (Figure 1F) (Martínez et al., 2022). Non-lesional skin also

showed upregulation IFN when compared to healthy controls, although this change was dwarfed

190   when compared to enrichment in lesional skin (Figure 1F). GSVA analysis of just healthy

controls and non-lesional skin in the absence of the high IFN enrichment from lesional skin revealed clear upregulation of the IFN gene set in non-lesional skin as well as de-enrichment of LC and keratinocyte gene sets (Figure 1G). These results were consistent with our previous GSVA analyses of other non-lesional cohorts (Martínez et al., 2022), suggesting along with

195   direct cellular analyses (Billi et al., 2022; Sarkar et al., 2018; Shipman et al., 2018) a loss of epidermal (LC and keratinocyte) function in the context of an IFN-rich proinflammatory environment. The IFN gene set upregulation in non-lesional skin was driven by IFNα, IFNκ, and IFNω gene set upregulation (Figure 1H). GSVA of these previously unanalyzed non-lesional DLE skin samples, then, reinforced previous gene expression analyses and revealed an IFN-I

200   signature in this cohort, all pointing to perturbed epidermal and skin homeostasis even in non-lesional skin.

We examined non-lesional ear skin from MRL/lpr and B6.Sle1yaa SLE mouse models by RNAseq. The photosensitive MRL/lpr model has a *Fas* gene mutation and develops a lupus-like phenotype especially well on the MRL and but not the B6 genetic background. It has reduced LC

205   ADAM17 sheddase function and mRNA, and topical EGFR ligand supplementation ameliorated photosensitivity (Shipman et al., 2018). Although the differences in gene expression between MRL/lpr mice and control MRL-MpJ (MRL/+) mice by RNAseq were less dramatic than those in the human cohort (Figures 1A-D and 2A-D) (Table S4), QuSAGE showed that the IFN-α/β and IFN-γ pathways were among the most highly expressed in MRL/lpr mice (Figure 2B and

210   Table S5). Consistent with the activation of IFN pathways, several IRF transcription factors and their targets were expressed at higher levels in MRL/lpr mice (Figure 2C-D and Figure S1B).

The B6.Sle1yaa model is driven by the Sle1 lupus susceptibility locus derived from lupus-prone NZB2410 mice in combination with the Y chromosome autoimmune accelerator locus

215 **Figure 2. Non-lesional skin of multiple photosensitive murine SLE models recapitulate gene expression patterns of human DLE non-lesional skin.** (A-I) Non-lesional skin gene expression from MRL/lpr (LPR) and control MRL/+ (MRL) mice (A-D, F-G) and B6.Sle1yaa and control B6 mice (E, H-I) were analyzed by RNA sequencing. (A, E) PCA using top 250 genes. (B) Differentially expressed pathways determined by QuSAGE (continued on next page)

Figure 2 (continued), pathway analysis against MSigDB. (C) Volcano plot of differentially expressed genes. Genes from IFN α/β (red), IFNγ (blue) and IRF transcription factor (green) pathways are marked. (D) Heatmap showing gene expression in the IFN α/β signaling pathway. (F, H) GSVA of gene sets relevant to lupus, adapted for murine models (Kingsmore et al., 2021). (G, I) GSVA of gene signatures specific to distinct IFN subtypes. (J) Imiquimod (IMQ) model non-lesional and lesional ears and control mice ears were analyzed by qPCR of select IFN-related genes that are upregulated in non-lesional skin of human DLE and MRL/lpr and B6.Sle1yaa models. (A, E-J) Each symbol represents one mouse (A, E-I) or one ear, with non-lesional and lesional ears coming from the same mice (J). (F-J) *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 by unpaired t-test.

whose effects are attributed to *Tlr7* duplication (Pisitkun et al., 2006; Subramanian et al., 2006).

These mice are photosensitive and have reduced LC *Adam17* mRNA expression (Shipman et al.,

2018). As expected, compared to B6 controls, B6.Sle1yaa mice expressed elevated levels of

235   several lupus related genes from the Yaa locus including Tlr7 (Figure S1C). Although

differences between the B6 and B6.Sle1yaa transcriptomes were modest (Figure 2E)(Table S6),

several genes involved in the regulation of and response to IFN were expressed at higher levels,

including *Irf7, Irf9, Isg15, Xaf1, Selp,* and *Ube2l6* (Figure S1D). Although LC numbers were not

different at 10 months of age when this model is fully diseased (Shipman et al., 2018), *Cd207*

240   expression in younger (5 month old) diseased mice was downregulated when compared to B6

controls (Figure S1D), similar to the findings in CLE skin (Figure 1C). By supervised gene set

analysis, both the MLR/lpr and B6.Sle1yaa murine models showed similarities to human non-

lesional skin.

By GSVA of lupus-relevant murine gene sets (Table S7), parallels between the MLR/lpr

245   and B6.Sle1yaa murine models and human lupus were even more dramatic. MRL/lpr mice

showed increased enrichment of a number of inflammatory cell and cytokine gene sets including

T cells, myeloid cells, neutrophils, pDCs, IFN, IL1, IL21, and TNF (Figure 2F). However, in

contrast to human DLE, non-lesional skin of MRL/lpr mice was enriched for the keratinocyte

gene signature. LC numbers in the skin of MRL/lpr mice are not downregulated when compared

250   to MRL/+ controls (Shipman et al., 2018), which was reflected in the absence of a decreased

dendritic cell signature. The IFN gene set upregulation was broad, with significant enrichment of

gene signatures induced by stimulation from multiple IFNs (Figure 2G). B6.Sle1yaa mice

showed less dramatic differences than the MRL/lpr model, but also showed upregulation of the

IFN gene set (although there was not a significant contribution from any one IFN signature) and

255   downregulation of the DC gene set, potentially reflective of decreased LC function (Figure 2H-

1). Together, our results showed a number of parallels to the human system, including an IFN

signature in these lupus models.

We also tested for expression of ISGs in the imiquimod (IMQ) lupus model. This model is

induced by 4-5 weeks of topical IMQ, a TLR 7/8 agonist, resulting in autoantibody production,

260   enlarged spleen and lymph nodes, nephritis, and photosensitivity characteristic of SLE (Ambler

et al., 2022; Goel et al., 2020; Yokogawa et al., 2014). The model can be induced in B6 mice,

allowing for the use of transgenic models to assess for effects on lupus features without having to

cross onto a different genetic background. We applied IMQ to the right ear to assess ISGs in both

the painted right "lesional" and unpainted left "non-lesional" ears. We performed qPCR for a set

265   of transcripts (*Gbp2, Ifi35, Ifit1, Ifit3, Ifit2, Irf7, Irf8, Isg15, Stat1,* and *Xaf1)* that were

upregulated across human and murine non-lesional skin datasets using our datasets (Figures

S1A, B, D) and that in (Billi et al., 2022) (Figure 2J). Most of the genes we examined showed

upregulation in the "lesional" ear, including transcripts for IRF target genes *Irf7, Irf8* and *Stat1*,

and the apoptosis marker *Xaf1* potentially reflecting tissue damage along with an IFN-rich

270   environment. Several transcripts also showed trends toward upregulation in non-lesional skin

when compared to mice that were not treated with IMQ (Figure 2J). Notably, *Isg15*, the only

transcript that we examined that was upregulated across all the datasets, was also upregulated in

the non-lesional ears of the IMQ mice. ISG15 has been shown to limit over-amplification of

IFN-I effects, and its deficiency predisposes to skin inflammation among other findings (Martin-

275   Fernandez et al., 2020; Zhang et al., 2015), suggesting that upregulation of *Isg15* may reflect an

attempt to minimize IFN-I-mediated inflammation. These results showed that non-lesional skin

from human lupus and 3 murine models were characterized by upregulation of IFN-associated

transcripts, albeit at lower levels than in lesional skin.

Turning our attention to common pathways in human and murine lupus non-lesional skin,

280 we performed ranked Gene Set Enrichment Analysis (GSEA) separately for DLE and MRL/lpr

datasets using negative log-transformed p-values as a ranking variable, constructed pathway

similarity networks, and compared selected pathways (FDR<0.005) in humans and mice.

Consistent with QuSAGE analysis (Figure 1B and 2B), IFN-related pathways were enriched in

human DLE and MRL/lpr non-lesional skin (nodes 3, 4, 8), along with several other pathways

285 often associated with lupus (Figure 3A): T helper pathway (nodes 26); natural killer cell-

mediated cytotoxicity (nodes 28); Toll-like receptor pathways (nodes 36,46); JAK-STAT

signaling pathway (node 43); and multiple pathways associated with complement activation

(nodes 6, 23, 29, 34, 52, 53, 58, 61, 62, 64, 75, 88, 94, 95). For the full list of pathways identified

by ranked GSEA, see Table S8. Direct comparison of differentially expressed genes in DLE and

290 MRL/lpr skin (Figure 3B) revealed a gene signature typical of nucleic acid-driven

interferonopathies dominated by IFN-stimulated transcription factors (IRF1,3,9 and STAT1),

IFN-induced viral restriction factors (BST2, SAMHD1, DDX60, IFIH1, ZBP1), neutrophilic

markers (HMOX1, S100A8, S100A9, LCN2, RSAD2), and apoptosis-related genes (XAF1,

TNFSF10). Several keratins (KRT16, 6B and 34) that are rapidly induced upon skin damage and,

295 in turn, downregulate inflammatory responses (Lessard et al., 2013), were highly expressed in

both human CLE and MRL/lpr skin, likely reflecting disturbed epidermal homeostasis.

Comparisons of the GSVA analyses of non-lesional skin among the datasets (Figure 3C)

echoed the GSEA analysis in emphasizing the prominent enrichment of IFN in human CLE and

both MRL/lpr and B6.Sle1yaa lupus-prone mice. In addition, de-enrichment of the LC/DC gene

REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM
REACTOME_INNATE_IMMUNE_SYSTEM
REACTOME_INTERFERON_SIGNALING
REACTOME_INTERFERON_ALPHA_BETA_SIGNALING
REACTOME_ADAPTIVE_IMMUNE_SYSTEM
KEGG_CYTOKINE_CYTOKINE_RECEPTOR_INTERACTION
REACTOME_INTERFERON_GAMMA_SIGNALING
REACTOME_SIGNALING_BY_INTERLEUKINS
REACTOME_NEUTROPHIL_DEGRANULATION
KEGG_PRIMARY_IMMUNODEFICIENCY
REACTOME_CHEMOKINE_RECEPTORS_BIND_CHEMOKINES
KEGG_CHEMOKINE_SIGNALING_PATHWAY
KEGG_CELL_ADHESION_MOLECULES_CAMS
REACTOME_LEISHMANIA_INFECTION
KEGG_HEMATOPOIETIC_CELL_LINEAGE
BIOCARTA_THELPER_PATHWAY
KEGG_LEISHMANIA_INFECTION
KEGG_NATURAL_KILLER_CELL_MEDIATED_CYTOTOXICITY
REACTOME_HEMOSTASIS
KEGG_VIRAL_MYOCARDITIS
REACTOME_COSTIMULATION_BY_THE_CD28_FAMILY
REACTOME_INTERLEUKIN_10_SIGNALING
REACTOME_TOLL_LIKE_RECEPTOR_CASCADES
REACTOME_GPCR_LIGAND_BINDING
REACTOME_CLASS_A_1_RHODOPSIN_LIKE_RECEPTORS
BIOCARTA_GRANULOCYTES_PATHWAY
KEGG_JAK_STAT_SIGNALING_PATHWAY
REACTOME_INTERLEUKIN_2_FAMILY_SIGNALING
KEGG_TOLL_LIKE_RECEPTOR_SIGNALING_PATHWAY
REACTOME_ANTIVIRAL_MECHANISM_BY_IFN_STIMULATED_GENES
REACTOME_SIGNALING_BY_GPCR
56 REACTOME_PEPTIDE_LIGAND_BINDING_RECEPTORS
58 REACTOME_FCGAMMA_RECEPTOR_FCGR_DEPENDENT_
   PHAGOCYTOSIS
61 REACTOME_PLATELET_ACTIVATION_SIGNALING_AND_AGGREGATION
63 REACTOME_ANTI_INFLAMMATORY_RESPONSE_FAVOURING_
   LEISHMANIA_PARASITE_INFECTION
65 REACTOME_INTERLEUKIN_4_AND_INTERLEUKIN_13_SIGNALING
69 KEGG_APOPTOSIS
72 BIOCARTA_NEUTROPHIL_PATHWAY
REACTOME_METAL_SEQUESTRATION_BY_ANTIMICROBIAL_PROTEINS
BIOCARTA_MHC_PATHWAY
BIOCARTA_HIVNEF_PATHWAY
BIOCARTA_MONOCYTE_PATHWAY
REACTOME_CREATION_OF_C4_AND_C2_ACTIVATORS
KEGG_B_CELL_RECEPTOR_SIGNALING_PATHWAY
REACTOME_CASPASE_ACTIVATION_VIA_DEATH_RECEPTORS_IN_
   THE_PRESENCE_OF_LIGAND
94 REACTOME_INITIAL_TRIGGERING_OF_COMPLEMENT

Figure 3. Analysis of human and murine lupus models show shared gene expression pathways. (A) Pathway similarity network built for the most enriched pathways (adjusted p<0.005) in DLE non-lesional skin determined by ranked GSEA (complete list in Table S8). Colors indicate network community clusters identified by the walktrap algorithm. Red borders indicate pathways that were also enriched in LPR mice (listed). (B) Comparison of gene expression changes in DLE vs MRL/lpr mice. (C) Heatmap of GSVA scores for shared gene sets between non-lesional skin from DLE patients, LPR mice, and B6.Sle1yaa mice. Asterisks indicate significant differences in GSVA scores compared to controls for each dataset.

310   set was present in both human DLE and B6.Sle1yaa mouse non-lesional skin (Figure 3C).

Multiple approaches to analyze gene expression of non-lesional skin from human DLE and

murine lupus models, then, showed upregulation of IFN-related transcripts and disturbance of

LC and epidermal homeostasis. These data are consistent with the idea that non-lesional skin is

primed for photosensitive responses and that LCs sit within an IFN-rich environment in both

315   human lupus and multiple murine models that may cause LC dysfunction.

IFN-I inhibits LC ADAM17 function

We next examined the effects of IFN-I on LC ADAM17 function and expression. As

previous (Shipman et al., 2018), we assessed LC ADAM17 function by quantifying UVR-

320   induced cell surface TNFR receptor 1 (TNFR1) shedding. We confirmed that UVR treatment

performed on a mixture of digested epidermal cells induced LC loss of surface TNFR1 in an LC

ADAM17-dependent manner (Figure S2A-B), similar to our results with isolated LCs (Shipman

et al., 2018). IFN-κ, an IFN-I that is highly expressed by lupus keratinocytes (Tsoi et al., 2019),

reduced LC ADAM17 activity by nearly 40% (Figure 4A). Using a validated anti-murine

325   ADAM17 antibody (Lora et al., 2021) (Figure S2A), we observed that IFN-I did not reduce

ADAM17 cell surface levels by the same magnitude (Figure 4B) (Lora et al., 2021), and LC

viability was not affected (Figure 4C). The IFN-κ-driven inhibition of ADAM17 activity was

abrogated by the JAK1/3 inhibitor tofacitinib without altering ADAM17 protein levels (Figure

4D-E) or cell viability (Figure 4F), supporting an IFNAR-mediated inhibition of LC ADAM17

330   sheddase activity rather than loss of ADAM17 cell surface expression. Application of IFN-κ to

the skin of wild-type mice reduced LC ADAM17 activity by 57% (Figure 4G) while cell surface

ADAM17 was reduced by 9% (Figure 4H). Together, these data indicate that IFN-I is sufficient

Figure 4. IFN-I inhibits LC ADAM17 sheddase activity. (A-K) Epidermal cell suspensions were treated ex vivo or mice were treated in vivo with IFN-I prior to assaying for UVR-induced LC ADAM17 sheddase activity, ADAM17 levels, and numbers. Cells from WT mice were treated with IFNκ or vehicle (A-C) with and without tofacitinib (D-F); IFN κ or vehicle was applied topically to ears of WT mice 16-20 hours prior to generation of epidermal cell suspensions (G-H); suction blister epidermal cell suspensions from healthy human donors were treated with IFNβ or vehicle (I-K). (A, D, G, I) LC ADAM17 sheddase activity as indicated by the extent of UVR-induced cell surface TNFR1 loss. Percent change in cell surface TNFR1 mean fluorescence intensity (MFI) after UVR (left); relative LC ADAM17 activity calculated by normalizing TNFR1 loss to that of vehicle controls (right). In (I), lines connect samples from the same donor. (B, E, H, J) Cell surface ADAM17 levels as indicated by MFI relative to that of vehicle controls. (C, F, K) Relative numbers of LCs per sample after indicated treatments. (A-K) Each symbol represents cells from a single mouse or donor, bars represent average values, and error bars are SD. n=3-10 per condition over 3-5 independent experiments. *p<0.05, **p<0.01, ***p<0.001, n.s.=not significant by unpaired t-test.

350   to downregulate LC murine ADAM17 function and does so primarily at the level of sheddase

activity ex vivo and in vivo.

We also asked how IFN-I affects human LCs. While LCs can be obtained from skin

biopsies or discarded tissues associated with surgical procedures, these tissues often require

lengthy enzymatic digestion to dissociate the cells (Shipman et al., 2018), which has the potential

355   to disrupt cell function. We found that epidermal roofs yielded from suction blistering could be

digested for a short amount of time to obtain an epidermal cell mixture containing LCs (Figure

S3) (Strassner et al., 2017). Addition of IFN-I reduced LC ADAM17 activity by 62% without

altering ADAM17 expression or LC viability (Figure 4I-K). These results together showed that

IFN-I can inhibit LC ADAM17 activity in both murine and human LCs and supported the idea

360   that high levels of IFN-I could potentially contribute to LC ADAM17 dysfunction in disease.

IFNAR is important for LC ADAM17 dysfunction in multiple lupus models

We next asked whether IFN-I was an important mediator of LC ADAM17 dysfunction in

lupus models. Consistent with previous findings (Shipman et al., 2018), MRL/lpr mice showed a

365   42% reduction in LC ADAM17 sheddase activity when compared to control MRL/+ mice

(Figure 5A). LC ADAM17 protein levels, however, were less affected, showing a 21% reduction

(Figure 5B). Intraperitoneal (IP) anti-IFNAR treatment at 500 µg/dose restored LC ADAM17

activity in MRL/lpr mice (Figure 5A) without increasing ADAM17 protein expression (Figure

5B). A lower 125 µg/dose of anti-IFNAR had similar effects in rescuing LC ADAM17 sheddase

370   function (Figure 5C-D). These results together suggested that IFN-I is important for inhibiting

LC ADAM17 sheddase activity in MRL/lpr mice.

Figure 5. Anti-IFNAR restores LC ADAM17 activity in multiple lupus models. (A-I)
MRL/lpr (A-D), B6.Sle1yaa (E-F), and IMQ (G-I) lupus model mice and their controls were
treated twice with anti-IFNAR or isotype control at indicated doses over 6 days prior to
collection of non-lesional epidermal cells. (A, C, E, G) UVR-induced LC ADAM17 sheddase
activity as in Figure 4. (B, D, F, H) Relative cell surface ADAM17 levels. (I) Relative LC
numbers per sample. (A-I) Each symbol represents a mouse, bars represent average values, and
error bars are SD. n=3-8 per condition over 3-4 independent experiments. *p<0.05, **p<0.01,
***p<0.001, n.s.=not significant by paired (A, C, E (left)) or unpaired ((A, C, E (right)), B, D, F-
I) t-test.

LCs from the B6.Sle1yaa model showed a 63% reduction in ADAM17 activity when compared to control mice, without a corresponding change in ADAM17 protein levels (Figure 5E-F). Similar to MRL/lpr mice, anti-IFNAR restored LC ADAM17 activity in B6.Sle1yaa mice without increasing ADAM17 protein expression (Figure 5E-F).

IMQ model mice showed reduced LC ADAM17 sheddase activity without altering cell surface ADAM17 protein levels (Figure 5G-H), similar to data from MRL/lpr and B6.Sle1yaa models. Interestingly, unlike the MRL/lpr and B6.Sle1yaa models where LC numbers were unchanged from controls (Shipman et al., 2018), skin LC numbers in the IMQ model were reduced by 35% (Figure 5I), echoing findings in human skin (Shipman et al., 2018). Similar to the MRL/lpr and B6.Sle1yaa models, anti-IFNAR restored LC ADAM17 activity in IMQ mice (Figure 5G). Together, the sufficiency of IFN-I in inhibiting LC ADAM17 activity and the restoration of LC ADAM17 activity by anti-IFNAR across 3 lupus models suggest that the IFN-I-rich environment drives LC ADAM17 dysfunction in non-lesional lupus skin.

Anti-IFNAR reduces photosensitivity in an EGFR and LC ADAM17-dependent manner

We asked whether the anti-IFNAR-mediated restoration of LC ADAM17 function also reduced photosensitive skin responses. Anti-IFNAR treatment at 500 μg/dose reduced ear swelling in MRL/lpr mice (Figure 6A-B) but did not reduce epidermal permeability, a readout of skin barrier function (Figure 6C). Furthermore, in control MRL/+ mice, anti-IFNAR actually increased UVR-induced ear swelling and epidermal permeability (Figure 6B-C), suggesting that anti-IFNAR increased photosensitivity in non-lupus mice. These results echoed findings by other groups that IFNAR deficiency in non-lupus B6 mice increased photosensitivity (Sontheimer et al., 2017) and reduced skin wound re-epithelialization (Gregorio et al., 2010), suggesting that the

Figure 6. Anti-IFNAR reduces photosensitivity in EGFR- and LC ADAM17- dependent manners. (A-I) MRL/lpr and (J-K) LCAd17 mice and controls were treated according to schematics in (A, G, J) and non-lesional ears were harvested. (B, D, H) Ear thickness. (C, E, I) Epidermal permeability as indicated by toluidine blue retention. (F, K) Neutrophils and monocytes per ear. (B-F, H-I, K). n=4-10 per condition over 4-6 independent experiments. *p<0.05, **p<0.01, ***p<0.001, n.s.=not significant by paired (B, D, H) and unpaired (C, E, F, I, K) t-test.

500 µg anti-IFNAR dose was disrupting intrinsic skin functions and potentially obscuring the

415   effects of LC ADAM17 activity restoration. Consistent with this idea, anti-IFNAR at 125

µg/dose still worsened MRL/+ ear swelling but no longer increased epidermal permeability

(Figure 6D-E), suggesting a decreased pathologic effect on homeostatic skin functions. This

lower dose, which was sufficient to restore LC ADAM17 activity in MRL/lpr mice (Figure 5C),

was sufficient to reduce both UVR-induced skin swelling and epidermal permeability (Figure

420   6D-E) and limit neutrophil and monocyte infiltration after UVR exposure (Figure 6F). At 125

µg/dose, then, anti-IFNAR rescued LC ADAM17 function and reduced UVR-induced skin

swelling, epidermal permeability, and inflammatory infiltrates in MRL/lpr mice.

We sought to understand the extent to which the anti-IFNAR-driven reduction in

photosensitivity reflected the rescue of LC ADAM17 activity. As LC ADAM17 limits

425   photosensitivity by stimulating EGFR signaling (Shipman et al., 2018), we blocked EGFR

activity upon anti-IFNAR treatment of MRL/lpr mice (Figure 6G). EGFR blockade with the

small molecule inhibitor PD16 abrogated the anti-IFNAR-mediated reduction in ear swelling and

epidermal permeability (Figure 6H-I), suggesting that anti-IFNAR requires EGFR signaling to

ameliorate photosensitivity in MRL/lpr mice. This is consistent with the idea that anti-IFNAR

430   reduces photosensitivity by restoring LC ADAM17 function.

In the IMQ model, anti-IFNAR also reduced inflammatory cell infiltrate in UVR-treated

skin (Figure 6J-K). In IMQ-treated LCAd17 mice lacking ADAM17 in LCs (Shipman et al.,

2018), however, anti-IFNAR failed to reduce neutrophil and monocyte accumulation (Figure

6K), suggesting that the ameliorative effects of anti-IFNAR on UVR-induced inflammation are

435   dependent on LC ADAM17. That anti-IFNAR reduced photosensitive responses across two

lupus models depended on EGFR signaling and LC ADAM17 provides strong support for the

idea that IFN-I contributes to photosensitivity at least in part by promoting LC ADAM17

dysfunction. These results also suggest that the beneficial effects of anifrolumab on SLE skin

may be attributable to the improvement of LC ADAM17 activity.

IFN-I inhibits UVR-induced LC ROS expression

UVR has been shown to trigger ADAM17 activity in fibroblasts via ROS generation

(Singh et al., 2009), and we asked if IFN-I modulation of ROS was associated with changes in

445 **Figure 7. LC ROS is reduced in lupus model mice and restored by anti-IFNAR.** (A-B)
Epidermal cell suspensions from indicated mice were exposed to UVR or not and LC ROS
content assayed by the ROS indicator CM-H2DCFDA. (A) WT mice; representative histograms
(left) and LC ROS content relative to non-exposed condition (right). Dashed "control" line
indicates signal without CM-H2DCFDA. (B) Control (-IMQ) and IMQ-treated mice (+IMQ)
450 were treated as in Figure 5G with anti-IFNAR or isotype IgG prior to harvest of non-lesional
skin; histograms (left) and fold change in LC ROS with UVR exposure (right). Each symbol
represents one mouse, bars represent average values, and error bars are SD. n=5-6 per condition
over 4-5 experiments. *p<0.05, **p<0.01, n.s.=not significant by unpaired t-test.

LC ADAM17 activity. We observed the expected UVR-induced ROS generation in LCs from

455   WT skin (Figure 7A). Skin from IMQ-treated mice showed a reduction in UVR-induced

LC ROS generation when compared to control skin, and anti-IFNAR treatment of IMQ mice

restored UVR-driven ROS generation (Figure 7B). These results suggest that IFN-I could

potentially limit UVR-induced LC ROS to inhibit LC ADAM17 function.

Together, our results suggested a model whereby the IFN-I-rich tissue microenvironment

460   in lupus skin causes dysfunction of UVR-induced LC ADAM17, leading to photosensitivity.

Anti-IFNAR restores this axis, suggesting that the therapeutic effects of anifrolumab on SLE

skin could, in part, be via restoration of LC function.

DISCUSSION

Here, we establish that multiple photosensitive lupus models resemble human lupus in showing upregulated IFN-I modules or IFN-regulated genes in non-lesional skin and that IFN-I in these lupus models causes LC ADAM17 dysfunction which contributes to photosensitive skin responses. These results suggest that the IFN-rich environment in lupus skin promotes

470 photosensitivity by causing Langerhans cell ADAM17 dysfunction and that the beneficial effects of anifrolumab on skin disease in SLE patients, in part, reflects restoration of LC function. Our results also suggest that photosensitivity in the MRL/lpr, B6.Sle1yaa, and IMQ lupus models shares pathogenic mechanisms with human lupus, underscoring the utility of these models for the study of lupus skin pathophysiology.

475 How the IFN-I-rich environment in lupus skin contributes to photosensitive responses has not been well understood. Kahlenberg and colleagues have shown that IFN-I can potentiate IL-6 expression and apoptosis of human keratinocytes (Sarkar et al., 2018; Stannard et al., 2017). Keratinocytes are likely a major local source of IFN-I to modulate LC function in non-lesional skin, as LCs are positioned in the suprabasal layer of the epidermis near basal and spinous

480 keratinocytes that demonstrate high IFN signaling (Billi et al., 2022). PDCs may also regulate LCs, as pDCs contribute to the IFN-rich environment in lupus skin (Furie et al., 2019; Karnell et al., 2021). Our results showing that anti-IFNAR restores LC ADAM17 function and reduces UVR-induced inflammation in an EGFR and LC ADAM17-dependent manner suggest that IFN-I contributes to photosensitivity by causing LC ADAM17 dysfunction. The accumulating

485 evidence points to a pathogenic feed-forward loop between LCs and keratinocytes whereby keratinocyte IFN contributes to LC ADAM17 dysfunction and LC ADAM17 dysfunction contributes to keratinocyte dysfunction and ultimate loss. It should be noted, however, that the

importance of IFN-I-mediated LC ADAM17 dysfunction does not rule out IFN-I-mediated keratinocyte-intrinsic mechanisms but points to multiple players in the ecosystem that maintain

490   skin health.

Our findings showing that anti-IFNAR treatment could increase UVR-induced skin inflammation in control MRL/+ mice underscores the idea that excessive IFN-I contributes to skin pathogenesis in lupus. IFNAR is important for re-epithelialization during wound healing (Gregorio et al., 2010) and our data support the idea that our treatments should not compromise

495   protective levels of IFNs. This notion is supported by the clinical findings that patients with high IFN-I scores responded to anifrolumab (Morand et al., 2020) and, when comparable, with a greater effect size than patients with lower IFN-I scores (Furie et al., 2017). Understanding the regulation and sources of the "excess" IFN-I in lupus will help to further develop therapies that ameliorate disease while preserving skin-intrinsic functions.

500   Our results extend our understanding of ADAM17 regulation as a contributor to disease. Increased ADAM17 activity can promote inflammation, such as by release of TNF or EGFR ligands within inflamed joints or lupus kidneys (Issuree et al., 2013; Qing et al., 2018) but loss of ADAM17 activity compromises barrier surfaces (Blaydon et al., 2011; Franzke et al., 2012). Here we identify IFN-I as a negative regulator of ADAM17 in LCs and our data suggest a

505   scenario whereby IFN-I reduces LC ROS generation to inhibit ADAM17 function. Further investigation will better delineate the locus and importance of LC ROS generation as well as contributions of other molecules such as the inactive Rhomboids (iRhoms) that regulate ADAM17 activity (Geesala et al., 2019).

High IFN-I states also include viral infections, genetic interferonopathies, and autoimmune

510   diseases rheumatoid arthritis and dermatomyositis (Muskardin and Niewold, 2018), in which

photosensitivity is found in the former and is a disease hallmark in the latter (Werth et al., 2004; Wysenbeek et al., 1989). Autoimmune disease-like manifestations that include photosensitivity are among the immune-related adverse events in cancer patients treated with checkpoint inhibitors (Brahmer et al., 2018), a situation where distinct IFN-I-mediated processes may have

515 specific roles in promoting anti-tumor immunity and treatment resistance (Benci et al., 2019). Our findings will benefit patients with lupus and a wide spectrum of diseases.

MATERIALS AND METHODS

Study design

520 The purpose of the study was to understand how LC function is regulated by the environment in non-lesional skin to contribute to photosensitivity. The subjects are both humans and mice. We used gene expression analyses to understand the gene expression profiles in human lupus and murine lupus models. We used ex-vivo flow cytometry-based assays, assessment of epidermal permeability, and skin swelling assays to understand the effects of IFN on LC function and

525 implications for photosensitivity. Sample sizes and replicate numbers are provided in each figure legend. Sample sizes were determined based on previous similar types of experiments. Mice were randomly assigned to treatment groups. No criteria were set for excluding certain data points and no data points were excluded. No specific confounder variables such as order of treatments or cage location were controlled for except that our experiments used multiple

530 independent cohorts of mice over replicate experiments. Assessments and analyses were not blinded. ARRIVE reporting guidelines (Percie du Sert et al., 2020) have been used.

Mice

8-16 week old mice were used unless otherwise specified. Mice were sex and age-matched. For

535 C57Bl/6 mice, LCAd17 (Langerin-cre; Adam17 f/f) mice, and wildtype littermate controls, males and females were used. Unless otherwise indicated, for MRL/+ and MRL/lpr mice, females were used at 8-12 weeks, while males were used at 12-16 weeks as females present with a disease phenotype earlier than males. For B6.Sle1yaa mice and their B6 controls, only males were used because the model depends on the autoimmune accelerator locus located on the Y

540 chromosome. B6 mice were either bred in-house or purchased from Jackson Laboratory (Bar

Harbor, ME). MRL/+, MRL/lpr mice, B6.Sle1yaa mice, and their B6 controls were either bred

in-house or purchased from Jackson Laboratory. LCAd17 mice (Shipman et al., 2018) were bred

in-house. Mice were kept in a specific pathogen-free barrier facility and all animal procedures

including protocols to reduce pain, suffering, and distress were performed in accordance with the

545 regulations of the Institutional Animal Care and Use Committee at Weill Cornell Medicine (New

York, NY).

Mouse treatments

For IFN-κ painting, 37.5 ng IFN-κ (R&D Systems, Minneapolis, MN) was dissolved in DMSO

550 (Sigma Aldrich, St. Louis, MO) and was applied topically onto each ear (18.75 ng on dorsal and

ventral sides, each). For anti-IFNAR1 treatments, mice were injected IP with 500 µg/dose or 125

µg/dose of anti-IFNAR1 (MAR1-5A3) or isotype control (MOPC-21) (Bio X Cell, West

Lebanon, NH). For UVR treatments, mice were exposed to 250 J/m^2 (25 mJ/cm^2)daily for four

consecutive days using a bank of four FS40T12 sunlamps that emitted a combination of UVA

555 and UVB radiation, similar to described (Shipman et al., 2018). Measurement using UVA and

UVB meters revealed that these lamps emit 2.1mW/cm2 UVA and 0.8mW/cm2 UVB. To

measure ear swelling after UVR exposure, a caliper (Mitutoyo, Aurora, IL) was used to take

three measurements throughout each ear, and the reported value is the average of both ears. Each

measurement was taken 22-24 hours after the previous measurement. For the IMQ-induced lupus

560 mouse model, mice were painted on the dorsal and ventral sides of both ears with 5% imiquimod

cream (42 mg/ear/mouse) 3x/week for 5-6 weeks (Yokogawa et al., 2014) prior to, and continued

through experiments. Indicated mice received 2.95 mg of the irreversible EGFR inhibitor

PD168393 (0.74 mg on dorsal and ventral sides, each) (Selleck Chem, Pittsburgh, PA).

Human subjects

For microarray analysis, seven discoid CLE patients were examined. Both lesional and non-lesional skin were collected and the gene expression data from lesional tissue of these patients have been published, and patient characteristics are described (Jabbari et al., 2014). Healthy controls include the three that were previously described (Jabbari et al., 2014). All human tissue collection and research use adhered to protocols approved by the Institutional Review and Privacy Boards at the Rockefeller University (IRB# AJA-00740) and New York University (IRB# 10-02117), where participants signed written informed consents.

For suction blistering, 1 male and 3 female healthy participants between the ages of 23 and 55 participated. All human tissue collection and research use adhered to protocols approved by the Institutional Review and Privacy Board at the Hospital for Special Surgery (IRB# 2019-1998), where participants signed written informed consents.

Human skin cell collection and preparation

Suction blistering was performed as previously described (27). Eight 5 mm suction blisters were generated with a negative pressure instrument (NP-4 model, Electronic Diversities, Finksburg, MD) on the arm of healthy donors over 30-60 minutes. The blister fluid was then collected by aspiration, and epidermal roofs were collected. After collection, epidermal roofs from suction blisters were digested with dispase (2.42 U/mL; Thermo Fisher Scientific, Waltham, MA), collagenase type II (616 U/mL; Worthington Biochemical Corporation, Lakewood, NJ), and DNAseI (80 μg/mL; Sigma Aldrich, St. Louis, MO) for 40 minutes to generate single cell suspensions that were used for the ADAM17 sheddase assay.

Skin cell collection and flow cytometry staining and quantification

For murine epidermal single cell suspensions, skin on the trunk was excised and subcutaneous

590   fat scraped off. Skin was incubated in dispase (2.42 U/mL) at 37°C. The epidermis was then

gently peeled, finely minced, digested in type II collagenase (616 U/mL) and DNAse I (80

µg/mL).

For collection of ear skin, ears were cut along the cartilage ridge, and the dorsal and ventral

sides were manually peeled. Ear skin was then finely minced and digested with a buffer

595   containing dispase, type II collagenase, and DNAse I as described above. For murine LC TNFR1

staining for the ADAM17 sheddase activity assay (see below), epidermal cells were preincubated

with Fc block and then stained so that we could gate on CD45+, CD3-, CD11c+, I-Ab+ for B6

mice and I-Ak+ for MRL mice, and EpCAM+ LCs and assess for TNFR1. LCs were sometimes

stained with rabbit-anti-ADAM17 (Lora et al., 2021) followed by donkey anti-rabbit IgG

600   (Jackson Immunoresearch, West Grove, PA). Monocytes and T cells were identified by gating

for CD45+, CD11b+, Ly6C+, and CD45+, CD11c-, I-Ak/I-Ab-, and CD3+ cells, respectively.

4',6-diamidino-2-phenylindole dihydrochloride (DAPI) was used to exclude dead cells and

debris (Sigma Aldrich, St. Louis, MO). All antibodies are from Biolegend, San Diego, CA unless

otherwise indicated.

605   For human LC TNFR1 staining for the ADAM17 sheddase activity assay, single cell

suspensions of the epidermal roofs of suction blisters and blister fluid were preincubated with Fc

block and subsequently stained for CD45+, CD11c+, CD14-, CD3- CD19-, CD56-, CD66b-,

EpCAM+, and HLA-DR+ to identify LCs. DAPI was used to exclude dead cells and debris, and

TNFR1 was also stained to assess ADAM17 activity. All antibodies are from Biolegend, San

Diego, CA unless otherwise indicated. Cells were analyzed using a FACS Canto or FACS Symphony A3 (BD Biosciences, San Jose, CA) and data were analyzed with FlowJo V10 Software (TreeStar, Ashland, OR).

For mean fluorescent intensity (MFI) calculations of TNFR1, each sample was assayed in triplicate, and the mean of the triplicates was recorded unless otherwise specified. The MFI of TNFR1 was divided by the MFI for the isotype control to generate a relative MFI. The percent TNFR1 MFI change was calculated by setting the relative MFI of the control sample to 1 and normalizing the value from the experimental samples to that control. In experiments with more than one control sample, the control values were averaged, and the experimental samples were each normalized to the averaged value.

For antibodies used, please see Table S9.

Ex vivo ADAM17 sheddase activity assay

For murine skin, cell suspensions of epidermis were suspended at 200,000-500,000 cells/mL medium. For MRL/LPR and B6.Sle1yaa skin, medium was RPMI with 2% FBS, HEPES buffer, L-glutamine, penicillin-streptomycin (all Thermo Fisher Scientific, Waltham, MA). For IMQ mice skin, medium was phenol red-free RPMI (Corning, Edison, New Jersey), no serum, with HEPES buffer, L-glutamine, penicillin-streptomycin. Cells were exposed to 500 (LPR and B6.Sle1yaa model mice) or 250 (IMQ model mice) J/m^2 UVR, similar to previous (Shipman et al., 2018), incubated for 45 minutes, and then stained for cell surface markers described above. For IFN-κ treatment, 3.125 ng/mL IFN-κ (R&D Systems, Minneapolis, MN) was added to the epidermal cell suspension that were incubated for 16-20 hours prior to UVR exposure. Indicated cells were treated with 1nM tofacitinib (Selleck Chem, Pittsburgh, PA) during IFN-κ incubation.

For human skin, cells were incubated for 1 hour with recombinant human IFN-β (35.7 ng/mL; activity=2.8×10^8 UI/mg) and then exposed to 500 J/m^2 UVR as above. Cells were further

635   incubated for 45 minutes and collected for preincubation with Fc block. Cells were then stained for cell surface markers described above.

Epidermal permeability assay

Epidermal barrier function was assessed as described (Shipman et al., 2018). Ear skin was

640   dehydrated and rehydrated in graded methanol and then incubated in 0.1% toluidine blue dye (Sigma Aldrich, St. Louis, MO). The dye was extracted with a buffer consisting of 95% methanol, 2.5% sulfuric acid, and 2.5% water and quantified with a colorimeter at 620 nm. Dye concentration was calculated using a standard curve.

645   **Ex vivo ROS assay**

Epidermal cell suspensions in phenol red-free RPMI were exposed to 250 J/m2 UVR and incubated with the ROS indicator 5-(and-6)-chloromethyl-2',7'-dichlorodihydrofluorescein diacetate, acetyl ester (CM-H2DCFDA) (Invitrogen, Waltham, MA) at 10uM for 45 minutes prior to cell surface marker staining for flow cytometry.

RNA extraction

Female MRL/+ and MRL/lpr 12-14 weeks old and 12 week old B6 and B6.Sle1yaa male mice were used for RNA sequencing. RNA was extracted from ear skin using an RNEasy Kit (Qiagen, Germantown, MD) and quality confirmed on a BioAnalyzer 2100 (Agilent Technologies, Santa

655 Clara, CA). For RT-qPCR of IMQ-treated mice, RNA was extracted with Trizol (Life

Technologies) following the manufacturer's protocol.

RNA sequencing

Libraries were prepared with non-stranded and poly-A selection TruSeq RNA Sample

660 Preparation kits. Sequencing libraries were sequenced by the Genomics Core Facility at Weill

Cornell Medicine using a HiSeq 2500 (MRL/lpr, MRL/+ samples) and NovaSeq (B6,

B6.Sle1yaa) at a depth of 15-30 million mappable paired-end reads per sample.

Computational analyses of gene expression data

665 Microarray analysis was performed using healthy controls from an existing GEO dataset

(GSE52471) obtained from GEO using GEOquery and combined with non-lesional skin sample

data kindly provided by the authors of the original study (Jabbari et al., 2014). Microarray data

were normalized with gcRMA normalization using the gcrma R package (Wu et al., 2021).

Differential gene expression analysis was performed with limma using the empirical bayes

670 method (Ritchie et al., 2015) and controlling for microarray hybridization kit and sex. Batch

correction was performed using ComBat (Leek et al., 2021). Visualizations were generated using

ComBat corrected data with plotly and ggplot2 in R.

 For RNA sequencing analysis, read quality filtering and adapter trimming were performed

using *fastp* (Chen et al., 2018). Filtered reads were mapped to the mouse genome (mm10), and

675 exonic reads were counted against GENCODE release 27 annotation with the STAR aligner

(Dobin et al., 2013) using a customized pipeline available

athttps://gitlab.com/hssgenomics/pipelines. Differential gene expression analysis was performed

with edgeR 3.30.2 using quasi-likelihood framework. Genes with low expression levels (<3

counts per million in at least one group) were filtered from all downstream analyses. P-values

680   were adjusted to correct for multiple testing using the Benhamini-Hochberg FDR procedure.

Genes with adjusted p-values < 0.01 were considered differentially expressed. Downstream

analyses were performed in R using a Shiny-driven visualization platform (RNAseq DRaMA)

developed at the HSS David Z. Rosensweig Genomics Research Center.

Differentially regulated pathways were identified using R QuSAGE 2.22.0 package (Yaari

685   et al., 2013) with MSigDB C2 set (curated gene sets). All gene sets with less than 10 genes were

excluded from the analysis. Pathways with Benhamini-Hochberg adjusted p-values <0.005 (non-

lesional skin versus healthy control) or <0.05 (MRL/lpr versus MRL/+) were used for further

analyses.

Ranked GSEA analysis was performed to define gene sets from MSigDB C2 enriched in

690   human non-lessional skin and MRL/lpr mice. Log-transformed fold changes in relevant

comparisons (non-lesional skin versus healthy control and MRL/lpr versus MRL/+) were used as

ranking variables. Pathways with Benhamini-Hochberg adjusted p-values <0.05 were selected

and the pathways' similarity network was built to visualize gene overlap between pathways

using the R/Bioconductor package *hypeR* (1.7.0) (Federico and Monti, 2020) and compared to

695   enriched pathways in human SLE patients and mouse models.

The R/Bioconductor package, GSVA (Hänzelmann et al., 2013)(v1.36.3), was utilized as

a non-parametric, unsupervised method to estimate enrichment of pre-defined gene sets in gene

expression data from human DLE skin and non-lesional skin from MRL/lpr and B6.Sle1yaa

mice. The inputs for GSVA were a matrix of expression values for all samples and curated gene

700   sets describing select immune/tissue cell types and inflammatory cytokines. Low-intensity

probes were filtered out if the interquartile range (IQR) of their expression values across all samples was not greater than 0. Enrichment scores were calculated using a Kolgomorov Smirnoff (KS)-like random walk statistic and represented the greatest deviation from zero for a particular sample and gene set. Scores across all samples were normalized to values between -1

705 (indicating no enrichment) and +1 (indicating enrichment). Gene sets used as input for GSVA are listed in Tables S3 and S7. The human gene sets were previously published (Martínez et al., 2022). The mouse plasma cell, T cell, myeloid cell, neutrophil, pDC, dendritic cell, and keratinocyte gene sets were derived from Mouse CellScan, a tool for the identification of cellular origin of mouse gene expression datasets. The mouse IFN gene sets have been previously

710 described (Kingsmore et al., 2021). The mouse cytokine signatures were generated by an iterative process involving derivation through literature mining and GO term definitions provided by the Mouse Genome Informatics (MGI) GO Browser (Bult et al., 2019).

qPCR

715 Both "lesional" and "non-lesional" ears were collected from each IMQ-treated and control mouse. Ears were flash-frozen with liquid nitrogen and stored at -80°C. Reverse transcription was performed using random primers and gene expression assessed by qPCR. PowerSYBR Green PCR Master Mix (Applied Biosystems) was used and assays were run on a QuantStudio 6 Pro (Applied Biosystems). Gene expression was normalized to the control for each target gene.

720 Primers used in the qPCR assay are listed in Table S10.

Statistical analyses

We determined the normality of data distribution using the Shapiro-Wilk test. For normally

distributed data, we used two-tailed unpaired and paired Student's t-tests for comparisons

725   between two conditions to evaluate p-values as indicated. For data that were not normally

distributed, the Mann-Whitney test was used for unpaired and Wilcoxon matched-pairs signed

rank test for paired comparisons. Significance was defined as $p<0.05$. GraphPad Prism software

was used. For figures showing normalized values, the control sample was set to one, and the

experimental samples were normalized relative to the control for that experiment. For

730   experiments that contained more than one control sample, the mean was obtained for the control

samples, and the individual control and experimental samples were calculated relative to this

mean.

Data availability

735   The murine RNAseq data have been deposited in GEO (accession ID : GSE222573). All other

data supporting the findings of this study are available within the paper and its Supplementary

Materials.

REFERENCES

Ambler, W., M.J. Howlader, M. Latha Somaraju, E. Seltzer, J. Sim, J. Shin, N. Schwartz, W.D. Shipman, D. Dasoveanu, C. Carballo, E. Sevim, S. Rodeo, D. Erkan, R.P. Kataru, B. Mehrara, and T. Lu. 2022. Lymphatic dysfunction lupus contributes to cutaneous photosensitivity and lymph node B cell responses. *bioRxiv : the preprint server for biology* 2022.2006.2013.495930.

Baechler, E.C., F.M. Batliwalla, G. Karypis, P.M. Gaffney, W.A. Ortmann, K.J. Espe, K.B. Shark, W.J. Grande, K.M. Hughes, V. Kapur, P.K. Gregersen, and T.W. Behrens. 2003. Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *PNAS* 100:2610-2615.

Benci, J.L., L.R. Johnson, R. Choa, Y. Xu, J. Qiu, Z. Zhou, B. Xu, D. Ye, K.L. Nathanson, C.H. June, E.J. Wherry, N.R. Zhang, H. Ishwaran, M.D. Hellmann, J.D. Wolchok, T. Kambayashi, and A.J. Minn. 2019. Opposing Functions of Interferon Coordinate Adaptive and Innate Immune Responses to Cancer Immune Checkpoint Blockade. *Cell* 178:933-948.e914.

Bennett, L., A.K. Palucka, E. Arce, V. Cantrell, J. Borvak, J. Banchereau, and V. Pascual. 2003. Interferon and Granulopoiesis Signatures in Systemic Lupus Erythematosus Blood. *J. Exp. Med.* 197:711-723.

Billi, A.C., F. Ma, O. Plazyo, M. Gharaee-Kermani, R. Wasikowski, G.A. Hile, X. Xing, C.M. Yee, S.M. Rizvi, M.P. Maz, C.C. Berthier, F. Wen, L.C. Tsoi, M. Pellegrini, R.L. Modlin, J.E. Gudjonsson, and J.M. Kahlenberg. 2022. Nonlesional lupus skin contributes to inflammatory education of myeloid cells and primes for cutaneous inflammation. *Science Translational Medicine* 14:eabn2263.

Blaydon, D.C., P. Biancheri, W. Di, V. Plagnol, R.M. Cabral, M.A. Brooke, D.A. van Heel, F. Ruschendorf, M. Toynbee, A. Walne, E.A. O'Toole, J.E. Martin, K. Lindley, T. Vulliamy, D.J. Abrams, T.T. MacDonald, J.I. Harper, and D.P. Kelsell. 2011. Inflammatory skin and bowel disease linked to ADAM17 deletion. *The New England Journal of Medicine* 365:1502-1508.

Brahmer, J.R., C. Lacchetti, B.J. Schneider, M.B. Atkins, K.J. Brassil, J.M. Caterino, I. Chau, M.S. Ernstoff, J.M. Gardner, P. Ginex, S. Hallmeyer, J. Holter Chakrabarty, N.B. Leighl, J.S. Mammen, D.F. McDermott, A. Naing, L.J. Nastoupil, T. Phillips, L.D. Porter, I. Puzanov, C.A. Reichner, B.D. Santomasso, C. Seigel, A. Spira, M.E. Suarez-Almazor, Y. Wang, J.S. Weber, J.D. Wolchok, J.A. Thompson, and N. National Comprehensive Cancer. 2018. Management of Immune-Related Adverse Events in Patients Treated With Immune Checkpoint Inhibitor Therapy: American Society of Clinical Oncology Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 36:1714-1768.

Bult, C.J., J.A. Blake, C.L. Smith, J.A. Kadin, and J.E. Richardson. 2019. Mouse Genome Database (MGD) 2019. *Nucleic Acids Res* 47:D801-d806.

Catalina, M.D., P. Bachali, N.S. Geraci, A.C. Grammer, and P.E. Lipsky. 2019. Gene expression analysis delineates the potential roles of multiple interferons in systemic lupus erythematosus. *Commun Biol* 2:140.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34:i884-i890.

Der, E., S. Ranabothu, H. Suryawanshi, K.M. Akat, R. Clancy, P. Morozov, M. Kustagi, M. Czuppa, P. Izmirly, H.M. Belmont, T. Wang, N. Jordan, N. Bornkamp, J. Nwaukoni, J. Martinez, B. Goilav, J.P. Buyon, T. Tuschl, and C. Putterman. 2017. Single cell RNA sequencing to dissect the molecular heterogeneity in lupus nephritis. *JCI Insight* 2:

Der, E., H. Suryawanshi, P. Morozov, M. Kustagi, B. Goilav, S. Ranabothu, P. Izmirly, R. Clancy, H.M. Belmont, M. Koenigsberg, M. Mokrzycki, H. Rominieki, J.A. Graham, J.P. Rocca, N. Bornkamp, N. Jordan, E. Schulte, M. Wu, J. Pullman, K. Slowikowski, S. Raychaudhuri, J. Guthridge, J. James, J. Buyon, T. Tuschl, and C. Putterman. 2019. Tubular cell and keratinocyte single-cell

transcriptomics applied to lupus nephritis reveal type I IFN and fibrosis relevant pathways. *Nat Immunol* 20:915-927.

790   Dobin, A., C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T.R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15-21.

Federico, A., and S. Monti. 2020. hypeR: an R package for geneset enrichment workflows. *Bioinformatics* 36:1307-1308.

Franzke, C.-W., C. Cobzaru, A. Triantafyllopoulou, S. Löffek, K. Horiuchi, D.W. Threadgill, T. Kurz, N. van
795   Rooijen, L. Bruckner-Tuderman, and C.P. Blobel. 2012. Epidermal ADAM17 maintains the skin barrier by regulating EGFR ligand-dependent terminal keratinocyte differentiation. *The Journal of Experimental Medicine* 209:1105-1119.

Furie, R., M. Khamashta, J.T. Merrill, V.P. Werth, K. Kalunian, P. Brohawn, G.G. Illei, J. Drappa, L. Wang, and S. Yoo. 2017. Anifrolumab, an Anti-Interferon-α Receptor Monoclonal Antibody, in
800   Moderate-to-Severe Systemic Lupus Erythematosus. *Arthritis Rheumatol* 69:376-386.

Furie, R., V.P. Werth, J.F. Merola, L. Stevenson, T.L. Reynolds, H. Naik, W. Wang, R. Christmann, A. Gardet, A. Pellerin, S. Hamann, P. Auluck, C. Barbey, P. Gulati, D. Rabah, and N. Franchimont. 2019. Monoclonal antibody targeting BDCA2 ameliorates skin lesions in systemic lupus erythematosus. *J Clin Invest* 129:1359-1371.

805   Geesala, R., P.D. Issuree, and T. Maretzky. 2019. Novel functions of inactive rhomboid proteins in immunity and disease. *J Leukoc Biol* 106:823-835.

Goel, R.R., X. Wang, L.J. O'Neil, S. Nakabo, K. Hasneen, S. Gupta, G. Wigerblad, L.P. Blanco, J.B. Kopp, M.I. Morasso, S.V. Kotenko, Z.X. Yu, C. Carmona-Rivera, and M.J. Kaplan. 2020. Interferon lambda promotes immune dysregulation and tissue inflammation in TLR7-induced lupus. *Proc*
810   *Natl Acad Sci U S A* 117:5409-5419.

Gregorio, J., S. Meller, C. Conrad, A. Di Nardo, B. Homey, A. Lauerma, N. Arai, R.L. Gallo, J. Digiovanni, and M. Gilliet. 2010. Plasmacytoid dendritic cells sense skin injury and promote wound healing through type I interferons. *J Exp Med* 207:2921-2930.

Hänzelmann, S., R. Castelo, and J. Guinney. 2013. GSVA: gene set variation analysis for microarray and
815   RNA-Seq data. *BMC Bioinformatics* 14:7.

Issuree, P.D., T. Maretzky, D.R. McIlwain, S. Monette, X. Qing, P.A. Lang, S.L. Swendeman, K.H. Park-Min, N. Binder, G.D. Kalliolias, A. Yarilina, K. Horiuchi, L.B. Ivashkiv, T.W. Mak, J.E. Salmon, and C.P. Blobel. 2013. iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. *J Clin Invest* 123:928-932.

820   Jabbari, A., M. Suárez-Fariñas, J. Fuentes-Duculan, J. Gonzalez, I. Cueto, A.G. Franks, Jr., and J.G. Krueger. 2014. Dominant Th1 and minimal Th17 skewing in discoid lupus revealed by transcriptomic comparison with psoriasis. *J Invest Dermatol* 134:87-95.

Karnell, J.L., Y. Wu, N. Mittereder, M.A. Smith, M. Gunsior, L. Yan, K.A. Casey, J. Henault, J.M. Riggs, S.M. Nicholson, M.A. Sanjuan, K.A. Vousden, V.P. Werth, J. Drappa, G.G. Illei, W.A. Rees, and J.N.
825   Ratchford. 2021. Depleting plasmacytoid dendritic cells reduces local type I interferon responses and disease activity in patients with cutaneous lupus. *Sci Transl Med* 13:

Kingsmore, K.M., P. Bachali, M.D. Catalina, A.R. Daamen, S.E. Heuer, R.D. Robl, A.C. Grammer, and P.E. Lipsky. 2021. Altered expression of genes controlling metabolism characterizes the tissue response to immune injury in lupus. *Sci Rep* 11:14789.

830   Kirou, K.A., C. Lee, S. George, K. Louca, I.G. Papagiannis, M.G. Peterson, N. Ly, R.N. Woodward, K.E. Fry, A.Y. Lau, J.G. Prentice, J.G. Wohlgemuth, and M.K. Crow. 2004. Coordinate overexpression of interferon-alpha-induced genes in systemic lupus erythematosus. *Arthritis Rheum* 50:3958-3967.

Lambers, W.M., K. de Leeuw, B. Doornbos-van der Meer, G.F.H. Diercks, H. Bootsma, and J. Westra. 2019. Interferon score is increased in incomplete systemic lupus erythematosus and correlates with myxovirus-resistance protein A in blood and skin. *Arthritis Research & Therapy* 21:260.

Leek, J.T., W.E. Johson, H.S. Parker, E.J. Fertig, A.E. Jaffe, Y. Zhang, J.D. Storey, and L. Callado Torres. 2021. sva: Surrogate Variable Analysis. . *R package version 3.40.0*

Lessard, J.C., S. Piña-Paz, J.D. Rotty, R.P. Hickerson, R.L. Kaspar, A. Balmain, and P.A. Coulombe. 2013. Keratin 16 regulates innate immunity in response to epidermal barrier breach. *Proc Natl Acad Sci U S A* 110:19537-19542.

Lora, J., G. Weskamp, T.M. Li, T. Maretzky, D.T.N. Shola, S. Monette, S.F. Lichtenthaler, T.T. Lu, C. Yang, and C.P. Blobel. 2021. Targeted truncation of the ADAM17 cytoplasmic domain in mice results in protein destabilization and a hypomorphic phenotype. *J Biol Chem* 296:100733.

Martin-Fernandez, M., M. Bravo García-Morato, C. Gruber, S. Murias Loza, M.N.H. Malik, F. Alsohime, A. Alakeel, R. Valdez, S. Buta, G. Buda, M.A. Marti, M. Larralde, B. Boisson, M. Feito Rodriguez, X. Qiu, M. Chrabieh, M. Al Ayed, S. Al Muhsen, J.V. Desai, E.M.N. Ferre, S.D. Rosenzweig, B. Amador-Borrero, L.Y. Bravo-Gallego, R. Olmer, S. Merkert, M. Bret, A.K. Sood, A. Al-Rabiaah, M.H. Temsah, R. Halwani, M. Hernandez, F. Pessler, J.L. Casanova, J. Bustamante, M.S. Lionakis, and D. Bogunovic. 2020. Systemic Type I IFN Inflammation in Human ISG15 Deficiency Leads to Necrotizing Skin Lesions. *Cell Rep* 31:107633.

Martínez, B.A., S. Shrotri, K.M. Kingsmore, P. Bachali, A.C. Grammer, and P.E. Lipsky. 2022. Machine learning reveals distinct gene signature profiles in lesional and nonlesional regions of inflammatory skin diseases. *Sci Adv* 8:eabn4776.

Morand, E.F., R. Furie, Y. Tanaka, I.N. Bruce, A.D. Askanase, C. Richez, S.C. Bae, P.Z. Brohawn, L. Pineda, A. Berglind, and R. Tummala. 2020. Trial of Anifrolumab in Active Systemic Lupus Erythematosus. *N Engl J Med* 382:211-221.

Muskardin, T.L.W., and T.B. Niewold. 2018. Type I interferon in rheumatic diseases. *Nature reviews. Rheumatology* 14:214-228.

Nickerson, K.M., J.L. Cullen, M. Kashgarian, and M.J. Shlomchik. 2013. Exacerbated Autoimmunity in the Absence of TLR9 in MRL.Fas[lpr] Mice Depends on Ifnar1. *The Journal of Immunology* 190:3889-3894.

Percie du Sert, N., V. Hurst, A. Ahluwalia, S. Alam, M.T. Avey, M. Baker, W.J. Browne, A. Clark, I.C. Cuthill, U. Dirnagl, M. Emerson, P. Garner, S.T. Holgate, D.W. Howells, N.A. Karp, S.E. Lazic, K. Lidster, C.J. MacCallum, M. Macleod, E.J. Pearl, O.H. Petersen, F. Rawle, P. Reynolds, K. Rooney, E.S. Sena, S.D. Silberberg, T. Steckler, and H. Würbel. 2020. The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *PLoS Biol* 18:e3000410.

Pisitkun, P., J.A. Deane, M.J. Difilippantonio, T. Tarasenko, A.B. Satterthwaite, and S. Bolland. 2006. Autoreactive B cell responses to RNA-related antigens due to TLR7 gene duplication. *Science* 312:1669-1672.

Qing, X., Y. Chinenov, P. Redecha, M. Madaio, J.J. Roelofs, G. Farber, P.D. Issuree, L. Donlin, D.R. McLlwain, T.W. Mak, C.P. Blobel, and J.E. Salmon. 2018. iRhom2 promotes lupus nephritis through TNF-alpha and EGFR signaling. *J Clin Invest* 128:1397-1412.

Reefman, E., H. Kuiper, P.C. Limburg, C.G. Kallenberg, and M. Bijl. 2008. Type I interferons are involved in the development of ultraviolet B-induced inflammatory skin lesions in systemic lupus erythaematosus patients. *Ann Rheum Dis* 67:11-18.

Ritchie, M.E., B. Phipson, D. Wu, Y. Hu, C.W. Law, W. Shi, and G.K. Smyth. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43:e47.

Sarkar, M.K., G.A. Hile, L.C. Tsoi, X. Xing, J. Liu, Y. Liang, C.C. Berthier, W.R. Swindell, M.T. Patrick, S. Shao, P.-S. Tsou, R. Uppala, M.A. Beamer, A. Srivastava, S.L. Bielas, P.W. Harms, S. Getsios, J.T.

Elder, J.J. Voorhees, J.E. Gudjonsson, and J.M. Kahlenberg. 2018. Photosensitivity and type I IFN responses in cutaneous lupus are driven by epidermal-derived interferon kappa. *Annals of the Rheumatic Diseases* 77:1653-1664.

885 Shipman, W.D., S. Chyou, A. Ramanathan, P.M. Izmirly, S. Sharma, T. Pannellini, D.C. Dasoveanu, X. Qing, C.M. Magro, R.D. Granstein, M.A. Lowes, E.G. Pamer, D.H. Kaplan, J.E. Salmon, B.J. Mehrara, J.W. Young, R.M. Clancy, C.P. Blobel, and T.T. Lu. 2018. A protective Langerhans cell–keratinocyte axis that is dysfunctional in photosensitivity. *Science Translational Medicine* 10:

Singh, B., M. Schneider, P. Knyazev, and A. Ullrich. 2009. UV-induced EGFR signal transactivation is
890 dependent on proligand shedding by activated metalloproteases in skin cancer cell lines. *International Journal of Cancer* 124:531-539.

Sontheimer, C., D. Liggitt, and K.B. Elkon. 2017. Ultraviolet B Irradiation Causes Stimulator of Interferon Genes–Dependent Production of Protective Type I Interferon in Mouse Skin by Recruited Inflammatory Monocytes. *Arthritis & Rheumatology* 69:826-836.

895 Stannard, J.N., T.J. Reed, E. Myers, L. Lowe, M.K. Sarkar, X. Xing, J.E. Gudjonsson, and J.M. Kahlenberg. 2017. Lupus Skin Is Primed for IL-6 Inflammatory Responses through a Keratinocyte-Mediated Autocrine Type I Interferon Loop. *J Invest Dermatol* 137:115-122.

Strassner, J.P., M. Rashighi, M. Ahmed Refat, J.M. Richmond, and J.E. Harris. 2017. Suction blistering the lesional skin of vitiligo patients reveals useful biomarkers of disease activity. *J Am Acad Dermatol*
900 76:847-855.e845.

Subramanian, S., K. Tus, Q.Z. Li, A. Wang, X.H. Tian, J. Zhou, C. Liang, G. Bartov, L.D. McDaniel, X.J. Zhou, R.A. Schultz, and E.K. Wakeland. 2006. A Tlr7 translocation accelerates systemic autoimmunity in murine lupus. *Proc Natl Acad Sci U S A* 103:9970-9975.

Tsoi, L.C., G.A. Hile, C.C. Berthier, M.K. Sarkar, T.J. Reed, J. Liu, R. Uppala, M. Patrick, K. Raja, X. Xing, E.
905 Xing, K. He, J.E. Gudjonsson, and J.M. Kahlenberg. 2019. Hypersensitive IFN Responses in Lupus Keratinocytes Reveal Key Mechanistic Determinants in Cutaneous Lupus. *The Journal of Immunology* 202:2121-2130.

Werth, V.P., M. Bashir, and W. Zhang. 2004. Photosensitivity in rheumatic diseases. *The journal of investigative dermatology. Symposium proceedings* 9:57-63.

910 Wu, J., R. Irizarry, J. MadDonald, and J. Gentry. 2021. gcrma: Background Adjustment Using Sequence Information. . *R package version 2.64.0*

Wysenbeek, A.J., D.A. Block, and J.F. Fries. 1989. Prevalence and expression of photosensitivity in systemic lupus erythematosus. *Annals of the Rheumatic Diseases* 48:461-463.

Yaari, G., C.R. Bolen, J. Thakar, and S.H. Kleinstein. 2013. Quantitative set analysis for gene expression: a
915 method to quantify gene set differential expression including gene-gene correlations. *Nucleic Acids Res* 41:e170.

Yokogawa, M., M. Takaishi, K. Nakajima, R. Kamijima, C. Fujimoto, S. Kataoka, Y. Terada, and S. Sano. 2014. Epicutaneous application of toll-like receptor 7 agonists leads to systemic autoimmunity in wild-type mice: a new model of systemic Lupus erythematosus. *Arthritis Rheumatol* 66:694-706.

920 Zhang, X., D. Bogunovic, B. Payelle-Brogard, V. Francois-Newton, S.D. Speer, C. Yuan, S. Volpi, Z. Li, O. Sanal, D. Mansouri, I. Tezcan, G.I. Rice, C. Chen, N. Mansouri, S.A. Mahdaviani, Y. Itan, B. Boisson, S. Okada, L. Zeng, X. Wang, H. Jiang, W. Liu, T. Han, D. Liu, T. Ma, B. Wang, M. Liu, J.Y. Liu, Q.K. Wang, D. Yalnizoglu, L. Radoshevich, G. Uzé, P. Gros, F. Rozenberg, S.Y. Zhang, E. Jouanguy, J. Bustamante, A. García-Sastre, L. Abel, P. Lebon, L.D. Notarangelo, Y.J. Crow, S.
925 Boisson-Dupuis, J.L. Casanova, and S. Pellegrini. 2015. Human intracellular ISG15 prevents interferon-α/β over-amplification and auto-inflammation. *Nature* 517:89-93.

ACKNOWLEDGEMENTS

Acknowledgements: The authors thank the members of the Lu Lab for helpful discussions and critical reading of the manuscript.

Funding:

Alpha Omega Alpha Carolyn L. Kuckein Student Research Fellowship (TML)

Erwin Schrodinger Fellowship J 4638-B FWF (VZ)

940 National Institutes of Health grant T32AR071302-01 to the Hospital for Special Surgery Research Institute Rheumatology Training Program (NS and WDS)

National Institutes of Health grant MSTP T32GM007739 to the Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-PhD Program (WDS)

Tow Foundation (YC and DJO)

National Institutes of Health grant K08 AR069111 to the University of Iowa Department of Dermatology (AJ)

Veterans Administration VA Merit I01 BX004907 (AJ),

Physician Scientist Career Development Award from the Dermatology Foundation (AJ)

955 National Institutes of Health grant R01AR077194 (AJ)

National Institutes of Health grant R01 DK099087 (IR)

National Institutes of Health grant R35GM134907 (CPB)

National Institutes of Health grant R01AI079178 (TTL)

National Institutes of Health grant R21 AR081493 (TTL)

965 Department of Defense grant W81XWH-21-LRP-IPA (TTL)

Lupus Research Alliance (TTL)

St. Giles Foundation (TTL)

Barbara Volcker Center for Women and Rheumatic Diseases (TTL)

A Lasting Mark Foundation (TTL)

975 National Institutes of Health Office of the Director grant S10OD019986 to Hospital for Special Surgery.

AUTHOR CONTRIBUTIONS

Conceptualization: TML, TTL

980 Performed experiments and acquired data: TML, VZ, ESS, MD, KRV, NS, AJ

Data analyses and interpretation: TML, VZ, ESS, MD, YC, ARD, DJO, IR, CPB, PEL, TTL

Methodology and reagents: JL, YL, WDS, WGA, JHZ, MR, NA, CPB

Supervision: SFT, KBO, JGK, IR, CPB, PEL, TTL

Writing—original draft: TML, VZ, MD, YC, AD, IR, PEL, TTL

985 Writing—review and editing: All authors

Competing interests:

Andrea R. Daamen and Peter E. Lipsky are employees of AMPEL BioSolutions, but have no financial conflicts of interest to report. Jonathan H. Zippin is a stockholder in YouV labs, Hoth Therapeutics, and FoxWayne Inc and is a paid consultant for Pfizer and Hoth Therapeutics, Inc.
990 Carl Blobel holds a patent on a method of identifying agents for combination with inhibitors of iRhoms. Carl Blobel and the Hospital for Special Surgery have identified iRhom2 inhibitors and have co-founded the start-up company SciRhom in Munich to commercialize these inhibitors. Theresa T. Lu is a paid consultant for Pfizer.

SUPPLEMENTARY MATERIALS

Figure S1. Further analyses of gene expression in human DLE and multiple murine lupus models.

Figure S2. ADAM17 expression levels in LCAd17 mice, UVR-induced cell surface TNFR1 loss on LCs in mixed epidermal cell suspensions is dependent on LC ADAM17.

Figure S3. Langerhans cell yield and gating with suction blistering of human skin.

Table S1. Differentially expressed genes in DLE vs healthy control non-lesional skin microarray

Table S2. Differentially expressed pathways in DLE vs healthy control non-lesional skin using QuSAGE

Table S3. Human gene sets used for GVSA

Table S4. Differentially expressed genes in MRL/lpr vs control MRL/+ non-lesional skin RNAseq

Table S5. Differential expressed pathways in MRL/lpr vs control MRL/+ non-lesional skin using QuSAGE

Table S6. Differentially expressed genes in B6.Sle1yaa vs control B6 non-lesional skin RNAseq

Table S7. Murine gene sets used for GVSA

Table S8. Ranked GSEA enriched pathways in DLE vs normal control dataset

Table S9. Antibodies used

Table S10. Primer sequences for qPCR