Environment dependent benefits of inter-individual variation in honey bee recruitment

Supraja Rajagopal[1,2], Axel Brockmann[3], Ebi Antony George[3*]

[1] Indian Institute of Science Education and Research, Kolkata, India

[2] University of Georgia, Athens, USA

[3] National Centre for Biological Sciences, Bangalore, India – 560065

ORCID
S.R.: 0000-0003-3322-2243
A.B.: 0000-0003-0201-9656
E.A.G.: 0000-0002-5533-5428

* corresponding author:

Ebi Antony George

Email ID: ebiantonyg@ncbs.res.in, ebi.antony.george@gmail.com

Abstract

Inter-individual differences in behaviour within the members of a social group can affect the group's productivity. In eusocial insects, individual differences amongst workers in a colony play a central role division of labour and task allocation. Extensive empirical and theoretical work has highlighted variation in response thresholds as a proximate mechanism underlying individual behavioural differences and hence division of labour. However, other response parameters, like response probability and intensity, can affect these differences. In this study, we first extended a previously published agent-based model on honey bee foraging to understand the relative importance of response (dance) probability and response (dance) intensity in the task of recruitment. Comparing variation obtained from the simulations with previously published empirical data, we found that response intensity plays a more important role than probability in producing consistent inter-individual differences in recruitment behaviour. We then explored the benefits provided by this individual variation in recruitment behaviour to the colony's collective foraging effort under different environmental conditions. Our results revealed that individual variation leads to a greater energetic yield per forager, but only when food is abundant. Our study highlights the need to consider all response parameters while studying division of labour and adds to the growing body of evidence linking individual variation in behavioural responses to the success of social groups.

Keywords

Division of labour, repeatability of behaviour, division of labour, recruitment, waggle dance, response thresholds, response probability, response intensity

Introduction

Individuals from a population can vary in their behavioural decisions, even when responding to the same stimuli (Dall et al., 2012). Behavioural syndromes refer to the phenomenon in which individual variation in behavioural response is consistent over time and correlated across contexts (Sih et al., 2004). Consistent inter-individual differences in behaviour have been observed in a wide variety of taxa and diverse behavioural contexts, like aggression, courtship, and foraging behaviour (Bell et al., 2009). Differences in behavioural responses lead to behavioural specialisation amongst individuals and thereby influences how populations respond to perturbations (Wolf and Weissing, 2012). In eusocial insects, inter-individual differences in behaviour have been studied extensively in the context of behavioural specialisation associated with division of labour (Dall et al., 2012; Loftus et al., 2021).

Division of labour contributes to the phenomenal success of eusocial insect societies (Hölldobler and Wilson, 2009). Eusocial insect colonies consist of highly related workers who perform different tasks, like brood care and foraging, necessary for the survival and growth of the colony (Wilson and Hölldobler, 2005). Moreover, individuals performing the same task show differences in task performance, which can be consistent over time (Jeanson and Weidenmüller, 2013). Individual differences in task performance are linked to variation in the perception of the stimulus intensity as well as differences in the behavioural threshold of responding to the stimulus intensity (Beshers and Fewell, 2001). Theoretical studies on response thresholds suggest that variation in these thresholds is essential to maintain division of labour and ensure efficient utilisation of the workers in social insect colonies (Bonabeau et al., 1996; Johnson, 2010; Theraulaz et al., 1998). Empirical studies on response thresholds in a variety of tasks, like brood care and thermoregulation, have confirmed the link between thresholds and division of labour (Modlmeier et al., 2012; Weidenmüller, 2004).

Unfortunately, the focus on response thresholds has come at the cost of neglecting other

response parameters that may be important for division of labour, like response probability

and intensity (Cook et al., 2020; Jeanson and Weidenmüller, 2013; Ulrich et al., 2021).

An individual's task performance for a given stimulus intensity is a combination of both its

probability of responding to the stimulus and the intensity of its response, either in terms of

the number or duration of response events (George and Brockmann, 2019; Weidenmüller,

2004). While thresholds determine only the minimum stimulus intensity at which individuals

start responding, probability and intensity determine its response at a particular stimulus

intensity. Individual variation in these response parameters can provide social insect colonies

with additional flexibility in allocating their workforce and in responding to perturbations.

Thus, a comprehensive understanding of the benefits of division of labour requires an

exploration of these response parameters in addition to response thresholds. Despite their

importance, empirical and theoretical studies on these response parameters in social insects

are limited (Cook et al., 2020; Jeanson and Weidenmüller, 2013). An important requirement

for such studies is a behavioural paradigm in which the responses of multiple individuals to

the same stimulus can be quantified repeatedly.

The honey bee recruitment process consists of individual foragers, repeatedly visiting the

same food source, using the highly stereotyped waggle dance behaviour to inform nestmates

of the location and reward offered at the food source (von Frisch, 1967). An individual

foragers motivation to recruit is dependent on both the reward offered by the food source as

well as the colony's nutritional state (De Marco and Farina, 2001; Nieh, 2010; Seeley, 1994;

Thom, 2003). Foragers active at the same food source experience the same stimulus intensity,

due to the same food reward and colony conditions, and hence the recruitment behaviour is

an excellent paradigm to explore individual differences in behavioural responses. Recent

work revealed that foragers show consistent inter-individual variation in their response

probability (number of dances per trip) and intensity (average number of waggle phases per

dance) of dance activity and, as a result, in their total recruitment activity for the same food

source (George and Brockmann, 2019). While empirical studies have sought to understand

how individual variation in recruitment behaviour benefits social insect colonies (Jones,

2004; Mattila and Seeley, 2010; Modlmeier et al., 2012), these are often difficult due to the

need to control for multiple sources of individual variation (Jeanson and Weidenmüller,

2013; Ulrich et al., 2021). Simulations of the recruitment process can be a useful approach to

understand how and under what conditions individual variation can potentially benefit social

insect colonies, and thereby help design further behavioural experiments.

In this study, we explored how individual variation in recruitment activity affects the

colony's foraging behaviour by expanding upon a previously published agent-based model of

the honey bee foraging process (Schürch and Grüter, 2014). We incorporated individual

variation in response probability and intensity of the dance behaviour in the foraging process

and perform two sets of experiments. In the first set of experiments, we estimated the relative

importance of response probability and intensity in maintaining individual variation in the

total recruitment activity of the agents. We did this by simulating a honey bee colony

foraging under conditions mimicking experimental studies and validated the individual-

specific parameters by comparing variation in the total recruitment activity in the simulations

with empirical data (George and Brockmann, 2019). In the second set of experiments, we

quantified the relative benefit of individual variation in recruitment activity under different

environmental food availability conditions. By examining individual variation in recruitment

behaviour from the perspective of both the cause of this variation in terms of response

parameters and the consequences of this variation for the colony's collective foraging

activity, our study provides important insights into the role of inter-individual variation in

social insect colonies.

Methods

We simulated an agent-based model (ABM) of honey bee nectar foragers with inter-individual variation in the probability and intensity of their recruitment behaviour. Our model is built on top of an earlier ABM (Schürch and Grüter, 2014), which we modified to incorporate individual-level parameters. We used NETLOGO 5.3.1 to implement the model, and follow the "Overview, Design Concepts and Details" protocol to describe it (Grimm et al., 2006).

Purpose

The overall purpose of the model was to explore the role of individual variation in the probability and intensity of recruitment behaviour amongst honey bee foragers in the colony's foraging activity. We first quantified the effect of individual variation in probability and intensity in maintaining consistent inter-individual variation in dance activity for a particular food source. We calibrated the individual level parameter values by comparing variation in the total dance activity obtained from the simulations with empirical data (George and Brockmann, 2019). We then explored if individual variation in recruitment behaviour benefited the colony's foraging efforts under different environmental conditions. Both probability and intensity (average number of waggle phases per dance) of dance activity is correlated with the forager's perception of the food reward (Barron et al., 2007; George and Brockmann, 2019; Seeley, 1994, 1989). In our model, we implemented two individual state parameters: a probability modulator and an intensity modulator. The probability modulator regulates the probability that a honey bee performs a waggle dance on returning to the hive after foraging. The intensity modulator regulates the number of waggle phases that a bee makes in each waggle dance for a given nectar reward.

State variables and scales

In all our model runs we simulated 300 foragers from one colony. The foragers were divided into two different types, scouts (n = 30) and recruits (n = 270), based on typical proportions of scouts and recruits in an *A. mellifera* colony (Seeley, 1995). We were interested in the consistency of recruitment activity to exploited food sources and hence focussed on the recruitment activity of the recruits. Scouts explore the environment for new food sources to exploit and usually abandon food sources even when they are still rewarding (Liang et al., 2012; Seeley, 1983). Recruits on the other hand get recruited to a food source (and can further recruit to the food source) and continue exploiting the food source till it becomes less rewarding (Biesmeijer and De Vries, 2001; Seeley, 1983). We were interested in individual variation in the recruitment activity of these recruits, as they bring in the bulk of the food necessary for the survival and growth of the colony. The states that individual agents could take in our model was the same as in the previously published model. Scouts could, at any time step, be in any of the 6 states (Fig. S1 *a*): idle in the colony, scouting for food sources, feeding at a food source, returning to the colony, recruiting of idle foragers to the newly discovered food source, returning to the nest without having discovered food. Recruits could be in any of the following 8 states (Fig. S1 *b*) at any time step: idle in the nest, waiting to be recruited, flying to a food source, feeding on a food source, returning to colony after feeding, recruiting new idle foragers, scouting for forage, returning to the colony without having discovered food. All agents were characterized by an identity number, a dance probability modulator, and a dance intensity modulator.

Our agents were located on a two-dimensional square grid with 201 x 201 patches. Following the previous model, the width of a patch corresponded to 100m in our model also. Thus, our agents could explore an area corresponding to a 20 km x 20 km square centred on the hive. Agents could occupy any position in this area and were not restricted to the edges of the

patch. The agents' hive was located at the centre patch and no food sources were present in

this patch throughout the simulations.

Two different sets of experiments were performed which differed in the number and

persistence of the food availability in the environment. In experiment 1, we simulated a

foraging environment for our agents with no temporal change in the distribution of nectar

sources for a short duration of 5 days. This was done to ensure that we could easily compare

our simulation results with previous experimental data (George and Brockmann, 2019). In

experiment 2, we simulated a more natural foraging environment with multiple transient food

sources for a longer duration of 51 days. Similar to the earlier published model (Schürch and

Grüter, 2014), we tested three different environmental conditions in this experiment: low

food density, medium food density and high food density.

Simulations were run in discrete time-steps. Each time step corresponded to 10 seconds in

real-time and each day consisted of 5000 steps, with agents active up to the 4000^{th} time step

on each day (corresponding to 11.11 hours of real time). The weather was not simulated, and

so the general foraging activity of the colony remained similar throughout the day.

Process overview and scheduling

Each simulation was run for 5 days (experiment 1) or 51 days (experiment 2), consisting of a

short non-experimental phase at the beginning (2 days for experiment 1 and 3 days for

experiment 2). The non-experimental phase allowed foragers to learn about the location of

food sources around their colony. Scouts and recruits followed similar processes while

foraging (Fig. S1 a and b). The recruitment process in our model was based on the earlier

model. After unloading the nectar, individual agents could dance to recruit one other agent to

the food source, with the probability of recruitment dependent on the duration of the dance.

The probability that a bee will dance on returning to the hive was determined by her

probability modulator, the nectar quality in terms of sugar concentration (Tautz and

Sandeman, 2003; Waddington, 1982), the distance of the nectar source from the hive (Seeley,

1994) and the current influx of returning foragers (Seeley, 1989). The quality of the food

sources and their distances from the hive was kept the same throughout, and the current

influx of returning foragers was the same for bees dancing at the same time step. Thus, in our

model, individual differences in dance probability for foragers visiting the same food source

were a direct result of individual differences in the probability modulator. If a bee decides to

dance, the intensity of her dance is determined by nectar source distance, nectar quality

(Seeley, 1994) and her dance modulator. Individual differences in dance intensity for foragers

visiting the same food source were a direct result of individual differences in the intensity

modulator.

Scouts and recruits became idle after dancing. Agents in an idle state could leave the nest

with probability p_{exit}. Scouts constantly searched for new food sources, while recruits showed

high patch fidelity (Al Toufailia et al., 2013; Moore and Doherty, 2009). Each day recruits,

that had previously foraged, decided to either continue foraging or abandon the patch. The

probability to abandon the patch depended on the individual probability modulator, nectar

quality and nectar source distance.

Experiment 1: Individual variation in probability and intensity and consistent differences in
recruitment behaviour

We quantified how much individual variation in probability and intensity of dancing

contributes to consistent inter-individual differences in the total dance activity of individual

foragers. To measure the consistency of dance activity, we ensured that foragers had access to

four food sources of the same quality (same sucrose concentration at the same distance from

the hive in the four cardinal directions). The food source was available throughout the day,

and we quantified the dance activity of foragers at this food source for 5 days (2 days to

acclimatize followed by a 3-day experimental phase). We quantified individual differences in

the dance activity in the experimental phase and obtained the repeatability estimates of the

total dance activity. This was then compared to empirical repeatability estimates obtained

from observations of groups of foragers trained to an ad-libitum food source over 3 days

(George and Brockmann, 2019). The comparison allowed us to calibrate the variation in

probability and intensity modulators to obtain empirical levels of variation in recruitment

activity and further quantify the relative importance of probability and intensity in

maintaining consistent differences in recruitment activity (through a sensitivity analysis).

Experiment 2: Individual variation in recruitment behaviour and colony foraging activity

Environmental conditions can affect the benefits of information transfer through the

recruitment process in social insects, with the greatest benefit often seen in low food density

conditions (Dornhaus et al., 2006; Goy et al., 2021; Schürch and Grüter, 2014). In this

experiment, we explored how inter-individual variation in the recruitment behaviour further

benefits the colony's foraging activity. We hypothesise that the benefits of inter-individual

variation in recruitment behaviour will be dependent on the environmental conditions. Inter-

individual variation in the recruitment behaviour was implemented as a function of individual

variation in both response probability and intensity, which we obtained from the previous

experiment. We tested the colony's foraging activity under three different environmental

conditions: low, medium, and high food density (d_{patch} corresponding to 0.01, 0.05 and 0.1

respectively).

In both experiments 1 and 2, we tested four different variations of our model:

Model 1: Honey bee forager agents were all similar with no variation in individual state

variables of probability modulator (I_p) and intensity modulator (I_i).

Model 2: Every agent was characterised by a different value of I_p and the same value of I_i.

Honey bee foragers showed inter-individual differences in their probability of dancing for

any given food reward. The probability modulator for each agent was obtained from a normal

distribution [Fig. S2 a; mean ± standard deviation (sd) of I_p = 0.255 ± 0.1].

Model 3: Every agent was characterised by a different value of I_i and the same value of I_p.

Honey bee foragers showed inter-individual differences in their intensity of dances for any

given food reward. The intensity modulator for each agent was obtained from a normal

distribution (Fig. S3 a; mean ± sd of I_i = 0.255 ± 0.1).

Model 4: Each agent was characterized by a different value of I_p and I_i, both of which were

obtained from normal distributions (mean ± sd = 0.255 ± 0.1). Honey bee foragers showed

inter-individual differences in both the probability and intensity of dancing for any given

food reward.

An agent with the mean values of I_p and I_i in models 2, 3 and 4 was similar to all agents in

model 1.

Design Concepts

In our model, individual honey bee foragers collected food from various food sources

offering different sucrose concentrations present throughout the environment around the hive

which was centrally located. The amount of food brought back is equivalent to the energy

procured by the colony and thus its fitness.

Environmental Stimuli: The honey bee foragers could evaluate whether the time of day was

suitable for foraging and the quality of the food sources. The probability and intensity of

dancing depended on this quality. Additionally, the probability of recruits to persist at the

food source also depended on this quality.

Biotic stimuli: Scouts and recruits could pass on the information about nectar source location

to other recruits. Recruits mostly flew out only after receiving this information.

Stochasticity: The probability and intensity modulators were assigned stochastically. They

were picked randomly from a normal distribution and assigned to the agents. Agent decisions

regarding leaving the hive, dancing for a food source, learning the location of a nectar source

and abandoning a nectar source were all probabilistically determined.

Observation: Different emergent properties of the model were observed in experiments 1 and

2. In experiment 1, the food patch visited and the number of dance circuits in each dance was

recorded and used to obtain the total number of dance circuits made by each agent for each

day. In experiment 2, the mean energy yield per forager, the median foraging distance, and

the mean search time in addition to the mean recruitment activity for each day were

quantified. The former three properties represent mutually non-exclusive mechanisms by

which inter-individual variation in recruitment can benefit the colony's foraging effort:

through an increase in the energy brought in by foragers, a reduction in the distance flown by

foragers to find a rewarding food source and a reduction in the time spent by foragers

searching for food sources independently (because of an increased probability of recruitment

to a rewarding food source).

Initialisation

All agents were idle at the start of the simulation. Environmental food availability at the start

of the model depended on the experiment. In experiment 1, four equally rewarding (0.8 M

sucrose concentration) nectar sources were available in the environment at a distance of 80

patches from the hive. The four patches were distributed around the nest with a 90° angle

between consecutive patches. These patches never got exhausted for the duration of the

simulations (5 days), similar to artificial feeders used in behavioural experiments with honey

bees (Seeley, 1995). In experiment 2, multiple food sources of varying quality were available

in the environment around the hive at the beginning of each simulation as initialised in the

previously published model (Schürch and Grüter, 2014). The number of food sources

depended on the environmental food density (d_{patch}). Each patch in the simulated world was

randomly allocated as a foraging patch or not with a probability equal to d_{patch}. The quality of

the food source was a random value from a normal distribution with a mean of q_{patch} and a

standard deviation of σ_{patch}. All food sources were finite with a maximum age (a_{max}) of 14

days. The current age of the patch at the beginning of the simulation was a random value

between 1 and 14 with equal probability.

Sub models

The Lévy flight

The movement of agents during a search followed an optimal Lévy flight pattern (Reynolds

et al., 2009). This sub model was obtained from Schürch and Grüter, 2014. Agents moved in

bouts after initially choosing a random direction. Before the start of the bout, a total travel

distance for the bout was calculated using the formula:

296
$$d_{L\acute{e}vy} = e^{\ln a * \frac{1}{1 - \mu_{L\acute{e}vy}}}$$

Here, *a* was obtained randomly from a uniform distribution bound by [0,1). The value for

$\mu_{L\acute{e}vy}$ was kept at 2.4 by default (Reynolds et al., 2009). Once the distance of the bout was

calculated, agents would then move in the chosen direction in subsequent time steps, till they

had reached the end of the distance. Then, the whole process was repeated till the agent found

a food source.

Individual decision to dance for food sources

On returning to the hive, successful scouts and recruits would first unload the nectar that they

had brought into the hive. They could then communicate information about the location of

the food source to their nest-mates by performing a waggle dance. The forager would decide

to dance only after the agent has finished unloading the nectar brought back from foraging.

The equation that linked the probability of dancing to the food reward was

308
$$p_{qt} = \frac{0.956937}{1 + e^{\left(\frac{I_p - \ln(q_t)}{0.2624014}\right)}}$$

309   Here I_p was the individual probability modulator and q_t was the energetic value of a food

310   source, given by

311
$$q_t = 1.02 + q_{patch} - 0.04915919 - 2.851457.10^{-5}.d + 3.174098.d^2$$

where q_{patch} was the sucrose concentration of the food source and d was the distance of the

food source from the hive.

Our equation was modified from the earlier model, which in turn used data from previous

empirical studies (Boch, 1956; Seeley et al., 1991). We incorporated individual probability

modulators, thereby giving each individual agent a different quantitative relationship between

food quality and dance probability. The mean value for the probability modulator was

obtained based on two assumptions: 1) all individual agents had the same qualitative

relationship between patch quality and dance probability as described in Schürch and Grüter,

2014 and 2) the probability modulators of the population of agents were normally distributed.

We chose the mean and standard deviation of the probability modulator based on the average

value of the curves fit for each agent, such that our two assumptions were met.

The dance probability also depended on the influx of other foragers,

324
$$p_{Influx} = \begin{cases} 1 - 0.035.N_{Influx} & N_{Influx} \leq 20 \\ 0.3 & N_{Influx} > 20 \end{cases}$$

325   Thus, the final dance probability was given by

326
$$p_d = p_{qt} \cdot p_{Influx}$$

327 *Individual decision about the intensity of dances*

328 Dance intensity is the number of waggle phases that a bee makes per dance. The relationship

329 between food quality and the number of waggle phases is linear (Seeley, 1994). In our model,

330 each bee was assigned a quantitatively different slope that predicted how much they would

331 dance after returning to the hive.

332
$$dance_{intensity} = 57 \times q_t \times I_i$$

Here I_i referred to the individual intensity modulator and q_t was the energetic value of a food

source.

To obtain the value of the intensity modulator, we made 4 assumptions: 1) all individual

agents had the same qualitative linear relationship between patch quality and dance intensity

as observed in Seeley, 1994, 2) the intensity modulators of the population of agents was

normally distributed, 3) all agents had the same intercept of zero for the linear link between

patch quality and dance intensity, i.e., there was no dance activity for a patch of zero quality

and 4) there is an upper limit to the number of waggle phases for the most rewarding food

source due to energetic constraints (Seeley, 1995). Based on these assumptions, we fit

individual curves for each agent and chose the mean and standard deviation of the intensity

modulator.

The intensity of the dance determines the duration of the dance. In the previously published

model, all dances had the same intensity and lasted for 10 seconds (equal to the minimum

duration for an agent to change its state). In our model, the more intense the dance, the longer

it lasted. Each dance circuit lasted for around 3.33 seconds such that within each discrete

time-step in the model, an agent could complete 3 dance circuits. As a result, agents with

steeper intensity curves (higher I_i values) spent more time performing the recruitment

behaviour. This had a corresponding effect on the agents recruited through the waggle dance.

In the previous model (Schürch and Grüter, 2014), each dance had a 25% chance to recruit a

forager, in accordance with empirical data (Dornhaus et al., 2006; Gould et al., 1970; Seeley

and Towne, 1992; Seeley and Visscher, 1988; Tautz, 1996). In our model, we adjust the

recruitment process so that every 10 dance circuits had a 25% chance to recruit a forager.

Thus, longer dances probabilistically recruited more agents.

Individual decision to abandon food sources

The probability of abandoning the nectar source was modelled analogously to the probability

of dancing (Schürch and Grüter, 2014; Seeley, 1989) with the probability modulator playing

a role in the decision to abandon a food source:

360
$$p_a = 1 - \frac{0.956937}{1 + e^{(\frac{I_p - \ln(q_t + 0.6687)}{0.2624014})}}$$

Thus, there was a correlation between an individual's decision to dance for a food source and

the decision to abandon the food source, tied to the individual's perception of the food reward

(George and Brockmann, 2019; Seeley, 1989).

Sensitivity Analysis

We ran each experimental simulation multiple times to account for the stochasticity in the

models. In experiment 1, each of the 4 variations of the model was run 100 times. In

experiment 2, each of the 12 combinations of model and food density variation (4 variations

of the model and 3 different food densities) were run 40 times.

Further, to test how sensitive our model was to changes in our two individual-level response

parameters, we ran a sensitivity analysis based on model 4 in experiment 1. We chose this

experiment as the controlled food availability allowed us to check the relative effect of

probability and intensity on consistent inter-individual variation in recruitment activity. We

chose this model as it incorporates individual variation in probability and intensity of

recruitment, unlike the other 3 models. We varied the distribution of the probability (Fig. S2

b) and intensity modulators (Fig. S3 b), to change the extent of inter-individual differences in

probability and intensity. The probability equation was modified as shown below to vary the

spread of the individual modulators:

378
$$p_a = 1 - \frac{0.956937}{1 + e^{(\frac{(m*I_p)-\ln(q_t)}{0.2624014})}}$$

where m took a value in between 0.25 and 2 in steps of 0.25. To change the spread of the

individual variation in intensity, we changed the standard deviation of the normal distribution

of I_i, while keeping the mean constant. The standard deviation was changed from 0.01 to 0.1

in steps of 0.01. We then quantified the consistency in inter-individual variation in the total

recruitment activity for each of the 80 combinations (8 levels of variation in I_p and 10 levels

of variation in I_i) and ran 10 simulations of each version for a total of 800 simulations. We

varied the spread of the two modulators in this way to keep the coefficient of variation of

both modulators similar. This allowed us to directly compare the effect of variation in

probability and intensity in maintaining consistent inter-individual differences in recruitment

behaviour.

Sensitivity analyses regarding other parameters in the model, like the colony size, probability

of scouts to exit the nest p_{exit}, Levi-flight parameter $\mu_{Lévy}$ and energy costs of recruiting were

explored in the previous study (Schürch and Grüter, 2014). The model used in their analysis

was analogous to model 1 from experiment 2 in this study with medium food density.

Statistical Analysis

Experiment 1

In this experiment, we were interested in quantifying the effect of individual variation in

probability and intensity of dance activity on maintaining consistent inter-individual

differences in recruitment behaviour. To do this, we first shortlisted foragers who consistently

visited the same patch over the three days to minimise any variation in dance activity due to a

forager switching states. As scouts abandoned food sources readily, they were automatically

removed in our analysis. We then used generalized linear mixed-effects models (GLMMs)

with the total dance activity in a day as the response variable and individual agent identity as

the random effect and a Poisson error distribution to quantify the repeatability of recruitment

behaviour. Repeatability estimates provide a quantification of the consistency in inter-

individual differences for a behaviour and are based on a comparison of intra-individual

variation to inter-individual variation (Nakagawa and Schielzeth, 2010). We obtained a single

repeatability estimate for each of the 100 simulations runs of each experiment. The

repeatability estimates were then compared using a beta regression model with the estimate

as the response variable and the model as the predictor (categorical variable of 4 levels,

corresponding to models 1, 2, 3 and 4). A beta regression model was used as the values of

repeatability estimates was a proportion between 0 and 1. We compared the 4 different

models against each other and corrected for multiple comparisons using Tukey's HSD.

Experiment 2

In this experiment, we were interested in quantifying the effect of individual variation in

recruitment behaviour on the colony's collective foraging activity. We observed 4 different

output parameters on each day for all the simulation runs: mean dance time per forager, mean

yield per forager, median foraging distance and mean search time per forager. We then

compared these parameters across the 4 different models and the 3 different environmental

food density conditions using linear mixed-effects models (LMMs). We built separate LMMs

for each parameter, with the parameter of interest as the response variable, an interaction

between the food density (categorical variable of 3 levels) and the model (categorical variable

of 4 levels) as the predictor and day nested within simulation number as the random effect.

We compared the effect of the different models against each other within each level of food

density and corrected for multiple comparisons using Tukey's HSD.

Sensitivity Analysis

For the sensitivity analysis, we first obtained the repeatability estimates for all 800 simulation

runs as in experiment 1. Then, a LOESS model was fit to this data with the mean

repeatability estimate for each of the 80 combinations of probability and intensity modulator

as the response variable and an interaction between the coefficient of variation in the

probability modulator and the coefficient of variation in the intensity modulator as the

predictor variable. We then predicted the repeatability estimates for other combinations of

variation in the probability and intensity modulator from this fitted model. Finally, contour

plots were used to visually examine the effect of the variation in probability and intensity

modulators on the repeatability of dance activity.

We built and ran the models in NETLOGO 5.3.1 (Wilensky, 1999). The model outputs were

cleaned up using the pandas package (McKinney and others, 2010) in Python (Van Rossum

and Drake, 2009). We used R software (R Core Team, 2018) along with the RStudio IDE

(RStudio Team, 2016) to perform all the statistical analyses. We built the GLMMs using the

rptR package in R (Stoffel et al., 2017). For the beta regression, we used the betareg package

(Cribari-Neto and Zeileis, 2010), for LMMs we used the glmmTMB package (Brooks et al.,

2017) and for multiple comparisons, we used the emmeans package (Lenth, 2019). We fit the

LOESS model using the base package in R. We validated the model assumptions for each of

the mixed-effects models using visual inspections of residual and normality plots through the

DHARMa package (Hartig, 2018). For data visualisation, we used the ggplot2 (Wickham,

2016) and cowplot (Claus O. Wilke, 2018) packages in R. All the code produced as part of

this manuscript can be found at the Zenodo repository (George et al., 2021).

Results

Experiment 1: Individual variation in probability and intensity and consistent differences in recruitment behaviour

Figure 1: Repeatability estimates of recruitment behaviour (i.e., total dance activity) of agents in the model. (a) Violin plots of the distribution of repeatability estimates from 1200 simulation runs of the four different models. Inset boxplots show the median (central black line), the first and third quartiles (lower and upper edges of the box) and 1.5 times the interquartile range between the first and third quartile (whiskers). Alphabets above each violin plot represent the results of the post-hoc pair-wise comparison of repeatability estimates between the different models, with different alphabets representing significant differences at the $p < 0.05$ level. (b) Comparison of the repeatability estimates obtained from models 3 and 4 with estimates obtained from empirical observations (data in George and Brockmann, 2019). The circles represent the repeatability estimate (mean of the estimates in the case of the models) and the error bars represent the 95% confidence interval. The grey shaded region represents the 95% confidence interval of the repeatability estimate from empirical observations.

Incorporating individual variation in probability and intensity increased the consistency of inter-individual differences in recruitment activity amongst the agents in the simulations (Table S2 and Fig. 1a, models 2, 3 and 4 all had significantly higher repeatability estimates compared to model 1 at the $p < 0.001$ level). However, repeatability estimates of the total recruitment activity were much higher in model 3 which only incorporated individual differences in intensity, as compared to model 2 which only incorporated individual

differences in probability (Table S2 and Fig. 1a, model 3 - model 2: repeatability estimate

difference = 0.524, z-ratio = 82.911, p < 0.001). Further model 4, which incorporated both

individual variation in probability and intensity, had lower repeatability estimates compared

to model 3, which had only individual differences in intensity (Table S2 and Fig. 1a, model 3

- model 4: repeatability estimate difference = 0.139, z-ratio = 18.152, p < 0.001). Both model

3 and model 4 had mean repeatability estimates comparable to empirical observations (Fig.

1b). However, estimates obtained from model 4 were the closest to empirically obtained

repeatability estimates.

Figure 2: Contour plot of the sensitivity analysis on the effect of individual variation in
probability and intensity of dancing on the repeatability estimates of recruitment behaviour.
The contour plot is coloured based on the value of the repeatability estimate predicted for
each combination of coefficient of variation in probability and intensity.

Our sensitivity analysis corroborated results from experiment 1. Individual variation in

intensity had a stronger effect on maintaining consistent inter-individual differences in dance

activity at the colony level as compared to a similar level of individual variation in

probability (Fig. 2).

Experiment 2: Individual variation in recruitment behaviour and colony foraging
activity

Incorporating inter-individual variation in recruitment behaviour affected the colony's

foraging dynamics in an environmental dependent manner, with the strongest effects at

medium and high food densities (Table S3 and Fig. 3). Inter-individual variation in

recruitment behaviour led to significantly shorter recruitment time at the group level under all

food density conditions (Fig. 3c, model 1 – model 4: low food density – difference estimate =

3, t-ratio = 10.08, p < 0.001; medium food density – difference estimate = 2.32, t-ratio = 7.77,

p < 0.001; high food density – difference estimate = 2.27, t-ratio = 7.63, p = < 0.001). There

was a similar trend in the opposite direction in the mean yield with individual variation in

recruitment behaviour leading to higher yield at medium and high food densities (Fig. 3d,

model 1 – model 4: low food density – difference estimate = 0.64, t-ratio = 0.74, p = 0.881;

medium food density – difference estimate = -3.08, t-ratio = -3.55, p = 0.002; high food

density – difference estimate = -3.44, t-ratio = -3.95, p < 0.001).

There was no significant difference in the median foraging distance between the model with

and without individual variation in probability and intensity in any of the food density

conditions (Fig. S4, model 1 – model 4: low food density – difference estimate = 0.001, t-

ratio = 0.08, p > 0.999; medium food density – difference estimate = -0.003, t-ratio = -0.2, p

= 0.997; high food density – difference estimate = -0.009, t-ratio = -0.66, p = 0.912). The

mean search time per forager in the model with and without individual variation in

recruitment activity followed the same pattern as the median foraging distance, with no

difference in any of the food density conditions (Fig. S4, model 1 – model 4: low food

density – difference estimate = -0.54, t-ratio = -0.22, p = 0.996; medium food density –

difference estimate = -0.001, t-ratio = -0.0005, p > 0.999; high food density – difference

estimate = 1.16, t-ratio = 0.48, p = 0.964).

Figure 3: Effect of individual variation in recruitment behaviour on the colony's foraging activity under different environmental conditions. (a and b) Line plots showing the change in (a) mean dance time per forager in seconds and (b) mean yield per forager in joules throughout the simulations. The central line represents the mean value of all agents on each day from 40 simulations while the shaded region represents the standard deviation around this mean. (c and d) Comparison of the (c) mean dance time and (d) mean yield per forager, averaged over the whole simulation duration for all 40 simulations, across models and food density conditions. Circles represent the mean and error bars represent the standard deviation. The lines and shaded region in (a) and (b) and the circles and error bars in (c) and (d) are

coloured according to the model: model 1 – brown, model 2 – green, model 3 – blue, model 4
– teal.

Incorporating individual variation in only one of the two response parameters did not produce

the same decrease in the mean time spent dancing per forager as incorporating individual

variation in both parameters did (Table S2 and Fig. 3, comparison of model 4 versus models

2 and 3). On the other hand, it produced similar levels of mean yield, especially in the high

food density conditions. Further, the median foraging distance and mean search time was

comparable across all models in the medium and high food density conditions, with a slight

increase in the median foraging distance and mean search time in the low food density

condition (Table S2 and Fig. S4).

Discussion

Our simulations revealed two important aspects of individual variation in recruitment activity

amongst honey bees. First, variation in response intensity contributed more to producing

consistent variation in recruitment activity as compared to variation in response probability.

At the same time, incorporating individual variation in both response parameters was

necessary to recapture empirically observed levels of consistent inter-individual variation in

recruitment activity. Second, individual variation in recruitment activity led to a decrease in

the average recruitment activity of the colony and benefitted the colony's foraging activity by

increasing the mean yield per forager. However, this benefit of individual variation in

recruitment activity was only apparent at medium and high food density conditions, and not

at low food density conditions. By focussing on the response parameters responsible for

individual variation in task performance, our study adds to the growing body of literature

highlighting the importance of response probability and intensity in addition to response

thresholds in social insect division of labour.

The three response parameters may play different roles in modulating individual differences

in task performance. Empirical work on individual variation in honey bee recruitment

behaviour showed that response intensity is weakly correlated with gustatory response

thresholds whereas response probability is not (George et al., 2019). Similarly, a study on

variation in recruitment behaviour at the level of patrilines revealed no correlation between

thresholds and probability (Mattila and Seeley, 2010). Our simulation results now reveal that

although individual variation in both response probability and intensity is necessary to obtain

empirical levels of inter-individual differences in recruitment activity, the latter is more

important. At the same time, there is evidence that response probability may play a vital role

in the social modulation of recruitment. Social interactions between foragers and receivers

who unload the nectar inform foragers of the nutritional state of the colony and can change

their recruitment behaviour (Cook et al., 2020; De Marco and Farina, 2001; Seeley, 1989).

We could not simulate these interactions in greater detail due to the different spatial scales of

these interactions (within the hive, over centimetres) and the foraging behaviour (outside the

hive, over kilometres). However, in a previous experiment, manipulating the nectar influx,

and thereby changing these social interactions of foragers (Seeley, 1989), affected the

probability but not the intensity of the recruitment behaviour (George and Brockmann, 2019).

Taken together, these results imply that response intensity and threshold is likely linked to an

internal behavioural state responsible for innate variation in the recruitment activity of

individual foragers. Modulation of response probability via social interactions allows the

colony further flexibility in utilising this basal level of inter-individual variation.

Variation in response thresholds is generally considered to be the major physiological basis

underlying task specialisation and division of labour in social insects (Beshers and Fewell,

2001; Page and Mitchell, 1998). The simplest models on division of labour assume that

individuals have fixed response thresholds, whereas more complex models involve flexible

individual thresholds which are modulated by factors like task performance and social

interactions (Theraulaz et al., 1998). In contrast, a few empirical studies have indicated that

individual differences in task performance within social insect colonies are often based on an

intricate relationship between thresholds and two other response parameters: response

probability and response intensity (Cook et al., 2020; Jeanson and Weidenmüller, 2013;

Ulrich et al., 2021; Weidenmüller, 2004). The relative importance of the response parameters

in task performance may vary depending on the task and the social context. In bumblebees,

for example, a combination of response probability and intensity was most predictive of

individual variation in task performance during thermoregulation (Garrison et al., 2018).

Response thresholds did not predict individual task performance and differed in the same

individuals when they were observed singly and in social groups. In honey bee recruitment,

response probability is sensitive to social signals, whereas response intensity remains

unaffected (George and Brockmann, 2019). Differences in the role of probability and

intensity were also apparent in our simulations, and a useful direction for future simulations

would be to incorporate individual variation in thresholds into our model.

Individual variation in recruitment behaviour can provide additional benefits to a social insect

colony on top of those provided by recruitment itself. Our results reveal potential trade-offs

during the collective foraging process in honey bees, specifically between recruitment to and

exploitation of a food source. Inter-individual differences in the waggle dance would allow

some foragers to recruit more foragers to a newly discovered food source while others can

continue exploiting it. This division of labour within groups of foragers active at the same

food source leads to an overall decrease in the average recruitment activity in our

simulations. Individual differences in recruitment should also ensure that recruitment is

further biased towards highly profitable food sources (Seeley, 1994, 1989, 1986). The high

reward offered by these food sources should also lead to continuous exploitation and lowered

abandonment by foragers. In line with this, there was an increase in the colony's productivity

in models incorporating individual variation in recruitment. At the same time, the benefits of

individual variation in recruitment are likely to be seen after the discovery of food sources

and not before. Consistent with this expectation, individual variation in recruitment did not

affect the median foraging distance of the foragers or the average time spent searching for

food sources by the foragers in our simulations.

The benefits of spatially directed recruitment, e.g., through the waggle dance, is more

apparent under heterogeneous environmental conditions with low food densities (Bailis et al.,

2010; Beekman and Lew, 2007; Dornhaus et al., 2006; Goy et al., 2021; Schürch and Grüter,

2014, but see I'Anson Price et al., 2019). When food is scarce, a decrease in recruitment

activity should lead to a decrease in the colony's productivity due to lower number of

foragers being directed to the most profitable food sources. On the other hand, when food

sources are plentiful, the energy and time costs of recruitment would likely outweigh the

benefits provided by this behaviour (Dornhaus et al., 2006; I'Anson Price and Grüter, 2015;

Schürch and Grüter, 2014). Our simulations reveal an under-appreciated role of inter-

individual differences in spatially directed recruitment at high food density conditions. When

food is plentiful, individual differences lead to a lowered average recruitment activity as well

as a corresponding increase in the energy gain for the colony. Thus, these differences can

offset the energy and time costs of recruitment and ensure that recruitment is still beneficial

to the colony under such environmental conditions. One important aspect of most studies on

the benefits of the waggle dance, including the current study, is that they are limited to *A.*

mellifera. The other extant honey bee species show marked differences in their behaviour and

ecology in comparison with *A. mellifera* (I'Anson Price and Grüter, 2015), particularly with

respect to the waggle dance (Beekman et al., 2015; E. A. George et al., 2021; Kohl et al.,

2020). Species in temperate regions (e.g., *A. mellifera* and some populations of *A. cerana*)

have a shorter time window for foraging throughout the year compared to species in tropical

regions (e.g., *A. florea*, *A. dorsata* and *A. cerana*) but must establish large winter food stores

(Ji et al., 2020). The role of individual variation under these different environmental

conditions is unknown and comparative studies on individual variation within the genus *Apis*

would be an important avenue for future research.

Individual variation in social groups is used as a bet-hedging strategy to buffer against

environmental variation (Honegger and de Bivort, 2018; Hopper, 1999). However, the

perspectives behind the studies on this variation have differed between eusocial groups and

other social groups (Dall et al., 2012). In eusocial insects, theoretical and empirical studies

across species (honey bees, bumblebees, and ants) and tasks (brood care, thermoregulation,

foraging and recruitment) have focussed on the role played by individual variation in task

allocation (Beshers and Fewell, 2001; Jeanson and Weidenmüller, 2013; Mattila and Seeley,

2010; Modlmeier et al., 2012; Weidenmüller et al., 2019; Westhus et al., 2013; Wiernasz et

al., 2008). On the other hand, in social groups, individual variation has been studied in terms

of the range of behavioural responses a group has access to, which can, in turn, affect its

productivity, e.g., in the case of foraging in birds and fishes (Aplin et al., 2014; Dyer et al.,

2009; Wolf and Weissing, 2012). Irrespective of these different perspectives, individual

variation in behavioural responses ultimately affects group success across social groups.

Intriguingly, a new theoretical framework proposes to integrate concepts from task allocation

in eusocial insects and animal personality studies in social groups (Loftus et al., 2021). A

careful exploration of such a framework (Pinter-Wollman, 2021) would enable a better

understanding of the link between consistent inter-individual differences in behavioural

responses and division of labour in social insects.

In conclusion, our study reveals both the relative importance of two response parameters in

producing consistent inter-individual differences in task performance during recruitment and

the effect of these differences on a honey bee colony's foraging activity. These results

corroborate results from previous experimental and theoretical studies on the benefits of

inter-individual variation for collective task performance in animal groups including social

insects (Aplin et al., 2014; Michelena et al., 2010; Seeley, 1994). They also highlight the role

played by individual differences in offsetting the costs of recruitment at the group level under

different environmental conditions (Beekman and Lew, 2007; Dornhaus et al., 2006; Schürch

and Grüter, 2014). Importantly, our study adds to the accumulating evidence on the

importance of multiple response parameters in driving task performance. While a theoretical

framework has been established for exploring the role of response thresholds in task

allocation and division of labour (Loftus et al., 2021; Weidenmüller et al., 2019), there is an

urgent need to incorporate different response parameters into this framework to obtain a

comprehensive understanding of how social insect groups function.

Data Accessibility

The NETLOGO code to run these simulations, the datasets obtained, and the Python and R

codes used for the analysis are available from Zenodo via the DOI: 10.5281/zenodo.5217630.

Funding

E.A.G. was supported by the National Centre for Biological Sciences – Tata Institute of

Fundamental Research Graduate school. A.B. was supported by National Centre for

Biological Sciences – Tata Institute of Fundamental Research institutional funds (12P4167).

A.B. also acknowledges the support of the Department of Atomic Energy, Government of

India, under 472 project no. 12-R&D-TFR-5.04-0800.

Author Contributions

Conceptualization: S.R., A.B., E.A.G.; Methodology: S.R., E.A.G.; Software: S.R., E.A.G.;

Formal analysis: E.A.G.; Investigation: S.R., E.A.G.; Writing – original draft: S.R., A.B.,

E.A.G.; Writing – review & editing: S.R., A.B., E.A.G.; Supervision: A.B., E.A.G.; Funding

acquisition: A.B.

Acknowledgements

References

Al Toufailia, H., Grüter, C., Ratnieks, F.L.W., 2013. Persistence to Unrewarding Feeding Locations by
Honeybee Foragers (Apis mellifera): the Effects of Experience, Resource Profitability and
Season. Ethology 119, 1096–1106. https://doi.org/10.1111/eth.12170
Aplin, L.M., Farine, D.R., Mann, R.P., Sheldon, B.C., 2014. Individual-level personality influences
social foraging and collective behaviour in wild birds. Proc. R. Soc. B Biol. Sci. 281, 20141016.
https://doi.org/10.1098/rspb.2014.1016
Bailis, P., Nagpal, R., Werfel, J., 2010. Positional communication and private information in honeybee
foraging models, in: Lecture Notes in Computer Science (Including Subseries Lecture Notes in
Artificial Intelligence and Lecture Notes in Bioinformatics). Springer, Berlin, Heidelberg, pp.
263–274. https://doi.org/10.1007/978-3-642-15461-4_23
Barron, A.B., Maleszka, R., Vander Meer, R.K., Robinson, G.E., 2007. Octopamine modulates honey
bee dance behavior. Proc. Natl. Acad. Sci. U. S. A. 104, 1703–7.
https://doi.org/10.1073/pnas.0610506104
Beekman, M., Lew, J.B., 2007. Foraging in honeybees-when does it pay to dance? Behav. Ecol. 19,
255–261. https://doi.org/10.1093/beheco/arm117
Beekman, M., Makinson, J.C., Couvillon, M.J., Preece, K., Schaerf, T.M., 2015. Honeybee linguistics - a
comparative analysis of the waggle dance among species of Apis. Front. Ecol. Evol. 3, 1–11.
https://doi.org/10.3389/fevo.2015.00011
Bell, A.M., Hankison, S.J., Laskowski, K.L., 2009. The repeatability of behaviour: a meta-analysis.
Anim. Behav. 77, 771–783. https://doi.org/10.1016/j.anbehav.2008.12.022
Beshers, S.N., Fewell, J.H., 2001. Models of division of labor in social insects. Annu. Rev. Entomol. 46,
413–40. https://doi.org/10.1146/annurev.ento.46.1.413
Biesmeijer, J.C., De Vries, H., 2001. Exploration and exploitation of food sources by social insect
colonies: A revision of the scout-recruit concept. Behav. Ecol. Sociobiol.
https://doi.org/10.1007/s002650000289
Boch, R., 1956. Die Tänze der Bienen bei nahen und fernen Trachtquellen. Z. Für Vgl. Physiol. 38,
136–167. https://doi.org/10.1007/BF00338624
Bonabeau, E., Theraulaz, G.U.Y., Deneubourg, J.-L., 1996. Quantitative study of the fixed threshold
model for the regulation of division of labour in insect societies. Proc. R. Soc. B 263, 1565–
1569. https://doi.org/10.1098/rspb.1996.0229
Brooks, M., E., Kristensen, K., Benthem, K., J. ,van, Magnusson, A., Berg, C., W., Nielsen, A., Skaug, H.,
J., Mächler, M., Bolker, B., M., 2017. glmmTMB Balances Speed and Flexibility Among
Packages for Zero-inflated Generalized Linear Mixed Modeling. R J. 9, 378.
https://doi.org/10.32614/RJ-2017-066
Claus O. Wilke, 2018. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2." R
Package Version 093.
Cook, C.N., Lemanski, N.J., Mosqueiro, T., Ozturk, C., Gadau, J., Pinter-Wollman, N., Smith, B.H.,
2020. Individual learning phenotypes drive collective behavior. Proc. Natl. Acad. Sci. 117,
17949–17956. https://doi.org/10/gg86wc
Cribari-Neto, F., Zeileis, A., 2010. Beta Regression in R. J. Stat. Softw. 34, 1–24.
https://doi.org/10.18637/jss.v034.i02

Dall, S.R.X., Bell, A.M., Bolnick, D.I., Ratnieks, F.L., 2012. An evolutionary ecology of individual
differences. Ecol. Lett. 15, 1189–1198. https://doi.org/10.1111/j.1461-
0248.2012.01846.x.An
De Marco, R.J., Farina, W.M., 2001. Changes in food source profitability affect the trophallactic and
dance behavior of forager honeybees (*Apis mellifera* L.). Behav. Ecol. Sociobiol. 50, 441–449.
https://doi.org/10.1007/s002650100382
Dornhaus, A., Klugl, F., Oechslein, C., Puppe, F., Chittka, L., Klu, F., Klügl, F., 2006. Benefits of
recruitment in honey bees: Effects of ecology and colony size in an individual-based model.
Behav. Ecol. 17, 336–344. https://doi.org/10.1093/beheco/arj036
Dyer, J.R.G., Croft, D.P., Morrell, L.J., Krause, J., 2009. Shoal composition determines foraging success
in the guppy. Behav. Ecol. 20, 165–171. https://doi.org/10.1093/beheco/arn129
Garrison, L.K., Kleineidam, C.J., Weidenmüller, A., 2018. Behavioral flexibility promotes collective
consistency in a social insect. Sci. Rep. 8, 1–11. https://doi.org/10.1038/s41598-018-33917-7
George, E., Rajagopal, S., Brockmann, A., 2021. Agent-based model for simulations of individual
variation in recruitment activity in honey bees. Zenodo.
https://doi.org/10.5281/zenodo.5217630
George, E.A., Brockmann, A., 2019. Social modulation of individual differences in dance
communication in honey bees. Behav. Ecol. Sociobiol. 73, 41.
https://doi.org/10.1007/s00265-019-2649-0
George, E.A., Bröger, A., Thamm, M., Brockmann, A., Scheiner, R., 2019. Inter-individual variation in
honey bee dance intensity correlates with expression of the *foraging* gene. Genes Brain
Behav. gbb.12592. https://doi.org/10.1111/gbb.12592
George, E.A., Thulasi, N., Kohl, P.L., Suresh, S., Rutschmann, B., Brockmann, A., 2021. Distance
estimation by Asian honey bees in two visually different landscapes. J. Exp. Biol. 224,
jeb242404. https://doi.org/10.1242/jeb.242404
Gould, J.L., Henerey, M., MacLeod, M.C., 1970. Communication of direction by the honey bee:
Review of previous work leads to experiments limiting olfactory cues to test the dance
language hypothesis. Science 169, 544–554.
Goy, N., Glaser, S.M., Grüter, C., 2021. The adaptive value of tandem communication in ants: Insights
from an agent-based model. J. Theor. Biol. 526, 110762.
https://doi.org/10.1016/j.jtbi.2021.110762
Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V., Giske, J., Goss-Custard, J., Grand, T.,
Heinz, S.K., Huse, G., Huth, A., Jepsen, J.U., Jørgensen, C., Mooij, W.M., Müller, B., Pe'er, G.,
Piou, C., Railsback, S.F., Robbins, A.M., Robbins, M.M., Rossmanith, E., Rüger, N., Strand, E.,
Souissi, S., Stillman, R. a., Vabø, R., Visser, U., DeAngelis, D.L., 2006. A standard protocol for
describing individual-based and agent-based models. Ecol. Model. 198, 115–126.
https://doi.org/10.1016/j.ecolmodel.2006.04.023
Hartig, F., 2018. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models.
Hölldobler, B., Wilson, E.O., 2009. The superorganism. W.W. Norton & Company, Inc.
Honegger, K., de Bivort, B., 2018. Stochasticity, individuality and behavior. Curr. Biol. 28, R8–R12.
https://doi.org/10.1016/j.cub.2017.11.058
Hopper, K.R., 1999. Risk-spreading and bet-hedging in insect population biology. Annu. Rev.
Entomol. 44, 535–560. https://doi.org/10.1146/annurev.ento.44.1.535
I'Anson Price, R., Grüter, C., 2015. Why, when and where did honey bee dance communication
evolve? Front. Ecol. Evol. 3, 1–7. https://doi.org/10.3389/fevo.2015.00125
Jeanson, R., Weidenmüller, A., 2013. Interindividual variability in social insects - proximate causes
and ultimate consequences. Biol. Rev. https://doi.org/10.1111/brv.12074
Ji, Y., Li, X., Ji, T., Tang, J., Qiu, L., Hu, J., Dong, J., Luo, S., Liu, S., Frandsen, P.B., Zhou, Xuguo, Parey,
S.H., Li, L., Niu, Q., Zhou, Xin, 2020. Gene reuse facilitates rapid radiation and independent

767   adaptation to diverse habitats in the Asian honeybee. Sci. Adv. 6, eabd3590.
768   https://doi.org/10/gmhdqd
769 Johnson, B.R., 2010. Division of labor in honeybees: Form, function, and proximate mechanisms.
770   Behav. Ecol. Sociobiol. 64, 305–316. https://doi.org/10.1007/s00265-009-0874-7
771 Jones, J.C., 2004. Honey Bee Nest Thermoregulation: Diversity Promotes Stability. Science 305, 402–
772   404. https://doi.org/10.1126/science.1096340
773 Kohl, P.L., Thulasi, N., Rutschmann, B., George, E.A., Steffan-Dewenter, I., Brockmann, A., 2020.
774   Adaptive evolution of honeybee dance dialects. Proc. R. Soc. B 20200190.
775   https://doi.org/10.1098/rspb.2020.0190
776 Lenth, R., 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means.
777 Liang, Z.S., Nguyen, T., Mattila, H.R., Rodriguez-Zas, S.L., Seeley, T.D., Robinson, G.E., 2012.
778   Molecular determinants of scouting behavior in honey bees. Science 335, 1225–1228.
779   https://doi.org/10.1126/science.1213962
780 Loftus, J.C., Perez, A.A., Sih, A., 2021. Task syndromes: linking personality and task allocation in social
781   animal groups. Behav. Ecol. 32, 1–17. https://doi.org/10.1093/beheco/araa083
782 Mattila, H.R., Seeley, T.D., 2010. Promiscuous honeybee queens generate colonies with a critical
783   minority of waggle-dancing foragers. Behav. Ecol. Sociobiol. 64, 875–889.
784   https://doi.org/10.1007/s00265-010-0904-5
785 McKinney, W., others, 2010. Data structures for statistical computing in python, in: Proceedings of
786   the 9th Python in Science Conference. Austin, TX, pp. 51–56.
787 Michelena, P., Jeanson, R., Deneubourg, J.-L., Sibbald, A.M., 2010. Personality and collective
788   decision-making in foraging herbivores. Proc. R. Soc. B Biol. Sci. 277, 1093–1099.
789   https://doi.org/10.1098/rspb.2009.1926
790 Modlmeier, A.P., Liebmann, J.E., Foitzik, S., 2012. Diverse societies are more productive: a lesson
791   from ants. Proc. R. Soc. B Biol. Sci. 279, 2142–2150. https://doi.org/10.1098/rspb.2011.2376
792 Moore, D., Doherty, P., 2009. Acquisition of a time-memory in forager honey bees. J. Comp. Physiol.
793   A 195, 741–51. https://doi.org/10.1007/s00359-009-0450-7
794 Nakagawa, S., Schielzeth, H., 2010. Repeatability for Gaussian and non-Gaussian data: A practical
795   guide for biologists. Biol. Rev. 85, 935–956. https://doi.org/10.1111/j.1469-
796   185X.2010.00141.x
797 Nieh, J.C., 2010. A negative feedback signal that is triggered by peril curbs honey bee recruitment.
798   Curr. Biol. 20, 310–5. https://doi.org/10.1016/j.cub.2009.12.060
799 Page, R.E., Mitchell, S.D., 1998. Self-organization and the evolution of division of labor. Apidologie
800   29, 171–190. https://doi.org/10.1051/apido:19980110
801 Pinter-Wollman, N., 2021. Proximate and ultimate processes may explain "task syndromes": a
802   comment on Loftus et al. Behav. Ecol. 32, 22–23. https://doi.org/10.1093/beheco/araa126
803 R Core Team, 2018. R: A Language and Environment for Statistical Computing. R Foundation for
804   Statistical Computing, Vienna, Austria.
805 Reynolds, A.M., Swain, J.L., Smith, A.D., Martin, A.P., Osborne, J.L., 2009. Honeybees use a Lévy flight
806   search strategy and odour-mediated anemotaxis to relocate food sources. Behav. Ecol.
807   Sociobiol. 64, 115–123. https://doi.org/10.1007/s00265-009-0826-2
808 RStudio Team, 2016. RStudio: Integrated Development Environment for R. Boston, MA.
809 Schürch, R., Grüter, C., 2014. Dancing bees improve colony foraging success as long-term benefits
810   outweigh short-term costs. PloS One 9, e104660.
811   https://doi.org/10.1371/journal.pone.0104660
812 Seeley, T.D., 1995. The Wisdom of the Hive, Harvard University Press.
813 Seeley, T.D., 1994. Honey bee foragers as sensory units of their colonies. Behav. Ecol. Sociobiol. 34,
814   51–62. https://doi.org/10.1007/BF00175458
815 Seeley, T.D., 1989. Social foraging in honey bees: how nectar foragers assess their colony's
816   nutritional status. Behav. Ecol. Sociobiol. 24, 181–199. https://doi.org/10.1007/BF00292101

817   Seeley, T.D., 1986. Social foraging by honeybees: how colonies allocate foragers among patches of
818   flowers. Behav. Ecol. Sociobiol. 19, 343–354. https://doi.org/10.1007/BF00295707
819   Seeley, T.D., 1983. Division of labor between scouts and recruits in honeybee foraging. Behav. Ecol.
820   Sociobiol. 12, 253–259. https://doi.org/10.1007/BF00290778
821   Seeley, T.D., Camazine, S., Sneyd, J., 1991. Collective decision-making in honey bees: how colonies
822   choose among nectar sources. Behav. Ecol. Sociobiol. 28, 277–290.
823   https://doi.org/10.1007/BF00175101
824   Seeley, T.D., Towne, W.F., 1992. Tactics of dance choice in honey bees: do foragers compare dances?
825   Behav. Ecol. Sociobiol. 30, 59–69. https://doi.org/10.1007/BF00168595
826   Seeley, T.D., Visscher, P.K., 1988. Assessing the Benefits of Cooperation in Honeybee Foraging:
827   Search Costs, Forage Quality, and Competitive Ability. Behav. Ecol. Sociobiol. 22, 229–237.
828   Sih, A., Bell, A., Johnson, J.C., 2004. Behavioral syndromes: An ecological and evolutionary overview.
829   Trends Ecol. Evol. 19, 372–378. https://doi.org/10.1016/j.tree.2004.04.009
830   Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2017. rptR: Repeatability estimation and variance
831   decomposition by generalized linear mixed-effects models. Methods Ecol. Evol.
832   https://doi.org/10.1111/2041-210X.12797
833   Tautz, J., 1996. Honeybee waggle dance: recruitment success depends on the dance floor. J. Exp.
834   Biol. 1381, 1375–1381.
835   Tautz, J., Sandeman, D.C., 2003. Recruitment of honeybees to non-scented food sources. J. Comp.
836   Physiol. A Neuroethol. Sens. Neural. Behav. Physiol. 189, 293–300.
837   https://doi.org/10.1007/s00359-003-0402-6
838   Theraulaz, G., Bonabeau, E., Deneubourg, J.-L., 1998. Response threshold reinforcements and
839   division of labour in insect societies. Proc. R. Soc. B Biol. Sci. 265, 327–332.
840   https://doi.org/10.1098/rspb.1998.0299
841   Thom, C., 2003. The tremble dance of honey bees can be caused by hive-external foraging
842   experience. J. Exp. Biol. 206, 2111–2116. https://doi.org/10.1242/jeb.00398
843   Ulrich, Y., Kawakatsu, M., Tokita, C.K., Saragosti, J., Chandra, V., Tarnita, C.E., Kronauer, D.J.C., 2021.
844   Response thresholds alone cannot explain empirical patterns of division of labor in social
845   insects. PLOS Biol. 19, e3001269. https://doi.org/10.1371/journal.pbio.3001269
846   Van Rossum, G., Drake, F.L., 2009. Python 3. CreateSpace, Scotts Valley, CA.
847   von Frisch, K., 1967. The dance language and orientation of bees. Harvard University Press,
848   Cambridge, Massachusetts.
849   Waddington, K.D., 1982. Honey bee foraging profitability and round dance correlates. J. Comp.
850   Physiol. A 148, 297–301. https://doi.org/10.1007/BF00679014
851   Weidenmüller, A., 2004. The control of nest climate in bumblebee (Bombus terrestris) colonies:
852   interindividual variability and self reinforcement in fanning response. Behav. Ecol. 15, 120–
853   128. https://doi.org/10.1093/beheco/arg101
854   Weidenmüller, A., Chen, R., Meyer, B., 2019. Reconsidering response threshold models—short-term
855   response patterns in thermoregulating bumblebees. Behav. Ecol. Sociobiol. 73, 112.
856   https://doi.org/10.1007/s00265-019-2709-5
857   Westhus, C., Kleineidam, C.J., Roces, F., Weidenmüller, a., 2013. Behavioural plasticity in the fanning
858   response of bumblebee workers: impact of experience and rate of temperature change.
859   Anim. Behav. 85, 27–34. https://doi.org/10.1016/j.anbehav.2012.10.003
860   Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis, 2nd ed, Springer International
861   Publishing. Springer. https://doi.org/10.1007/978-3-319-24277-4
862   Wiernasz, D.C., Hines, J., Parker, D.G., Cole, B.J., 2008. Mating for variety increases foraging activity
863   in the harvester ant, Pogonomyrmex occidentalis: MULTIPLE MATING AND COLONY
864   ACTIVITY. Mol. Ecol. 17, 1137–1144. https://doi.org/10.1111/j.1365-294X.2007.03646.x
865   Wilensky, U., 1999. NetLogo. Center for Connected Learning and Computer-Based Modeling,
866   Northwestern University, Evanston, Illinois, USA.

867   Wilson, E.O., Hölldobler, B., 2005. Eusociality: Origin and Consequences. Proc. Natl. Acad. Sci. 102,
868   13367–13371. https://doi.org/10.1073/pnas.0505858102
869   Wolf, M., Weissing, F.J., 2012. Animal personalities: Consequences for ecology and evolution. Trends
870   Ecol. Evol. 27, 452–461. https://doi.org/10.1016/j.tree.2012.05.001
871