Gene expression in African Americans and Latinos reveals ancestry-specific patterns of genetic architecture

Angel C.Y. Mak[1†*], Linda Kachuri[2†], Donglei Hu[1], Celeste Eng[1], Scott Huntsman[1], Jennifer R. Elhawary[1], Namrata Gupta[3], Stacey Gabriel[3], Shujie Xiao[4], Hongsheng Gui[4], Kevin L. Keys[1,5§], Akinyemi Oni-Orisan[6,7,8], Sam S. Oh[1], Max A. Seibold[9,10], José R. Rodríguez-Santana[11], Michael LeNoir[12], L. Keoki Williams[4,13], Luisa N. Borrell[14], Christopher R. Gignoux[15], Noah A. Zaitlen[16,17], Esteban G. Burchard[1,7‡], Elad Ziv[1,8,18‡]

1. Department of Medicine, University of California San Francisco, San Francisco, CA, USA
2. Department of Epidemiology & Biostatistics, University of California San Francisco, San Francisco, CA, USA
3. Broad Institute of MIT and Harvard, Cambridge, MA, USA
4. Center for Individualized and Genomic Medicine Research, Henry Ford Health System, Detroit, MI, USA
5. Berkeley Institute for Data Science, University of California, Berkeley, CA, USA
6. Department of Clinical Pharmacy, University of California San Francisco, San Francisco, CA, USA
7. Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA, USA
8. Institute for Human Genetics, University of California San Francisco, San Francisco, CA, USA
9. Department of Pediatrics, National Jewish Health, Denver, CO, USA
10. Division of Pulmonary Sciences and Critical Care Medicine, University of Colorado School of Medicine, Aurora, CO, USA
11. Centro de Neumología Pediátrica, San Juan, Puerto Rico
12. Bay Area Pediatrics, Oakland, CA, USA
13. Department of Internal Medicine, Henry Ford Health System, Detroit, MI, USA
14. Department of Epidemiology and Biostatistics, Graduate School of Public Health and Health Policy, City University of New York, New York, NY, USA
15. Colorado Center for Personalized Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA
16. Department of Neurology, University of California Los Angeles, Los Angeles, CA, USA
17. Department of Computational Medicine, University of California Los Angeles, Los Angeles, CA, USA
18. Division of General Internal Medicine, Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA, USA

§ Current address: Department of Bioinformatics and Computational Biology, Ambys Medicines, San Francisco, CA, USA

† These authors contributed equally to this work

‡ These authors jointly supervised this work

Corresponding authors:

Angel C. Y. Mak
Department of Medicine
University of California San Francisco
Box 2911, 1550 4th Street,
San Francisco, CA 94143

Tel: 415-514-9931
Email: ChiYan.Mak@ucsf.edu

Esteban G. Burchard
Department of Medicine
University of California San Francisco
Box 2911, 1550 4th Street,
San Francisco, CA 94143

Tel: 415-514-9677
Email: Esteban.Burchard@ucsf.edu

ABSTRACT

We explored the role of genetic ancestry in shaping the genetic architecture of whole blood gene expression using whole genome and RNA sequencing data from 2,733 African American and Hispanic/Latino children. We find that heritability of gene expression significantly increases with greater proportion of genome-wide African ancestry and decreases with higher levels of Indigenous American ancestry. Fine-mapping of expression quantitative trait loci (eQTLs) in individuals with predominantly African or Indigenous American ancestry revealed ancestry-specific eQTLs in over 30% of heritable genes. We leveraged our data to train genetically derived transcriptome prediction models, which identified significantly more associated genes when applied to 28 traits from a multi-ancestry population. Our findings underscore the importance of increasing representation from ancestrally diverse populations in genomic studies to enable new discoveries and ensure their equitable translation.

INTRODUCTION

Gene expression has been extensively studied as a trait affected by genetic variation in humans[1]. Expression quantitative trait loci (eQTLs) have been identified in most genes[2–4] and extensive analyses across multiple tissues have demonstrated both tissue-specific and shared eQTLs[2]. Genome-wide association studies (GWAS) tend to identify loci that are enriched for eQTLs[5]. Colocalization of eQTLs with GWAS has become an important element of identifying causal genes and investigating the biology underlying genetic susceptibility to disease[6]. More recently, transcriptome-wide association studies (TWAS) have been developed to systematically leverage eQTL data by imputing transcriptomic profiles in external datasets, which has led to the discovery of trait-associated genes that were often missed by GWAS[7,8].

GWAS have identified thousands of loci for hundreds of diseases and disease-related phenotypes in human populations[9]. However, non-European ancestry populations are significantly under-represented in GWAS[10,11] and in studies of gene expression and eQTLs. We and others have shown that gene expression prediction models trained in predominantly European ancestry reference datasets, such as the Genotype-Tissue Expression (GTEx) project[2], have substantially lower accuracy to predict gene expression levels when applied to populations of non-European ancestry[3,12,13]. The importance of having ancestry-matched training datasets for prediction accuracy is also reflected by the limited cross-population portability of other multi-SNP prediction models, such as polygenic risk scores (PRS)[14,15]. Therefore, the limited diversity in genetic association studies and reference datasets is a major obstacle for applying existing integrative genomic studies to non-European populations.

To address this gap, we leveraged whole genome and RNA sequencing data from 2,733 African American and Latino children from the Genes-environments and Admixture in Latino Americans (GALA II) study and the Study of African Americans, Asthma, Genes, and Environments (SAGE) to characterize the genetic architecture of whole blood eQTLs. The diversity within the GALA II/SAGE population enabled us to evaluate how genetic ancestry relates to the heritability of gene expression, and systematically quantify the prevalence of ancestry-specific eQTLs. Lastly, we

93   developed a powerful set of TWAS models to facilitate genetic association analyses in multi-
94   ancestry populations.

RESULTS

Demographic characteristics of GALA II and SAGE participants

We analyzed data from a total of 2,733 participants from the GALA II and SAGE asthma case-
control studies, including 757 self-identified African Americans (AA), 893 Puerto Ricans (PR), 784
Mexican Americans (MX), and 299 other Latinos (LA) who did not self-identify as Mexican
American or Puerto Rican (Table 1, Table S1). Participants identified all four grandparents as
Latino for GALA II or African American for SAGE. The median age of the participants varied from
13.2 (PR) to 16.0 (AA) years old. About 50% of the participants were female and 45% (MX) to 62%
(PR) had physician-diagnosed asthma. For each participant we estimated genome-wide genetic
ancestry (global ancestry) proportions, visualized in Figure 1. Median global African ancestry was
highest in AA (82.6%), followed by PR (19.7%), and lowest in MX (3.5%).

Variability of gene expression explained by common genetic variation increases with
African ancestry

We compared the heritability (h^2) and genetic variance (V_G) of whole blood gene expression
across self-identified race/ethnicity groups (AA, PR, MX) and populations defined based on
genetic ancestry. V_G corresponds to the observed variance in transcriptome levels attributed to
common genetic variation (minor allele frequency [MAF] ≥0.01) within the *cis*-region (see
Methods). Heritability (h^2) was calculated as the ratio of V_G to total phenotypic variance. There
was an overarching trend whereby increasing proportion of African ancestry was positively
associated with h^2 and V_G (Figure 2a-b, Table S2). *Cis*-heritability was significantly higher in AA
($h^2=0.158$) compared to PR ($h^2=0.135$; Wilcoxon rank sum test: $p=2.2\times10^{-50}$) and MX ($h^2=0.123$;
$p=3.3\times10^{-134}$), as well as PR compared to MX ($p=2.2\times10^{-25}$).

In examining genetic ancestry, we considered both global ancestry and local ancestry. Global
ancestry is defined as the genome-wide average ancestry for each individual and local ancestry is

defined as the number of alleles (0, 1 or 2) derived from distinct ancestral populations at a given

locus. The same pattern was observed in the distribution of h^2 and V_G when we assigned

participants to high global African ancestry (AFR_{high}, AFR > 50%) and low global African ancestry

(AFR_{low}, AFR < 10%) groups (Table S1-2, Figure S2c-d). *Cis*-heritability and V_G were significantly

higher in AFR_{high} compared to AFR_{low} groups (Wilcoxon rank sum test: $p=9.6\times10^{-126}$ for h^2; V_G:

$p=7.6\times10^{-106}$ for V_G). The opposite was observed for Indigenous American (IAM) ancestry, with

higher h^2 and V_G in IAM_{low} (<10% IAM ancestry) compared to IAM_{high} (≥50% IAM ancestry) groups

($p=3.1x10^{-103}$ for h2; $p=4.5x10^{-159}$ for V_G).

Next, we refined the observed relationship with heritability by examining ancestry at the locus

level. For each gene, individuals with homozygous local African ancestry (AFR/AFR) were

compared to those with heterozygous local African and European ancestry (AFR/EUR) at the

transcription start site. Consistent with global ancestry, h^2 and V_G were significantly higher in

AFR/AFR homozygotes compared to AFR/EUR and lower in IAM/IAM compared to IAM/EUR

heterozygotes (Figure 2e-h). Within the pooled GALA/SAGE II population, V_G was significantly

higher in AFR/AFR compared to AFR/EUR group (Wilcoxon rank sum test: $p=2.00\times10^{-7}$; Table S3).

Conversely, the V_G in IAM/IAM was significantly lower compared to EUR/EUR ($p=1.76\times10^{-7}$ for

Pooled;) or IAM/EUR ($p=1.62\times10^{-8}$ for Pooled; Table S3). All associations remained statistically

significant in analyses stratified by self-reported race/ethnicity (Table S3; $p=5.70x10^{-5}$ for AFR/AFR

vs. AFR/EUR in AA; $p=2.98x10^{-4}$ for IAM/IAM vs. EUR/EUR in MX; $p=7.67x10^{-11}$ for IAM/IAM vs.

IAM/EUR in MX).

As a parallel approach for evaluating the role of genetic ancestry in the context of whole blood

gene expression, we tabulated the number of heritable genes for which global and/or local

ancestry was significantly associated (FDR<0.05) with transcript levels. The proportion of global

AFR ancestry was associated with the expression of 326 (2.4%) and 589 (4.5%) of heritable genes

in AA and PR, respectively (Table S4). Associations with local, but not global AFR ancestry were

more common (8.9% in AA; 10.9% in PR). Few genes were associated with both global and local

AFR ancestry (1.5% in AA and 2.5% in PR). Among genes associated with both global and local AFR

ancestry in AA, global AFR ancestry explained a median of 1.8% of variation in gene expression,

while local AA accounted for 3.8% of variation (Figure S1). Local IAM ancestry was significantly

associated with the expression of 9.8% of genes in MX, compared to 2.8% for global. The variation

in gene expression accounted for by local IAM ancestry was also higher (3.5%) than global (1.8%)

in genes associated with both global and local IAM ancestry.

Identification of ancestry-specific eQTLs

We next sought to understand patterns of eQTLs in the GALA and SAGE datasets and identify

potential novel population-specific eQTLs. We identified 19,568 genes with at least one eQTL

(eGenes) in the pooled samples and 17,351, 16,980, and 15,975 eGenes in African American,

Puerto Rican, and Mexican American samples, respectively (Table S5, Figure S2).

To characterize ancestry-related differences in the genetic determinants of gene expression, we

built a framework to identify eQTLs with potential ancestry-specific effects on gene expression

levels. We then classified them based on their ancestry-specific properties, including allele

frequency and effect size (Figure 3a). We termed these eQTLs ancestry-specific eQTLs (anc-

eQTLs). To identify anc-eQTLs, we compared the lead eQTL signals of *cis*-heritable genes (Table

S6) between high and low ancestry groups (AFR$_{high}$ vs AFR$_{low}$; IAM$_{high}$ vs IAM$_{low}$). We identified the

lead eQTL signal of each gene in each ancestry group (AFR$_{high}$, AFR$_{low}$, IAM$_{high}$ and IAM$_{low}$) by fine-

mapping 95% credible set of causal variants using CAVIAR[16].

We divided anc-eQTLs into three tiers of ancestry-specificity in multiple steps, first based on

overlap in the 95% credible sets for each gene, followed by analyses accounting for differences in

eQTL effect size, allele frequency, and linkage disequilibrium (LD) between ancestry groups (see

Methods). Over 30% (n=2,961) and 8% (n=679) of heritable genes had at least one AFR$_{high}$-eQTL

and IAM$_{high}$-eQTL, respectively. We define the three tiers of ancestry-associated eQTLs below.

Tier 1 anc-eQTLs had non-overlapping 95% credible sets because they were only common (MAF

≥ 0.01) in high ancestry groups and were thus considered to be the most ancestry specific. Over

28% (n=2,695) of genes contained at least one Tier 1 AFR$_{high}$ lead eQTL, while 7% (n=562) of genes

contained a Tier 1 IAM_{high} eQTL (Figure 3a, left branch; Table S7). Tier 2 anc-eQTLs also had non-

overlapping 95% credible sets for each gene but were commonly segregating (MAF ≥ 0.01) in both

high and low ancestry groups. This class of anc-eQTLs was identified using PESCA[17], which

accounted for population-specific LD patterns. We identified 109 genes (1.1%) with eQTLs having

a posterior probability (PP) > 80% of being specific to AFR_{high} and 33 genes (0.4%) matching the

same criteria for IAM_{high}. Lastly, we found that over 50% of genes (5,058 for AFR and 5,355 for

IAM) contained eQTLs that were shared between high and low ancestry groups. Among these

shared signals, only a small proportion of eQTLs exhibited effect size heterogeneity, as assessed

using Cochran's Q test. We refer to these as Tier 3 anc-eQTLs (2.0% for AFR; 1.0% for IAM) and

note this tier has the lowest level of ancestry-specificity.

Next, we explored whether the ancestry-specific eQTLs were previously associated with

diseases/traits based on the NHGRI-EBI GWAS catalog[18]. We found 70 unique anc-eQTLs in 61

genes associated with 84 diseases/traits reported in 65 publications (Table S8, supplementary file

1). Representative examples of disease-associated anc-eQTLs are shown in (Figure 3b-d). A Tier 1

AFR_{high}-specific eQTL of the *ABAT* gene (rs1641047) was associated with systemic sclerosis (SSc)

in a study that compared European and African American SSc patients[19]. It is common in African

Americans in SAGE (MAF = 0.04) but rare in Mexican Americans in GALA II (MAF = 2×10^{-3}, Figure

3b) and Europeans (MAF = 2×10^{-4}, n=15,666). A Tier 2 IAM_{high}-specific eQTL of *SPSB2* gene

(rs11064437) was previously reported as a risk variant for colorectal cancer in East Asian

populations[20], likely due to the fact that it is present at higher frequencies in this population (East

Asian MAF = 0.26). In our dataset, the lead *SPSB2* eQTL in the IAM_{high} group (rs11064437) was

independent from that of the IAM_{low} group (rs6489739), as indicated by low LD (r^2=0.16, Figure

3d).A Tier 3 AFR_{high}-specific eQTL of *KCNK17* gene (rs34246110) was associated with type 2

diabetes in two independent studies performed in Japanese and multi-ancestry (European,

African American, Hispanic and Asian) populations[21,22]. The detection of this association in

multiple populations was consistent with the characteristics of Tier 3 anc-eQTLs in that they share

lead eQTL signals with other populations but likely exhibit heterogenous effect sizes.

Gene expression prediction models from admixed populations increase power for

gene discovery

We generated gene expression imputation models from GALA II and SAGE following the PrediXcan approach[7]. We used the pooled population (n=2,733) to generate models with significant prediction (see Methods) for 11,830 heritable genes with mean cross-validation (CV) R^2=0.157 (Table S9 and Figure S3). We also generated population-specific models for African Americans (10,090 genes, CV R^2=0.180), Puerto Ricans (9,611 genes, CV R^2=0.163), and Mexican Americans (9,084 genes, CV R^2=0.167). In sensitivity analyses adjusting for local ancestry, we did not observe gains in predictive performance (Table S10). For genes with local ancestry estimates, adjusting for local ancestry did not result in improved model performance (AA: CV R^2=0.177; PR: CV R^2=0.154; MX: CV R^2=0.159).

Our TWAS models were successfully validated in the Study of Asthma Phenotypes and Pharmacogenomic Interactions by Race-Ethnicity (SAPPHIRE)[23], an independent adult population of African Americans (Figure S4). As expected, both pooled and AA-specific TWAS models, which contained the highest number of ancestry-matched individuals, had significantly highest gene expression prediction accuracy in SAPPHIRE compared to other models (correlation R^2=0.026 for AA compared to 0.023 for PR [Wilcoxon rank sum test: p=2.39 x 10^{-5}], 0.020 for MX [p=1.19 x 10^{-17}] and 0.015 for GTEx v8 [p=5.43 x 10^{-85}] models).

To evaluate the potential of TWAS models built from GALA II and SAGE pooled samples (hereafter referred to as GALA/SAGE models) to improve gene discovery in admixed populations, we applied our models to GWAS summary statistics for 28 traits from the multi-ancestry Population Architecture using Genomics and Epidemiology (PAGE) study[24] and conducted parallel analyses using TWAS models based on GTEx v8[2,7] and the Multi-Ethnic Study of Atherosclerosis (MESA)[3]. GTEx v8 whole blood models are based on 670 subjects of predominantly European ancestry (85%) with median age between 50-60 years old[2]. MESA models impute monocyte gene expression[3] based on a sample of African American and Hispanic/Latino individuals (MESA$_{AFHI}$:

n=585). As such, MESA more closely resembles the ancestry composition of our population as
well as subjects included in the PAGE study.

The total number of genes examined was 39% to 82% higher in GALA/SAGE compared to GTEx
(n=7,249) and MESA$_{AFHI}$ (n=5,555), with an intersection of 3,146 genes (Figure 4a, left). Restricting
to genes shared across all three models, CV R^2 was significantly higher in GALA/SAGE compared
to GTEx (p=5.1×10^{-159}) and MESA$_{AFHI}$ (p=9.9×10^{-65}), which is expected based on the substantially
larger sample size of GALA/SAGE (Figure 4a). More importantly, we observed the same trend with
an intersection of 3,003 genes using GALA/SAGE TWAS models built in African Americans only
(n=757), which was a population of comparable sample size (Figure 4b). Association results across
28 PAGE traits demonstrate that TWAS using GALA/SAGE (pooled) models identified a larger
number of significant gene-trait pairs (n=380, FDR<0.05), followed by MESA$_{AFHI}$ (n=303), and GTEx
(n=268), with only 30 genes (35 gene-trait pairs) significant in all three analyses (Figure 4c).

GALA/SAGE models yielded a larger number of associated genes than MESA in 80% of analyses
(binomial test: p=0.012) and 79% compared to GTEx (p=0.019). Of the 330 genes with FDR<0.05
in GALA/SAGE, 143 (43%) were not present in GTEx and 199 (60%) were not present in MESA$_{AFHI}$.
For genes that were significant in at least one TWAS, z-scores in GALA/SAGE were highly
correlated with GTEx (Figure 4c; r=0.74, p=3.5×10^{-64}) and MESA$_{AFHI}$ (Figure 4d; r=0.55, p=
p=8.5×10^{-27}), suggesting that most genes have concordant effects even if they fail to achieve
statistical significance in both analyses (Figure 4c). Despite the higher correlation with GTEx TWAS
z-scores, we observed a higher proportion of gene-trait pairs significant (FDR<0.05) in GALA/SAGE
that were not even nominally associated (p<0.05) in GTEx (33%), compared to 18% in MESA$_{AFHI}$.

The high-density lipoprotein (HDL) cholesterol trait exhibited one of the largest differences in
TWAS associations, with over 60% more significant genes identified using GALA/SAGE models
(n=29) than GTEx predictions (n=11; Figure 4c). Models for several associated genes, including
CETP which has been consistently implicated in HDL cholesterol transport, were not available in
GTEx. The top-ranking HDL cholesterol-associated gene, *CD36* (z-score= -10.52, p=6.9×10^{-26}) had
Tier 1 AFR$_{high}$-specific eQTLs (rs3211938) that were not present at an appreciable frequency in

populations with low African ancestry (MAF in European = 1.3×10^{-4}). The difference in MAF may

explain why *CD36* was not detected using GTEx (z-score=0.057, p=0.95), even though all 43

variants used in GTEx prediction model were available in PAGE summary statistics. In addition to

HDL cholesterol levels, *CD36* expression was also associated with levels of C-reactive protein (z-

score= 5.30, $p=1.1 \times 10^{-7}$).

In addition to multi-ancestry analyses, we also demonstrate the utility of ancestry-specific gene

expression prediction models for the discovery of trait-gene association using TWAS. We

conducted a TWAS of neutrophil counts in African ancestry subjects using population-specific

summary statistics from Chen et al.[25] and ancestry-matched GALA/SAGE models, which identified

139 genes. Eleven of the 139 genes identified were outside of the Duffy locus on 1q23 with

FDR<0.05 (Figure 5b). Novel findings included *CREB5* ($p=1.5 \times 10^{-14}$), *DARS* ($p=2.9 \times 10^{-8}$), *CD36*

($p=1.1 \times 10^{-5}$), *PPT2* ($p=1.3 \times 10^{-5}$), *SSH2* ($p=4.7 \times 10^{-5}$), *TOMM5* ($p=2.9 \times 10^{-4}$), and *ARF6* ($p=3.4 \times 10^{-4}$).

DISCUSSION

Our comprehensive analysis in a uniquely large and diverse multi-racial/ethnic population

revealed new insight into the role of genetic ancestry in shaping the genetic architecture of whole

blood gene expression that may have broad implications for other complex traits. We found that

cis-heritability of gene expression increased with higher proportions of African ancestry. We also

demonstrated that in populations of mixed ancestry, *cis*-heritability was highest with increasing

local African ancestry. The consistency of this pattern in analyses considering global and local

ancestry, as well as in pooled and stratified analyses within populations (African Americans or

Puerto Ricans), suggests this is unlikely to be attributed to confounding by social or environmental

factors. Parallel analyses examining Indigenous American Ancestry revealed an inverse

relationship – with genetic variance and *cis*-heritability of gene expression decreasing in

individuals with higher levels Indigenous American ancestry compared to European ancestry.

These observations are consistent with the overall pattern of heterozygosity of these populations.

African populations consistently show the highest heterozygosity since all other populations

passed through a bottleneck during their migration out of Africa[26,27]. Indigenous American

populations have passed through additional bottlenecks which possibly date to, or prior to, the

settling of the Americas[28,29]. The finding that heritability tracks with heterozygosity across human

populations may be expected from fundamental assumptions that there are more segregating

causal variants in populations with higher heterozygosity[30]. However, to our knowledge, this has

not yet been systematically demonstrated for whole blood gene expression, particularly in a

sufficiently large racially/ethnically diverse sample size.

A second major finding of our work is that over 30% of heritable genes have ancestry-specific

eQTLs and these are predominantly comprised of Tier 1 variants that were rare (MAF < 0.01) or

not polymorphic in another population. Tier 2 ancestry eQTLs are a less striking, albeit interesting

class of variants which were present at a sufficient frequency (MAF>0.01) in both high and low

ancestry groups but did not map to the same eQTL signal. Tier 2 eQTLs could arise due to

differences in environmental effects on gene expression, multiple causal variants at the same

locus that are in different degrees of LD with each other, or *trans*-effects, which are contributed

by genetic variation outside of the *cis*-region. Additional research is needed to elucidate the

underlying reasons for more subtle ancestry-driven differences in eQTL effects.

Given genetic architecture of complex traits is, to variable degree, mirrored or driven by the

genetics of gene expression[31], our work has several important implications. Observations

regarding heritability suggest genetic prediction of complex traits in populations with at least 50%

global African ancestry should be at least as accurate, if not more effective, than in populations

with low proportion of African ancestry. However, for most complex traits the accuracy of

polygenic prediction models in populations of predominantly African ancestry significantly lags

behind other populations[15], particularly those of European ancestry. Although there are caveats

associated with using heritability to gauge the future potential of genetic risk prediction models[32],

our results imply that current deficits in predictive performance in admixed populations are likely

to be a consequence of insufficient sample sizes and systematic underrepresentation. The

substantial prevalence of ancestry-specific eQTLs implies that analytic approaches alone will yield

limited improvements in the generalizability of genetic prediction models, including European-

ancestry derived polygenic risk scores (PRS). For instance, fine-mapping approaches that rely on

accounting for differential LD tagging to identify causal variants will recover some deficits in

predictive performance, but will not compensate for missing genetic predictors. Our results

reinforce the conclusion that developing truly generalizable genetic prediction models requires

capturing the full spectrum of genetic variance in non-European populations. As such, access to

sufficiently large and racially/ethnically and ancestrally diverse populations is the main rate

limiting step.

Previous work by Gay *et al.* assessed the amount of variance in gene expression explained by

global and/or local ancestry[33]. It was reported that local ancestry explains at least 7% of the

variance in residual expression for 1% of expressed genes in an analysis of 117 admixed individuals

from GTEx. Using sample sizes ranging from 757 (AA) to 893 (PR) in GALA II/SAGE, we found that

local ancestry was a significant predictor of transcript levels for at least 10% of heritable genes,

explaining between 2.1% (PR) and 5.1% (AA) of residual variance. Consistent with Gay et al., we

observed that local ancestry explains a larger proportion of variance in gene expression corrected

for global ancestry. However, contrary to their findings, we found that, on average, local ancestry

also explains more variance in residual gene expression than global ancestry.

The importance of comprehensively accounting for genetic determinants of trait variation in

multi-ethnic populations is illustrated in our TWAS results for 28 traits from the PAGE study. TWAS

models trained in the racially/ethnically and ancestrally diverse GALA II/SAGE study identified

significantly more trait-associated genes than GTEx and MESA respectively. This is consistent with

the findings of Geoffroy et al.[13] using GTEx and MESA models, as well as other observations[12,13]

that ancestry-matched models improve power for gene discovery in admixed populations. Over

40% of significantly associated TWAS genes detected using GALA/SAGE models were not available

in GTEx, which underscores how biologically meaningful associations may be overlooked in

studies that exclusively rely on European ancestry-based predictions. The top two HDL

cholesterol-associated genes, *CETP* in 16q13 and *CD36* in 7q21, with established effects on lipid

metabolism[34–37], were not detected in TWAS using GTEx due differences in eQTLs. The finding for

CD36 is compelling since this gene was associated with multiple phenotypes and contains Tier 1

anc-eQTLs that are specific to individuals with high African ancestry, consistent with earlier

findings that evolutionary pressures have elevated genetic divergence at this locus[38,39]. *CD36*

encodes a transmembrane protein that binds many ligands, including collagen, thrombospondin,

and long-chain fatty acids, and serves as a negative regulator of angiogenesis[40]. Beyond lipid

metabolism, its most extensively studied function involves mediating the adherence of

erythrocytes infected with *Plasmodium falciparum*, the parasite responsible for severe

malaria[41,42].

An even more striking example of ancestry-specific genetic architecture involves the Duffy

antigen receptor gene (*ACKR1*) on 1q23, which is responsible for persistently lower white blood

cell and neutrophil counts in African ancestry individuals[43,44]. Common African-derived alleles at

this locus confer a selective advantage against *Plasmodium vivax* malaria and are extremely rare

in European populations. Expression of *ACKR1* could not be imputed using GTEx or MESA, but this

causal gene was captured by the pooled and AA-specific GALA/SAGE TWAS models. Furthermore,

we identified 11 genes associated with neutrophil counts outside of the Duffy locus, including

associations not reported in the source GWAS by Chen *et al*[25]. We replicated the *PSMD3* signal in

17q21[45], which was previously identified in African Americans, as well as *CREB5*, *SSH2*, and *PPT2,*

which were discovered in European ancestry populations[46]. We also identified several novel

genes that have not previously been linked to hematologic traits: *DARS1* in 2q31.1 modulates

reactivity to mosquito antigens[47], while *TOMM5* has been implicated in lipoprotein phospholipase

A2 activity[48].

Taken together, our TWAS findings suggest that models trained in racially/ethnically and

ancestrally diverse populations improve power for identifying trait-associated genes in shared

biological pathways. When applied to admixed populations, GALA II/SAGE imputation models

benefit from having more similar allele frequency profiles to the target datasets, such as PAGE,

as well as more accurate modeling of LD. However, we note that adjustment for local ancestry

did not improve the predictive performance of our TWAS models. Previous analyses of GTEx data

indicated that local ancestry adjustment increases power for discovery in cis-eQTL mapping[33,49],

however this has not been assessed explicitly within the TWAS framework. It is possible that the

slight decrease in CV R^2 occurs due to overadjustment, since local ancestry may serve as a proxy

for information already captured by genetic variants in the *cis*-region.

Interestingly, the correlation of TWAS summary statistics was higher with GTEx than MESA$_{AFHI}$,

which may partly reflect the lack of neutrophils present in monocyte gene expression in MESA$_{AFHI}$

compared to whole blood in GALA/SAGE and GTEx. However, the scarcity of discordant TWAS

findings (null in one analysis or significant associations with opposite directions of effect) or

missing genes between GALA II/SAGE and MESA, indicate that similarity in ancestry may partly

compensate for differences in the cell type analyzed. Furthermore, GALA II/SAGE and GTEx

conducted whole-genome sequencing, whereas MESA TWAS models are based on imputed

genotype data, although we this effect should be limited as we filter to common variants. While

our study population is comprised of pediatric studies, TWAS analyses of medically relevant traits

and chronic conditions in PAGE identified more genes than adult-derived prediction models. This

implies that the power gained from ancestry-matched prediction and large sample size of the

training set may outweigh substantial differences in age between participants in our cohort and

participants in PAGE studies that range across adulthood.

In evaluating the contributions of our work, a few limitations should be acknowledged. Although

our analyses were limited to a single tissue, we note that whole blood is one of the most clinically-

informative and commonly-collected samples, and for over 60% of genes whole blood

transcriptomes can significantly capture expression levels in other tissues[50]. Thus, our

observations regarding the genetic architecture of whole blood eQTLs in non-European ancestry

populations are likely generalizable to other tissues. Our approach for ancestry-specific eQTLs

may result in an underestimation of the number of these loci. We assumed that each gene had

one causal eQTL locus and focused all comparisons on the corresponding 95% credible set. This

assumption is likely violated for genes with multiple independent eQTLs, which would limit our

ability to assess the ancestry-specificity of all signals. In general, we believe this is a conservative

assumption that would lead us to potentially miss some ancestry-specific eQTLs. Detection of Tier

2 anc-eQTLs by the PESCA method relies on having regions that are approximately LD

independent in both populations to estimate the proportion of causal variants. This estimate may

be biased if there is residual LD between regions, which is particularly a challenge in admixed

populations with longer-range LD. Lastly, our comparison of TWAS models may be slightly biased

against GTEx since we did not apply MASHR models, which predict a larger number of genes using

fine-mapped eQTLs[51]. We chose to compare with elastic net GTEx models instead because

GALA/SAGE TWAS models were developed using the same analytic pipeline.

Despite these limitations, our study leveraged a uniquely large and diverse sample of 2,733

African American and Latino youths to advance our understanding of the interplay between

genetic ancestry and genetic determinants of gene expression. Our systematic approach for

evaluating the ancestry-specificity of whole blood eQTLs revealed that most of this signal is due

to Tier 1 anc-eQTLs, which are only polymorphic in predominantly AFR and IAM ancestry

populations. These ancestry-specific eQTLs are present in a large fraction of genes, which

highlights the need for larger genomic studies in globally representative populations to detect

functional genetic variants and evaluate their role in disease susceptibility. We also developed

genetic prediction models of whole blood transcriptomes and demonstrated that they covered

more genes and, in TWAS based on multi-ethnic populations, performed better than similar

models built in European populations. In summary, the results of our study have implications not

only for the genetic prediction of gene expression, but for other complex traits, and ultimately

for the equitable translation of precision medicine efforts to all populations worldwide.

METHODS

Study population

This study examined African American, Puerto Rican and Mexican American children between 8-

21 years of age with or without physician-diagnosed asthma from the Genes-environments and

Admixture in Latino Americans II (GALA II) study and the Study of African Americans, Asthma,

Genes & Environments (SAGE). The inclusion and exclusion criteria are previously described in

detail[52,53]. Briefly, participants were eligible if they were 8-21 years of age and identified all four

grandparents as Latino for GALA II or African American for SAGE. Study exclusion criteria included

the following: 1) any smoking within one year of the recruitment date; 2) 10 or more pack-years

of smoking; 3) pregnancy in the third trimester; 4) history of lung diseases other than asthma (for

cases) or chronic illness (for cases and controls).

The local institutional review board from the University of California San Francisco Human

Research Protection Program approved the studies (IRB# 10-02877 for SAGE and 10-00889 for

GALA II). All subjects and their legal guardians provided written informed consent.

Whole genome sequencing data and processing

Genomic DNA samples extracted from whole blood were sequenced as part of the Trans-Omics

for Precision Medicine (TOPMed) whole genome sequencing (WGS) program[54] and the Centers

for Common Disease Genomes of the Genome Sequencing Program. WGS was performed at the

New York Genome Center and Northwest Genomics Center on a HiSeq X system (Illumina, San

Diego, CA) using a paired-end read length of 150 base pairs (bp), with a minimum of 30x mean

genome coverage. DNA sample handling, quality control, library construction, clustering, and

sequencing, read processing and sequence data quality control are previously described in

detail[54]. All samples were jointly genotyped by the TOPMed Informatics Research Center. Variant

calls were obtained from TOPMed data freeze 8 VCF files generated based on the GRCh38

assembly. Variants with a minimum read depth of 10 (DP10) were used for analysis unless

otherwise stated.

RNA sequencing data generation and processing

Total RNA was isolated from PAXgene tube using MagMax™ for Stabilized Blood Tubes RNA

Isolation Kit (Applied Biosystem, P/N 4452306). Globin depletion was performed using

GLOBINcleasr™ Human (Thermo Fisher Scientific, cat. no. AM1980). RNA integrity and yield were

assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA).

Total RNA was quantified using the Quant-iT™ RiboGreen® RNA Assay Kit and normalized to

5ng/ul. An aliquot of 300ng for each sample was transferred into library preparation which was

an automated variant of the Illumina TruSeq™ Stranded mRNA Sample Preparation Kit. This

method preserves strand orientation of the RNA transcript. It uses oligo dT beads to select mRNA

from the total RNA sample. It is followed by heat fragmentation and cDNA synthesis from the RNA

template. The resultant cDNA then goes through library preparation (end repair, base 'A' addition,

adapter ligation, and enrichment) using Broad-designed indexed adapters substituted in for

multiplexing. After enrichment the libraries were quantified with qPCR using the KAPA Library

Quantification Kit for Illumina Sequencing Platforms and then pooled equimolarly. The entire

process is in 96-well format and all pipetting is done by either Agilent Bravo or Hamilton Starlet.

Pooled libraries were normalized to 2nM and denatured using 0.1 N NaOH prior to sequencing.

Flowcell cluster amplification and sequencing were performed according to the manufacturer's

protocols using the HiSeq 4000. Each run was a 101bp paired-end with an eight-base index

barcode read. Each sample was targeted to 50M reads. Data was analyzed using the Broad Picard

Pipeline which includes de-multiplexing and data aggregation.

RNA-seq reads were further processed using the TOPMed RNA-seq pipeline for Year 3 and Phase

5 RNA-seq data (supplementary file 2 obtained from https://topmed.nhlbi.nih.gov/sites/default/

files/TOPMed_RNAseq_pipeline_COREyr3.pdf) . Count-level data were generated using GRCh38

human reference genome and GENCODE 30 for transcript annotation.

Count-level quality control (QC) and normalization were performed following the Genotype-

Tissue Expression (GTEx) project v8 protocol (https://github.com/hakyimlab/PredictDB_Pipeline_

GTEx_v7). Sample-level QC included removal of RNA samples with RIN < 6, genetically related

samples (equal or more related than third degree relative), and sex-discordant samples based on

reported sex and their *XIST* and *RPS4Y1* gene expression profiles. Count distribution outliers were

detected as follows: (i) Raw counts were normalized using the trimmed mean of M values (TMM)

method in edgeR[55] as described in GTEx v8 protocol. (ii) The log2 transformed normalized counts

at the 25th percentile of every sample were identified (count$_{q25}$). (iii) The 25th percentile (Q25)

of count$_{q25}$ was calculated. (iv) Samples were removed if their count$_{q25}$ was lower than -4 as

defined by visual inspection (Figure S6).

Estimation of hidden confounding factors

To account for hidden confounding factors such as batch effects, technical and biological variation

in the sample preparation, and sequencing and/or data processing procedures, latent factors

were estimated using the Probabilistic Estimation of Expression Residuals (PEER) method[56].

Optimization was performed according to approach adopted by GTEx with the goal to maximized

eQTL discovery[57]. A total of 50 (for AA, PR, MX, pooled samples) and 60 (for AFR$_{high}$, AFR$_{low}$,

IAM$_{high}$, IAM$_{low}$) PEER factors were selected for downstream analyses (Figure S7).

Estimation of global and local genetic ancestry

Genetic principal components (PCs), global and local ancestry, and kinship estimation on genetic

relatedness were computed using biallelic single nucleotide polymorphisms (SNPs) with a PASS

flag from TOPMed freeze 8 DP10 data as described previously[58,59]. Briefly, genotype data from

European (CEU), African (YRI) and Indigenous American (IAM) ancestral populations were used as

the reference panels for global and local ancestry estimation assuming three ancestral

populations. Global ancestry was estimated by ADMIXTURE[60] in supervised while local ancestry

was estimated by RFMIX version 2 with default settings[61]. Throughout this study, local ancestry

of a gene was defined as the number of ancestral alleles (0, 1, or 2) at the transcription start site.

Comparative analyses were performed based on two different sample grouping strategies, by

self-identified race/ethnicity or by global ancestry. Self-identified race/ethnicity included four

groups – African Americans (AA), Puerto Ricans (PR), Mexican Americans (MX), and the pooling

of AA, PR, MX and other Latinos (pooled). For groups defined by global ancestry, samples were

grouped into high (> 50%, AFR_{high} or IAM_{high}) or low (< 10%, AFR_{low} or IAM_{low}) global African or

Indigenous American ancestry (Figure S5). The sample size for each group is shown in Table S1.

Cis-heritability of gene expression

The genetic region of *cis*-gene regulation was defined by 1MB region flanking each side of the

transcription start site (*cis*-region). *Cis*-heritability (h^2) of gene expression was estimated using

unconstrained GREML[62] analysis (--reml-no-constrain), and estimation was restricted to common

autosomal variants (MAF ≥ 0.01). Inverse-normalized gene expression was regressed on PEER

factors, and the residuals were used as the phenotype for GREML analysis. Sex and asthma case-

control status was used as categorical covariates, while age at blood draw and the first 5 genetic

PCs were used as quantitative covariates. *Cis*-heritability was estimated separately for each self-

identified race/ethnicity group (AA, PR, MX and pooled) and groupings based on global (AFR_{high},

AFR_{low}, IAM_{high} and IAM_{low}) and local ancestry (described below). Differences in the distribution

of h^2 and genetic variance (V_G) between groups were tested using two-sided Wilcoxon tests.

These analyses included all genes and were not restricted to *cis*-heritable genes. For analyses

based on local ancestry, we also tested whether the proportion of genes with higher V_G in the

AFR/AFR group was higher than that in AFR/EUR group using a two-sided binomial test. Parallel

analyses were also conducted for Indigenous American ancestry (IAM/IAM vs. EUR/EUR and

IAM/IAM vs. IAM/EUR).

Association of global and local ancestry with gene expression

Methods from Gay *et al* (2020)[33] was modified to identify genes associated with global and local

ancestry (Figure S1). In step 1, inversed normalized gene expression was regressed on age, sex

and asthma status (model 0). In step 2, the residuals from model 0 were regressed on global

ancestry (model 1). In step 3, the residuals from model 1 were regressed on local ancestry (model

2) to identify genes that are associated with local ancestry. False discovery rate of 0.05 (R package

qvalue) was applied to step 2 and 3 separately to identify genes that were significantly associated

with global and/or local ancestry. Step 1 to step 3 were run separately for African and Indigenous

American ancestry. For heritable genes that were associated with global and/or local ancestry, a

joint model of regressing global and local ancestry from residuals from model 0 was also examined

to assess the percentage of variance of gene expression explained by global and/or local ancestry.

Identification of eGenes, cis-eQTLs and ancestry-specific cis-eQTLs

Raw gene counts were processed and eQTLs were identified using FastQTL[63] according to the

GTEx v8 pipeline (https://github.com/broadinstitute/gtex-pipeline). Age, sex, asthma status, first

5 genetic PCs, and PEER factors were used as covariates for FastQTL analysis. *Cis*-eQTLs were

defined as genetic variants that have nominal p values less than the nominal p value threshold

estimated from the permutation step of FastQTL. eGenes are defined as genes with at least one

eQTL.

Ancestry-specific eQTL (anc-eQTL) mapping was performed in participants stratified by high and

low global African and Indigenous American ancestry (see "Grouping samples by self-identified

race/ethnicity or global ancestry"). We developed a framework to identify anc-eQTLs by focusing

on the lead eQTL signal for each gene and compared these between high and low global ancestry

groups (AFR$_{high}$ vs AFR$_{low}$; IAM$_{high}$ vs IAM$_{low}$) (Figure 3a).

Anc-eQTLs were classified into three tiers as described below based on differential effect size and

allele frequency, while accounting for linkage disequilibrium (LD) between eQTLs (Figure 3a). For

every protein coding and heritable (GCTA h^2 LRT $p < 0.05$) eGene (Table S6), the lead eQTL signal

was identified using CAVIAR[16] assuming one causal locus (c=1). The 95% credible set of the high

and low global ancestry group were compared to determine if there was any overlap of the 95%

credible set (overlapping or non-overlapping lead eQTL signals).

For genes with non-overlapping lead eQTL signals (Figure 3a, left branch), eQTLs from the high

group were categorized into Tier 1 anc-eQTLs if they were common variants (MAF >= 0.01) in the

high group but rare or absent in the low group. If the eQTLs were common variants in both the

high and low groups, their specificity was further assessed using PESCA[17] for differential effect

size while accounting for LD between eQTLs. LD pruning on eQTLs with r2 > 0.95 was performed

prior to PESCA analysis. All eQTL pairs with r2 > 0.95 were identified in both the high and low

groups and only those pairs common to both groups were removed. In the PESCA analysis, each

eQTL obtained a posterior probability whether it was specific to the high group (PP_{high}), low group

(PP_{low}) or shared between the two groups (PP_{shared}). All eQTLs in the credible set of the high group

were required to obtain the highest posterior probability in the high group ($PP_{high} > PP_{low}$ and

$PP_{high} > PP_{shared}$) and pass the $PP_{high} > 0.8$ cutoff to be considered Tier 2 ancestry-specific eQTLs.

For genes with overlapping lead eQTL signals (Figure 3a, right branch), Cochran's Q statistics were

used to test for effect size heterogeneity between high and low ancestry groups. An eQTL with

significant heterogeneity in effect size was defined as having a p value of Cochran's Q smaller than

the bonferroni threshold (0.05/nGene, where nGene was the number of genes tested and was

5,058 for AFR and 5,355 for IAM ancestry, see Table S7 row C). Since we assume the 95% credible

set corresponds to a single lead eQTL signal, all eQTLs in the credible set of the high group were

required to have a significant heterogeneous effect size to be qualified into Tier 3 anc-eQTLs.

553 *Development of gene prediction models and transcriptome-wide association*
554 *analyses*

Gene prediction models for *cis*-gene expression were generated using common variants and

elastic net modeling implemented in the PredictDB v7 pipeline (https://github.com/hakyimlab/

PredictDB_Pipeline_GTEx_v7). Models were filtered by nested cross validation (CV) prediction

performance and heritability p value (rho_avg > 0.1, zscore_pval < 0.05 and GCTA h^2 LRT p value

< 0.05). Sensitivity analysis was performed by generating gene prediction models by including

additional number of ancestral alleles as covariates for local ancestry adjustments. For AA,

number of African ancestral allele count was used while European and Indigenous American

ancestral alleles were used as covariates.

Out-of-sample validation of the gene expression prediction models were done using the African

American asthma cohort, Study of Asthma Phenotypes and Pharmacogenomic Interactions by

Race-Ethnicity (SAPPHIRE)[23]. Predicted gene expression from SAPPHIRE genotypes was generated

using the predict function from MetaXcan. Genotypes of SAPPHIRE samples were generated by

whole genome sequencing through the TOPMed program and were processed the same way as

GALA II and SAGE. RNA-seq data from SAPPHIRE were generated as previously described[64] and

were normalized using TMM in edgeR. Predicted and normalized gene expression data were

compared to generate correlation R^2.

To assess the performance of the resulting GALA/SAGE models we conducted transcriptome-wide

association studies (TWAS) of 28 traits using GWAS summary statistics from the Population

Architecture using Genomics and Epidemiology (PAGE) Consortium study by Wojcik et al[24].

Analyses were performed using S-PrediXcan with whole blood gene prediction models from GALA

II and SAGE (GALA/SAGE models), GTEx v8, and monocyte gene expression models from the Multi-

Ethnic Study of Atherosclerosis (MESA) study[3].

Data availability

TOPMed WGS and RNA-seq data from GALA II and SAGE are available on dbGaP under accession

number phs000920.v4.p2 and phs000921.v4.p1, respectively. TOPMed WGS data from SAPPHIRE

are available under the dbGaP accession number phs001467.v1.p1. Models for performing

transcriptome wide association studies developed from GALA II and SAGE are freely available

upon request from the corresponding authors. Due to use restrictions and prohibition of public

posting of genomic summary results from GALA II and SAGE, please email the corresponding

authors to set up data transfer.

ACKNOWLEDGEMENTS

Generation of molecular data for the TOPMed (Trans-Omics in Precision Medicine) program was supported by the National Heart, Lung and Blood Institute (NHLBI). RNA sequencing for "NHLBI TOPMed: Gene-Environment, Admixture and Latino Asthmatics Study" (phs000920, GALA II) and "NHLBI TOPMed: Study of African Americans, Asthma, Genes and Environments" (phs000921, SAGE) was performed at Broad Institute Genomics Platform (HHSN268201600034I). Whole genome sequencing (WGS) for the same studies were performed at the New York Genome Center (NYGC, 3R01HL117004-02S3) and Northwest Genomics Center (NWGC, HHSN268201600032I). Core support including centralized genomic read mapping and genotype calling, along with variant quality metrics and filtering were provided by the TOPMed Informatics Research Center (3R01HL-117626-02S1; contract HHSN268201800002I). Core support including phenotype harmonization, data management, sample-identity QC, and general program coordination were provided by the TOPMed Data Coordinating Center (R01HL-120393; U01HL-120393; contract HHSN268201800001I).

Whole genome sequencing of part of GALA II was performed by the New York Genome Center under The Centers for Common Disease Genomics of the Genome Sequencing Program (GSP) Grant (UM1 HG008901). The GSP Coordinating Center (U24 HG008956) contributed to cross-program scientific initiatives and provided logistical and general study coordination. GSP is funded by the National Human Genome Research Institute, the National Heart, Lung, and Blood Institute, and the National Eye Institute.

This work and EGB were supported in part by the Sandler Family Foundation, the American Asthma Foundation, the RWJF Amos Medical Faculty Development Program, Harry Wm. and Diana V. Hind Distinguished Professor in Pharmaceutical Sciences II, the National Heart, Lung, and Blood Institute (R01HL117004, R01HL135156, X01HL134589, U01HL138626), the National Institute of Health and Environmental Health Sciences (R01ES015794), the National Institute on Minority Health and Health Disparities (R56MD013312, P60MD006902), the Tobacco-Related Disease Research Program (24RT-0025, 27IR-0030), and the National Human Genome Research

Institute (U01HG009080). LK was supported by funding from National Cancer Institute
(K99CA246076). CRG was supported by funding from the National Human Genome Research
Institute (R01HG010297), CRG and NAZ were supported by funding from the National Heart, Lung,
and Blood Institute (R01HL151152) and the National Human Genome Research Institute
(R01HG011345). We gratefully acknowledge the studies and participants who provided biological
samples and data for TOPMed. We thank Soren Germer, Michael C. Zody, Lara Winterkorn and
Catherine Reeves from NYGC and Deborah A. Nickerson from NWGC for overseeing the
production of GALA II and SAGE whole genome sequencing data. We thank Huwenbo Shi, Kathryn
S. Burch and Bodgdan Pasaniuc for their technical support on the PESCA software.

The content is solely the responsibility of the authors and does not necessarily represent the
official views of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

ACYM, LK, KLK, SSO, MAS, CRG, NZ, EGB, EZ contributed to the conception or design of the work.

ACYM, LK, DH, CE, SH, JRE, NG, SG, SX, HG, AOO, JRS, MAL, LKW, LNB, CRG, NZ, EGB, EZ contributed to the acquisition, analysis, or interpretation of data. ACYM, LK, JRE, KLK, AOO, SSO, LNB, CRG, NZ, EGB, EZ have drafted the work or substantively revised it. All authors approved the submission of this manuscript.

REFERENCES

1. Majewski, J. & Pastinen, T. The study of eQTL variations by RNA-seq: from SNPs to phenotypes. *Trends Genet* **27**, 72–79 (2011).

2. The GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318–1330 (2020).

3. Mogil, L. S. *et al.* Genetic architecture of gene expression traits across diverse populations. *PLoS Genet* **14**, e1007586 (2018).

4. Wen, X., Luca, F. & Pique-Regi, R. Cross-population joint analysis of eQTLs: fine mapping and functional annotation. *PLoS Genet* **11**, e1005176 (2015).

5. Kim-Hellmuth, S. *et al.* Cell type–specific genetic regulation of gene expression across human tissues. *Science* **369**, eaaz8528 (2020).

6. Porcu, E. *et al.* Mendelian randomization integrating GWAS and eQTL data reveals genetic determinants of complex and clinical traits. *Nat Commun* **10**, 3300 (2019).

7. Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference transcriptome data. *Nature Genetics* **47**, 1091–1098 (2015).

8. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet* **48**, 245–252 (2016).

9. Tam, V. *et al.* Benefits and limitations of genome-wide association studies. *Nat Rev Genet* **20**, 467–484 (2019).

10. Sirugo, G., Williams, S. M. & Tishkoff, S. A. The Missing Diversity in Human Genetic Studies. *Cell* **177**, 26–31 (2019).

11. Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature News* **538**, 161 (2016).

12. Keys, K. L. *et al.* On the cross-population generalizability of gene expression prediction models. *PLOS Genetics* **16**, e1008927 (2020).

13. Geoffroy, E., Gregga, I. & Wheeler, H. E. Population-Matched Transcriptome Prediction Increases TWAS Discovery and Replication Rate. *iScience* **23**, 101850 (2020).

14. Martin, A. R. *et al.* Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. *The American Journal of Human Genetics* **100**, 635–649 (2017).

15. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health disparities. *Nature Genetics* **51**, 584–591 (2019).

16. Hormozdiari, F., Kostem, E., Kang, E. Y., Pasaniuc, B. & Eskin, E. Identifying Causal Variants at Loci with Multiple Signals of Association. *Genetics* **198**, 497–508 (2014).

17. Shi, H. *et al.* Localizing Components of Shared Transethnic Genetic Architecture of Complex Traits from GWAS Summary Data. *The American Journal of Human Genetics* **106**, 805–817 (2020).

18. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* **47**, D1005–D1012 (2019).

19. Gorlova, O. Y. *et al.* Gene-level association analysis of systemic sclerosis: A comparison of African-Americans and White populations. *PLOS ONE* **13**, e0189498 (2018).

20. Zeng, C. *et al.* Identification of Susceptibility Loci and Genes for Colorectal Cancer Risk. *Gastroenterology* **150**, 1633–1645 (2016).

21. Suzuki, K. *et al.* Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. *Nat Genet* **51**, 379–386 (2019).

22. Vujkovic, M. *et al.* Discovery of 318 new risk loci for type 2 diabetes and related vascular outcomes among 1.4 million participants in a multi-ancestry meta-analysis. *Nature Genetics* **52**, 680–691 (2020).

23. Levin, A. M. *et al.* Nocturnal asthma and the importance of race/ethnicity and genetic ancestry. *American journal of respiratory and critical care medicine* **190**, 266–273 (2014).

24. Wojcik, G. L. *et al.* Genetic analyses of diverse populations improves discovery for complex traits. *Nature* **570**, 514–518 (2019).

25. Chen, M.-H. *et al.* Trans-ethnic and Ancestry-Specific Blood-Cell Genetics in 746,667 Individuals from 5 Global Populations. *Cell* **182**, 1198-1213.e14 (2020).

26. Rosenberg, N. A. *et al.* Genetic Structure of Human Populations. *Science* **298**, 2381–2385 (2002).

27. Henn, B. M., Cavalli-Sforza, L. L. & Feldman, M. W. The great human expansion. *Proc Natl Acad Sci U S A* **109**, 17758–17764 (2012).

28. Reich, D. *et al.* Reconstructing Native American population history. *Nature* **488**, 370–374 (2012).

29. Wall, J. D. *et al.* Genetic variation in Native Americans, inferred from Latino SNP and resequencing data. *Mol Biol Evol* **28**, 2231–2237 (2011).

30. Lin, M., Park, D. S., Zaitlen, N. A., Henn, B. M. & Gignoux, C. R. Admixed Populations Improve Power for Variant Discovery and Portability in Genome-Wide Association Studies. *Front Genet* **12**, 673167 (2021).

31. Cookson, W., Liang, L., Abecasis, G., Moffatt, M. & Lathrop, M. Mapping complex disease traits with global gene expression. *Nat Rev Genet* **10**, 184–194 (2009).

32. Chatterjee, N., Shi, J. & García-Closas, M. Developing and evaluating polygenic risk prediction models for stratified disease prevention. *Nat Rev Genet* **17**, 392–406 (2016).

33. Gay, N. R. *et al.* Impact of admixture and ancestry on eQTL analysis and GWAS colocalization in GTEx. *Genome Biology* **21**, 233 (2020).

34. Barter, P. J. *et al.* Cholesteryl ester transfer protein: a novel target for raising HDL and inhibiting atherosclerosis. *Arterioscler Thromb Vasc Biol* **23**, 160–167 (2003).

35. Armitage, J., Holmes, M. V. & Preiss, D. Cholesteryl Ester Transfer Protein Inhibition for Preventing Cardiovascular Events: JACC Review Topic of the Week. *J Am Coll Cardiol* **73**, 477–487 (2019).

36. Dewey, F. E. *et al.* Distribution and clinical impact of functional variants in 50,726 whole-exome sequences from the DiscovEHR study. *Science* **354**, aaf6814 (2016).

37. Klarin, D. *et al.* Genetics of blood lipids among ~300,000 multi-ethnic participants of the Million Veteran Program. *Nat Genet* **50**, 1514–1523 (2018).

38. Fry, A. E. *et al.* Positive selection of a CD36 nonsense variant in sub-Saharan Africa, but no association with severe malaria phenotypes. *Hum Mol Genet* **18**, 2683–2692 (2009).

39. Bhatia, G. *et al.* Genome-wide comparison of African-ancestry populations from CARe and other cohorts reveals signals of natural selection. *Am J Hum Genet* **89**, 368–381 (2011).

40. Silverstein, R. L. & Febbraio, M. CD36, a scavenger receptor involved in immunity, metabolism, angiogenesis, and behavior. *Sci Signal* **2**, re3 (2009).

41. Oquendo, P., Hundt, E., Lawler, J. & Seed, B. CD36 directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes. *Cell* **58**, 95–101 (1989).

42. Hsieh, F.-L. *et al.* The structural basis for CD36 binding by the malaria parasite. *Nat Commun* **7**, 12837 (2016).

43. Nalls, M. A. *et al.* Admixture mapping of white cell count: genetic locus responsible for lower white blood cell count in the Health ABC and Jackson Heart studies. *Am J Hum Genet* **82**, 81–87 (2008).

44. Reich, D. *et al.* Reduced neutrophil count in people of African descent is due to a regulatory variant in the Duffy antigen receptor for chemokines gene. *PLoS Genet* **5**, e1000360 (2009).

45. Reiner, A. P. *et al.* Genome-Wide Association Study of White Blood Cell Count in 16,388 African Americans: the Continental Origins and Genetic Epidemiology Network (COGENT). *PLOS Genetics* **7**, e1002108 (2011).

46. Astle, W. J. *et al.* The Allelic Landscape of Human Blood Cell Trait Variation and Links to Common Complex Disease. *Cell* **167**, 1415-1429.e19 (2016).

47. Jones, A. V. *et al.* GWAS of self-reported mosquito bite size, itch intensity and attractiveness to mosquitoes implicates immune-related predisposition loci. *Hum Mol Genet* **26**, 1391–1406 (2017).

48. Yeo, A. *et al.* Pharmacogenetic meta-analysis of baseline risk factors, pharmacodynamic, efficacy and tolerability endpoints from two large global cardiovascular outcomes trials for darapladib. *PLoS One* **12**, e0182115 (2017).

49. Zhong, Y., Perera, M. A. & Gamazon, E. R. On Using Local Ancestry to Characterize the Genetic Architecture of Human Traits: Genetic Regulation of Gene Expression in Multiethnic or Admixed Populations. *Am. J. Hum. Genet.* **104**, 1097–1115 (2019).

50. Basu, M., Wang, K., Ruppin, E. & Hannenhalli, S. Predicting tissue-specific gene expression from whole blood transcriptome. *Science Advances* **7**, eabd6991 (2021).

51. Barbeira, A. N. *et al.* Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. *Nat Commun* **9**, 1825 (2018).

52. Oh, S. S. *et al.* Effect of secondhand smoke on asthma control among black and Latino children. *The Journal of allergy and clinical immunology* **129**, 1478–83.e7 (2012).

53. White, M. J. *et al.* Novel genetic risk factors for asthma in African American children: Precision Medicine and the SAGE II Study. *Immunogenetics* **68**, 391–400 (2016).

54. Taliun, D. *et al.* Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature* **590**, 290–299 (2021).

55. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

56. Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of expression residuals (PEER) to obtain increased power and interpretability of gene expression analyses. *Nat Protoc* **7**, 500–507 (2012).

57. Aguet, F. *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).

58. Mak, A. C. Y. *et al.* Lung Function in African American Children with Asthma Is Associated with Novel Regulatory Variants of the KIT Ligand KITLG/SCF and Gene-By-Air-Pollution Interaction. *Genetics* **215**, 869–886 (2020).

59. Lee, E. Y. *et al.* Whole-Genome Sequencing Identifies Novel Functional Loci Associated with Lung Function in Puerto Rican Youth. *Am J Respir Crit Care Med* **202**, 962–972 (2020).

60. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome research* **19**, 1655–1664 (2009).

61. Maples, B. K., Gravel, S., Kenny, E. E. & Bustamante, C. D. RFMix: A Discriminative Modeling Approach for Rapid and Robust Local-Ancestry Inference. *The American Journal of Human Genetics* **93**, 278–288 (2013).

62. Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nature Genetics* **42**, 565–569 (2010).

63. Ongen, H., Buil, A., Brown, A. A., Dermitzakis, E. T. & Delaneau, O. Fast and efficient QTL mapper for thousands of molecular phenotypes. *Bioinformatics* **32**, 1479–1485 (2016).

64. Levin, A. M. *et al.* Integrative approach identifies corticosteroid response variant in diverse populations with asthma. *Journal of Allergy and Clinical Immunology* **143**, 1791–1802 (2019).

Figure 1. Study overview.

Figure 1. Study overview. This study included TOPMed whole genome sequencing and whole transcriptome data generated from whole blood samples of SAGE African American and GALA II Latino individuals (n=2,733). Global ancestry proportions for all individuals are shown in the top panel. **Middle row:** we studied the genetic architecture of gene expression including *cis*-heritability and genetic component of gene expression variation (see Figure 2 for details), and association of gene expression with global and/or local ancestry (see Figure S1 for details). We also identified eQTLs, eGenes and ancestry-specific eQTLs (see Figure 3 for details). **Bottom:** we generated gene prediction models (left panel) and applied them in TWAS based on GWAS summary statistics generated from the PAGE study. We also compared TWAS results generated from GTEx v8 whole blood and MESA monocyte-derived models.

Figure 2. *Cis*-heritability of gene expression and variance explained by genetic component.

Figure 2. *Cis*-heritability (h^2) of gene expression and genetic component of gene expression variance (V_G). **a, b|** Estimated h^2 and V_G of whole blood gene expression in self-identified race/ethnicity groups . **c, d|** Estimated h^2 and V_G in groups defined by genetic ancestry, comparing subjects with > 50% global ancestry (high) to < 10% of the same ancestry (low). **e, f|** Estimated h^2 and V_G in groups defined by local ancestry at gene's transcriptional start site (TSS), comparing subjects with 100% (AFR/AFR) to 50% (AFR/EUR) local African ancestry. **g, h|** Similar to e and f but for 100% (IAM/IAM) to 50% (IAM/EUR) local Indigenous ancestry at the TSS. Two-sided p-values are based on the Wilcoxon rank sum test.

Figure 3.   Identification and examples of ancestry-specific eQTLs.

Figure 3. Identification of ancestry-specific eQTLs (anc-eQTL). **a|** Decision tree for the identification of anc-eQTLs. The lead eQTL signal of each heritable gene, represented by the 95% credible set, was identified by CAVIAR assuming 1 causal variant in each global ancestry high and low group separately (AFR_{high}, AFR_{low}, IAM_{high} and IAM_{low}). Sample size and global ancestry cutoff criteria used to classify individuals into each group are shown in the grey box at the top left corner. In genes with non-overlapping eQTL lead signal (left branch), Tier 1 anc-eQTLs were identified based on differences in allele frequency between groups (MAF $\geq$ 0.01 in high and < 0.01 in low). Tier 2 anc-eQTLs were identified with the consideration of different effect size and LD pattern. To be considered Tier 2 ancestry-specific, all credible eQTLs in the high group were required to show the highest posterior probability to be causal specific to the high group (PP_{high}) and at least one of these eQTLs have to attain PP_{high} > 0.8. In genes with overlapping credible sets (right branch), Tier 3 anc-eQTLs were identified based on different effect size assessed by Cochran's Q. All credible eQTLs in the high group were required to have a Cochran's Q p-value smaller than the bonferroni threshold. Number of genes remaining for each ancestry (AFR and IAM) after each filtering step is indicated alongside each branch and in Table S6. The procedures are described in detail in the method section. **b|** An example of a Tier 1 AFR_{high}-specific eQTL (rs1641047) of the *ABAT* gene that was reported in GWAS catalog to be associated with systemic sclerosis. The minor allele, T allele, was associated with systemic sclerosis and was common (MAF $\geq$ 0.01) in African Americans but rare in Puerto Ricans and Mexican Americans. **c|** An example of Tier 3 AFR_{high}-specific eQTLs (rs34247110 and rs3734618) of the *KCNK17* gene. Both eQTLs were shared between the 95% credible set (CAVIAR) of AFR_{high} and AFR_{low} groups. The eQTL, rs34247110, was reported in GWAS catalog by two different

studies to be associated with type 2 diabetes in European, African American, Hispanic and Asian

populations. The detection of rs34247110 from multiple populations was consistent with the

nature of overlapping lead eQTL signals for Tier 3 anc-eQTLs. Black and open white points and

their whiskers indicate eQTL effect size and their 95% confidence interval (CI) in the AFR_{high} and

AFR_{low} group, respectively. The two eQTLs show non-overlapping 95% CI of effect sizes between

the AFR_{high} and AFR_{low} groups, supporting their identity as Tier 3 anc-eQTLs, which have

significantly different effect size between the two groups (Cochran's Q p < bonferroni

threshold). Diamond and rectangle indicate effect size and their 95% confidence interval from

meta-analysis using METASOFT. **d**| An example of a Tier 2 IAM_{high}-specific, colorectal cancer-

associated eQTL (rs11064437) of the *SPSB2* gene. The top 2 panels on the left show non-

overlapping 95% credible set (CAVIAR) between the IAM_{high} (black outline triangle) and IAM_{low}

(grey outline triangle). The fill color of the triangle indicates LD of an eQTL with the eQTL

marked by the vertical dotted line. The CAVIAR 95% credible eQTLs in IAM_{high} and IAM_{low} were

in different LD blocks, as indicated by low LD represented by the fill of the triangle and LD plots

on the right. LD plots showed eQTLs in the same order as the top two panels on the left. The

bottom three panels on the left show the left eQTL had the highest posterior probability

(PESCA) of being IAM-specific while the right eQTL had highest posterior probability of being

shared between two groups. In the PESCA analysis, eQTLs were pruned by R^2 of 0.95 to remove

eQTLs that have almost perfect LD with each other.

Figure 4. Comparison of transcriptome imputation model performance and transcriptome-wide association study (TWAS) results.

Figure 4. Comparison of transcriptome imputation model performance and transcriptome-wide association study (TWAS) results. **a,b|** Internal cross-validation R2 values for each TWAS model restricting to the subset of overlapping genes. GALA/SAGE models were trained by pooled (a) or AA (b) samples. **c|** Summary of TWAS results across 28 traits from the PAGE GWAS meta-analysis. Venn diagrams depict the number of genes analyzed in each TWAS and the number significant (FDR<0.05) findings across all traits. **d,e|** Correlation in z-scores for the union of genes that were statistically significant in at least one TWAS model. Correlation between GALA/SAGE pooled models with GTEx (d) or MESA (e).

Figure 5.   Transcriptome-wide association study (TWAS) results.

Figure 5. Transcriptome-wide association study (TWAS) results for **a|** HDL cholesterol using summary statistics form the PAGE multi-ancestry GWAS meta-analysis (N=33,063) and **b|** neutrophil counts from a GWAS meta-analysis of African ancestry individuals (N=13,476). Genes that achieved TWAS significance (FDR<0.05) are indicated by circles with a black border and are labeled, except for chromosome 1 for neutrophil counts due to the large number of associations. Significantly associated genes for which expression levels could not be predicted using GTEx v8 elastic net models are indicated in red.

Table 1. Demographic characteristics of GALA II and SAGE participants included in this study.

	African American (N=757)	Puerto Rican (N=893)	Mexican (N=784)	Other Latino (N=299)	Pooled (N=2,733)
Age (years)	16.0 [6.6]	13.2 [4.8]	13.8 [6.5]	13.7 [5.7]	14.0 [6.3]
Sex					
Female	405 (53.5%)	451 (50.5%)	427 (54.5%)	158 (52.8%)	1,441 (52.7%)
Asthma status					
Case	433 (57.2%)	549 (61.5%)	351 (44.8%)	156 (52.2%)	1,489 (54.5%)
Genetic ancestry (%)					
African	82.6 [9.4]	19.7 [13.3]	3.5 [2.7]	8.3 [14.8]	17.5 [61.8]
Indigenous American	0.3 [0.9]	9.9 [3.6]	55.3 [23.2]	42.3 [43.2]	10.7 [45.2]
European	16.5 [9.5]	69.5 [13.6]	40.3 [21.9]	45.9 [20.8]	44.2 [43.8]
Recruitment center					
SF Bay Area	757 (100%)	0 (0%)	348 (44.4%)	109 (36.5%)	1,214 (44.4%)
Chicago	0 (0%)	31 (3.5%)	247 (31.5%)	52 (17.4%)	330 (12.1%)
Puerto Rico	0 (0%)	837 (93.7%)	0 (0%)	8 (2.7%)	845 (30.9%)
New York City	0 (0%)	22 (2.5%)	36 (4.6%)	86 (28.8%)	144 (5.3%)
Houston	0 (0%)	3 (0.3%)	153 (19.5%)	44 (14.7%)	200 (7.3%)

Summary statistics of quantitative variables are reported as median [interquartile range] while

categorial variables were reported as count (percentage).