Title: CRISPR-based kinome-screening revealed MINK1 as a druggable player to rewire 5FU-resistance in OSCC through AKT/MDM2/p53 axis

Running title: MINK1 promotes 5FU resistance via AKT/MDM2/p53 axis

Sibasish Mohanty[#1,2], Pallavi Mohapatra[#1,2], Omprakash Shriwas[1], Shamima Azma Ansari[1,2], Manashi Priyadarshini[1,3], Swatismita Priyadarsini[1], Rachna Rath[4], Mahesh Sultania[5], Saroj Kumar Das Majumdar[6], Rajeeb Kumar Swain[1], Rupesh Dash[1*]

1. Institute of Life Sciences, Bhubaneswar, Odisha, India-751023

2. Regional Center for Biotechnology, Faridabad, India.

3. KIIT School of Biotechnology, KIIT University, Bhubaneswar, India.

4. Dept. of Oral & Maxillofacial Pathology, SCB Dental College & Hospital, Cuttack, Odisha, India- 753007

5. Department of Surgical Oncology, All India Institute of Medical Sciences, Bhubaneswar, Odisha, India-751019

6. Department of Radiotherapy, All India Institute of Medical Sciences, Bhubaneswar, Odisha, India-751019

[#] Authors contributed equally

[*] Corresponding authors

Rupesh Dash, Institute of Life Sciences, Nalco Square, Chandrasekharpur, Bhubaneswar-751023, Odisha, India. Phone: +91-674-2301460, Fax: +91-674-2300728, E-mail: rupesh.dash@gmail.com , rupeshdash@ils.res.in

The authors declare no potential conflicts of interest.

Financial support: Grant support: This work is supported by ICMR (5/13/9/2019-NCD-III),

Institute of Life Sciences, Bhubaneswar intramural support and DBT BT/ INF/22/SP28293/2018

(for imaging facility).

Keywords: MINK1, 5Fluorouracil, AKT, MDM2, p53, Lestaurtinib, Patient derived cells

Abstract

Cisplatin, 5FU and docetaxel (TPF) are the most common chemotherapy regimen used for

advanced OSCC. However, many cancer patients experience relapse, continued tumor growth,

and spread due to drug resistance, which leads to treatment failure and metastatic disease. Here,

using a CRISPR/Cas9 based kinome knockout screening, Misshapen-like kinase 1 (MINK1) is

identified as an important mediator of 5FU resistance in OSCC. Analysis of clinical samples

demonstrated significantly higher MINK1 expression in the tumor tissues of chemotherapy non-

responder as compared to chemotherapy responders. The nude mice and zebrafish xenograft

experiments indicate that knocking out MINK1 restores 5FU mediated cell death in

chemoresistant OSCC. An antibody based phosphorylation array screen revealed MINK1 as a

negative regulator of p53. Mechanistically, MINK1 modulates AKT phosphorylation at Ser473,

which enables p-MDM2 (Ser 166) mediated degradation of p53. We also identified lestaurtinib

as a potent inhibitor of MINK1 kinase activity. The patient derived TPF resistant cell based

xenograft data suggest that lestaurtinib restores 5FU sensitivity and facilitates a significant

reduction of tumor burden. Overall, our study suggests that MINK1 is a major driver of 5FU

resistance in OSCC. The novel combination of MINK1 inhibitor lestaurtinib and 5FU needs

further clinical investigation in advanced OSCC.

Introduction

Majority of head and neck cancer is originated from mucosal epithelium collectively termed as Oral squamous cell carcinomas (OSCC) [1]. It is the most prevalent neoplasm in developing country like India with approximately 80000 new cases diagnosed every year [2]. Unfortunately, most of the patients present with advanced OSCC are without having any preclinical history of pre malignant lesions. The treatment modalities for advanced OSCC includes surgical removal of tumor followed by concomitant chemoradiotherapy. Neoadjuvant chemotherapy is frequently prescribed for surgically unresectable OSCC tumors [3]. However, despite of having all these treatment modalities the 5-year survival rate of advanced tongue OSCC is less than 50%. Chemoresistance is one of the major causes of treatment failure in OSCC[4]. The chemotherapeutic regimen used for OSCC are cisplatin, 5FU and Docetaxel (TPF)[3]. Though chemotherapy drugs show initial positive response, tumor acquires resistance gradually and patients experience continued tumor growth and metastatic disease.

Reprogramming resistant cells to undergo drug induced cell death is a viable way to overcome drug resistance. This can be achieved by identifying the causative factors for acquired chemoresistance and discovering novel agents to target critical causative factors, which will restore drug-induced cell death in chemoresistant OSCC. Kinases, which transfer a reversible phosphate group to proteins, play important role in regulating several phenotypes of carcinogenesis including growth, proliferation, angiogenesis, metastasis and evasion of antitumor immune responses [5]. There are approximately 538 known kinases in human which are known to regulate different kinase signaling. A few of them are also known to regulate drug resistance in HNSCC. A kinome study revealed, microtubule-associated serine/threonine kinase 1 (MAST1) is a major driver of cisplatin resistance in HNSCC. MAST1 inhibitor lestaurtinib efficiently

sensitized chemoresistant cells to cisplatin. Overall, the study suggests that MAST1 is a viable

target to overcome cisplatin resistance [6]. Ectopic overexpression of receptor tyrosine kinase in

HNSCC mediates acquired resistance against cetuximab. For example, hyper activation of AXL

was observed in clinical samples those are resistant to cetuximab [7]. It was also found that RAS-

MAPK are key mediator for cetuximab resistance in OSCC [8]. However, very limited studies are

available about the kinases those mediate 5FU resistance in OSCC.

MINK1 belongs to germinal center kinase (GCK) family and it is involved in regulation of

several important signaling cascades [9]. Recently, it is reported that MINK1 can regulate the

planner cell polarity, which is essential for spreading of cancer cells. The PRICKLE1 encodes

the planner cell polarity protein that binds to MINK1 and RICTOR (a member in mTOR2

complex) and this complex regulates the AKT meditated cell migration. Selectively targeting

either of MINK1, PRICKLE1 or RICTOR can significantly decrease the migration of cancer cell

in breast carcinomas [10]. Ste20-related kinase, misshapen (msn), a Drosophila homolog of

MINK1 regulates embryonic dorsal closure through activation of c-jun amino-terminal kinase

(JNK) [11].

The goal of this study is to find out the potential kinase(s) those are major driver(s) of 5FU

resistance in OSCC, for which a CRISPR based kinome screening was employed on 5FU

resistant OSCC lines. The top ranked protein MINK1 was selected for validation in multiple cell

lines and patient derived cells. In addition to this, lestaurtinib was identified to inhibit MINK1

kinase activity, which can reverse 5FU mediated cell death in chemoresistant OSCC lines.

Ultimately, we demonstrated that MINK1 regulates the p53 in 5FU resistant OSCC. MINK1

activates AKT by phosphorylation at Ser473, which phosphorylates MDM2 at Ser166, the later

in turn triggers degradation of p53.

Materials and Methods

Cell culture: The human tongue OSCC lines (H357, SCC4 and SCC9) were obtained from Sigma Aldrich, sourced from European collection of authenticated cell culture. All OSCC cell lines were cultured and maintained in DMEM F12 supplemented with 10% FBS (Thermo Fisher Scientific), penicillin–streptomycin (Pan Biotech) and 0.5 ug/ml sodium hydrocortisone succinate. HEK 293T cells were maintained in DMEM supplemented with 10% FBS and penicillin–streptomycin (Pan Biotech).

High Content Screening: 1000 cells/ well were seeded in black flat bottom 96 well plate (Thermo Scientific™ Nunc) and divided into two experimental groups, one without 5FU treatment and the other with 5FU treatment. A CRISPR based kinome-wide screening was performed using a lentiviral sgRNA library (LentiArray™ Human Kinase CRISPR Library, Thermo Fisher Scientific, M3775) that knocks out 840 kinase and kinase related genes individually with total number of 3214 sgRNA constructs. Transduction of lentiviruses (MOI:2) containing pooled sg RNAs (up to 4) targeting each of 840 genes along with positive and negative control lentiviruses into individual wells was carried out in presence of polybrene (8μg/ml). At 48 hours post transduction, selection with puromycin (0.5μg/ml) was performed for next 2-3 days, followed by treatment of vehicle control and 5FU at sub lethal dose respectively in both groups for 48h. Finally, cells were stained with LIVE/DEAD™ Viability/Cytotoxicity Kit (Thermo Fisher Scientific Cat # L3224) and high content screening was performed using CellInsight CX7 High-Content Screening (HCS) Platform. The green fluorescence indicates the living cells and red fluorescence indicates dead cells. Images from 20 fields per well were acquired using 10X objective lens. Two different fluorescent channels (excitation wavelengths - 488nm and 561nm) were used for acquiring images. Image analysis was performed using the

HCS Studio software. A threshold value for each channel was set once and used for the entire

screening. To identify the cells, segmentation was done. Some of the clumped and poorly

segmented cells were excluded from further analysis on the basis of area, shape and intensity. On

the basis of intensity, number of live and dead cells were counted and an objective mask (blue

lines in the images) was created around each cell. For positive control, Cas9 over expressing

cells were transduced with lentiviruses expressing sgRNA targeting human hypoxanthine

phosphoribosyltransferase 1 (HPRT1) (LentiArray™ CRISPR Positive Control Lentivirus,

human HPRT, Thermo Fisher Scientific Cat # A32829). HPRT1 knockout cells shows resistance

to 6-thioguanine (6TG) induced cell death. For negative control, Cas9 over expressing cells were

transduced with lentiviruses expressing gRNA with no sequence homology to any region of the

human genome (LentiArray™ CRISPR Negative Control Lentivirus, Thermo Fisher Scientific,

Cat # A32327).

Patient Derived Xenograft: BALB/C-nude mice (6-8 weeks, male, NCr-Foxn1nu athymic)

were purchased from Hylasco Bio-Technology Pvt. Ltd. For xenograft model early passage of

patient-derived cells (PDC2) established from chemo non-responder patient (treated with TPF

without having any response) was considered. Two million cells were suspended in phosphate-

buffered solution-Matrigel (1:1, 100 μl) and transplanted into upper flank of mice. The PDC

MINK1WT cells were injected in right upper flank and PDC MINK1KO cells were injected in

the left upper flank of same mice. These mice were randomly divided into 2 groups (n=5) once

the tumors reached a volume of 50 mm^3 and injected with vehicle control or 5FU (10mg/kg)

intraperitonially twice a week. In another experimental set up, PDC2 WT cells were injected in

right upper flank of mice. These mice were randomly divided into 4 groups (n=5) after the

tumors have reached a volume of 50 mm^3 and injected with vehicle control, 5FU (10mg/kg),

Lestaurtinib (20mg/kg) and 5FU (10mg/kg) and Lestaurtinib (20mg/kg) respectively in each

individual group, intraperitonially twice a week. Tumor size was measured using digital Vernier

caliper twice a week until the completion of experiments. Tumor volume was determined using

the following formula: Tumor volume (mm^3) = (minimum diameter)^2 × (maximum diameter)/2.

Phospho-Protein Profiling: For performing the Phospho Explorer Antibody Array (Full Moon

Biosystems Cat# PEX10), $5X10^6$ number of H3575FUR MINK1WT and H3575FUR

MINK1KO cells were seeded and lysates were isolated, after which the protein samples were

labelled with biotin as described by manufacturer. Biotinylated protein samples were further

blocked in skim milk and were subjected to coupling with the 1318 number of antibodies present

on the array slides. Then for the detection of expression of phospho proteins, Cy3-streptavidin

(Sigma, Cat#S6402) was added. Array slides were scanned at Fullmoon Biosystems array

scanning service and the image analysis was done using ImageJ software.

In vitro **MINK1 kinase activity assays:** MINK1 Kinase Enzyme System (Promega Cat No#

V3911) and ADP-Glo™ Kinase Assay (Promega Cat No#V9101) were procured to perform the

in vitro kinase assay. Selected compounds (10 µM) were incubated at room temperature for 1 hr

with recombinant human MINK1 kinase along with substrate MBP and ATP to perform the

kinase reaction using 1X kinase buffer. Further ADP-Glo™ Reagent was added at room

temperature for 40 mins to stop the kinase reaction and deplete the unconsumed ATP, hence

leaving only ADP. Then Kinase detection reagent was added and incubated at room temperature

for 30 mins to convert ADP to ATP and introduce luciferase and luciferin to detect ATP. Finally,

the luminescence was measured using VICTOR® Nivo™ Multimode Plate Reader (Perkin

Elmer).

Statistical analysis: All data points are presented as mean and standard deviation and Graph Pad

Prism 9.0 was used for calculation. The statistical significance was calculated by one-way

variance (one-way ANOVA), Two-Way ANOVA and considered significance at $P \leq 0.05$.

Study approval: This study was approved by the Institute review Board and Human Ethics

committees (HEC) of Institute of Life Sciences, Bhubaneswar (110/HEC/21) and All India

Institute of Medical Sciences (AIIMS), Bhubaneswar (T/EMF/Surg.Onco/19/03). The animal

related experiments were performed in accordance to the protocol approved by Institutional

Animal Ethics Committee of Institute of Life Sciences, Bhubaneswar (ILS/IAEC-204-AH/DEC-

20). Approved procedures were followed for patient recruitment and after receiving written

informed consent from each patient, tissues samples were collected. Institutional biosafety

committee (IBSC) approved all related experiments.

Results

Establishment and characterization of 5FU resistant OSCC lines: The 5FU resistant OSCC

lines were established by prolonged treatment of 5FU to OSCC cell lines as described in

materials and methods. Monitoring the cell viability of 5FU sensitive (5FUS) and resistant

(5FUR) pattern of H357, SCC4 and SCC9 cell lines by MTT assay suggest that 5FUR cells

achieved acquired resistance (Fig. S1A,B). Enhanced cancer like stem cells (CSCs) and elevated

expression of ATP-binding cassette (ABC) transporters are the hallmarks of chemoresistant cells.

qRT-PCR data suggest that CSC markers (SOX2, OCT4 and NANOG) as well as majority of

ABC transporters expression were elevated in 5FUR cells as compared to 5FUS cells (Fig. S1C,

D).

Kinome wide screening identifies MINK1 as a potential driver of 5FU resistance in OSCC:

To identify the kinases those play important role in 5FU resistance, a CRISPR based kinome-wide screening was performed using a lentiviral sgRNA library knocking out 840 kinases individually with a total number of 3214 sgRNA constructs. To target the individual kinase, upto 4 sgRNA lentiviral constructs were pooled together. For kinome screening, Cas9 overexpressing 5FU resistant OSCC lines were established (Fig. S2), which showed similar drug resistant pattern with parental 5FUR OSCC lines(Fig. S3A). We also determined the polybrene and puromycin tolerance concentration in Cas9 overexpressing clones (Fig. S3B, C). The 5FUS and 5FUR lines were treated with 5FU and cell death was measured in high content analyzer using a fluorescent cell viability dye, the data suggest significantly lower cell death in 5FUR cells as compared to 5FUS cells, which is in harmony with the previously measured 5FU tolerance in both lines (Fig. S4A-C). The screening protocol was optimized using appropriate positive and negative control. When 6TG (6-Thioguanine) was treated to HPRT1 (Hypoxanthine Phosphoribosyltransferase 1) KO lines, which is used as positive control for screening, it showed resistance to cell death, whereas HPRT1 WT cells were sensitive to 6TG, suggesting optimized screening protocol (Fig. S4D, E). As a negative control, lentivirus expressing scrambled sgRNA was used.

For primary screening, the 5FU resistant line (H357 5FUR) was transduced with a lentivirus containing sgRNAs targeting each of the 840 individual kinases, after which sub lethal dose of 5FU was treated for 48h followed by measuring cell death in high content analyzer using a fluorescent cell viability dye (Fig. 1A). From primary screening, 334 kinases out of 840 were selected for further consideration by rejecting rest of sgRNA clones which alone induced cell death more than 30 % (Fig. 1B, C). The 60 candidate kinases having lowest survival fraction

score were evaluated in the secondary screening using three more chemoresistant lines i.e SCC4

5FUR, SCC9 5FUR and H357 CisR. From the primary and secondary screening, MINK1, SBK1

and FKBP1A emerged as the only three common kinases among the 5FUR lines with MINK1

having the lowest survival fraction score, which sensitize the chemoresistant cell to 5FU

mediated cell death the most (Fig. 1 C-E). In secondary screening, MINK1 knock out showed

minimal efficacy in sensitizing cisplatin resistant cell lines to cisplatin (Fig. 1D), indicating the

specific role of MINK1 towards acquired 5FU resistance. Next, monitoring the expression of

MINK1 in OSCC resistant lines, we found the expression of MINK1 is significantly higher in

5FUR lines as compared to 5FUS lines of OSCC (Fig. 1F). With the evaluation of clinical

samples, the expression of MINK1 was found to be elevated in tumor tissues of chemotherapy

non-responders as compared to chemotherapy responders (Fig. G, H). We also evaluated the

MINK1 expression in drug-naive and post-CT nonresponder paired tumor samples from the

same patient and observed that the post–CT-treated tumor samples showed higher MINK1

expression (Fig. I, J).

MINK1 is an important target to overcome 5FU resistance in OSCC: To confirm the

finding from the kinome screening, MINK1KO (knock out) clones were generated, using

lentivirus expressing two different sgRNAs, in Cas9 overexpressing 5FUR OSCC lines and

patient derived line 2 (PDC2) (Fig. S5A). PDC2 was isolated and characterized earlier from

tumor of chemotherapy-non-responder patient, who was treated with neoadjuvant TPF without

any response [22]. The colony forming, MTT and spheroid assay data suggest that knocking out

MINK1 significantly reduced the cell viability when the chemoresistant cells were treated with

5FU (Fig. S5B, C and Fig.2A). Here onwards we used sgRNA 1 for rest of the experiments.

Similarly, knocking out MINK1 induced 5FU mediated cell death in chemoresistant cells (Fig.

2B). Enhanced p-H2AX and cleaved PARP was observed in MINK1KO cells followed by

treatment with 5FU indicating the potential role of MINK1 in mediating 5FU resistance (Fig.

2C,D). Further, to test the in vivo efficacy of the kinome screening data, we implanted PDC2

MINK1WT cells into right upper flank and PDC2 MINK1KO cells into the left upper flank of

nude mice followed by treatment with 5FU. Treatment with 5FU (10 mg/kg) significantly

reduced the tumor burden in the MINK1KO   but not in MINK1WT group (Fig. 2E-G).

Immunohistochemistry data suggests markedly decreased cell proliferation signal (Ki67) in 5FU-

treated MINK1KO tumors (Fig. 2H). Earlier it is known that selective knockdown of MINK1

decreases the migration of human breast cancer lines [10]. To evaluate whether depletion of

MINK1 also reduces migration of chemoresistant OSCC lines and PDC2, Boyden chamber

assays and scratch/wound healing assays were performed. The data suggest that knock out of

MINK1 followed by treatment with 5FU significantly reduces the relative number of migrated

cells (Fig. S6A). Similarly, scratch area analysis suggest that percentage of scratch area is

significantly higher when 5FU is treated to MINK1KO drug resistant cells (Fig. S6B, C). Next,

we used Zebrafish (Danio rerio) [Tg(fli1:EGFP)] tumor xenograft model to further validate our

findings. Equal number of the WT and PDC2MINK1 KO cells were stained with Dil (1,1'-

Dioctadecyl-3,3,3',3'-Tetramethylindocarbocyanine Perchlorate) and injected into perivitelline

space of 48-hour post fertilized zebrafish embryos. After 3 days of injection, embryos were

treated with vehicle control or 5FU (500μM). After 5 days of injection, the primary tumors and

metastatic distribution of cancer cells were documented using a fluorescence microscope. The

tumor growth, as measured by fluorescence intensity of primary tumors, was found to be

significantly reduced in the MINK1 KO group with the treatment of 5FU (Fig. 2I, J). Also,

cancer cells showed reduced distal migration from the primary site in the case of MINK1 KO

5FU treated group (Fig. 2K). These data indicate MINK1 dependency of 5FU resistant OSCC.

Ectopic expression of MINK1 promotes 5FU resistance in OSCC: To confirm the potential

role of MINK1 in 5FU resistance, we performed gain of function study. For this, using a

lentiviral approach we generated MINK1ShRNA stable clones in 5FUR lines and PDC2

(MINK1UTRKD), where the shRNA targets the 3'UTR of MINK1 mRNA. For ectopic

overexpression of MINK1, the MINK1UTRKD cells were transfected with pDESTCMV/TO

MINK1 vector (Fig. 3A). The cell viability and cell death data suggest that knocking down

MINK1 in 5FUR cells result in sensitizing the resistant cells to 5FU, however ectopic

overexpression of MINK1 rescues the 5FU resistant phenotype (Fig. 3B, C). Similarly,

immunostaining data suggest enhanced p-H2AX signal in MINK1UTRKD cells, whereas ectopic

overexpression of MINK1 reduces the p-H2AX signal indicating rescue of 5FU resistance in

OSCC cells (Fig. 3D). We also observed the rescue of cleaved PARP with ectopic expression of

MINK1 suggesting reduced cell death (Fig. 3E). Finally, when MINK1 was overexpressed in

OSCC sensitive lines, cells showed resistance to 5FU induced cell death (Fig. 3F).

MINK1 downregulates the expression of p53 in chemoresistant OSCC through activation

of AKT and MDM2: To understand the specific role of MINK1 in 5FU resistant OSCC, we

performed high-throughput phosphorylation profiling with 1,318 site-specific antibodies from

over 30 signaling pathways in 5FUR cells stably expressing MINK1KO and MINK1WT. From

this study, phosphorylation of p53 at Ser33 and Ser15 were found to be significantly up

regulated in MINK1KO cells as compared to MINK1WT cells. In addition to this,

phosphorylation of AKT at Ser473 and phosphorylation of MDM2 at Ser166 were found to be

down regulated in MINK1KO cells as compared to MINK1WT (Fig. 4A). Further,

immunoblotting was performed to validate the finding of phosphorylation profiling antibody

array. The data suggest that phosphorylation of p53 at Ser15 and Ser33 is significantly

upregulated and phosphorylation of MDM2 at Ser166 is profoundly downregulated in

MINK1KO cells as compared to MINK1WT chemoresistant cells (Fig. 4B). OSCC lines H357

and SCC4 have mutant p53, whereas MCF7 and HEK 293 have wild type p53 expression.

Further, p53 expression was also found to be inversely correlated with MINK1 irrespective of its

mutation status. (Fig. 4B, S7). Next, when MINK1 was ectopically overexpressed in MINK1KD

(shRNA targeting 3'UTR) clones, downregulation of p53, p-p53 (Ser33) and p-p53 (Ser15) were

observed in chemoresistant OSCC lines (Fig. 4C). In harmony to our finding of phosphorylation

array, p-AKT(Ser473) was found to be down regulated in MINK1 depleted cells, which was

rescued with ectopic overexpression of MINK1 (Fig. 4 D, E). To confirm the potential role of

AKT in modulating MINK1 mediated p53 regulation, we ectopically overexpressed

constitutively active AKT (myrAKT) in MINK1KO cells. The immunoblotting data suggest that

expression of p53, p-p53 (Ser33) and p-p53 (Ser15) were downregulated when MyrAKT was

overexpressed in MINK1KO cells (Fig. 4F). Similarly, when MINK1 over expressing cells were

treated with AKT inhibitor (Akti-1/2) , the p53 expression was rescued along with

downregulation of p-MDM2 (Ser166) (Fig. 4G). p53 target genes were also evaluated in MINK1

KO cells and the immunoblotting data suggest that expression of p21, NOXA and TIGAR in

MINK1KO clones were upregulated as compared to MINK1WT clones (Fig. 4H).

Evaluation of Lestaurtinib as MINK1 inhibitor to reverse 5FU resistance in OSCC: From

the screening data, we observed that MINK1 expression is elevated in chemoresistant OSCC and

genetic inhibition of the same sensitizes drug resistant lines to 5FU induced cell death. Hence,

MINK1 can be a potential therapeutic target to overcome chemoresistance in OSCC. Very

limited information on the inhibitors of MINK1 is available in the literature. Hence, we looked

for the potential MINK1 inhibitors in the international union of basic and clinical pharmacology

(IUPHAR) database, where a screen of 72 inhibitors against 456 human kinases binding activity

is provided. Among the potential twelve MINK1 inhibitors, we tested the MINK1 inhibitory

activity of three inhibitors i.e., staurosporine, pexmetinib and lestaurtinib. The kinase assay data

suggest that lestaurtinib and pexametinib have highest inhibitory activity for MINK1 (Fig. 5A).

The 50% MINK1 inhibitory activity was observed at concertation of 100 nM in case of

lestaurtinib and 10 µM for pexmetinib (Fig. 5B). Next, cell viability assay was performed to

select a dose of lestaurtinib and pexmetinib that does not affect cell viability when treated alone

(viability > 80%) in 5FU resistant OSCC lines (Fig. 5C, D). Further, the cell viability, spheroid

assay and cell death data suggest that the selected sub lethal dose of lestaurtinib (50nM) and

pexmetinib (500µM) can efficiently restore 5FU mediated cell death in chemoresistant OSCC

lines and PDC2 (Fig. 5 E, F and S8A). The IC50 value of 5FU in H3575FUR is 20.49µM,

however combination of lestaurtinib (50nM) decreases the IC50 value to 4.82 µM and

combination of pexmetinib (500µM) lowers the IC50 value of 5FU to 7.08 µM (Fig. 5 E). As

lestaurtinib, with a much lower concentration (50nM) as compared to pexmetinib (500nM)

sensitize 5FU to chemoresistant cells, from here on lestaurtinib was considered for rest of the

study. Enhanced expression of p-H2AX and cleaved PARP was observed only in combination

group with lestaurtinib and 5FU indicating programmed cell death (Fig. 5G and S8B). Boyden

chamber assays data suggests that combinatorial treatment of lestaurtinib and 5FU

significantly reduces the relative number of migrated cells (Fig S8C). In harmony to the

observation made by knockout of MINK1 in chemoresistant cells, we also found that lestaurtinib

significantly decreased the phosphorylation of MDM2(Ser166) and AKT(Ser473) and elevated

the expression of p53 in chemoresistant cells (Fig. 5H). Further, we found that lestaurtinib failed

to sensitize 5FU mediated cell death in MINK1 knocked out 5FUR lines (Fig. 5I), which

suggests that lestaurtinib conferred 5FU sensitivity by inhibiting MINK1 kinase activity. To

check the in vivo efficacy of this novel combination, nude mice xenograft model was performed

using patient derived cells (PDC2). The in vivo data suggest that the combination of lestaurtinib

(20mg/kg) and 5FU (10mg/kg) profoundly reduced the tumor burden as compared to treatment

with either of the single agents (Fig. 6A-C). Immunohistochemistry data suggest significant

reduction in CD44 and Ki67 expression along with increased expression of cleaved caspase 3 in

combination group (Fig. 6D). Finally, we performed combinatorial anti-tumor effect of non-

cytotoxic extremely low dose of cisplatin (1μM), 5FU (1μM) and lestaurtinib (50nM) in PDC2.

The cell viability, cell death, western blotting and colony forming assay data suggest

significantly higher cell death in cisplatin, 5FU and lestaurtinib combinatorial group, as

compared to any other possible combinatorial group, i.e. 5FU and lestaurtinib or cisplatin and

lestaurtinib or cisplatin and 5FU (Fig. S8).

Discussion: The hallmark chemoresistant phenotypes of cancer cells are reduced apoptosis,

altered metabolic activity, enhanced cancer stem cells like population, increased drug efflux and

decreased drug accumulations. However, the causative factors which are responsible for acquired

chemoresistance is largely not known. It is well known that kinases play key role in various

processes of carcinogenesis and kinase inhibitors are established as potential anti-tumor agents.

In this study, for the first time, we have performed a kinome screening in drug resistant cancer

cells to explore the potential kinase(s) those mediate 5FU resistance in OSCC. From the primary

and secondary kinome screening, MINK1 was found to be top ranked kinase that can re-sensitize

drug resistant cells to 5FU. Overall, MINK1 is known to regulate cell senescence, cell motility

and migration. Till date the potential role of MINK1 in modulating chemoresistanace is still

unknown. Here in this study, we found the novel function of MINK1 by which it regulates 5FU

resistance in OSCC.

To understand the mechanism by which MINK1 regulates 5FU resistance, we performed a high-

throughput phosphorylation profiling in 5FUR cells stably expressing MINK1sgRNA. From this

study, we found p- p53 (Ser33) and p-53(Ser15) to be significantly up-regulated in MINK1KO

cells and p-AKT (Ser473) and p-MDM2 (Ser166) were found to be down-regulated in

MINK1KO cells as compared to MINK1WT. The tumor suppressor p53 is phosphorylated at

various amino acids by different kinases, which tightly regulates its stability [23]. It is well known

that MDM2 (a E3 ubiquitin ligase) acts as a negative regulator of p53. MDM2 forms a complex

with p53 and facilitates the recruitment of ubiquitin molecules for its degradation [24]. Earlier, it

was established that insulin induced activated AKT (Ser473) phosphorylates MDM2 at Ser 166

and Ser 186, which can lead to MDM2 mediated proteasomal degradation of p53 in cytoplasm

as well as in nucleus [25, 26]. These events lead to blocking of p53-mediated transcription of genes

those generally involve in apoptosis, cell cycle regulation and senescence. In addition to this,

p53 is phosphorylated at Ser15 by ATM, DNA-PK and ATR in response to DNA damage [27, 28,

29]. Hence, phosphorylation of Ser15 and Ser33 leads to activation and stabilization of p53 as they

attenuate the MDM2 mediated degradation of p53 [30, 31]. Overall, in this study we found that

MINK1 regulates the expression of p53 through activation of AKT which in turns triggers p-

MDM2 (Ser 166) (Figure 6 E).

Jin et al 2018 performed a kinome screening in cisplatin resistant cells to explore the potential

kinases those confer cisplatin resistance in HNSCC. The data suggests that microtubule-

associated serine/threonine-protein kinase 1 (MAST1) mediates cisplatin resistance in HNSCC

by phosphorylating MEK1, triggering cRaf-independent activation of MEK1, which led to down regulation of BH3 only protein BIM. Jin et al 2018 also found that lestaurtinib to be a potent inhibitor of MAST1. Lestaurtinib successfully restores the cisplatin induced cell death in cisplatin resistant cells [6]. Lestaurtinib not only inhibits MAST1 activity but also known as an inhibitor of JAK2, Trk and FLT3 [32, 33]. In this study, we found that lestaurtinib inhibits activity of MINK1 and lestaurtinib can resensitize the drug resistant OSCC to 5FU. The most common chemotherapy regimen for OSCC is the combination of cisplatin, 5FU and Docetaxel (TPF). Finally, our data suggests that combination of extremely low dose of cisplatin (1μM), 5FU (1μM) and lestaurtinib (50nM) can overcome chemoresistance in OSCC (Fig. S8). Currently lestaurtinib alone or in combination with other chemotherapy drug is under clinical investigation (phase II) for patients having AML and it is well tolerated in human beings [34].

Overall, our data suggests that MINK1 is a mediator of 5FU resistance in OSCC. Besides this, though we have demonstrated that MINK1 negatively regulates P53 through AKT/MDM2 axis in 5FU resistant OSCC, the 5FU specificity of this MINK1 driven signalling cascade remains to be fully elucidated. Further, genetic or pharmacological (Lestaurtinib) inhibition of MINK1 successfully resensitized chemo resistant lines to 5FU. This novel combination of 5FU and Lestaurtinib needs further clinical investigation.

Acknowledgements: PM is a CSIR-SRF, SM is UGC-SRF, SAA is a UGC-JRF.

Author contribution: SM, PM, OS, MP, SAA and SP performed experiments, and analyzed the data, under the direction of R.D. RR, MS and S.K.M. performed part of experiments. SM, PM, RKS and R.D. designed experiments and supervised the study. R.D wrote the manuscript. All authors approved the final version.

Competing interests: The authors declare no conflict of interest.

References

1.   Johnson DE, Burtness B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head and

neck squamous cell carcinoma. *Nat Rev Dis Primers* 2020, **6**(1)**:** 92.

2.   Chaturvedi A, Husain N, Misra S, Kumar V, Gupta S, Akhtar N*, et al.* Validation of the

Brandwein Gensler Risk Model in Patients of Oral Cavity Squamous Cell Carcinoma in

North India. *Head Neck Pathol* 2020, **14**(3)**:** 616-622.

3.   Vishak S, Rangarajan B, Kekatpure VD. Neoadjuvant chemotherapy in oral cancers:

Selecting the right patients. *Indian journal of medical and paediatric oncology : official*

journal of Indian Society of Medical & Paediatric Oncology 2015, **36**(3)**:** 148-153.

4.   Maji S, Panda S, Samal SK, Shriwas O, Rath R, Pellecchia M*, et al.* Bcl-2 Antiapoptotic

Family Proteins and Chemoresistance in Cancer. *Advances in cancer research* 2018, **137:**

37-75.

5.   Gross S, Rahal R, Stransky N, Lengauer C, Hoeflich KP. Targeting cancer with kinase

inhibitors. *J Clin Invest* 2015, **125**(5)**:** 1780-1789.

6. Jin L, Chun J, Pan C, Li D, Lin R, Alesi GN*, et al.* MAST1 Drives Cisplatin Resistance in Human Cancers by Rewiring cRaf-Independent MEK Activation. *Cancer cell* 2018, **34**(2)**:** 315-330 e317.

7. Brand TM, Iida M, Stein AP, Corrigan KL, Braverman CM, Luthar N*, et al.* AXL mediates resistance to cetuximab therapy. *Cancer research* 2014, **74**(18)**:** 5152-5164.

8. Rampias T, Giagini A, Siolos S, Matsuzaki H, Sasaki C, Scorilas A*, et al.* RAS/PI3K crosstalk and cetuximab resistance in head and neck squamous cell carcinoma. *Clin Cancer Res* 2014, **20**(11)**:** 2933-2946.

9. Dan I, Watanabe NM, Kobayashi T, Yamashita-Suzuki K, Fukagaya Y, Kajikawa E*, et al.* Molecular cloning of MINK, a novel member of mammalian GCK family kinases, which is up-regulated during postnatal mouse cerebral development. *FEBS Lett* 2000, **469**(1)**:** 19-23.

10. Daulat AM, Bertucci F, Audebert S, Serge A, Finetti P, Josselin E*, et al.* PRICKLE1 Contributes to Cancer Cell Dissemination through Its Interaction with mTORC2. *Developmental cell* 2016, **37**(4)**:** 311-325.

11. Su YC, Treisman JE, Skolnik EY. The Drosophila Ste20-related kinase misshapen is
required for embryonic dorsal closure and acts through a JNK MAPK module on an
evolutionarily conserved signaling pathway. *Genes Dev* 1998, **12**(15)**:** 2371-2380.

12. Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for
CRISPR screening. *Nat Methods* 2014, **11**(8)**:** 783-784.

13. Tzelepis K, Koike-Yusa H, De Braekeleer E, Li Y, Metzakopian E, Dovey OM*, et al.* A
CRISPR Dropout Screen Identifies Genetic Vulnerabilities and Therapeutic Targets in
Acute Myeloid Leukemia. *Cell Rep* 2016, **17**(4)**:** 1193-1205.

14. Shriwas O, Arya R, Mohanty S, Mohapatra P, Kumar S, Rath R*, et al.* RRBP1 rewires
cisplatin resistance in oral squamous cell carcinoma by regulating Hippo pathway. *Br J*
Cancer 2021, **124**(12)**:** 2004-2016.

15. Moffat J, Grueneberg DA, Yang X, Kim SY, Kloepfer AM, Hinkle G*, et al.* A lentiviral
RNAi library for human and mouse genes applied to an arrayed viral high-content screen.
Cell 2006, **124**(6)**:** 1283-1298.

16. Samal SK, Routray S, Veeramachaneni GK, Dash R, Botlagunta M. Ketorolac salt is a
newly discovered DDX3 inhibitor to treat oral cancer. *Scientific reports* 2015, **5:** 9982.

17. Shriwas O, Priyadarshini M, Samal SK, Rath R, Panda S, Das Majumdar SK*, et al.*
DDX3 modulates cisplatin resistance in OSCC through ALKBH5-mediated m(6)A-
demethylation of FOXM1 and NANOG. *Apoptosis* 2020, **25**(3-4)**:** 233-246.

18. Mohapatra P, Shriwas O, Mohanty S, Ghosh A, Smita S, Kaushik SR*, et al.* CMTM6
drives cisplatin resistance by regulating Wnt signaling through the ENO-
1/AKT/GSK3beta axis. *JCI Insight* 2021, **6**(4).

19. Johannessen CM, Boehm JS, Kim SY, Thomas SR, Wardwell L, Johnson LA*, et al.* COT
drives resistance to RAF inhibition through MAP kinase pathway reactivation. *Nature*
2010, **468**(7326)**:** 968-972.

20. Campeau E, Ruhl VE, Rodier F, Smith CL, Rahmberg BL, Fuss JO*, et al.* A versatile
viral system for expression and depletion of proteins in mammalian cells. *PloS one* 2009,
4(8)**:** e6529.

21. Ramaswamy S, Nakamura N, Vazquez F, Batt DB, Perera S, Roberts TM*, et al.*
Regulation of G1 progression by the PTEN tumor suppressor protein is linked to
inhibition of the phosphatidylinositol 3-kinase/Akt pathway. *Proc Natl Acad Sci U S A*
1999, **96**(5)**:** 2110-2115.

22.   Maji S, Shriwas O, Samal SK, Priyadarshini M, Rath R, Panda S, *et al.* STAT3- and

GSK3beta-mediated Mcl-1 regulation modulates TPF resistance in oral squamous cell

carcinoma. *Carcinogenesis* 2019, **40**(1)**:** 173-183.

23.   Maclaine NJ, Hupp TR. The regulation of p53 by phosphorylation: a model for how

distinct signals integrate into the p53 pathway. *Aging (Albany NY)* 2009, **1**(5)**:** 490-502.

24.   Moll UM, Petrenko O. The MDM2-p53 interaction. *Mol Cancer Res* 2003, **1**(14)**:** 1001-

1008.

25.   Gottlieb TM, Leal JF, Seger R, Taya Y, Oren M. Cross-talk between Akt, p53 and

Mdm2: possible implications for the regulation of apoptosis. *Oncogene* 2002, **21**(8)**:**

1299-1303.

26.   Mayo LD, Donner DB. A phosphatidylinositol 3-kinase/Akt pathway promotes

translocation of Mdm2 from the cytoplasm to the nucleus. *Proc Natl Acad Sci U S A*

2001, **98**(20)**:** 11598-11603.

27.   Khanna KK, Keating KE, Kozlov S, Scott S, Gatei M, Hobson K*, et al.* ATM associates

with and phosphorylates p53: mapping the region of interaction. *Nat Genet* 1998, **20**(4)**:**

398-400.

28. Lakin ND, Hann BC, Jackson SP. The ataxia-telangiectasia related protein ATR mediates

DNA-dependent phosphorylation of p53. *Oncogene* 1999, **18**(27)**:** 3989-3995.

29. Shieh SY, Ikeda M, Taya Y, Prives C. DNA damage-induced phosphorylation of p53

alleviates inhibition by MDM2. *Cell* 1997, **91**(3)**:** 325-334.

30. Sakaguchi K, Herrera JE, Saito S, Miki T, Bustin M, Vassilev A*, et al.* DNA damage

activates p53 through a phosphorylation-acetylation cascade. *Genes Dev* 1998, **12**(18)**:**

2831-2841.

31. Yogosawa S, Yoshida K. Tumor suppressive role for kinases phosphorylating p53 in

DNA damage-induced apoptosis. *Cancer Sci* 2018, **109**(11)**:** 3376-3382.

32. Hexner EO, Serdikoff C, Jan M, Swider CR, Robinson C, Yang S*, et al.* Lestaurtinib

(CEP701) is a JAK2 inhibitor that suppresses JAK2/STAT5 signaling and the

proliferation of primary erythroid cells from patients with myeloproliferative disorders.

Blood 2008, **111**(12)**:** 5663-5671.

33. Shabbir M, Stuart R. Lestaurtinib, a multitargeted tyrosine kinase inhibitor: from bench

to bedside. *Expert Opin Investig Drugs* 2010, **19**(3)**:** 427-436.

34.   Knapper S, Burnett AK, Littlewood T, Kell WJ, Agrawal S, Chopra R, *et al.* A phase 2

515   trial of the FLT3 inhibitor lestaurtinib (CEP701) as first-line treatment for older patients

516   with acute myeloid leukemia not considered fit for intensive chemotherapy. *Blood* 2006,

517   **108**(10)**:** 3262-3270.

Figure Legends

Figure 1: CRISPR based Kinome screening revealed MINK1 as a potential mediator for

5FU resistance in OSCC. A) Schematic presentation of approach for CRSPR/Cas9 based

kinome knockout screening to discover the potential kinase responsible for 5FU resistance in

OSCC. **B-C)** Primary screening of 840 kinases was performed with sublethal dose of 5FU

(8μM). The kinases (n=506 nos) whose knockout alone induced significantly higher cell death (>

30%) depicted in red were excluded. From the rest of the kinases (n=334 nos) depicted in green,

the survival fraction (5FU treated/ Vehicle Control) was determined and top 60 candidates

having lowest survival fraction were considered for secondary screening. **D)** For secondary

screening with top 60 kinases, four cell lines were considered i.e., H357 5FUR, SCC4 5FUR,

SCC9 5FUR and H357CisR. After overlapping all three 5FUR cell lines, MINK1, SBK1,

FKBP1A were found to be the common kinases among them. MINK1 was selected as a potential

kinase target purely based on having the lowest survival fraction among all common candidates.

E) The fluorescent images acquired from high content analyzer with indicated treated group

during kinome screening. **F)** Lysates were collected from indicated cells and immunoblotting

was performed with indicated antibodies. **G)** Protein expression of MINK1 was analyzed by IHC

in chemotherapy- responder and chemotherapy-non-responder OSCC tumors. Scale bars: 50 μm.

H) IHC scoring for MINK1 from panel G (Q Score =Staining Intensity $\times$ % of Staining),

(Median, n=11 for chemotherapy-responder and n=23 for chemotherapy-non-responder) *$P <$

0.05 by 2-tailed Student's t test. **I)** Protein expression of MINK1 was analyzed by

immunohistochemistry (IHC) in pre- and post-TPF treated paired tumor samples from

chemotherapy-non-responder patients. Scale bars: 50 μm. **J)** IHC scoring for MINK1 from panel

I (Q Score =Staining Intensity × % of IHC Staining). *$P < 0.05$ by 2-tailed Student's t test.

Figure 2: Selectively targeting MINK1 restores 5FU induced cell death in chemoresistant

OSCC: A) 5FU resistant cells stably expressing MINK1sgRNA (#1 and #2) and NTsgRNA

were treated with 5FU for 48h and cell viability was determined by MTT assay (n=3 and 2-way

ANOVA). **B)** Indicated cells were treated with 5FU for 48h, after which cell death was

determined by annexin V/7AAD assay using flow cytometer. Bar diagrams indicate the

percentage of cell death (early and late apoptotic) with respective treated groups (Mean ±SEM,

n=3, Two-way ANOVA). **C)** Indicated cells were treated with 5 μM of 5FU for 48h, after which

immunostaining was performed for γ-H2AX. **D)** Indicated cells were treated with 5FU for 48h

and immunoblotting was performed with indicated antibodies. **E)** PDC2 MINK1WT cells were

implanted in right upper flank of athymic male nude mice and PDC2 MINK1KO cells were

implanted in left upper flank, after which they were treated with 5FU at indicated concentration.

At the end of the experiment mice were euthanized, tumors were isolated and photographed

(n=5). **F)** Tumor growth was measured in indicated time points using digital slide caliper and

plotted as a graph (mean ± SEM, n = 5). Two-way ANOVA. **G)** Bar diagram indicates the tumor

weight measured at the end of the experiment (mean ± SEM, n = 5). Two-way ANOVA. **H)**

After completion of treatment, tumors were isolated and paraffin-embedded sections were

prepared as described in materials and methods to perform immunohistochemistry with indicated

antibodies. Scale bars: 50 μm. **I, J)** Lateral view of fluorescent transgenic [Tg(fli1:EGFP)]

zebrafish embryos at Day 0 and Day 5 injected with Dil-Red stained PDC2 control and MINK1

KO cells with and without treatment of 5FU **(J)**. The tumor growth and migration was assessed

by an increase in fluorescence intensity on the 5th day compared to the day of injection. n=6.

The quantitation of fluorescence intensity **(I)** was performed using ImageJ software and

represented as fold change of fluorescence intensity where day 0 reading was taken as baseline.

K) Zoomed image of distal part of embryo (5days post injection) to monitor migration of tumor

cells.

Figure 3: Ectopic overexpression of MINK1 rescued the drug resistant phenotype in

MINK1KD drug resistant OSCC: A) Using a lentiviral approach, 5FU resistant OSCC lines

and PDC2 were stably transfected with ShRNA which targets 3'UTR of MINK1 mRNA

(MINK1 UTRKD). For ectopic overexpression, pLenti CMV/TO Puro DEST MINK1 and

control vector were transiently transfected to indicated MINK1 UTRKD cells and

immunoblotting (n=3) was performed with indicated antibodies. **B)** MINK1 was ectopically

overexpressed in 5FUR cells stably expressing MINK1ShRNA targeting UTR and treated with

5FU at indicated concentration for 48 h, after which cell viability was determined by MTT assay

(n=3), 2-way ANOVA. **C)** Cells were treated as indicated in B panel and cell death (early and

late apoptotic) was determined by annexin V/7AAD assay using flow cytometer. Bar diagrams

indicate the percentage of cell death with respective treated groups (Mean ±SEM, n=3), 2-way

ANOVA. **D)** MINK1 was overexpressed in 5FUR cells stably expressing MINK1ShRNA

targeting UTR and treated with 5FU for 48h, after which immunostaining was performed for γ-

H2AX as described in materials and methods. **E)** MINK1 was overexpressed in chemoresistant

cells stably expressing MINK1ShRNA targeting 3' UTR, followed by 5FU treatment for 48

hours, and immunoblotting (n = 3) was performed with indicated antibodies. **F)** 5FU sensitive

OSCC lines were transfected with pLenti CMV/TO Puro DEST MINK1 followed by treatment

with 5FU at indicated concentration for 48h, after which cell viability was determined by MTT

assay (n=3), 2-way ANOVA.

Figure 4: MINK1 regulates the expression of p53 in 5FU resistant OSCC lines through

AKT/MDM2

A) High-throughput phosphorylation profiling with 1,318 site-specific antibodies from over 30

signaling pathways was performed in the lysates of MINK1KO and MINK1WT clones of H357

5FUR cells as described in methods. The top 20 upregulated phosphoproteins (MINK1

KO/MINK1WT) is represented in left panel, whereas top 20 downregulated phosphorylated

proteins is represented in right panel. The upregulated targets considered in the study is marked

in green red box, whereas downregulated targets in green box. **B)** Lysates were collected from

indicated cells and immunoblotting was performed with indicated antibodies. **C)**

pDESTCMV/TO MINK1 (ectopic overexpression of MINK1) was transiently transfected in

5FUR lines stably expressing MINK1 ShRNA (targeting 3'UTR) and immunoblotting was

performed with indicated antibodies. **D)** Lysates were collected from indicated cells and

immunoblotting was performed with indicated antibodies. **E)** MINK1 was ectopically

overexpressed in 5FUR lines stably expressing MINK1 ShRNA (targeting 3'UTR) and

immunoblotting was performed with indicated antibodies. **F)** pLNCX myr HA Akt1 (ectopic

overexpression of myr AKT) was transiently transfected in indicated MINK1KO cells and

immunoblotting was performed with indicated antibodies.   **G)** MINK1 was ectopically

overexpressed in 5FUR lines stably expressing MINK1ShRNA (UTRKD) as described in panel

C and treated with AKT inhibitor (Akti-1/2) for 24h and immunoblotting was performed with

indicated antibodies. **H)** Lysates were collected from indicated cells and immunoblotting was

performed with indicated antibodies.

Figure 5: Evaluation of Lestaurtinib as a MINK1 inhibitor to restore 5FU sensitivity in drug resistant OSCC: A) *In vitro* MINK1 kinase assay was performed using three compounds potentially binding to MINK1 (based on IUPHAR database). All compounds (10 μM) were incubated with recombinant human MINK1 along with substrate MBP and ATP and further subjected to ADP-Glo™ Kinase Assay as described in materials and methods section. **B)** Determination of EC50 value for kinase activity of top two MINK1 inhibitors selected from panel (A). **C-D)** Selection of highest dose of Lestaurtinib and Pexmetinib that does not affect cell viability when treated alone (viability > 80%) in 5FU resistant OSCC lines (n=3), 2-way ANOVA. **E)** 5FU resistant cells were treated with indicated dose of MINK1 inhibitor (50 nM Lestaurtinib, 500 nM Pexmetinb) in combination with increasing concentrations of 5FU for 48 h, after which cell viability was determined by MTT assay (n=3), 2-way ANOVA. **F)** 5FU resistant OSCC lines and PDC2 cells were treated indicated dose of MINK1 inhibitor (50 nM Lestaurtinib, 500 nM Pexmetinb) in combination with increasing concentrations of 5FU for 48 h, after which cell death (early and late apoptotic) was determined by annexin V/7AAD assay using flow cytometer. Bar diagrams indicate the percentage of cell death with respective treated groups (Mean ±SEM, n=3). Two-way ANOVA. **G)** Indicated 5FU resistant OSCC lines and PDC2 cells were treated with 5 μM of 5FU and/or 50nM of Lestaurtinib for 48h, after which immunostaining was performed for γ-H2AX as described in materials ad methods. **H)** Indicated 5FU resistant OSCC lines and PDC2 cells were treated with Lestaurtinib for 48h, after which immunoblotting was performed with indicated antibodies. **I)** Effect on 5FU IC50 upon Lestaurtinib treatment in cells with or without MINK1 knockout in indicated 5FU resistant OSCC lines and PDC2 cells (n=3), $*P < 0.05$ by 2-way ANOVA.

Figure 6: Lestaurtinib and 5FU synergistically reduced tumor burden *in vivo* in drug resistant OSCC: A) Patient-derived cells (PDC2) were earlier established from tumor of chemotherapy (TPF) non-responder patient. PDC2 were implanted in the right upper flank of athymic male nude mice, after which they were treated with 5FU and/or Lestaurtinib at indicated concentrations. At the end of the experiment mice were euthanized, and tumors were isolated and photographed (n = 5). **B)** Bar diagram indicates the tumor weight measured at the end of the experiment (mean ± SEM, n = 5). Two-way ANOVA. **C)** Tumor growth was measured at the indicated time points using digital slide caliper and plotted as a graph (mean ± SEM, n = 5). Two-way ANOVA. **D)** After completion of treatment, tumors were isolated, and paraffin-embedded sections were prepared as described in Methods to perform IHC with indicated antibodies. Scale bars: 50 μm. **E)** Schematic presentation of the mechanism by which MINK1 regulates p53 expression through AKT/MDM2 axis**.**

Figure 7: Evaluation of combinatorial anti-tumor effect of low dose of cisplatin, 5FU and lestaurtinb in TPF resistant patient derived cells (PDC2). A-B) PDC2 cells were treated with indicated concentrations of cisplatin, 5FU and lestaurtinib for 48h and cell viability was measured by MTT assay(n=3 and *P < 0.05 by 2-way ANOVA). **C)** PDC2 cells were treated with indicated concentrations of cisplatin, 5FU and lestaurtinib for 48h after which cell death was determined by annexin V/7AAD assay using flow cytometer. Bar diagrams indicate the percentage of cell death (early and late apoptotic) with respective treated groups (Mean ±SEM, n=3 by Two-way ANOVA). **D)** PDC2 cells were treated with indicated concentrations of cisplatin, 5FU and lestaurtinib for 48h and immunoblotting was performed with indicated antibodies. **E)** PDC2 cells were treated with indicated concentrations of cisplatin, 5FU, lestaurtinib for 12 days and colony forming assays were performed as described in method

651   section. Left panel: Bar diagram indicate the relative colony number (n=3 and *P < 0.05 by 2-

652   way ANOVA). Right panel: representative photographs of colony forming assay in each group.

Figure 01

A

Strategy for CRISPR based kinome knock out screening in 5FU resistant OSCC lines

B

- cell viability of sgRNA alone < 70%
- cell viability of sgRNA alone > 70%

Selected (n = 334)

Rejected (n = 506)

C

MINK1 SBK1 FKBP1A

D

H357 5FUR   SCC4 5FUR
SCC95FUR   H357 CisR

↓

Lenti sgRNA targeting top 60 kinases
from primary screen
+
Sublethal dose
of 5-Fluorouracil

↓

Overlapping candidates
among 5FUR cell lines

H357 5FUR

ROCK2
STK16
IKBKB
SGK223

ROR2   MORN1
MINK1   TEK

NTRK3   SH3BP4
NTRK2   SBK1   NOL9
NLK   FKBP1A   CNTLN
SQSTM1   CDK19
MAPK3   GTF2H2   PKN1
        GTF2H4

SCC4 5FUR   SCC9 5FUR

Relative cell viability
Low [gradient] High

Secondary kinome screening in 5FU resistant OSCC

E

MINK1 KO (-5FU)   MINK1 KO (+5FU)

SBK1 KO (-5FU)   SBK1 KO (+5FU)

FKBP1A KO (-5FU)   FKBP1A KO (+5FU)

-VE CONTROL (-5FU)   -VE CONTROL (+5FU)

F

150 kD — MINK1

42 kD — β-ACTIN

G

Chemotherapy responder

Chemotherapy non-responder

H

Chemotherapy Responder (n=11)   Chemotherapy Non responder (n=23)

I

Before Chemotherapy

After Chemotherapy

J

- Before chemotherapy
- After chemotherapy

Patient 1   Patient 2   Patient 3   Patient 4   Patient 5

Figure 02

Figure 03

Figure 04

A

Phosphorylated proteins	Phosphorylation ratio of MINK1 KO/ WT
p53 (Phospho-Ser33)	2.387529
ALK (Phospho-Tyr1507)	2.187586
GSK3 beta (Phospho-Ser9)	2.172644
Gab2 (Phospho-Ser643)	2.016842
AKT1S1 (Phospho-Thr246)	2.008168
p53 (Phospho-Ser15)	1.920548
c-Jun (Phospho-Ser63)	1.897019
Catenin beta (Phospho-Ser654)	1.799196
Keratin 8 (Phospho-Ser432)	1.757952
IL-13R/CD213a1 (Phospho-Tyr405)	1.646146
Dok-2 (Phospho-Tyr299)	1.604889
VEGFR2 (Phospho-Tyr1175)	1.60054
GSK3 alpha (Phospho-Ser21)	1.588892
Androgen Receptor (Phospho-Ser213)	1.566496
Tau (Phospho-Thr181)	1.565912
IGF2R (Phospho-Ser2409)	1.56576
eIF2A (Phospho-Ser51)	1.559497
CDK5 (Phospho-Tyr15)	1.547983
14-3-3 zeta (Phospho-Ser58)	1.522109
4E-BP1 (Phospho-Thr45)	1.505419

Phosphorylated proteins	Phosphorylation ratio of MINK1 KO/ WT
HDAC1 (Phospho-Ser421)	0.32937
AKT1 (Phospho-Ser473)	0.351857
LYN (Phospho-Tyr507)	0.356728
Abl1 (Phospho-Tyr204)	0.403395
HSL (Phospho-Ser554)	0.406255
ATP1A1/Na+K+ ATPase1 (Phospho-Ser23)	0.454414
EEF2 (Phospho-Thr56)	0.47177
TYK2 (Phospho-Tyr1054)	0.506242
JAK2 (Phospho-Tyr1007)	0.532182
Chk2 (Phospho-Ser516)	0.538779
FLT3 (Phospho-Tyr599)	0.558251
EGFR (Phospho-Tyr869)	0.560672
GluR1 (Phospho-Ser849)	0.581223
AurB (Phospho-Thr232)	0.590025
AKT2 (Phospho-Ser474)	0.594925
Integrin beta-1 (Phospho-Thr788)	0.595157
DYN1 (Phospho-Ser774)	0.596312
MDM2 (Phospho-Ser166)	0.61482
IL-2RA/CD25 (Phospho-Ser268)	0.627742
Caspase 9 (Phospho-Tyr153)	0.630743

Figure 05

A — Relative MINK1 Kinase Activity for DMSO, Staurosporine (10μM), Pexmetinib (10μM), Lestaurtinib (10μM)

B — Relative MINK1 Kinase Activity

Lestaurtinib (μM): 0 .01 .02 .05 .1 .2 .5 1 10 - - - - - - - - -
Pexmetinib (μM): - - - - - - - - - 0 .01 .02 .05 .1 .2 .5 1 10

C — % Cell Viability vs Lestaurtinib (nM): H357 5FUR, SCC4 5FUR

D — % Cell Viability vs Pexmetinib (nM): H357 5FUR, SCC4 5FUR

E

H357 5FUR — ● 5FU ■ 5FU+Lestaurtinib 50nM ▲ 5FU+Pexmetinib 500nM

SCC4 5FUR — ● 5FU ■ 5FU+Lestaurtinib 50nM ▲ 5FU+Pexmetinib 500nM

Treatment groups	5FU alone	5FU+Lestaurtinib 50nM	5FU+Pexmetinib 500nM
5FU IC50 (μM)	20.49 ± 0.84	4.82 ± 0.27	7.08 ± 0.68

Treatment groups	5FU alone	5FU+Lestaurtinib 50nM	5FU+Pexmetinib 500nM
5FU IC50 (μM)	22.85 ± 0.17	5.36 ± 0.33	9.45 ± 0.26

F — % Cell death (AnnexinV+7AAD): H357 5FUR, SCC4 5FUR, PDC2

G — DAPI / γH2AX / MERGE: H357 5FUR, SCC4 5FUR, PDC2

5FU (5μM) / Lestaurtinib (50nM)

H — Western blot: p-MDM2 Ser 166 (90 kD), MDM2 (90 kD), p-AKT Ser 473 (60 kD), AKT (60 kD), P53 (53 kD), β-ACTIN (42 kD); Lestaurtinib (100nM) − + for H357 5FUR and SCC4 5FUR

I — 5-FU IC50 (uM): H357 5FUR, SCC4 5FUR, PDC2

Lestaurtinib (50nM) / MINK1 sgRNA #1

Figure 06

A

PDC2 Vehicle Control

PDC2 5FU(10mg/kg)

PDC2 Lestaurtinib (20mg/kg)

PDC2 5FU (10mg/kg) + Lestaurtinib (20mg/kg)

B

- PDC2 Vehicle Control
- PDC2 5FU (10mg/kg)
- PDC2 Lestaurtinib (20mg/kg)
- PDC2 5FU (10mg/kg) + Lestaurtinib (20mg/kg)

C

- PDC2 Vehicle Control
- PDC2 5FU (10mg/kg)
- PDC2 Lestaurtinib (20mg/kg)
- PDC2 5FU(10mg/kg) + Lestaurtinib (20mg/kg)

D

PDC2 Vehicle Control | PDC2 5FU | PDC2 Lestaurtinib | PDC2 5FU + Lestaurtinib

CD44

Cleaved Caspase 3

Ki67

E

5FU Resistant OSCC cells

Figure 07

A

B

C

D

E