1

New relationships are not necessarily more variable: female degus are more consistent with

strangers, male degus are less

Short title: Variability of interactions in new peer relationships

Amber Thatcher[1], Nathan Insel[1*]

Department of Psychology, University of Montana, Missoula, Montana, USA

*correspondence should be addressed to nathan.insel@umontana.edu, University of Montana,

Skaggs Rm 143, 32 Campus Dr., Missoula, MT, USA

Abstract

When an organism explores a new environment or stimulus it varies its behavior to ensure proper sampling. As contingencies are learned, variance can give-way to routines and stereotypies. This phenomenon is common across species but has not been well studied in the social domain, in which the stimulus an animal investigates, another individual, may react negatively to unexpected behaviors. Here we investigate the effects of social familiarity on interaction variability in degus, female members of which are predisposed to form relationships with new, same-sex individuals. Degus were presented with a series of 20 minute, dyadic "reunion" sessions across days, interleaving exposures to familiar and unfamiliar same-sex conspecifics. We found that session-to-session variability in males was higher between strangers compared with cagemates, suggesting males may establish relationships by testing different social roles. In contrast, following an initial exposure, female strangers showed lower session-to-session change compared with cagemates, potentially establishing new relationships by maintaining behavioral norms. Social novelty did not appear to affect variability of interaction timing within a session. Given ecological pressures on female degus to form large, stable social networks, the data are consistent with the notion that higher behavioral variability across encounters is maladaptive for establishing cooperative peer relationships.

Keywords

degu, exploration, learning, peer, relationship, social

Introduction

Novelty evokes exploratory behaviors in animals ranging from invertebrates to mammals [1–3]. During exploration, animals vary their behavior to sample across the dimensions of a stimulus or environment: hands are moved around to rotate and grip new objects, variable eye movements help view a new scenes, and bodies are moved through the landmarks and edges of a new place. As the structure and values of a stimulus space become learned, behavioral variability decreases, movements become more restricted to sample only the relevant features and can, eventually, become stereotyped habits.

Social novelty can also increase investigative behavior, but the effects of novelty on behavioral variability have not been well studied. By analogy with exploration of space or other stimuli, strangers may initially try-out different behaviors, later converging to particular sets and sequences of interactions as animals learn how best to respond to one-another. There is indirect evidence across animal taxa that interaction variability decreases as relationships form. When fruit flies lose a fight with a particular conspecific, for example, they alter the types of fighting movements used with that individual in the future [4]. Similarly, the formation of dominance relationships in male mice can be observed as an increasingly stable asymmetry of aggressive versus subordinate behaviors within the dyad [5]. These particular examples involve competitive relationships, raising the question of whether social familiarity is also associated with reduced variability in cooperative social relationships. Directly comparing competitive and cooperative relationships can be difficult, in that traditional methods used to investigate relationship formation rely on measures specific to the type of relationship and species studied. It is therefore important to construct metrics that generalize across relationship types, allowing for comparisons between them.

The present study was designed to examine the effects of social novelty on interaction variability in degus (*Octodon degus*), female members of which are predisposed to form and maintain new, cooperative peer relationships. Unrelated female degus nest together and nurse one-another's young [6–10], and reproductive success is associated with larger, more stable

social groups [10,11]. Unrelated male degus are, in contrast, less likely to cohabitate, and

reproductive success is not linked to group size [11]. In laboratory settings, female degus

interact with strangers more than familiar peers [12], even when they have the option to

choose between the two [13]. They also quickly become affiliative with stranger peers in ways

that have not been observed in other laboratory rodents [12–14]. Degus therefore offer a

unique case study to investigate behavioral patterns associated with the formation of social

relationships, one that allows a direct contrast between potentially more competitive

relationship types in males and more cooperative in females.

The principal hypothesis of the present work is that animals are more variable in their social

interactions when exploring novel conspecifics, and that this holds irrespective of social

relationship type. We predict that both male and female degus, with potentially more

competitive and cooperative relationships respectively, show decreases in variability as long-

term relationships are established. To assess this, we expose same-sex strangers to one-

another multiple times over 20 to 40 days and apply measures of interaction variability within

and between the reunions. For controls, we also examine the interactions of already-familiar

cagemate dyads during the same time period, and subsequently "new cagemates" and "new

strangers". Contrary to expectations, we find that relative variability in strangers versus

cagemates are in the opposite direction in female compared with male degus, suggesting that

behavioral consistency between encounters depend on the nature of an animal's relationship.

The present study focuses on ethogram-based classification of social interactions; however, one

goal of this work has been to establish measures that can be flexibly applied across modes of

behavioral observation, species, and setting.

Results

Degus interact more with strangers

To help establish validity of the dataset and assess general effects of social novelty, we first

tested whether degus interact more with strangers [12,13,15] Degus were separated from a

cagemate partner for 24 hours and then united with either a cagemate or stranger in a 50 x 50

cm experimental recording chamber for 20 minutes while video was collected (a "reunion"

session, **Figure 1A**). The procedure was repeated 5 times each for cagemates and strangers in

an interleaved way, counterbalancing the order between dyads (**Figure 1B**). Consistent with

previous findings, degus interacted more with strangers than cagemates (3-factor mixed

ANOVA across individuals with exposure day and strangers/cagemate as repeated measures,

sex as between-subject; effect of stranger/cagemate: $F(1,94)=34.18$, $p=7.2 \times 10^{-8}$; **Figure 1C**).

Interaction levels were also higher in males across both stranger and cagemate groups (effect

of sex: $F(1,94)=12.47$, $p=6.4 \times 10^{-4}$). Interactions were subdivided into five types: agonistic,

allogrooming, rear-sniffing, face-to-face, and body-sniffing. There was relatively high variability

in the levels and types of interactions both within and across animals over the five cagemate

and stranger exposures (**Figure 1D**). However, across the five exposures female degus

interacting with strangers showed overall higher levels of agonistic interactions, rear-sniffing,

and face-to-face compared against cagemates (2-factor mixed ANOVA, repeated measures over

days, false discovery rate correction for 5 tests, agonistic: $F(1,51)=8.70$, $p=0.0048$, rear-sniffing:

$F(1,51)=12.37$, $p=9.3 \times 10^{-4}$, and face-to-face: $F(1,51)=10.32$, $p=0.0023$; **Figure 1E**, left panels). In

males, only agonistic behaviors consistently distinguished strangers and cagemates across

exposures ($F(1,43)=17.44$, $p=1.4 \times 10^{-4}$, **Figure 1E**, right panels).

Figure 1. Design and interaction levels. A) Degus received a repeated sequence of 24 hr isolation, 20 minute reunion recording sessions, and 24 hr co-habitation with cagemates. CAG = cagemate reunion, STR = stranger. B) Example sequence of recording sessions. Following interleaved cagemate and stranger reunions, female strangers were co-housed creating "new cagemates" (STR->NC). After all sessions degus were tested with a new stranger (NST). C) Total interaction levels for females (left) and males (right) across the 5 initial exposures (mean ± SEM; statistics are stranger vs. cagemate in 2-way ANOVA). D) Colormaps of interaction types for each degu in 2 s bins across the five exposures. Individual degus (rows) are sorted according to their peak levels of agonistic interactions (AG, red; other abbreviations: GR = allogrooming, RS = rear-sniffing, BS = body sniffing, FF = face-to-face). Higher concentration of face-to-face at session starts helps demarcate the 5 sessions. E) Breakdown of interaction levels across different types of interactions. Higher levels of agonistic, rear-sniffing, and face-to-face were observed in female strangers relative to cagemates (left panels). In males (lower panels), some interaction types decreased and others increased over days. F) Interaction levels over days in female degus that interacted more with strangers (SC-HIGH, top) and those that did not (SC-LOW, bottom). Levels of social interaction

139 **between new cagemates (exposures 6 through 9), as well as levels in new strangers (blue),**
140 **were consistent with those observed in initial strangers on exposures 1 through 5.**
141

142 We predicted that the difference in interaction levels between strangers and cagemates would

143 diminish as initial strangers became more familiar. This was confirmed by a statistical

144 interaction between stranger/cagemate and exposure day (stranger/cagemate x exposure:

145 $F(4,376)=3.16$, $p=0.014$). The pattern was particularly apparent in males, in which face-to-face

146 interactions were more prevalent between strangers during initial exposures

147 (stranger/cagemate x exposure number, false discovery correction for 5 behaviors:

148 $F(4,172)=4.00$, $p=0.0040$, **Figure 1E**), with other, non-agonistic behaviors showing similar trends

149 (e.g., rear-sniffing: $F(4,172)=2.36$, $p=0.056$). Female degus, in contrast, did not show evidence of

150 diminishing stranger-cagemate differences across days (all interactions: $F(4,204)=0.48$, $p=0.75$;

151 individual interaction types: $p >> 0.05$).

152

153 To further examine the absence of familiarity effects in females, we subdivided female degus

154 into those that showed significantly higher levels of interaction with strangers ("SC-HIGH", 16 of

155 52 degus, determined using a paired t-test on each animal across 5 stranger, 5 cagemate

156 sessions, $p < 0.1$) and those that did not ("SC-LOW", 36 of 52 degus). There was no evidence

157 that stranger-cagemate interaction levels changed over days in either the SC-HIGH animal (2-

158 factor repeated-measures ANOVA stranger/cagemate: $F(1,15)=83.40$, $p=1.63 \times 10^{-7}$, day:

159 $F(4,60)=0.89$, $p=0.47$, stranger/cagemate x day: $F(4,140)=0.20$, $p=0.94$) or the SC-LOW animals

160 (stranger/cagemate: $F(1,35) = 8.0 \times 10^{-4}$, $p=0.98$, day: $F(4,140)=1.49$, $p=0.22$, stranger/cagemate

161 x day: $F(4,140)=0.20$, $p=0.94$; **Figure 1E**). A possible explanation is that the 20 min reunion

162 sessions provided insufficient time for partners to become fully familiarized with one-another.

163 To test this, a subgroup of 40 female degus were co-housed with their stranger partners for 1

164 day creating new cagemates, then tested in two more reunions, creating "new cagemates". A

165 smaller subgroup was further co-housed an additional 7 days and tested again twice.

166 Unexpectedly, interactions between SC-HIGH strangers, as new cagemates, remained higher

167 than long-term cagemates following either 1 day (paired t-test, averaging together all cagemate

168 sessions and both new cagemate sessions, $t_9=5.50$, $p=3.80 \times 10^{-4}$) or 7 days of co-housing

(t_7=2.98, p=0.021). To investigate whether interaction levels were due to genetic relatedness,

we compared sibling versus non-sibling pairs. As only two of the cagemate dyads were non-

siblings, this could not be evaluated statistically; however, both members of one, non-sibling

cagemate dyad interacted more with strangers (i.e., they were also SC-HIGH animals) while the

members of the other dyad did not (were SC-LOW animals). Overall, therefore, female degus

exhibited higher interaction levels with strangers than cagemates but there was no evidence

these attenuated with familiarity and; consistent with research from other labs [15], this was

unlikely to be due exclusively to genetic relatedness.

In males, the differences between interactions with strangers versus cagemates decreased with

familiarity, though this pattern could have been influenced by familiarity with the protocol and

experimental apparatus. As a final control condition, degus were exposed to a new, unfamiliar

conspecific ("new stranger"). In both males and SC-HIGH females, levels of interaction with new

strangers were higher than with cagemates (paired t-test between within-animal averages in

cagemates and new-strangers, males: t_{43}=5.13, p=6.66x10^{-6}, SC-HIGH females: t_{15}=4.17, p=

8.18x10^{-4}) and were not lower than interactions with the original strangers (males: t_{43}=1.76,

p=0.086 with new stranger > old stranger; SC-HIGH females: t_{15}=1.13, p=0.28, **Figure 1F**).

In summary, degus tended to interact more with strangers than with cagemates, though in

females this was only the case for one-third of the animals (SC-HIGH group), which continued to

interact more even after strangers were co-housed to become new cagemates. Both male

degus and SC-HIGH females also interacted more with new strangers than with cagemates.

In males, strangers have higher session-to-session variability that cagemates; in females,

strangers have lower session-to-session variability

The overarching hypothesis of this study is that social interactions are more variable between

unfamiliar individuals. Variability can be assessed at multiple timescales, and we began by

examining the consistency of a dyad's interactive behavior between 20 minute reunion

sessions. To measure this, we computed interaction vectors: social interaction time budgets

across each of 5 interaction types (agonistic, allogrooming, rear-sniffing, body-sniffing, and

face-to-face), in each of the two individuals in a dyad. We then computed two, complementary

measures: distances (dissimilarity) between interaction vectors, and dyad classification success

(the ability of a support vector machine, SVM, to correctly identify which dyad an interaction

vector belongs to). While the distance-based method allowed observation of overall variability,

the classifier-based method established whether dyads could be distinguished from one-

another using combinations of interaction vector elements (**Figures 2A&B**). Notably, the

classifier method measures within-dyad variability relative to differences between dyads, which

allows specific behaviors to be ignored if inconsistent within a dyad, thereby honing-in on

behavior combinations that more selectively define the social relationship.

Figure 2. Variability across sessions. A) Diagram illustrating analysis methods. Each column of
boxes represents the interaction vector from a single session (abbreviations same as in Figure
1D, body-sniffing omitted for illustration). To the right, vectors are plotted in cartesian
coordinates. Top right shows SVM-based classification success, with dashed line representing
a hyperplane separating groups in the relevant feature-space. Bottom panel shows
interaction vector distances, with points placed in 3 dimensions to emphasize that non-
relevant features contribute to the measure. B) Plot of all female dyad interaction vectors
using the first two principal components. Dark black dots show all 5 sessions from one
example cagemate dyad, red dots from one stranger dyad. In this example, there is higher
clustering (less session-to-session variance) in the stranger relative to cagemate. C) Swarm
plots showing distribution of within-dyad interaction vector distances (top: Euclidean,
bottom: cosine). F- and P-values refer to stranger/cagemate x sex interaction in a 2-way
ANOVA. D) Success rate of SVM classifiers discriminating pairs of dyads within each of the
four groups (errorbars are mean ± SEM). E) Within-dyad interaction vector distances across
exposure numbers (e.g., "exposure 1" only includes distances between exposure 1 interaction

vectors and other days). Values are across-dyad averages of within-dyad medians, ±SEM, F-
and P-values refer to stranger/cagemate x exposure number interaction using 2-way mixed
ANOVA. F) Relative success rates for different days (e.g., "exposure 1" only includes models
trained on exposures 2 through 5, then accuracy tested on exposure 1).
230
Across-session variability was found to differ between cagemates and strangers, but the

direction of this difference depended on sex. This could be observed using both interaction

vector distances and classifier success rates. Within-dyad distances tended to be lower than

between-dyad distances (within- versus between-dyad ratio of 0.73 and 0.69 for females and

males respectively), suggesting that individual dyads exhibited unique profiles. Within-dyad,

Euclidean distances between interaction vectors were found to be significantly higher in male

relative to female degus (2-way ANOVA, effect of sex: $F(1,92)=5.5$, $p=0.021$) significantly higher

in strangers compared with cagemates (effect of stranger/cagemate: $F(1,92)=32.1$, $p=1.7 \times 10^{-7}$)

but relatively higher in male strangers compared with female stranger (stranger/cagemate x sex

interaction: $F(1,92)=6.8$, $p=0.011$; **Figure 2C, top**). A concern with using Euclidean distance is its

sensitivity to total interaction levels; therefore, cosine distance was also used. Cosine

interaction vector distances showed differences between males and females (2-way ANOVA,

effect of sex: $F(1,92)=6.6$, $p=0.012$) and a statistical trend for stranger-cagemate differences to

differ between males and females (stranger/cagemate x sex interaction: $F(1,92)=3.4$, $p=0.070$;

Figure 2C, bottom). As described below, trend met criteria for statistical significance when

considering exposure number (**Figure 2E**). Contrasting with hypotheses, female strangers

appeared to show less variability (more consistency) across sessions relative to female

cagemates, male cagemates, or male strangers. This observation females did not appear to be

restricted to only dyads with SC-HIGH or only SC-LOW degus (**Supplementary Figure 1A&B**).

The interaction between sex and social context depended most on allogrooming and rear-

sniffing, as effects decreased when removing either of these components from the interaction

vectors, but increased when other interaction types were removed (2-way ANOVA of cosine

distances, stranger/cagemate x sex interaction after removing agonistic: $F(1,92)=4.3$, $p=0.040$;

allogrooming: $F(1,92)=2.1$, $p=0.15$; rear-sniffing: $F(1,92)=1.65$, $p=0.20$; face-to-face:

$F(1,92)=4.52$, $p=0.036$; body-sniffing, $F(1,92)=6.2$, $p=0.015$; **Supplementary Figure 2A**).

Variability differences between males and females appeared to depend more on face-to-face

and body-sniffing behaviors (effect of sex after removing agonistic: $F_{(1,92)}=7.2$, p=0.0085;

allogrooming, $F_{(1,92)}=8.3$, p=0.0050; rear-sniffing, $F_{(1,92)}=7.3$, p=0.0083; face-to-face:

$F_{(1,92)}=3.0$, p=0.086; body-sniffing: $F_{(1,92)}=2.7$, p=0.10).

Dyad classification success was evaluated using SVM models trained to discriminate the five,

full-session interaction vectors of one dyad from the five interaction vectors of another dyad

within the same group (for example, one SVM model might be trained to distinguish one

cagemate dyad from another cagemate dyad). Each model was repeatedly trained and tested

using a leave-one-out cross validation method. The average success rate was higher than

chance, confirming that individual dyads expressed behavioral patterns that were

distinguishable from other dyads (binomial probability distribution predicts ~5% probability of

success classifying at least 7 of 10 sessions, mean rates for female cagemates = 7.1, female

strangers = 7.8, male cagemates = 7.6, male stranger = 7.3). Rates significantly differed between

groups, with classification success higher in female strangers than cagemates, but lower in male

strangers than male cagemates (2-way ANOVA effect of stranger-cagemate: $F_{(1,1108)}=7.1$,

p=0.0078, effect of sex: $F_{(1,1108)}=0.03$, p=0.86, stranger-cagemate x sex interaction:

$F_{(1,1108)}=23.6$, $p=1.4 \times 10^{-6}$; **Figure 2D**). The effects were not due to any one interaction type, as

the sex difference remained even after removal of any specific type from the interaction

vectors (stranger-cagemate x sex interaction for each: F >20, p <<0.01, **Supplementary Figure**

2B). Higher classification success in strangers was found when considering only female dyads

with SC-HIGH degus, and when classifying between dyads with only SC-LOW degus

(stranger/cagemate: $F_{(1,313)}=6.2$, p=0.012, SC-HIGH vs. SC-LOW: $F_{(1,313)}=5.4$, p=0.020,

stranger/cagemate x SC-HIGH/LOW interaction: $F_{(1,313)}=0.11$, p=0.74; **Supplementary Figure**

1C&D).

We next asked how variability changed over exposures. In females, there was a significant

increase in stranger-cagemate differences over days, with variability in stranger dyads

decreasing after the first exposure (2-way mixed ANOVA, repeated measures over days, cosine

distance: cagemate-stranger: $F_{(1,47)}=2.19$, p=0.14, exposure day $F_{(4,188)}=1.9$, p=0.10,

cagemate-stranger x exposure day: F(4,188)=2.75, p=0.030); classification success: cagemate-

stranger: F(1,699)=29.6, p=7.4x10^{-8}, exposure day: F(4,2796)=6.6, p=2.53 x 10^{-5}, cagemate-

stranger x exposure day: F(4,2796)=5.65, p=1.6x10^{-4} ; **Figures 2E&F, left panels**). In males,

classification success rates appeared to increase in cagemates after the first exposure, while

strangers showed less evidence of change (2-way mixed ANOVA, stranger-cagemate:

F(1,503)=1.5, p=0.22, exposure day: F(4,2012)=6.1, p=7.74x10^{-5}, stranger-cagemate x exposure

day: F(4,2012)=4.54, p=0.0011; **Figures 2E&F, right panels**). These patterns were

counterintuitive, in that strangers did not converge to become more like cagemates with

familiarity; instead, after an initial social exposure in the experimental chamber both female

strangers and male cagemates began interacting similarly across days, while female cagemates

and male strangers either remained or became less consistent.

Within-sessions, interaction patterns were not strongly affected by social context

There are no standard measures for variability of social interaction. We selected four measures

that provide estimates of variation in the sequencing of actions, the timing between actions,

and in blocks of interactions within sessions.

To assess sequencing between actions, we computed transition matrices describing the

probability that a "reference" interaction type will be followed by another interaction type. The

majority of interactions showed statistically significant relationships (t-test across sessions of

probabilities minus chance, baseline probabilities, Benjamini & Hochberg false-discovery

correction across 20 comparisons for same-degu interaction pairings and 25 between-degu

interaction pairings; **Figure 3A**). The most notable exception was an absence of association

between agonistic in one individual with allogrooming in the other. In males, there were also no

observable pairings between rear-sniffing and agonistic behaviors across two members of a

dyad.

Figure 3. A) Transition matrices between interaction types, averaged across all degus (top:
female, bottom: male). Left matrices show probability of the next action (columns) given a
reference action (rows) performed by the same degu, right matrices shows probability that
the other degu will act, given a reference action in the first degu (matrices computed
independently). All cells showed a significant, non-zero probability aside from those labeled
"n.s." Diagonal black lines indicate same-action transitions not included in analysis. Acronyms
are the same as in Figure 1D. B) Sequence randomness of interactions across exposures. No
significant changes were observed over exposures, nor differences observed between dyad
groups in either female (left) or male (right) dyads. C) Example interaction cross-
correlograms. The top four emphasize cases of potential difference between cagemates and
strangers (thick lines are averages across dyads, thin lines SEM). Top two panels show male
levels of allogrooming before and after face-to-face (left) and levels of face-to-face before
and after allogrooming (right). Middle panels show female levels of body-sniffing, with
cagemates potentially more likely to body-sniff before and after agonistic encounters (left)
and less likely than strangers before and after allogrooming (right). Bottom panels show
levels of face-to-face interactions around rear-sniffing (left) and body-sniffing (right) in both
male (blue) and female (red) dyads. D) Local variance measures examining the consistency of

temporal intervals between specific interaction types. No differences between strangers and
cagemates were detected.

Although the transition matrices revealed structure in the sequencing of interactions, there was

no compelling evidence for overall variability differences between cagemate and stranger

groups. This was measured using a sequence randomness score, an index into the overall

degree to which behaviors can be predicted given a current behavioral state. Sequence

randomness was evaluated across exposures for same-degu and between-degu transition

matrices of both sexes. No apparent effects of stranger/cagemate or exposure were observed

(mixed 2-way ANOVA, between-subject variability of stranger/cagemate dyad, repeated

measures over exposures; female same-degu stranger/cagemate: $F(1,4)=0.32$, $p=0.57$,

stranger/cagemate x exposure: $F(4,128)=0.79$, $p=0.53$; female between-degu,

stranger/cagemate: $F(1,42)=1.74$, $p=0.19$, stranger/cagemate x exposure: $F(4,168)=1.36$,

$p=0.25$; **Figure 3B**). When female degus were split into pairs containing SC-HIGH animals and

those with only SC-LOW animals, a trend was observed in SC-HIGH for increased, between-degu

sequence randomness among strangers ($F(1,84)=3.40$, $p=0.080$; **Supplementary Figure 3**). The

low effect size and lack of generalizability across sexes, or across all females, emphasized that

any differences were very subtle.

Putting aside variability measures, there was also little evidence that stranger and cagemate

groups systematically differed in specific interaction transitions (two-sample t-tests, false

discovery corrections as above, corrected p-values >0.05). These results did not change when

only the first exposure day was considered, nor were differences found in the second-order

transition matrix (i.e., the probabilities that one interaction type leads to another, subsequent

to the next). Cross-correlations were used to more precisely evaluate the interaction pairings.

As with the transition matrices, and as previously observed [12] most pairs of actions showed

temporal relationships; however, differences between strangers and cagemates did not exceed

threshold for statistical significance (**Supplementary Figure 4**). It is possible that the correction

for multiple comparisons occluded subtle differences between cagemate and stranger

interaction sequencing. Taking a data exploration (hypothesis-generation) angle, a few specific

interaction pairings may have been less likely in strangers: 1) Male strangers may have been

less likely to follow face-to-face encounters with allogrooming (two-sample t-test, p=0.020;

Figure 3C, top-panels), consistent with observations made in females in a prior study ("nuzzling

after grooming", [12]. 2) Male strangers also may have been less likely to precede agonistic

behaviors with body-sniffing (p=0.008). 3) female strangers may have been less likely to body-

sniff before and after agonistic encounters (p=0.038 and p=0.030), but 3) more likely than

cagemates to body-sniff before and after allogrooming (p=0.021 and p=0.025).

Variability of the intervals between interactions of the same type were also analyzed. For

example: if face-to-face interactions are a way for two individuals to "check-in" after a period of

time had elapsed, these interactions may be more consistently paced in animals who know

one-another compared with those who do not. This was measured as the Local Variance of

inter-behavior-intervals (IBI's) for each of the 5 main interaction types There was no evidence

for differences in the Local Variance for any type of interaction (two-sample t-tests, false-

discovery rate correction, p >> 0.05; **Figure 3D**). The same was true when only the first day of

interaction was considered.

A set of final tests examined within-session variability in sets of interactions, by measuring

interaction vector distances across blocks of "n" interactions (where n was varied from 1 to 30).

As illustrated by **Supplementary Figure 5**, the analysis had fundamental limitations such that

distance scores appear higher for sessions with more interactions. When this was controlled

for, the data offered no evidence for higher levels of variability in strangers compared with

cagemates for either sex.

Discussion

The present studies were designed to test whether relationship formation involves an initial

period of higher variability in the interactions between two individuals. Previous evidence

suggests that male and female degus are predisposed to different types of peer relationships,

offering an opportunity to compare variability patterns associated with the formation of

different types of relationships. We found that interaction variability differed between new and

established relationships, but the differences were not consistent between male and female

groups. Male stranger dyads showed evidence for higher variability across reunions compared

with cagemates—particularly when estimated using dyad classification success. Female

stranger dyads, however, were less variable than cagemates across reunions, a reduction that

became apparent after either the first or second exposure session when measured using dyad

classification success and interaction vector distance respectively. Analyses examining behavior

at shorter timescales, within sessions, revealed a structure that is consistent with previous

observations [12]; however, only subtle differences were observed between cagemate and

stranger groups, with no compelling evidence for higher variability in the latter. As discussed

below, degus did appear to interact more with strangers than with cagemates. Together, the

results suggest that social exploration may not be expressed as much by behavioral variability,

but by higher levels of engagement, in ways that conform to a familiar behavioral structure.

A primary question raised by the results of this study is why, after an initial social exposure,

female cagemate dyads exhibited higher levels of session-to-session variability than female

strangers. It is possible that the tendency to maintain consistent behavioral patterns helped

establish roles in the new relationships. Conversely, it may be that when female degus were

placed with a cagemate, they felt less constrained or attentive to the social situation, resulting

in behavior that depended more on day-to-day circumstantial factors (e.g., hormonal

fluctuations, ambient odors, room temperature, etc.). These explanations are consistent with

the notion that being unpredictable is less conducive to forming cooperative relationships.

Many primates use specific sets of vocal and physical interactions when encountering strangers,

often in stereotyped and ritual sequences. The circumstances in which these behaviors take

place—e.g., between similar-ranking males, younger individuals greeting older, and when

approaching pregnant females—suggest that the behaviors signal predictability, and function to

reduce uncertainty and tension [16–20]. To put another way, reduced variability may help avert

high-cost aggressive or competitive interactions. Male degus, unlike females, are not believed

to benefit from larger or more stable social groups [11]. Based on the present findings, we

hypothesize that a predisposition for more consistent interactions with strangers goes hand-in-

hand with a predisposition to form cooperative relationships with new individuals.

One way in which female and male degus were similar is that both interacted more with

strangers than with cagemates; however, this increase took-on different patterns. Male

strangers showed more greeting (face-to-face) interactions, and a trend for increased

investigation (rear-sniffing) during the first few exposures that subsequently attenuated

(agonistic interactions with strangers notably became more common after the first social

exposure). In female strangers, higher face-to-face, rear-sniffing, and agonistic encounters

persisted across reunions (agonistic again being more evident after the first day). Higher levels

of interaction with strangers is not surprising, and has been previously observed in our lab and

others [12,13,15]. It is also unsurprising given the species' gregarious behavior in natural

habitats: degu burrows are found near one-another, and members of different burrows greet

one-another frequently [9,21]. But if female degus form stable, cooperative peer relationships

with unrelated individuals, as ecological data suggest [6–10], it is difficult to understand why, in

the present study, new cagemates differed from old cagemates. We cannot rule-out the

possibility that the structure of the protocol played a role; for example, extended time away

from the original cagemates may have sensitized effects of the 24 hour isolation period. It is

also possible that degus form fundamentally different bonds with non-siblings encountered in

adulthood compared with conspecifics reared together. There are some hints of this from other

studies. Females in the wild have been reported to show preferences for members of their own

burrow groups during daytime foraging and social greetings [9]. In the lab, genetic kinship can

influence social behavior—just not as strongly as familiarity [15]. Future investigation using

similar protocols in young animals may help establish whether there are critical periods of

development in which certain types of peer bonds take place.

Another notable feature of the present data is that in both males and females, the structure of

social behavior was largely preserved between cagemates and strangers. There was no strong

evidence for differences in variability in sequencing or temporal relationships of interactions

within sessions. Some of the common structure of behavior between groups may be due to the

physics of interaction—e.g., two socially attentive degus are more likely to encounter one-

another's faces before body or tail. Or it may be due to the logic of investigative order: the

mouth may offer the most important information about another individual, the anogenital

region the next most, etc. Additionally, the sequencing and timing of behaviors might be

supported by genetic and early-life learning to ensure coordinated activity between individuals

of a species. The effects of juvenile social isolation on adult social behavior has been looked at

in detail in other species (e.g., chimpanzees: [22,23], rats: [24,25]). While early social

deprivation has been shown to have effects on social motivation, the types of interactions used,

and an animal's tendency toward aggression, the sequencing between interactions appears to

be relatively resilient to early-life social isolation [25]. Early-life social deprivation may still have

subtle effects on the timing of interactions, as well as how vocalizations are used [26], which

may make an animal less predictable and thereby destabilizing relationships to greater

escalation of aggressive/agonistic encounters [24]. Further work examining interaction

sequencing and the effect of early-life isolation in female degus will be valuable for

understanding the animals' abilities to engage constructively with strangers.

Several methodological factors may be critical to interpreting the present results. One of these

is the relatively unfamiliar setting of the reunion chamber. Although degus were individually

habituated to the chamber over five days prior to the start of the experiment, the room and

chamber were still relatively novel, and this may have impacted social behavior between both

cagemates and strangers, particularly on the first exposure day. Reactivity of social behavior to

environmental novelty has been previously reported in rats [27]. A second, potentially

important factor for understanding the results is the relatively few ethogram categories used to

track degu interactions. Face-to-face, for example, comprises both nose-to-nose and nose-to-

mouth contact, while rear-sniffing includes both anal and genital sniffing. Social interactions

can include a high dimensionality of subtle but relevant movements, such as flattening or

raising of ears, immobility versus rapid escape, and hesitant versus unrestrained approach.

Measuring these additional variables could provide a richer understanding of social dynamics

and moment-to-moment metrics of variability. On the other hand, the behaviors that are

measured—agonistic (primarily mounting), allogrooming, rear-sniffing, face-to-face, and body-

sniffing—are categories that likely have distinct social/ethological relevance for the animals.

The low numbers of interactions also simplified the process of identifying which were most

responsible for the general patterns. Thus, while the ethogram measures fail to capture the

depth of social interaction patterns, they may have helped streamline identification of the

similarities and differences between dyads.

While the present project focused its examination on the development of social relationships in

degus, an underlying goal was to advance methods that could generalize across species and

behavioral measurement tools. In some cases, it may only be possible to interpret an animal's

behavior through its specific ethogram. But while degus are unique in their sociality and peer

structures, these same methods could be applied to better understand the formation of

cooperative social relationships across species, sexes, and developmental stages. In this way,

the study of interaction variability joins a wider movement to apply standardized metrics

toward analysis of animal social behavior (e.g., see [28]). As the results demonstrate, however,

while distance metrics and classification success can be useful, they cannot be taken in

isolation.

Materials and Methods

Subjects

Fifty-two adult female degus aged seven to twenty-nine months (mean = 11.9 mo, median = 11

mo) and forty-four male degus aged six to twenty months (mean = 10.4, median = 10) were

used. A higher number of female degus were used than males due to a late addition of the

"new cagemate" groups in females (see Testing). All animals were housed in same-sex pairs

50.8 x 40.6 x 21.6 cm plastic cages in a breeding vivarium at the University of Montana. Degus

were fed a 1:1 mixture of chinchilla and guinea pig "Teklad" feeds (Envigo; Indianapolis, IN).

Animals were housed on a 12:12 h light/dark cycle, with all tests occurring during the light

(active) cycle. Each cage was also equipped with additional enrichment (e.g., hay, cardboard

enclosures, plastic "bones", and wooden blocks) and, prior to the start of the study, provided

with occasional dust baths and handling sessions. Cagemate dyads were created by pairing two

same-sex individuals together at the time of weaning. Animals used for stranger dyads were the

same as those for cagemates. Pairings were selected with an attempt to keep animals matched

for age. Strangers inhabited the same vivarium but in no cases had they been in physical

contact.

Apparatus

All sessions recorded video and audio using a Logitech HD Pro webcam C920 USB2 camera

recorded at 30 frames/s. Recording chamber was 50 x 50 x 50 cm painted and sealed wood.

Following each social exposure the chamber was cleaned using 70% ethanol then dried.

Testing

Prior to the reunion procedure, each animal was exposed to the testing chamber without other

animals present for at least 5 minutes each day over five days. Twenty-four hours before the

first testing session, cagemates were separated into individual cages in the vivarium. Testing

began after degus were transported to the testing room and the backside of one degu in each

pair was marked for identification (Pet Paint; Camarillo, CA). A pair of either cagemates or

strangers was then placed in the chamber and allowed to interact freely for 20 minutes, after

which the session ended and animals were returned to home cages in the vivarium with their

previous cagemate for 24 hours (**Figure 1A**). By placing cagemates together following each

reunion session, animals had the opportunity to "recalibrate" their relationships, preserving it

over the weeks of experiment. This cycle of co-housing, isolation, and testing was repeated,

with exposures to strangers and cagemates interleaved in a pseudo-randomized order

(alternating partners after either one or two reunions with the same partner) for a total of 5

cagemate and 5 stranger sessions (counterbalancing stranger-first and cagemate-first across

dyads; **Figure 1B**).

Following 10 reunion sessions, female stranger dyads were co-housed to create "new

cagemates". This was not performed with males to avoid aggression-related injury. Of the 26

stranger female dyads, 24 were made new cagemate and tested after 24 hours of co-housing

and 24 hours of isolation, and again after a second 48 hour cycle. A subset of 14 dyads was

further tested twice more after 7 days of co-housing. To prevent sampling bias, statistical

comparisons that included new-cagemates omitted animals that did not receive this control

condition.

Following re-housing reunion sessions in females, and the initial 10 reunion sessions in males,

degus were subsequently returned to their original cagemates and then, following 24 hours of

isolation, tested with a new stranger. The purpose of the "new stranger" reunions was to

establish whether changes in stranger behavior across repeated exposures were due to

increased social familiarity, rather than acclimation to the recording chamber or testing

protocol.

Behavioral Scoring

Scoring of physical behavior was performed using BORIS (Behavioral Observation Interactive

Research Software; [32]), which allows users to log events during video playback. All analyses

were custom-written in MATLAB and are freely available on GitHub.

Filenames for video recordings were changed to ensure raters were blinded to conditions.

Raters coded the start and end times of each observed behavior, the type of behavior, and the

animal that initiated the behavior. Although the ethogram comprised 17 individual behaviors,

for simplicity and consistency with our prior work [12], analyses collapsed behavior types into

the following categories: agonistic (mounting, biting, wrestling, boxing, marking, rear-push, tail

shaking), allogrooming (sniffing of neck or body with small, repetitive movements), rear-sniffing

(anogenital sniffing), face-to-face (nose-to-nose and nose-to-mouth sniffing/contact), and body

sniffing (sniffing toward neck and body). Approximately 52% of sessions were scored by author

AT and the remainder by 7 undergraduate students all directly trained by AT. Inter-rater

reliability (IRR) was estimated using sessions scored both by AT and another student, applying

the same transformations and measures used for the primary analyses. Total interaction levels

could vary up to a maximum of 20% between raters; however, follow-up examination revealed

that inter-rate variance was unlikely to contribute to the primary reported results as 1) sessions

were distributed randomly across students and 2) the magnitude of inter-rater error scaled

with interaction levels, and analysis controls were used to ensure that interaction levels could

not account for interaction variability patterns.

Measures

All measures were computed from ethogram-based scoring of behavior, performed manually in

BORIS. Examination of total interaction time across sessions revealed a tailed distribution that

closely resembled a gaussian distribution following a cube root transformation (**Supplementary**

Figure 7). To ensure validity of parametric statistical tests and distance metrics, a cube root

transform was applied to all "interaction time" measures before performing statistical tests.

Interaction time (time budget analysis) allowed comparison with previous datasets and

assessment of overall interaction level changes across days. Interaction time was computed as

a proportion of session time, and was considered both as a sum total across behavioral types

(agonistic, face-to-face, etc.), and separately between behaviors.

Between-session variability was estimated using "interaction vector distances" and "dyad
classification success".
595
Interaction vector distances were computed by fisrt calculating the proportion of time that each
animal spent using a particular type of interactive behavior during a session (an "interaction
vector") and then computing the distance between the vectors associated with different
sessions. The types of interactions used for interaction vectors were agonistic, allogrooming,
rear-sniffing, body-sniffing, and face-to-face contact. Total vector length was 9 elements, as
face-to-face interactions were considered mutual (not assigned to one or the other degu). To
compute distances between the interaction vectors, both Euclidean and cosine distance were
used. Euclidean distance takes into account total interaction levels, while cosine distance does
not. Euclidean and cosine distance are relatively simple, but have advantages over alternatives:
e.g., Jensen-Shannon distance (based on Kullback-Leibler divergence) ignores zero elements,
which would have misrepresented distances; the Pearson correlation is, like cosine distance,
scale invariant, and likely would yield similar values but ignore changes common across all
interaction types.
609
Dyad classification success was computed by finding the degree to which a group of interaction
vectors could be discriminated from another group. Classification success conflates two
variables: within-dyad variability, and between-dyad diversity. The advantage relative to
interaction vector distances is that a trained model should ignore types of social interactions
that are not useful for discriminating between two dyads, emphasizing those combinations of
interactions that are useful. Classification success used a binary support vector machine
classifier (SVM, "fitcsvm" in Matlab). SVM works by finding boundaries (i.e., hyperplanes)
through n-dimensional space that best differentiate between two sets of data. It's rate of
success in discriminating two sets of data offers a measure of within- versus between-set
variability. We used a leave-one-out cross validation method, in which the model is trained on
all vectors but one (of 10 interaction vectors, the model is trained on 9), and classification is
made on the remaining, repeated for each of the ten interaction vectors across the two dyads.

A linear kernel was used in favor of a radial basis function due to ceiling effects in the latter.

SVM classifiers were used in favor of neural network classifiers due to the algorithm's relative

tolerance for small training sets.

Within-session variability was estimated by examining action-to-action transitions and

timing, as well as by comparing interaction vector distances across sets of actions.

Action-to-action transition variability was computed using "sequence randomness". Sequence

randomness is inversely related to how well an animal's next action can be predicted, given its

previous action. Transition matrices were first computed describing the probability that one

type of interaction would lead to one of another type within a 10 s interval (10 s was chosen

based on temporal patterns observed in the interaction cross correlograms). Independent

transition matrices were computed for interactions performed by the same individual and

those performed by the other; for example, if one begins to rear-sniff, the "self" transition

matrix estimates the probability that the same animal would subsequently body-sniff,

allogroom, etc, while the "other" (between-animal) matrix estimates the probability that the

other individual would subsequently perform a body-sniff, allogroom, etc. Rransition matrices

did not include probabilities of "no-interaction". Sequence randomness was the Shannon

entropy for each row, averaged across rows:

$$SR = 1/n \sum_{p=1}^{n} \sum_{a=1}^{n} -P(x_a|y_p) log P(x_a|y_p)$$

Where x_a is interaction type "a", y_p is the immediately preceding interaction type, and "n" is the

number of interaction types.

Variability in action-to-action timing was estimated using Local Variance and cross correlations.

Local variance (LV) was used to compute inter-behavior-interval (IBI) variance, a measure

commonly used for variance in neuron spike intervals [33]. LV was defined as

$LV = 1/(n-1) \sum_{i=1}^{n-1} 3(T_i - T_{i+1})^2 / (T_i + T_{i+1})^2$

Where T_i is the time at which the ith interaction took place and "n" the number of spikes.

Local variance makes no assumptions about the distribution of time intervals between

interactions. Empirical evaluation of the IBIs indicate distributions conformed most closely to an

exponential (**Supplementary Figure 7**); i.e., behaviors often come in bursts, similar to what has

been described in previous reports [5].

Action-to-action interactions and timing were also evaluated using cross correlations (see also

[12]). Cross-correlations can be thought of as a histogram where, in this case, times preceding

zero (left side of the x-axis) showed the incidence of a target action relative to the initiation

time of another, reference action, and times following zero (right side of axis) showed the

incidence relative to reference action termination. To make cross-correlations more

interpretable, values were normalized by dividing by the proportion of time during the session

that a target action took place. Therefore, values of 1 in the cross correlation meant no

observable temporal relationship between two interaction types, and y-axis units were

normalized rates rather than standard correlation coefficients.

Within-session, interaction vector distances were computed using count-based interaction

vectors, where each vector element was the sum of interactions of each interaction type over a

block of n interactions. If block size n = 10, for example, the vector described how many of 10

consecutive interactions were rear-sniffing events performed by degu A, allogrooming events

performed by degu B, face-to-face bouts, etc. This measure helped identify the degree to which

dyads changed the set of interactions they were using after a number of behaviors had been

performed. Exploratory methods were used to investigate cagemate versus stranger

differences across block sizes (1 to 30 interactions per vector), across days. As described in

results, many sessions lacked sufficiently high numbers of interactions to yield a single distance

value.

Statistics

Statistical evaluation often involved ANOVA and Student's t-tests, and only after data were

inspected and recognized to approximate gaussian distributions. When assessing interaction

levels across exposures, a 2-way repeated measures ANOVA was used (exposure number and

cagemate/stranger conditions). When assessing interaction variability changes across

exposures, each dyad was considered independent, and mixed models were used (exposure

number as a within-dyad repeated measure variable, and cagemate/stranger as a between-

subjects variable). In several cases, multiple comparisons had to be taken into account and

corrected for. This includes tests that separately examined each interaction type (5 types

tested), and when testing significance of temporal relationships between interactions (20 types

tested). Significance values are corrected using the Benjamini and Hochberg false discovery rate

procedure [36].

Ethical Note

All protocols were approved by the Institutional Animal Care and Use Committee at University

of Montana (protocol number 036-18NIPSYC-061918). Degus were housed with extra space and

enrichment and no invasive procedures were performed prior to or during the course of the

experiment. Original sample sizes were determined by power analysis performed in G*Power

[37] and were based on preliminary effect sizes obtained from within-dyad behavioral patterns

across conditions in a prior experiment [12]. Following the experiment, degus were transferred

to other experimental protocols; when retired from the laboratory, animals were transferred

either to other labs, adopted locally, or humanely euthanized, with post-mortem tissue often

used for other research.

Acknowledgements

Experiment was supported by NIH grant 1R15MH117611-01A1. We would like to thank Stephen Cooke, Kendra Kuehn, and Kendall Butler for help scoring behavior, Kinsey Webb, Janelle Shamp, and Dani Crandell, for help with the degu colony, and Travis Wheeler and Joshua Starmer for initial review of the manuscript and comments on analysis.

References

1.   Berlyne DE. 1966 Curiosity and Exploration. *Science* **153**, 25–33. (doi:10.1126/science.153.3731.25)

2.   Hughes RN. 1997 Intrinsic exploration in animals: motives and measurement. *Behavioural Processes* **41**, 213–226. (doi:10.1016/S0376-6357(97)00055-7)

3.   Hills TT, Todd PM, Lazer D, Redish AD, Couzin ID. 2015 Exploration versus exploitation in space, mind, and society. *Trends in Cognitive Sciences* **19**, 46–54. (doi:10.1016/j.tics.2014.10.004)

4.   Yurkovic A, Wang O, Basu AC, Kravitz EA. 2006 Learning and memory associated with aggression in Drosophila melanogaster. *Proceedings of the National Academy of Sciences* **103**, 17519–17524. (doi:10.1073/pnas.0608211103)

5.   Lee W, Fu J, Bouwman N, Farago P, Curley JP. 2019 Temporal microstructure of dyadic social behavior during relationship formation in mice. *PLoS ONE* **14**, e0220596. (doi:10.1371/journal.pone.0220596)

6.   Ebensperger LA, Veloso C, Wallem P. 2002 Do female degus communally nest and nurse their pups? *Journal of Ethology* **20**, 143–146. (doi:10.1007/s10164-002-0063-x)

7.   Ebensperger LA *et al.* 2012 Ecological drivers of group living in two populations of the communally rearing rodent, Octodon degus. *Behav. Ecol. Sociobiol. (Print)* **66**, 261–274. (doi:10.1007/s00265-011-1274-3)

8.   Ebensperger LA, Villegas Á, Abades S, Hayes LD. 2014 Mean ecological conditions modulate the effects of group living and communal rearing on offspring production and survival. *Behavioral Ecology* **25**, 862–870. (doi:10.1093/beheco/aru061)

9.   Ebensperger LA, Hurtado M, Soto-Gamboa M, Lacey EileenA, Chang AnnT. 2004 Communal nesting and kinship in degus (Octodon degus). *Naturwissenschaften* **91**. (doi:10.1007/s00114-004-0545-5)

10. Hayes LD, Chesh AS, Castro RA, Tolhuysen LO, Burger JR, Bhattacharjee J, Ebensperger LA. 2009 Fitness consequences of group living in the degu Octodon degus, a plural breeder

rodent with communal care. *Animal Behaviour* **78**, 131–139.
(doi:10.1016/j.anbehav.2009.03.022)

11. Ebensperger LA, Correa LA, León C, Ramírez-Estrada J, Abades S, Villegas Á, Hayes LD. 2016
The modulating role of group stability on fitness effects of group size is different in females
and males of a communally rearing rodent. *J Anim Ecol* **85**, 1502–1515. (doi:10.1111/1365-
2656.12566)

12. Lidhar NK, Thakur A, David A-J, Takehara-Nishiuchi K, Insel N. 2021 Multiple dimensions of
social motivation in adult female degus. *PLoS ONE* **16**, e0250219.
(doi:10.1371/journal.pone.0250219)

13. Insel N, Shambaugh KL, Beery AK. 2020 Female degus show high sociality but no preference
for familiar peers. *Behav. Processes* **174**, 104102. (doi:10.1016/j.beproc.2020.104102)

14. Beery AK, Shambaugh KL. 2021 Comparative Assessment of Familiarity/Novelty Preferences
in Rodents. *Front. Behav. Neurosci.* **15**, 648830. (doi:10.3389/fnbeh.2021.648830)

15. Villavicencio CP, Márquez IN, Quispe R, Vásquez RA. 2009 Familiarity and phenotypic
similarity influence kin discrimination in the social rodent Octodon degus. *Animal Behaviour*
78, 377–384. (doi:10.1016/j.anbehav.2009.04.026)

16. Aureli F, Schaffner CM. 2007 Aggression and conflict management at fusion in spider
monkeys. *Biol. Lett.* **3**, 147–149. (doi:10.1098/rsbl.2007.0041)

17. Dias PAD, Rodríguez Luna E, Canales Espinosa D. 2008 The functions of the "Greeting
Ceremony" among male mantled howlers (Alouatta palliata) on Agaltepec Island, Mexico.
American Journal of Primatology **70**, 621–628. (doi:10.1002/ajp.20535)

18. Kutsukake N, Suetsugu N, Hasegawa T. 2006 Pattern, Distribution, and Function of Greeting
Behavior Among Black-and-White Colobus. *Int J Primatol* **27**, 1271–1291.
(doi:10.1007/s10764-006-9072-x)

19. Smuts BB, Watanabe JM. 1990 Social relationships and ritualized greetings in adult male
baboons (Papio cynocephalus anubis). *International Journal of Primatology* **11**, 147–172.
(doi:10.1007/BF02192786)

20. Whitham JC, Maestripieri D. 2003 Primate Rituals: The Function of Greetings between Male
Guinea Baboons. *Ethology* **109**, 847–859. (doi:10.1046/j.0179-1613.2003.00922.x)

21. Fulk GW. 1976 Notes on the Activity, Reproduction, and Social Behavior of Octodon degus.
Journal of Mammalogy **57**, 495–505. (doi:10.2307/1379298)

22. Kalcher-Sommersguter E, Preuschoft S, Crailsheim K, Franz C. 2011 Social competence of
adult chimpanzees (Pan troglodytes) with severe deprivation history: I. An individual
approach. *Developmental Psychology* **47**, 77–90. (doi:10.1037/a0020783)

23. Turner CH, Davenport RK, Rogers CM. 1969 The Effect of Early Deprivation on the Social
Behavior of Adolescent Chimpanzees. *AJP* **125**, 1531–1536. (doi:10.1176/ajp.125.11.1531)

24. Himmler SM, Himmler BT, Pellis VC, Pellis SM. 2016 Play, variation in play and the
development of socially competent rats. *Behav* **153**, 1103–1137. (doi:10.1163/1568539X-
00003307)

25. Van Den Berg CL, Van Ree JM, Spruijt BM. 1999 Sequential Analysis of Juvenile Isolation-
Induced Decreased Social Behavior in the Adult Rat. *Physiology & Behavior* **67**, 483–488.
(doi:10.1016/S0031-9384(99)00062-1)

26. Keesom SM, Finton CJ, Sell GL, Hurley LM. 2017 Early-Life Social Isolation Influences Mouse
Ultrasonic Vocalizations during Male-Male Social Encounters. *PLoS ONE* **12**, e0169705.
(doi:10.1371/journal.pone.0169705)

27. Varlinskaya E. 2008 Social interactions in adolescent and adult Sprague–Dawley rats: Impact
of social deprivation and test context familiarity. *Behavioural Brain Research* **188**, 398–405.
(doi:10.1016/j.bbr.2007.11.024)

28. Silk J, Cheney D, Seyfarth R. 2013 A practical guide to the study of social relationships: A
Practical Guide to the Study of Social Relationships. *Evol. Anthropol.* **22**, 213–225.
(doi:10.1002/evan.21367)

29. Hummer DL, Jechura TJ, Mahoney MM, Lee TM. 2007 Gonadal hormone effects on
entrained and free-running circadian activity rhythms in the developing diurnal rodent
Octodon degus. *Am J Physiol Regul Integr Comp Physiol* **292**, R586-597.
(doi:10.1152/ajpregu.00043.2006)

30. Mahoney MM, Rossi BV, Hagenauer MH, Lee TM. 2011 Characterization of the Estrous
Cycle in Octodon degus. *Biology of Reproduction* **84**, 664–671.
(doi:10.1095/biolreprod.110.087403)

31. Lee TM. 2004 Octodon degus: a diurnal, social, and long-lived rodent. *ILAR J* **45**, 14–24.
(doi:10.1093/ilar.45.1.14)

32. Friard O, Gamba M. 2016 BORIS⊠: a free, versatile open-source event-logging software for
video/audio coding and live observations. *Methods Ecol Evol* **7**, 1325–1330.
(doi:10.1111/2041-210X.12584)

33. Shinomoto S, Miura K, Koyama S. 2005 A measure of local variation of inter-spike intervals.
Biosystems **79**, 67–72. (doi:10.1016/j.biosystems.2004.09.023)

34. Benjamini Y, Hochberg Y. 1995 Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B
(Methodological)* **57**, 289–300. (doi:10.1111/j.2517-6161.1995.tb02031.x)

35. Faul F, Erdfelder E, Lang A-G, Buchner A. 2007 G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*
Methods **39**, 175–191. (doi:10.3758/BF03193146)

809