# Somatic mutation rates scale with lifespan across mammals

**Authors**

Alex Cagan[1+]*, Adrian Baez-Ortega[1]*, Natalia Brzozowska[1], Federico Abascal[1], Tim H. H. Coorens[1], Mathijs A. Sanders[1,2], Andrew R. J. Lawson[1], Luke M. R. Harvey[1], Shriram Bhosle[1], David Jones[1], Raul E. Alcantara[1], Timothy M. Butler[1], Yvette Hooks[1], Kirsty Roberts[1], Elizabeth Anderson[1], Sharna Lunn[1], Edmund Flach[3], Simon Spiro[3], Inez Januszczak[3,4], Ethan Wrigglesworth[3], Hannah Jenkins[3], Tilly Dallas[3], Nic Masters[3], Matthew W. Perkins[5], Robert Deaville[5], Megan Druce[6,7], Ruzhica Bogeska[6,7], Michael D. Milsom[6,7], Björn Neumann[8,9], Frank Gorman[10], Fernando Constantino-Casas[10], Laura Peachey[10,11], Diana Bochynska[10,12], Ewan St. John Smith[13], Moritz Gerstung[14], Peter J. Campbell[1], Elizabeth P. Murchison[10], Michael R. Stratton[1], Iñigo Martincorena[1+]

* These authors contributed equally

[+] Correspondence to: ac36@sanger.ac.uk (A.C.) and im3@sanger.ac.uk (I.M.)

**Author information**

(1) Cancer, Ageing and Somatic Mutation (CASM), Wellcome Sanger Institute, Hinxton, CB10 1SA, UK

(2) Department of Hematology, Erasmus MC Cancer Institute, 3015 CN Rotterdam, Netherlands

(3) Wildlife Health Services, Zoological Society of London, Regent's Park, London, NW1 4RY, UK

(4) The Natural History Museum, Cromwell Road, London, SW7 5BD, UK

(5) Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY, UK

(6) Division of Experimental Hematology, German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany

(7) Heidelberg Institute for Stem Cell Technology and Experimental Medicine GmbH (HI-STEM), 69120 Heidelberg, Germany

(8) Wellcome Trust-Medical Research Council Cambridge Stem Cell Institute, University of Cambridge, Cambridge, CB2 0AW, UK

(9) Department of Clinical Neuroscience, University of Cambridge, Cambridge, CB2 0QQ, UK

(10) Department of Veterinary Medicine, University of Cambridge, Cambridge, CB3 0ES, UK

(11) Bristol Veterinary School, Faculty of Health Sciences, University of Bristol, Langford, BS40

5DU, UK

(12) (12) Pathology Department, Faculty of Veterinary Medicine, Universitatea de Stiinte

Agricole si Medicina Veterinara, Cluj-Napoca 400372, Romania

(13) Department of Pharmacology, University of Cambridge, Cambridge CB2 1PD, UK

(14) European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Hinxton, CB10 1SD, UK

**Abstract**

The rates and patterns of somatic mutation in normal tissues are largely unknown outside of

humans. Comparative analyses can shed light on the diversity of mutagenesis across species and

on long-standing hypotheses regarding the evolution of somatic mutation rates and their role in

cancer and ageing. Here, we used whole-genome sequencing of 208 intestinal crypts from 56

individuals to study the landscape of somatic mutation across 16 mammalian species. We found

somatic mutagenesis to be dominated by seemingly endogenous mutational processes in all species,

including 5-methylcytosine deamination and oxidative damage. With some differences, mutational

signatures in other species resembled those described in humans, although the relative contribution

of each signature varied across species. Remarkably, the somatic mutation rate per year varied

greatly across species and exhibited a strong inverse relationship with species lifespan, with no

other life-history trait studied displaying a comparable association. Despite widely different life

histories among the species surveyed, including ~30-fold variation in lifespan and ~40,000-fold

variation in body mass, the somatic mutation burden at the end of lifespan varied only by a factor

of ~3. These data unveil common mutational processes across mammals and suggest that somatic

mutation rates are evolutionarily constrained and may be a contributing factor in ageing.

**Introduction**

Somatic mutations accumulate in healthy cells throughout life. They underpin cancer

development[1,2] and, for decades, have been speculated to contribute to ageing[3–5]. Directly studying

somatic mutations in normal tissues has been challenging due to the difficulty of detecting

mutations present in single cells or small clones in a tissue. Only recent technological

developments, such as *in vitro* expansion of single cells into colonies[6,7], microdissection of

histological units[8,9] or single-cell or single-molecule sequencing[10–12], are beginning to enable the

study of somatic mutation in normal tissues.

Over the last few years, studies in humans have started to provide a detailed understanding of

somatic mutation rates and the contribution of endogenous and exogenous mutational processes

across normal tissues[6–8,13,14]. These studies are also revealing how, as we age, some human tissues

are colonised by mutant cells carrying cancer-driving mutations, and how this clonal composition

changes with age and disease. With the exception of some initial studies, far less is known about

somatic mutation in other species[15–20]. Yet, comparative analyses of somatic mutagenesis would

shed light on the diversity of mutagenic processes across species, and on long-standing questions

regarding the evolution of somatic mutation rates and their role in cancer and ageing.

A decades-long hypothesis on the evolution of somatic mutation rates pertains to the relationship

between body mass and cancer risk. Some models predict that the risk of cancer should increase

proportionally to the number of cells at risk of transformation. However, there appears to be no

correlation between body mass and cancer risk across species[21–23]. This observation, known as

Peto's paradox, suggests that the evolution of larger body sizes likely requires the evolution of

stronger cancer suppression mechanisms[23,24]. Whether evolutionary reduction of cancer risk across

species is partly achieved by a reduction of somatic mutation rates remains unknown[25].

A second long-standing hypothesis on the evolution of somatic mutation rates relates to the

proposed role of somatic mutations in ageing. Multiple forms of molecular damage, including

somatic mutations, telomere attrition, epigenetic drift or loss of proteostasis, have been proposed

to contribute to ageing, but their causal roles and relative contributions remain debated[26,27].

Evolutionary theory predicts that species will evolve protection or repair mechanisms against life-

threatening damage to minimise death from intrinsic causes, but that selection is too weak to delay

ageing far beyond the typical life expectancy of an organism in the wild[28–32] (**Supplementary Note**

**1**). If somatic mutations contribute to ageing, theory predicts that somatic mutation rates may

inversely correlate with lifespan across species[30,33]. This long-standing prediction has remained

largely untested due to the difficulty of measuring somatic mutation rates across species.

**Detection of somatic mutations across species**

The study of somatic mutations with standard whole-genome sequencing requires isolating clonal

groups of cells recently derived from a single cell[6–8]. To study somatic mutations across a diverse

set of mammals, we isolated 208 individual intestinal crypts from 56 individuals across 16 species

with a wide range of lifespans and body sizes: black-and-white colobus monkey, cat, cow, dog,

ferret, giraffe, harbour porpoise, horse, human, lion, mouse, naked mole-rat, rabbit, rat, ring-tailed

lemur, and tiger (**Supplementary Table 1**). We chose intestinal crypts for several reasons. First,

they are histologically identifiable units that line the epithelium of the colon and small intestine

and are amenable to laser microdissection. Second, human studies have confirmed that individual

crypts become clonally derived from a single stem cell and show a linear accumulation of mutations

with age, enabling the estimation of somatic mutation rates through genome sequencing of single

crypts[8]. Third, in most human crypts, the majority of somatic mutations are caused by endogenous

mutational processes common to other tissues, rather than by environmental mutagens[8,12].

A colon sample was collected from each individual, with the exception of a ferret from which only

a small intestine sample was available. This sample was included because results in humans have

shown the mutation rate of colorectal and small intestine epithelium to be similar[7,14] (**Extended**

**Data Figure 1**). We then used laser microdissection on histological sections to isolate individual

crypts for whole-genome sequencing with a low-input library preparation method[34] (**Fig. 1a,**

**Extended Data Figure 2, Supplementary Table 2**), with the exception of human crypts, for which

sequencing data were obtained from a previous study[8]. A bioinformatic pipeline was developed to

call somatic mutations robustly in all these species despite variable quality of their genome

assemblies (**Methods**). The distribution of variant allele fractions (VAFs) of the mutations detected

in each crypt confirmed that crypts are clonal units in all species, enabling the study of somatic

mutation rates and signatures (**Extended Data Figure 3**).

We found substantial variation in the number of somatic single-base substitutions (SBS) across
species and across individuals within each species (**Fig. 1b**). For five species with samples from
multiple individuals, linear regression confirmed a clear accumulation of somatic mutations with
age (**Fig. 1c, Extended Data Figure 4, Supplementary Table 3**). All linear regressions were also
consistent with a non-significant intercept. This resembles observations in humans[14] and suggests
that the time required for a single stem cell to drift to fixation within a crypt is a small fraction of
the lifespan of a species. This facilitates the estimation of somatic mutation rates across species by
dividing the number of mutations in a crypt by the age of the individual (**Supplementary Table
4**). The number of somatic insertions and deletions (indels) was consistently lower than that of SBS
in all crypts (**Fig. 1b**), in agreement with previous findings in humans[8].

**Similar mutational signatures across mammals**

Somatic mutations can be caused by different mutational processes, including multiple forms of
DNA damage and repair. Different processes cause characteristic frequencies of base substitution
types and indels at different sequence contexts, often referred to as mutational signatures, which
can be inferred from mutation data[35,36]. Across species, the mutational spectra showed clear
similarities, with a dominance of cytosine-to-thymine (C>T) substitutions at CpG sites, as observed
in human colon, but with considerable variation in the frequency of other substitution types (**Fig.
2a**). To quantify the contribution of different mutational processes to the observed spectra, we
applied mutational signature decomposition[8,35]. We used a Bayesian model to infer mutational
signatures *de novo*, while accounting for differences in genome sequence composition across
species, and using the COSMIC human signature SBS1 (C>T substitutions at CpG sites) as a fixed
prior to ensure its complete deconvolution[37] (**Methods**). This approach identified two signatures
beyond SBS1, labelled SBSB and SBSC, which resemble COSMIC human signatures SBS5 and
SBS18, respectively (cosine similarities 0.93 and 0.91) (**Fig. 2b**).

This analysis suggests that the same three signatures that dominate somatic mutagenesis in the
human colon are dominant in other mammals: SBS1, which is believed to result from the
spontaneous deamination of 5-methylcytosine[8,38]; SBSB/5, a common signature across human
tissues that may result from endogenous damage and repair[12,39]; and SBSC/18, dominated by C>A
substitutions and attributed to oxidative DNA damage[36]. Signature SBSC contains a minor

component of T>A substitutions (resembling COSMIC SBS34), which appear to be the result of

DNA polymerase slippage at the boundaries between adjacent adenine and thymine homopolymer

tracts but that could also reflect assembly errors at those sites[39]. While all species surveyed shared

the three mutational signatures, their contributions varied substantially across species (**Fig. 2c**).

SBSC was particularly prominent in mouse and ferret, and the ratio of SBS1 to SBSB (SBS5)

varied from approximately 1.2 in rat or rabbit to 6.4 in tiger. In several species with data from

multiple individuals, separate linear regressions for each signature confirmed that mutations from

all three signatures accumulate with age (**Fig. 2d, Extended Data Figure 5**).

Although signature deconvolution identified three signatures active across species, we noticed

some differences in the mutational profile of signature SBSB among species. To explore this

further, we inferred independent versions of SBSB from each species, while accounting for

differences in genome sequence composition (**Methods**). This revealed inter-species variability in

the mutational profile of this signature, particularly in the C>T component (**Extended Data Figure**

**6**). Species-specific versions of SBSB showed different similarities to the related human signatures

SBS5 and SBS40. For example, SBSB inferred from the human data showed a stronger similarity

with the reference SBS5 human signature (cosine similarities with SBS5 and SBS40: 0.93 and

0.84), whereas SBSB from rabbit more closely resembled the reference human SBS40 signature

(0.87 and 0.91). These observations are consistent with the hypothesis that SBS5 and SBS40 result

from a combination of correlated mutational processes, with some variation across human

tissues[12,39] and across species.

Analysis of the indel mutational spectra revealed a dominance of the human indel signatures ID1

and ID2, characterised by single-nucleotide insertions and deletions at A/T homopolymers, and

probably caused by strand slippage during DNA replication[36] (**Extended Data Figure 7**). The ratio

of insertions (ID1) to deletions (ID2) appears to vary across species, possibly reflecting a

differential propensity for slippage of the template and nascent DNA strands[36]. In addition, the

indel spectra suggest a potential contribution of signature ID9 (whose aetiology remains unknown)

to human, colobus, cow, giraffe and rabbit. Analysis of indels longer than 1 base pair (bp) also

suggested the presence of a signature of 4-bp insertions at tetrameric repeats, particularly prevalent

in mouse and tiger; a pattern of ≥5-bp insertions at repeats in colobus; and a pattern of ≥5-bp

deletions at repeats prominent in rabbit, which resembles ID8 (a signature possibly caused by

double-strand break repair via non-homologous end joining[36]) (**Extended Data Figure 7**).

**Other mutational processes and selection**

The apparent lack of additional mutational signatures is noteworthy. A previous study of 445

colorectal crypts from 42 human donors found many crypts to be affected by a novel signature that

was later attributed to colibactin, a genotoxin produced by *pks+* strains of *Escherichia coli*[8,40,41].

Analysing the original human data and our non-human data with the same methodology, we found

evidence of colibactin mutagenesis in 21% of human crypts but only uncertain evidence of

colibactin in one non-human crypt (0.6%) (**Extended Data Figure 8, Methods**). This revealed a

significant depletion of colibactin mutagenesis in the non-human crypts studied (Fisher's exact

tests, $P=7\times10^{-14}$). The apparent difference in colibactin mutagenesis observed between species, or

between the cohorts studied, might result from a different prevalence of *pks+ E. coli* strains[42] or a

different expression of colibactin by *pks+ E.coli* across species[43]. Finally, we also searched for

evidence of APOBEC signatures (SBS2/13), which have been reported in a small number of human

crypts and are believed to be caused by APOBEC DNA-editing cytidine deaminases. We detected

APOBEC signatures in 2% (n = 9) of human crypts and only uncertain evidence in one non-human

crypt ($P=0.30$).

Beyond single-base substitutions and indels, crypts from the eight species with chromosome-level

genome assemblies were inspected for large-scale copy number changes ($\geq$1 megabase, Mb)

(**Methods**). Studies in humans have found large-scale copy number changes to be relatively rare

in normal tissues, including colorectal epithelium[8]. Consistent with these results, we only identified

four large copy number changes across the 162 crypts included in this analysis: two megabase-

scale deletions in two crypts from the same cow, the loss of a chromosome X in a female mouse

crypt, and a 52-Mb segment with copy-neutral loss of heterozygosity in a human crypt (**Extended**

**Data Figure 9, Methods**). These results suggest that large-scale somatic copy number changes in

normal tissues are also rare in other mammalian species.

Previous analyses in humans have shown that most somatic mutations in colorectal crypts

accumulate neutrally, without clear evidence of negative selection against non-synonymous

mutations and with a low frequency of positively selected cancer-driver mutations[8]. To study somatic selection in our data, we calculated the exome-wide ratio of non-synonymous to synonymous substitution rates (dN/dS) in each of the 12 species with available genome annotation. To do so and to detect genes under positive selection, while accounting for the effects of trinucleotide sequence context and mutation rate variation across genes, we used the dNdScv model[44] (**Methods**). Although the limited number of coding somatic mutations observed in most species prevented an in-depth analysis of selection, exome-wide dN/dS ratios for somatic substitutions were not significantly different from unity in any species, in line with previous findings in humans[8] (**Extended Data Figure 10**). Gene-level analysis did not find genes under significant positive selection in any species, although larger studies are likely to identify rare cancer-driver mutations[8].

**Somatic mutation rates and life-history traits**

Whereas similar mutational processes operate across the species surveyed, the mutation rate per genome per year varied widely. Across the 15 species with age information, we found that substitution rates per genome ranged from 47 SBS/year in humans to 796 SBS/year in mice, and indel rates from 2.5 to 158 indels/year, respectively (**Fig. 3a, Supplementary Table 4, Methods**).

To explore the relationship between somatic mutation rates, lifespan and other life-history traits, we first estimated the lifespan of each species using survival curves. We used a large collection of mortality data from animals in zoos to minimise the impact of extrinsic mortality (**Extended Data Figure 11**). We defined lifespan as the age at which 80% of individuals reaching adulthood have died, to reduce the impact of outliers and variable cohort sizes that affect maximum lifespan estimates[45,46] (**Methods**). Remarkably, we found a tight anticorrelation between somatic mutation rates per year and lifespan across species (**Fig. 3b**). A log-log allometric regression yielded a strong linear anticorrelation between mutation rate per year and lifespan (fraction of variance explained, FVE=0.85, $P$=1x10$^{-6}$), with a slope close to and non-significantly different from -1. This supports a simple model in which somatic mutation rates per year are inversely proportional to the lifespan of a species (rate ∝ 1/lifespan), such that the number of somatic mutations per cell at the end of the lifespan (the end-of-lifespan burden, ELB) is similar in all species (**Fig. 3a-c**).

To further study the relationship between somatic mutation rates and life-history variables, we used
linear mixed-effects regression models. These models account for the hierarchical structure of the
data (with multiple crypts per individual and multiple individuals per species), as well as the
heteroscedasticity of somatic mutation rate estimates across species (**Methods**). Using these
models, we estimated that the inverse of lifespan explained 82% of the inter-species variance in
somatic substitution rates (*rate = k/lifespan*) ($P=2.9\times10^{-9}$; **Fig. 3c**), with the slope of this regression
(*k*) representing the mean estimated ELB across species (3206.4 substitutions per genome per crypt,
95% confidence interval 2683.9–3728.9). Strikingly, despite uncertainty in the estimates of both
somatic mutation rates and lifespans, and despite the diverse life histories of the species surveyed,
including ~30-fold variation in lifespan and ~40,000-fold variation in body mass, the estimated
mutation load per cell at the end of lifespan varied by only ~3-fold across species (**Table 1**).
Analogous results were obtained repeating the analysis with protein-coding mutation rate
estimates, which may be a better proxy for the functional impact of somatic mutations (85% of
variance explained, ELB: 31 coding substitutions per crypt) (**Extended Data Figure 12**,
**Methods**).
269
We next explored the association between somatic mutation rates and adult body mass, which is
known to be a common confounder in correlations involving lifespan[47,48]. An anticorrelation
between somatic mutation rates and body mass may be expected if the modulation of cancer risk
across species of vastly different sizes has been a major factor in the evolution of somatic mutation
rates. We observed that log-transformed adult body mass was less strongly associated with the
somatic substitution rate (allometric regression FVE=0.21, **Fig. 3d**; linear mixed-effect regression
FVE=0.44, **Fig. 3e**). Given that lifespan is correlated with body mass, we performed two tests to
assess whether body mass explained any variation in somatic mutation rates not explained by
lifespan. First, including both lifespan and log-transformed adult body mass in the regression model
suggested that body mass does not explain a significant amount of variance in somatic mutation
rates across species after accounting for the effect of lifespan (likelihood ratio tests: *P*=0.16 for
body mass on a model with lifespan, $P<10^{-4}$ for lifespan on a model with body mass; **Fig. 3f,**
**Methods**). Second, partial correlation analyses using allometric regressions further confirmed that
the association between somatic mutation rates and lifespan is unlikely to be mediated by the effect
of body mass on both variables (lifespan residuals: $P=3.2\times10^{-6}$, FVE=0.82, **Fig. 3b**; body mass
residuals: *P*=0.39, FVE=0.06, **Fig. 3d**; **Methods**).

The fact that the variation in somatic mutation rates across species appears to be dominated by

lifespan rather than body size is also apparent when looking at particularly informative species.

Giraffe and naked mole-rat, for instance, have similar somatic mutation rates (99 and 93

substitutions/year), in line with their similar lifespans (80th percentiles: 24 and 25 years), despite

a ~23,000-fold difference in adult body mass (**Fig. 3d,e**). Cows, giraffes and tigers weigh much

more than an average human, and yet have somatic mutation rates several fold higher, in line with

expectation from their lifespans but not their body mass. Altogether, the weak correlation between

body mass and somatic mutation rates after correction for lifespan suggests that the evolution of

larger body sizes may have relied on alternative or additional strategies to limit cancer risk, as has

been speculated (**Supplementary Note 2**)[23,49–51]. Of note, the low somatic mutation rate of naked

mole-rats, unusual for their body mass but in line with their long lifespan (**Fig. 3d,e**), might

contribute to the exceptionally low cancer incidence rates of this species[52,53].

We found similar results for other life-history variables that have been proposed to correlate with

lifespan, namely basal metabolic rate (BMR) and litter size[54,55] (**Fig. 3f**). With the caveat that

estimates for these variables vary in quality, they showed weaker correlations with the somatic

mutation rate as single predictors, and small non-significant increases in explanatory power when

considered together with lifespan (likelihood ratio tests, $P$=0.92 for litter size, P=0.083 for log-

BMR, $P$=0.79 for allometric BMR residuals; **Fig. 3f, Methods**). We note that the results above are

robust to the use of alternative measures of the somatic mutation rate, including the rate per exome

or mutations/Mb (**Extended Data Figure 12**, **Methods**); alternative estimates of lifespan,

including maximum lifespan (**Extended Data Figure 13**, **Methods**); alternative regression

models, including a Bayesian hierarchical model and a phylogenetic generalised least-squares

regression, which accounts for the effect of phylogenetic relationships (**Extended Data Figure**

**14a,b**, **Methods**); and bootstrapping analyses at the level of individuals or species (**Extended Data**

**Figure 14c**, **Methods**).

**Mutational processes and lifespan**

To explore whether a single biological process could drive the association between somatic

mutation rates and lifespan, we analysed each mutational signature separately. SBS1, SBSB/5 and

SBSC/18 are believed to result from different forms of DNA damage and are expected to be subject to different DNA repair pathways[12,39,56]. They also appear to differ in their association with the rate of cell division in humans, with SBS1 being more common in fast-proliferating tissues, such as colon and embryonic or foetal tissues, and SBS5 dominating in post-mitotic cells in the absence of cell division[7,12,14]. Overall, we found clear anticorrelations between mutation rates per year and lifespan for the three SBS signatures and for indels, suggesting that a single biological process or DNA repair pathway is unlikely to be responsible for this association (**Fig. 4**). The total mutation burden also appears to show a closer fit with lifespan than individual mutational processes, as measured by the range of end-of-lifespan-burden (ELB) for each process across species (**Fig. 4**). This may be expected if the observed anticorrelation were the result of evolutionary pressure on somatic mutation rates.

DNA damage and somatic mutations in the mitochondrial genome have also attracted considerable interest in the ageing field[57,58]. Our whole-genome sequencing of individual crypts provided high coverage of the mitochondrial genome, ranging from 2,188 to 29,691-fold. Normalised against the nuclear coverage, these data suggest that colorectal crypts contained on the order of ~100–2,000 mitochondrial genomes per cell (**Extended Data Figure 15**). Using a mutation calling algorithm sensitive to low-frequency variants, we found a total of 261 mitochondrial mutations across 199 crypts (**Methods**). The mutational spectra across species appeared broadly consistent with that observed in humans, with a dominance of C>T and A>G substitutions believed to result from mtDNA replication errors rather than DNA damage[59] (**Extended Data Figure 16**). While the low number of mitochondrial mutations detected per species precludes a detailed analysis, the estimated number of somatic mutations per copy of mtDNA also appears to show a significant anticorrelation with lifespan. Across species, we obtained an average of 0.23 detectable mutations per copy of the mitochondrial genome by the end of lifespan (**Fig. 4**, **Methods**), a considerable burden given the coding-sequence density and functional relevance of the mitochondrial genome.

**Discussion**

Using whole-genome sequencing of 208 colorectal crypts from 56 individuals, we provide insights into the somatic mutational landscape of 16 mammalian species. Despite their different diets and life histories, we found remarkable similarities in their mutational spectra. Three main mutational

signatures explain the spectra across species, albeit with varying contributions and subtle variations

in the profile of signature SBSB. These results suggest that, at least in the colorectal epithelium, a

conserved set of mutational processes dominate somatic mutagenesis across mammals.

The most striking finding of this study is the inverse scaling of somatic mutation rates with lifespan,

a long-standing prediction of the somatic mutation theory of ageing[4,30,33]. Considering evolutionary

and mechanistic models of ageing together provides a framework for discussing the possible

implications of these results for ageing (**Supplementary Note 1**). Jointly, these models predict

ageing to be multifactorial, with multiple forms of molecular and cellular damage contributing to

organismal ageing due to evolutionary limits to selection acting on the rates of these processes.

The inverse scaling of somatic mutation rates and lifespan is consistent with somatic mutations

contributing to ageing and with somatic mutation rates being evolutionarily constrained, although

we discuss alternative explanations below. This interpretation is also supported by studies reporting

more efficient DNA repair in longer-lived species[60–62]. Somatic mutations could contribute to

ageing in different ways. Traditionally, they have been proposed to contribute to ageing through

deleterious effects on cellular fitness[4,63], but recent findings question this assumption

(**Supplementary Note 1**). Instead, the discovery of widespread clonal expansions in ageing human

tissues[13,64–66] raises the possibility that some somatic mutations contribute to ageing by increasing

the fitness of mutant cells at a cost to the organism[63,67,68]. Recent examples include the possible

links between clonal haematopoiesis and cardiovascular disease[69], between mutations in liver

disease and insulin resistance [70], and between driver mutations in cavernomas and brain

haemorrhages[63,67,68]. Detailed studies on the extent and impact of somatic mutations and clonal

expansions in age-related diseases may help clarify the precise role, if any, of somatic mutations

in ageing. Even if clear causal links between somatic mutations and ageing are established, ageing

is likely to be multifactorial. Other forms of molecular damage involved in ageing could be

expected to display similar anticorrelations with lifespan and, indeed, such anticorrelations have

been reported for telomere shortening and protein turnover[71,72].

Alternative non-causal explanations for the observed anticorrelation between somatic mutation

rates and lifespan need to be considered. One alternative explanation is that cell division rates scale

with lifespan and explain the observed somatic mutation rates. Available estimates of cell division

rates, though imperfect and limited to a few species, do not readily support this argument

(**Methods**). More importantly, studies in humans have shown that cell division rates are not a major

determinant of somatic mutation rates across human tissues[7,12]. Another alternative explanation for

the observed anticorrelation might be that selection acts to reduce germline mutation rates in

species with longer reproductive spans, in turn causing an anticorrelation of somatic mutation rates

and lifespan. While selective pressure on germline mutation rates could influence somatic mutation

rates, it is unlikely that germline mutation rates determine somatic mutation rates: somatic mutation

rates in humans are 10–20 times higher than germline mutation rates, show variability across cell

types, and are influenced by additional mutational processes[12,14]. Overall, the strong scaling of

somatic mutation rates with lifespan across mammals, despite variable contributions of different

mutational processes, suggests that somatic mutation rates themselves have been evolutionarily

constrained, possibly through selection on multiple DNA repair pathways. Alternative explanations

need to be able to explain the strength of the scaling despite these differences.

Altogether, this study provides an unprecedented description of somatic mutation across mammals,

identifying common and variable features and shedding light on long-standing hypotheses. Scaled

across the tree of life and across tissues, in species with vastly different physiologies, life histories,

genome compositions and mutagenic exposures, similar studies promise to transform our

understanding of somatic mutation and its impact on evolution, ageing, and disease.

**References**

1. Nowell, P. C. The clonal evolution of tumor cell populations. *Science* **194**, 23–28 (1976).

2. Stratton, M. R., Campbell, P. J. & Futreal, P. A. The cancer genome. *Nature* **458**, 719–724 (2009).

3. Szilard, L. ON THE NATURE OF THE AGING PROCESS. *Proc. Natl. Acad. Sci. U. S. A.* **45**, 30–

45 (1959).

4. Morley, A. A. The somatic mutation theory of ageing. *Mutat. Res.* **338**, 19–23 (1995).

5. Vijg, J. & Dong, X. Pathogenic Mechanisms of Somatic Mutation and Genome Mosaicism in Aging.

*Cell* **182**, 12–23 (2020).

6. Welch, J. S. *et al.* The origin and evolution of mutations in acute myeloid leukemia. *Cell* **150**, 264–

278 (2012).

7.    Blokzijl, F. *et al.* Tissue-specific mutation accumulation in human adult stem cells during life. *Nature*

**538**, 260–264 (2016).

8.    Lee-Six, H. *et al.* The landscape of somatic mutation in normal colorectal epithelial cells. *Nature*

**574**, 532–537 (2019).

9.    Moore, L. *et al.* The mutational landscape of normal human endometrial epithelium. *Nature* **580**,

640–646 (2020).

10.  Lodato, M. A. *et al.* Aging and neurodegeneration are associated with increased mutations in single

human neurons. *Science* **359**, 555–559 (2018).

11.  Zhang, L. *et al.* Single-cell whole-genome sequencing reveals the functional landscape of somatic

mutations in B lymphocytes across the human lifespan. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 9014–

9019 (2019).

12.  Abascal, F. *et al.* Somatic mutation landscapes at single-molecule resolution. *Nature* (2021)

doi:10.1038/s41586-021-03477-4.

13.  Martincorena, I. *et al.* Tumor evolution. High burden and pervasive positive selection of somatic

mutations in normal human skin. *Science* **348**, 880–886 (2015).

14.  Moore, L. *et al.* The mutational landscape of human somatic and germline cells. *Cold Spring Harbor*

*Laboratory* 2020.11.25.398172 (2020) doi:10.1101/2020.11.25.398172.

15.  Garcia, A. M. *et al.* Age- and temperature-dependent somatic mutation accumulation in Drosophila

melanogaster. *PLoS Genet.* **6**, e1000950 (2010).

16.  Behjati, S. *et al.* Genome sequencing of normal cells reveals developmental lineages and mutational

processes. *Nature* **513**, 422–425 (2014).

17.  Milholland, B. *et al.* Differences between germline and somatic mutation rates in humans and mice.

*Nat. Commun.* **8**, 15183 (2017).

18.  Schmid-Siegert, E. *et al.* Low number of fixed somatic mutations in a long-lived oak tree. *Nat Plants*

**3**, 926–929 (2017).

19.  Orr, A. J. *et al.* A phylogenomic approach reveals a low somatic mutation rate in a long-lived plant.

*Proc. Biol. Sci.* **287**, 20192364 (2020).

20.   López, E. H. & Palumbi, S. R. Somatic Mutations and Genome Stability Maintenance in Clonal

Coral Colonies. *Mol. Biol. Evol.* **37**, 828–838 (2020).

21.   Peto, R., Roe, F. J., Lee, P. N., Levy, L. & Clack, J. Cancer and ageing in mice and men. *Br. J.*

*Cancer* **32**, 411–426 (1975).

22.   Caulin, A. F. & Maley, C. C. Peto's Paradox: evolution's prescription for cancer prevention. *Trends*

*Ecol. Evol.* **26**, 175–182 (2011).

23.   Tollis, M., Boddy, A. M. & Maley, C. C. Peto's Paradox: how has evolution solved the problem of

cancer prevention? *BMC Biol.* **15**, 60 (2017).

24.   Peto, R. Epidemiology, multistage models, and short-term mutagenicity tests. *Int. J. Epidemiol.* **45**,

621–637 (2016).

25.   Caulin, A. F., Graham, T. A., Wang, L.-S. & Maley, C. C. Solutions to Peto's paradox revealed by

mathematical modelling and cross-species cancer gene analysis. *Philos. Trans. R. Soc. Lond. B Biol.*

*Sci.* **370**, (2015).

26.   López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging.

*Cell* **153**, 1194–1217 (2013).

27.   Schumacher, B., Pothof, J., Vijg, J. & Hoeijmakers, J. H. J. The central role of DNA damage in the

ageing process. *Nature* **592**, 695–703 (2021).

28.   Medawar, P. B. UNSOLVED problem of biology. *Med. J. Aust.* **1**, 854–855 (1953).

29.   Williams, G. C. Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution* **11**, 398–

411 (1957).

30.   Kirkwood, T. B. & Holliday, R. The evolution of ageing and longevity. *Proc. R. Soc. Lond. B Biol.*

*Sci.* **205**, 531–546 (1979).

31.   Partridge, L. & Barton, N. H. Optimality, mutation and the evolution of ageing. *Nature* **362**, 305–311

(1993).

32.   Hughes, K. A. & Reynolds, R. M. Evolutionary and mechanistic theories of aging. *Annu. Rev.*

*Entomol.* **50**, 421–445 (2005).

33.  Burnet, M. Intrinsic mutagenesis. (1974) doi:10.1007/978-94-011-6606-5.

34.  Ellis, P. *et al.* Reliable detection of somatic mutations in solid tissues by laser-capture

microdissection and low-input DNA sequencing. *Nat. Protoc.* **16**, 841–871 (2021).

35.  Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415–421

(2013).

36.  Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature* **578**, 94–101

(2020).

37.  Gori, K. & Baez-Ortega, A. sigfit: flexible Bayesian inference of mutational signatures. *bioRxiv*

372896 (2020) doi:10.1101/372896.

38.  Lindahl, T. & Nyberg, B. Heat-induced deamination of cytosine residues in deoxyribonucleic acid.

*Biochemistry* **13**, 3405–3410 (1974).

39.  Zou, X. *et al.* A systematic CRISPR screen defines mutational mechanisms underpinning signatures

caused by replication errors and endogenous DNA damage. *Nat Cancer* **2**, 643–657 (2021).

40.  Wilson, M. R. *et al.* The human gut bacterial genotoxin colibactin alkylates DNA. *Science* **363**,

(2019).

41.  Pleguezuelos-Manzano, C. *et al.* Mutational signature in colorectal cancer caused by genotoxic pks+

E. coli. *Nature* **580**, 269–273 (2020).

42.  Smati, M. *et al.* Quantitative analysis of commensal Escherichia coli populations reveals host-

specific enterotypes at the intra-species level. *Microbiologyopen* **4**, 604–615 (2015).

43.  Oliero, M. *et al.* Oligosaccharides increase the genotoxic effect of colibactin produced by pks+

Escherichia coli strains. *BMC Cancer* **21**, 172 (2021).

44.  Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* **171**,

1029–1041.e21 (2017).

45.  Moorad, J. A., Promislow, D. E. L., Flesness, N. & Miller, R. A. A comparative assessment of

univariate longevity measures using zoological animal records. *Aging Cell* **11**, 940–948 (2012).

46. Tidière, M. *et al.* Comparative analyses of longevity and senescence reveal variable survival benefits

of living in zoos across mammals. *Sci. Rep.* **6**, 36361 (2016).

47. Speakman, J. R. Correlations between physiology and lifespan--two widely ignored problems with

comparative studies. *Aging Cell* **4**, 167–175 (2005).

48. de Magalhães, J. P., Costa, J. & Church, G. M. An analysis of the relationship between metabolism,

developmental schedules, and longevity using phylogenetic independent contrasts. *J. Gerontol. A*

*Biol. Sci. Med. Sci.* **62**, 149–160 (2007).

49. Risques, R. A. & Promislow, D. E. L. All's well that ends well: why large species have short

telomeres. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373**, (2018).

50. Tollis, M., Schneider-Utaka, A. K. & Maley, C. C. The Evolution of Human Cancer Gene

Duplications across Mammals. *Mol. Biol. Evol.* **37**, 2875–2886 (2020).

51. Vazquez, J. M. & Lynch, V. J. Pervasive duplication of tumor suppressors in Afrotherians during the

evolution of large bodies and reduced cancer risk. *Elife* **10**, (2021).

52. Buffenstein, R. Negligible senescence in the longest living rodent, the naked mole-rat: insights from

a successfully aging species. *J. Comp. Physiol. B* **178**, 439–445 (2008).

53. Smith, E. S. J., Schuhmacher, L.-N. & Husson, Z. The naked mole-rat as an animal model in

biomedical research: current perspectives. *Open Access Anim. Physiol.* 137 (2015)

doi:10.2147/oaap.s50376.

54. Millar, J. S. & Zammuto, R. M. Life Histories of Mammals: An Analysis of Life Tables. *Ecology* **64**,

631–635 (1983).

55. Speakman, J. R. Body size, energy metabolism and lifespan. *J. Exp. Biol.* **208**, 1717–1730 (2005).

56. Sanders, M. A. *et al.* Life without mismatch repair. *bioRxiv* 2021.04.14.437578 (2021)

doi:10.1101/2021.04.14.437578.

57. Lane, R. K., Hilsabeck, T. & Rea, S. L. The role of mitochondrial dysfunction in age-related

diseases. *Biochim. Biophys. Acta* **1847**, 1387–1400 (2015).

58. Kauppila, T. E. S., Kauppila, J. H. K. & Larsson, N.-G. Mammalian Mitochondria and Aging: An

Update. *Cell Metab.* **25**, 57–71 (2017).

59.   Ju, Y. S. *et al.* Origins and functional consequences of somatic mitochondrial DNA mutations in

human cancer. *Elife* **3**, (2014).

60.   Hall, K. Y., Hart, R. W., Benirschke, A. K. & Walford, R. L. Correlation between ultraviolet-induced

DNA repair in primate lymphocytes and fibroblasts and species maximum achievable life span.

*Mech. Ageing Dev.* **24**, 163–173 (1984).

61.   Tian, X. *et al.* SIRT6 Is Responsible for More Efficient DNA Double-Strand Break Repair in Long-

Lived Species. *Cell* **177**, 622–638.e22 (2019).

62.   Zhang, L. *et al.* Maintenance of genome sequence integrity in long- and short-lived rodent species.

*Sci Adv* **7**, eabj3284 (2021).

63.   Smith, J. M. Review Lectures on Senescence - I. The causes of ageing. *Proceedings of the Royal*

*Society of London. Series B. Biological Sciences* **157**, 115–127 (1962).

64.   Jaiswal, S. *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N. Engl. J.*

*Med.* **371**, 2488–2498 (2014).

65.   Martincorena, I. *et al.* Somatic mutant clones colonize the human esophagus with age. *Science* **362**,

911–917 (2018).

66.   Yokoyama, A. *et al.* Age-related remodelling of oesophageal epithelia by mutated cancer drivers.

*Nature* **565**, 312–317 (2019).

67.   Jaiswal, S. & Ebert, B. L. Clonal hematopoiesis in human aging and disease. *Science* **366**, (2019).

68.   Ren, A. A. *et al.* PIK3CA and CCM mutations fuel cavernomas through a cancer-like mechanism.

*Nature* **594**, 271–276 (2021).

69.   Jaiswal, S. & Libby, P. Clonal haematopoiesis: connecting ageing and inflammation in

cardiovascular disease. *Nat. Rev. Cardiol.* **17**, 137–144 (2020).

70.   Ng, S. W. K. *et al.* Convergent somatic mutations in metabolism genes in chronic liver disease.

*Nature* **598**, 473–478 (2021).

71.   Swovick, K. *et al.* Interspecies Differences in Proteome Turnover Kinetics Are Correlated With Life

542   Spans and Energetic Demands. *Mol. Cell. Proteomics* **20**, 100041 (2021).

543  72. Whittemore, K., Vera, E., Martínez-Nevado, E., Sanpera, C. & Blasco, M. A. Telomere shortening

544   rate predicts species life span. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 15122–15127 (2019).

545  73. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-*

546   *bio.GN]* (2013).

547  74. Jones, D. *et al.* cgpCaVEManWrapper: Simple Execution of CaVEMan in Order to Detect Somatic

548   Single Nucleotide Variants in NGS Data. *Curr. Protoc. Bioinformatics* **56**, 15.10.1–15.10.18 (2016).

549  75. Raine, K. M. *et al.* cgpPindel: Identifying Somatically Acquired Insertion and Deletion Events from

550   Paired End Sequencing. *Curr. Protoc. Bioinformatics* **52**, 15.7.1–15.7.12 (2015).

551  76. Gerstung, M., Papaemmanuil, E. & Campbell, P. J. Subclonal variant calling with multiple samples

552   and prior knowledge. *Bioinformatics* **30**, 1198–1204 (2014).

553  77. Gerstung, M. *et al.* Reliable detection of subclonal single-nucleotide variants in tumour cell

554   populations. *Nat. Commun.* **3**, 811 (2012).

555  78. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.

556   *Bioinformatics* **26**, 841–842 (2010).

557  79. Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic Acids Res.*

558   **46**, D1083–D1090 (2018).

559  80. Conde, D. A. *et al.* Data gaps and opportunities for comparative and conservation biology. *Proc.*

560   *Natl. Acad. Sci. U. S. A.* **116**, 9658–9664 (2019).

561  81. Blokzijl, F., Janssen, R., van Boxtel, R. & Cuppen, E. MutationalPatterns: comprehensive genome-

562   wide analysis of mutational processes. *Genome Med.* **10**, 33 (2018).

563  82. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines, Timetrees,

564   and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

565  83. Snippert, H. J. *et al.* Intestinal crypt homeostasis results from neutral competition between

566   symmetrically dividing Lgr5 stem cells. *Cell* **143**, 134–144 (2010).

567  84. Rijke, R. P., Plaisier, H. M. & Langendoen, N. J. Epithelial cell kinetics in the descending colon of

the rat. *Virchows Arch. B Cell Pathol. Incl. Mol. Pathol.* **30**, 85–94 (1979).

85.  Potten, C. S., Kellett, M., Rew, D. A. & Roberts, S. A. Proliferation in human gastrointestinal

epithelium using bromodeoxyuridine in vivo: data for different sites, proximity to a tumour, and

polyposis coli. *Gut* **33**, 524–529 (1992).

86.  Bach, S. P., Renehan, A. G. & Potten, C. S. Stem cells: the intestinal stem cell as a paradigm.

*Carcinogenesis* **21**, 469–476 (2000).

**Acknowledgements**

We are grateful to the animals whose tissues were used in this study. We thank the staff of

Wellcome Sanger Institute Sample Logistics, Sequencing and Informatics facilities for their

contribution; CASM IT and administrative staff for their help processing the data. This research

was made possible by the worldwide information network of zoos and aquariums, which are

members of Species360 and is authorized by Species360 research data use and grant agreement

#60633. We thank Dalia Conde and Johanna Staerk for help accessing Species360 data.

**Funding**

This research was funded by Wellcome (grant number 206194). For the purpose of Open Access,

the author has applied a CC BY public copyright licence to any Author Accepted Manuscript

version arising from this submission.

**Author Contributions**

A.C., E.P.M., M.R.S. and I.M. conceived the project. I.M., E.P.M., and M.R.S. supervised the

project. E.F., S.S., I.J., E.W., N.M., R.D., M.W.P., M.D., R.B., M.D.M., F.G., F.C.-C., L.P., D.B.,

E.St.J.S., B.N., and E.P.M. performed and facilitated sample collection. A.C. performed the laser

capture microdissection. A.C., L.M.R.H., A.R.J.L., Y.H., K.R., E.A., S.L., M.M, P.S., J.F., L.O.,

and C.L processed the samples. A.C., A.B.-O., F.A., N.B., T.H.H.C., M.A.S., D.J., R.A. and K.R

processed the data. J.S. and D.C provided animal longevity records. A.C., A.B.-O. and N.B. led

the analysis with help from F.A., T.H.H.C., M.A.S., A.R.J.L., T.M.B., T.D., H.J., E.P.M. and I.M.

The manuscript was written by A.C., A.B.-O., N.B. and I.M. with input from all the authors.

## Competing Interests

600     The authors declare no competing interests.

**Figures**

**Figure 1. Somatic mutation burden in mammalian colorectal crypts. a,** Histology images from horse, lion, naked mole-rat and rat colon samples, with one colorectal crypt marked in each. **b,** Burden of somatic single-base substitutions and indels per genome in each colorectal crypt sample (corrected for the size of the analysable genome). Samples are grouped by individual, with samples from the same individual coloured in the same shade. Species, and individuals within each species, are sorted by mean mutation burden **c,** Linear regression of somatic substitution burden (corrected for analysable genome size) on individual age for dog, human, mouse and naked mole-rat samples. Samples from the same individual are shown in the same colour. Regression was performed using mean mutation burden per individual. Shaded areas indicate 95% confidence intervals of the regression line. BW: black-and-white, H: harbour, N: naked, RT: ring-tailed.

**Figure 2. Mutational processes in the mammalian colon. a**, Mutational spectra of somatic
single-base substitutions in each species. The *x*-axis presents 96 mutation types on a trinucleotide
context, coloured by base substitution type. **b**, Mutational signatures inferred from (SBSB, SBSC)
or fitted to (SBS1) the species mutational spectra shown in **a**, and normalised to the human genome
trinucleotide frequencies. **c**, Estimated contribution of each signature to each sample. Samples are
arranged horizontally as in **Fig. 1b**. **d**, Regression of signature-specific mutation burdens on
individual age for human, mouse and naked mole-rat samples. Regression was performed using
mean mutation burden per individual. Shaded areas indicate 95% confidence intervals of the
regression line. BW, black-and-white; H, harbour; N, naked; RT, ring-tailed.

627
**Figure 3. Associations between somatic mutation rates and life-history variables. a**, Somatic
mutation rate per year (top) and expected end-of-lifespan substitution burden in each crypt.
Samples are grouped and sorted as in **Fig. 1b**; harbour porpoise samples were excluded due to the

unknown age of the individual. **b**, Allometric (log-log) regression of somatic mutation rate on
lifespan (left), and regression of body-mass-adjusted residuals for somatic mutation rates and
lifespan (partial correlation, **Methods**). Regressions were performed using a simple linear model
on the mean mutation rate per species. Shaded areas represent 95% confidence intervals (CI) of the
regression lines. Fraction of inter-species variance explained by the model (FVE) and *p*-value (*P*)
are indicated (note that for simple linear regression FVE = $R^2$). Dashed line denotes a reference
slope of –1. **c**, Zero-intercept linear mixed-effects (LME) regression of somatic mutation rate on
inverse lifespan (1/lifespan), presented on the scale of untransformed lifespan (*x*-axis). For
simplicity, the *y*-axis represents mean mutation rate per species, although mutation rates per crypt
were used in the regression. Darker shaded area indicates 95% CI of the regression line; lighter
shaded area marks a two-fold deviation from the regression line. The point estimate and 95% CI
of the slope (*k*), the FVE, and the range of end-of-lifespan burden (ELB) are shown. **d**, Allometric
regression and linear regression of lifespan-adjusted residuals for somatic mutation rate and body
mass. Elements are as described in **b**. **e**, Free-intercept LME regression of somatic mutation rate
on log-transformed adult mass. The *y*-axis represents mean mutation rate per species, although
mutation rates per crypt were used in the regression. Shaded area indicates the 95% bootstrap
interval of the regression line. Point estimates of the regression intercept and slope coefficients,
and model FVE, are indicated. **f**, Comparison of FVE values for free-intercept LME models using
1/lifespan and other life-history variables (alone or in combination with 1/lifespan) as explanatory
variables. Error bars indicate 95% bootstrap intervals. BMR, basal metabolic rate; BW, black-and-
white; N, naked; RT, ring-tailed.
652

**Figure 4. Association between mutation rate subtypes and species lifespan.** Zero-intercept linear mixed-effects regression of somatic rates of signature-specific substitutions, indels and mtDNA mutations (top to bottom) on inverse lifespan (1/lifespan), presented on the scale of untransformed lifespan (*x*-axis). For simplicity, y-axes represent mean mutation rate per species, although mutation rates per crypt were used in the regressions. Darker shaded areas indicate 95% confidence intervals (CI) of the regression lines; lighter shaded areas mark a two-fold deviation from the regression lines. Point estimates of the regression slope coefficient (*k*), fraction of inter-species variance explained by each model (FVE), and ranges of end-of-lifespan burden (ELB) are shown.

**Tables**

**Table 1. Variation in adult mass, lifespan, mutation rate and end-of-lifespan burden across the 16 mammalian species surveyed.** Species-level estimates are provided in Supplementary Tables 3 and 6**.**

| Variable | Minimum | Maximum | Fold variation |
|---|---|---|---|
| **Adult mass (g)** | 20.50 | 800,000.00 | 39,024.39 |
| **Lifespan (yr)** | 2.75 | 83.67 | 30.44 |
| **Mutation rate per year (SBS/genome)** | 47.12 | 796.42 | 16.90 |
| **End-of-lifespan burden (SBS/genome)** | 1828.08 | 5378.73 | 2.94 |

**METHODS**

**1. Sample collection**

We obtained colorectal epithelium and skin samples from a range of sources (**Supplementary Table 1**). For comparability across species a ~1 cm biopsy of the colorectal epithelium was taken from the terminal colon during necropsy. All necropsies occurred as soon as possible post-mortem to minimise tissue and DNA degradation. Tissue samples taken later than 24 hours post-mortem typically showed extensive degradation of the colorectal epithelium, making identification of colorectal crypts challenging. These samples were also associated with poor DNA yields and so were not included in his study. Sampled tissue was fixed in PAXgene FIX (PreAnalytiX, Hombrechtikon, Switzerland), a commercially available fixative, during the necropsy. After 24 hours in the fixative at room temperature samples were transferred into the PAXgene STABILIZER and stored at -20°C until further processing.

**2. Sample processing**

Samples were processed using a workflow designed for detection of somatic mutations in solid tissues by laser-capture microdissection (LCM) using low-input DNA sequencing. For a more detailed description see the paraffin workflow described in Ellis et al[34]. Briefly, PAXgene-fixed tissue samples of the colorectal epithelium were paraffin-embedded using a Sakura Tissue-Tek VIP tissue processor. Sections of 16 µm were cut using a microtome, mounted on PEN-membrane slides, and stained with Gill's haematoxylin and eosin by sequential immersion in the following: xylene (two minutes, twice), ethanol (100%, 1 minute, twice), deionised water (1 minute, once), Gill's haematoxylin (10 seconds, once), tap water (20 seconds, twice), eosin (5 seconds, once), tap water (20 seconds, once), ethanol (70%, 20 seconds, twice) and xylene or Neo-Clear, a xylene substitute (20 seconds, twice).

High-resolution scans were obtained from representative sections of each species. Example images are shown in **Fig. 1a** and **Extended Data Figure 2**. Individual colorectal crypts were isolated from sections on polyethylene naphthalate (PEN) membrane slides by laser-capture microdissection with a Leica LMD7 microscope. Haematoxylin and eosin histology images were reviewed by a veterinary pathologist. For some samples we also cut a section of muscle tissue from below the colorectal epithelium of the section to use as a germline control for variant calling

(**Supplementary Table 2**). Pre- and post-microdissection images of the tissue were recorded for

each crypt and muscle sample taken. Each microdissection was collected in a separate well of a

96-well plate.

Crypts were lysed using the Arcturus PicoPure Kit (Applied Biosystems) as previously

described[8,34]. Each crypt then underwent DNA library preparation, without a quantification step

to avoid loss of DNA, following the protocol described in Ellis et al.[34]. For some animals a

PAXgene fixed bulk skin biopsy was used as the germline control. For these skin samples, DNA

was extracted using the DNeasy Blood & Tissue Kit (Qiagen).

**3. Library preparation and sequencing**

Libraries from microdissected samples were prepared using enzymatic fragmentation, adapter

ligation and whole-genome sequencing following the method described in Ellis et al.[34]. Libraries

from skin samples were prepared using standard Illumina whole-genome library preparation.

Samples were multiplexed and sequenced using Illumina XTEN and Novaseq 6000 machines to

generate 150 base pair (bp) paired-end reads. Samples were sequenced to ~30× depth

(**Supplementary Table 2**).

**4. Sequence read alignment**

For each species sequences were aligned to a reference assembly (**Supplementary Table 2**)

using the BWA-MEM algorithm[73] as implemented in BWA version 0.7.17-r1188, with options '-

T 30 -Y -p -t 8'. The aligned reads were sorted using the bamsort tool from the biobambam2

package, version 2.0.86 (gitlab.com/german.tischler/biobambam2), with options 'fixmates=1

level=1 calmdnm=1 calmdnmrecompindetonly=1 calmdnmreference=<reference_fasta>

outputthreads=7 sortthreads=7'. Duplicate reads were marked using the bammarkduplicates2 tool

from biobambam2, with option 'level=0'.

**5. Variant calling**

Identification of somatic single-base substitutions (SBS) and short insertions and deletions

(indels) was divided into two steps: variant calling, and variant filtering to remove spurious calls

(see 'Variant filtering' below). For human colorectal crypts, we obtained previously sequenced

and mapped reads from a study where colorectal crypts were isolated by laser-capture

microdissection[8], and processed them using the sample variant calling and filtering process that
was applied to the non-human samples.

Substitutions were identified using the Cancer Variants through Expectation Maximization
(CaVEMan) algorithm[74], version 1.13.15. CaVEMan uses a naive Bayesian classifier to perform
a comparative analysis of the sequence data from a target and control sample from the same
individual to derive a probabilistic estimate for putative somatic substitutions at each site. The
copy number options were set to 'major copy number = 5' and 'minor copy number = 2', as in
our experience this maximises the sensitivity to detect substitutions in normal tissues. CaVEMan
identifies and excludes germline variants shared in the target (colorectal crypt) and matched
normal (skin or muscle tissue) samples, and produces a list of putative somatic mutations present
only in the target sample. CaVEMan was run separately for each colorectal crypt, using either
bulk skin or muscle microdissected from the sample colorectal biopsy as the matched normal
control (**Supplementary Table 2**). For two human donors where an alternative tissue was not
available, a colonic crypt not included as a target sample was used as the matched normal control.

Indels were identified using the Pindel algorithm[75], version 3.3.0, using a second sample from the
same individual as a matched control. The indel calls produced by Pindel were subsequently re-
genotyped using the vafCorrect tool ([github.com/cancerit/vafCorrect](github.com/cancerit/vafCorrect)), which performs a local
sequence assembly to address alignment errors for indels located at the end of sequence reads,
and produces corrected counts of sequence reads supporting the indel and corrected estimates of
variant allele fraction (VAF).

**6. Variant filtering**
A number of post-processing filters were applied to the variant calls to remove false positives
(**Extended Data Figure 17a,b**).
**Quality flag filter.** CaVEMan and Pindel annotate variant calls using a series of quality flags,
with the 'PASS' flag denoting no quality issues affecting the call[74,75]. Variant calls presenting
any flag other than 'PASS' were discarded.
**Alignment quality filter.** Variants were excluded if more than half of the supporting reads were
clipped. The library preparation methods create short insert size libraries that can result in reads
overlapping. To avoid the risk of double-counting mutant reads we used fragment-based

statistics. Variants without at least four high-quality fragments (alignment score $\geq$ 40 and base

Phred quality score $\geq$ 30) were excluded. Variants were excluded if reads supporting the variant

had a secondary alignment score that was greater than the primary alignment score. This filter

was not applied to indel calls.

**Hairpin filter.** To remove variants introduced by erroneous processing of cruciform DNA during

the enzymatic digestion we applied a custom filter to remove variants in inverted repeats[34]. This

filter was not applied to indel calls.

**Chromosome and contig filter.** For species with chromosome-level assemblies, we discarded

variants located in non-chromosomal contigs, including the mitochondrial genome (calling of

mitochondrial variants is described in the section 'Mitochondrial variant calling and filtering').

For males, variants on the Y chromosome were excluded for species where the Y chromosome

was annotated in the assembly.

**N-tract and contig-end filter.** To reduce artefactual calls due to read misalignment, we

discarded variants located within 1 kilobase (kb) of a tract of 50 or more consecutive N bases in

the reference assembly, as well as variants within 1 kb of the start or end of a contig (this implies

discarding all variants in contigs shorter than 2 kb).

**Sequencing coverage filter.** A sample-specific read depth filter was designed to exclude sites

with coverage above the 99th coverage percentile in the sample or its matched normal control, as

well as sites with coverage < 10× in the sample or its matched normal control.

**Allelic strand bias filter.** We discarded variants without any supporting reads on either the

forward or reverse strand.

**Indel proximity filter.** We discarded variants for which the total number of reads supporting the

presence of an indel within 10 bp of the variant was more than 3 times larger than the number of

reads supporting the presence of the variant. This filter was not applied to indel calls.

**Spatial clustering filter.** Visual assessment of variant calls and mutational spectra showed

spatially clustered variants to be highly enriched for artefacts. Therefore, we discarded groups of

two or more variants located within 1 kb of each other.

**Beta-binomial filter.** For each crypt, an artefact filter based on the beta-binomial distribution

was applied, which exploits read count information in other crypts from the same individual.

More specifically, for each sample, we fitted a beta-binomial distribution to the variant allele

counts and sequencing depths of somatic variants across samples from the same individual. The

beta-binomial distribution was used to determine whether read support for a mutation varies

across samples from an individual, as expected for genuine somatic mutations but not for

artefacts. Artefacts tend to be randomly distributed across samples and can be modelled as drawn

from a binomial or a lowly overdispersed beta-binomial distribution. True somatic variants will

be present at a high VAF in some samples, but absent in others, and are hence best captured by a

highly overdispersed beta-binomial. For each variant site, the maximum likelihood estimate of

the overdispersion factor ($\rho$) was calculated using a grid-based method, with values ranging

between $10^{-6}$ and $10^{-0.05}$. Variants with $\rho > 0.3$ were considered to be artefactual and discarded.

The code for this filter is based on the Shearwater variant caller[76]. We found this to be one of the

most effective filters against spurious calls (**Extended Data Figure 17b**).

**Minimum VAF filter.** For each sample, we discarded variants whose variant allele fraction

(VAF) was less than half the median VAF of variants passing the beta-binomial filter (see above)

in that sample.

**Maximum indel VAF filter.** For each sample, we discarded indels presenting VAF > 0.9, as

such indels were found to be highly enriched in spurious calls in some species. This filter was not

applied to SBS calls.

To validate our variant calling strategy, we used LCM to microdissect two sections from the

same mouse colorectal crypt. We expected to detect a high fraction of shared somatic variants in

these two sections, since their cells should be derived from the same ancestral epithelial stem cell.

Both sections were submitted for independent library preparation, genome sequencing, variant

calling and filtering using our pipeline. The majority of SBS variant calls (2742 of 2933, 93.5%)

were shared between both sections (**Extended Data Figure 17c**). In contrast, when comparing

five separate crypts from a mouse, a maximum of two variants were shared between two crypts,

and no variants were shared by three or more crypts (**Extended Data Figure 17d**).

**7. Sample filtering**

Our method for estimation of mutation rates assumes monoclonality of colorectal crypt samples.

This assumption can be violated due to several causes, including contamination from other

colorectal crypts during microdissection or library preparation, contamination with non-epithelial

cells located in or near the crypt, insufficient time for a stem cell to drift to clonality within the

crypt, or the possibility that in some species, unlike in humans[8], polyclonal crypts are the norm.

Therefore, a truncated binomial mixture model was applied in order to remove crypts that showed

evidence of polyclonality, or for which the possibility of polyclonality could not be excluded. An expectation–maximization (EM) algorithm was employed to determine the optimal number of variant allele fraction (VAF) clusters within each crypt sample, as well as each cluster's location and relative contribution to the overall VAF distribution. The algorithm considered a range of numbers of clusters (1–5), with the optimal number being that which minimised the Bayesian Information Criterion (BIC). As the minimum number of supporting reads to call a variant was 4, the binomial probability distribution was truncated to incorporate this minimum requirement for the number of successes, and subsequently re-normalised. The EM algorithm returned the inferred optimal number of clusters, the mean VAF (location) and mixing proportion (contribution) of each clone, and an assignment of each input variant to the most likely cluster. After applying this model to the somatic substitutions identified in each sample, sample filtering was performed on the basis of the following three criteria.

**Low mutation burden.** We discarded samples presenting fewer than 50 somatic variants, which was indicative of low DNA quality or sequencing issues.

**High mutation burden.** We discarded samples with a number of somatic variants greater than three times the median burden of samples from the same individual (excluding samples with less than 50 variants). This served to exclude a small minority of samples presenting evident sequencing quality problems (such as low sequencing coverage), but which did not fulfill the low-VAF criterion for exclusion (see below).

**Low VAF.** We discarded samples in which less than 70% of the somatic variants were assigned to clusters with VAF $\geq 0.3$. However, this rule was not applied to those cases in which all the samples from the same individual had primary clusters with mean VAF $< 0.3$; this was done to prevent the removal of samples from individuals presenting high fractions of non-epithelial cells, but whose crypts were nonetheless dominated by a single clone.

These criteria led to the exclusion of 41 out of 249 samples. On the basis of visual assessment of sequencing coverage and VAF distributions, we decided to preserve three samples (ND0003c_lo0004, ND0003c_lo0011, TIGRD0001b_lo0010) which we considered to be clonal, but which would have been discarded based on the criteria above.

**8. Mitochondrial variant calling and filtering**

For six species whose reference genome assemblies did not include the mitochondrial sequence, mitochondrial reference sequences were obtained from the GenBank database (**Supplementary**

**Table 5**). For each species, alignment to the reference genome was performed using BWA

version 0.7.17-r1188, as described above (see 'Sequence read alignment'). Pileup files were

generated for mtDNA genomes using the 'bam2R' function in the deepSNV (v1.32.0) R

package[76,77]. The mapping quality cut-off was set to 0, taking advantage of the fact that the

mitochondrial genome coverage for most samples was >100-fold higher than the nuclear genome

coverage, and hence most reads with poor mapping scores should be of mitochondrial origin.

Mitochondrial variants were called using the Shearwater algorithm[76] (deepSNV package

v1.32.0). Multiple rounds of filtering were applied to identify and remove false positives. The

first set of filters removed germline polymorphisms, applied a maximum false discovery rate

(FDR) threshold of $q > 0.01$, required that mismatches should be supported by at least one read

on both the forward and reverse strands, and merged consecutive indel calls. Further filtering

steps were as follows.

**Minimum VAF filter.** Only variants with VAF > 0.01 were considered for analysis, based on the

quality of the mutational spectra.

**Sequencing coverage filter.** Due to species-specific mtDNA regions of poor mappability, we

discarded sites with read coverage < 500×.

**D-loop filter.** Analysis of the distribution of mutations along the mitochondrial genome revealed

clusters of mutations within the hypervariable region of mtDNA known as the D-loop. To obtain

estimates of the mutation burden in mtDNA unaffected by hypermutation of the D-loop,

mutations in the D-loop region (coordinates MT:1–576 and MT:16,024–16,569 in human) were

excluded from this analysis

**High mutation burden.** We discarded samples having a number of somatic mtDNA variants

greater than four times the mean mtDNA burden across all samples. This served to exclude a

small minority of samples that were suspected of enrichment in false positive calls. Visual

inspection of these samples in a genome browser confirmed the presence of high numbers of

variants found on sequence reads with identical start positions and/or multiple base mismatches,

suggestive of library preparation or sequencing artefacts.

We examined the mutational spectra of somatic mtDNA substitutions on a trinucleotide sequence

context (**Extended Data Figure 16**). The specificity of the filtered variant calls was supported by

the observation that the mutational spectra across species were broadly consistent with those

previously observed in studies of human tissues[59], with a dominance of C>T and T>C

transversions and a strong replication strand bias.

**9. Mitochondrial copy number analysis**

Sequence reads from each sample were separately mapped to the species-specific mtDNA

reference sequence in order to estimate average mtDNA sequencing coverage. Excluding nuclear

reference sequences from the alignment enabled obtaining even coverage across the

mitochondrial genome by preventing mismapping of sequence reads to inherited nuclear

insertions of mitochondrial DNA (known as NuMTs). Next, coverage information from

individual mtDNA and whole-genome alignment (BAM) files was obtained using the genomecov

tool in the bedtools suite (v2.17.0)[78]. Mitochondrial copy number was calculated according to the

formula

$$depth_{mtDNA} \times ploidy \ / \ depth_{gDNA},$$

where $depth_{mtDNA}$ and $depth_{gDNA}$ are the mean coverage values for mtDNA and the nuclear

genome, respectively, and $ploidy = 2$ (assuming normal somatic cells to be diploid). For

simplicity, the sex chromosomes were excluded from the calculation of the mean nuclear genome

coverage.

**10. Analysable genome size calculation**

To estimate the somatic mutation rate, it was first necessary to establish the size of the analysable

nuclear genome (i.e. the portion of the genome where variant calling could be performed reliably)

for each sample (**Supplementary Table 4**). For both single-base substitutions and indels, the

analysable genome of a sample was defined as the complement of the union of the following

genomic regions: regions reported as 'not analysed' by the CaVEMan variant caller; regions

failing the 'chromosome and contig' filter; regions failing the 'N-tract and contig-end' filter; and

regions failing the 'sequencing coverage' filter (see 'Variant filtering'). For the analysis of

mitochondrial variants, the analysable genome of a sample was defined as the portion of mtDNA

satisfying the 'sequencing coverage' filter (see 'Mitochondrial variant calling and filtering'), after

subtracting the hypervariable region (D-loop).

**11. Life history data**

Obtaining accurate lifespan estimates is challenging; while point estimates of maximum lifespan

are available for many species, their veracity is often difficult to assess and estimates can vary

widely for the same species (**Supplementary Table 6**). There can be many causes for this

variation, including errors in recording and real variation in longevity between populations (i.e.

captive *versus* wild). As we were interested in whether the somatic mutation burden has an

association with lifespan in the absence of extrinsic mortality, we sought to obtain estimates of

longevity from individuals under human care, to minimise the impact of external factors such as

predation or infection.

Mortality records for 14 species were obtained from the Species360 database, authorized by

Species360 research data use agreement #60633 [Species360 Zoological Information

Management System (ZIMS) (2020), zims.Species360.org]. This database contains lifespan data

of zoo animals from international zoo records. Using records from 1980 to the present, we

excluded animals for which the date of birth or death was unknown or uncertain. To avoid infant

mortality influencing the longevity estimates for each species, we removed animals that died

before the age of female sexual maturity, as defined by the AnAge database[79]. This resulted in a

mean of 2,681 animal lifespan records per species for the species in the study (minimum 309,

maximum 8403; **Supplementary Table 6**). For the domestic dog, we combined records for

domestic dogs (*Canis lupus familiaris*) and wolves (*Canis lupus*) because of the paucity of

records for domestic dogs in Species360. Although the data are curated, they are still vulnerable

to the presence of inaccurate records, which can bias the lifespan estimates. To reduce the impact

of these outliers, for each species lifespan was estimated as the age at which 80% of the adults

from that species had died (**Supplementary Table 6**)[46].

Human longevity estimates were obtained using census birth and death record data from

Denmark, (1900–2020), Finland (1900–2019) and France (1900–2018), retrieved from the

Human Mortality Database [University of California, Berkeley (USA), and Max Planck Institute

for Demographic Research (Germany); www.mortality.org, www.humanmortality.de]. We

selected these countries because they had census records going back at least 100 years. To

remove the impact of infant mortality, we excluded individuals who died before the age of 13.

For each country, we selected the cohort born in 1900 and calculated the age at which 80% of the

individuals had died (Denmark, 87 years; Finland, 83 years; France, 81 years). We then used the

mean of the three countries as our estimate of the human 80% lifespan (83.7 years)

(**Supplementary Table 6**).

To test the impact of different estimates of lifespan on our results, we also obtained maximum

longevity estimates for each species from a range of databases[80] and a survey of the literature

(**Supplementary Table 6**). Other life-history metrics were obtained from the AnAge database[79]

(**Supplementary Table 6**).

**12. Mutational signature analysis**

Mutational signatures of single-base substitutions on a trinucleotide sequence context were

inferred from sets of somatic mutation counts using the sigfit (v2.0) R package[37]. Initially,

signature extraction was performed *de novo* for a range of numbers of signatures ($N = 2, ..., 10$),

using counts of mutations grouped per sample, per individual and per species. To account for

differences in sequence composition across samples, and especially across species, mutational

opportunities per sample, per individual and per species were calculated from the reference

trinucleotide frequencies across the analysable genome of each sample (see 'Analysable genome

size calculation'), and supplied to the 'extract_signatures' function in sigfit. The

'convert_signatures' function in sigfit was subsequently used to transform the extracted

signatures to a human-relative representation (**Fig. 2b**), by scaling the mutation probability values

using the corresponding human genome trinucleotide frequencies. The best-supported number of

signatures, on the basis of overall goodness-of-fit[37] and consistency with known COSMIC

signatures (cancer.sanger.ac.uk/signatures), was found to be $N = 3$. The cleanest deconvolution of

the three signatures was achieved when using the mutation counts grouped by species, rather than

by sample or individual. The three extracted signatures (labelled SBSA, SBSB, SBSC) were

found to be highly similar to COSMIC signatures SBS1 (cosine similarity 0.96), SBS5 (0.89),

and SBS18 (0.91), respectively. These signatures were independently validated using the

MutationalPatterns (v1.12.0) R package[81], which produced comparable signatures (respective

cosine similarities 0.999, 0.98 and 0.89).

This *de novo* signature extraction approach, however, failed to deconvolute signatures SBSA and

SBSB entirely from each other, resulting in a general overestimation of the exposure to SBSA

(**Extended Data Figure 18**). To obtain more accurate estimates of signature exposure, the

deconvolution was repeated using an alternative approach that combines signature fitting and

extraction in a single inference process[37]. More specifically, the 'fit_extract_signatures' function

in sigfit was used to fit COSMIC signature SBS1 (retrieved from the COSMIC v3.0 signature

catalogue; cancer.sanger.ac.uk/signatures) to the mutation counts grouped by species (with

species-specific mutational opportunities), while simultaneously extracting two additional

signatures *de novo* (SBSB and SBSC). Before this operation, COSMIC SBS1 was transformed

from its human-relative representation to a genome-independent representation using the

'convert_signatures' function in sigfit. By completely deconvoluting SBS1 and SBSB, this

approach yielded a version of SBSB that was more similar to COSMIC SBS5 (cosine similarity

0.93); the similarity of SBSC to COSMIC SBS18 was the same under both approaches (0.91).

Finally, the inferred signatures were re-fitted to the mutational spectra of mutations in each

sample (using the 'fit_signatures' function in sigfit with sample-specific mutational

opportunities) to estimate the exposure of each sample to each signature. The fitting of the three

signatures yielded spectrum reconstruction similarity values (measured as the cosine similarity

between the observed mutational spectrum and a spectrum reconstructed from the inferred

signatures and exposures) with median 0.98 and interquartile range 0.96–0.99. Although the

purely *de novo* extraction approach and the 'fitting and extraction' approach yielded comparable

versions of signatures SBSB and SBSC, the fixing of COSMIC SBS1 in the latter approach

resulted in lower SBS1 exposures and higher SBSB exposures in the majority of samples, due to

the cleaner deconvolution of these two signatures (**Fig. 2, Extended Data Figure 18**).

To examine potential variation in the spectrum of signature SBS5 across species, the following

procedure was conducted for each species: individual-specific mutation counts and mutational

opportunities were calculated for each individual in the species, and the 'fit_extract_signatures'

function was used to fit COSMIC signatures SBS1, SBS18 and SBS34 (transformed to a

genome-independent representation using the 'convert_signatures' function) to the mutational

spectra of each individual, while simultaneously inferring one additional signature

(corresponding to signature SBS5 as manifested in that species; **Extended Data Figure 6**).

To assess the presence in non-human colorectal crypts of mutational signatures caused by

APOBEC or colibactin, which have been previously observed in human crypts[8], we used an

expectation–maximisation algorithm for signature fitting, in combination with likelihood ratio

tests (LRTs). More specifically, for each non-human sample, we tested for exposure to colibactin

(signature SBS88, COSMIC v3.2) by comparing the log-likelihoods of (*i*) a model fitting

COSMIC signatures SBS1, SBS5, SBS18, SBS34 and SBS88, and (*ii*) a reduced model fitting

only the first four signatures. Benjamini–Hochberg multiple-testing correction was applied to the

*p*-values resulting from the LRTs, and colibactin exposure was considered significant in a sample

if the corresponding corrected *q*-value was less than 0.05. We followed the same approach to

assess exposure to APOBEC (SBS2 and SBS13), using two separate sets of LRTs for models

including either SBS2 or SBS13, in addition to SBS1, SBS5, SBS18 and SBS34. APOBEC

exposure was considered significant in a sample if its *q*-values for the models including SBS2

and SBS13 were both less than 0.05. This analysis identified 1/180 crypts with significant

exposure to each of colibactin and APOBEC (although the evidence for the presence of the

relevant signatures in these two crypts was not conclusive). To test for depletion of colibactin or

APOBEC exposure in non-human crypts relative to human crypts, we first applied the LRT-

based method described above to a published set of 445 human colorectal crypts[8], identifying 92

colibactin-positive and 9 APOBEC-positive crypts. We then compared the numbers of colibactin-

and APOBEC-positive crypts in the human and non-human sets using two separate Fisher's exact

tests ('fisher.test' function in R). This revealed the difference in colibactin exposure to be highly

significant ($P=7\times10^{-14}$), unlike the difference in APOBEC exposure ($P=0.30$).

Mutational spectra of somatic indels identified in each species were generated using the

'indel.spectrum' function in the Indelwald tool for R (24/09/2021 version;

github.com/MaximilianStammnitz/Indelwald).

**13. Selection analysis**

Evidence of selection was assessed using the ratio of nonsynonymous to synonymous substitution

rates (dN/dS) in the somatic mutations called in each species. The dNdScv (v0.0.1.0) R package[44]

was used to estimate dN/dS ratios for missense and truncating substitutions in each species

separately. Reference CDS databases for the dNdScv package were built for those species with

available genome annotation in Ensembl (www.ensembl.org; **Supplementary Table 2**), using

the 'buildref' function in dNdScv. For each species, the 'dndscv' function was applied to the list

of somatic substitutions called in samples of that species, after de-duplicating any substitutions

that were shared between samples from the same individual in order to avoid counting shared

somatic mutations multiple times. In addition, the analysis was restricted to genes that were fully

contained in the analysable genomes of all samples from the species (a condition satisfied by the

vast majority of protein-coding genes). Genome-wide and gene-specific dN/dS ratios were

obtained for missense and truncating substitutions in each species; no genes with statistically

significant dN/dS $\neq$ 1 were observed.

**14. Copy number analysis**

For species with chromosome-level assemblies (cat, cow, dog, horse, human, mouse, rabbit, rat),

total and allele-specific copy number (CN) were assessed in each sample adapting a likelihood

model previously applied to the detection of subclonal CN changes in normal human skin[13]. This

method exploits two sources of evidence: relative sequencing coverage and B-allele fraction

(BAF; the fraction of reads covering a heterozygous SNP that support one of the alleles). Human

samples PD36813x15 and PD36813x16 were excluded from this analysis due to the poor quality

of their SNP data.

For each sample, sequencing coverage was measured in non-overlapping 100-kilobase (kb) bins

along the species' reference genome, using the coverageBed tool in the bedtools suite (v2.17.0)[78].

For each bin, the coverage per base pair was calculated by dividing the number of reads mapping

to the bin by the bin length, and multiplying the result by the read length (150 bp). A normalised

sample–normal coverage ratio was then calculated for each bin by dividing the bin coverage in

the sample by the corresponding coverage in its matched normal control (see 'Sample

processing'). Heterozygous SNPs were isolated for each sample by selecting germline SNPs with

a BAF between 0.4 and 0.6 in the matched normal sample, and a coverage of at least 15 reads in

both the target sample and its matched normal sample. After assigning each SNP to its

corresponding 100-kb genome bin, the bins in each sample were divided into two sets: (*i*) bins

with coverage $\geq$ 10 in both the target sample and its matched normal, and at least one

heterozygous SNP; and (*ii*) bins with coverage $\geq$ 10 in both the target sample and its matched

normal, and no heterozygous SNPs. For the first set, estimates of total and allele-specific CN

were inferred by maximising the joint likelihood of a beta-binomial model for BAF and a

negative binomial model for relative coverage, as previously described[13]. The most likely

combination of allele CN values was obtained for each bin by conducting an exhaustive search of

CN values between 0 and 4, and selecting the combination maximising the joint likelihood
(calculated on the basis of expected BAF and relative coverage values). A penalty matrix was
used to penalise more complex solutions over simpler ones, as previously described[13]. For the
second set of bins (bins without SNPs), only estimates of total CN were inferred, by maximising
the likelihood of a negative binomial model for relative coverage. The most substantial
differences between these methods and the one previously published are: (*i*) SNPs were obtained
from the variant calling output, instead of from a public database; (*ii*) relative coverage was
calculated per 100-kb bin, rather than per SNP; (*iii*) SNPs were not phased within each gene, but
within each bin; (*iv*) no reference bias was assumed (i.e. the underlying BAF of heterozygous
SNPs was assumed to be 0.5); (*v*) the minimum sample purity was raised to 0.85; (*vi*) putative
CN changes were not subjected to significance testing, but selected according to their likelihood,
and subsequently filtered by means of a segmentation algorithm (see below).
426
Estimates of total and allele-specific CN per bin were merged into CN segments, which were
defined as contiguous segments composed of five or more bins with identical CN states.
Segmentation was performed separately for total and allele-specific CN estimates in each sample.
After this process, any pair of adjacent segments with the same CN assignment, and separated by
a distance shorter than 5 bins, was merged into a single segment. Finally, within each species,
segments presenting CN values other than 2 (or 1/1 for allele-specific CN), and being either
shorter than 10 bins (1 megabase), or shared among two or more samples, were discarded,
resulting in the removal of nearly all spurious CN changes.
435
**15. Mutation rate estimation**
For each sample, the somatic mutation density (mutations per bp) was calculated by dividing the
somatic mutation burden (total number of mutations called) by the analysable genome size for the
sample (see 'Analysable genome size calculation'). The adjusted somatic mutation burden
(number of mutations per whole genome) was then calculated by multiplying the mutation
density by the total genome size of the species (see below). The somatic mutation rate per year
(mutations per genome per year) was obtained by dividing this adjusted mutation burden by the
age of the individual, expressed in years (**Supplementary Table 2**). The expected end-of-
lifespan burden (ELB) for each sample was calculated by multiplying the somatic mutation rate
by the estimated lifespan of the species (see 'Life history data').

The total genome size of a species was estimated as the total size of its reference genome

assembly. Across species, the mean genome size was 2.67 gigabases (Gb), ranging between 2.41

Gb and 3.15 Gb and with a standard deviation of 221 megabases (**Supplementary Table 4**). This

suggests that inter-species variation in genome size should not have a substantial influence on the

somatic mutation rate estimates. For an assessment of alternative measures of mutation rate, see

'Association of mutation rate and end-of-lifespan burden with lifespan'.

**16. Association of mutation rate with life-history traits**

The association of the somatic mutation rate with different life-history traits was assessed using

linear mixed-effects (LME) models. In particular, associations with the following traits were

examined: lifespan (in years), adult mass (or adult weight, in grams), basal metabolic rate (BMR,

in watts), and litter size (see 'Life history data'). Associations for lifespan, adult mass and BMR

were assessed using the following transformed variables: $1/\text{Lifespan}$, $\log_{10}(\text{Adult mass})$, and

$\log_{10}(\text{BMR})$. To account for the potentially confounding effect of the correlation between

metabolic rate and body mass, the residuals of a fitted allometric regression model of BMR on

adult mass (equivalent to a simple linear regression of $\log_{10}(\text{BMR})$ on $\log_{10}(\text{Adult mass})$) were

employed as a mass-adjusted measure of metabolic rate, referred to as 'BMR residuals'.

For each variable, an LME model was implemented for the regression of somatic mutation rates

per sample on the variable of interest, using the 'lme' function in the nlme R package (v3.1-137;

[cran.r-project.org/package=nlme](cran.r-project.org/package=nlme)). To account for non-independence of the samples, both at the

individual level and at the species level, the model included fixed effects (intercept and slope

parameters) for the variable of interest, and random effects (slope parameters) at the individual

and species levels. In addition, to account for the heteroscedasticity of mutation rate estimates

across species, the usual assumption of constant response variance was replaced with explicit

species-specific variances, to be estimated within the model.

To determine the fraction of inter-species variance in mutation rate explained by each life-history

variable individually, the LME model described above was used to produce predictions of the

mean mutation rate per species; only fixed effects were employed when obtaining these

predictions, random effects being ignored. The variance of these predictions was then compared

to the variance in observed mean mutation rates; the latter were calculated for each species as the

mean of the observed mean rates per individual, to avoid individuals with larger numbers of

samples exerting a stronger influence on the species mean. The fraction of inter-species variance

explained by the model was calculated using the standard formula for the coefficient of

determination,

483
$$R^2 = \text{ESS} / (\text{ESS} + \text{RSS}),$$

484    where ESS is the explained sum of squares, and RSS is the residual sum of squares:

485
$$\text{ESS} = \sum_i (\hat{y}_i - \bar{y})^2, \quad \text{RSS} = \sum_i (y_i - \hat{y}_i)^2.$$

486    In this formulation, $y_i$ and $\hat{y}_i$ denote the observed and predicted mutation rates for species $i$,

487    respectively, and $\bar{y}$ is the overall mean rate. This definition of $R^2$ coincides with the fraction of

488    variance explained (FVE), defined as 1 minus the fraction of variance unexplained (FVU):

489
$$\text{FVE} = 1 - \text{FVU} = 1 - [\text{RSS} / (\text{ESS} + \text{RSS})] = \text{ESS} / (\text{ESS} + \text{RSS}) = R^2.$$

As the predicted and observed values correspond to mean mutation rates per species, rather than

mutation rates per sample, FVE provides a measure of the fraction of inter-species variance

explained by the fixed effects of the LME model. Among the variables considered, 1/Lifespan

was found to have the greatest explanatory power (FVE = 0.84, using a free-intercept model).

To compare the explanatory power of variables other than 1/Lifespan when considered either

individually or in combination with 1/Lifespan, the method described above was also applied to

two-variable combinations of 1/Lifespan and each of the remaining variables, using an LME

model with fixed effects for both variables and random effects for 1/Lifespan only. The $R^2$

formula above was used to measure the fraction of inter-species variance explained by each

model. In addition, to test whether the inclusion of a second explanatory variable was justified by

the increase in model fit, likelihood ratio tests between each two-variable LME model and a

reduced LME model including only 1/Lifespan were performed using the 'anova' function in the

nlme R package.

To further assess the potential effects of body mass and lifespan on each other's association with

the somatic mutation rate, allometric regression models (equivalent to simple linear models under

logarithmic transformation of both variables) were fitted to the mean somatic mutation rate per

species, using either adult mass or lifespan as the explanatory variable. In addition, the

'allometric residuals' of mutation rate, adult mass and lifespan (i.e. the residuals of pairwise

allometric regressions among these three variables) were used to examine the associations

between somatic mutation rate and either body mass or lifespan, after accounting for the effect of

the third variable (partial correlation analysis). For instance, to account for the potential influence

of body mass on the relationship between somatic mutation rate and lifespan, the residuals of an

allometric regression between mutation rate and adult mass, and the residuals of an allometric

regression between lifespan and adult mass, were analysed using simple linear regression. This

analysis supported a strong association between somatic mutation rate and lifespan

(independently of the effect of mass; FVE=0.82, $P$=3.2×10$^{-6}$; **Fig. 3c**), and a non-significant

association between somatic mutation rate and body mass (independently of the effect of

lifespan). Therefore, the relationship between somatic mutation rate and lifespan does not appear

to be mediated by the effect of body mass on both variables. Importantly, this result remains after

excluding naked mole-rat: after removing this species, partial correlation analysis still reveals a

strong association between somatic mutation rate and lifespan (FVE=0.77, $P$=4.1×10$^{-5}$), and a

non-significant association between somatic mutation rate and body mass ($P$=0.84). This

demonstrates that the observed relationships are not dependent on the presence of naked mole-rat

in the study.

To assess the robustness of the LME regression analyses described above, we performed

bootstrap analysis on each LME model, at the level of both individuals and species. More

specifically, for each level we used each of the LME models to perform regression on 10,000

bootstrap replicates, produced by resampling either species or individuals with replacement. We

then assessed the distributions of FVE across bootstrap replicates (**Extended Data Figure 14c**).

Additionally, we performed a similar bootstrap analysis using a collection of maximum longevity

estimates obtained from the literature (see section 'Life history data'). We applied the zero-

intercept LME model described above (for regressing mutation rate on inverse lifespan) on a set

of 5,000 bootstrap replicates, each of which used a set of species lifespan estimates randomly

sampled from the collection of literature-derived estimates (**Extended Data Figure 13**).

The results obtained with the LME models were additionally validated using an independent

hierarchical Bayesian model, in which the mean somatic mutation burden of each individual was

modelled as following a normal distribution with mean defined as a linear predictor containing a

species-specific slope parameter and a multiplicative offset (corresponding to the individual's

age; inclusion of this offset minimises the heteroscedasticity of the mutation rate across species,

which results from dividing mutation burdens by age). Species-specific slope parameters were in

turn modelled as normally distributed around a global slope parameter, equivalent to the fixed-

effect slope estimated by the LME model. This hierarchical model produced very similar results

to those of the LME model for all life-history variables (**Extended Data Figure 14a**).

We note that samples CATD0002b_lo0003 and MD6267ab_lo0003 were excluded from all

regression analyses, due to the fact that each shared the majority of its somatic variants with

another sample from the same individual (indicating, in each case, that both samples were closely

related), hence violating the assumption of independence among samples. The inclusion of these

two samples, however, had no effect on the outcome of the analyses.

**17. Association of mutation rate and end-of-lifespan burden with lifespan**

The relationship between somatic mutation rate and species lifespan was further explored by

adapting the LME model described in the previous section to perform constrained (zero-

intercept) regression of the adjusted mutation rate per year on the inverse of lifespan, 1/Lifespan

(see 'Life history data', 'Mutation rate estimation' and 'Association of mutation rate with life-

history traits'). The use of zero-intercept regression was motivated by the prediction that, if

somatic mutation is a determinant of maximum lifespan, then it would be expected for all species

to end their lifespans with a similar somatic mutation burden. Indeed, this was confirmed via

simple linear regression of the species mean end-of-lifespan mutation burden against species

lifespan (slope $P$=0.39). Thus, if $m$ is the mutation rate per year, and $L$ is the species' lifespan, the

expected relationship is of the form

$$m\,L \approx k,$$

where $k$ is a constant representing the typical end-of-lifespan mutation burden across species.

According to this relationship, the mutation rate per year is linearly related to the inverse of

lifespan,

$$m \approx k\,(1/L).$$

Therefore, the cross-species average end-of-lifespan burden ($k$), can be estimated as the slope

parameter of a zero-intercept linear regression model with the mutation rate per year ($m$) as the

dependent variable, and the inverse of lifespan (1/$L$) as the explanatory variable. To this purpose,

the LME model described in the previous section was altered by removing the fixed-effect

intercept parameter, thus considering only fixed- and random-effect slope parameters for

1/Lifespan.

The zero-intercept LME model estimated a value of $k = 3210.52$ (95% confidence interval

2686.89–3734.15). The fraction of inter-species variance explained by the zero-intercept model

(FVE) was 0.82, while the LME model described in the previous section (which estimated $k =$

2869.98, and an intercept of 14.76) achieved FVE = 0.84 (see 'Association of mutation rate with

life-history traits'). To test whether the increase in model fit justifies the inclusion of an intercept,

both models were compared using a likelihood ratio test (as implemented by the 'anova' function

in the nlme R package [v3.1-137]). This yielded $P=0.23$, indicating that the free-intercept model

does not achieve a significantly better fit than the zero-intercept model. Similarly, the zero-

intercept model yielded lower values for both the Bayesian information criterion (BIC) and the

Akaike information criterion (AIC). Notably, equivalent analyses using somatic mutation rates

per megabase and per protein-coding exome (instead of per whole genome) yielded comparable

results (**Extended Data Figure 12**).

To investigate the possibility of phylogenetic relationships between the species sampled

confounding the analysis, a phylogenetic generalised linear model was used to regress the mean

mutation rate of each species on the inverse of its lifespan ($1/L$), while accounting for the

phylogenetic relationships among species. A phylogenetic tree of the 15 species examined was

obtained from the TimeTree resource[82], and the phylogenetic linear model was fitted using the

'pgls' function in the caper R package (v1.0.1; cran.r-project.org/package=caper). The estimates

produced by zero-intercept regression of mean mutation rates per species on 1/Lifespan were

compared between this phylogenetic generalised linear model and a simple linear model ('lm'

function in R). The use of this simple model, as well as the use of mean mutation rates per

species (rather than mutation rates per sample), was necessary due to the impossibility of

replicating the heteroscedastic mixed-effects structure of the LME model employed for the main

association analyses (see 'Association of mutation rate with life-history traits') within the

phylogenetic linear model. Both the phylogenetic linear model and the simple linear model

produced similar estimates (**Extended Data Figure 14b**), suggesting that phylogenetic non-

independence of the samples does not have a substantial effect on the association analyses.

**18. Cell division analysis**

To investigate the extent to which differences in cell division rate could explain differences in mutation rate and burden across species, we obtained estimates of intestinal crypt cell division rates from mouse[83], rat[84] and human[85,86] (**Supplementary Table 7**). Using these cell division rates, our lifespan estimates and the observed SBS mutation rates, we calculated the number of cell divisions at the end of lifespan and the corresponding number of mutations per cell division expected under a simple model assuming that all mutations occur during cell division (**Supplementary Table 7**).

We investigated whether differences in the number of cell divisions among species could explain the observed differences in mutation burden. Although colorectal cell division rate estimates are lacking for most species, existing estimates from mouse, rat and human indicate that the total number of stem cell divisions per crypt in a lifetime varies greatly across species, for example with ~6–31 fold more divisions per intestinal stem cell in a human than a rat over their respective lifetimes, depending on the cell division rate estimates used (**Supplementary Table 7**). Mouse intestinal stem cells are estimated to divide once every 24 hours[83], while estimates of the human intestinal stem cell division rate vary from once every 48 hours[85] to once every 264 hours[86]. Thus mouse intestinal stem cells divide 2–11 times faster than human intestinal stem cells. By the end of lifespan, an intestinal stem cell is predicted to have divided ~1351 times in a mouse, ~486 times in a rat and 2774–15,257 times in a human (depending on the cell division rate estimate used). Applying our somatic mutation burden and lifespan data, this implies that the somatic mutation rate per cell division in a mouse is ~1.5–8.4 fold higher than in a human. However, the observed fold difference in somatic mutation rate between these two species is 16.9 (**Table 1**). Therefore, differences in cell division rate appear unable to fully account for the observed differences in mutation rate across species. Nevertheless, we note that accurate cell division rate estimates for basal intestinal stem cells are lacking for the majority of species.

**19. Data reporting**

No statistical methods were used to predetermine sample size. The experiments were not randomised and the investigators were not blinded to allocation during experiments and outcome assessment.

**20. Ethics statement**

All animal samples were obtained with the approval of the local ethical review committee

(AWERB) at the Wellcome Sanger Institute and those at the holding institutions.

**21. Data and code availability**

DNA sequence data have been deposited in the European Genome-Phenome Archive (ega-
archive.org) under overarching accession EGAD00001008032. The computer code and data files
used in the analyses are available on Zenodo (doi.org/10.5281/zenodo.5554801;
doi.org/10.5281/zenodo.5554777) and GitHub (github.com/baezortega/CrossSpecies2021).

656 **Extended Data Figures**

658 **Extended Data Figure 1. Somatic mutational spectra of the human colon and small**

659 **intestine.** Trinucleotide-context mutational spectra of somatic single-base substitutions from

660 human adult stem cells in colon (top) and small intestine, using mutation calls obtained from

661 Blokzijl *et al.*[7].

    **Extended Data Figure 2. Histology images of intestinal crypts across species.** Histological

667    images of the colorectal or intestinal (ferret) epithelium for each non-human species. Scale bars

668    are provided at the bottom of each image.

670
671
672
673

674 **Extended Data Figure 3. Somatic VAF distributions per species.** Distributions of variant
675 allele fraction (VAF) for somatic substitutions in each crypt for each species. Each distribution
676 refers to the variants in a single sequenced crypt.
677

678
679

**Extended Data Figure 4. Somatic mutation accumulation across species.** Each panel presents
somatic substitution burdens per genome for a given species. Each dot represents a crypt sample,
with samples from the same individual sharing the same colour. For species with two or more
individuals, the estimated regression line from a simple linear regression model on individual
mean burdens is shown. For species with three or more individuals, blue shaded regions indicate
the 95% confidence intervals of the regression line. Harbour porpoise samples were excluded
because the individual was of unknown age.

**Extended Data Figure 6. Profiles of signature SBSB as inferred from different species.**

Trinucleotide-context mutational spectra of signature SBSB, as inferred independently from

somatic mutations in crypts from four representative species (top to bottom): human, naked mole-

rat, rat, and rabbit (Methods). Signatures are shown in a human-genome-relative representation.

Cosine similarities between each signature and the human COSMIC SBS5 and SBS40 are

provided.

**Extended Data Figure 8. Colibactin exposure in non-human and human colorectal crypts.**

Exposures to mutational signatures SBS1, SBS5, SBS18, SBS34 and SBS88, as inferred via

expectation–maximisation, for 180 non-human crypts (top) and 445 human crypts sequenced in a

previous study[8]. Asterisks indicate samples with statistically significant colibactin (SBS88)

exposure, based on a likelihood ratio test (Methods). BW, black-and-white; H, harbour; N, naked;

RT, ring-tailed.

**Extended Data Figure 9. Identified copy number changes.** Somatic copy number changes in
cow (**a**, **b**), mouse (**c**) and human (**d**) colorectal crypts. For each case, chromosomes are presented
along the horizontal axis, and each point represents a 100-kb genomic bin. The top panel presents
the ratio between observed and expected sequencing coverage per bin; the middle panel shows
the median B-allele fraction (BAF) of heterozygous germline SNPs per bin; and the bottom panel
presents the inferred segments of total copy number (green) and allele-specific copy number
(red/blue). Regions of copy number change are highlighted in pink. The sparsity of BAF and
allele-specific copy number values in the mouse crypt (**c**) are due to the fact that mouse samples
generally had very low numbers of germline SNPs.

**Extended Data Figure 10. Somatic dN/dS.** Estimates of dN/dS for somatic missense and

truncating mutation in each of the species with available genome annotation. Dots and error bars

represent maximum likelihood estimates and 95% confidence intervals, respectively. Note the

logarithmic scale of the vertical axis.

**Extended Data Figure 11. Kaplan-Meier curves of longevity in captivity.** Kaplan Meier

survival curves for each species using captive lifespan data from Species360 for non-human

species and census record data for humans (**Methods**). The red shaded areas represent 95%

confidence intervals of the curve. A horizontal grey bar indicates the 80th percentile, which was

employed as a robust estimate of species lifespan.

**Extended Data Figure 12. Associations between life-history variables and alternative**

**measures of somatic mutation rate.** The figure presents the same analyses as **Fig. 3d,e**, but

employing somatic mutation rates per megabase (**a**), or per protein-coding exome (**b**), rather than

per genome (**Methods**). Leftmost panels show zero-intercept linear mixed-effects (LME)

regressions of somatic mutation rates on inverse lifespan (1/Lifespan), presented on the scale of

untransformed lifespan (horizontal axes). Vertical axes present mean mutation rate per species,

although mutation rates per crypt were used in the regressions. Darker shaded areas indicate 95%

confidence interval (CI) of the regression lines; lighter shaded areas mark a two-fold deviation

from the regression line. Point estimate and 95% CI of the regression slope coefficient (*k*),

fraction of inter-species variance explained by the model (FVE), and range of end-of-lifespan

burden (ELB) are provided. Rightmost panels show comparisons of FVE values achieved by

free-intercept LME models using inverse lifespan and other life-history variables (alone or in

combination with inverse lifespan) as explanatory variables. BMR, basal metabolic rate; BW,

black-and-white; Mb, megabase; N, naked; RT, ring-tailed.

**Extended Data Figure 14. Comparison of regression models for somatic mutation rates. a**,

Constrained-intercept regression of somatic substitution rates on the inverse of lifespan

(1/Lifespan), using a linear mixed-effects (LME) model applied to the rates per crypt (left) and a

Bayesian hierarchical normal regression model applied to the mean rates per individual. For

simplicity, black dots present mean mutation rates per species. Darker shaded area indicates 95%

confidence/credible interval (CI) of the regression line; lighter shaded area marks a two-fold

deviation from the regression line. Point estimates and 95% CI of the regression slopes ($k$) and

fraction of inter-species variance explained (FVE) are provided. **b**, Comparison of regression

lines for the of somatic substitution rates on 1/Lifespan (left; zero intercept) and log-transformed

adult mass (right; free intercept), using simple linear models (dark and light blue), phylogenetic

generalised least-squares models (orange and yellow), Bayesian hierarchical normal models

(green) and LME models (red). Point estimates of the regression coefficients for each model are

provided. **c**, Distributions of regression FVE under individual- and species-level bootstrapping.

For the LME models regressing somatic mutation rates on inverse lifespan (zero intercept; left)

and log-mass (free intercept), the curves present distributions of FVE from 10,000 bootstrap

replicates, obtained through random resampling of either individuals (blue) or species (orange)

(**Methods**). Vertical lines indicate the FVE values obtained using the entire data set.

 **Extended Data Figure 15. mtDNA mutation burden and copy number. a**, Total somatic

801    mtDNA mutations (substitutions and indels) called in each sample. **b**, Somatic mutation burden

802    per mitochondrial genome copy per sample. **c**, Estimated mtDNA copy number per sample.

803    Samples are arranged as in **Fig. 1b**, with samples from the same individual coloured in the same

804    shade of grey.

806
807
808

809    **Extended Data Figure 16. Mutational spectra of mtDNA substitutions in each species.**

810    Horizontal axis presents 96 mutation types on a trinucleotide context, coloured by base

811    substitution type. Mutations on the upper and lower halves of the spectrum represent substitutions

812    in which the pyrimidine base is located in the heavy and light strands of mtDNA, respectively.

**Extended Data Figure 17. Assessment of variant calling and filtering. a**, Spectra of SBS calls
before (left) and after application of the final eight variant filters, across all giraffe samples. Note
that the set of 'unfiltered' variants (left) has gone through the three early filters named 'quality
flag filter', 'alignment quality filter' and 'hairpin filter' (Methods). **b**, Spectra of calls flagged as
artefactual by each of the final eight variant filters, across all giraffe samples. Sets of variants
flagged by different filters are not mutually exclusive. **c**, Venn diagram showing the number of
variant calls shared between two LCM sections from the same mouse colorectal crypt. **d**, Venn
diagram showing the numbers of variant calls shared between five different colorectal crypts
from the same mouse.

**Extended Data Figure 18. Mutational signatures and exposures inferred *de novo*. a**,

Mutational signatures inferred *de novo* from the species mutational spectra shown in **Fig. 2a**.

Signatures are shown in a human-genome-relative representation. SBSA is the *de novo*

equivalent of COSMIC signature SBS1 (**Fig. 2b**). **b**, Exposure of each sample to each of the

mutational signatures shown in **a**. Samples are arranged horizontally as in **Fig. 1b**. **c**, Regression

of signature-specific mutation burdens on individual age for human, mouse and naked mole-rat

samples. Regression was performed using mean mutation burden per individual. Shaded areas

indicate 95% confidence intervals of the regression line. BW, black-and-white; H, harbour; N,

naked; RT, ring-tailed.

**Supplementary Tables**

**Supplementary Table 1. Species information.** For each of the (sub)species in the study, the table provides: common name, scientific name, number of individuals in the study, number of colorectal crypts sequenced, range of individual ages, and source institution.

**Supplementary Table 2. Sample information.** For each colorectal crypt sample in the study, the table provides: sample ID, individual ID, species name, matched normal sample ID, matched normal sample type, and median sequencing depth.

**Supplementary Table 3. Mutation rate and burden regression coefficients per species.** For each species in the study (except harbour porpoise), the table provides the mean observed values of the rate of somatic substitutions per genome per year, and point estimates and 95% confidence intervals for simple linear regression of mean substitution burdens per individual on individual ages. The estimated regression slopes correspond to the estimated mutation rate per year for each species. Estimates are provided for constrained-intercept linear models applied to all species, and for free-intercept linear models applied to the eight species with at least three individuals.

**Supplementary Table 4. Somatic mutation burdens, rates and signature exposures.** For each colorectal crypt sample in the study, the table provides: sample ID, individual ID, species name, individual age, total genome size, coding genome size, analysable genome size, analysable mtDNA size, mutational signature exposures (SBS1, SBSB, SBSC); somatic mutation burdens per genome for single-base substitutions, indels, signature-specific substitutions (SBS1, SBSB, SBSC), and mtDNA mutations; and somatic mutation rates per genome per year for single-base substitutions, indels, signature-specific substitutions (SBS1, SBSB, SBSC), and mtDNA mutations.

**Supplementary Table 5. Reference genome information.** For each species in the study, the table provides: reference genome version used, reference mtDNA sequence used, Ensembl genome annotation version used (where applicable), reference genome file source, reference mtDNA file source, reference genome file URL, reference mtDNA file URL.

**Supplementary Table 6. Life history data.** For each species in the study (except harbour

porpoise), the table provides: adult mass (g), basal metabolic rate (W), litter/clutch size,

maximum longevity (years); and maximum likelihood estimate and 95% confidence limits for the

estimated 80% lifespan, together with the corresponding sample size. The source of each estimate

is given in brackets.

**Supplementary Table 7. Number of cell divisions per lifespan.** For mouse, rat and human, the

table provides: estimated rate of colorectal cell division (hours), estimated lifespan (years),

estimated number of cell division at the end of lifespan, estimated number of mutations per cell

division (obtained using our mutation rate estimates), and the reference for the cell division rate.

For humans, two estimates of cell division rate are included.

887