Selective estrogen receptor modulators limit alphavirus infection by targeting the viral capping enzyme nsP1

Rajat Mudgal[1], Chandrima Bharadwaj[2], Aakriti Dubey[1], Shweta Choudhary[1], Perumal Nagarajan[2], Megha Aggarwal[1], Yashika Ratra[2], Soumen Basak[2*], Shailly Tomar[1*]

[1]Department of Biosciences and Bioengineering, Indian Institute of Technology Roorkee, India

[2]National Institute of Immunology, New Delhi, India

Correspondence:

*S. Tomar, Department of Biosciences and Bioengineering,

Indian Institute of Technology Roorkee,

Roorkee 247667, Uttarakhand, India

Tel: +91 1332285849

E-mail: shailly.tomar@bt.iitr.ac.in

*S. Basak, Systems Immunology Laboratory,

National Institute of Immunology,

Aruna Asaf Ali Marg, New Delhi 110067, India

Email: sobasak@nii.ac.in

Abstract

Alphaviruses cause animal or human diseases that are characterized by febrile illness, debilitating arthralgia, or encephalitis. Selective estrogen receptor modulators (SERMs), a class of FDA-approved drugs, have been shown to possess antiviral activities against multiple viruses, including Hepatitis C virus, Ebola virus, dengue virus, and vesicular stomatitis virus. Here, we evaluated three SERM compounds, namely 4-hydroxytamoxifen, tamoxifen, and clomifene, for plausible antiviral properties against two medically important alphaviruses, chikungunya virus (CHIKV) and Sindbis virus (SINV). In cell culture settings, these SERMs displayed potent activity against CHIKV and SINV at non-toxic concentrations with EC_{50} values ranging between 400 nM and 3.9 µM. Further studies indicated that these compounds inhibit a post-entry step of the alphavirus life cycle, while enzymatic assays involving purified recombinant proteins confirmed that these SERMs target the enzymatic activity of non-structural protein 1 (nsP1), the capping enzyme of alphaviruses. Finally, tamoxifen treatment restrained CHIKV growth in the infected mice and diminished musculoskeletal pathologies. Combining biochemical, cell culture-based studies, and *in vivo* analyses, we strongly argue that SERM compounds, or their derivatives, may provide for attractive therapeutic options against alphaviruses.

Topic: Alphavirus, chikungunya virus, Sindbis virus, SERMs, tamoxifen, clomifene, antiviral, nsP1, murine model

Introduction

Alphaviruses are enveloped arboviruses and possess a positive-sense (+ve sense), single-stranded RNA genome. Symptomatic manifestations of alphavirus infection in humans vary from fever or rash to incapacitating polyarthralgia and intense asthenia persisting for months (1). These viruses can be broadly classified into New World alphaviruses, such as Venezuelan equine encephalomyelitis virus (VEEV), and Old World alphaviruses, such as chikungunya virus (CHIKV) and Sindbis virus (SINV). While Old World viruses mainly cause polyarthritis, New World viruses are predominantly associated with encephalitis in the infected host (2). In particular, numerous re-emergence cases of CHIKV infections were reported in the past 50 years; the most severe outbreak was observed in the year 2005 in La Réunion and parts of the Indian subcontinent resulting in significant morbidities, economic losses, and stress on the health care infrastructure (3, 4). On the other hand, SINV has been attributed to recurrent outbreaks in Scandinavia, Africa, Europe, and the Middle East, since 1952 (5). Despite significant human health relevance, there are currently no licensed antiviral drugs or vaccines to combat the alphavirus menace. Treatment of infection is palliative and consists of the administration of analgesics or antipyretics to impart symptomatic relief.

Post-infection, the alphaviral genome is translated to produce a non-structural and a structural polyprotein. Proteolytic cleavage of these polyproteins generates functional viral polypeptides (2, 6). Proteolytic cleavage of the non-structural polyprotein generates four non-structural proteins (nsPs) viz. nsP1, nsP2, nsP3, and nsP4, which play essential roles in viral RNA (vRNA) synthesis within host cells (7). In particular, nsP1, which is a membrane-associated enzyme, directs the capping of genomic and subgenomic vRNAs with its methyltransferase (MTase) and guanylyltransferase (GTase) activities (8). Capping of the vRNA is crucial for translation

initiation, protecting vRNAs from host exonucleases, and masking viral nucleic acids from host-encoded pathogen recognition receptors, such as MDA5 and RIG-I (9). Due to a distinctive and non-canonical capping mechanism, nsP1 is thought to be a promising antiviral target for alphaviruses (10–12).

Identification and evaluation of novel antiviral therapeutics is a challenging and time-consuming process. However, repurposing drugs already in clinical use may expedite antiviral development against alphaviruses (13). 4-hydroxytamoxifen (4-OHT), tamoxifen, and clomifene belong to the triphenylethylene class of molecules and are known estrogen receptor antagonists (14, 15). Tamoxifen is typically used as a maintenance agent in estrogen-receptor-positive tumor patients (16). 4-OHT is a metabolite of tamoxifen (17). Clomifene is used for treating female infertility. Both tamoxifen and clomifene have also found their off-label use in treating male infertility and idiopathic gynecomastia (18, 19). These compounds, along with their analogs, are generally referred to as selective estrogen receptor modulators (SERMs) (15). Previous reports revealed that SERMs are active against multiple viruses, including the hepatitis C virus (HCV), herpes simplex type 1 virus (HSV-1), Ebola virus, dengue virus, vesicular stomatitis virus (VSV), etc (20–25). However, the plausible role of SERMs in limiting alphavirus infections has not been examined.

Here, we report potent antiviral activity of three SERMs, namely, 4-OHT, tamoxifen, and clomifene, against CHIKV and SINV in both cell culture-based settings and the preclinical mouse model. Our biochemical studies further established that SERMs inhibit the enzymatic activity of nsP1 from multiple alphaviruses. Taken together, SERM compounds may offer an effective means for antiviral therapy against the alphavirus genus.

Materials and Methods:

Cells, animals, viruses, and compounds

Vero cells (NCCS, Pune, India) were maintained in DMEM (HiMedia, India) supplemented with 10% heat-inactivated FBS (Gibco), 100 units of penicillin, and 100 mg/ml of streptomycin at 37°C and in a 5% CO_2 incubator. Stocks of CHIKV (accession number, KY057363.1; strain, 119067) and SINV were maintained using standard viral adsorption techniques (26, 27). (Z)-4-OHT, tamoxifen, and clomifene citrate were purchased from Cayman chemicals and dissolved in 100% dimethyl sulfoxide (DMSO). WT C57BL/6 mice were housed at the National Institute of Immunology (NII) and used adhering to the institutional guidelines (Approval number – IAEC 545/20).

In vitro **antiviral assays:** Cytotoxicity of various compounds was evaluated in MTT assay and LDH assay (see the supporting information). For inhibitor treatment of the infected cells, sub-confluent Vero cells were infected for 1.5 h at MOI:1 in a 24-well plate at a density of $1x10^5$ cells/well. The media was then removed, and compound dilutions, lower than their respective toxic doses, prepared in low-serum media were added and cells were incubated at 37°c. At 24 h post-infection (hpi), culture supernatant was harvested and subjected to plaque-forming assay (see a detailed description of the plaque assay in the supporting information). The reduction in viral titer relative to DMSO-treated cells was measured and graphs were analysed using Graph Pad Prism for determining the half-maximal effective concentration (EC_{50}) values. The antiviral activity of various compounds was also examined in the indirect immunofluorescence assay, where intracellular viral particles were stained using anti-CHIKV antibodies (see the supporting information for details). Cells were harvested at various time points and the abundance of intracellular and extracellular vRNA was quantified in the qRT-PCR assay. A detailed

description of the luciferase-based antiviral assays, qRT-PCR protocol, and primers are available in the supporting information.

nsP1 expression, purification, and enzymatic assays: CHIKV nsP1 and VEEV nsP1 were expressed and purified from *E. coli* as described in a previous study (12). Reaction conditions and data analysis for capillary electrophoresis (CE)-based MTase assay and enzyme-linked immunosorbent assay (ELISA)-based guanylation assay were described earlier (12, 28).

CHIKV infection in mice and tamoxifen treatment: Six weeks old male mice were subcutaneously (SC) injected on the ventral side of the right footpad with 10^6 PFU of CHIKV (strain no: 119067) (29). CHIKV-infected mice were injected with tamoxifen dissolved in sesame oil at day+1, day+3, and day+5 post-infection. Control mice were injected with an equal volume of oil. Infected mice were further examined for body weight changes at the indicated days post-infection. The swelling of the infected footpad was quantified by estimating the increase in the footpad upon infection using Insize digital Vernier caliper (30). In certain sets, mice were sacrificed at day six, and tissues from decalcified footpads were subjected to H&E staining for scoring pathologies associated with musculoskeletal inflammation. At the indicated days post-infection, blood was collected through the retro-orbital route, and the footpad was dissected from the infected mice. Total RNA was isolated from blood or tissue samples and the abundance of vRNA was determined by qRT-PCR using primers specific for the +ve sense genome (31). Separately, we prepared a serial dilution of CHIKV stock, whose titer was determined using the standard plaque assay. RNA was isolated and qRT-PCR was performed using these serial diluents. By plotting C_T values against the known CHIKV titer, a standard curve was generated. We used this standard curve for estimating the number of copies of RNA

genomes present in samples from infected mice. Alternatively, the viral titer in these tissues was determined by the $TCID_{50}$ method (see supporting information) (32).

Results

SERMs inhibit alphaviral propagation

As SERMs were active against multiple viruses, we sought to examine if this family of drugs also display antiviral properties against alphaviruses. Prior to the evaluation of antiviral activity, cell viability of BHK-21 and Vero cells in the presence of compounds was determined by MTT [3-(4,5- Dimethylthiazol- 2-yl)-2,5- Diphenyltetrazolium Bromide] assay (Figure S1A-C and S2A-C). An additional cytotoxicity assay, Lactate Dehydrogenase assay (LDH) assay, which assesses the membrane integrity of cells, was also performed in Vero cells (Figure S2D-F). The calculated half-maximal cytotoxic concentration (CC_{50}) values are presented in Table 1. A luciferase reporter system, SINV-FLuc, that has a *firefly luciferase* gene fused at the 3´end of the SINV E1, was used for primary evaluation of antiviral activity of compounds (Figure 1A). The compounds were tested in BHK-21 cells at 2.5 µM concentration at which there was no observable cytotoxicity. All the three compounds showed a significant decrease in luciferase activity that was used as a surrogate of viral replication in the infected cells (Figure 1A).

The quantitative analysis of the antiviral activity of the compounds was done by the titer determination of the progeny virus released 24 hpi in the supernatant of the treated or untreated cells infected with CHIKV or SINV. All three compounds demonstrated a marked reduction in infectious virus production in a dose-dependent manner (Figure 1B). CHIKV-infected Vero cells treated with 4-OHT or tamoxifen showed maximally 1 log_{10} reduction relative to mock-treated

cells while clomifene treatment reduced viral titer by 2 $\log_{10}$ at higher concentrations. Similarly, SINV titers were reduced maximally by 2 $\log_{10}$ when infected cells were treated with clomifene or tamoxifen. Even greater viral titer reduction (>3 $\log_{10}$) was observed upon 4-OHT treatment of SINV-infected Vero cells. Selectivity index values calculated from EC_{50} and CC_{50} suggested that within an order of magnitude, these drugs displayed largely comparable selectivity against CHIKV and SINV (Table 1). The anti-CHIKV and anti-SINV activities of 4-OHT, clomifene, and tamoxifen were further validated by qRT-PCR. There was a significant reduction in intracellular vRNA in the cells treated with the tested compounds 24 hpi. (Figure 2A). Corroborating our plaque assay, immunostaining cells with an anti-alphavirus antibody confirmed that treatment of CHIKV- or SINV-infected cells with 4-OHT, tamoxifen, or clomifene substantially suppress the abundance of viral proteins at 36 hpi (Figure 2B).

Table 1: The CC_{50}, EC_{50}, and selectivity index values of 4-OHT, tamoxifen, and clomifene in BHK-21 and Vero cells.

Compound	BHK-21 cells	Vero cells						
	CC_{50}-MTT (µM)	CC_{50}-MTT (µM)	CC_{50}-LDH (µM)	CHIKV		SINV		
				EC_{50} (µM)	SI^1	EC_{50} (µM)	SI^2	
4-OHT	8.83	14.63	14.92	0.93	15.7	0.40	36.5	
Tamoxifen	12.25	25.40	19.61	3.95	6.4	3.69	6.9	
Clomifene	8.54	26.73	20.43	3.16	8.5	1.24	21.5	

[1] Selectivity index (CC_{50}-MTT/ EC_{50} CHIKV)
[2] Selectivity index (CC_{50}-MTT/ EC_{50} SINV)

2 **Inhibition of alphaviral replication at a post-entry step**

3 To gather insight into the mode of action of these compounds, time-of-addition studies were

4 performed (Figure 3A-D). When cells were pretreated with the compounds before virus

infection, minimal inhibitory effect on the viral yield was observed. A cotreatment assay was performed to examine if the compounds can affect viral binding and entry into the cell. Viral yield remained largely unaffected when compounds were present during infection suggesting that compounds do not interfere with virus particle stability or its attachment to the host cell. Moderate dose-dependent inhibition of viral replication observed in cotreatment assay can be attributed to the possible effects of the residual amount of inhibitors retained by the cells during cotreatment (Figure 3B). All three compounds were most effective against both CHIKV and SINV when they were added during an early post-entry phase of viral replication in the infected cells suggesting a similar antiviral mechanism of the inhibitors in these two alphaviruses (Figure 3C-D).

A SINV reporter replicon was designed to investigate the inhibitory effect of the compounds (Figure 3E). The tested compounds showed a moderate inhibitory effect against SINV replicon indicating a weak suppression of synthesis of viral genome and proteins by nsPs (Figure 3E). Additionally, to examine if the compounds exert an inhibitory effect on vRNA replication and to minimize the contribution of multiple rounds of viral infection, intracellular vRNA from virus-infected cells was quantified at 6 hpi. A statistically significant but modest decrease in cell-associated vRNA was observed 6 hpi for CHIKV and SINV (Figure 3F and Figure 3G). To further examine the mechanistic function of these SERMs against alphaviruses, SINV vRNA in the extracellular fractions was quantified (Figure 3H). Intriguingly, there was markedly less reduction in extracellular virus-associated vRNA as compared to the infectious virions released at 24 hpi (Figure 3H and Figure 1B).

Further, CHIKV was serially passaged in the presence of gradually increasing concentrations of compounds. Viral-induced cytopathic effects (CPE) were monitored and no change in viral

susceptibility to the inhibitors was observed even after multiple passages. Only partial resistance

to 4-OHT could be observed after 15 passages which suggested that a high resistance is indeed

difficult to generate against these compounds. Plaque-purified variant harboring all four

compensatory mutations also displayed an altered plaque phenotype compared to the wild-type

virus but did not offer resistance against tamoxifen or clomifene (Figure 4A-B). This partially-

resistant variant displayed ~4-fold resistance (EC_{50} 4-OHT-resistance variant [3.99 µm]/EC_{50}

wild-type [0.93 µM]) relative to wild-type virus (Figure S3) The sequencing data of partially-

resistant isolates revealed four non-synonymous mutations in the nsP-coding region (Figure 4B).

Two mutations lie in the zinc-binding and macro domains of the nsP3-coding region, one is in

disordered region of the nsP4-coding region, and one lies in the MTase/GTase domain of nsP1.

While the mutations in nsP3 and nsP4 fall in the enzymatically-inactive regions, the point of

mutation in nsP1 lies in the catalytic domain of the enzyme indicating that the likely molecular

target of SERM compounds is alphaviral nsP1. Additionally, mutations identified within nsP3

and nsP4 proteins correlate with domains of the proteins that are known to, or have strong

evidence for, the protein-protein interaction (33–35).

SERM compounds inhibit alphaviral nsP1 activity

Results of the studies described above prompted us to evaluate the effect of these compounds on

the enzymatic activity of alphaviral nsP1. A non-radioactive CE-based MTase assay that

measures the amount of SAH generated during CHIKV and VEEV nsP1 MTase reactions was

utilized (12). Further, an ELISA-based assay that measures the m[7]GMP-nsP1 adduct formed in

guanylation reaction was used to determine the inhibitory activity of compounds on CHIKV

nsP1 (28). We observed that the compounds inhibited the MTase activity of both CHIKV and

VEEV (Figure 4C). The ELISA-based assay also showed a concentration-dependent reduction in

CHIKV nsP1 enzymatic activity in the presence of tested SERMs (Figure 4D).

Tamoxifen alleviates CHIKV-induced pathologies *in vivo*

Next, we set out to examine if the anti-alphaviral property of SERMs was preserved *in vivo*. To

this end, we adopted the previously published CHIKV footpad infection model involving adult

wild-type mice that recapitulates certain arthritogenic phenotypes observed in human subjects

(29). As described earlier, subcutaneous CHIKV infection into the footpad caused footpad

swelling peaking at day two and again at day six (Figure S4) (29). Using tamoxifen as a

representative drug, we examined the efficacy of SERMs in reducing these inflammatory

musculoskeletal pathologies. We optimized a therapeutic regime where CHIKV-infected mice

were successively injected with 20 mg/kg of tamoxifen on every alternate day beginning at day

one post-infection. We found that compared to the vehicle control, tamoxifen treatment

significantly improved the footpad swelling at day six post-infection, while somewhat

moderately restraining footpad pathology at day two (Figure 5A-B).

Because tamoxifen also directly impedes inflammatory pathways, we asked if the curative

tamoxifen effect was indeed associated with reduced viremia *in vivo* (36). Consistent with the

reduced musculoskeletal pathology, tamoxifen treatment led to a more than one log decrease in

the titer of plaque-forming CHIKV in the infected footpad at both day two and day six post-

infection (Figure 5C). Similarly, we captured an approximately five-fold decrease in the level of

vRNA in the blood of tamoxifen-treated mice at day two (Figure 5D). In parallel, tamoxifen

treatment ensured a close to one log reduction in the level of viral genome RNA in the footpad at

both day two and day six (Figure 5E). Histopathological analyses of tissues derived from the

joint footpad revealed that CHIKV infection triggered marked subcutaneous edema, profound

infiltration of mononuclear immune cells, and local necrosis at day six (Figure 5F) (29).

Tamoxifen treatment regime considerably ameliorated these signatures of CHIKV-induced

inflammation. In sum, our analyses confirmed that SERMs are capable of restraining viremia and

mitigating immunopathologies in alphavirus-infected mice.

Discussion

Based on the antiviral effect exhibited by multiple SERMs against other virus families, we

attempted to characterize the anti-alphaviral activity of three SERMs: 4-OHT, tamoxifen, and

clomifene. As evident by plaque reduction assay, RT-PCR studies, and immunostaining, all these

three compounds effectively abrogated the replication of SINV and CHIKV in cell culture.

Results of the time-of addition studies and sequence analysis of partial-4-OHT resistant

phenotype suggested that SERM compounds likely target nsP1 capping enzyme, a component of

the vRNA replication machinery. In the presence of SERMs, a marked difference was observed

between the number of infectious virions released and extracellular vRNA. Capping of vRNA in

the infected cells is orchestrated by a functional alphaviral nsP1 and inhibition of nsP1 by SERM

compounds would release a significant number of noncapped virus particles. The discrepancy in

the magnitude of effect on extracellular vRNA and infectious units released is supported by

previous studies which demonstrated that noncapped vRNAs are infectious but at significantly

lower levels as compared to the capped RNAs (Figure 6) (37, 38). Though, SINV RNA has been

shown to undergo translation and encapsidation without a cap structure, noncapped vRNAs are

found to be highly unstable (38). Concordantly, the enzymatic activity of CHIKV nsP1 and

VEEV nsP1 was inhibited by all the three compounds tested in the MTase activity assay. A

secondary orthogonal ELISA-based assay also validated the anti-nsP1 activity of the tested

SERM compounds.

Replicon assay using SINV-REP which does not involve viral entry or release indicated that

these SERMs interfere with the viral replication machinery and subsequently affect viral protein

synthesis. The data from the SINV-FLuc virus also suggested a decrease in vRNA replication

and/or translation when the infected cells were treated with these compounds. The observed

decrease in RNA translation can be a direct consequence of the reduction in capping activity by

nsP1 or a result of perturbation of RNA replication machinery due to inhibitor binding to nsP1.

We further demonstrated that these compounds marginally suppress vRNA replication in

infected cells when vRNA was quantified 6 hpi. Apart from vRNA capping, nsP1 has also been

found to be involved in the synthesis of negative strand of vRNA (39, 40). However, we cannot

rule out the possibility that reduction of vRNA levels or subgenomic translation is due to the

degradation of noncapped vRNAs (generated because of the compound-mediated inhibition of

nsP1) by host RNA decay machinery or plausible modulation of nsP1 interaction with other nsPs

due to inhibitor binding.

Besides engaging estrogen receptor signaling, SERMs also modulate numerous other cellular

factors and functions; they inhibit immune-regulatory NF-κB signaling, repress the protein

kinase C activity, control the calmodulin pathway, and restrict acidification of lysosomes and

endosomes (36, 41–43). Previous studies established broad-spectrum antiviral activities of

SERMs against a number of human pathogens. Interestingly, antiviral activities of SERMs were

found to be mostly independent of estrogen receptors and involved other cellular or viral targets

(14, 20, 23–25, 44, 45). For example, tamoxifen inhibited the binding and penetration of HSV-1

by targeting the chloride channels (46). A recent study revealed that tamoxifen and clomiphene

could suppress the spike protein-induced membrane fusion which is an important early step for

severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection (47). Endolysosomal

calcium accumulation upon treatment with SERMs obstructed cellular entry by the Ebola virus

(45). On the other hand, tamoxifen diminished VSV infection wherein the inhibition was

mediated by type-1 interferon (25). Interestingly in flaviviruses, the target for SERMs are the

RNA translation and/or replication of the virus (20). In this study anti-CHIKV studies were

performed using Vero cells, which lack the expression of estrogen receptor α and produce only a

low level of estrogen receptor β (23). Of note, Vero cells also do not secrete type-1 interferon.

However, this study has not ruled out that SERMs might influence other host pathways or target

other viral proteins of the alphaviruses.

In our mouse infection studies, the therapeutic application of tamoxifen efficiently restricted

CHIKV propagation and persistence *in vivo*. We further noticed a modest but significant

improvement of the footpad swelling in CHIKV-infected mice administered with a moderate

dose of tamoxifen (20 mg/kg). Local inflammation triggered upon CHIKV infection orchestrates

the influx of leukocytes, including macrophages and CD4 T cells, in the infected tissue leading

to musculoskeletal pathologies (29). Importantly, alphaviruses engage the cellular NF-κB

signaling pathway, which promotes inflammatory gene expressions (48, 49). It has been recently

shown that besides promoting inflammation, NF-κB signaling also augments replication of SINV

in infected neurons (49). Curiously, tamoxifen, independent of estrogen receptor signaling, also

potently suppresses the NF-κB pathway and restrains inflammation (36, 50). It is also possible

that SERMs relied, at least in part, on the down-modulation of NF-κB signaling for mitigating

alphavirus virus-induced disease pathologies. Taken together, we argue that the therapeutic

effect of tamoxifen observed in our preclinical model could have been attributed to its both anti-

nsP1 activity and generic anti-inflammatory functions. Though we have demonstrated nsP1

inhibition by SERM compounds as a probable mechanism for alphaviral inhibition by the

compounds, further studies are warranted to fully dissect the contribution of host factors and

viral proteins in generating anti-alphaviral effects of SERMs.

While our results assure that adaptive mutations may necessarily not pose a threat to the

translation potential of SERM regimes, future studies ought to more extensively characterize the

SERM-alphavirus interaction interface. There is a possibility that adaptive mutations at multiple

sites are required to circumvent the inhibitory effect of the compounds. Though our *in vitro*

enzymatic assays confirmed inhibition of enzymatic activity of purified CHIKV and VEEV nsP1

proteins, computational analysis of protein-ligand binding (including extensive molecular

dynamic simulations) and site-directed mutagenesis combined with *in vitro* activity studies will

be required to accurately assess the nsP1-SERM interaction and elucidate the binding

preferences of similar compounds to the alphaviral capping enzyme. Nonetheless, our current

work offers a proof of concept that SERM compounds can act as direct acting antivirals by

impairment of alphaviral nsP1 thereby providing another estrogen receptor signalling-

independent mechanism for antiviral effect of SERMs.

In conclusion, this report identifies and characterizes SERMs as promising lead candidates for

further development of targeted anti-alphaviral therapy. Dose-dependent inhibition of CHIKV

and SINV by all three tested SERMs, and inhibition of CHIKV nsP1 and VEEV nsP1 enzymatic

activity affirms the broad-spectrum antiviral activity of SERMs against alphaviruses. A number

of SERMs are FDA-approved, have good oral availability, established pharmacological and

safety profiles, and are already in clinical practice for half a century. With the effective inhibition

of alphaviral replication in cell culture and mice model by SERM compounds, a possible new

approach of using SERMs and their superior analogs can be further investigated for the

identification and development of pan-alphavirus therapeutic agents.

Acknowledgements

We thank Dr. Richard Kuhn at Purdue University, USA for providing SINV-FLuc plasmid.

R.M., S.C., and M.A. are thankful to council of scientific and industrial research, India. A.D.

thanks Department of Biotechnology, India (DBT), and Y.R. thanks Indian Council for Medical

Research (ICMR), Govt. of India for research fellowship. Authors thank Biotech Consortium

India Limited, DBT, Ministry of Science and Technology, Government of India, New Delhi,

India for their assistance in filing the patent for "An antiviral composition against

alphaviruses" (Patent file no. 201911049130, dated 29th November 2019).

Funding

This work was supported by a research grant ISRM/12(46)/2020 to S.T. from ICMR. Virus

research in the S.B.'s laboratory is funded by the S.N. Ramachandran National Bioscience

Award for Career Development award to S.B. by DBT, Govt. of India, and NII-core.

Transparency declarations

None to declare.

References:

1. Mostafavi H, Abeyratne E, Zaid A, Taylor A. 2019. Arthritogenic Alphavirus-Induced

Immunopathology and Targeting Host Inflammation as A Therapeutic Strategy for

Alphaviral Disease. 3. Viruses 11:290.

2. Strauss JH, Strauss EG. 1994. The alphaviruses: gene expression, replication, and

evolution. Microbiol Rev 58:491–562.

3. Pialoux G, Gaüzère B-A, Jauréguiberry S, Strobel M. 2007. Chikungunya, an epidemic

arbovirosis. Lancet Infect Dis 7:319–327.

4. Soumahoro M-K, Boelle P-Y, Gaüzere B-A, Atsou K, Pelat C, Lambert B, Ruche GL,

Gastellu-Etchegorry M, Renault P, Sarazin M, Yazdanpanah Y, Flahault A, Malvy D,

Hanslik T. 2011. The Chikungunya Epidemic on La Réunion Island in 2005–2006: A Cost-

of-Illness Study. PLoS Negl Trop Dis 5:e1197.

5. Adouchief S, Smura T, Sane J, Vapalahti O, Kurkela S. 2016. Sindbis virus as a human

pathogen-epidemiology, clinical picture and pathogenesis. Rev Med Virol 26:221–241.

6. Strauss EG, Rice CM, Strauss JH. 1984. Complete nucleotide sequence of the genomic

RNA of Sindbis virus. Virology 133:92–110.

7. Rupp JC, Sokoloski KJ, Gebhart NN, Hardy RW. 2015. Alphavirus RNA synthesis and

non-structural protein functions. J Gen Virol 96:2483–2500.

8. Ahola T, Kääriäinen L. 1995. Reaction in alphavirus mRNA capping: formation of a

covalent complex of nonstructural protein nsP1 with 7-methyl-GMP. Proc Natl Acad Sci

92:507–511.

9.   Hyde JL, Diamond MS. 2015. Innate immune restriction and antagonism of viral RNA

lacking 2′-O methylation. Virology 0:66–74.

10. Tomar S, Narwal M, Harms E, Smith JL, Kuhn RJ. 2011. Heterologous production,

purification and characterization of enzymatically active Sindbis virus nonstructural protein

nsP1. Protein Expr Purif 79:277–284.

11. Feibelman KM, Fuller BP, Li L, LaBarbera DV, Geiss BJ. 2018. Identification of small

molecule inhibitors of the Chikungunya virus nsP1 RNA capping enzyme. Antiviral Res

154:124–131.

12.   Inhibition of Chikungunya virus by an Adenosine Analog targeting the SAM☐dependent

nsP1 methyltransferase - Mudgal - - FEBS Letters - Wiley Online Library.

13. Pushpakom S, Iorio F, Eyers PA, Escott KJ, Hopper S, Wells A, Doig A, Guilliams T,

Latimer J, McNamee C, Norris A, Sanseau P, Cavalla D, Pirmohamed M. 2019. Drug

repurposing: progress, challenges and recommendations. 1. Nat Rev Drug Discov 18:41–

58.

14. Montoya MC, Krysan DJ. 2018. Repurposing Estrogen Receptor Antagonists for the

Treatment of Infectious Disease. mBio 9:e02272-18.

15. Plouffe L. 2000. Selective estrogen receptor modulators (SERMs) in clinical practice. J Soc

Gynecol Investig 7:S38-46.

16. Legha SS. 1988. Tamoxifen in the Treatment of Breast Cancer. Ann Intern Med 109:219–

228.

17.   Squirewell EJ, Qin X, Duffel MW. 2014. Endoxifen and Other Metabolites of Tamoxifen

224   Inhibit Human Hydroxysteroid Sulfotransferase 2A1 (hSULT2A1). Drug Metab Dispos

225   42:1843–1850.

226   18.   Homburg R. 2005. Clomiphene citrate—end of an era? a mini-review. Hum Reprod

227   20:2043–2051.

228   19.   Wibowo E, Pollock PA, Hollis N, Wassersug RJ. 2016. Tamoxifen in men: a review of

229   adverse events. Andrology 4:776–788.

230   20.   Eyre NS, Kirby EN, Anfiteatro DR, Bracho G, Russo AG, White PA, Aloia AL, Beard MR.

231   2020. Identification of Estrogen Receptor Modulators as Inhibitors of Flavivirus Infection.

232   Antimicrob Agents Chemother 64:e00289-20, /aac/64/8/AAC.00289-20.atom.

233   21.   Watashi K, Inoue D, Hijikata M, Goto K, Aly HH, Shimotohno K. 2007. Anti-hepatitis C

234   virus activity of tamoxifen reveals the functional association of estrogen receptor with viral

235   RNA polymerase NS5B. J Biol Chem 282:32765–32772.

236   22.   Tohma D, Tajima S, Kato F, Sato H, Kakisaka M, Hishiki T, Kataoka M, Takeyama H, Lim

237   C-K, Aida Y, Saijo M. 2019. An estrogen antagonist, cyclofenil, has anti-dengue-virus

238   activity. Arch Virol 164:225–234.

239   23.   Johansen LM, Brannan JM, Delos SE, Shoemaker CJ, Stossel A, Lear C, Hoffstrom BG,

240   DeWald LE, Schornberg KL, Scully C. 2013. FDA-approved selective estrogen receptor

241   modulators inhibit Ebola virus infection. Sci Transl Med 5:190ra79-190ra79.

24.   Zheng K, Chen M, Xiang Y, Ma K, Jin F, Wang X, Wang X, Wang S, Wang Y. 2014.

Inhibition of herpes simplex virus type 1 entry by chloride channel inhibitors tamoxifen and

NPPB. Biochem Biophys Res Commun 446:990–996.

25.   Cham LB, Friedrich S-K, Adomati T, Bhat H, Schiller M, Bergerhausen M, Hamdan T, Li

F, Machlah YM, Ali M, Duhan V, Lang KS, Friebus-Kardash J, Lang J. 2019. Tamoxifen

Protects from Vesicular Stomatitis Virus Infection. Pharmaceuticals 12:142.

26.   Singh H, Mudgal R, Narwal M, Kaur R, Singh VA, Malik A, Chaudhary M, Tomar S.

2018. Chikungunya virus inhibition by peptidomimetic inhibitors targeting virus-specific

cysteine protease. Biochimie 149:51–61.

27.   Kaur R, Neetu null, Mudgal R, Jose J, Kumar P, Tomar S. 2019. Glycan-dependent

chikungunya viral infection divulged by antiviral activity of NAG specific chi-like lectin.

Virology 526:91–98.

28.   Kaur R, Mudgal R, Narwal M, Tomar S. 2018. Development of an ELISA assay for

screening inhibitors against divalent metal ion dependent alphavirus capping enzyme. Virus

Res https://doi.org/10.1016/j.virusres.2018.06.013.

29.   Gardner J, Anraku I, Le TT, Larcher T, Major L, Roques P, Schroder WA, Higgs S,

Suhrbier A. 2010. Chikungunya virus arthritis in adult wild-type mice. J Virol 84:8021–

8032.

30.   Teo T-H, Chan Y-H, Lee WWL, Lum F-M, Amrun SN, Her Z, Rajarethinam R, Merits A,

Rötzschke O, Rénia L, Ng LFP. 2017. Fingolimod treatment abrogates chikungunya virus–

induced arthralgia. Sci Transl Med 9.

31. Agarwal A, Singh AK, Sharma S, Soni M, Thakur AK, Gopalan N, Parida MM, Rao PVL,

Dash PK. 2013. Application of real-time RT-PCR in vector surveillance and assessment of

replication kinetics of an emerging novel ECSA genotype of Chikungunya virus in Aedes

aegypti. J Virol Methods 193:419–425.

32. Sourisseau M, Schilte C, Casartelli N, Trouillet C, Guivel-Benhassine F, Rudnicka D, Sol-

Foulon N, Roux KL, Prevost M-C, Fsihi H, Frenkiel M-P, Blanchet F, Afonso PV, Ceccaldi

P-E, Ozden S, Gessain A, Schuffenecker I, Verhasselt B, Zamborlini A, Saïb A, Rey FA,

Arenzana-Seisdedos F, Desprès P, Michault A, Albert ML, Schwartz O. 2007.

Characterization of Reemerging Chikungunya Virus. PLoS Pathog 3.

33. Rana J, Rajasekharan S, Gulati S, Dudha N, Gupta A, Chaudhary VK, Gupta S. 2014.

Network mapping among the functional domains of Chikungunya virus nonstructural

proteins. Proteins 82:2403–2411.

34. Sreejith R, Rana J, Dudha N, Kumar K, Gabrani R, Sharma SK, Gupta A, Vrati S,

Chaudhary VK, Gupta S. 2012. Mapping interactions of Chikungunya virus nonstructural

proteins. Virus Res 169:231–236.

35. Žusinaite E, Tints K, Kiiver K, Spuul P, Karo-Astover L, Merits A, Sarand I. 2007.

Mutations at the palmitoylation site of non-structural protein nsP1 of Semliki Forest virus

attenuate virus replication and cause accumulation of compensatory mutations. J Gen Virol

88:1977–1985.

36. Takada Y, Bhardwaj A, Potdar P, Aggarwal BB. 2004. Nonsteroidal anti-inflammatory

agents differ in their ability to suppress NF-κB activation, inhibition of expression of

cyclooxygenase-2 and cyclin D1, and abrogation of tumor cell proliferation. Oncogene

23:9247–9258.

37. Rice CM, Levis R, Strauss JH, Huang HV. 1987. Production of infectious RNA transcripts

from Sindbis virus cDNA clones: mapping of lethal mutations, rescue of a temperature-

sensitive marker, and in vitro mutagenesis to generate defined mutants. J Virol 61:3809–

3819.

38. Sokoloski KJ, Haist KC, Morrison TE, Mukhopadhyay S, Hardy RW. 2015. Noncapped

Alphavirus Genomic RNAs and Their Role during Infection. J Virol 89:6080–6092.

39. Wang Y-F, Sawicki SG, Sawicki DL. 1994. Alphavirus nsP3 functions to form replication

complexes transcribing negative-strand RNA. 10. J Virol 68:6466–6475.

40. Wang YF, Sawicki SG, Sawicki DL. 1991. Sindbis virus nsP1 functions in negative-strand

RNA synthesis. J Virol 65:985–988.

41. Wiseman H, Laughton MJ, Arnstein HR, Cannon M, Halliwell B. 1990. The antioxidant

action of tamoxifen and its metabolites. Inhibition of lipid peroxidation. FEBS Lett

263:192–194.

42. O'Brian CA, Housey GM, Weinstein IB. Specific and Direct Binding of Protein Kinase C

to an Immobilized Tamoxifen Analogue 5.

43.   Tamoxifen inhibits acidification in cells independent of the estrogen receptor | PNAS.

44.   Murakami Y, Fukasawa M, Kaneko Y, Suzuki T, Wakita T, Fukazawa H. 2013. Selective

estrogen receptor modulators inhibit hepatitis C virus infection at multiple steps of the virus

life cycle. Microbes Infect 15:45–55.

45.   Fan H, Du X, Zhang J, Zheng H, Lu X, Wu Q, Li H, Wang H, Shi Y, Gao G, Zhou Z, Tan

D-X, Li X. 2017. Selective inhibition of Ebola entry with selective estrogen receptor

modulators by disrupting the endolysosomal calcium. 1. Sci Rep 7:41226.

46.   Zheng K, Chen M, Xiang Y, Ma K, Jin F, Wang X, Wang X, Wang S, Wang Y. 2014.

Inhibition of herpes simplex virus type 1 entry by chloride channel inhibitors tamoxifen and

NPPB. 4. Biochem Biophys Res Commun 446:990–996.

47.   Zu S, Luo D, Li L, Ye Q, Li R-T, Wang Y, Gao M, Yang H, Deng Y-Q, Cheng G. 2021.

Tamoxifen and clomiphene inhibit SARS-CoV-2 infection by suppressing viral entry.

Signal Transduct Target Ther 6:1–4.

48.   Bais SS, Ratra Y, Khan NA, Pandey R, Kushawaha PK, Tomar S, Medigeshi G, Singh A,

Basak S. Chandipura Virus Utilizes the Prosurvival Function of RelA NF-κB for Its

Propagation. J Virol 93:e00081-19.

49.   Yeh JX, Park E, Schultz KLW, Griffin DE. NF-κB Activation Promotes Alphavirus

Replication in Mature Neurons. J Virol 93:e01071-19.

50.   Behjati S, Frank MH. 2009. The effects of tamoxifen on immunity. Curr Med Chem

16:3076–3080.

 Figure Legends

 Figure 1: Primary evaluation of anti-alphaviral activity of SERMs. (A) Schematic diagram

 of SINV-FLuc (top panel) wherein FLuc is placed downstream of an internal ribosomal entry site

 (IRES). BHK-21 were infected with SINV-FLuc at an MOI of 0.1 and treated with SERM

 compounds or 0.1% DMSO. Luciferase activity of the cell lysate was measured 12 hpi using

 firefly luciferase assay kit. Values are mean and error bars are standard deviation (n=3). FLuc:

 Firefly luciferase; hpi: hours post-infection. Statistical significance of the difference in luciferase

 activity levels between treated and vehicle control-treated cells was assessed by one-way

 ANOVA test and Dunnett's posttest. ***, $p<0.001$; **, $p<0.01$. Values are mean and error bars

 are standard deviation (n=3). (B) Plaque assay revealing changes in CHIKV (top three panels) or

 SINV (bottom three panels) titer on the treatment of virus-infected cells with an increasing

 concentration of 4-OHT, tamoxifen, and clomifene. The viral titer in the culture supernatant was

 determined at 24 hpi. The viral titer in corresponding vehicle-treated cells was considered 100%.

 Data were normalized using Graph pad's non-linear regression curve fit and the calculated 50%

 effective concentration (EC_{50}) values are summarized in Table 1. Values are mean from three

 independent experiments and error bars are standard deviation.

 Figure 2: Antiviral activity of SERMs against CHIKV and SINV in cell culture. (A)

 Barplots showing fold-reduction in the abundance of vRNA in CHIKV or SINV infected Vero

 cells treated with SERMs compared to vehicle control (VC). At 24 hpi, RNA was isolated and

 analyzed by qRT-PCR for the expression of the E1 region of the viral genome. Statistical

 significance of the difference between intracellular vRNA in treated cells and vehicle control-

 treated cells was assessed by one-way ANOVA test and Dunnett's posttest.***, $p<0.001$. Values

 are mean and error bars are standard deviation (n=3). (B) Immunofluorescence microscopy of

CHIKV infection in Vero cells upon treatment with SERMs. Virus infection and treatment with

SERMs were performed as mentioned in the supporting information. At 36 hpi, the infected cells

were fixed and stained using an anti-alphavirus primary antibody and FITC-conjugated

secondary antibody. DAPI was used for counter-staining cell nuclei. The image represents three

independent biological replicates. Bars, 200 µM.

Figure 3: Characterizing SERM-mediated inhibition of viral growth. (A) Schematic

representation of the time course of viral infection and treatment with SERM compounds. Vero

cells were treated with compounds at different time points (in hours) before, during, or after

virus infection and the viral titer (PFU/ml) was determined for CHIKV (B and C) and SINV (B

and D). For Cotreatment studies, compounds were added simultaneously with the virus

inoculum. For pre-treatment assay, Vero cells were treated with the inhibitors 2 h prior to viral

adsorption. For post-treatment assay, cells were incubated at 37°C with viral infection medium

which was replaced by maintenance medium at different time-points containing 5 µM 4-OHT or

7.5 µM tamoxifen/clomifene. Supernatants were collected 24 hpi for the quantification of viral

titer by plaque assay. 0.1% DMSO was used as vehicle control. For all the experimental groups,

Vero cells were infected with CHIKV or SINV at an MOI of 1. VC: vehicle control; hpi: hours

post-infection; PFU: plaque-forming unit. (E) SINV minigenome replicon assay capturing the

luciferase activity in lysate derived from BHK21 cells transfected with capped RNA encoding

the SINV-REP reporter replicon. Cells were subjected to 6 h of treatment with VC or 2.5 µM of

the indicated SERMs. A schematic diagram of SINV-REP has also been presented (top). (F and

G) Changes in vRNA levels in CHIKV and SINV infected cells at 6 hpi. Vero cells were infected

with CHIKV and SINV at MOI of 5 followed by treatment with 5 µM of the compounds. vRNA

levels were assessed by qRT-PCR and the fold reduction in vRNA relative to VC-treated cells is

plotted. Actin: endogenous control; E1: target gene. (H) Quantitative determination of

extracellular SINV vRNA levels at 24 hpi. Vero cells were infected with SINV at MOI of 1 and

were treated with indicated concentrations of the SERM compounds. Quantification of the

vRNA in the culture supernatant was done using qRT-PCR as discussed in the material and

methods section. Statistical significance of the difference in viral titer/luciferase activity/vRNA

levels between treated and vehicle control-treated cells was assessed by one-way ANOVA test

and Dunnett's posttest. ***, $p<0.001$; **, $p<0.01$; *, $p<0.05$. Values are mean and error bars are

standard deviation from triplicate experiments.

Figure 4: nsP1 capping enzyme is implicated in alphavirus inhibition by SERMs. Altered

plaque phenotype of the partially-resistant variant of CHIKV against 4-OHT. (A) Standard

plaque assay performed on the indicated virus. Vero cells were overlaid with 1% carboxymethyl

cellulose-containing medium either without or with 5 µM 4-OHT. Plaques were visualised 48

hpi using crystal violet. Plaque-purified variant harboring all four mutations was used for

phenotypic characterization. The resistant variant replicates more efficiently than wild-type

CHIKV both in terms of number and size of plaques in the presence of 4-OHT. (B) Non-

synonymous mutations in the nsP-coding region of partially-resistant CHIKV variant. Mutations

were identified with the sequencing of three independent plaque-purified viruses. nsP: non-

structural proteins. (C) nsP1 MTase activity inhibition by tested compounds. SERM compounds

were tested at 50 µM concentration in the CHIKV nsP1 (10 µM) and VEEV nsP1 (20 µM)

MTase reaction mixture. The reaction was carried out as described previously by Mudgal et al.,

2020. The MTase activity is shown relative to the enzyme activity in the reaction mixture in the

presence of vehicle control. (D) Dose-dependent inhibition of guanylation of nsP1 in the

presence of SERMs. Relative guanylation by CHIKV nsP1 was determined using an ELISA

assay that measures the amount of m^7GMP-nsP1 adduct formed in the reaction. Reaction

conditions were similar to those described earlier by Kaur et al., 2018. Data points represent

mean value and error bars are standard deviation from two independent experiments.

Figure 5: Testing therapeutic efficacy of tamoxifen in CHIKV-infected mice. CHIKV-

infected mice were injected successively with 20 mg/kg of tamoxifen or vehicle control at day

one, three, and five post-infection. Subsequently, changes in the footpad area (A) was measured

at the indicated days. (B) Representative image of CHIKV infected joint footpad of mice

subjected to tamoxifen therapeutic regime. (C) Dot plots showing the load of infectious virus

particles in the footpad at the indicated days post-infection, as determined by $TCID_{50}$ method.

Graphs revealing the abundance of vRNA in the blood (D) and the infected footpad (E) at the

indicated day post-infection. (F) Representative immunohistochemistry images revealing H&E-

stained sections of the joint footpad. Mice were either uninfected (left panel) or infected with

CHIKV six days prior to image acquisition (middle panel) or infected similarly with CHIKV and

then subjected to a therapeutic regime (right panel). Footpad sections from two animals per set

and three fields/sections were examined. E denotes edema; * denotes infiltrates; B signifies bone.

Images are representative of three mice per group. Data were quantified from the indicated number

of mice from two independent experimental replicates and presented as mean ± SEM. Two-tailed

Student's t-test was performed to assess statistical significance. **P<0.01; *P<0.05.

Figure 6: Possible mechanism of anti-alphaviral action of the SERM compounds mediated

via nsP1 inhibition. vRNA is released in the cytoplasm upon entry of SINV/CHIKV in the

infected cell. Subsequent translation of viral replication proteins generates viral RNA synthetic

complex that in turn synthesizes numerous copies of vRNA. nsP1 (PDB ID: 6z0v) plays a critical

role in 5□ capping of progeny genome. Putative inhibition of nsP1 by SERM compounds viz. 4-

OHT, tamoxifen, and clomifene produces significant amount of noncapped nascent vRNAs that

are encapsidated and subsequently released as mature virus particles. Upon entry into the host

cell cytoplasm, these noncapped vRNAs undergo degradation by host RNA decay machinery

following nucleocapsid disassembly and vRNA release.

A

Concentration of 4-OHT

Wild-type CHIKV | 4-OHT-resistant variant

0 μM

5 μM

B

Non-synonymous mutations	Region	Frequency
A155T	**nsP1**- MTase/GTase domain	100%
H280Q	**nsP3**- Macro Domain	33%
L490M	**nsP3**- Zinc Binding Domain	33%
E39K	**nsP4**- Disordered Region	100%

C

D