1 **Title:**
2 Long-term translocation and hatchery release explain population genetic structure of a
3 recreationally fished iconic species in Japan: A review and reanalysis
4
5

6 Shuichi Kitada
7
8 Tokyo University of Marine Science and Technology, Tokyo 108-8477, Japan

9
10
Short running title: Effects of long-term translocation

Abstract
Ayu (*Plecoglossus altivelis altivelis*) occur widely in China, Korea, Japan, Taiwan and
Vietnam. It is an important freshwater fisheries resource and popular recreational fishing
species in Japan, lives for only one year and has a single breeding season. To supplement
increased recreational fishing demand, huge numbers of wild-born landlocked juvenile have
been translocated from Lake Biwa into most Japanese rivers for more than 50 generations.
Hatchery-born juveniles have also been extensively released for more than 30 generations.
Previous studies reported that landlocked and amphidromous forms of Ayu easily hybridise,
but survival of landlocked larvae is very low in seawater, leading to a general consensus that
the reproductive success of landlocked Ayu is very low, or even 0 in translocated rivers.
However, there is limited information on the reproductive success of landlocked Ayu in
translocated rivers and no study evaluated effects of translocation and hatchery release on
genetic population structure in the distribution range. To provide an up-to-date review on the
key management question, I performed a comprehensive literature search and analysed
published microsatellite data sets collected from the distribution of this species in Japan. The
analyses provide strong evidence for very high gene flow between populations, but population
structure was retained in several regions, and several populations were nested. Genetic
diversity was surprisingly homogeneous. Bayesian admixture analysis estimated hybrid
proportion between both forms to be 37%, uncovered widespread hybridisation in Japanese
rivers. Results are discussed in relation to the conservation and management of this species.
32
33 **Keywords:** amphidromous, Ayu, hatchery, hybrid proportion, landlocked,
34 Lake Biwa

Introduction

Anthropogenic forces often break reproductive barriers, providing an opportunity for hybridisation between closely related species that might not otherwise exist under natural conditions (Grabenstein & Taylor 2018). A geographic area where natural hybridisation occurs is referred to as a hybrid zone, while an area where two divergent populations or closely related species occur and potentially hybridise is called a contact zone (Barton & Hewitt 1985; Hewitt 1988; Johannesson et al. 2020). Hybrid zones work as selective filters, enabling determination of the evolutionary roles of reproductive barriers at various stages of speciation (Barton & Hewitt 1985; Hewitt 1988; Martinsen *et al.* 2001). Repeat backcrossing and recombination of genotypes in hybrid zones results in genetic changes beyond the first hybrid generation, allowing neutral and adaptive alleles to spread among populations (Barton 1979; Barton & Hewitt 1985; Hewitt 1988; Johannesson *et al.* 2020). Hybridisation could also be beneficial to species adaptation in a changing environment, so a case-by-case evaluation is needed to elucidate its consequences (Hirashiki *et al.* 2021).

Translocation of organisms across geographic boundaries has been often used in conservation and management of wildlife; however, because it removes reproductive barriers, and can lead to hybridisation of introduced and resident populations (Seddon *et al.* 2014; Berger-Tal *et al.* 2020; Taylor *et al.* 2021). Translocation can disturb the ecological and evolutionary processes of a species (Conant 1988; Moritz 1995), and unintended, negative outcomes of hybridisation can occur because of altered offspring fitness (Ricciardi & Simberloff 2009; Bucharova 2017). In endangered fish recovery plans, translocations are prevalent, but consequences are often poorly documented (Healy *et al.* 2020). Quantitative evaluation on empirical cases is needed to determine their conservation efficacy and future management efforts (Minckley 1995; George *et al.* 2009; Healy et al. 2020).

Herein, a case study on Ayu (*Plecoglossus altivelis altivelis*), which occur widely in coastal areas of China, Korea, Japan, Taiwan (Kwan *et al.* 2012), and Vietnam (Tran *et al.* 2017). Ayu is an iconic fish species targeted by recreational fishers in Japanese rivers, is presented. Ayu—an essentially amphidromous species, except for some land-locked populations (landlocked form) (Nishida 1985)—can be found throughout Japan, and is one of the most well studied fish species in the country. Landlocked Ayu have been extensively translocated from Lake Biwa into amphidromous populations in rivers throughout Japan since 1970s. In addition, hatchery-born amphidromous fish (including amphidromous and landlocked form hybrids) have been extensively released. Below I briefly review the knowledge of the species and describe the aim of the present study.

Fisheries, translocation of landlocked juveniles, and hatchery supplementation

Ayu is an important fisheries resource in Japanese freshwaters, and a most popular recreational fishing species. Government statistics (e-Stat, https://www.e-stat.go.jp/) indicate that Ayu is caught in almost all prefectures in Japan, with characteristically high catches along the central and southern Pacific coast (Figure S1a). The commercial catch increased markedly to reach a maximum in 1991 but declined thereafter (Figure S1b). While catch statistics exclude recreational catch, this is likely to be substantial, with recreational catch estimated at ~ 5.3×10^6 fish (260 t) over 485×10^3 days (day × fishers) in Nakagawa River, Tochigi Prefecture, alone (Kitada & Tezuka 2002)—a tonnage similar to the commercial catch in the same year in one of the top production areas in Japan (Figure S1a). The fishing season for ayu begins on June 1st and closes at the end of October in this river, which is a typical fishing period in Japan.

Translocation of wild-born landlocked Ayu juveniles from Lake Biwa into Tama River, Tokyo, began in 1912, and this process rapidly expanded throughout Japan in the 1950s and 1970s to supplement increased recreational fisher demand (Takamura 2009). The catch of

wild-born juveniles of the landlocked form in Lake Biwa increased markedly and reached a
maximum in 1991, but this also declined thereafter (Figure S1b). In the 1990s, many local
fisheries cooperatives gradually changed to using hatchery-born amphidromous juveniles
and/or hybrids of the two forms instead of relying on landlocked juveniles for two main
reasons: an outbreak of cold-water disease in Lake Biwa (Izumi & Wakabayashi 1997), and
because landlocked forms translocated into rivers were not expected to reproduce successfully
(e.g., Pastene *et al.* 1991; Seki & Taniguchi 1998; Tago 2004; Takeshima *et al.* 2009).
In recent years, three types of juveniles have been released: hatchery-born
amphidromous fish (including amphidromous and landlocked form hybrids), which have been
released in the largest quantity, followed by wild-born landlocked juveniles from Lake Biwa,
and wild-born amphidromous fish collected in coastal waters (Figure S2a). Most wild-born
juveniles are caught in Shiga Prefecture (the landlocked Ayu collected in Lake Biwa, Figure
S2b), and most hatchery-reared juveniles (hatchery fish) are produced in Tochigi Prefecture,
followed by Miyazaki, Tokushima, Yamaguchi, and Wakayama prefectures, implying that
landlocked juveniles and hatchery fish have been translocated to non-natal rivers.
100
Life history
Ayu live for one year, and have a single breeding season (Nishida 1986). The amphidromous
form of Ayu spawns in the lower reaches of rivers during autumn. Hatched larvae
immediately migrate downstream into brackish waters, then enter the sea where they
overwinter. From spring to early summer, juveniles migrate upstream. In autumn, they mature
and again descend downstream to spawn, then die (Nishida 1986). Ayu in the Lake Biwa
landlocked population—the largest of landlocked populations—spawn in autumn in the lower
reaches of rivers that flow into the lake, after which larvae migrate downstream into the lake
and overwinter. This form is differentiated from amphidromous populations in ecology,
morphology, and molecular characteristics (e.g. Kawanabe 1958; Nishida 1985; Nishida 1986;
Tsukamoto & Kajihara 1987). A previous study using allozymes inferred that genetic distance
between the landlocked form in Lake Biwa and an amphidromous population was correspond
to a divergence time approximately 100 k years before present (Nishida 1985).
114
The reproductive success of translocated Ayu
In the early 1990s, experiments demonstrated that amphidromous and landlocked forms of
Ayu interbred naturally in outdoor ponds, leading to concerns that the fitness of hybrid
progeny would be reduced (Iguchi & Ito 1994). Hybridisation in transplanted populations has
remained a concern for Ayu management and conservation (e.g. Taniguchi *et al.* 1983; Seki *et*
al. 1994; Iguchi *et al.* 1997; Otake *et al.* 2002; Iwata *et al.* 2007). Strontium/calcium ratios of
otoliths suggest that spawning of translocated Ayu from Lake Biwa represents about 20% of
the total spawning of Ayu in natural spawning grounds, with a 35% hybridisation probability
(Otake *et al.* 2002). Genetic stock identification using allozyme markers identified
translocated landlocked Ayu to have mixed in populations, with a high proportion encountered
during the high fishing season (Seki *et al.*1994; Kaewsangk *et al.* 2000; Taniguchi *et al.* 2002),
but a lower proportion in the spawning season. Studies using allozyme and mtDNA markers
have concluded that released landlocked Ayu have little effect on the genetic characteristics of
native populations (Pastene *et al.* 1991; Seki & Taniguchi 1998).
The mixing proportion of the landlocked form in drifting larvae sampled in
September was first estimated using single site of the mtDNA control region as the gene
marker at ~25% in Sho River, Toyama Prefecture (Iwata *et al.* 2007). The study also estimated
that mixing proportions in drifting larvae sampled in October/December and marine juveniles
sampled in February/April. The proportion of larvae of the landlocked form in samples was <
~5% and/or close to 0, with 95% confidence intervals including 0 (Iwata *et al.* 2007),

indicating that Ayu juveniles released from Lake Biwa reproduced in rivers into which they
were stocked, but that the survival rate of progeny was extremely low or 0 in seawater.
Experimental tests on salinity tolerance confirm the survival rates of landlocked-form larvae
to be lower than those of the amphidromous form in seawater (Tabata & Azuma 1986; Iguchi
& Yamaguchi 1994), though salinity tolerance of hybrids between both forms has not been
tested. These results have led to a general consensus among Japanese scientists and hatchery
managers that the reproductive success of landlocked Ayu is very low, or even 0 in
translocated rivers (Takeshima *et al.* 2009). With the exception of Iwata *et al.* (2007), however,
there is no direct evidence for the reproductive success of landlocked Ayu. No study has
identified introgression of landlocked Ayu into amphidromous populations in Japan using
neutral markers in the nuclear genome. Thus, translocations and hatchery releases of Ayu have
regarded as put-and-take fisheries (e.g. Pawson MG 1982; Ross & Loomis 2001; Lorenzen et
al. 2012; Beard & Mann 2020), which provide intra-generational benefits to Ayu commercial
and recreational fisheries during early summer and fall (Tsuboi *et al.* 2019). However,
inter-generational genetic effects of translocations and hatchery releases and its risk
assessment have received little attention (see, e.g. Waples & Drake 2004; Araki & Schmid
2010; Laikre *et al.* 2010; Lorenzen et al. 2012; Christie *et al.* 2014; Taranger *et al.* 2015;
Glover *et al.* 2017; Kitada 2018, references therein).
153
Study aims
Extensive translocation of landlocked Ayu has continued for at least 50 generations (Figure
S1b), with hatchery releases using captive-reared parents for more than 30 generations (Iguchi
et al. 1999; Ikeda *et al.* 2005). Hatchery fish include the amphidromous form, and hybrids of
the amphidromous and landlocked forms. Repeat hybridisation over many generations in
translocated rivers may cause introgression of genes from the landlocked form into
amphidromous populations, leading to reduced fitness of stocked populations because of
lowered survival of progeny of the landlocked form in seawater (Tabata & Azuma 1986;
Iguchi & Yamaguchi 1994). Therefore, inferring hybridisation between the two forms is
central to future management and conservation of this iconic species. However, there is
limited information on the reproductive success of landlocked Ayu in translocated rivers and
no study inferred hybridisation in translocated rivers. Moreover, no study evaluated effects of
translocation and hatchery release on genetic population structure in the distribution range.
To assist with conserving and managing Ayu, the current status of populations is
assessed from a population genetics perspective, and introgression of the landlocked form into
amphidromous populations is inferred throughout its Japanese distribution. Hybridisation and
hybrid individuals in contact zones can be identified by Bayesian clustering methods using
highly polymorphic molecular markers such as microsatellites (Beaumont *et al.* 2001;
Pierpaoli *et al.* 2003; Manel *et al.* 2005; Vähä & Primmer 2006). Using existing (published)
microsatellite data, the population structure, gene flow, geographic distribution of genetic
diversity, and admixture proportions of putative ancestry populations throughout Japan, are
analysed. Using a Bayesian clustering method, magnitudes of introgression of landlocked Ayu
are inferred from hybrid proportions. Results are discussed in relation to the future
conservation and management of this species.
178
Materials and Methods
Genetic data screening
Published genetic data were screened using the Google Scholar search system with keyword
searches of "Ayu", "genetic diversity", "landlocked", "*Plecoglossus*" and "population
structure". Studies that were known to us were also added. The population genetic structure of
Ayu has been studied extensively in Japan. Initially (in the 1980s), molecular markers

included allozymes (Taniguchi *et al.* 1983; Nishida 1985; Seki *et al.* 1988), but with
technological developments this expanded from the late 1990s to include mitochondrial DNA
(mtDNA) (Iguchi *et al.* 1997; Iguchi & Nishida 2000; Iguchi *et al.* 2002; Ikeda & Taniguchi
2002; Takeshima *et al.* 2005) and microsatellite markers (Takagi *et al.* 1999; Ikeda *et al.*
2005; Yamamoto *et al.* 2007; Takeshima *et al.* 2009; Yamaguchi *et al.* 2016) to estimate
genetic characteristics of populations at regional scales. Microsatellite markers have been
recently used to describe this species' population structure throughout its distribution based on
a comprehensive sampling of Japanese rivers (Takeshima *et al.* 2016a). No population
genetics study has used single nucleotide polymorphism (SNPs).
The data screening found that it is the only publicly available one and covered the
distribution range. In this study, the genotypes of Ayu at 12 microsatellite markers collected
throughout Japan were analysed (118 populations, *n*=4,746) (Takeshima *et al.* 2016a,b).
Samples of the amphidromous form comprised upstream migrants and/or marine juveniles
collected from 98 rivers/streams and or coastal waters, and individuals of the landlocked form
collected from 20 rivers flowing into Lake Biwa. The overall independence of the 12
microsatellite loci was confirmed, and all loci were selectively neutral, and deviations from
Hardy–Weinberg equilibrium were observed at the *Pag-070* locus but excluding this locus did
not change results (Takeshima *et al.* 2016a). These data are the most suitable to address the
objectives of this current study; their quality have been checked and they cover the
distribution of this species throughout Japan. Two samples of Ryukyu Ayu (*P. a. ryukyuensis*)
were excluded from the data in the subsequent analysis. In addition, previous results using the
same 12 microsatellite markers for different samples of upstream migrants in the Yodo River
system were analysed, which is the sole outlet to flow from Lake Biwa and the contact zone
(Takeshima *et al.* 2009).
209
Population structure and genetic diversity
Sample site coordinates were retrieved from original studies, but approximate coordinates for
river mouths of sampling sites from Lake Biwa and Toyama Bay were obtained from Google
Maps. Sample locations were plotted on a map using the 'sf' package in R. Genotype data
were converted into Genepop format (Rousset 2008) for implementation in the R package
FinePop2_v0.2 at CRAN. Expected heterozygosity (*He*) for each population was
automatically estimated in the 'read.GENEPOP' function. Genome-wide (averaged over loci)
global F_{ST} values of Weir & Cockerham (1984) were computed using the 'globalFST'
function. Pairwise F_{ST} values from the G_{ST} moment estimator (Nei & Chesser 1983) were
computed using the 'pop_pairwiseFST' function. Sampling locations were connected with
pairwise F_{ST} values smaller than a given criterion by yellow lines to visualise gene flow.
Wright's island model assumes $F_{ST} \approx \frac{1}{4N_e m + 1}$, where N_e is the effective population size and
m is the average migration rate between populations (Slatkin 1987). For example, a criterion
of $F_{ST} = 0.005$ corresponds to $4N_e m \approx 199$ migrants per generation (see Whitlock &
Mccauley 1999; Waples & Gaggiotti 2006). An unrooted neighbour-joining (NJ) tree (Saitou
& Nei 1987) was created for the pairwise F_{ST} distance matrix using the 'nj' function in the R
package 'ape.' The performance of the F_{ST} analysis applied here is evaluated by simulations
and empirical data analysis (Kitada *et al.* 2021).
Sample points were visualised using a colour gradient of populations *He* values. The
colour of population *i* was rendered as RGB $(1 - H_{e0,i}, 0, H_{e0,i})$, where $H_{e0,i} = (H_{e,i} -$
$\min H_e)/(\max H_e - \min H_e)$. This conversion represents the standardised magnitude of an *He*
value at the sampling point, with a continuous colour gradient ranging from blue (lowest) to
red (highest) for *He* values. Populations were grouped into nine geographic areas: Hokkaido,
northern Sea of Japan, western Sea of Japan, northern Pacific, central Pacific, southern Pacific,

Kyushu (facing the East China Sea), and Lake Biwa. Differences in average *He* values
between all pairs of the nine groups were tested by one-way ANOVA using the 'anova'
function, with the *p*-values being adjusted for multiple comparisons with a 5% significance
using the 'TukeyHSD' function in R.
238
Admixture proportions
STRUCTURE (Pritchard *et al.* 2000) and NEWHYBRIDS (Anderson & Thompson 2002)
have been applied to detect hybrids in populations (Vähä & Primmer 2006). STRUCTURE
assigns probabilities for individuals to have *K* putative ancestry populations (individual
q-values), while NEWHYBRIDS estimates the probability of individuals belonging to distinct
hybrid or purebred classes. Both methods produced mostly consistent results when applying
empirical data with 14 microsatellite markers (Steyer *et al.* 2018). Here, the Ayu dataset of 12
microsatellites was analysed using STRUCTURE. Genotype data in Genepop format were
converted into STRUCTURE format using PGDSpider v2.1.1.5 (Lischer & Excoffier 2012).
STRUCTURE 2.3.4 was run with no prior information about sampling locations (Pritchard *et*
al. 2000), assuming admixture and corelated allele frequencies between populations (Falush
et al. 2003). A burn-in of 50,000 sweeps followed by 100,000 MCMC sweeps was used for
the number of original populations *K* = 1–5 with 10 runs for each *K*. Longer runs with
500,000 MCMC sweeps following a burn-in period of 100,000 sweeps were initially tested,
but very similar results were obtained, and shorter sweep length were used. The most likely *K*
was estimated based on $\ln(P|D)$ for each *K*. A threshold *q*-value of 0.1 (Vähä & Primmer
2006; Devitt *et al.* 2011; Fogelqvist *et al.* 2015) and/or 0.2 (Geraldes *et al.* 2014;
Karamanlidis *et al.* 2021; Larison *et al.* 2021) has been used often to identify hybrids of two
species and/or distinct populations; a 0.20 threshold *q*-value is used herein, where individuals
with landlocked ancestral genes ($0.8 > q > 0.2$) are classified as hybrids, those with $q \geq 0.8$ as
the purebred landlocked form, and those for which $q \leq 0.2$ as the purebred amphidromous
form (Vähä & Primmer 2006).
261
Isolation-by-distance
Isolation-by-distance relationships between populations are examined. Great circle geographic
distances between all pairs of populations were computed using the 'distm' function in the
geosphere R package, and the linear relationship between geographic distances and genetic
distances (as measured by pairwise F_{ST}) was estimated using the 'lm' function in R. The
landlocked Ayu have been translocated form Lake Biwa into almost all rivers and major seed
producers have sold hatchery-born juveniles to other prefectures (Figure S2b), waypoints
(Ramachandran *et al.* 2005; Kitada *et al.* 2017), which account for natural migration paths
were not assumed. A Mantel correlation test was performed using the 'mantel' function in
vegan R package.
272
Results
Sampling locations were plotted on a map (Figure S3). The global F_{ST} (± standard error) was
0.0162 (± 0.0074) for all populations, but after excluding populations from Lake Biwa, this
dropped to 0.0058 ± 0.0038. The mean of pairwise F_{ST} was 0.0088 ± 0.0099 (SD) for all
population pairs. Using the criterion of pairwise $F_{ST} < 0.005$ ($4N_e m \approx 199$), substantial gene
flow between populations was visualised, with very high gene flow between populations and
geographic areas (Figure 1a).

280
281
282 **Figure 1** Ayu population structure throughout Japan. (a) Map visualising geneflow between
283 populations using 12 microsatellite genotypes from 118 populations ($n = 4746$). Yellow lines
284 connect population pairs with pairwise $F_{ST} < 0.005$. (b) Unrooted neighbour-joining (NJ) tree
285 based on the pairwise F_{ST} matrix. Numbers: 67, Tone River, and 98, Yodo River (see Table
286 S1 and Figure S3 for sampling information)
287

288
289
Figure 2 Expected heterozygosity (*He*) of Ayu throughout Japan. (a) Geographic distributions
of population-specific *He* averaged over 12 microsatellite loci in 118 populations ($n = 4746$).
The colour of each population reflects the magnitude of *He* values, with a continuous colour
gradient ranging from blue to red for the lowest and highest *He* values, respectively (see text).
(b) Geographical distribution of *He*. Differences between areas ($p < 0.05$) are indicated by
different lowercase letters
296
The unrooted NJ tree (Figure 1b, Figure S4) indicates clear differences between

amphidromous and landlocked forms, with a sample from Yodo River (#98) included in the
landlocked cluster. Tone River (#67) was located slightly away from the amphidromous
cluster in a direction towards the landlocked cluster. Hokkaido and the northern Sea of Japan
group fall into one cluster, and Kyushu and the southern Pacific form into another.
Populations from the western Sea of Japan, northern Pacific, central Pacific, and Seto Inland
Sea were located relatively close to each other in the centre of the tree and retained regional
characteristics, with a number of nested populations. Contrarily, the NJ tree (Figure S5)
created using the F_{ST} matrix (Takeshima *et al.* 2009) located upstream migrants between Lake
Biwa and amphidromous populations, being consistent with the original study.
The mean ± standard deviation (SD) of *He* values was 0.66 ± 0.013, with a very
narrow range between 0.62 and 0.68. No clear pattern in geographic distribution of *He* was
apparent (Figure 2a). Excluding Lake Biwa, *He* values were homogeneous (Figure 2b);
average *He* values in the northern and western Sea of Japan, and northern and central Pacific
were significantly higher than those for Lake Biwa, while those in Hokkaido, the southern
Pacific, Kyushu, and the Seto Inland Sea did not differ from those in Lake Biwa.
313

314
315
316 **Figure 3** STRUCTURE analyses. (a) Coloured bars of individual *q*-values illustrate
317 admixture proportions of inferred ancestries at *K* = 3 (*n* = 4746). Numbers in parentheses refer
318 to sample sizes. (b) Geographic distribution of admixture proportions in 118 populations
319
STRUCTURE suggests the most likely *K* to be 3 for all replicate runs (Figure S6). A
bar plot of individual *q*-values for *K* = 3 (Figure 3a) depicts Ayu in Japan to have three
common putative ancestral origins, with similar admixture patterns within amphidromous and
landlocked populations. Blue and green colours refer to putative amphidromous ancestries,
and orange to the landlocked one. The amphidromous population in Yodo River has a similar
admixture pattern to that of Lake Biwa. Figure 3b depicts the geographic distribution of

admixture patterns. Box plots of individual q-values of landlocked ancestry in 118
populations illustrate that amphidromous and landlocked populations to have similar patterns
with outliers (Figure S7); the average of individual q-values for amphidromous populations is
0.21, and 0.79 for landlocked ones. The geographic distribution of proportions of hybrid,
purebred landlocked, and amphidromous forms using the threshold q-value of 0.2 (Figure 4,
Table S1) reveals hybrids to have expanded into all amphidromous populations with mean ($\pm$
SD) proportion of 37% $\pm$ 10%. A maximum 60% of individuals were hybrid in Tone River
(#67), and a minimum of 15% in Sho River (#30). The average proportion of purebred
landlocked individuals was $1 \pm 6\%$, and that of purebred amphidromous individuals $62 \pm 12\%$.
In Yodo River, the proportion of hybrids was 39%, and that of the purebred landlocked form
61.0%, with no purebred amphidromous individuals. A similar result was obtained for Lake
Biwa, where the proportion of purebred landlocked individuals was $62 \pm 8\%$, while that of
hybrids was $38 \pm 8\%$; similarly, no purebred amphidromous individuals were recognised.
339

340
341 **Figure 4** Inferred proportions of hybrid and purebred classes in 118 populations ($n = 4746$).
342 Individuals with landlocked ancestral genes ($0.8 > q > 0.2$) ae classified as hybrids; those with
343 $q \geq 0.8$ ae classified as purebred landlocked form, and $q \leq 0.2$ as purebred amphidromous
344 form (Table S1). Numbers: 30, Sho River; 67, Tone River (see text)
345
The relationship between great circle geographic distances and genetic distances as
measured by pairwise F_{ST} is depicted in Figure 5. The linear regression of F_{ST} on geographic
distance produced an isolation-by-distance relationship within amphidromous populations
(black dots). The Mantel correlation was significant ($r = 0.32$, $p = 0.001$), and R^2 was 0.10.
Contrarily, the scatter plot (coloured dots) for Lake Biwa and amphidromous populations
indicates that the linear relationship between geographic distance and genetic distance was

disrupted. Many amphidromous populations in the
within-amphidromous-population-comparison are overlapped by those in the between-Lake
Biwa-comparison.
355

356
Figure 5 Scatterplot of geographic distance and pairwise F_{ST}. Black dots depict
within-amphidromous-populations comparison, with a regression line [$\hat{F}_{ST} = 1.518 \times$
$10^{-3} + 4.184 \times 10^{-6} \times$ (geographic distance)] in red, with R^2 0.01; the Mantel correlation
between F_{ST} and geographic distance is 0.32 ($p = 0.001$). Orange dots depict the samples from
within the Lake Biwa population; other colours are for between Lake Biwa and
amphidromous populations comparison
363

Discussion

Reanalysis of published microsatellite data for Ayu throughout its distribution in Japan
presents strong evidence for very high gene flow between populations and regions, and for
population structure to be retained in several regions, with some nested populations apparent
(Figures 1, S4). Genetic diversity in terms of *He* was surprisingly homogeneous (Figure 2),
and admixtures of landlocked and amphidromous ancestral populations occurred throughout
Japan (Figure 3). Analyses also reveal that hybrid individuals have expanded their distribution
throughout Japan, with a relatively high mean hybrid proportion (37%) in populations (with
mean values for the purebred amphidromous form (62%), and purebred landlocked (1%) in

amphidromous populations (Figure 4, Table S1)). In the Lake Biwa population, the estimated
purebred landlocked form accounts for 62% of individuals and the hybrid proportion 38%,
with no purebred amphidromous individuals. A very similar result to Lake Biwa was found in
Yodo River. The linear regression of pairwise F_{ST} on geographic distance produced an
isolation-by-distance relationship within amphidromous populations, but many
amphidromous populations were overlapped by those in the between-Lake Biwa-comparison.
Contrarily, an isolation-by-distance relationship was disrupted in the comparison of Lake
Biwa and amphidromous populations.
381
Population structure and effects of translocations and hatchery releases
Analysed sample data was obtained from upstream migrant and/or marine juvenile Ayu
(Takeshima *et al.* 2016a). With the exception of the sample from Yodo River, no
amphidromous population was included in the cluster of Lake Biwa, and the proportions of
the purebred landlocked form were 0 in most amphidromous populations (Figures 1, 4, Table
S1). This indicates that the probability of samples containing released landlocked juveniles is
very low.
The population structure of Ayu was maintained in Hokkaido, the northern Sea of
Japan, Kyushu, and the southern Pacific (Figure 1a), despite very high gene flow between
populations and areas. Juvenile Ayu spend months at sea over winter (Nishida 1986), and drift
in sea currents and/or move to natal and adjacent river mouths (Iguchi *et al.* 2006). Juveniles
commence their upstream migration in spring at ~ 60 mm (Tsukamoto & Kajihara 1987);
those individuals that stray, that do not return to natal rivers, contribute to gene exchange
among rivers. Natural recruitment of amphidromous Ayu depends primarily on the coastal
environment, particularly that of estuaries (Takahashi *et al.* 1998; Tago 2002). Long-range
dispersal to offshore waters might reduce survival of marine juveniles, resulting in
environmental-induced natal river fidelity (Iguchi *et al.* 2006). Temporally stable and
differentiated structures with high gene flow were reported for Atlantic cod (*Gadus morhua*)
populations (F_{ST} = 0.0023; Knutsen *et al.*2003), Atlantic herring (*Clupea harengus*) in
Swedish waters (F_{ST} = 0.002–0.003; Larsson *et al.* 2010), and Pacific herring (*C. pallasii*) in
British Columbia (F_{ST} = 0.003; Beacham *et al.* 2008). Results for Ayu population structure in
some regions, despite the high gene flow (F_{ST} = 0.0058), are comparable to these findings for
cod and herring. The isolation-by-distance relationship within amphidromous populations
(Figure 5) agrees with the maintenance of population structure in several regions.
Genetic differentiation was very low in the western Sea of Japan, northern and
central Pacific, and Seto Inland Sea, with many populations nested among clusters (Figures
1b, S4). The coefficient of variation (CV) of *He* values was surprisingly low at only 2%,
indicating the homogeneous distribution of *He* in Japan (Figure 2). The nested population
structure, very high gene flow, and homogeneous distribution of genetic diversity, suggest that
translocated landlocked juveniles and hatchery reared juveniles have reproduced with native
amphidromous fish in Japanese rivers. The isolation-by-distance relationship (Figure 5)
supports these results.
414
Hybridisation between landlocked and amphidromous form
The sample from Yodo River (#98) was included in the landlocked cluster, with the proportion
of hybrid and purebred landlocked individuals being very similar to that of Lake Biwa, in that
there were no purebred amphidromous individuals. Upstream migration of Ayu along Yodo
River, the sole outlet to flow from Lake Biwa, was interrupted by the construction of the
Amagase Dam in 1954. Well-developed larvae have been supplied from Lake Biwa into this
contact zone every year, and they might survive in coastal waters and return to migrate up
Yodo River (Takeshima *et al.* 2009). Results agree with Takeshima et al.'s speculation, in that

sample #98 collected from Yodo River was supplied from Lake Biwa. However, the unrooted
NJ tree based on different samples (Figure S5) located samples of upstream migrants between
Lake Biwa and amphidromous populations, suggesting hybridization occurred in the Yodo
River system. Contrarily, hybrid proportions (Figure 4) in Lake Biwa should reflect the
demographic history of the landlocked form, which might have been most recently, naturally
invaded by amphidromous individuals about 100 k years before present (Nishida 1985).
Tone River (#67), located a short distance from the amphidromous cluster in the NJ
tree (Figure 1b), had the highest hybrid proportion (60%) of all 118 amphidromous
populations. Each year throughout the 1970s more than 1×10^6 landlocked Ayu juveniles
were translocated from Lake Biwa into a tributary of Tone River (Takamura 2009). The
substantial catch in this area (Figure S1) suggests an appropriate estuarine environment for
juvenile survival, which might enhance hybrid fitness. Marine juveniles collected in coastal
waters in late March and early April (samples 37, 38) comprised 33% and 37% hybrids,
respectively (Table S1), indicating that juvenile hybrids do survive at sea. In Sho River (#30),
where the estimated proportion of landlocked form individuals was close to 0 (with 95%
confidence intervals including 0 (Iwata et al., 2007)), the hybrid proportion was the lowest
(15%) and the purebred amphidromous form was 85%. The single-site marker using the
mtDNA control region could detect a maternal but not paternal effect on reproduction of the
landlocked form (Iwata *et al.* 2007). mtDNA plays a key role in aerobic ATP production and
the ability for endurance exercise training (reviewed by Stefàno *et al.* 2019), and the mtDNA
control region functions in dNTP metabolism (Nicholls & Minczuk 2014) and oxygen
consumption (Kong *et al.* 2020). Non neutral mtDNA control region might affect their mixing
proportion estimates.
Pre-translocation population structure is unknown because no genetic data exist, and
translocation and hatchery releases have been performed throughout Japanese rivers.
Therefore, a BACI design (before-after-control impact, Smith, 2002) is not possible. However,
analyses consistently explain present-day population structure. Our results suggested
substantial backcrossing of landlocked and amphidromous forms throughout Japanese rivers,
while purebred landlocked fish were not survived (Figure 4). Crossing experiments of
hatchery and wild salmon demonstrated that most of the genes have an additive effect and
showed intermediate gene expression (Christie *et al.* 2016). The hybrid Ayu produced from
hatchery-reared and wild fish had intermediate susceptibility to bacterial cold-water disease
(Nagai & Sakamoto 2006). However, the genetic background of gene transcription looked to
be largely nonadditive (Gibson *et al.* 2004; Hedgecock *et al.* 2007; Roberge *et al.* 2008).
Hybrid individuals can have nonadditive inheritance of quantitative traits, such as outbreeding
depression (Gilk *et al.* 2004) and heterosis (Lippman & Zamir 2007). The effect of
hybridisation on genes controlling fitness traits of Ayu is unknown, but nation-wide
hybridisation between landlocked and amphidromous forms (Figure 4) should reduce fitness
of populations in Japanese rivers, because pure landlocked Ayu show very low survival in
seawater (Tabata & Azuma 1986; Iguchi & Yamaguchi 1994).
463
Implications for conservation and management
Translocation of landlocked Ayu may reduce fitness of amphidromous populations because of
their lower fitness. Therefore, an important action to ensure the sustainability of this
freshwater species would be to avoid the use of translocated landlocked Ayu juveniles to stock
rivers. In hatcheries, parent fish have been captive-reared for generations, mainly because of
difficulties experienced collecting sufficient mature wild fish for synchronous spawning each
year (Iguchi & Mogi 2007). Consequently, genetic diversity of hatchery-born juveniles has
decreased over generational (4–31) time (Iguchi *et al.* 1999; Ikeda *et al.* 2005; Tsuboi *et al.*
2019). Continued release of captive-reared juveniles could reduce the effective population

size of the entire population (hatchery + wild) and replace wild genes (Ryman & Laikre 1991).
Moreover, evidence exists for reduced reproductive success of hatchery-reared salmonids
when they breed in the wild (Fleming *et al.* 2000; McGinnity *et al.* 2003; Araki *et al.* 2007;
Christie *et al.* 2014). Field experiments (Tsuboi *et al.* 2019) found that hatchery-reared Ayu
born from captive-reared parents for more than seven generations had low survival,
maladapted behaviour of migration, and poor vulnerability to fishing when they were released
in the wild. Apparently, reducing hatchery releases is crucial to reduce genetic effects of
hatchery releases. As recommended by Takeshima and colleagues (2016a), translocation of
hatchery reared juveniles should also be limited within regions.
Because the migration range of marine juveniles is expected to be limited, individual
rivers might be inappropriate management units. Rivers or areas where there was no release of
landlocked or hatchery-reared Ayu should be established within each prefecture to limit gene
flow and dilute any genetic effects of translocations and hatchery releases through
backcrosses with wild individuals (Anderson 1949; McGinnity *et al.* 2003; Hindar *et al.* 2006;
Tymchuk *et al.* 2006; Roberge *et al.* 2008; Kitada *et al.* 2019). Healthy stocks of wild fish are
necessary for no-release rivers and/or areas to succeed as a management option. Fortunately,
wild fish genes remain dominant in Ayu (Figures 4).
Suitable spawning and nursery habitat is essential for reproduction of marine and
freshwater species. A rigorous monitoring of red sea bream (*Pagrus major*) demonstrated that
both increased seaweed communities in nursery habitat and reduced fishing effort were
primarily responsible for wild population recovery, while the effects of hatchery released
fishes decreased because of the exponentially reduced survival rates of juveniles produced
from captive-reared parents (Kitada *et al.* 2019). Releases of hatchery reared juveniles may be
useful for short-term management objectives, but management of spawning escapement, and
improving spawning and nursery habitat are the keys for the long-term sustainability of
fishery resources in Japan (Kitada 2020). Management began by the Shubutogawa Fishers
Cooperatives, Hokkaido, where Ayu juveniles had been translocated from Honshu since 1984;
this practice was stopped in 2013 to avoid genetic risks to the conservation of this species at
its northern margin. Efforts to create and/or restore spawning grounds in this river have
continued in collaboration with fishers and locals (https://www.shubutogawa.com/).
503

Acknowledgements
This research was made possible by access to microsatellite genotype data for Ayu
maintained by Hirohiko Takeshima and his colleagues. Gratitude is also expressed to the
researchers and staff responsible for the time-consuming sampling of Ayu throughout its
distribution in Japan that enabled this dataset to be generated. This study was supported by the
Japan Society for the Promotion of Science Grant-in-Aid for Scientific Research (KAKENHI)
No. 18K0578116.

511
512 **Conflict of interest**
513 The author has no conflict of interest to declare.
514
515 **Data availability statement**
516 All source data used are in the public sector, and links to their online sources are specified in
517 the text or in Supporting information.
518

References

Anderson E. Introgressive Hybridization. New York, USA: Wiley; 1949

Anderson EC, Thompson EA (2002) A model-based method for identifying species hybrids
using multilocus genetic data. *Genetics* **160**: 1217–1229.
https://doi.org/10.1093/genetics/160.3.1217

Araki H, Cooper B, Blouin MS (2007) Genetic effects of captive breeding cause a rapid,
cumulative fitness decline in the wild. *Science* **318**: 100–103.
https://doi.org/10.1126/science.1145621

Araki H, Schmid C (2010) Is hatchery stocking a help or harm?: Evidence, limitations and
future directions in ecological and genetic surveys. *Aquaculture* **308**: S2–S11.
https://doi.org/10.1016/j.aquaculture.2010.05.036

Barton NH (1979) The dynamics of hybrid zones. *Heredity* **43**: 341–359.
https://doi.org/10.1038/hdy.1979.87

Barton NH, Hewitt GM (1985) Analysis of hybrid zones. *Ann. Rev. Ecol. System.* **16**:113–148.
https://doi.org/10.1146/annurev.es.16.110185.000553

Beacham TD, Schweigert JF, MacConnachie C, Le KD, Flostrand L (2008) Use of
microsatellites to determine population structure and migration of Pacific herring in
British Columbia and adjacent regions. *Trans. Am. Fish. Soc.* **137**: 1795–1811.
https://doi.org/10.1577/T08-033.1

Beard ZS, Mann RD (2020) Retracted: Factors influencing angler satisfaction and catch rates
in Arizona stream put-and-take trout fisheries. *N. Am. J. Fish. Manag.* **40**: 521–521.
https://doi.org/10.1002/nafm.10322

Beaumont M, Barratt EM, Gottelli D, Kitchener AC, Daniels MJ, Pritchard JK *et al.* (2001)
Genetic diversity and introgression in the Scottish wildcat. *Molec. Ecol.* **10**: 319–336.
https://doi.org/10.1046/j.1365-294X.2001.01196.x

Berger-Tal O, Blumstein DT, Swaisgood RR (2020) Conservation translocations: a review of
common difficulties and promising directions. *Anim. Conserv.* **23**: 121–131.
https://doi.org/10.1111/acv.12534

Bucharova A (2017) Assisted migration within species range ignores biotic interactions and
lacks evidence. *Restor. Ecol.* **25**: 14–18. https://doi.org/10.1111/rec.12457

Christie MR, Ford MJ, Blouin MS (2014) On the reproductive success of early-generation
hatchery fish in the wild. *Evol. Appl.* **7**: 883–896. https://doi.org/10.1111/eva.12183

Christie MR, Marine ML, Fox SE, French RA, Blouin MS (2016) A single generation of
domestication heritably alters the expression of hundreds of genes. *Nature*
Communications **7**: 1–6. https://doi.org/10.1038/ncomms10676

Conant S (1988) Saving endangered species by translocation: are we tinkering with
evolution?. *BioSci.* **38**: 254–257. https://doi.org/10.2307/1310848

Devitt TJ, Baird SJ, Moritz C (2011) Asymmetric reproductive isolation between terminal
forms of the salamander ring species *Ensatina eschscholtzii* revealed by fine-scale
genetic analysis of a hybrid zone. *BMC Evol. Biol.* **11**: 1–15.
https://doi.org/10.1186/1471-2148-11-245

Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus
genotype data: linked loci and correlated allele frequencies. *Genetics* **164**: 1567–1587.
https://doi.org/10.1111/j.1471-8286.2007.01758.x

Fleming IA, Hindar K, MjÖlnerÖd IB, Jonsson B, Balstad T, Lamberg A (2000) Lifetime
success and interactions of farm salmon invading a native population. *Proc. Royal Soc.*
Lond. B **267**: 1517–1523. https://doi.org/10.1098/rspb.2000.1173

Fogelqvist J, Verkhozina AV, Katyshev AI, Pucholt P, Dixelius C, Rönnberg-Wästljung AC *et*
al. (2015) Genetic and morphological evidence for introgression between three species
of willows. *BMC Evol. Biol.* 15: 1–10. https://doi.org/10.1186/s12862-015-0461-7

George AL, Kuhajda BR, Williams JD, Cantrell MA, Rakes PL, Shute JR (2009) Guidelines for propagation and translocation for freshwater fish conservation. *Fisheries* **34**: 529–545. https://doi.org/10.1577/1548-8446-34.11.529

Geraldes A, Farzaneh N, Grassa CJ, McKown AD, Guy RD, Mansfield SD *et al.* (2014) Landscape genomics of Populus trichocarpa: the role of hybridization, limited gene flow, and natural selection in shaping patterns of population structure. *Evolution* **68**: 3260–3280. https://doi.org/10.1111/evo.12497

Gibson G, Riley-Berger R, Harshman L, Kopp A, Vacha S, Nuzhdin S *et al.* (2004) Extensive sex-specific nonadditivity of gene expression in Drosophila melanogaster. *Genetics*, 167, 1791–1799. https://doi.org/10.1534/genetics.104.026583

Gilk SE, Wang IA, Hoover CL, Smoker WW, Taylor SG, Gray AK. *et al.* (2004) Outbreeding depression in hybrids between spatially separated pink salmon, *Oncorhynchus gorbuscha*, populations: marine survival, homing ability, and variability in family size. *Environ. Biol. Fish.* **69**: 287–297. https://doi.org/10.1023/B:EBFI.0000022888.28218.c1

Glover KA, Solberg MF, McGinnity P, Hindar K, Verspoor E, Coulson MW. *et al.* (2017) Half a century of genetic interaction between farmed and wild Atlantic salmon: status of knowledge and unanswered questions. *Fish Fish.***18**: 890–927. https://doi.org/10.1111/faf.12214

Grabenstein KC, Taylor SA (2018) Breaking barriers: causes, consequences, and experimental utility of human-mediated hybridization. *Trends Ecol. Evol.* **33**: 198–212. https://doi.org/10.1016/j.tree.2017.12.008

Healy BD, Omana Smith EC, Schelly RC, Trammell MA, Nelson C B (2020) Establishment of a reproducing population of endangered Humpback Chub through translocations to a Colorado River tributary in Grand Canyon, Arizona. *N. Am. J. Fish. Manag.* **40**: 278–292. https://doi.org/10.1002/nafm.10408

Hedgecock D, Lin JZ, DeCola S, Haudenschild CD, Meyer E, Manahan DT *et al.* (2007) Transcriptomic analysis of growth heterosis in larval Pacific oysters (*Crassostrea gigas*). *Proc. Natl. Acad. Sci.* **104**: 2313–2318. https://doi.org/10.1073/pnas.0610880104

Hewitt GM (1988) Hybrid zones-natural laboratories for evolutionary studies. *Trends in Ecol. Evol.* **3**:158–167. https://doi.org/10.1016/0169-5347(88)90033-X

Hindar K, Fleming IA, McGinnity P, Diserud O (2006) Genetic and ecological effects of salmon farming on wild salmon: modelling from experimental results. *ICES J. Mar. Sci.* **63**: 1234–1247. https://doi.org/10.1016/j.icesjms.2006.04.025

Hirashiki C, Kareiva P, Marvier M (2021) Concern over hybridization risks should not preclude conservation interventions. *Conserv. Sci. Prac.* **3(4)**: e424. https://doi.org/10.1111/csp2.424 https://doi.org/10.1111/j.1467-2979.2008.00299.x

Iguchi K, Ito F (1994) Occurrence of cross-mating in Ayu: Amphidromous× land-locked forms, and diploid× triploid. *Fish. Sci.* **60**: 653–655. https://doi.org/10.2331/fishsci.60.653

Iguchi K, Yamaguchi M (1994) Adaptive significance of inter-and intrapopulational egg size variation in ayu *Plecoglossus altivelis* (Osmeridae). *Copeia*: 184–190. https://doi.org/10.2307/1446684

Iguchi K, Tanimura Y, Nishida M (1997) Sequence divergence in the mtDNA control region of amphidromous and landlocked forms of ayu. *Fish. Sci.* **63**: 901–905. https://doi.org/10.2331/fishsci.63.901

Iguchi K, Watanabe K, Nishida M (1999) Reduced mitochondrial DNA variation in hatchery populations of ayu (*Plecoglossus altivelis*) cultured for multiple generations. *Aquaculture* **178**: 235–243.https://doi.org/10.1016/S0044-8486(99)00133-7

Iguchi K, Nishida M (2000) Genetic biogeography among insular populations of the amphidromous fish *Plecoglossus altivelis* assessed from mitochondrial DNA analysis.

Conser. Genet. **1**: 147–156. https://doi.org/10.1023/A:1026582922248

Iguchi K, Ohkawa T, Nishida M (2002) Genetic structure of land-locked ayu within the Biwa Lake system. _Fish. Sci._ **68**: 138–143. https://doi.org/10.1046/j.1444-2906.2002.00399.x

Iguchi K, Konishi M, Takeshima H (2006) Early dispersal of ayu during marine stages as inferred from geographic variation in the number of vertebrae. _Fish. Sci._ **72**: 737–741. https://doi.org/10.1111/j.1444-2906.2006.01212.x

Iguchi K, Mogi M (2007) Effect of introducing wild paternity on stock performance of hatchery-reared ayu. _Fish. Sci._ **73**: 845–850. https://doi.org/10.1111/j.1444-2906.2007.01405.x

Ikeda M, Taniguchi N (2002) Genetic variation and divergence in populations of ayu _Plecoglossus altivelis_, including endangered subspecies, inferred from PCRRFLP analysis of the mitochondrial DNA D-loop region. _Fish. Sci._ **68**: 18–26. https://doi.org/10.1046/j.1444-2906.2002.00384.x

Ikeda M, Takagi S, Taniguchi N (2005) Relationships between genetic diversity and number of successive generations in hatchery populations of ayu _Plecoglossus altivelis_ assessed by microsatellite DNA polymorphism. _Nippon Suisan Gakkaishi_ **71**:768–774. (In Japanese with English abstract) https://doi.org/10.2331/suisan.71.768

Iwata H, Takeshima H, Tago Y, Watanabe K, Iguchi K, Nishida M (2007) Post-stocking dynamics of landlocked ayu _Plecoglossus altivelis_ traced by a mitochondrial SNP marker. _Nippon Suisan Gakkaishi_ **73**: 278–283. (In Japanese with English abstract) https://doi.org/10.2331/suisan.73.278

Izumi S, Wakabayashi H (1997) Use of PCR to detect _Cytophaga psychrophila_ from apparently healthy juvenile ayu and coho salmon eggs. _Fish Pathol._ **32**: 169-173. https://doi.org/10.3147/jsfp.32.169

Johannesson K, Le Moan A, Perini S, André C (2020) A Darwinian laboratory of multiple contact zones. _Trends Ecol. Evol._ **35**: 1021–1036. https://doi.org/10.1016/j.tree.2020.07.015

Kaewsangk K, Hayashizaki K, Asahida T, Ida H (2000) An evaluation of the contribution of stocks in the supplementation of ayu _Plecoglossus altivelis_ in the Tohoku area, using allozyme markers. _Fish. Sci._ **66**: 915–923. https://doi.org/10.1046/j.1444-2906.2000.00147.x

Karamanlidis AA, Skrbinšek T, Amato G, Dendrinos P, Gaughran S. et al. 2021 Genetic and demographic history define a conservation strategy for earth's most endangered pinniped, the Mediterranean monk seal Monachus monachus. _Sci. Rep._ **11**: 373. https://doi.org/10.1038/s41598-020-79712-1

Kawanabe H (1958) On the Significance of the Social Structure for the Mode of Density Effect in a Salmon-like Fish," Ayu", _Plecoglossus altivelis_ TEMMINCK et SCHLEGEL. _Mem. Coll. Sci. Univ. Kyoto Series B_ **25**: 171–180.

Kitada S (2018) Economic, ecological and genetic impacts of marine stock enhancement and sea ranching: a systematic review. _Fish Fish._ **19**: 511–532. https://doi.org/10.1111/faf.12271

Kitada S (2020) Lessons from Japan marine stock enhancement and sea ranching p programmes over 100 years. _Rev. Aquacult._ **12**: 1944–1969. https://doi.org/10.1111/raq.12418

Kitada S, Tezuka K (2002) Longitudinal logbook survey designs for estimating recreational fishery catch, with application to ayu (_Plecoglossus altivelis_). _Fish. Bull._ **100**: 228–243. http://fishbull.noaa.gov/1002/07kitada.pdf

Kitada S, Nakamichi R, Kishino H (2017) The empirical Bayes estimators of fine-scale population structure in high gene flow species. Molec. Ecol. Res. **17**: 1210–1222. https://doi.org/10.1111/1755-0998.12663

Kitada S, Nakajima K, Hamasaki K, Shishidou H, Waples RS, Kishino H (2019) Rigorous
monitoring of a large-scale marine stock enhancement program demonstrates the need
for comprehensive management of fisheries and nursery habitat. *Sci. Rep.* **9**: 1–10.
https://doi.org/10.1038/s41598-019-39050-3

Kitada S, Nakamichi R, Kishino H (2021) Understanding population structure in an
evolutionary context: population-specific F_{ST} and pairwise F_{ST}. *G3:*
Genes|*Genomes*|*Genetics* (accepted), bioRxiv. 927186v4.
https://doi.org/10.1101/2020.01.30.927186

Knutsen H, Jorde PE, André C, Stenseth NC (2003) Fine-scaled geographical population
structuring in a highly mobile marine species: the Atlantic cod. *Molec. Ecol.* **12**: 385–394.
https://doi.org/10.1046/j.1365-294X.2003.01750.x

Kong M, Xiang H, Wang J, Liu J, Zhang X, Zhao X (2020) Mitochondrial DNA Haplotypes
Influence Energy Metabolism across Chicken Transmitochondrial Cybrids. *Genes* **11**:
100. https://doi.org/10.3390/genes11010100

Kwan YS, Song HK, Lee HJ, Lee W O, Won YJ (2012) Population genetic structure and
evidence of demographic expansion of the Ayu (*Plecoglossus altivelis*) in East Asia.
Animal Syst. Evol. Divers. **28**: 279-290. https://doi.org/10.5635/ASED.2012.28.4.279

Laikre L, Schwartz MK, Waples RS, Ryman N (2010) Compromising genetic diversity in the
wild: unmonitored large-scale release of plants and animals. *Trends Ecol. Evol.* **25**:
520–529. https://doi.org/10.1016/j.tree.2010.06.013

Larison B, Lindsay AR, Bossu C, Sorenson MD, Kaplan JD, Evers DC *et al.* (2021)
Leveraging genomics to understand threats to migratory birds. *Evol. Appl.* 1**4**:
1646–1658. https://doi.org/10.1111/eva.13231

Larsson LC, Laikre L, André C, Dahlgren TG, Ryman N (2010) Temporally stable genetic
structure of heavily exploited Atlantic herring (*Clupea harengus*) in Swedish waters.
Heredity **104**: 40–51. https://doi.org/10.1038/hdy.2009.98

Lippman ZB, Zamir D (2007) Heterosis: revisiting the magic. *Trends Genet.* **23**: 60–66.
https://doi.org/10.1016/j.tig.2006.12.006

Lischer HEL, Excoffier L (2012) PGDSpider: An automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics* **28**: 298–299.
https://doi.org/10.1093/bioinformatics/btr642

Lorenzen K, Beveridge MC, Mangel M (2012) Cultured fish: integrative biology and
management of domestication and interactions with wild fish. *Biol. Rev.* 87: 639–660.
https://doi.org/10.1111/j.1469-185X.2011.00215.x

Manel S, Gaggiotti OE, Waples RS (2005) Assignment methods: matching biological
questions with appropriate techniques. *Trends Ecol. Evol.* **20**: 136–142.
https://doi.org/10.1016/j.tree.2004.12.004

Martinsen GD, Whitham TG, Turek RJ, Keim P (2001) Hybrid populations selectively filter
gene introgression between species. *Evolution* **55**: 1325–1335.
https://doi.org/10.1111/j.0014-3820.2001.tb00655.x

McGinnity P, Prodöhl P, Ferguson A, Hynes R, ó Maoiléidigh N, Baker N *et al.* (2003) Fitness
reduction and potential extinction of wild populations of Atlantic salmon, *Salmo salar*, as
a result of interactions with escaped farm salmon. *Proc. Royal Soc. Lond. B* **270**:
2443–2450. https://doi.org/10.1098/rspb.2003.2520

Minckley WL (1995) Translocation as a tool for conserving imperiled fishes: experiences in
western United States. *Biol. Conserv.* **72**: 297–309.
https://doi.org/10.1016/0006-3207(94)00091-4

Moritz C (1995) Uses of molecular phylogenies for conservation. *Philos. Trans. R. Soc. B*
Biol. Sci. **349**: 113–118. https://doi.org/10.1098/rstb.1995.0097

Nagai T, Sakamoto T (2006) Susceptibility and immune response to Flavobacterium psychrophilum between different stocks of ayu *Plecoglossus altivelis*. *Fish Pathol.* **41**: 99–104. https://doi.org/10.3147/jsfp.41.99

Nei M, Chesser RK (1983) Estimation of fixation indices and gene diversities. *Ann. Hum. Genet.* **47**: 253–259. https://doi.org/10.1111/j.1469-1809.1983.tb00993.x

Nicholls TJ, Minczuk M (2014) In D-loop: 40 years of mitochondrial 7S DNA. *Exp. Gerontol.* **56**: 175–181. https://doi.org/10.1016/j.exger.2014.03.027

Nishida M (1985) Substantial genetic differentiation in Ayu *Plecoglossus altivelis* of the Japan and Ryukyu Islands. *Bull. Jap. Soc. Sci. Fish.* **51**: 1269–1274. https://doi.org/10.2331/suisan.51.1269

Nishida M (1986) Geographic variation in the molecular, morphological and reproductive characters of the ayu *Plecoglossus altivelis* (Plecoglossidae) in the Japan-Ryukyu Archipelago. *Jap. J. Ichthy.* **33**: 232–248. https://doi.org/10.11369/jji1950.33.232

Otake T, Yamada C, Uchida K (2002) Contribution of stocked ayu (*Plecoglossus altivelis altivelis*) to reproduction in the Nagara River, Japan. *Fish. Sci.* **68**: 948–950. https://doi.org/10.1046/j.1444-2906.2002.00516.x

Pastene LA, Numachi K, Tsukamoto K (1991) Examination of reproductive success of transplanted stocks in an amphidromous fish, *Plecoglossus altivelis* (Temmink et Schlegel) using mitochondrial DNA and isozyme markers. *J. Fish Biol.* **39**: 93–100. https://doi.org/10.1111/j.1095-8649.1991.tb05072.x

Pawson MG (1982) Recapture rates of trout in a put-and-take fishery: analysis and management implications. *Aquacult. Res.* **13**: 19–31. https://doi.org/10.1111/j.1365-2109.1982.tb00028.x

Pierpaoli M, Biro ZS, Herrmann M, Hupe K, Fernandes M, Ragni B *et al.* (2003). Genetic distinction of wildcat (*Felis silvestris*) populations in Europe, and hybridization with domestic cats in Hungary. *Molec. Ecol.* **12**: 2585–2598. https://doi.org/10.1046/j.1365-294X.2003.01939.x

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959. https://doi.org/10.1093/genetics/155.2.945

Ramachandran S, Deshpande O, Roseman C C, Rosenberg NA, Feldman MW, Cavalli-Sforza LL (2005) Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. *Proc. Natl. Acad. Sci.* **102**: 15942–15947. https://doi.org/10.1073/pnas.0507611102

Ricciardi A, Simberlof, D (2009) Assisted colonization is not a viable conservation strategy. *Trends Ecol. Evol.* **24**: 248–253. https://doi.org/10.1016/j.tree.2008.12.006

Roberge C, Normandeau E, Einum S, Guderley H, Bernatchez L (2008) Genetic consequences of interbreeding between farmed and wild Atlantic salmon: insights from the transcriptome. *Molec. Ecol.* **17**: 314–324. https://doi.org/10.1111/j.1365-294X.2007.03438.x

Ross MR, Loomis DK (2001) Put-and-take fisheries: investigating catch and retention assumptions. *Fisheries* **26**: 13-18. https://doi.org/10.1577/1548-8446(2001)026<0013:PFICAR>2.0.CO;2

Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molec. Ecol. Resour.* **8**: 103–106. https://doi.org/10.1111/j.1471-8286.2007.01931.x

Ryman N, Laikre L (1991) Effects of supportive breeding on the genetically effective population size. *Conserv. Biol.* **5**: 325–329. https://doi.org/10.1111/j.1523-1739.1991.tb00144.x

Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Molec. Biol. Evol.* **4**: 406–425.
https://doi.org/10.1093/oxfordjournals.molbev.a040454
Seddon PJ, Griffiths CJ, Soorae PS, Armstrong DP (2014) Reversing defaunation: restoring
species in a changing world. *Science* **345**: 406–412.
https://doi.org/10.1126/science.1251818
Seki S, Taniguchi N, Jeon SR (1988) Genetic divergence among natural populations of Ayu
from Japan and Korea. *Nippon Suisan Gakkaishi* **54**: 559–568. (In Japanese with English
abstract) https://doi.org/10.2331/suisan.54.559
Seki S, Taniguchi N, Murakami N, Takamichi A, Takahashi I (1994) Seasonal changes in the
mixing rate of restocked ayu-juveniles and assessment of native stock using an allozyme
marker. *Fish. Sci.* **60**: 31–35. https://doi.org/10.2331/fishsci.60.31
Seki S, Taniguchi N (1998) Traking the released Ayu of landlocked form by isozyme markers.
Nippon Suisan Gakkaishi **54**: 745–749. (In Japanese with English abstract)
https://doi.org/10.2331/suisan.54.745
Slatkin M (1987) Gene flow and the geographic structure of natural populations. *Science* **236**:
787–792. https://doi.org/10.1126/science.3576198
Smith EP. BACI design. In: El-Shaarawi AH, Piegorsc WW, eds. *Encyclopedia of*
Environmetrics, 1, Wiley; 2002: 141–148.
Stefàno E, Marsigliante S, Vetrugno C, Muscella A (2019) Is mitochondrial DNA profiling
predictive for athletic performance?. *Mitochondrion* **47**: 125–138.
https://doi.org/10.1016/j.mito.2019.06.004
Steyer K, Tiesmeyer A, Muñoz-Fuentes V, Nowak C (2018) Low rates of hybridization
between European wildcats and domestic cats in a human-dominated landscape. *Ecol.*
Evol. **8**: 2290–2304. https://doi.org/10.1002/ece3.3650
Tabata K, Azuma M (1986) Survival of three types of ayu fish *Plecoglossus altivelis* fry in sea
water rearing. *Bull. Hyogo Pref. Fish. Exp. Stn*. **24**: 29–34. (In Japanese with English
abstract)
Tago Y (2002) Larval distribution of ayu *Plecoglossus altivelis* in the surface layer of estuary
regions in Toyama Bay. *Nippon Suisan Gakkaishi* **68**: 61–71. (In Japanese with English
abstract) https://doi.org/10.2331/suisan.68.61
Tago Y (2004) Relationship between body size of Ayu migrating up rivers flowing into
Toyama Bay and water temperature. *Suisanzoshoku* **52**: 315–323. (In Japanese with
English abstract) https://doi.org/10.11233/aquaculturesci1953.52.315
Takagi M, Shoji E, Taniguchi N (1999) Microsatellite DNA polymorphism to reveal genetic
divergence in ayu, *Plecoglossus altivelis*. *Fish. Sci.* **65**: 507–512.
https://doi.org/10.2331/fishsci.65.507
Takahashi I, Azuma K, Fujita S, Kinoshita I (1998) Spatial distribution of larval ayu
Plecoglossus altivelis in the Shimanto estuary, Japan. *Fish. Sci.* **64**: 522–525.
https://doi.org/10.2331/fishsci.64.522
Takamura K (2009) Invasive species from the highly endemic fish fauna of Lake Biwa
threatening freshwater fish in rivers of the Kanto region. Jap. J. Limnol. **70**: 249–253. (In
Japanese with English abstract) https://doi.org/10.3739/rikusui.70.249
Takeshima H, Iguchi K, Nishida M (2005) Unexpected ceiling of genetic differentiation in the
control region of the mitochondrial DNA between different subspecies of the ayu
Plecoglossus altivelis. *Zool. Sci.* **22**: 401–410. https://doi.org/10.2108/zsj.22.401
Takeshima H, Iguchi K, Nishida M (2009) Ayu (*Plecoglossus altivelis*) in a contact zone
between amphidromous and landlocked forms: genetic analyses of populations in the
Yodo River system. *Zool. Sci.* **26**: 536–542. https://doi.org/10.2108/zsj.26.536
Takeshima H, Iguchi K, Hashiguchi Y, Nishida M (2016a) Using dense locality sampling

resolves the subtle genetic population structure of the dispersive fish species
Plecoglossus altivelis. *Molec. Ecol.* **25**: 3048–3064. https://doi.org/10.1111/mec.13650
Takeshima H, Iguchi K, Hashiguchi Y, Nishida M (2016b) Data from: Using dense locality
sampling resolves the subtle genetic population structure of the dispersive fish species
Plecoglossus altivelis, Dryad, Dataset, https://doi.org/10.5061/dryad.s55f5
Taniguchi N, Seki S, Inada Y (1983) Genetic variability and differentiation of amphidromous,
landlocked, and hatchery populations of ayu *Plecoglossus altivelis*. *Bull. Jap. Soc. Sci.*
Fish. **49**: 1655–1663. (In Japanese with English abstract)
https://doi.org/10.2331/suisan.49.1655
Taniguchi N, Dong S, Kondo K, Imai S (2002) Mixed ratio of the released landlocked form
and distribution of the native amphidromous form of Ayu, *Plecoglossus altivelis* in
Yoshino River estimated by genetic markers. *Suisanzosyoku* **50**: 17–24. (In Japanese
with English abstract) https://doi.org/10.11233/aquaculturesci1953.50.17
Taranger GL, Karlsen Ø, Bannister RJ, Glover KA, Husa V, Karlsbakk E *et al.* (2015) Risk
assessment of the environmental impact of Norwegian Atlantic salmon farming. *ICES J.*
Mar. Sci. **72**: 997–1021. https://doi.org/10.1093/icesjms/fsu132
Taylor LU, Benavides E, Simmons JW, Near TJ (2021) Genomic and phenotypic divergence
informs translocation strategies for an endangered freshwater fish. *Molec. Ecol.*
https://doi.org/10.1111/mec.15947
Tran HD, Iida M, Maeda K (2017) Downstream migration of newly-hatched ayu
(*Plecoglossus altivelis*) in the Tien Yen River of northern Vietnam. *Environ. Biol. Fish.*
100:1329–1341. https://doi.org/10.1007/s10641-017-0646-6
Tsuboi J, Kaji K, Baba S, Arlinghaus R (2019) Trade-offs in the adaptation towards hatchery
and natural conditions drive survival, migration, and angling vulnerability in a territorial
fish in the wild. *Can. J. Fish. Aquat. Sci.* **76**: 1757–1767.
https://doi.org/10.1139/cjfas-2018-0256
Tsukamoto K, Kajihara T (1987) Age determination of ayu with otolith. *Nippon Suisan*
Gakkaishi **53**: 1985–1997. https://doi.org/10.2331/suisan.53.1985
Tymchuk WE, Biagi C, Withler R, Devlin RH (2006) Growth and behavioral consequences of
introgression of a domesticated aquaculture genotype into a native strain of coho salmon.
Trans. Am. Fish. Soc. **135**: 442–455. https://doi.org/10.1577/T05-181.1
Vähä JP, Primmer CR (2006) Efficiency of model‐based Bayesian methods for detecting
hybrid individuals under different hybridization scenarios and with different numbers of
loci. *Molec. Ecol.* **15**: 63–72. https://doi.org/10.1111/j.1365-294X.2005.02773.x
Waples RS, Drake J. In: Leber K, Kitada S, Blankenship HL, Svåsand T, eds. *Stock*
enhancement and sea ranching, developments pitfalls and opportunities. Blackwell;
2004: 260–306.
Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some
genetic methods for identifying the number of gene pools and their degree of
connectivity. *Molec. Ecol.* **15**: 1419–1439.
https://doi.org/10.1111/j.1365-294X.2006.02890.x
Weir BS, Cockerham CC (1984) Estimating *F*-statistics for the analysis of population
structure. Evolution **38**: 1358–1370. https://doi.org/10.2307/2408641
Whitlock MC, Mccauley DE (1999) Indirect measures of gene flow and migration:
$F_{ST} \neq 1(4Nm + 1)$. *Heredity* **82**: 117–125.
https://doi.org/10.1046/j.1365-2540.1999.00496.x
Yamaguchi K, Otomo Y, Kaniwa J, Umezawa K, Dairiki K, Nakajima M (2016) Genetic
relationships and variability of Ayu (*Plecoglossus altivelis altivelis*) populations
migrating upstream in the Rivers Edo, Arakawa, and Tamagawa, which flow into Tokyo
Bay. *Fish Genet. Breed. Sci.* **46**: 41–48.

Yamamoto S, Uchida K, Sato T, Katsura K, Takasawa T (2007) Population genetic structure
and effective population size of ayu (*Plecoglossus altivelis*), an amphidromous fish. *J. of*
Fish Biol. **70**: 191–201. https://doi.org/10.1111/j.1095-8649.2007.01396.x

Supporting Information
Additional Supporting Information may be found in the Supporting Information section at the
end of this article.

Table S1 Ayu sampling locations and dates, and expected heterozygosities and hybrid
proportions
Figure S1 Ayu catch and release statistics in Japan
Figure S2 Juvenile release and production statistics
Figure S3 Ayu sampling locations throughout Japan
Figure S4 Unrooted neighbour-joining (NJ) trees for Ayu in Japan
Figure S5 Unrooted neighbour-joining (NJ) trees for Ayu in the Yodo River system
Figure S6 Results from STRUCTURE analysis for optimal K
Figure S7 Distribution of individual q-values of landlocked ancestry in 118 populations