RNA-Seq is not required to determine stable reference genes for qPCR normalization

Nirmal Kumar Sampathkumar*, ‡, a, b, f, Venkat Krishnan Sundaram*, ‡, c, Prakroothi S Danthi e, Rasha Barakat d, Shiden Solomon a, b, Mrityunjoy Mondal a, b, Ivo Carre a, b, Tatiana El Jalkh c, Aïda Padilla-Ferrer c, Julien Grenier c, Charbel Massaad c, Jacqueline C Mitchell b.

Affiliations

a UK-Dementia Research Institute, King's College London, London, UK.

b Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, Basic and Clinical Neuroscience Institute, King's College London, London UK.

c INSERM UMRS 1124, Université de Paris, Faculty of Basic and Biomedical Sciences, Paris, France.

d INSERM U1016, Institut Cochin, Paris, France.

e McGill Group for Suicide Studies, Douglas Mental Health University Institute, McGill University, Montréal, Canada.

f Present address: Alzheimer's Research UK Oxford Drug Discovery Institute, Centre for Medicines Discovery, University of Oxford, Oxford, UK

‡ Correspondence to nirmal.sampathkumar@cmd.ox.ac.uk and venkatkrishnan.sundaram@u-paris.fr

*These authors contributed equally.

Author contributions:

Conceptualization: NKS, VKS.
Methodology: NKS, VKS
Experimentation: VKS, VKS, RB, SMS, IC, MM, TEJ, APF.
Data analysis: VKS, NKS, PSD.
Validation: JM, CM.
Visualization: JG, MM.
Writing – original draft: NKS, VKS.
Writing – review & editing: All Authors.

Acknowledgements:

The authors thank the Animal House Core Facility and the Cyto2BM Molecular Biology Platform of BioMedTech Facilities (INSERM US36/CNRS UMS2009) for the animals and research services pertaining to the generation of qPCR data. We also thank Benjamin SAINTPIERRE and Dr. Franck LETOURNEUR from the Genomic Platform at Institut Cochin, Paris for the quality control of RNA samples and for their assistance in the analysis of RNAseq data. NKS would like to thank Prof. Christopher E SHAW and Tanisha LEWIS from the UK Dementia Research Institute, King's College London, for their support towards this work.

Data availability:

The original contributions presented in the study are included in the article & Supplementary Data. The metadata for the RNA-Seq and qPCR experiments can be accessed at https://doi.org/10.6084/m9.figshare.15169104.v1. The datasets available include the normalized count matrices for the RNA-Seq data, qPCR Cq values and differential expression analysis, primer sequences, Coefficient of Variation and NormFinder analysis. Further inquiries can be directly addressed to the corresponding authors.

Funding:

No specific funding for the study. VKS and APF were funded by a PhD fellowship from the French Ministry of Research and Innovation and Université de Paris. RB was funded by a doctoral fellowship from INSERM U1016, Institut Cochin, Paris. TEJ was funded by a doctoral fellowship from Université de Paris. SS and IC were funded by a PhD fellowship from the UK Dementia Research Institute. MM and NKS were funded by a postdoctoral fellowship from the UK Dementia Research Institute.

Abstract

Assessment of differential gene expression by qPCR is heavily influenced by the choice of reference genes. Although numerous statistical approaches have been proposed to determine the best reference genes, they can give rise to conflicting results depending on experimental conditions. Hence, recent studies propose the use of RNA-Seq to identify stable genes followed by the application of different statistical approaches to determine the best set of reference genes for qPCR data normalization. In this study, we demonstrate that the statistical approach to determine the best reference genes from randomly selected candidates is more important than the preselection of 'stable' candidates from RNA-Seq data. Using a qPCR data normalization workflow that we have previously established; we show that qPCR data normalization using randomly chosen conventional reference genes renders the same results as stable reference genes selected from RNA-Seq data. We validated these observations in two distinct cross-sectional experimental conditions involving human iPSC derived microglial cells and mouse sciatic nerves. These results taken together show that given a robust statistical approach for reference gene selection, stable genes selected from RNA-Seq data do not offer any significant advantage over commonly used reference genes for normalizing qPCR assays.

Keywords: qPCR, Reference Genes, RNA-Seq, Normalization Factor, Statistical Methods

Introduction

Normalization of relative gene expression using qPCR assays relies crucially on the use of reference genes that exhibit minimal variation across experimental conditions (Bustin *et al*, 2009; Thellin *et al*, 1999; Eisenberg & Levanon, 2013; Jacob *et al*, 2013; Chapman & Waldenström, 2015). Thus, the choice of reference genes has a significant bearing on the normalized profiles of target genes. In the last two decades, many statistical approaches have been proposed to help researchers identify stable reference genes from a given set of candidates (Vandesompele *et al*, 2002; Pfaffl *et al*, 2004; Andersen *et al*, 2004; De Spiegelaere *et al*, 2015; Silver *et al*, 2006; Boda *et al*, 2009). However, the calculations and assumptions employed in these approaches fundamentally differ and could give rise to conflicting results (Sundaram *et al*, 2019). To address these concerns, we recently reviewed these approaches and following a comparison of existing methods including Coefficient of Variation (CV) analysis, GeNorm, NormFinder, pairwise ΔCT method and Best Keeper, we proposed an effective qPCR analysis workflow (Sundaram *et al*, 2019). Our approach combining visual representation and statistical testing of intrinsic variation, CV analysis for identifying overall reference gene variation and the NormFinder algorithm, proved to be effective in determining stable reference genes from a given set of candidates.

However, more recently, a growing number of studies have proposed the use of bulk RNA-Seq to screen for stable reference genes followed by the application of existing statistical methods to obtain the "best" reference genes to normalize qPCR assays (Carmona *et al*, 2017; Zhan *et al*, 2014; Hoang *et al*, 2017; Yang *et al*, 2014; Chim *et al*, 2017; Dai *et al*, 2021; Pombo *et al*, 2017; Zhou *et al*; Hernandez-Segura *et al*, 2019). In these studies, the intrinsic validity of screening for reference genes directly from the bulk RNA-Seq data is clear, as they have used RNA-Seq to identify stable reference genes in an effort to standardize the use of certain genes for their experimental setting. Although bulk RNA-Seq data analysis pipelines quantify differential expression of target genes, performing a qPCR confirmation is often prudent as genes that exhibit shorter transcript lengths and lower expression levels exhibit discordant results between RNA-Seq and qPCR (Coenye, 2021; Everaert *et al*, 2017). This is largely because frequently used RNA-Seq normalization strategies are prone to overall and sample specific transcript-length bias wherein longer transcripts are attributed with more counts regardless of expression levels (Oshlack & Wakefield, 2009; Mandelboum *et al*, 2019). Moreover, in a standard RNA-Seq experiment employing 3 – 4 biological replicates per condition, a vast majority of the reads arise from a small set of highly expressed genes and thus there is an inherent discrimination towards genes that are less expressed in the system (Berge *et al*, 2019; Liu *et al*, 2014). These factors are mostly likely at play when discordant differential expression results are obtained between qPCR and RNA-Seq (Everaert *et al*, 2017). Thus, the selection of qPCR reference genes from RNA-Seq data is not the best strategy as it can potentially lead to sub-optimal candidates. Moreover, the cost-benefit trade off in performing RNA-Seq to identify stable reference genes for qPCR assays is negligible. This approach is also not always feasible, especially in cases of sample scarcity or when dealing with poor yields of RNA. It is particularly relevant when we are interested in only a small number of target

genes whose expression can be assessed solely by qPCR. However, although qPCR is not prone to the same biases as RNA-Seq, the strongest impediment to performing reliable qPCR data analysis is poor reference gene selection for a given experimental setting. Thus, the approach used to validate reference genes is of utmost importance and it should ideally not be dependent on a specific source of "stable" candidates. It is also advantageous if suitable reference genes can be identified by the same technology rather than RNA-Seq.

In this study, we address these concerns by demonstrating that the statistical approach employed to validate reference genes is more important than the preselection of stable genes from bulk RNA-Seq data. Specifically, with the right statistical approach, reference genes filtered from any random set of candidates generate the same differential expression results when compared to stable genes preselected from bulk RNA-Seq data. Furthermore, the results obtained are also comparable with the fold changes observed in RNA-Seq for the genes that we tested in qPCR. These results taken together demonstrate that the preselection of candidate reference genes is not of any significance for data normalization and hence, RNA-Seq data is not an essential requisite to obtain robust reference genes for qPCR data normalization.

Materials and Methods:

Sample Procurement:

iPSC Microglia:
The CRISPR/Cas gene edited TREM2 knock-out iPSC line BIONi010-C-17 (TREM2 KO) and its isogenic control BIONi010-C (TREM2 WT) were purchased from EBISC and maintained at 37°C 5% CO2 with E8-Flex medium. Microglia were differentiated from these lines as previously described (Haenseler *et al*, 2017). In short, using AggreWell 800 (34850, StemCell Technologies), embryonic bodies (EBs) were prepared from iPSCs in E8-Flex medium supplemented with 50 ng/ml VEGF (PHC9394, ThermoFisher), 50 ng/ml BMP4 (120-05ET, Peprotech), and 20 ng/ml SCF (300-07, PeproTech) for 3 days with 75% medium change each day. EBs were transferred to a T75 flask in X-VIVO15 (BE02-060F, Lonza) medium supplemented with 25 ng/ml IL-3 (PHC0031, ThermoFisher), 2 mM Glutamax (35050061, ThermoFisher), 100 ng/ml M-CSF (300-25, PeproTech), and 0.055 mM β-mercaptoethanol (31350-010, ThermoFisher). Precursor cells started to emerge from the EBs approximately 4 weeks later, and were collected by gently tapping the flask. These precursor cells are differentiated to microglia over 7-days in DMEM F:12, Neuronal basal medium plus (1:1) supplemented with 100 ng/ml IL-34, 10 ng/ml GM-CSF, and 100 ng/ml M-CSF.

Sciatic Nerves:
To assess the stability of commonly used reference genes during myelination of the sciatic nerves, P3 and P21 C57BL6/J mice (3 Males and 4 females) were dissected and the sciatic nerves were harvested for RNA extraction. The RNA-seq data for these time points was mined from a publicly available dataset as explained in the *"Bulk RNA-seq and data analysis"* section below. All aspects of animal care and animal experimentation were performed in accordance with the relevant guidelines and regulations of INSERM, Université de Paris, and approved by the French National Committee of Animal experimentation and ethics.

Total RNA extraction :

iPSC Microglia:
Approximately $2x10^5$ iPSC derived microglial cells were harvested using TRIzol reagent (Ambion Life Technologies 15596018) for total RNA isolation. Total RNA was isolated using Direct-Zol™ RNA microprep columns (Zymo research R2062) according to the manufacturer's protocol.

Sciatic Nerves:
Total RNA was extracted from sciatic nerves using 1 mL of TRIzol reagent (Ambion Life Technologies 15596018) on ice using the manufacturer's instructions with slight modifications. Briefly, 100% ethanol was substituted for isopropanol to reduce the precipitation of salts. In addition, RNA precipitation was carried out overnight at -20°C in the presence of glycogen (0.02mg/mL final concentration). The following day, precipitated RNA

188   was pelleted by centrifugation and washed at least 3 times with 70% Ethanol to eliminate any
189   residual contamination. Tubes were then spin-dried in a vacuum centrifuge for 5 minutes and
190   RNA was resuspended in 20 µL of RNase resuspension buffer containing 0.1mM EDTA, pH
191   8. RNA was then stored at -80°C till RTqPCR.
192
193
194   **RNA Quality, Integrity, and Assay:**
195
iPSC Microglia:
RNA quantity was assayed using UV spectrophotometry on Nanodrop One (Thermo
Scientific). Optical density absorption ratios A260/A280 & A260/A230 of the iPSC microglia
samples were 2.0 (±0.1 SD) and 2.1 (±0.1 SD) respectively. RNA integrity was verified using
Agilent bioanalyzer. All the samples exhibited a RIN score ≥ 9 and were subsequently used
for downstream analysis including paired-end bulk RNA-seq and qPCR.
202
Sciatic Nerves:
RNA quantity was assayed using UV spectrophotometry on Nanodrop One (Thermo
Scientific). Optical density absorption ratios A260/A280 & A260/A230 of sciatic nerve
samples were 1.84 (±0.04 SD) and 1.54 (±0.53 SD) respectively. RNA integrity was verified
using Agilent bioanalyzer at the Genomics Platform at Institut Cochin, Paris. All samples
exhibited a RIN score ≥ 8.8 and were used for qPCR analysis.
209
210   **Bulk RNA-seq and data analysis:**
211
For the iPSC samples, total RNA was extracted and sent to Genewiz, Germany for paired-end
bulk RNA-seq. The raw data, final processed data and the metadata are available in the GEO
database (GSE178924). For the sciatic nerves, bulk RNA-Seq data was obtained from a
previously published study (Ackerman *et al*, 2018). The raw data can be accessed at the Zenodo
repository (https:// zenodo.org/record/1154250). Only the WT datasets at P3 and P21 were
mined and re-analyzed. Fastq files were aligned using the STAR algorithm (version 2.7.6a).
Reads were then counted using RSEM (v1.3.1) and the statistical analyses on the read counts
were performed with the DESeq2 package (DESeq2_1.28.1) to determine the proportion of
differentially expressed genes between the two experimental conditions (Love *et al*, 2014). The
standard DESeq2 normalization method (DESeq2's median of ratios with the DESeq function)
was used, with a pre-filter of reads and genes (reads uniquely mapped on the genome, or up to
10 different loci with a count adjustment, and genes with at least 10 reads in at least 3 different
samples). The biomaRt package (v2.44.4) was used to substitute gene names for the Ensembl
IDs in the count matrix (Durinck *et al*, 2009). The DESeq2 pipeline was used to fit a
generalized linear model (GLM) for the expression of each gene relative to the experimental
groups. To determine the variation in gene expression, the dispersion value computed by the
GLM was tabulated into the results data frame. Square root of dispersion was calculated using
sqrt function from base functions in R (v4.0.2). The coefficient of dispersion (CVfromDisp)
was calculated by multiplying square root of dispersion by 100.
231

Reference gene selection from RNA-Seq:

The data frame was subjected to 4 filtration methods to select stable genes in both the iPSC and Sciatic nerve datasets. First filtration was done by retaining all the genes with padj value above 0.05. The resulting data frame was ordered based on the log2FoldChange (log2FC) column and the second filtration was performed by retaining all the genes with log2FC between -0.1 to 0.1 implying negligible intergroup variation. Third filtration was done by retaining all the genes that lie in the CVfromDisp range between 10 & 20. The fourth and last filtration was performed by retaining all the genes that exhibited a Basemean value greater than or equal to 500 implying sufficient expression for qPCR detection.

Once these criteria were applied, 683 genes were obtained (basemean values ranging from 500 to 123000) in the iPSC dataset and 42 genes were obtained (basemean values ranging from 500 to 48000) in the sciatic nerve dataset. To obtain stable reference genes with both low and high expression, the final list of candidates was partitioned into quartiles based on basemean expression. 3 reference genes from each of the first three quartiles and 1 reference gene from the last quartile were chosen; thereby generating 10 candidates. This selection strategy resulted in candidate reference genes that exhibit Cq values between 18 cycles (high expression) and 26 cycles (relatively lower expression) in qPCR.

Conventional reference gene selection:

All conventional reference genes were randomly chosen. They do not have any proven stability for the experimental setting in question. For the iPSC dataset, the conventional human reference genes chosen were *ACTB, GAPDH, GUSB, HPRT, PGK1, PPIA, RPL13A, TBP,* and *UBC*. For the sciatic nerve dataset, the conventional mouse reference genes chosen were *Actb, Gapdh, Tbp, Sdha, Pgk1, Ppia, Rpl13a, Hsp60, Mrpl10, Rps26*. These mouse reference genes have been previously used to establish the qPCR data analysis workflow (Sundaram *et al*, 2019)

Target gene selection:

Differentially expressed genes that exhibited Padj<0.05 were first retained from the results dataframe. Next, genes that exhibited log2FC values above +0.6 and below -0.6 were retained and partitioned into 2 separate lists. 3 target genes were randomly chosen from each of these lists for differential expression analysis in the iPSC dataset. In the Sciatic nerve dataset, 3 upregulated genes and 3 downregulated genes were chosen based on the recently published Sciatic Nerve Atlas https://www.snat.ethz.ch (Gerber *et al*, 2021).

Primer Design:

All primers used in the study were designed using the Primer 3 plus software (https://primer3plus.com/cgi-bin/dev/primer3plus.cgi). Splice variants and the protein-coding sequence of the genes were identified using the Ensembl database (www.ensembl.org).

276 Constitutively expressed exons among all splice variants were then identified using the
277 ExonMine database (Mollet *et al*, 2010). Primers that spanned two subsequent constitutively
278 expressed exons were then designed using the Primer 3 plus software. The amplicon size of all
279 primers was between 88bp – 200bp. For detailed information on Primer sequences refer to
280 Supplementary data.
281
282 **Amplification Efficiencies:**
283
The amplification efficiencies of primers were calculated using serial dilution of cDNA
molecules. Briefly, cDNA from both the experimental groups of the sciatic nerve (for mouse
primers) and iPSC (for human primers) were serially diluted four times by a factor of 10 (1,
1:10, 1:100 & 1:1000). qPCR was then performed using these dilutions and the results were
plotted as a standard curve against the respective concentration of cDNA. Amplification
efficiency (E) was calculated by linear regression of standard curves using the following
equation: $E = 10^{(-1/\text{Slope of the standard curve})}$. Primer pairs that exhibited an Amplification Efficiency
(E) of 1.9 to 2.1 (95% - 105%) and an R^2 value (Determination Coefficient) of 0.99 or above
were chosen for this study. The Cq values of all genes across different groups were well within
the range of the standard dilution curve.
294
295
296 **RT-qPCR:**
297
iPSC Microglia:
1000 ng of total RNA was first subjected to DNase digestion (Thermo scientific EN0525) at
37°C for 30 minutes to eliminate contaminating genomic DNA. Next, DNase activity was
stopped using EDTA (Thermo scientific) and cDNA synthesis was done using iSCRIPT^{TM}
cDNA synthesis kit from BioRad (1708891) to a total volume of 20 µl. cDNA was diluted 1:10
with nuclease free water for qPCR analysis. qPCR was performed using Takyon ROX SYBR
2X MasterMix (Eurogentec UF-RSMT-B0701) as a fluorescent detection dye. All reactions
were carried out in a final volume of 8 µl in 384 well plates with 300 nM gene-specific primers,
around 5 ng of cDNA (at 100% RT efficiency) and 1X SYBR Master Mix in each well. Each
reaction was performed in triplicates. All qPCR experiments were performed using
ThermoFisher Scientific QuantStudio 7 with a No-Template-Control (NTC) to check for
primer dimers and a No-RT-Control (NRT) to check for any genomic DNA contamination.
310
311
Sciatic Nerves:
500 ng of total RNA was reverse transcribed with Random Primers (Promega C1181) and
MMLV Reverse Transcriptase (Sigma M1302) according to prescribed protocols. qPCR was
performed using Absolute SYBR ROX 2X qPCR mix (Thermo AB1162B) as a fluorescent
detection dye. All reactions were carried out in a final volume of 7 µl in 384 well plates with
300 nM gene-specific primers, around 3.5 ng of cDNA (at 100% RT efficiency) and 1X SYBR
Master Mix in each well. Each reaction was performed in triplicates. All qPCR experiments

were performed on BioRad CFX384 with a No-Template-Control (NTC) to check for primer
dimers and a No-RT-Control (NRT) to check for any genomic DNA contamination.
321

qPCR statistical analysis and data visualization:

qPCR readouts were analyzed in Precision Melt Analysis Software v1.2 (Sciatic nerve
samples) and QuantStudio Real-Time PCR Software v1.7.1 (iPSC samples). The amplicons
were subjected to Melt Curve analysis and were verified for a single dissociation peak at a
Melting Temperature (Tm) > 75°C as expected from the primer constructs. The Cq data was
exported to Microsoft Excel for further calculations. Each biological sample had three technical
replicates thereby generating three individual Cq values. The arithmetic mean of the triplicates
was taken to be the Cq representing the biological sample. The standard deviation (SD) of the
triplicates was also calculated and samples that exhibited SD > 0.20 were considered
inconsistent. In such cases, one outlier Cq was removed to have at least duplicate Cq values
for each biological sample and an SD < 0.20.

Reference gene validation was performed according to our qPCR data analysis workflow
(Sundaram *et al*, 2019). Visual representation of potential intrinsic variation in reference genes
was identified by plotting the raw expression profiles ($2^{-\Delta Cq}$) of all candidate reference genes.
Due to reduced sample sizes, a non-parametric Mann Whitney U-Test was performed to assess
statistically significant expression variation between experimental groups. The alpha value was
set at 0.05 for statistical significance. The genes were then screened using Coefficient of
variation (CV) analysis to eliminate genes that exhibited CV>50% as they could impede the
robustness of the Normfinder algorithm (Sundaram *et al*, 2019). Normfinder was then used to
determine the best pair of reference genes (least S value) to compute the Normalization Factor
(NF) for qPCR data normalization (Andersen *et al*, 2004). From the resulting pair of stable
reference genes, the NF was then calculated as the arithmetic mean of the Cq values of the 2
genes for each sample. Relative expression of target genes was quantified using the $2^{-\Delta\Delta Ct}$
method using the NF as the reference gene (Schmittgen & Livak, 2008; Livak & Schmittgen,
2001). Data was then analysed and visualised using R studio. For the fold change comparison
between qPCRs and RNA-Seq, the non-parametric Kruskal-Wallis ANOVA was performed.
Multiple group comparisons were performed using the non-parametric Dunn's Post Test. The
alpha value was set at 0.05 for statistical significance.

Results:

Reference gene selection from RNA-Seq of microglia (WT vs TREM2KO)

A growing body of literature promotes the use of RNA-Seq to identify stable reference genes for qPCR data normalization in different experimental conditions. However, we believe that this approach is superfluous and that stable reference genes can readily be identified by starting out with any random set of candidates and adopting the right statistical approach to determine the best reference genes for a given experimental condition. To test this hypothesis, we first performed paired-end bulk RNA-Seq on iPSC-derived WT and TREM2KO microglia, an experimental model used to study the role of microglia in neuroinflammation and neurodegeneration (McQuade *et al*, 2020; Andreone *et al*, 2020). Based on the criteria detailed in the Materials and Methods section (see *Reference gene selection from RNA-Seq*), we shortlisted 10 candidate reference genes (*ANXA7, APIP, CCNT2, CNBP, DDX42, FOXK1, KIF13A, MRPL37, PPP1R10,* and *USP5*) from our RNA-Seq data. The base mean values of these genes and other RNA-Seq data features are detailed in Supplementary Table 1.

Reference gene validation of RNA-Seq derived reference genes (WT vs TREM2KO)

The reference genes selected from RNA-Seq were then subjected to validation using the qPCR workflow that we developed recently (Sundaram *et al*, 2019). We first computed the intrinsic variation of reference genes by linearizing the Cq values (2^{-Cq}) followed by visually representing the non-normalized expression levels ($2^{-\Delta Cq}$) of the genes in both experimental groups (**Figure 1**). The WT group was used as the experimental calibrator. Subsequently, statistical testing was performed using a non-parametric Mann Whitney Test to determine significant differences between the groups. Among the 10 genes tested, only 2 genes (MRPL37 and PPP1R10) showed significant variation in non-normalized expression levels between the WT and TREM2KO groups (**Figure 1**).

*Figure 1. Non-normalized expression profiles ($2^{-\Delta Cq}$) of reference genes derived from RNA-Seq data of iPSC derived microglia. WT group was used as the experimental calibrator. (a) ANXA7, (b) APIP, (c) CCNT2, (d) FOXK1, (e) KIF13A, (f) USP5, (g) CNBP, (h) DDX42, (i) MRPL37, (j) PP1R10 and (k) Normalization Factor (NF) assessed by NormFinder that combines CNBP & KIF13A. Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05.*

In accordance with our workflow, we next performed Coefficient of Variation analysis on linearized Cq values (2^{-Cq}) to assess the overall variation (both groups included) of the reference genes (**Table 1**). The genes tested exhibited CV values between 5.99% and 15.75% and they were ranked as shown in **Table 1**. As all genes exhibited CV<50%, they could therefore be screened using the NormFinder algorithm to determine the best combination of

reference genes to be used for data normalization. This is a crucial step of reference gene validation as genes that exhibit CV>50% can compromise the robustness of the algorithm (Sundaram *et al*, 2019)**.** NormFinder determines the stability of a reference gene by factoring the intergroup as well as the intragroup variation and computes a Stability S Score (Andersen *et al*, 2004). The lower the S score, the higher the stability of the gene across all experimental groups. The algorithm also suggests the best pair of reference genes that can be used for robust data normalization. NormFinder results of the selected reference genes from RNA-seq are represented in **Table 1** along with their respective S Scores and ranks. The algorithm further revealed CNBP and KIF13A to be the best combination of genes with a grouped Stability S score of 0.05 (**Table 1**). The Cq values of these two genes were then combined to compute the Normalization Factor (NF). Evidently and as expected, the non-normalized expression level of the NF does not vary significantly between the two groups (**Figure 1K**). It is to be noted that the combination of the two best reference genes exhibits lesser intergroup and intragroup variation (**Figure 1K versus the rest**) including the top ranked CNBP (**Figure 1G**). This is an inherent feature of the algorithm.

Table 1. CV and NormFinder analysis

Expression stability of candidate reference genes derived from bulk RNA-Seq data in iPSC-derived microglia were evaluated using Coefficient of Variation (CV) Analysis & NormFinder.

CV Analysis			NormFinder		
Gene	**%CV**	**Rank**	**Gene**	**Stability S**	**Rank**
CNBP	5.99	1	*CNBP*	0.07	1
KIF13A	6.36	2	*CCNT2*	0.08	2
PPP1R10	6.87	3	*KIF13A*	0.1	3
CCNT2	7.46	4	*PPP1R10*	0.1	4
MRPL37	7.74	5	*USP5*	0.1	5
ANXA7	10.58	6	*MRPL37*	0.11	6
USP5	10.81	7	*DDX42*	0.12	7
DDX42	11.41	8	*ANXA7*	0.15	8

APIP	14.61	9	*FOXK1*	0.15	9
FOXK1	15.75	10	*APIP*	0.16	10

Normfinder Best Pair: CNBP/KIF13A
Grouped Stability: 0.05

Reference gene validation of conventional human reference genes (WT vs TREM2KO)

In addition to the reference genes derived from the RNA-Seq data, we next sought to validate commonly used human reference genes chosen arbitrarily. We selected 10 reference genes (*ACTB*, *GAPDH*, *GUSB*, *PPIA*, *RPLP0*, *TBP*, *HPRT*, *PGK1*, *UBC*, *and RPL13A*) and assessed their intergroup variation by plotting their non-normalized profiles (**Figure 2**)**.** This was followed by statistical testing for significant differences between the two experimental groups. Among the 10 genes tested, none of the genes exhibited significant variation in their non-normalized expression levels (**Figure 2**).

Figure 2. Non-normalized expression profiles ($2^{-\Delta Cq}$) of conventional human reference genes in iPSC derived microglia. WT group is used as the experimental calibrator. (a) *ACTB*, (b) *GAPDH*, (c) *GUSB*, (d) *PPIA*, (e) *RPLP0*, (f) *TBP*, (g) *HPRT*, (h) *PGK1*, (i) *UBC*, (j) *RPL13A*, and (k) *Normalization Factor (NF) assessed by NormFinder that combines RPLP0 & GUSB. Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

These genes were then subjected to CV analysis to assess the overall variation (**Table 2**). They exhibited CV values ranging from 5.29% to 32.37%. However, all genes satisfied the CV<50% criterion for NormFinder screening. NormFinder results for these genes are also shown in **Table 2.** Interestingly and in contrast to the reference genes derived from the RNA-Seq data (**Table 1**), the Stability S score of the conventional genes are largely lower than the genes derived from RNA-Seq, thus indicating better stability. The first 6 ranks from RPL13A to

GAPDH have S scores less than 0.1 whereas in the previous dataset (**Table 1**), only the first 2 ranks (CNBP & CCNT2) exhibit scores less than 0.1. Finally, among all the conventional reference genes screened, NormFinder suggested the use of RPLP0 and GUSB as the best pair for data normalization with a grouped stability of 0.03 (**Table 2**); which in comparison, is lesser than the Stability Score of 0.05 from CNBP and KIF13A (**Table 1**). In principle, these results show that randomly selected reference genes could potentially exhibit better stability than stable reference genes filtered from RNA-Seq data.

Table 2: CV and NormFinder analysis of conventional reference genes. Expression stability of randomly chosen candidate reference genes were evaluated using Coefficient of Variation (CV) Analysis, NormFinder. Best pair chosen is *RPLP0* and *GUSB* with a Stability S score of 0.03.

CV Analysis			NormFinder		
Gene	**%CV**	**Rank**	**Gene**	**Stability S**	**Rank**
GUSB	5.29	1	*RPL13A*	0.03	1
TBP	7.56	2	*GUSB*	0.04	2
RPL13A	8.16	3	*RPLP0*	0.05	3
RPLP0	8.18	4	*TBP*	0.05	4
PGK1	9.56	5	*PGK1*	0.06	5
GAPDH	11.35	6	*GAPDH*	0.07	6
PPIA	13.75	7	*PPIA*	0.10	7
ACTB	14.31	8	*UBC*	0.11	8
UBC	17.47	9	*ACTB*	0.14	9
HPRT	32.27	10	*HPRT*	0.18	10

NormFinder Best Pair: RPLP0/GUSB
Grouped Stability S: 0.03

Differential expression of target genes by qPCR and RNA-Seq (WT vs TREM2KO)

Following reference gene validation of conventional and RNA-Seq-derived candidates, we picked 6 target genes that were differentially expressed based on the RNA-Seq results (*see Target gene selection* from *Materials & Methods*). We included 3 target genes (*TREM2, P2RY12* and *CD52*) that were significantly downregulated and 3 others (*ACSS3, SYDE1* and *TBX10*) that were significantly upregulated. The relative expression of these genes was then computed by qPCR using the Normalization Factor (NF) from conventional candidate reference genes (*RPLP0 & GUSB* or Conventional NF) or from RNA-Seq-derived candidate reference genes (*CNBP & KIF13A* or RNA-Seq derived NF) (**Figure 3**). To compare our qPCR results, we also computed the fold change assessed by RNA-Seq using the Normalized counts of these genes.

Regarding the downregulated genes, *TREM2* expression in the TREM2KO group, when assessed through qPCRs, exhibited expression fold change values of 0.31 ± 0.05 (mean FC $\pm$ SD, conventional NF, **Figure 3A**) and 0.35 ± 0.04 (RNA-Seq derived NF, **Figure 3B**). In RNA-Seq, the fold change values were 0.23 ± 0.03 (**Figure 3C**). *P2RY12* fold change values in the TREM2KO group were 0.04 ± 0.02 when assessed using qPCRs (both conventional NF and RNA-Seq derived NF, **Figure 3D and 3E**). However, RNA-Seq fold change values of *P2RY12* were comparatively higher at 0.62 ± 0.19 (**Figure 3F**). *CD52* fold change in the TREM2KO group when assessed by qPCR were 0.38 ± 0.06 (Conventional NF, **Figure 3G**) and 0.40 ± 0.05 (RNA-Seq derived NF, **Figure 3H**). RNA-Seq fold changes were comparatively lower at 0.16 ± 0.01 (**Figure 3I**).

For upregulated genes, *ACSS3* in the TREM2KO group exhibited fold change values of 3.72 ± 0.79 (Conventional NF, **Figure 3J**) and 3.40 ± 0.88 (RNA-Seq derived NF, **Figure 3K**). The fold changes in RNA-Seq were 4.54 ± 1.25 (**Figure 3L**). *SYDE1* fold change values assessed by qPCR were at 3.58 ± 1.18 (Conventional NF, **Figure 3M**) and 3.29 ± 1.28 (RNA-Seq derived NF, **Figure 3N**). RNA-Seq expression levels of *SYDE1* were at 3.62 ± 0.34 (**Figure 3O**). Finally, *TBX10* expression fold changes assessed by qPCR were at 3.06 ± 1.51 (Conventional NF, **Figure 3P**) and 2.82 ± 1.60 (RNA-Seq derived NF, **Figure 3Q**). RNA-Seq fold changes were at 3.24 ± 0.61 (**Figure 3R**). These results taken together with the visual representation of differential expression described in **Figure 3** show that qPCR normalization using conventional NF or RNA-Seq derived NF render the same results. They do not always concur with RNA-Seq fold changes in magnitude but do so in tendency.

Figure 3: Differential expression of target genes between WT and TREM2KO iPSC cells assessed by qPCRs and RNA-Seq. *The target genes (TREM2, P2RY12, CD52, ACSS3, SYDE1, TBX10) have been normalized with the normalization factor (NF) computed from conventional reference genes RPLP0 & GUSB in the first column (A, D, G, J, M, P) or from RNA-Seq derived reference genes CNBP & KIF13A in the second column (B, E, H, K, N, Q). The third column contains the Fold Changes computed from RNA-Seq (C, F, I, L, O, R). The normalized counts for each gene in each group were used to calculate the fold changes in RNA-Seq. The WT group was used as the experimental control. The P_{adj} value of RNA-Seq is indicated in the TREM2KO group for each gene in the RNA-Seq column. For the qPCRs, Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05.*

Comparison of fold changes between qPCRs and RNA-Seq (WT vs TREM2KO):

We next investigated in detail if the Fold Change distributions in the TREM2KO experimental group computed across the 3 methods differed significantly from one another. To this end, we compared the distributions by a non-parametric ANOVA (Kruskal Wallis Test) of ordinal distributions followed by Dunn's multiple comparison post-test (**Supplementary Figure 1**). qPCR Fold changes computed using Conventional NF were used as the control for multiple comparisons. Therefore, the first comparison was made between the conventional and RNA-Seq derived NF groups (difference in qPCRs, hereafter referred to as Comparison 1) and the second comparison was made between the Conventional NF group and RNA-Seq differential expression (difference between qPCR and RNA-Seq, hereafter referred to as Comparison 2).

The WT experimental group was omitted from this analysis as it was used as the experimental calibrator for differential expression. The mean Fold Change of the WT group is always at 1 regardless of the gene/method in question and therefore it is redundant to test for statistical significance of the WT fold change levels across different methods for each gene.

Non-parametric ANOVA revealed no significant difference between the FC distributions of the upregulated genes *ACSS3, SYDE1* and *TBX1* (**Supplementary Figure 1D, 1E and 1F**) suggesting that the qPCR fold changes concur with RNA-Seq fold changes in these genes. As for the downregulated genes, *TREM2* FC distributions varied significantly between the groups (Kruskal Wallis $P < 0.5$); however, Dunn's post test revealed no significant difference in Comparison 1 and Comparison 2 (**Supplementary Figure 1A**). *P2RY12* FC distributions varied significantly between the groups (Kruskal Wallis $P<0.01$). Multiple comparisons revealed no difference in Comparison 1 but a significant difference in Comparison 2 (Dunn's Post Test $P<0.05$) (**Supplementary Figure 1B**) CD*52* FC distributions varied significantly between the groups (Kruskal Wallis $P < 0.5$). Multiple comparisons, yet again, revealed no difference in Comparison 1 but a significant difference in Comparison 2 (Dunn's Post Test $P<0.05$) (**Supplementary Figure 1C**). These results show firstly that no significant difference exists between the FC distributions of all the genes between the qPCRs (Comparison 1). Secondly, FC distributions of qPCRs and RNA-Seq do not show any difference for the upregulated genes *ACSS3*, *SYDE1* and *TBX10* (Comparison 2). However, for the downregulated genes, although no differences exist between qPCRs (Comparison 1); *P2RY12* & *CD52* FC distributions differ significantly between qPCR and RNA-Seq. In summation, these results prove beyond reasonable doubt that differential expression assessed by qPCR using randomly chosen conventional reference genes and "stable" reference genes filtered from RNA-Seq render similar expression profiles of target genes without any significant difference among the two normalization strategies. However, concordance with RNA-Seq results was only observed in the upregulated genes and could not be achieved in the downregulated genes.

Reference gene selection from RNA-Seq of mouse sciatic nerves (P3 vs P21)

To rule out any circumstantial evidence that may support our observations in the iPSC dataset, we sought to validate our results using a completely different experimental setting. We wanted to rule out the possibility that the conventional reference genes that we chose in the iPSC dataset were indeed stable reference genes by mere happenstance and thus they rendered the same profiles of differential expression as the reference genes filtered from RNA-Seq.

Therefore, in a confirmatory experimental approach we mined and re-analyzed RNA-Seq data that was previously generated from mouse sciatic nerves at Postnatal day 3 (P3) and P21 (Ackerman *et al*, 2018). This dataset permits us to study RNA expression changes in the sciatic nerves during postnatal myelination. We applied the same criteria to select 10 candidate reference genes (*Ppp3ca, Fkbp4, Vcp, Lama2, Ank2, Coq9, Chmp2a, Laptm5, Leprotl1, Supt4a*). The basemean values of these genes and other RNA-Seq data features are detailed in Supplementary Table 2.

Reference gene validation of RNA-Seq derived reference genes from mouse sciatic nerves (P3 vs P21)

We evaluated the intrinsic intergroup variation of these reference genes by visual representation of their non-normalized profiles ($2^{-\Delta Cq}$) using P3 samples as the calibrator (Figure 4). As previously described, statistical inference was then assessed using the non-parametric Mann Whitney U test owing to reduced sample sizes. Interestingly, and in contrast to the iPSC dataset, 6 out of 10 genes (*Fkbp4, Lama2, Coq9, Chmp2a, Leprotl1 & Supt4a*) exhibit significant intergroup variation across the two experimental groups (**Figure 4A, 4B, 4C, 4D, 4E & 4F**).

Figure 4. Raw expression profiles of reference genes obtained from RNA-Seq analysis of developing sciatic nerve of age Post-natal day 3 (P5) and 21 (P21), expressed as fold changes (2-ΔCq). P5 group is the experimental calibrator. (A) *Fkbp4*, (B) *Lama2*, (C) *Leprotl4*, (D) *Supt4a*, (E) *Coq9*, (F) *Chmp2a*, (G) *Ank2*, (H) *Vcp*, (I) *Laptm5*, (J) *Ppp3ca*, and (K) Normalisation Factor (*Ppp3ca + Coq9*). *Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

We next performed the CV analysis to assess overall variation (**Table 3**). The genes tested exhibited CV values between 12.32% and 36.24% and they were ranked as shown in **Table 3**. As all genes exhibited CV<50%, they could therefore be screened using the NormFinder algorithm. NormFinder analysis further revealed *Ppp3ca* and *Coq9* to be the best combination of genes with a grouped Stability S score of 0.04 (**Table 3**). Although Coq9 exhibited significant intergroup variation (**Figure 4E**), the combination of the gene with *Ppp3ca*

resulting in the Normalization Factor (NF) does not exhibit any (**Figure 4K**). Interestingly, the
RNA-Seq data features of the 6 genes that exhibit significant intergroup variation (**Figure 4A,**
4B, 4C, 4D, 4E & 4F) show that they are not differentially expressed. However, if these genes
were to be normalized using the NF computed herein, some of them could potentially exhibit
differential expression in qPCR as the NF is stable across experimental groups (natural
deduction, data not shown). It is a striking example to demonstrate the discordance of RNA-
Seq and qPCR results in this dataset. However, the computation of a stable NF despite the
presence of genes with significant intrinsic variation confirms the strength of the NormFinder
algorithm and the validity of using the two best reference genes proposed by the method.

Table 3. CV and NormFinder analysis of reference genes derived from RNA-Seq

CV Analysis			NormFinder		
Gene	%CV	Rank	Gene	Stability S	Rank
Laptm5	12.32	1	*Ank2*	0.07	1
Ppp3ca	12.96	2	*Ppp3ca*	0.08	2
Supt4a	14.79	3	*Coq9*	0.08	3
Coq9	16.26	4	*Laptm5*	0.16	4
Ank2	16.31	5	*Supt4a*	0.16	5
Vcp	19.59	6	*Leprotl1*	0.23	6
Leprotl1	24.49	7	*Fkbp4*	0.26	7
Fkbp4	25.86	8	*Vcp*	0.27	8
Lama2	29.96	9	*Chmp2a*	0.47	9
Chmp2a	36.24	10	*Lama2*	0.56	10

584
585 *Normfinder Best combination: Ppp3ca + Coq9*
586 *Grouped Stability: 0.04*
587

Reference gene validation of conventional mouse reference genes (P3 vs P21)

To obtain stable reference genes from a set of conventional genes, we used 10 candidate mouse reference genes that we had previously used to establish our workflow (*Actb, Gapdh, Tbp, Sdha, Pgk1, Ppia, Rpl13a, Hsp60, Mrpl10, Rps26*) (Sundaram *et al*, 2019). As explained above, visual representation followed by inferential statistics were performed and the results are described in **Figure 5**. 4 out of 10 genes (*Gapdh, Sdha, Pgk1 & Rps26*) exhibited significant intrinsic variation between the experimental groups (**Figure 5G ,5H, 5I & 5J**)

Figure 5. Raw expression profiles of conventional reference genes of developing sciatic nerve of age Post-natal day 3 (P3) and 21 (P21), expressed as fold changes ($2^{-\Delta Cq}$). P3 group is the experimental calibrator. (A) Actb, (B) Mrpl10, (C) Hsp60, (D) Ppia, (E) Rpl13a, (F) Tbp, (G) Gapdh, (H) Pgk1, (I) Rps26, (J) Sdha, and (K) Normalisation Factor (Hsp60 + Ppia). Non-parametric Mann Whitney U test was used to assess differences between the groups. The

CV analysis was then performed on these genes and the results are depicted in **Table 4**. CV
values ranged from 12.3% to 30.44% and the genes were ranked accordingly. As all genes
exhibited CV<50%, we screened these genes through the NormFinder algorithm to determine
the best combination of reference genes for qPCR normalization. NormFinder determined
Hsp60 and *Ppia* to be the best combination of reference genes with a grouped Stability S Score
of 0.06 (**Table 6**). Both *Hsp60* (**Figure 5C**) & *Ppia* (**Figure 5D**) did not exhibit any intergroup
variation and the same was true for the NF computed by combining these genes (**Figure 5K**).

Table 4. CV and NormFinder analysis of conventional reference genes

CV Analysis			NormFinder		
Gene	**%CV**	**Rank**	**Gene**	**Stability S**	**Rank**
Mrpl10	12.30	1	*Tbp*	0.03	1
Gapdh	13.06	2	*Mrpl10*	0.04	2
Hsp60	14.36	3	*Hsp60*	0.05	3
Tbp	15.24	4	*Ppia*	0.05	4
Pgk1	15.83	5	*Pgk1*	0.06	5
Ppia	19.36	6	*Actb*	0.07	6
Actb	21.07	7	*Rpl13a*	0.10	7
Rps26	22.04	8	*Gapdh*	0.11	8
Rpl13a	26.78	9	*Rps26*	0.14	9
Sdha	30.44	10	*Sdha*	0.18	10

612
613   *Normfinder Best combination: Hsp60 + Ppia*
614   *Grouped stability: 0.06*
615

Differential expression of target genes by qPCR and RNA-Seq (P3 vs P21)

Following reference gene validation of conventional and RNAseq-derived candidates, we picked 6 target genes that were known to be differentially expressed during postnatal development of the sciatic nerve (Gerber *et al*, 2021). 3 target genes (*Mpz, Mbp and Cd90*) that were chosen are significantly upregulated and the other 3 target genes (*p75NTR, Mki67* and *Sox2*) are significantly downregulated at these time points. The relative expression of these genes was then computed by qPCR using the Normalization Factor (NF) from conventional candidate reference genes (*Hsp60 & Ppia* or Conventional NF) or from RNAseq-derived candidate reference genes (*Ppp3ca & Coq9* or RNAseq derived NF) (**Figure 6**). To compare our qPCR results, we also computed the fold change assessed by RNA-Seq using the Normalized counts of these genes.

Regarding the downregulated genes, *Mki67* expression in the P21 group, when assessed through qPCRs, exhibited fold change values of 0.14 ± 0.10 (mean FC $\pm$ SD, conventional NF, **Figure 6A**) and 0.12 ± 0.08 (RNA-Seq derived NF, **Figure 6B**). In RNA-Seq, the fold change values were 0.08 ± 0.01 (**Figure 6C**). *p75NTR* expression fold change in the P21 group were 0.30 ± 0.04 (conventional NF, **Figure 6D**) and 0.26 ± 0.02 (RNA-Seq derived NF, **Figure 6E**) when assessed using qPCRs. However, RNA-Seq fold change values of *p75NTR* were comparatively higher at 0.63 ± 0.18 (**Figure 6F**). *Sox2* expression fold change in the P21 group when assessed by qPCR were 0.49 ± 0.05 (Conventional NF, **Figure 6G**) and 0.42 ± 0.03 (RNA-Seq derived NF, **Figure 6H**). RNA-Seq changes were at 0.39 ± 0.09 (**Figure 6I**).

For the upregulated genes, *Mbp* expression levels in the P21 group exhibited fold change values of 1.73 ± 0.15 (Conventional NF, **Figure 6J**) and 1.49 ± 0.19s (RNA-Seq derived NF, **Figure 6K**). The fold changes in RNA-Seq were 2.02 ± 0.62 (**Figure 6L**). *Mpz* expression levels assessed by qPCR were at 1.76 ± 0.38 (Conventional NF, **Figure 6M**) and 1.53 ± 0.35 (RNA-Seq derived NF, **Figure 6N**). However, RNA-Seq expression fold change of *Mpz* were significantly higher at 3.49 ± 0.26 (**Figure 6O**). Finally, *Cd90* expression fold change assessed by qPCR were at 1.83 ± 0.35 (Conventional NF, **Figure 6P**) and 1.57 ± 0.30 (RNA-Seq derived NF, **Figure 6Q**). RNA-Seq fold changes were at 4.81 ± 2.77 (**Figure 6R**). Yet again, these results taken together with the visual representation of differential expression described in **Figure 6** show that qPCR normalization using conventional NF or RNA-Seq derived NF render very similar results. We observe again that they do not always concur with RNA-Seq fold changes in magnitude, but they do so in tendency.

Figure 6: Differential expression of target genes in P3 and P21 mouse sciatic nerves by qPCRs and RNA-Seq. *The target genes (Mki67, p75NTR, Sox2, Mbp, Mpz, Cd90) have been normalized with the normalization factor (NF) computed from conventional reference genes Hsp60 & Ppia in the first column (A, D, G, J, M, P) or from RNA-Seq derived reference genes Ppp3ca and Coq9 in the second column (B, E, H, K, N, Q). The third column contains the Fold Changes computed from RNA-Seq (C,F,I,L,O,R). The normalized counts for each gene in each group were used to calculate the fold changes in RNA-Seq. The WT group was used as the experimental control. The P_{adj} value of RNA-Seq is indicated in the P21 group for each gene*

in the RNA-Seq column. For the qPCRs, Non-parametric Mann Whitney U test was used to
assess differences between the groups. The alpha value was set at 0.05 and P values are
*annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

Comparison of differential expression between qPCRs and RNA-Seq (P3 vs P21):

Similar to the analysis performed in the iPSC dataset, we next investigated in detail if the Fold
Change distributions in the P21 experimental group computed across the 3 methods differed
significantly from one another. We used non-parametric ANOVA (Kruskal Wallis Test) of
ordinal distributions followed by Dunn's multiple comparison post-test (**Supplementary**
Figure 2). As shown in the iPSC dataset, we performed multiple comparisons between qPCRs
(Comparison 1) and also between qPCR and RNA-Seq (Comparison 2).

Non-parametric ANOVA revealed no significant difference between the FC distributions of
Mki67 and *Mbp* (**Supplementary Figure 2A and 2E**) suggesting that the qPCR fold changes
concur with RNA-Seq fold changes in these genes. *Sox2, p75NTR* and *Mpz* FC distributions
varied significantly between the groups (Kruskal Wallis $P < 0.05$, $P < 0.01$ and $P<0.01$
respectively); however, Dunn's post test revealed no significant difference in both Comparison
1 and Comparison 2 (**Supplementary Figure 2B, 2C & 2F**). *Cd90* FC distributions varied
significantly between the groups (Kruskal Wallis $P<0.01$). Multiple comparisons revealed no
difference in Comparison 1 and Comparison 2 (**Supplementary Figure 2D**). In summation,
all the genes tested in the sciatic nerve dataset show no significant differences between
Conventional NF and RNA-Seq-derived NF results. In summation, these results show once
again that stable reference genes filtered from RNA-Seq data do not give any added advantage
to qPCR data normalization and that not all genes exhibit comparable Fold Changes when
comparing qPCR results and RNA-Seq.

Discussion:

The central premise of this article was to demonstrate that stable reference genes for qPCR data normalisation can be obtained from any random set of candidates provided the statistical approach of reference gene validation is sound and consistent. To this end, using our qPCR workflow we have shown using two separate and unique datasets that randomly chosen reference genes can render the same results of differential expression when compared to stable references selected from RNA-Seq data.

The advantage of this approach in filtering the best candidates from any random set lies in combining 3 methods that are complementary to each other. Although multiple statistical methods have been proposed before, we have previously highlighted the potential pitfalls and assumptions of these computational methods (Sundaram *et al*, 2019). We thus devised this workflow combining the well-known NormFinder method with CV analysis and the visual representation of non-normalized expression followed by statistical testing. Indeed, it should be noted the core method of this approach is the NormFinder algorithm. However, this method was first described and validated with larger sample sizes (typically between 10 - 20 or more) and with a high number of candidate genes chosen from microarrays (typically between 15 - 20) exhibiting very little overall variation (Andersen *et al*, 2004). In our previous study, we identified that lower sample sizes and candidate genes with high overall variation can skew the results of this method because of the algorithm's construct (Sundaram *et al*, 2019). Although scores of studies have used this algorithm without due diligence to the prerequisites, our combined approach resolves these issues. The CV analysis helps in the identification of genes with high overall variance and their successive elimination from analysis gives rise to more robust results from NormFinder. Visual representation of the non-normalized profiles ($2^{-\Delta Cq}$) followed by statistical testing further validates the results of NormFinder as the Normalization Factor computed from the two best reference genes does not exhibit any significant differences between the experimental groups regardless of experimental conditions (**Figure 1K, 2K, 4K & 5K**). Thus, in combining CV analysis and statistical testing of non-normalized profiles, we extrapolated the NormFinder approach to be used for experimental setups involving lower sample sizes similar to those used in this study.

The cut off at 50% CV to qualify for NormFinder analysis is rooted in the fact that at 50%, the standard deviation of the expression distribution is half the mean expression level. Previous datasets that we published have indicated that the Normalization Factor (NF) computed by NormFinder when including genes that exhibit CV>50% exhibit significant intergroup variation in the non-normalized profiles (Sundaram *et al*, 2019). Below this cut off, NF expression levels are stable across experimental groups. Indeed, the user can be more stringent by using a lower CV cut off but our data from this study and the previous one suggests that this is not necessary. However, this criterion gives rise to a theoretical limitation of our approach. In a given experimental setup, if all the candidate reference genes tested exhibit CV values of above 50%, then our approach will fail and NormFinder results would no longer be reliable. In such cases, a new list of candidates is required. However, our experience suggests

723 that as little as 6 genes after eliminating genes exhibiting CV>50% can still provide a sound
724 NF for qPCR data normalization (Sundaram *et al*, 2019).
725

Although not the primary objective of the study, the comparison of qPCR results with
RNA-Seq merits discussion. In the iPSC dataset, comparisons made between qPCR and RNA-
Seq results revealed that 4 out of 6 genes (*ACSS3, SYDE1, TREM2* and *TBX10*) showed no
significant difference between the fold changes computed by both the methods
(**Supplementary Figures 1D, 1E, 1F & 1C**). However, changes in the expression of *CD52*
and *P2RY12* significantly differed between the two methods (**Figure 3 and Supplementary**
Figure 1A and 1B). In the sciatic nerve dataset, 5 out of 6 genes tested show no difference
between qPCR and RNA-Seq data in multiple comparisons (**Supplementary Figure 2A, 2B,**
2C, 2E, 2F). However, this analysis is not optimal as the number of samples used in qPCR is
much higher (N=7) than the number of samples in RNA-Seq data that we mined (N=3). In
reality, only 3 genes (*Mki67, Sox2 & Mbp*) exhibited overlapping fold change distributions
across the methods (**Supplementary Figure 2A, 2B & 2E**). However, regardless of the
magnitudes of the change observed, the pattern of expression changes was always conserved
across qPCR and RNA-Seq for the genes that we tested. Indeed, it is possible that if we include
more target genes, we can potentially notice disagreements in expression patterns as well.
However, as stated earlier, our objective is restricted to demonstrating the futility of selecting
reference genes from RNA-Seq data for qPCR data normalization. The comparison of the two
methodologies falls beyond the scope of this study but it has been amply addressed elsewhere
(Griffith *et al*, 2010; Rajkumar *et al*, 2015; Everaert *et al*, 2017).
745

These studies have indeed tried to validate RNA-Seq data with RTqPCR for a large set
of target genes typically ranging from a few hundred to the entire transcriptome. The consensus
from these studies is that the extent of correlation between RTqPCR and RNAseq data is
dependent on how the sequencing data was aligned, mapped and counted based on existing
protocols. Additionally, though not discussed in these articles, we believe that this would
equally depend on how reference genes were selected for qPCR assays that validate the
RNAseq data. Overlooking these technicalities, these studies also show that around 85% of all
genes tested show concordant differential expression (comparable magnitudes with the same
tendency) between the two methodologies with correlation coefficients typically above 0.8.
Although these numbers are relatively reassuring at the outset, from an absolute perspective,
the implications raise important concerns. For a hypothetical bulk RNA-Seq dataset of around
18000 protein-coding transcripts, 85% concordance leaves 2700 genes whose differential
expression is discordant between RNA-Seq and qPCR. Of note, the study conducted by
Everaert and colleagues (Everaert *et al*, 2017) conclude that discordant genes typically are
smaller, have fewer exons and are expressed at lower levels. Indeed, and as mentioned in the
introduction, these observations can be possibly substantiated by the transcript-length bias and
the bias against genes with lower expression in RNA-Seq. In our study, the discordant genes
CD52 and *P2RY12* in the microglia dataset have only 2 exons. The other 4 concordant genes
have between 4 – 13 exons and therefore are relatively larger. The basemean values (and
therefore the expression levels) of the 2 discordant genes are largely higher than the 4
concordant genes. Therefore, number of exons appears to be more important than expression

levels for concordance between qPCR and RNA-Seq for these target genes. On the contrary,
we did not observe any correlation between discordance and small gene size or lower
expression in the sciatic nerve samples. These observations show that an objective approach to
identify discordant genes is lacking in literature. As RNA-Seq data analysis is particularly
prone to certain biases while computing differential gene expression, it would be valuable to
computationally suggest qPCR validation for specific genes that are susceptible to these
normalization biases based on their transcript length, expression levels and other unidentified
parameters. Ideally, these suggestions should be integrated into the data analysis pipelines and
represented in the differential gene expression data frames. This would provide the foundation
for an objective and structured validation of Sequencing data rather than an obtuse overall
validation of RNA-Seq results by qPCR which is indeed burdensome. If these computational
suggestions are achieved, the methodology used in our study would aptly address the necessity
of performing robust and reproducible qPCR assays using any random set of reference genes
that are independent of RNA-Seq results.

References:

Ackerman SD, Luo R, Poitelon Y, Mogha A, Harty BL, D'Rozario M, Sanchez NE,
Lakkaraju AKK, Gamble P, Li J, *et al* (2018) GPR56/ADGRG1 regulates development
and maintenance of peripheral myelin. *J Exp Med* 215: 941–961
Andersen C, Jensen J & Orntoft T (2004) Normalization of Real - Time Quantitative Reverse
Transcription - PCR Data : A Model - Based Variance Estimation Approach to Identify
Genes Suited for Normalization , Applied to Bladder and Colon Cancer Data Sets.
Cancer Res 64: 5245
Andreone BJ, Przybyla L, Llapashtica C, Rana A, Davis SS, van Lengerich B, Lin K, Shi J,
Mei Y, Astarita G, *et al* (2020) Alzheimer's-associated PLCγ2 is a signaling node
required for both TREM2 function and the inflammatory response in human microglia.
Nat Neurosci 23: 927–938
Berge K Van den, Hembach KM, Soneson C, Tiberi S, Clement L, Love MI, Patro R &
Robinson MD (2019) RNA Sequencing Data: Hitchhiker's Guide to Expression
Analysis. *https://doi.org/101146/annurev-biodatasci-072018-021255* 2: 139–173
Boda E, Pini A, Hoxha E, Parolisi R & Tempia F (2009) Selection of Reference Genes for
Quantitative Real-time RT-PCR Studies in Mouse Brain. *J Mol Neurosci* 37: 238–253
Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T,
Pfaffl MW, Shipley GL, *et al* (2009) The MIQE Guidelines: Minimum Information for
Publication of Quantitative Real-Time PCR Experiments.
Carmona R, Arroyo M, Jiménez-Quesada MJ, Seoane P, Zafra A, Larrosa R, Alché J de D &
Claros MG (2017) Automated identification of reference genes based on RNA-seq data.
Biomed Eng Online 16
Chapman JR & Waldenström J (2015) With Reference to Reference Genes: A Systematic
Review of Endogenous Controls in Gene Expression Studies. *PLoS One* 10: e0141853
Chim SSC, Wong KKW, Chung CYL, Lam SKW, Kwok JSL, Lai C-Y, Cheng YKY, Hui
ASY, Meng M, Chan O-K, *et al* (2017) Systematic Selection of Reference Genes for the
Normalization of Circulating RNA Transcripts in Pregnant Women Based on RNA-Seq
Data. *Int J Mol Sci* 18: 1709
Coenye T (2021) Do results obtained with RNA-sequencing require independent
verification? *Biofilm* 3: 100043
Dai Y, Cao Y, Köhler J, Lu A, Xu S & Wang H (2021) Unbiased RNA-Seq-driven
identification and validation of reference genes for quantitative RT-PCR analyses of
pooled cancer exosomes. *BMC Genomics* 22: 27
Durinck S, Spellman PT, Birney E & Huber W (2009) Mapping Identifiers for the Integration
of Genomic Datasets with the R/Bioconductor package biomaRt. *Nat Protoc* 4: 1184
Eisenberg E & Levanon EY (2013) Human housekeeping genes, revisited. *Trends Genet* 29:
569–74
Everaert C, Luypaert M, Maag JL V, Cheng QX, Dinger ME, Hellemans J & Mestdagh P
(2017) Benchmarking of RNA-sequencing analysis workflows using whole-
transcriptome RT-qPCR expression data. *Sci Rep* 7: 1559
Gerber D, Pereira JA, Gerber J, Tan G, Dimitrieva S, Yánguez E & Suter U (2021)
Transcriptional profiling of mouse peripheral nerves to the single-cell level to build a
sciatic nerve atlas (Snat). *Elife* 10
Griffith M, Griffith OL, Mwenifumbo J, Goya R, Morrissy AS, Morin RD, Corbett R, Tang
MJ, Hou YC, Pugh TJ, *et al* (2010) Alternative expression analysis by RNA sequencing.
Nat Methods 7: 843–847
Haenseler W, Sansom SN, Buchrieser J, Newey SE, Moore CS, Nicholls FJ, Chintawar S,
Schnell C, Antel JP, Allen ND, *et al* (2017) A Highly Efficient Human Pluripotent Stem

Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports* 8: 1727–1742

Hernandez-Segura A, Rubingh R & Demaria M (2019) Identification of stable senescence-associated reference genes. *Aging Cell* 18: e12911

Hoang VLT, Tom LN, Quek X-C, Tan J-M, Payne EJ, Lin LL, Sinnya S, Raphael AP, Lambie D, Frazer IH, *et al* (2017) RNA-seq reveals more consistent reference genes for gene expression studies in human non-melanoma skin cancers. *PeerJ* 5: e3631

Jacob F, Guertler R, Naim S, Nixdorf S, Fedier A, Hacker NF & Heinzelmann-Schwarz V (2013) Careful Selection of Reference Genes Is Required for Reliable Performance of RT-qPCR in Human Normal and Cancer Cell Lines. *PLoS One* 8: e59180

Liu Y, Zhou J & White KP (2014) RNA-seq differential expression studies: more sequence or more replication? *Bioinformatics* 30: 301–304

Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^-ΔΔCT Method. *Methods* 25: 402–408

Love MI, Huber W & Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol 2014 1512* 15: 1–21

Mandelboum S, Manber Z, Elroy-Stein O & Elkon R (2019) Recurrent functional misinterpretation of RNA-seq data caused by sample-specific gene length bias. *PLOS Biol* 17: e3000481

McQuade A, Kang YJ, Hasselmann J, Jairaman A, Sotelo A, Coburn M, Shabestari SK, Chadarevian JP, Fote G, Tu CH, *et al* (2020) Gene expression and functional deficits underlie TREM2-knockout microglia responses in human models of Alzheimer's disease. *Nat Commun* 11: 1–17

Mollet IG, Ben-Dov C, Felício-Silva D, Grosso AR, Eleutério P, Alves R, Staller R, Silva TS & Carmo-Fonseca M (2010) Unconstrained mining of transcript data reveals increased alternative splicing complexity in the human transcriptome. *Nucleic Acids Res* 38: 4740–4754

Oshlack A & Wakefield MJ (2009) Transcript length bias in RNA-seq data confounds systems biology. *Biol Direct 2009 41* 4: 1–10

Pfaffl MW, Tichopad A, Prgomet C & Neuvians TP (2004) Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pair-wise correlations. *Biotechnol Lett* 26: 509–515

Pombo MA, Zheng Y, Fei Z, Martin GB & Rosli HG (2017) Use of RNA-seq data to identify and validate RT-qPCR reference genes for studying the tomato- Pseudomonas pathosystem. *Sci Rep* 7: 44905

Rajkumar AP, Qvist P, Lazarus R, Lescai F, Ju J, Nyegaard M, Mors O, Børglum AD, Li Q & Christensen JH (2015) Experimental validation of methods for differential gene expression analysis and sample pooling in RNA-seq. *BMC Genomics* 16: 1–8

Schmittgen TD & Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3: 1101–8

Silver N, Best S, Jiang J & Thein SL (2006) BMC Molecular Biology Selection of housekeeping genes for gene expression studies in human reticulocytes using real-time PCR.

De Spiegelaere W, Dern-Wieloch J, Weigel R, Schumacher V, Schorle H, Nettersheim D, Bergmann M, Brehm R, Kliesch S, Vandekerckhove L, *et al* (2015) Reference gene validation for RT-qPCR, a note on different available software packages. *PLoS One* 10: 1–13

Sundaram VK, Sampathkumar NK, Massaad C & Grenier J (2019) Optimal use of statistical methods to validate reference gene stability in longitudinal studies. *PLoS One* 14: e0219440

881   Thellin O, Zorzi W, Lakaye B, De Borman B, Coumans B, Hennen G, Grisar T, Igout A &
882   Heinen E (1999) Housekeeping genes as internal standards: use and limits
883   Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A & Speleman F
884   (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric
885   averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034
886   Yang H, Liu J, Huang S, Guo T, Deng L & Hua W (2014) Selection and evaluation of novel
887   reference genes for quantitative reverse transcription PCR (qRT-PCR) based on genome
888   and transcriptome data in Brassica napus L. *Gene* 538: 113–122
889   Zhan C, Zhang Y, Ma J, Wang L, Jiang W, Shi Y & Wang Q (2014) Identification of
890   reference genes for qRT-PCR in human lung squamous-cell carcinoma by RNA-Seq.
891   *Acta Biochim Biophys Sin (Shanghai)* 46: 330–337
892   Zhou Z, Cong P, Tian Y & Zhu Y Using RNA-seq data to select reference genes for
893   normalizing gene expression in apple roots.
894