RNA-Seq is not required to determine stable reference genes for qPCR normalization

Nirmal K Sampathkumar*, ‡, a, b, f, Venkat Krishnan Sundaram*, ‡, c, Prakroothi S Danthi [e],
Rasha Barakat [c d], Shiden Solomon [a, b], Mrityunjoy Mondal [a, b], Ivo Carre [a, b], Tatiana El Jalkh[c],
Aïda Padilla-Ferrer [c], Julien Grenier[c], Charbel Massaad[c], Jacqueline C Mitchell [b].

Affiliations

[a] UK-Dementia Research Institute, King's College London, London, UK.

[b] Department of Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and
Neuroscience, Basic and Clinical Neuroscience Institute, King's College London, London UK.

[c] INSERM UMRS 1124, Université de Paris, Faculty of Basic and Biomedical Sciences, Paris,
France.

[d] INSERM U1016, Institut Cochin, Paris, France.

[e] McGill Group for Suicide Studies, Douglas Mental Health University Institute, McGill
University, Montréal, Canada.

[f] Present address: Alzheimer's Research UK Oxford Drug Discovery Institute, Centre for
Medicines Discovery, University of Oxford, Oxford, UK

‡Correspondence to nirmal.sampathkumar@cmd.ox.ac.uk and
venkatkrishnan.sundaram@u-paris.fr

*These authors contributed equally.

Author contributions:

Conceptualization: NKS, VKS.
Formal analysis: VKS, NKS, PSD
RNA-Seq data analysis pipeline: PSD.
Methodology: NKS, VKS
Experimentation: VKS, NKS, RB, SMS, IC, MM, TEJ, APF.
Validation: JM, CM.
Visualization: JG, MM.
Writing – original draft: NKS, VKS.
Writing – review & editing: All Authors.

Acknowledgements:

The authors thank the Animal House Core Facility and the Cyto2BM Molecular Biology Platform of BioMedTech Facilities (INSERM US36/CNRS UMS2009) for the animals and research services pertaining to the generation of qPCR data. We also thank Benjamin SAINTPIERRE and Dr. Franck LETOURNEUR from the Genomic Platform at Institut Cochin, Paris for the quality control of RNA samples and for the analysis of RNAseq data. NKS would like to thank Prof. Christopher E SHAW and Tanisha LEWIS from the UK Dementia Research Institute, King's College London, for their support towards this work.

Data availability:

The original contributions presented in the study are included in the article & Supplementary Data. The metadata for the RNA-Seq and qPCR experiments can be accessed at https://doi.org/10.6084/m9.figshare.15169104.v1. The datasets available include the normalized count matrices for the RNAseq data, qPCR Cq values and differential expression analysis, primer sequences, Coefficient of Variation and Normfinder analysis.   Further inquiries can be directly addressed to the corresponding author(s).

Funding:

No specific funding for the study. VKS and APF were funded by a PhD fellowship from the French Ministry of Research and Innovation and Université de Paris. RB was funded by a doctoral fellowship from INSERM U1016, Institut Cochin, Paris. TEJ was funded by a doctoral fellowship from Université de Paris. SS and IC were funded by a PhD fellowship from the UK Dementia Research Institute. MM and NKS were funded by a postdoctoral fellowship from the UK Dementia Research Institute.

Abstract

Assessment of differential gene expression by qPCR is heavily influenced by the choice of reference genes. Although numerous statistical approaches have been proposed to determine the best reference genes, they can give rise to conflicting results depending on experimental conditions. Hence, recent studies propose the use of RNA-Seq to identify stable genes followed by the application of different statistical approaches to determine the best set of reference genes for qPCR data normalization. In this study, we demonstrate that the statistical approach to determine the best reference genes from randomly selected candidates is more important than the preselection of 'stable' candidates from RNA-Seq data. Using a qPCR data normalization workflow that we have previously established; we show that qPCR data normalization using randomly chosen conventional reference genes renders the same results as stable reference genes selected from RNA-Seq data. Furthermore, the differential expression of target genes assessed by qPCR using our normalization strategy is comparable to RNA-Seq results. We validated these observations in two distinct cross-sectional experimental conditions involving human iPSC derived microglial cells and mouse sciatic nerves. These results taken together show that given a robust statistical approach for reference gene selection, stable genes selected from RNA-Seq data do not offer any significant advantage over commonly used reference genes for normalizing qPCR assays.

Key words: qPCR, Reference Genes, RNA-Seq, Normalization Factor, Statistical Methods

Introduction

Normalization of relative gene expression using qPCR assays relies crucially on the use of reference genes that exhibit minimal variation across experimental conditions (Bustin *et al*, 2009; Thellin *et al*, 1999; Eisenberg & Levanon, 2013; Jacob *et al*, 2013; Chapman & Waldenström, 2015). Thus, the choice of reference genes has a significant bearing on the normalized profiles of target genes. In the last two decades, many statistical approaches have been proposed to help researchers identify stable reference genes from a given set of candidates (Vandesompele *et al*, 2002; Pfaffl *et al*, 2004; Andersen *et al*, 2004; De Spiegelaere *et al*, 2015; Silver *et al*, 2006; Boda *et al*, 2009). However, the calculations and assumptions employed in these approaches fundamentally differ and could give rise to conflicting results. To address these concerns, we recently reviewed these approaches and following a comparison of existing methods including CV analysis, GeNorm, Normfinder, pairwise dCT method and BestKeeper, we proposed an effective qPCR analysis workflow (Sundaram *et al*, 2019). Our approach employing visual representation of intrinsic variation, CV analysis for identifying overall reference gene variation and NormFinder, proved to be effective in determining stable reference genes from a given set of candidates.

More recently, a growing number of studies have proposed the use of bulk RNA-Seq to screen for stable reference genes followed by the application of existing statistical methods to obtain the "best" reference genes to normalize qPCR assays (Carmona *et al*, 2017; Zhan *et al*, 2014; Hoang *et al*, 2017; Yang *et al*, 2014; Chim *et al*, 2017; Dai *et al*, 2021; Pombo *et al*, 2017; Zhou *et al*; Hernandez-Segura *et al*, 2019). In these studies, the intrinsic validity of screening for reference genes directly from the bulk RNA-Seq data is clear, as they have used RNA-Seq to identify stable reference genes in an effort to standardize the use of certain genes for their experimental setting. Although these efforts are commendable, the cost-benefit trade off in performing RNA-Seq to identify stable reference genes for qPCR assays is negligible. Moreover, this approach is not always feasible, especially in cases of sample scarcity or when dealing with poor yields of RNA. This is particularly relevant when we are interested in only a small number of target genes whose expression can be assessed solely by qPCR. Furthermore, although bulk RNA-Seq quantifies differential expression of target genes, a qPCR confirmation is often recommended, especially when dealing with genes of lower sizes and expression levels (Coenye, 2021; Everaert *et al*, 2017).Preselection of candidate reference genes from RNA-Seq data is further complicated by the fact that the differential expression (fold change) of genes assessed by RNA-Seq can differ significantly from qPCR results in expression tendencies (Coenye, 2021) . Although this can partially be attributed to fundamental differences in the normalization strategy of RNA-Seq and qPCR, it still proves to be an important hurdle in substantiating RNA-Seq data with corresponding qPCR assays. These observations, are systematically overlooked while publishing RNA-Seq data with or without supporting qPCR assays. However, they pose important questions pertaining to the robustness and reproducibility of mRNA expression data across different methods and studies employing similar experimental conditions.

In this study, we address these concerns by demonstrating that the statistical approach employed to filter reference genes is more important than the preselection of stable genes from bulk RNA-Seq data. Specifically, with the right statistical approach,

reference genes filtered from any random set of candidates generate the same differential
expression results when compared to stable genes preselected from bulk RNA-Seq data.
Furthermore, the results obtained are also comparable with the fold changes observed in
RNA-Seq. These results taken together demonstrate that the preselection of candidate
reference genes is not of any significance for data normalization and hence, RNA-Seq data is
not an essential requisite to obtain stable reference genes for qPCR data normalization.
140

Materials and Methods:

141
142
143 **Sample Procurement:**
144
iPSC Microglia:
The CRISPR/Cas gene edited TREM2 knock-out iPSC line BIONi010-C-17 (TREM2 KO) and its
isogenic control BIONi010-C (TREM2 WT) were purchased from EBISC and maintained at
37°C 5% $CO2$ with E8-Flex medium. Microglia were differentiated from these lines as
previously described (Haenseler *et al*, 2017). In short, using AggreWell 800 (34850, StemCell
Technologies), embryonic bodies (EBs) were prepared from iPSCs in E8-Flex medium
supplemented with 50 ng/ml VEGF (PHC9394, ThermoFisher), 50 ng/ml BMP4 (120-05ET,
Peprotech), and 20 ng/ml SCF (300-07, PeproTech) for 3 days with 75% medium change
each day. EBs were transferred to a T75 flask in X-VIVO15 (BE02-060F, Lonza) medium
supplemented with 25 ng/ml IL-3 (PHC0031, ThermoFisher), 2 mM Glutamax (35050061,
ThermoFisher), 100 ng/ml M-CSF (300-25, PeproTech), and 0.055 mM β-mercaptoethanol
(31350-010, ThermoFisher). Precursor cells started to emerge from the EBs approximately 4
weeks later, and were collected by gently tapping the flask. These precursor cells are
differentiated to microglia over 7-days in DMEM F:12, Neuronal basal medium plus (1:1)
supplemented with 100 ng/ml IL-34, 10 ng/ml GM-CSF, and 100 ng/ml M-CSF.
160
Sciatic Nerves:
To assess the stability of commonly used reference genes during myelination of the sciatic
nerves, P3 and P21 C57BL6/J mice (3 Males and 4 females) were dissected and the sciatic
nerves were harvested for RNA extraction. The RNA-seq data for these time points was
mined from a publicly available dataset as explained in the *"Bulk RNA-seq and data*
analysis" section below. All aspects of animal care and animal experimentation were
performed in accordance with the relevant guidelines and regulations of INSERM, Université
de Paris, and approved by the French National Committee of Animal experimentation and
ethics.
170
171 **Total RNA extraction:**
172
173 *iPSC Microglia:*
174 Approximately $2x10^5$ iPSC derived microglial cells were harvested using TRIzol reagent
175 (Ambion Life Technologies 15596018) for total RNA isolation. Total RNA was isolated using
176 Direct-Zol^{TM} RNA microprep columns (Zymo research R2062) according to the
177 manufacturer's protocol.
178
Sciatic Nerves:

Total RNA was extracted from sciatic nerves using 1 mL of TRIzol reagent (Ambion Life
Technologies 15596018) on ice using the manufacturer's instructions with slight
modifications. Briefly, 100% ethanol was substituted for isopropanol to reduce the
precipitation of salts. In addition, RNA precipitation was carried out overnight at -20°C in the
presence of glycogen (0.02mg/mL final concentration). The following day, precipitated RNA
was pelleted by centrifugation and washed at least 3 times with 70% Ethanol to eliminate
any residual contamination. Tubes were then spin-dried in a vacuum centrifuge for 5
minutes and RNA was resuspended in 20 µL of RNase resuspension buffer containing 0.1mM
EDTA, pH 8. RNA was then stored at -80°C till RTqPCR.
189
190
191 **RNA Quality, Integrity, and Assay:**
192
iPSC Microglia:
RNA quantity was assayed using UV spectrophotometry on Nanodrop One (Thermo
Scientific). Optical density absorption ratios A260/A280 & A260/A230 of the iPSC microglia
samples were 2.0 (±0.1 SD) and 2.1 (±0.1 SD) respectively. RNA integrity was verified using
Agilent bioanalyzer. All the samples exhibited a RIN score ≥ 9 and were subsequently used
for downstream analysis including paired-end bulk RNA-seq and qPCR.
199
Sciatic Nerves:
RNA quantity was assayed using UV spectrophotometry on Nanodrop One (Thermo
Scientific). Optical density absorption ratios A260/A280 & A260/A230 of sciatic nerve
samples were 1.84 (±0.04 SD) and 1.54 (±0.53 SD) respectively. RNA integrity was verified
using Agilent bioanalyzer at the Genomics Platform at Institut Cochin, Paris. All samples
exhibited a RIN score ≥ 8.8 and used for qPCR analysis.
206
207 **Bulk RNA-seq and data analysis:**
208
For the iPSC samples, total RNA was extracted and sent to Genewiz, Germany for paired-end
bulk RNA-seq. The raw data, final processed data and the metadata are available in the GEO
database (GSE178924). For the sciatic nerves, bulk RNA-Seq data was obtained from a
previously published study (Ackerman *et al*, 2018). The raw data can be accessed at the
Zenodo repository (https:// zenodo.org/record/1154250). Only the WT datasets at P3 and
P21 were mined and re-analyzed. Fastq files were aligned using the STAR algorithm (version
2.7.6a). Reads were then counted using RSEM (v1.3.1) and the statistical analyses on the
read counts were performed with the DESeq2 package (DESeq2_1.28.1) to determine the
proportion of differentially expressed genes between the two experimental conditions (Love
et al, 2014). The standard DESeq2 normalization method (DESeq2's median of ratios with
the DESeq function) was used, with a pre-filter of reads and genes (reads uniquely mapped
on the genome, or up to 10 different loci with a count adjustment, and genes with at least
10 reads in at least 3 different samples). The biomaRt package (v2.44.4) was used to
substitute gene names for the Ensembl IDs in the count matrix (Durinck *et al*, 2009). The
DESeq2 pipeline was used to fit a generalized linear model (GLM) for the expression of each
gene relative to the experimental groups. To determine the variation in gene expression,
the dispersion value computed by the GLM was tabulated into the results data frame.
Square root of dispersion was calculated using sqrt function from base functions in R

(v4.0.2). The coefficient of dispersion (CVfromDisp) was calculated by multiplying square
root of dispersion by 100.

Reference gene selection from RNA-Seq:
230
The data frame was subjected to 4 filtration methods to select stable genes in both the iPSC
and Sciatic nerve datasets. First filtration was done by retaining all the genes with padj value
above 0.05. The resulting data frame was ordered based on the log2FoldChange (log2FC)
column and the second filtration was performed by retaining all the genes with log2FC
between -0.1 to 0.1 implying negligible intergroup variation. Third filtration was done by
retaining all the genes that lie in the CVfromDisp range between 10 & 20. The fourth and
last filtration was performed by retaining all the genes that exhibited a Basemean value
greater than or equal to 500 implying sufficient expression for qPCR detection.
239
Once these criteria were applied, 683 genes were obtained (basemean values ranging from
500 to 123000) in the iPSC dataset and 42 genes were obtained (basemean values ranging
from 500 to 48000) in the sciatic nerve dataset. To obtain stable reference genes with both
low and high expression, the final list of candidates was partitioned into quartiles based on
basemean expression. 3 reference genes from each of the first three quartiles and 1
reference gene from the last quartile were chosen; thereby generating 10 candidates. This
selection strategy resulted in candidate reference genes that exhibit Cq values between 18
cycles (high expression) and 26 cycles (relatively lower expression) in qPCR.
248
249   **Conventional reference gene selection:**
250
All conventional reference genes were randomly chosen. They do not have any proven
stability for the experimental setting in question. For the iPSC dataset, the conventional
human reference genes chosen were *ACTB, GAPDH, GUSB, HPRT, PGK1, PPIA, RPL13A, TBP,*
and *UBC*. For the sciatic nerve dataset, the conventional mouse reference genes chosen
were *Actb, Gapdh, Tbp, Sdha, Pgk1, Ppia, Rpl13a, Hsp60, Mrpl10, Rps26*. These mouse
reference genes have been previously used to establish the qPCR data analysis workflow
(Sundaram *et al*, 2019)
258
259   **Target gene selection:**
260
Differentially expressed genes that exhibited Padj<0.05 were first retained from the results
dataframe. Next, genes that exhibited log2FC values above +0.6 and below -0.6 were
retained and partitioned into 2 separate lists. 3 target genes were randomly chosen from
each of these lists for differential expression analysis in the iPSC dataset. In the Sciatic nerve
dataset, 3 upregulated genes and 3 downregulated genes were chosen based on the
recently published Sciatic Nerve Atlas https://www.snat.ethz.ch (Gerber *et al*, 2021).
267
268   **Primer Design:**
269
All primers used in the study were designed using the Primer 3 plus software
(https://primer3plus.com/cgi-bin/dev/primer3plus.cgi). Splice variants and the protein-
coding sequence of the genes were identified using the Ensembl database
(www.ensembl.org). Constitutively expressed exons among all splice variants were then
identified using the ExonMine database (Mollet *et al*, 2010). Primers that spanned two
subsequent constitutively expressed exons were then designed using the Primer 3 plus

software. The amplicon size of all primers was between 88bp − 200bp. For detailed
information on Primer sequences refer to Supplementary data.
278
279 **Amplification Efficiencies:**
280
The amplification efficiencies of primers were calculated using serial dilution of cDNA
molecules. Briefly, cDNA from both the experimental groups of the sciatic nerve (for mouse
primers) and iPSC (for human primers) were serially diluted four times by a factor of 10 (1,
1:10, 1:100 & 1:1000). qPCR was then performed using these dilutions and the results were
plotted as a standard curve against the respective concentration of cDNA. Amplification
efficiency (E) was calculated by linear regression of standard curves using the following
equation: $E = 10^{(-1/Slope\ of\ the\ standard\ curve)}$. Primer pairs that exhibited an Amplification Efficiency
(E) of 1.9 to 2.1 (95% - 105%) and an R^2 value (Determination Coefficient) of 0.99 or above
were chosen for this study. The Cq values of all genes across different groups were well
within the range of the standard dilution curve.
291
292
293 **RT-qPCR:**
294
iPSC Microglia:
1000 ng of total RNA was first subjected to DNase digestion (Thermo scientific EN0525) at
37°C for 30 minutes to eliminate contaminating genomic DNA. Next, DNase activity was
stopped using EDTA (Thermo scientific) and cDNA synthesis was done using iSCRIPT™ cDNA
synthesis kit from BioRad (1708891) to a total volume of 20 μl. cDNA was diluted 1:10 with
nuclease free water for qPCR analysis. qPCR was performed using Takyon ROX SYBR 2X
MasterMix (Eurogentec UF-RSMT-B0701) as a fluorescent detection dye. All reactions were
carried out in a final volume of 8 μl in 384 well plates with 300 nM gene-specific primers,
around 5 ng of cDNA (at 100% RT efficiency) and 1X SYBR Master Mix in each well. Each
reaction was performed in triplicates. All qPCR experiments were performed using
ThermoFisher Scientific QuantStudio 7 with a No-Template-Control (NTC) to check for
primer dimers and a No-RT-Control (NRT) to check for any genomic DNA contamination.
307
308
Sciatic Nerves:
500 ng of total RNA was reverse transcribed with Random Primers (Promega C1181) and
MMLV Reverse Transcriptase (Sigma M1302) according to prescribed protocols. qPCR was
performed using Absolute SYBR ROX 2X qPCR mix (Thermo AB1162B) as a fluorescent
detection dye. All reactions were carried out in a final volume of 7 μl in 384 well plates with
300 nM gene-specific primers, around 3.5 ng of cDNA (at 100% RT efficiency) and 1X SYBR
Master Mix in each well. Each reaction was performed in triplicates. All qPCR experiments
were performed on BioRad CFX384 with a No-Template-Control (NTC) to check for primer
dimers and a No-RT-Control (NRT) to check for any genomic DNA contamination.
318
319 **qPCR statistical analysis and data visualization:**
320
qPCR readouts were analyzed in Precision Melt Analysis Software v1.2 (Sciatic nerve
samples) and QuantStudio Real-Time PCR Software v1.7.1 (iPSC samples). The amplicons

were subjected to Melt Curve analysis and were verified for a single dissociation peak at a
Melting Temperature (Tm) > 75°C as expected from the primer constructs. The Cq data was
exported to Microsoft Excel for further calculations. Each biological sample had three
technical replicates thereby generating three individual Cq values. The arithmetic mean of
the triplicates was taken to be the Cq representing the biological sample. The standard
deviation (SD) of the triplicates was also calculated and samples that exhibited SD > 0.20
were considered inconsistent. In such cases, one outlier Cq was removed to have at least
duplicate Cq values for each biological sample and an SD < 0.20.
331
Reference gene validation was performed according to our qPCR data analysis workflow
(Sundaram *et al*, 2019). Visual representation of potential intrinsic variation in reference
genes was identified by plotting the raw expression profiles ($2^{-\Delta Cq}$) of all candidate reference
genes. Due to reduced sample sizes, a non-parametric Mann Whitney U-Test was performed
to assess statistically significant expression variation between experimental groups. The
alpha value was set at 0.05 for statistical significance. The genes were then screened using
Coefficient of variation (CV) analysis to eliminate genes that exhibited CV>50% as they could
impede the robustness of the Normfinder algorithm (Sundaram *et al*, 2019). Normfinder
was then used to determine the best pair of reference genes (least S value) to compute the
Normalization Factor (NF) for qPCR data normalization (Andersen *et al*, 2004). From the
resulting pair of stable reference genes, the NF was then calculated as the arithmetic mean
of the Cq values of the 2 genes for each sample. Relative expression of target genes was
quantified using the $2^{-\Delta\Delta Ct}$ method using the NF as the reference gene (Schmittgen & Livak,
2008; Livak & Schmittgen, 2001). Data was then analysed and visualised using R studio. For
the fold change comparison between qPCRs and RNA-Seq, the non-parametric Kruskal-
Wallis ANOVA was performed. Multiple group comparisons were performed using the non-
parametric Dunn's Post Test. The alpha value was set at 0.05 for statistical significance.
349

Results:

Reference gene selection from RNA-Seq of microglia (WT vs TREM2KO)

A growing body of literature promotes the use of RNA-Seq to identify stable reference genes for qPCR data normalization in different experimental conditions. However, we believe that this approach is superfluous and that stable reference genes can readily be identified by starting out with any random set of candidates and adopting the right statistical approach to determine the best reference genes for a given experimental condition. To test this hypothesis, we first performed paired-end bulk RNA-Seq on iPSC-derived WT and TREM2KO microglia, an experimental model used to study the role of microglia in neuroinflammation and neurodegeneration (McQuade *et al*, 2020; Andreone *et al*, 2020). Based on the criteria detailed in the Materials and Methods section (see *Reference gene selection from RNA-Seq*), we shortlisted 10 candidate reference genes (*ANXA7, APIP, CCNT2, CNBP, DDX42, FOXK1, KIF13A, MRPL37, PPP1R10,* and *USP5*) from our RNA-Seq data. The base mean values of these genes and other RNA-Seq data features are detailed in Supplementary Table 1.

Reference gene validation of RNA-Seq derived reference genes (WT vs TREM2KO)

The reference genes selected from RNA-Seq were then subjected to validation using the qPCR workflow that we developed recently (Sundaram *et al*, 2019). We first computed the intrinsic variation of reference genes by linearizing the Cq values (2^{-Cq}) followed by visually representing the non-normalized expression levels ($2^{-\Delta Cq}$) of the genes in both experimental groups (**Figure 1**). The WT group was used as the experimental calibrator. Subsequently, statistical testing was performed using a non-parametric Mann Whitney Test to determine significant differences between the groups. Among the 10 genes tested, only 2 genes (MRPL37 and PPP1R10) showed significant variation in non-normalized expression levels between the WT and TREM2KO groups (**Figure 1**). This significant difference in the non-normalized profiles should not be misconstrued as differential expression but it is rather an empirical estimate of intergroup variation (or the lack of thereof) in qPCR.

Figure 1. Non-normalized expression profiles (2$^{-\Delta Cq}$) of reference genes derived from RNA-Seq data of iPSC derived microglia. WT group was used as the experimental calibrator. (a) ANXA7, (b) APIP, (c) CCNT2, (d) FOXK1, (e) KIF13A, (f) USP5, (g) CNBP, (h) DDX42, (i) MRPL37, (j) PP1R10 and (k) Normalization Factor (NF) assessed by NormFinder that combines CNBP & KIF133A. Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05.

In accordance with our workflow, we next performed Coefficient of Variation analysis on linearized Cq values (2^{-Cq}) to assess the overall variation (both groups included) of the reference genes (**Table 1**). The genes tested exhibited CV values between 5.99% and

15.75% and they were ranked as shown in **Table 1**. As all genes exhibited CV<50%, they
could therefore be screened using the NormFinder algorithm to determine the best
combination of reference genes to be used for data normalization. This is a crucial step of
reference gene validation as genes that exhibit CV>50% can compromise the robustness of
the algorithm (Sundaram *et al*, 2019)**.** NormFinder determines the stability of a reference
gene by factoring the intergroup as well as the intragroup variation and computes a Stability
S Score (Andersen *et al*, 2004). The lower the S score, the higher the stability of the gene
across all experimental groups. The algorithm also suggests the best pair of reference genes
that can be used for robust data normalization. NormFinder results of the selected
reference genes from RNA-seq are represented in **Table 1** along with their respective S
Scores and ranks. The algorithm further revealed CNBP and KIF13A to be the best
combination of genes with a grouped Stability S score of 0.05 (**Table 1**). The Cq values of
these two genes were then combined to compute the Normalization Factor (NF). Evidently
and as expected, the non-normalized expression level of the NF does not vary significantly
between the two groups (**Figure 1K**).

Table 1. CV and NormFinder analysis

Expression stability of candidate reference genes derived from bulk RNA-Seq data in iPSC-
derived microglia were evaluated using Coefficient of Variation (CV) Analysis &
NormFinder.
409

CV Analysis			NormFinder		
Gene	**%CV**	**Rank**	**Gene**	**Stability S**	**Rank**
CNBP	5.99	1	*CNBP*	0.07	1
KIF13A	6.36	2	*CCNT2*	0.08	2
PPP1R10	6.87	3	*KIF13A*	0.1	3
CCNT2	7.46	4	*PPP1R10*	0.1	4
MRPL37	7.74	5	*USP5*	0.1	5
ANXA7	10.58	6	*MRPL37*	0.11	6
USP5	10.81	7	*DDX42*	0.12	7
DDX42	11.41	8	*ANXA7*	0.15	8

APIP	14.61	9	FOXK1	0.15	9
FOXK1	15.75	10	APIP	0.16	10

Normfinder Best Pair: CNBP/KIF13A
Grouped Stability: 0.05

Reference gene validation of conventional human reference genes (WT vs TREM2KO)

In addition to the reference genes derived from the RNA-Seq data, we next sought to validate commonly used human reference genes chosen arbitrarily. We selected 10 reference genes (*ACTB, GAPDH, GUSB, PPIA, RPLP0, TBP, HPRT, PGK1, UBC, and RPL13A*) and assessed their intergroup variation by plotting their non-normalized profiles (**Figure 2**). This was followed by statistical testing for significant differences between the two experimental groups. Among the 10 genes tested, none of the genes exhibited significant variation in their non-normalized expression levels (**Figure 2**).

Figure 2. Non-normalized expression profiles ($2^{-\Delta Cq}$) of conventional human reference genes in iPSC derived microglia. WT group is used as the experimental calibrator. (a) *ACTB*, (b) *GAPDH*, (c) *GUSB*, (d) *PPIA*, (e) *RPLP0*, (f) *TBP*, (g) *HPRT*, (h) *PGK1*, (i) *UBC*, (j) *RPL13A*, and (k) *Normalization Factor (NF) assessed by NormFinder that combines RPLP0 & GUSB. Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

These genes were then subjected to CV analysis to assess the overall variation (**Table 2**). They exhibited CV values ranging from 5.29% to 32.37%. However, all genes satisfied the CV<50% criterion for NormFinder screening. NormFinder results for these genes are also shown in **Table 2**. Interestingly and in contrast to the reference genes derived from the RNA-Seq data (**Table 1**), the Stability S score of the conventional genes are largely lower than the genes derived from RNA-Seq, thus indicating better stability. The first 6 ranks from RPL13A to GAPDH have S scores less than 0.1 whereas in the previous dataset (**Table 1**), only the first 2 ranks (CNBP & CCNT2) exhibit scores less than 0.1. Finally, among all the

conventional reference genes screened, NormFinder suggested the use of RPLP0 and GUSB
as the best pair for data normalization with a grouped stability of 0.03 (**Table 2**); which in
comparison, is lesser than the Stability Score of 0.05 from CNBP and KIF13A (**Table 1**). In
principle, these results show that randomly selected reference genes could potentially
exhibit better stability than stable reference genes filtered from RNA-Seq data.

Table 2: CV and NormFinder analysis of conventional reference genes. Expression stability
of randomly chosen candidate reference genes were evaluated using Coefficient of
Variation (CV) Analysis, NormFinder. Best pair chosen is *RPLP0* and *GUSB* with a Stability S
score of 0.03.

CV Analysis			NormFinder		
Gene	%CV	Rank	Gene	Stability S	Rank
GUSB	5.29	1	*RPL13A*	0.03	1
TBP	7.56	2	*GUSB*	0.04	2
RPL13A	8.16	3	*RPLP0*	0.05	3
RPLP0	8.18	4	*TBP*	0.05	4
PGK1	9.56	5	*PGK1*	0.06	5
GAPDH	11.35	6	*GAPDH*	0.07	6
PPIA	13.75	7	*PPIA*	0.10	7
ACTB	14.31	8	*UBC*	0.11	8
UBC	17.47	9	*ACTB*	0.14	9
HPRT	32.27	10	*HPRT*	0.18	10

449 ***NormFinder Best Pair: RPLP0/GUSB***
450 ***Grouped Stability S: 0.03***

Differential expression of target genes by qPCR and RNA-Seq (WT vs TREM2KO)

Following reference gene validation of conventional and RNA-Seq-derived candidates, we picked 6 target genes that were differentially expressed based on the RNA-Seq results (*see Target gene selection* from *Materials & Methods*). We included 3 target genes (*TREM2*, *P2RY12* and *CD52*) that were significantly downregulated and 3 others (*ACSS3, SYDE1* and *TBX10*) that were significantly upregulated. The relative expression of these genes was then computed by qPCR using the Normalization Factor (NF) from conventional candidate reference genes (*RPLP0* & *GUSB* or Conventional NF) or from RNA-Seq-derived candidate reference genes (*CNBP* & *KIF13A* or RNA-Seq derived NF) (**Figure 3**). To compare our qPCR results, we also computed the fold change assessed by RNA-Seq using the Normalized counts of these genes.

Regarding the downregulated genes, *TREM2* expression in the TREM2KO group, when assessed through qPCRs, exhibited expression fold change values of 0.31 ± 0.05 (mean FC ± SD, conventional NF, **Figure 3A**) and 0.35 ± 0.04 (RNA-Seq derived NF, **Figure 3B**). In RNA-Seq, the fold change values were 0.23 ± 0.03 (**Figure 3C**). *P2RY12* expression fold change in the TREM2KO group were 0.04 ± 0.02 folds lower when assessed using qPCRs (both conventional NF and RNA-Seq derived NF, **Figure 3D and 3E**). However, RNA-Seq fold change values of *P2RY12* were comparatively higher at 0.62 ± 0.19 (**Figure 3F**). *CD52* fold change in the TREM2KO group when assessed by qPCR were 0.38 ± 0.06 (Conventional NF, **Figure 3G**) and 0.40 ± 0.05 (RNA-Seq derived NF, **Figure 3H**). RNA-Seq fold changes were comparatively lower at 0.16 ± 0.01 (**Figure 3I**).

For upregulated genes, *ACSS3* expression fold change in the TREM2KO group exhibited fold change values of 3.72 ± 0.79 (Conventional NF, **Figure 3J**) and 3.40 ± 0.88 (RNA-Seq derived NF, **Figure 3K**). The fold changes in RNA-Seq were 4.54 ± 1.25 (**Figure 3L**). *SYDE1* expression levels assessed by qPCR were at 3.58 ± 1.18 (Conventional NF, **Figure 3M**) and 3.29 ± 1.28 (RNA-Seq derived NF, **Figure 3N**). RNA-Seq expression levels of *SYDE1* were at 3.62 ± 0.34 (**Figure 3O**). Finally, *TBX10* expression fold changes assessed by qPCR were at 3.06 ± 1.51 (Conventional NF, **Figure 3P**) and 2.82 ± 1.60 (RNA-Seq derived NF, **Figure 3Q**). RNA-Seq fold changes were at 3.24 ± 0.61 (**Figure 3R**). These results taken together with the visual representation of differential expression described in **Figure 3** show that qPCR normalization using conventional NF or RNA-Seq derived NF render the same results. They do not always concur with RNA-Seq fold changes in magnitude but do so in tendency.

Figure 3: Differential expression of target genes between WT and TREM2KO iPSC cells assessed by qPCRs and RNA-Seq. *The target genes (TREM2, P2RY12, CD52, ACSS3, SYDE1, TBX10) have been normalized with the normalization factor (NF) computed from conventional reference genes RPLP0 & GUSB in the first column (A, D, G, J, M, P) or from RNA-Seq derived reference genes CNBP & KIF13A in the second column (B, E, H, K, N, Q). The third column contains the Fold Changes computed from RNA-Seq (C, F, I, L, O, R). The normalized counts for each gene in each group were used to calculate the fold changes in RNA-Seq. The WT group was used as the experimental control. The P_{adj} value of RNA-Seq is indicated in the TREM2KO group for each gene in the RNA-Seq column. For the qPCRs, Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05.*

Comparison of fold changes between qPCRs and RNA-Seq (WT vs TREM2KO):

We next investigated in detail if the Fold Change distributions in the TREM2KO experimental group computed across the 3 methods differed significantly from one another. To this end, we compared the distributions by a non-parametric ANOVA (Kruskal Wallis Test) of ordinal distributions followed by Dunn's multiple comparison post-test (**Supplementary Figure 1**). qPCR Fold changes computed using Conventional NF were used as the control for multiple comparisons. Therefore, the first comparison was made between the conventional and RNA-Seq derived NF groups (difference in qPCRs, hereafter referred to as Comparison 1) and the second comparison was made between the Conventional NF group and RNA-Seq differential expression (difference between qPCR and RNA-Seq, hereafter referred to as Comparison 2).

The WT experimental group was omitted from this analysis as it was used as the experimental calibrator for differential expression. The mean Fold Change of the WT group is always at 1 regardless of the gene/method in question and therefore it is redundant to test for statistical significance of the WT fold change levels across different methods for each gene.

Non-parametric ANOVA revealed no significant difference between the FC distributions of the upregulated genes *ACSS3, SYDE1* and *TBX1* (**Supplementary Figure 1D, 1E and 1F**) suggesting that the qPCR fold changes concur with RNA-Seq fold changes in these genes. As for the downregulated genes, *TREM2* FC distributions varied significantly between the groups (Kruskal Wallis $P < 0.5$); however, Dunn's post test revealed no significant difference in Comparison 1 and Comparison 2 (**Supplementary Figure 1A**). *P2RY12* FC distributions varied significantly between the groups (Kruskal Wallis $P<0.01$). Multiple comparisons revealed no difference in Comparison 1 but a significant difference in Comparison 2 (Dunn's Post Test $P<0.05$) (**Supplementary Figure 1B**) CD*52* FC distributions varied significantly between the groups (Kruskal Wallis $P < 0.5$). Multiple comparisons, yet again, revealed no difference in Comparison 1 but a significant difference in Comparison 2 (Dunn's Post Test $P<0.05$) (**Supplementary Figure 1C**). These results show firstly that no significant difference exists between the FC distributions of all the genes between the qPCRs (Comparison 1). Secondly, FC distributions of qPCRs and RNA-Seq do not show any difference for the upregulated genes *ACSS3, SYDE1* and *TBX10* (Comparison 2). However, for the downregulated genes, although no differences exist between qPCRs (Comparison 1); *P2RY12* & *CD52* FC distributions differ significantly between qPCR and RNA-Seq. In summation, these results prove beyond reasonable doubt that differential expression assessed by qPCR using randomly chosen conventional reference genes and "stable" reference genes filtered from RNA-Seq render similar expression profiles of target genes without any significant difference among the two normalization strategies. However, a perfect concordance with RNA-Seq results was only observed in the upregulated genes and could not be achieved in the downregulated genes.

Reference gene selection from RNA-Seq of mouse sciatic nerves (P3 vs P21)

To rule out any circumstantial evidence that may support our observations in the iPSC dataset, we sought to validate our results using a completely different experimental setting. We wanted to rule out the possibility that the conventional reference genes that we chose

in the iPSC dataset were indeed stable reference genes by mere happenstance and thus
they rendered the same profiles of differential expression as the reference genes filtered
from RNA-Seq.

Therefore, in a confirmatory experimental approach we mined and re-analyzed RNA-Seq
data that was previously generated from mouse sciatic nerves at Postnatal day 3 (P3) and
P21 (Ackerman *et al*, 2018). This dataset permits us to study RNA expression changes in the
sciatic nerves during postnatal myelination. We applied the same criteria to select 10
candidate reference genes (*Ppp3ca, Fkbp4, Vcp, Lama2, Ank2, Coq9, Chmp2a, Laptm5,*
Leprotl1, Supt4a). The basemean values of these genes and other RNA-Seq data features are
detailed in Supplementary Table 2.

Reference gene validation of RNA-Seq derived reference genes from mouse sciatic
nerves (P3 vs P21)

We evaluated the intrinsic intergroup variation of these reference genes by visual
representation of their non-normalized profiles ($2^{-\Delta Cq}$) using P3 samples as the calibrator
(Figure 4). As previously described, statistical inference was then assessed using the non-
parametric Mann Whitney U test owing to reduced sample sizes. Interestingly, and in
contrast to the iPSC dataset, 6 out of 10 genes (*Fkbp4, Lama2, Coq9, Chmp2a, Leprotl1 &*
Supt4a) exhibit significant intergroup variation across the two experimental groups (**Figure**
4A, 4B, 4C, 4D, 4E & 4F). However, it should be noted that this is an empirical estimate of
intrinsic variation as we have represented the non-normalized profiles. This does not
necessarily imply that these genes are differentially expressed between the two groups.

Figure 4. Raw expression profiles of reference genes obtained from RNA-Seq analysis of developing sciatic nerve of age Post-natal day 3 (P5) and 21 (P21), expressed as fold changes (2-ΔCq). P5 group is the experimental calibrator. (A) *Fkbp4*, (B) *Lama2*, (C) *Leprotl4*, (D) *Supt4a*, (E) *Coq9*, (F) *Chmp2a*, (G) *Ank2*, (H) *Vcp*, (I) *Laptm5*, (J) *Ppp3ca*, and (K) Normalisation Factor (*Ppp3ca + Coq9*). *Non-parametric Mann Whitney U test was used to assess differences between the groups. The alpha value was set at 0.05 and P values are annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

We next performed the CV analysis to assess overall variation (**Table 3**). The genes tested exhibited CV values between 12.32% and 36.24% and they were ranked as shown in **Table 3**. As all genes exhibited CV<50%, they could therefore be screened using the NormFinder algorithm. NormFinder analysis further revealed *Ppp3ca* and *Coq9* to be the best combination of genes with a grouped Stability S score of 0.04 (**Table 3**). Although Coq9 exhibited significant intergroup variation (**Figure 4E**), the combination of the gene with *Ppp3ca* resulting in the Normalization Factor (NF) does not exhibit any (**Figure 4K**). This

577 confirms the strength of the NormFinder algorithm and the validity of using the two best
578 reference genes proposed by the method.

579 **Table 3. CV and NormFinder analysis of reference genes derived from RNA-Seq**

CV Analysis			NormFinder		
Gene	%CV	Rank	Gene	Stability S	Rank
Laptm5	12.32	1	*Ank2*	0.07	1
Ppp3ca	12.96	2	*Ppp3ca*	0.08	2
Supt4a	14.79	3	*Coq9*	0.08	3
Coq9	16.26	4	*Laptm5*	0.16	4
Ank2	16.31	5	*Supt4a*	0.16	5
Vcp	19.59	6	*Leprotl1*	0.23	6
Leprotl1	24.49	7	*Fkbp4*	0.26	7
Fkbp4	25.86	8	*Vcp*	0.27	8
Lama2	29.96	9	*Chmp2a*	0.47	9
Chmp2a	36.24	10	*Lama2*	0.56	10

581
582 ***Normfinder Best combination: Ppp3ca + Coq9***
583 ***Grouped Stability: 0.04***
584
585

Reference gene validation of conventional mouse reference genes (P3 vs P21)

To obtain stable reference genes from a set of conventional genes, we used 10 candidate
mouse reference genes that we had previously used to establish our workflow (*Actb, Gapdh,*
Tbp, Sdha, Pgk1, Ppia, Rpl13a, Hsp60, Mrpl10, Rps26) (Sundaram *et al*, 2019). As explained
above, visual representation followed by inferential statistics were performed and the
results are described in **Figure 5**. 4 out of 10 genes (*Gapdh, Sdha, Pgk1 & Rps26*) exhibited
significant intrinsic variation between the experimental groups (**Figure 5G ,5H, 5I & 5J**)

Figure 5. Raw expression profiles of conventional reference genes of developing sciatic nerve of age Post-natal day 3 (P5) and 21 (P21), expressed as fold changes (2-ΔCq). P5 group is the experimental calibrator. (A) *Actb*, (B) *Mrpl10*, (C) *Hsp60*, (D) *Ppia*, (E) *Rpl13a*, (F) *Tbp*, (G) *Gapdh*, (H) *Pgk1*, (I) *Rps26*, (J) Sdha, and (K) Normalisation Factor (*Hsp60 + Ppia*). Non-parametric Mann Whitney U Test was performed to assess differences between the means of all groups. Statistical significance is denoted by '*' representing P values: **P<0.01.

CV analysis was then performed on these genes and the results are depicted in **Table 4**. CV values ranged from 12.3% to 30.44% and the genes were ranked accordingly. As all genes exhibited CV<50%, we screened these genes through the NormFinder algorithm to determine the best combination of reference genes for qPCR normalization. NormFinder determined *Hsp60* and *Ppia* to be the best combination of reference genes with a grouped Stability S Score of 0.06 (**Table 6**). Both Hsp60 (**Figure 5C**) & Ppia (**Figure 5D**) did not exhibit

607 any intergroup variation and the same was true for the NF computed by combining these
608 genes (**Figure 5K**).

609 **Table 4. CV and NormFinder analysis of conventional reference genes**

CV Analysis			NormFinder		
Gene	%CV	Rank	Gene	Stability S	Rank
Mrpl10	12.30	1	*Tbp*	0.03	1
Gapdh	13.06	2	*Mrpl10*	0.04	2
Hsp60	14.36	3	*Hsp60*	0.05	3
Tbp	15.24	4	*Ppia*	0.05	4
Pgk1	15.83	5	*Pgk1*	0.06	5
Ppia	19.36	6	*Actb*	0.07	6
Actb	21.07	7	*Rpl13a*	0.10	7
Rps26	22.04	8	*Gapdh*	0.11	8
Rpl13a	26.78	9	*Rps26*	0.14	9
Sdha	30.44	10	*Sdha*	0.18	10

611
612 *Normfinder Best combination: Hsp60 + Ppia*
613 *Grouped stability: 0.06*
614

Differential expression of target genes by qPCR and RNA-Seq (P3 vs P21)

Following reference gene validation of conventional and RNAseq-derived candidates, we
picked 6 target genes that were known to be differentially expressed during postnatal
development of the sciatic nerve (Gerber *et al*, 2021). 3 target genes (*Mpz, Mbp and Cd90*)
that were chosen are significantly upregulated and the other 3 target genes (*p75NTR, Mki67*
and *Sox2*) are significantly downregulated at these time points. The relative expression of
these genes was then computed by qPCR using the Normalization Factor (NF) from

conventional candidate reference genes (*Hsp60* & *Ppia* or Conventional NF) or from
RNAseq-derived candidate reference genes (*Ppp3ca* & *Coq9* or RNAseq derived NF) (**Figure**
6). To compare our qPCR results, we also computed the fold change assessed by RNA-Seq
using the Normalized counts of these genes.

Regarding the downregulated genes, *Mki67* expression in the P21 group, when assessed
through qPCRs, exhibited fold change values of 0.14 ± 0.10 (mean FC ± SD, conventional NF,
Figure 6A) and 0.12 ± 0.08 (RNA-Seq derived NF, **Figure 6B**). In RNA-Seq, the fold change
values were 0.08 ± 0.01 (**Figure 6C**). *p75NTR* expression fold change in the P21 group were
0.30 ± 0.04 (conventional NF, **Figure 6D)** and 0.26 ± 0.02 (RNA-Seq derived NF, **Figure 6E**)
when assessed using qPCRs. However, RNA-Seq fold change values of *p75NTR* were
comparatively higher at 0.63 ± 0.18 (**Figure 6F**). *Sox2* expression fold change in the P21
group when assessed by qPCR were 0.49 ± 0.05 (Conventional NF, **Figure 6G**) and 0.42 ±
0.03 (RNA-Seq derived NF, **Figure 6H**). RNA-Seq changes were at 0.39 ± 0.09 (**Figure 6I**).

For the upregulated genes, *Mbp* expression levels in the P21 group exhibited fold change
values of 1.73 ± 0.15 (Conventional NF, **Figure 6J**) and 1.49 ± 0.19s (RNA-Seq derived NF,
Figure 6K). The fold changes in RNA-Seq were 2.02 ± 0.62 (**Figure 6L**). *Mpz* expression levels
assessed by qPCR were at 1.76 ± 0.38 (Conventional NF, **Figure 6M**) and 1.53 ± 0.35 (RNA-
Seq derived NF, **Figure 6N**). However, RNA-Seq expression fold change of *Mpz* were
significantly higher at 3.49 ± 0.26 (**Figure 6O**). Finally, *Cd90* expression fold change assessed
by qPCR were at 1.83 ± 0.35 (Conventional NF, **Figure 6P**) and 1.57 ± 0.30 (RNA-Seq derived
NF, **Figure 6Q**). RNA-Seq fold changes were at 4.81 ± 2.77 (**Figure 6R**). Yet again, these
results taken together with the visual representation of differential expression described in
Figure 6 show that qPCR normalization using conventional NF or RNA-Seq derived NF render
very similar results. We observe again that they do not always concur with RNA-Seq fold
changes in magnitude, but they do so in tendency.

Figure 6: Differential expression of target genes in P3 and P21 mouse sciatic nerves by qPCRs and RNA-Seq. *The target genes (Mki67, p75NTR, Sox2, Mbp, Mpz, Cd90) have been normalized with the normalization factor (NF) computed from conventional reference genes Hsp60 & Ppia in the first column (A, D, G, J, M, P) or from RNA-Seq derived reference genes Ppp3ca and Coq9 in the second column (B, E, H, K, N, Q). The third column contains the Fold Changes computed from RNA-Seq (C,F,I,L,O,R). The normalized counts for each gene in each group were used to calculate the fold changes in RNA-Seq. The WT group was used as the experimental control. The P_{adj} value of RNA-Seq is indicated in the P21 group for each gene in the RNA-Seq column. For the qPCRs, Non-parametric Mann Whitney U test was used to*

assess differences between the groups. The alpha value was set at 0.05 and P values are
annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.

Comparison of differential expression between qPCRs and RNA-Seq (P3 vs P21):

Similar to the analysis performed in the iPSC dataset, we next investigated in detail if the
Fold Change distributions in the P21 experimental group computed across the 3 methods
differed significantly from one another. We used non-parametric ANOVA (Kruskal Wallis
Test) of ordinal distributions followed by Dunn's multiple comparison post-test
(**Supplementary Figure 2**). As shown in the iPSC dataset, we performed multiple
comparisons between qPCRs (Comparison 1) and also between qPCR and RNA-Seq
(Comparison 2).

Non-parametric ANOVA revealed no significant difference between the FC distributions of
Mki67 and *Mbp* (**Supplementary Figure 2A and 2E**) suggesting that the qPCR fold changes
concur with RNA-Seq fold changes in these genes. *Sox2, p75NTR* and *Mpz* FC distributions
varied significantly between the groups (Kruskal Wallis $P < 0.05$, $P < 0.01$ and $P < 0.01$
respectively); however, Dunn's post test revealed no significant difference in both
Comparison 1 and Comparison 2 (**Supplementary Figure 2B, 2C & 2F**). *Cd90* FC distributions
varied significantly between the groups (Kruskal Wallis $P < 0.01$). Multiple comparisons
revealed no difference in Comparison 1 and Comparison 2 (**Supplementary Figure 2D**). In
summation, all the genes tested in the sciatic nerve dataset show no significant differences
between Conventional NF and RNA-Seq-derived NF results. In summation, these results
show once again that stable reference genes filtered from RNA-Seq data do not give any
added advantage to qPCR data normalization and that not all genes exhibit comparable Fold
Changes when comparing qPCR results and RNA-Seq.

Discussion:

The central premise of this article was to demonstrate that stable reference genes
for qPCR data normalisation can be obtained from any random set of candidates provided
the statistical approach of reference gene validation is sound and consistent. To this end,
using our qPCR workflow we have shown using two separate and unique datasets that
randomly chosen reference genes can render the same results of differential expression
when compared to stable references selected from RNA-Seq data. It therefore follows that
our statistical approach can provide reference genes that are as powerful as stable genes
selected from RNA-Seq data.

The advantage of this approach in filtering the best candidates from any random set
lies in combining 3 methods that are complementary to each other. Although multiple
statistical methods have been proposed before, we have previously highlighted the
potential pitfalls and assumptions of these computational methods (Sundaram *et al*, 2019).
We thus devised this workflow combining the well-known NormFinder method with CV
analysis and the visual representation of non-normalized expression followed by statistical
testing. Indeed, it should be noted the core method of this approach is the NormFinder
algorithm. However, this method was first described and validated with larger sample sizes
(typically between 10 - 20 or more) and with a high number of candidate genes chosen from
microarrays (typically between 15 - 20) exhibiting very little overall variation (Andersen *et*
al, 2004). In our previous study, we identified that lower sample sizes and candidate genes
with high overall variation can skew the results of this method because of the algorithm's
construct (Sundaram *et al*, 2019). Although scores of studies have used this algorithm
without due diligence to the prerequisites, our combined approach resolves these issues.
The CV analysis helps in the identification of genes with high overall variance and their
successive elimination from analysis gives rise to more robust results from NormFinder.
Visual representation of the non-normalized profiles ($2^{-\Delta Cq}$) followed by statistical testing
further validates the results of NormFinder as the Normalization Factor computed from the
two best reference genes does not exhibit any significant differences between the
experimental groups regardless of experimental conditions (**Figure 1K, 2K, 4K & 5K**). Thus, in
combining CV analysis and statistical testing of non-normalized profiles, we extrapolated the
NormFinder approach to be used for experimental setups involving lower sample sizes
similar to those used in this study.

The cut off at 50% CV to qualify for NormFinder analysis is rooted in the fact that at
50%, the standard deviation of the expression distribution is half the mean expression level.
Previous datasets that we published have indicated that the Normalization Factor (NF)
computed by NormFinder when including genes that exhibit CV>50% exhibit significant
intergroup variation in the non-normalized profiles (Sundaram *et al*, 2019). Below this cut
off, NF expression levels are stable across experimental groups. Indeed, the user can be
more stringent by using a lower CV cut off but our data from this study and the previous one
suggests that this is not necessary. However, this criterion gives rise to a theoretical
limitation of our approach. In a given experimental setup, if all the candidate reference
genes tested exhibit CV values of above 50%, then our approach will fail and NormFinder
results would no longer be reliable. In such cases, a new list of candidates is required.

However, our experience suggests that as little as 6 genes after eliminating genes exhibiting
CV>50% can still provide a sound NF for qPCR data normalization (Sundaram *et al*, 2019).
728
Although not the primary objective of the study, the comparison of qPCR results
with RNA-Seq merits discussion. In the iPSC dataset, comparisons made between qPCR and
RNA-Seq results revealed that 4 out of 6 genes (*ACSS3, SYDE1, TREM2* and *TBX10*) showed
no significant difference between the fold changes computed by both the methods
(**Supplementary Figures 1D, 1E, 1F & 1C**). However, changes in the expression of *CD52* and
P2RY12 significantly differed between the two methods (**Figure 3 and Supplementary
Figure 1A and 1B**). In the sciatic nerve dataset, 5 out of 6 genes tested show no difference
between qPCR and RNA-Seq data in multiple comparisons (**Supplementary Figure 2A, 2B,
2C, 2E, 2F**). However, this analysis is not optimal as the number of samples used in qPCR is
much higher (N=7) than the number of samples in RNA-Seq data that we mined (N=3). In
reality, only 3 genes (*Mki67, Sox2 & Mbp*) exhibited overlapping fold change distributions
across the methods (**Supplementary Figure 2A, 2B & 2E**). However, regardless of the
magnitudes of the change observed, the pattern of expression changes was always
conserved across qPCR and RNA-Seq.
743
There are several reasons for these discrepancies in magnitudes. In our study,
samples that were subjected to RNA-Seq and qPCR were different in both datasets. In the
iPSC dataset, although the number of biological replicates were the same, qPCR analysis was
performed at a later cell passage number than the RNA-Seq analysis. Similarly, in the sciatic
nerve dataset, the RNA-Seq data was mined from a previous study that employed only 3
biological replicates at P21. These differences in sampling could be responsible for the
differences in expression assessed by these methods. A more thorough approach to
compare qPCR and RNA-Seq data is to subject the same samples to both methodologies and
then comparing their results bearing in mind the requirement of larger sample sizes for
statistical testing of differential expression in qPCR assays. Previous studies have indeed
tried to validate RNAseq data with RTqPCR for a large set of target genes typically ranging
from a few hundred to the entire transcriptome (Griffith *et al*, 2010; Rajkumar *et al*, 2015;
Everaert *et al*, 2017). The consensus from these studies is that the extent of correlation
between RTqPCR and RNAseq data is dependent on how the sequencing data was aligned,
mapped and counted based on existing protocols. Additionally, though not discussed in
these articles, we believe that this would equally depend on how reference genes were
selected for qPCR assays that validate the RNAseq data. Overlooking these technicalities,
these studies also show that around 85% of all genes tested show concordant differential
expression (comparable magnitudes with the same tendency) between the two
methodologies with correlation coefficients typically above 0.8. Although these numbers are
relatively reassuring at the outset, from an absolute perspective, the implications raise
important concerns. For a hypothetical bulk RNA-Seq dataset of around 18000 protein-
coding transcripts, 85% concordance leaves 2700 genes whose differential expression is
discordant between RNA-Seq and qPCR. Of note, the study conducted by Everaert and
colleagues (Everaert *et al*, 2017) conclude that discordant genes typically are smaller, have
fewer exons and are expressed at lower levels. In our study, the discordant genes *CD52* and
P2RY12 in the microglia dataset have only 2 exons. The other 4 concordant genes have
between 4 – 13 exons and therefore are relatively larger. The basemean values (and
therefore the expression levels) of the 2 discordant genes are largely higher than the 4

concordant genes. Therefore, number of exons appears to be more important than
expression levels for concordance between qPCR and RNA-Seq. On the contrary, we did not
observe any correlation between discordance and small gene size/expression in the sciatic
nerve samples. These observations show that the parameters are subjective and there is no
objective definition or correlation between transcript sizes and expression levels in different
tissues. Therefore, they prove beyond reasonable doubt that it is prudent to substantiate
RNA-Seq results by qPCR assays at least for the subset of genes or specific pathways that are
important for the study at hand. The methodology used in our study aptly addresses this
necessity by demonstrating that robust and reproducible qPCR assays can be performed
using any random set of reference genes lest the sequencing data not be substantiated due
to poor and sub-optimal qPCR normalization.
784
785   In conclusion, our data shows that the qPCR workflow that we employ can render
786 the most stable reference genes from any list of randomly chosen candidates regardless of
787 their provenance. Differential expression assessed using this workflow can thus be used
788 either as rigorous standalone results or as reliable confirmatory results to substantiate RNA-
789 seq data.

References:

Ackerman SD, Luo R, Poitelon Y, Mogha A, Harty BL, D'Rozario M, Sanchez NE, Lakkaraju AKK, Gamble P, Li J, *et al* (2018) GPR56/ADGRG1 regulates development and maintenance of peripheral myelin. *J Exp Med* 215: 941–961

Andersen C, Jensen J & Orntoft T (2004) Normalization of Real - Time Quantitative Reverse Transcription - PCR Data : A Model - Based Variance Estimation Approach to Identify Genes Suited for Normalization , Applied to Bladder and Colon Cancer Data Sets. *Cancer Res* 64: 5245

Andreone BJ, Przybyla L, Llapashtica C, Rana A, Davis SS, van Lengerich B, Lin K, Shi J, Mei Y, Astarita G, *et al* (2020) Alzheimer's-associated PLCγ2 is a signaling node required for both TREM2 function and the inflammatory response in human microglia. *Nat Neurosci* 23: 927–938

Boda E, Pini A, Hoxha E, Parolisi R & Tempia F (2009) Selection of Reference Genes for Quantitative Real-time RT-PCR Studies in Mouse Brain. *J Mol Neurosci* 37: 238–253

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, *et al* (2009) The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments.

Carmona R, Arroyo M, Jiménez-Quesada MJ, Seoane P, Zafra A, Larrosa R, Alché J de D & Claros MG (2017) Automated identification of reference genes based on RNA-seq data. *Biomed Eng Online* 16

Chapman JR & Waldenström J (2015) With Reference to Reference Genes: A Systematic Review of Endogenous Controls in Gene Expression Studies. *PLoS One* 10: e0141853

Chim SSC, Wong KKW, Chung CYL, Lam SKW, Kwok JSL, Lai C-Y, Cheng YKY, Hui ASY, Meng M, Chan O-K, *et al* (2017) Systematic Selection of Reference Genes for the Normalization of Circulating RNA Transcripts in Pregnant Women Based on RNA-Seq Data. *Int J Mol Sci* 18: 1709

Coenye T (2021) Do results obtained with RNA-sequencing require independent verification? *Biofilm* 3: 100043

Dai Y, Cao Y, Köhler J, Lu A, Xu S & Wang H (2021) Unbiased RNA-Seq-driven identification and validation of reference genes for quantitative RT-PCR analyses of pooled cancer exosomes. *BMC Genomics* 22: 27

Durinck S, Spellman PT, Birney E & Huber W (2009) Mapping Identifiers for the Integration of Genomic Datasets with the R/Bioconductor package biomaRt. *Nat Protoc* 4: 1184

Eisenberg E & Levanon EY (2013) Human housekeeping genes, revisited. *Trends Genet* 29: 569–74

Everaert C, Luypaert M, Maag JL V, Cheng QX, Dinger ME, Hellemans J & Mestdagh P (2017) Benchmarking of RNA-sequencing analysis workflows using whole-transcriptome RT-qPCR expression data. *Sci Rep* 7: 1559

Gerber D, Pereira JA, Gerber J, Tan G, Dimitrieva S, Yánguez E & Suter U (2021) Transcriptional profiling of mouse peripheral nerves to the single-cell level to build a sciatic nerve atlas (Snat). *Elife* 10

Griffith M, Griffith OL, Mwenifumbo J, Goya R, Morrissy AS, Morin RD, Corbett R, Tang MJ, Hou YC, Pugh TJ, *et al* (2010) Alternative expression analysis by RNA sequencing. *Nat Methods* 7: 843–847

Haenseler W, Sansom SN, Buchrieser J, Newey SE, Moore CS, Nicholls FJ, Chintawar S, Schnell C, Antel JP, Allen ND, *et al* (2017) A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports* 8: 1727–1742

Hernandez-Segura A, Rubingh R & Demaria M (2019) Identification of stable senescence-

associated reference genes. *Aging Cell* 18: e12911

Hoang VLT, Tom LN, Quek X-C, Tan J-M, Payne EJ, Lin LL, Sinnya S, Raphael AP,
Lambie D, Frazer IH, *et al* (2017) RNA-seq reveals more consistent reference genes for
gene expression studies in human non-melanoma skin cancers. *PeerJ* 5: e3631

Jacob F, Guertler R, Naim S, Nixdorf S, Fedier A, Hacker NF & Heinzelmann-Schwarz V
(2013) Careful Selection of Reference Genes Is Required for Reliable Performance of
RT-qPCR in Human Normal and Cancer Cell Lines. *PLoS One* 8: e59180

Livak KJ & Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2^-ΔΔCT Method. *Methods* 25: 402–408

Love MI, Huber W & Anders S (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol 2014 1512* 15: 1–21

McQuade A, Kang YJ, Hasselmann J, Jairaman A, Sotelo A, Coburn M, Shabestari SK,
Chadarevian JP, Fote G, Tu CH, *et al* (2020) Gene expression and functional deficits
underlie TREM2-knockout microglia responses in human models of Alzheimer's
disease. *Nat Commun* 11: 1–17

Mollet IG, Ben-Dov C, Felício-Silva D, Grosso AR, Eleutério P, Alves R, Staller R, Silva TS
& Carmo-Fonseca M (2010) Unconstrained mining of transcript data reveals increased
alternative splicing complexity in the human transcriptome. *Nucleic Acids Res* 38:
4740–4754

Pfaffl MW, Tichopad A, Prgomet C & Neuvians TP (2004) Determination of stable
housekeeping genes, differentially regulated target genes and sample integrity:
BestKeeper-Excel-based tool using pair-wise correlations. *Biotechnol Lett* 26: 509–515

Pombo MA, Zheng Y, Fei Z, Martin GB & Rosli HG (2017) Use of RNA-seq data to identify
and validate RT-qPCR reference genes for studying the tomato- Pseudomonas
pathosystem. *Sci Rep* 7: 44905

Rajkumar AP, Qvist P, Lazarus R, Lescai F, Ju J, Nyegaard M, Mors O, Børglum AD, Li Q
& Christensen JH (2015) Experimental validation of methods for differential gene
expression analysis and sample pooling in RNA-seq. *BMC Genomics* 16: 1–8

Schmittgen TD & Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T)
method. *Nat Protoc* 3: 1101–8

Silver N, Best S, Jiang J & Thein SL (2006) BMC Molecular Biology Selection of
housekeeping genes for gene expression studies in human reticulocytes using real-time
PCR.

De Spiegelaere W, Dern-Wieloch J, Weigel R, Schumacher V, Schorle H, Nettersheim D,
Bergmann M, Brehm R, Kliesch S, Vandekerckhove L, *et al* (2015) Reference gene
validation for RT-qPCR, a note on different available software packages. *PLoS One* 10:
1–13

Sundaram VK, Sampathkumar NK, Massaad C & Grenier J (2019) Optimal use of statistical
methods to validate reference gene stability in longitudinal studies. *PLoS One* 14:
e0219440

Thellin O, Zorzi W, Lakaye B, De Borman B, Coumans B, Hennen G, Grisar T, Igout A &
Heinen E (1999) Housekeeping genes as internal standards: use and limits

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A & Speleman F
(2002) Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034

Yang H, Liu J, Huang S, Guo T, Deng L & Hua W (2014) Selection and evaluation of novel
reference genes for quantitative reverse transcription PCR (qRT-PCR) based on genome
and transcriptome data in Brassica napus L. *Gene* 538: 113–122

Zhan C, Zhang Y, Ma J, Wang L, Jiang W, Shi Y & Wang Q (2014) Identification of
reference genes for qRT-PCR in human lung squamous-cell carcinoma by RNA-Seq.

Acta Biochim Biophys Sin (Shanghai) 46: 330–337
Zhou Z, Cong P, Tian Y & Zhu Y Using RNA-seq data to select reference genes for
normalizing gene expression in apple roots.
893

Supplementary data

Supplementary Figure 1: Comparison of fold changes in TREM2 KO group among the three
analysed methods (qPCR FC computed using NF Conv and NF RNA-Seq and finally RNA-Seq).
Non-parametric ANOVA was performed using Kruskal Wallis test (red line comparing the
three groups). Multiple comparisons were performed using the Dunn's post hoc test using
the NF conv. group as the control condition (green lines). *The alpha value was set at 0.05*
*and P values are annotated as follows: * P<0.05, ** P<0.01, *** P<0.001.*

Supplementary Figure 2: Comparison of fold changes in TREM2 KO group among the three analysed methods (qPCR FC computed using NF Conv and NF RNA-Seq and finally RNA-Seq). Non-parametric ANOVA was performed using Kruskal Wallis test (red line comparing the three groups). Multiple comparisons were performed using the Dunn's post hoc test using the NF conv. group as the control condition (Blue lines). *The alpha*

*value was set at 0.05 and P values are annotated as follows: * P<0.05, ** P<0.01, ****

P<0.001.

Supplementary Table 1. Reference genes chosen from RNA-Seq data WT vs TREM2KO

ENSEMBL ID	Gene Name	baseMean	log2FC	pValue	pAdj	CVfrom Disp
ENSG00000169714	CNBP	14427.26	0.05	0.68	0.84	12.41
ENSG00000138279	ANXA7	6823.55	0.09	0.43	0.66	11.66
ENSG00000198231	DDX42	6558.35	-0.01	0.95	0.98	12.44
ENSG00000111667	USP5	5356.63	-0.03	0.82	0.92	11.50
ENSG00000204569	PPP1R10	4105.63	0.10	0.42	0.66	11.84
ENSG00000137177	KIF13A	3615.09	-0.02	0.85	0.94	12.32
ENSG00000164916	FOXK1	2502.17	-0.04	0.73	0.87	11.56
ENSG00000149089	APIP	1996.63	0.04	0.72	0.87	12.00
ENSG00000082258	CCNT2	1652.96	0.05	0.72	0.87	12.22

ENSG0000 0116221	MRPL37	1521.71	0.04	0.76	0.89	12.06

922 **Supplementary Table 2. Reference genes chosen from RNA-Seq data P3 vs P21 Sciatic**
923 **nerves**

ENSEMBL ID	Gene Name	baseMean	log2FC	pValue	pAdj	CVfromDisp
ENSMUSG000 00020485	Supt4a	878.49	0.03	0.81	0.88	12.93
ENSMUSG000 00031513	Leprotl1	964.45	-0.01	0.94	0.96	12.80
ENSMUSG000 00032826	Ank2	966.76	0.04	0.78	0.86	12.57
ENSMUSG000 00028581	Laptm5	1020.98	-0.03	0.86	0.92	12.89
ENSMUSG000 00031782	Coq9	1049.41	-0.06	0.68	0.79	11.99
ENSMUSG000 00030357	Fkbp4	1095.09	0.08	0.57	0.71	12.76
ENSMUSG000 00028161	Ppp3ca	1937.16	-0.05	0.74	0.84	12.76
ENSMUSG000 00033916	Chmp2a	3261.45	0.08	0.56	0.70	12.26
ENSMUSG000 00019899	Lama2	5070.82	0.00	0.99	1.00	12.25

ENSMUSG000 00028452	Vcp	6228.91	-0.10	0.48	0.63	12.34

Metadata to be submitted during review

Table 1: CV and Normfinder Metadata (WT vs TREM2KO IPSC)

Table 2: CV and Normfinder Metadata (P3 vs P21 SN)

Table 3: Primer sequences and amplification efficiencies

Table 4: Differential expression Raw data (WT vs TREM2KO IPSC)

Table 5: Differential expression Raw data (P3vs P21 SN)