A multiscale spatiotemporal model including a switch from aerobic to anaerobic metabolism reproduces succession in the early infant gut microbiota

David M. Versluis,[a] Ruud Schoemaker,[b] Ellen Looijesteijn,[b] Daniël Muysken,[d],†
Prescilla V. Jeurink,[b] Marcel Paques,[b] Jan M. W. Geurts,[b] Roeland M. H.
Merks,[a,c,*]

Leiden University, Institute of Biology, Leiden, The Netherlands[a]; FrieslandCampina, Amersfoort, the Netherlands[b]
Leiden University, Mathematical Institute, Leiden, The Netherlands[c]; Centrum Wiskunde & Informatica (CWI),
Amsterdam, The Netherlands[d];

ABSTRACT   The human intestinal microbiota starts to form immediately after birth, and is important for the health of the host. During the first days facultatively anaerobic bacterial species generally dominate, such as Enterobacteriaceae. These are succeeded by strictly anaerobic species, particularly *Bifidobacterium* species. An early transition to *Bifidobacterium* species is associated with health benefits: for example, *Bifidobacterium* species repress growth of pathogenic competitors and modulate the immune response. Succession to *Bifidobacterium* is thought to be due to consumption of intracolonic oxygen present in newborns by facultative anaerobes, including Enterobacteriaceae. To study if oxygen depletion suffices for the transition to *Bifidobacterium* species, here we introduce a multiscale mathematical model that considers metabolism, spatial bacterial population dynamics and cross-feeding. Using publicly available metabolic network data from the AGORA collection, the model simulates *ab initio* the competition of strictly and facultatively anaerobic species in a gut-like environment under the influence of lactose and oxygen. The model predicts that individual differences in intracolonic oxygen in newborn infants can explain the observed individual variation in succession to anaerobic species, in particular *Bifidobacterium* species. *Bifidobacterium* species become dominant in the model by using the bifid shunt, which allows *Bifidobacterium* to switch to sub-optimal yield metabolism with fast growth at high lactose concentrations as predicted here using flux-balance analysis. The computational model allows us to test the plausibility of hypotheses for bacterial colonization and succession in the infant colon.

IMPORTANCE   The composition of the infant microbiota has a great impact on infant health, but its controlling factors are still incompletely understood. The frequently dominant anaerobic *Bifidobacterium* species benefit health, e.g., because they can keep harmful competitors under control and modulate the intestinal immune response. Controlling factors could include nutritional composition and intestinal mucus composition, as well as environmental factors, such as antibiotics. We introduce a modeling framework of a metabolically realistic intestinal microbial ecology in which hypothetical scenarios can be tested and compared. We present simulations that suggest that greater levels of intra-intestinal oxygenation more strongly delay the dominance of

Bifidobacterium species, explaining the observed variety of microbial composition and
demonstrating the use of the model for hypothesis generation. The framework al-
lows us to test a variety of controlling factors, including intestinal mixing and transit
time. Future versions will also include detailed modeling of oligosaccharide and mucin
metabolism.

KEYWORDS: Infant microbiota, Microbial ecology, Flux balance analysis.

INTRODUCTION
The human infant microbiota starts forming directly after birth, and differs greatly
from the adult microbiota ([1]). Three main community types are observed in the infant:
(a) a Proteobacteria-dominated microbiota; (b) an Actinobacteria-dominated micro-
biota; and, less frequently (12-14% in one study ([2])), (c) a Bacilli-dominated microbiota
([1]). The three main community types are established after an ecological succession of
early communities. This ecological succession is thought to be controlled by nutrition
and early oxygen in the colon. To develop hypotheses on the potential mechanisms
and controlling factors of the initial development of the human microbiota, in particu-
lar the role of early oxygen, we introduce a computational model.
In the first 24 to 48 hours after birth, Proteobacteria, including *Escherichia coli* and
Enterobacter cloacae, and Bacilli, including *Streptococcus*, *Lactobacillus*, and *Staphylo-*
coccus, are the most common ([3], [1]). In the following weeks Proteobacteria are often
replaced by anaerobic Actinobacteria, mainly *Bifidobacterium* species, whereas Bacilli
are succeeded by either Proteobacteria or Actinobacteria. Other anaerobic species
are also found, but typically do not dominate. Actinobacteria generally persist as the
dominant group until weaning or the cessation of breastfeeding ([1]). A possible trigger
for the replacement of the Enterobacteriacea by *Bifidobacterium* spp., is depletion of a
hypothesized initial amount of oxygen ([4], [5]). Oxygen diffuses into the gut lumen from
the body and is taken up by bacteria, by colonocytes, and by non-biological chemi-
cal processes in the cecal contents ([6], [7], [8]) leading to oxygen depletion. The relative
importance of these processes is still under debate ([7], [8]).
Bifidobacterium species ([1]) are associated with a range of health benefits. For ex-
ample, early succession to a *Bifidobacterium* dominated microbiota is correlated with
reduced probabilities to be underweight at 18 months of age or to experience colic
([2], [9]). Acetate and lactate produced by *Bifidobacterium* also contribute to the acidifica-
tion of the infant gut, which suppresses many pathogens ([10]). A key question, there-
fore, is if and how *Bifidobacterium* can be promoted. The composition and develop-
ment of the infant microbiota is thought to be determined by many factors, including
nutrition ([11], [12], [13]), i.e. human milk or infant formula. The presence of lactose, the
most abundant carbohydrate in human milk and most infant formulas, is essential for
acquiring high levels of *Bifidobacterium* ([14]). This suggests that nutrition can play a
crucial role in promoting *Bifidobacterium*.
To give insight into the ecological and metabolic mechanisms underlying bacterial
colonization and succession in the infant colon, here we propose a multiscale, spatial
computational model of the infant gut microbiota. Earlier work has introduced similar
models of the adult gut microbiota ([15], [16], [17], [18]). These make use of flux balance
analysis (FBA) to model metabolism. Given a number of constraints including sub-
strate availability and enzyme availability, FBA uses genome-scale metabolic network
models (GEMs) to predict optimal biomass or energy production and the exchange
of metabolites with the environment. In spatial FBA, FBA models representing small

subpopulations of bacteria are coupled together to model a microbial, metabolizing ecosystem. While some approaches assume the whole system is in steady-state, allowing them to use the optimization approach for the whole ecosystem [16], here we build upon dynamic approaches such as BacArena [17], COMETS [19], and the approach by Van Hoek and Merks [18]. In these models, the metabolite exchange rates given by FBA are used to dynamically change metabolite concentrations in the environment, and predict the resulting effects on the bacterial populations. These properties make these dynamic, spatial FBA approaches highly suitable for our purpose of modeling the formation of the infant gut microbiota.

To analyze succession of bacterial species in a dynamic ecosystem such as the infant gut, it is key to accurately describe the dependency of growth rate and metabolite exchange on environmental metabolite concentrations. Previous work has shown that these are well described by constraints on the metabolic capacity of bacterial cells. These are given by an 'enzymatic constraint' which poses a limit on the summed metabolic flux through the whole cell [20], or through more advanced methods such as flux balance analysis with molecular crowding (FBAwMC) [21] which weighs the flux constraint according to the efficiency and volume of the individual enzymes [22]. Our model is, therefore, based upon the previous approach by Van Hoek and Merks [18] that uses FBAwMC. However, because such detailed data on enzyme efficiency and volume is unavailable for the majority of GEMs, here we have replaced FBAwMC with the simpler 'enzymatic constraint' approach [20]. The model represents the first 21 days of development of the infant microbiota in a spatially extended, gut-like environment, and considers three scales. At the microscopic scale, the model simulates the metabolism and growth of local populations of bacteria as a function of the local availability of metabolites. At the mesoscopic scale, the local environment dictates the metabolites available to the bacterial model, and receives the metabolites it deposits. At the macroscopic scale, a diffusion model simulates the spread of metabolites among adjacent lattice sites. Bacterial populations are also mixed locally, and can form new populations in their immediate environment. The metabolites advect distally. The system represents an ecosystem of 15 bacterial species, which are simulated using 15 GEMs taken from the AGORA collection [23], a database of 773 semi-automatically reconstructed GEMs of human gut bacteria that includes all major infant species [23].

The infant gut microbiota is partially limited by carbohydrates. Prebiotic carbohydrates increase population sizes *in vitro* and in mice inoculated with infant gut bacteria [24, 25]. The primary carbohydrate in human milk and nearly all infant formula is lactose, and infant formulas without lactose lead to a different infant microbiota with greatly reduced *Bifidobacterium* abundance [14]. We therefore focus on the carbohydrate metabolism and currently take lactose as the only input carbon source in the model, and ignore key nutrient sources such as human milk oligosaccharides, fats, protein residues, or intestinal mucus.

The model demonstrates the consistency of the hypothesis [4, 5] that a transition from an Enterobacteriaceae-dominated microbiota to a *Bifidobacterium* spp.-dominated microbiota is explained through gradual depletion of oxygen leading to anaerobiosis. The model suggests that high levels of initial oxygen can lead to prolonged dominance by *Enterobacteriaceae*, even after oxygen has been depleted, possibly explaining observed variation in the composition of the infant microbiota [1, 13], and reveals possible cross-feeding interactions between the species. Altogether the model shows to be a useful tool for qualitatively evaluating hypotheses on the dynamics of the infant gut microbiota.

FIG 1 The multi-scale metabolic model
(A) Schematic of the model at a system level. Circles represent bacterial populations, color represents species. Flow through the tube is from left (proximal) to right (distal). Lactose is placed at the proximal side. Output metabolites are examples, and depend on bacterial metabolism. Lattice dimensions and ratio are schematic. (B) Screenshot of the bacterial layer of the model at a single time point. Color indicates species, brightness indicates growth rate on the current timestep. (C,D) Screenshots showing, respectively, lactose and lactate concentration of the model corresponding to B. Intensity indicates concentration.

RESULTS

Model outline The model of the infant colon is discussed in detail in Section Methods. Briefly, given a typical infant colon length of 45 cm (26) and diameter of approximately 1.5 cm (27) the colon is described on a regular square lattice of 225×8 boxes of 2 $mm \times 2$ mm each, thus reaching a good balance between computational efficiency and accuracy (Fig. 1A). Because bacterial populations can vary at a much smaller scale (28, 29), the boxes are purely used as a computational discretization. Each lattice site can contain a population of one species, representing only the locally dominant bacterial species. It is described by a GEM from a list of 15 species prevalent in the earliest infant gut microbiota ((30, 11) selected from the AGORA collection (23) (Table 1, Methods section Species composition).

A simulation proceeds as follows. We initialize the system by placing approximately 540 small populations of the 15 species randomly across the lattice (Fig. 1B). We run the model in timesteps representing three minutes of real time, which proceed as follows: To mimic carbohydrates entering the colon from the small intestine, a pulse of lactose is introduced every 60 timesteps to the lattice sites of the six most proximal columns (Fig. 1A-1). Then we predict the metabolism of local bacterial populations using FBA. To mimic metabolic trade-offs, we use an enzymatic constraint that sets the maximum summed flux for each FBA solution. The uptake bounds for each population are set according to the local concentrations of metabolites (Fig. 1C&D). FBA returns a solution that maximizes the rate of ATP production, which is a suitable proxy for the rate of biomass production (31) given the focus on carbohydrate metabolism. This biomass production rate is used to update the local population size (indicated by

brightness in Fig. 1B). Populations spread into an available adjacent lattice site once
the population size has exceeded a threshold. The FBA solution also returns a set of
metabolite exchange rates, which are used to update the local metabolite concentra-
tions (Fig. 1D). In total, 723 extracellular metabolites are described in the model, but
only 17 of these are typically found in more than micromolar amounts in our simu-
lations (S1 Table). Bacterial populations diffuse through swaps of the content of ad-
jacent lattice sites (Fig. 1A-2). Metabolites diffuse between adjacent lattice sites (Fig.
1A-3) and advect to more distal sites (Fig. 1A-4) over one lattice per timestep, reaching
transit times of effectively 11 hours (32, 33). Metabolites and populations that reach
the distal side of the system (Fig. 1A-5) are removed. To mimic the appearance of new
bacterial populations (e.g. from ingested spores or introduction from intestinal crypts
(34)) each empty lattice site now has a small probability to be filled with a new bacterial
population selected at random from all available species (Table 1). This completes the
timestep, after which we record the metabolite concentrations and bacterial popula-
tion in the whole model. All parameters are given in Section Parameters. We have also
performed a sensitivity analysis for several parameters around the default parameter
set shown in Table 2 (Section Sensitivity analysis).

TABLE 1 Species included in our model, all from the AGORA collection(23). Color indicates color used in figures

Species	Family	Anaerobic? (23)	Reactions
Bacteroides vulgatus	Bacteroidaceae	Yes	2474
Bifidobacterium breve	Bifidobacteriaceae	Yes	1987
Bifidobacterium longum ssp. *infantis*	Bifidobacteriaceae	Yes	1005
Bifidobacterium longum ssp. *longum*	Bifidobacteriaceae	Yes	2043
Blautia hansenii	Lachnospiraceae	Yes	2055
Collinsella aerofaciens	Coriobacteriaceae	Yes	920
Dorea formicigenerans	Lachnospiraceae	Yes	2072
Parabacteroides distasonis	Porphyromonadaceae	Yes	2519
Ruminococcus gnavus	Lachnospiraceae	Yes	2225
Veillonella dispar	Veillonellaceae	Yes	1124
Enterococcus faecalis	Enterococcaceae	Facultative	1174
Enterobacter cloacae	Enterobacteriaceae	Facultative	1800
Escherichia coli SE11	Enterobacteriaceae	Facultative	2356
Lactobacillus gasseri	Lactobacillaceae	Facultative	1163
Streptococcus salivarius	Streptococcaceae	Facultative	1190

We performed several checks to ensure biochemical and thermodynamic plausi-
bility in the model (Methods section Checking the validity of the GEMs). After correc-
tions to the GEMs (S2 Table), no atoms were generated or lost by the FBA solutions,
there was no flux through the objective function without a substrate, and 99.999% of
all fluxes of Fig. 3 contained less free energy in the output than in the input (S1 Figure).
We concluded that the system conserved mass and energy.

Enzymatic constraint reproduces metabolic switching in *Bifidobacterium* Many
microorganisms switch between metabolic pathways of higher and lower yield de-
pending on substrate availability. These switches can be reproduced in many species
through an enzymatic constraint (20, 35, 36, 22) that limits the summed metabolic flux
of a bacterium. *Bifidobacterium* also displays a metabolic switch, where it is known as
the bifid shunt. At low extracellular concentrations of sugars *Bifidobacterium* produces
a mixture of acetate, ethanol, and formate in a 4:1:2 ratio, and at high concentrations
it produces acetate and lactate in a 3:2 ratio (37, 38).

To test if the enzymatic constraint suffices to explain the metabolic switch in *Bifi-*

FIG 2 Enzymatic constraint reproduces metabolic switching in *Bifidobacterium*
(A) Production of metabolites per timestep by a *B. longum infantis* population of $5 \cdot 10^9$
with access to one lattice site (0.05ml) using our FBA with enzymatic constraint method.
(B) Growth per timestep by lactose concentration for populations of $5 \cdot 10^9$ **bacteria with**
access to one lattice site (0.05ml) of some major bacterial species using our FBA with
enzymatic constraint method. *Bifidobacterium longum infantis* **is used as the *Bifidobac-***
***terium* representative.**

dobacterium, we performed simulations of a single timestep in increasing lactose con-
centrations (Fig. 2A). These correctly predicted the metabolic switch. The ratios be-
tween the metabolites before and after the metabolic switch, as well as the transient
between the two regimes match the experimental observations. Simulations without
the constraint only reproduced the situation at low concentrations of lactose (S2 Fig-
ure A), showing that the enzymatic constraint is required. The model also predicts
an unobserved metabolic switch on lactose in *E. coli* (S2 Figure B) and reproduced
the metabolic switch in *Bacteroides vulgatus* (39) (S2 Figure C). Another effect of the
enzymatic constraint is that the growth rate saturates at a species-dependent concen-
tration (Fig. 2B, S2 Figure D). As a consequence the relative growth rates of the species
depend on the lactose concentration.

Model predicts *Bifidobacterium* dominance through metabolism under anaer-
obic conditions After establishing the usefulness of the enzymatic constraint, we stud-
ied the behavior of the mathematical model for anaerobic conditions. After 21 days
(10080 timesteps) *Bifidobacterium* spp. had become the most abundant in 27 of 30 sim-
ulations (Fig. 3A, S1 Video), a typical situation found in infants of a corresponding age
(11, 2). Also in agreement with *in vivo* data, the model predicted the presence of pop-
ulations of *E. coli*. Our model also predicted populations of *Blautia hansenii*, which is
only occasionally present in the infant colon (11). We will investigate the niche of this
species further in the section 'Non-*Bifidobacterium* species benefit from lactate con-
sumption'. All species other than the aforementioned had abundances either lower
or not significantly higher than their initial abundance after 21 simulated days (p>0.05,
two-sample t-test). To test if these species were required for the formation of the com-
munity we ran an additional 30 simulations in which we only included species that had
an abundance significantly larger than their initial abundance at the end of other sim-
ulations. This led to a very similar set of outcomes (S3 Figure). We decided to continue

FIG 3 *Bifidobacterium* spp become dominant anaerobically and mostly proximally with a unique metabolic profile

(A) Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (B) Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation. n=30. (C,D) Visualisation of metabolic interactions in the sample run of S2 Video. Green lines represent production, yellow lines represent consumption. Line width is relative to quantity exchanged with the environment. Data from hour 90 to 93 (step 1800 to 1860) for C and last 3 hours (60 timesteps) for D (E) Spatial distribution of lactate over the last two days (960 timesteps) of the simulation. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (F) Abundance per location for grouped species over the last two days (960 timesteps). Curve shows mean value and shaded area one standard deviation over n=30 simulations. (G) Metabolism of lactose by grouped species. All data from last two days. n=30. (H) Metabolism of lactate by grouped species. All data from last two days. n=30.

including these species to prevent an initial bias, and allow for the possibility that they become abundant under alternate conditions.

We next analysed the metabolites diffusing or advecting out of the distal end of the gut over the last two days of the simulations of fig. 3A. We found considerable amounts of acetate, formate, lactate, hydrogen, ethanol, and succinate (Fig. 3B). All other metabolites were present in much smaller quantities. The high abundance of acetate and lower abundance of lactate agree with the fecal composition of three month old formula-fed infants (40), and is characteristic of microbiota dominated by *Bifidobacterium*. Formate, hydrogen, ethanol and succinate are also all reported in infant feces (41, 42, 43, 44).

To examine how this metabolic pattern is formed we analysed the metabolic production and uptake in a sample run (Fig. 3C). *Bifidobacterium* converts lactose into a mix of acetate, lactate, ethanol, and formate, as predicted (Fig. 2A), with a higher lactate production than ethanol or formate. Some of this lactate is reabsorbed and converted into the three other metabolites by *Bifidobacterium*. A mixture of lactose and crossfeeding products is converted by *E. coli* and *B. hansenii* into the remaining metabolites of Fig. 3B. The main crossfeeding nutrient absorbed is lactate, produced by *Bifidobacterium*, but there is also uptake of ethanol, formate, acetate, and succinate. These are also produced by the crossfeeders. The network has become simpler at the end of this run, as *B. hansenii* is at a very low abundance and *E. coli* now only consumes lactose, lactate, and ethanol, which it does not produce (Fig. 3D).

Crossfeeding emerges in simulations The observed crossfeeding in the sample run (Fig. 3C&D) suggested that production of metabolites by *Bifidobacterium* spp. may be crucial for *E. coli* and *B. hansenii*. To further analyse such crossfeeding interactions we next analysed the spatial distribution of the primary Bifidobacterial metabolites. From day 19 to 21, lactose was present mostly proximally (S4 Figure A), lactate was most abundant in the middle of the colon (Fig. 3E), and acetate was most abundant distally (S4 Figure B). To analyse the role of *Bifidobacterium* spp. in shaping this metabolic pattern, we analysed the location of populations from day 19 to 21 (Fig. 3F). *Bifidobacterium* spp. was located proximally, whereas the other species were located more distally. These results suggested that separated metabolic niches had formed in the simulated colon. To quantify crossfeeding interactions we summed the flux of lactose and lactate for *Bifidobacterium* and for all non-*Bifidobacterium* species grouped together. *Bifidobacterium* spp. was the largest consumer of lactose (Fig. 3G) and both the largest producer and consumer of lactate (Fig. 3H). However, the non-*Bifidobacterium* groups consumed more lactate relative to their lactose consumption. We performed additional simulations in which *Bifidobacterium* was not capable of consuming lactate, by blocking it from converting lactate to pyruvate. In these simulations *Bifidobacterium* was the largest single group in 20 of the 30 runs, with the largest average abundance (S5 Figure A). From this, we hypothesized that *Bifidobacterium*'s primary niche is as primary consumer, consuming lactose, while for other species niches exist as primary and/or secondary consumer around consuming lactose or lactate. Combined with the observation (Fig. 3D) that lactate concentration is lower in the distal colon and the recorded fluxes in our network visualisation (S2 Video), we can conclude that the population at the distal end of the simulated colon consists at least partially of such secondary consumers, i.e. crossfeeders.

Lactose uptake through the bifid shunt is essential for *Bifidobacterium* dominance Next we determined whether the bifid shunt is essential to *Bifidobacterium* dominance in the model. We disabled the bifid shunt by blocking all flux through the reactions associated with fructose-6-phosphate phosphoketolase and ran 30 simula-

FIG 4 *Bifidobacterium* and *E. coli* require lactose, but *B. hansenii* requires lactate
(A) Abundances for grouped species over 21 days. The bifid shunt was disabled for all *Bifidobacterium* species. One day is 480 timesteps.Curve shows mean value and shaded area one standard deviation over n=30 simulations. (B) Abundances for grouped species over 21 days, but with lactose disabled for non-*Bifidobacterium* species from 5040 timesteps (10.5 days) onwards. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (C,D) Visualisation of metabolic interactions in a sample run. Green lines represent production, yellow lines represent consumption. Line width is relative to quantity exchanged with the environment. Data from hour 90 to 93 (step 1800 to 1860) for C and last 3 hours (60 timesteps) for D. (E) Growth per timestep by lactate concentration for populations $5 \cdot 10^9$ with access to one lattice site (0.05ml) of some major bacterial species (F) Abundances for grouped species, with lactate consumption disabled for all species. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

tions. Fructose-6-phosphate phosphoketolase is unique to and necessary for the bifid
shunt, and characteristic of *Bifidobacterium* as a genus (45). *Bifidobacterium* spp. were
dominant in none of these simulations (Fig. 4B, S3 Video). Instead, *E. coli*, *B. vulgatus*,
and *B. hansenii* were dominant. This indicates that the bifid shunt is crucial to the dom-
inance of *Bifidobacterium* spp. in our model. We also ran 30 simulations in which only
the production of lactate was disabled in *Bifidobacterium* (S6 Figure). These simula-
tions still led to *Bifidobacterium* dominance in 27 out of 30 simulations. This indicates
that the metabolic switch to lactate production was not essential to *Bifidobacterium*
dominance in our model.

B. hansenii but not E. coli can sustain itself on crossfeeding We observed that
the non-*Bifidobacterium* species did not occur as strongly proximally as *Bifidobacterium* spp.
(Fig. 3F), but did consume lactose (Fig. 3D&G), and analyzed how these non-*Bifidobacterium*
species could persist in the model. More distal parts of the colon have a lower lac-
tose concentration compared to the proximal part (S4 Figure A), and in the presence
of low concentrations of lactose *E. coli* produces more ATP per timestep than *Bifi-*
dobacterium spp. (Fig. 2B). We therefore hypothesized that these non-*Bifidobacterium*
species outcompete *Bifidobacterium* on low-concentration lactose in the model.To test
this, we ran 30 simulations in which lactose consumption was disabled for non-*Bifidobacterium*
species at the midway point of the simulation (10.5 days). By disabling it after 10.5
days we rule out that the non-*Bifidobacterium* species die out before *Bifidobacterium*
has produced crossfeeding metabolites. There was a sharp population decrease of
non-*Bifidobacterium* species from the point where lactose uptake was disabled (Fig.
4B). *E. coli* had a near-zero abundance at the end of these simulations. *B. hansenii* also
had a significantly lower abundance compared to the baseline (p<0.01, two-sample
t-test), but still had a presence (S3 Video). This indicates that some amount of primary
consumption is essential for *E. coli*, but not for *B. hansenii*.

Non-Bifidobacterium species benefit from lactate consumption To further anal-
yse how *B. hansenii* could sustain a population as a purely secondary consumer, we
investigated the substrates used by non-*Bifidobacterium* species in the model. Analysis
of the flow of metabolites in a different run from fig. 3A showed the uptake of lactose
by *B. hansenii* decreasing over time, while uptake of lactate increased (Fig. 4C&D). *E.*
coli and *B. hansenii* also produce more ATP from lactate than *Bifidobacterium* at any
concentration (Fig. 4E). Combined with our earlier observations of crossfeeding on
lactate (Fig. 3D&H) we hypothesized that *Bifidobacterium* cannot compete with *E. coli*
and *B. hansenii* species on lactate. Consumption of lactate by *B. hansenii* could be es-
sential to its ability to sustain populations without lactose uptake.

To investigate if lactate is essential in feeding secondary consumer populations,
we blocked the lactate uptake reaction for all species and ran 30 model simulations.
(Fig. 4F, S3 Video). At 21 days *B. hansenii* populations were much smaller than in sim-
ulations with functional lactate uptake, not significantly larger than the initial popu-
lations (p>0.05, two-sample t-test). *E. coli* populations were also significantly smaller
than those in simulations with lactate uptake, but also significantly larger than their
initial population at the start of the run (p<0.05, two-sample t-test). *Bifidobacterium*
populations were similar to those in the first simulations (Fig. 3A). Together with the
results in Fig. 4B this indicates that *E. coli* does not require lactate as a crossfeeding
substrate like *B. hansenii* does.

E. coli outcompetes Bifidobacterium by taking up early oxygen During the first
days Enterobacteriaceae are dominant in the *in vivo* infant gut microbiota, after which
Bifidobacterium spp. are often dominant (1, 11). Initially present oxygen in the gut
may explain this pattern (5, 3). Oxygen could have diffused through the gut wall (7)

and accumulated in the newborn infant gut *in utero* in the absence of microbes (46).
Intracolonic oxygen diffusion depends on host tissue oxygenation and colonic blood
flow (47, 48). In the absence of precise data on the concentration and variation of oxy-
gen levels in the infant gut, we examined this hypothesis by testing oxygen values from
0.1 to 10 μmol initial oxygen per lattice site, which covers a wide range of outcomes
in our model. Oxygen was diffused, but was not moved distally every timestep, thus
mimicking an early presence of oxygen. 0.1 μmol initial oxygen sufficed to reproduce
the pattern observed *in vivo*: *E. coli* is initially dominant and *Bifidobacterium* becomes
dominant after a few days or weeks of simulated time (Fig. 5A, S4 Video, S5 Video).
The initial presence of oxygen also strongly affected the other species in the model -
B. hansenii remained nearly entirely absent, while *B. vulgatus* had a higher abundance
than it did without initial oxygen. In the network analysis of a sample run a similar pat-
tern occurs as in the networks of figure 3C, but with different early species (Fig. 5B).
Here the Enterobacteriaceae *E. coli* and *Enterobacter cloacea* take up more of the early
lactose, along with oxygen. By the end of the run *Bifidobacterium* again took the pri-
mary consumer and lactate producing role and *E. coli* had transitioned to a stronger
secondary consumer role, feeding largely on lactate (Fig. 5C).

At 1 or 10 μmol initial oxygen per lattice site *E. coli* and *E. cloacea* remained dom-
inant and the main primary consumers for longer (Fig. 5D, S6 Video), and succession
to *Bifidobacterium* took much longer, or did not occur at all within the simulated time-
frame (Fig. 5E). At 10 μmol oxygen per lattice site *Bifidobacterium* either dominated
the population or remained much smaller, with few intermediate cases, leading to
a bimodal distribution (Fig. 5F). This prediction matches *in vivo* observations (1, 13).
The proportion of infants dominated by Enterobacteriaceae at 21 days of age varies
depending on the study population, but can be 22% (2) or 25% (49) after 21 days, an in-
termediate value between the 10% (Fig. 5D) and the 60% (Fig. 5E) that we observe with
different oxygen levels. We also initialised four sets of simulations in which oxygen
was released from the gut wall every timestep (S8 Figure). This led to an increasingly
stronger stimulation of *E. coli* with higher levels of oxygen, but did not lead to any
of the temporal effects observed *in vivo*. In a separate set of simulations we ended
oxygen release at the midpoint of our simulations. We observed an increase of *Bifi-
dobacterium* spp. and a decrease of *E. coli* after that point (S9 Figure A), leading to a
bimodal outcome similar to the runs with initial oxygen (S9 Figure B).

We further examined the causes of *E. coli* dominance over *Bifidobacterium* in the
presence of oxygen in our model. Fig. 5G shows that when oxygen is available *E. coli*
has a much higher growth per timestep in our model than *Bifidobacterium* spp. for all
concentrations of lactose, even though *Bifidobacterium* spp. also produce more ATP
than under anaerobic conditions. *B. hansenii* and *S. salivarius*, in contrast, grew slower
than *Bifidobacterium* for all concentrations in the presence of oxygen. *E. coli* does not
use overflow metabolism in our model, though *E. cloacea* does, and *E. coli* does when
in the presence of glucose (S7 Figure).

To test if the direct uptake of oxygen was indeed responsible for *E. coli*'s domi-
nance of the microbiota, we disabled the oxygen uptake reaction of *E. coli* for 30 simu-
lations with 1 μmol initial oxygen per lattice site. Under these conditions, *E. coli* failed
to dominate the population (Fig. 5H). Other species, primarily *E. cloacae*, became dom-
inant early in the simulations as a primary consumer (S6 Video). These species are
replaced in all simulations by a population composition similar to that from the simu-
lations without oxygen. Altogether, the model predicts that *E. coli* relies on direct con-
sumption of oxygen to dominate the microbiota under initially oxygenated conditions
in the model. It also indicates that *E. cloacea* does not have an anaerobic metabolism

FIG 5 Initial oxygen leads to initial *E. coli* dominance and succession by *Bifidobacterium* (A,D,E) Abundances for grouped species. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. Initial oxygen per lattice site in μmol is (A) 0.1 (D) 1 (E) 10 (B,C) Visualisation of metabolic interactions in a sample run. Green lines represent production, yellow lines represent consumption. Line width is relative to quantity exchanged with the environment. Data from hour 30 to 33 (step 600 to 660) for B and last 3 hours (60 timesteps) for C (F) Distribution of total *Bifidobacterium* abundance at 21 days (10080 timesteps) performed with 10 μmol initial oxygen per lattice site n=30. (G) Growth per timestep by lactose concentration in the presence of abundant oxygen for populations $5 \cdot 10^9$ with access to one lattice site (0.05ml) of some major bacterial species (H) Abundances for grouped species with oxygen uptake disabled for *E. coli*. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

competitive enough to sometimes become dominant over *Bifidobacterium* spp. as *E.*
coli does.

Sensitivity analysis A number of key parameters in our model were based on
reasonable estimates. To test the effects of these parameter values on the simulation
results, we performed a sensitivity analysis under anaerobic conditions, which best
represents the situation at the end of the simulations (Fig. 6). As shown in Fig. 2B and
S2 Figure D, after relaxing the enzymatic constraint *Bifidobacterium* spp. loses its com-
petitive advantage at high lactose concentrations (S7 Video). This leads to reduced
dominance or extinction in whole-gut simulations (Fig. 6A). After tightening the enzy-
matic constraint only *Bifidobacterium* spp. remained, with a much smaller population
size. Thus the enzymatic constraint is crucial for a diverse microbiota in which *Bifi-*
dobacterium can become dominant. The ATP required for a unit of population growth
had a linear effect on population size, as expected, but had little effect on the relative
abundance of *Bifidobacterium* (Fig. 6B). The rate of colonization had little effect on the
simulation outcomes: at increased colonization rates, *Bifidobacterium* spp. still dom-
inates the microbiota and its relative abundance remains unchanged (Fig. 6C). Even
eliminating colonization altogether hardly affected *Bifidobacterium* spp. dominance in
most simulations, but led to complete extinction of all *Bifidobacterium* species in 5/30
simulations and of all 12 non-Bifidobacterium species in 9/30 simulations. Thus we
kept a moderate amount of colonization in the model. We also ran examined the ef-
fect of placing new populations after initialization only in the first column of the lattice,
with a probability increased to match the lower number of eligible squares (S10 Fig-
ure). This also led to a slightly larger but still comparatively small population for the
non-*Bifidobacterium* species. To determine the optimal spatial discretization we ran
the model on more coarse and on more refined lattices, scaling the diffusion, advec-
tion, and initialization and colonization parameters accordingly (Fig. 6D; see Methods).
Lattice refinement did not change the dominance of *Bifidobacterium* nor the spatial
distribution of metabolites and populations (S11 Figure A-D), but on a more coarse lat-
tice the smaller number of subpopulations seemed to increase the rate of extinctions
and variability of metabolite distributions (S11 Figure E-H). The diffusion of metabo-
lites has a larger effect on the simulation results (Fig. 6E). At low diffusion rates, the
populations tended to die out, because they had less opportunity to take up lactose
as they were exposed only as it 'passed by' due to advection. At high diffusion rates
cross-feeders and *Bifidobacterium* consumed all lactate (S12 Figure); in infant feces
substantial amounts of lactate are found ((40, 50)). We therefore decided to choose
a value for which the system produced the observed concentration of lactate. Finally
the amount of mixing of bacteria was varied (6F). *Bifidobacterium* remained dominant
regardless of the diffusion parameter, even when the bacteria were fully mixed (i.e.
placed at random locations) each timestep. Without mixing the populations could not
spread, for lack of empty space. We have maintained bacterial diffusion at the base-
line to allow for a similar level of diffusion of metabolites, while giving each lattice site
one movement chance.

DISCUSSION

To develop new hypotheses for possible external effects on the initial phases of the in-
fant gut microbiota, we have developed a dynamic multiscale model. Our simulations
reproduce the initial dominance of Enterobacteriaceae, particularly *E. coli*, over *Bifi-*
dobacterium species, and the subsequent dominance of *Bifidobacterium* species, out
of a broad consortium of infant species. Moreover, our simulations suggest that this
could be due to an initial presence of oxygen, followed by microbial depletion. These

FIG 6 Sensitivity analysis for estimated parameters
Abundances for *Bifidobacterium* spp. and other species at the end of day 21 with different parameters. n=30 for each condition. **(A) Effect of enzymatic constraint around the default value of 4 (B) Effect of amount of ATP required for growth around the default value of 10 (C) Effect of variation in continued influx of populations per timestep per empty square around the default of 0.00005 (D) Effect of variation in width and height of lattice, including adjustment of associated parameters, around the default of 8x225 (E) Effect of variation in diffusion for metabolites away from the baseline of** $4.4 \cdot 10^{-5}$ **(F) Effect of variation in diffusion of bacterial populations away from the baseline of** $5.7 \cdot 10^{-5}$

predictions on oxygen suggest that higher levels of initial oxygen may explain the absence or strong delay of *Bifidobacterium* development that is observed in some infants (2). A continuous input of oxygen did not explain the pattern observed *in vivo* (S8 Figure), but stopping the input of oxygen at the midway point of the simulation did lead to a similar pattern (S9 Figure).

While it is known that there is variation in colonic oxygenation (47, 48), the range of this variation in newborns is not known. In premature infants a positive association exists between the number of days of supplemental oxygen and *Klebsiella* sp. abundance (51). *Klebsiella* was the primary representative of the Enterobacteriaceae in this study population. However, only very few of these infants developed a *Bifidobacterium* dominated microbiota, regardless of the duration of supplemental oxygen. Nonetheless, it may be desirable to lower oxygen concentration faster to achieve the positive health effects of *Bifidobacterium* more consistently. As the direct oxidation of lipids and other organic substrates may contribute to creating anaerobic conditions in the colon (7), altered nutrition may allow for a decrease in oxygen in the colon without stimulating facultative anaerobic bacteria such as *E. coli*. There are also indications that short-chain fatty acids produced by *Bifidobacterium* may decrease oxygen levels by stimulating the oxygen use of colonocytes (8). Although oxygen use by colonocytes may play a role, our model predicts that *Bifidobacterium* is not abundant until oxygen levels have decreased, and so do not produce large amounts of short chain fatty acids under these conditions.

Our simulations predict that the dominance of *Bifidobacterium* spp. in the anaerobic infant system could be explained from its consumption of lactose through its bifid shunt metabolism. Our model can correctly represent bifid shunt metabolism due to the implementation of an enzymatic constraint on the maximum total metabolic flux in the bacteria (Fig. 2). Both the low and high yield metabolism, and the switch between them based on nutrient availability, have been extensively described *in vitro* (38). The FBA with enzymatic constraint method we use is similar to flux balance analysis with molecular crowding (FBAwMC) (21). As in FBAwMC we consider the enzymatic constraint to represent the limited availability of space and production capacity for enzymes within a bacterial cell. However, in the absence of good reaction-specific thermodynamic information for our current model, we do not utilize the reaction-specific technique of FBAwMC. We instead base an enzymatic constraint only on the size of the local population, which is sufficient for modeling a metabolic switch. This is equivalent to using FBAwMC and placing all crowding coefficients at the same positive number, which has been done previously to model metabolic switches (21). More broadly, our method is similar to other existing methods that model metabolic switches by including limits on fluxes that represent some physical limit on metabolism, such as proteome constraints or membrane occupancy (52, 35, 53, 54). It has been shown that metabolic switches in microbes can be modelled in many ways, provided that two simultaneous constraints are in place (22). In our case, these are the concentration of substrate and the enzymatic constraint.

The metabolic switch of *Bifidobacterium* in our model determines the production of its metabolites. The two most abundant, acetate and lactate, occur in the simulated feces at a median ratio of 3.5:1 (Fig. 3B), similar to that found in formula-fed infant feces (40, 50). We use ratios here because the absolute measures, per gram or per milliliter, are not available in our model. The median ethanol:acetate ratio of 1:21 is similar to the ratios of 1:18 and 1:10 reported in term infants (43) and the 1:50 reported in pre-term infants (55). We observe a lower quantity of ethanol in the metabolic output of the simulations where lactate uptake by *Bifidobacterium* was disabled (S5 Figure

B).

Bifidobacterium takes up lactate in our model and converts it into pyruvate, cat-alyzed by lactate dehydrogenase. However, strong lactate uptake by *Bifidobacterium* spp. has not been observed *in vitro* (in e.g. ([56])). Thus, while the enzymes used for lactate consumption exist in *Bifidobacterium*, it is not known if the pathway is used *in vivo* to the extent predicted by the model. The small difference in abundance between the condition with and without lactate uptake by *Bifidobacterium* (S5 Figure A) suggests that it may occur as a purely primary consumer *in vivo*. Similarly, though growth on lactate by *E. coli* in the presence of oxygen is well documented, this is not the case for the anaerobic condition ([57]). Lactate uptake is not essential for *E. coli*, as it can exist in a more purely primary consumer role (Fig. 4F). The third consumer of lactate, *B. hansenii*, is present only marginally *in vivo* ([11]). This matches with our results for initially oxygenated conditions, in which *B. hansenii* is also largely absent. However, other species that could fill the same lactate-consuming secondary consumer niche as *B. hansenii* in our model are common *in vivo* ([58]). Our selection of species may be further improved to allow the correct species to emerge in this secondary consumer niche. For example, it might be appropriate to add *Eubacterium hallii*, a common infant gut bacterium known to consume *Bifidobacterium*-produced lactate and acetate *in vitro* ([59]).

More generally, our model predicts a lower diversity overall in both bacterial species and metabolites compared to what is observed *in vivo* ([11]). While the abundant species in our model largely match with the abundant species *in vivo*, many other species have a lower abundance in our model compared to the *in vivo* conditions. We have decided to keep all less abundant species in our simulations, to show how in most cases the correct species out of a broad consortium become dominant. There were no large differences when we ran the simulation without these smaller species (S3 Figure). Particularly the lack of representation of the Bacilli species in our model outcomes is notable. While some sources report a portion (e.g. 12-14% ([2])) of their subjects to be dominated by Bacilli, mainly *Streptococcus*, the *Streptococcus* species in our model never became dominant. We showed that *Streptococcus* has an inferior metabolism on lactose and lactate compared to other species in our model (Fig. 2A, Fig. 4E, Fig. 5G).

The discrepancy between our model and the *in vivo* data in the abundance of less common species such as *Streptococcus* may partially be explained through the focus of our model on carbon dissimilation. Though it would be preferable to represent the whole metabolism of the infant microbiota, including factors such as consumption of amino acids, oligosaccharides and intestinal mucus, the initial version presented here only considers carbon metabolism from lactose. While a more extensive metabolism might have allowed more mechanisms and niches to be discerned, it would also introduce additional free parameters, as there is no clear data on the concentration and uptake of these nutrients. The uptake bounds would have to be set arbitrarily, with uniform values or random sampling. Many substances such as fatty acids or protein residues are not even included as metabolites in the database we use for bacterial metabolism ([23]), further complicating any introduction. By focusing on carbon metabolism, using ATP production as a proxy for growth rate, and only using lactose as an input nutrient, we can circumvent these problems. ATP production has been shown to be a good proxy for biomass production in *E. coli* ([31]). In addition, the influence of prebiotic carbohydrates on the *in vitro* and model organism infant gut microbiota has been shown to be large ([24], [25]). This indicates that carbon metabolism is a limiting factor, especially in the absence of prebiotics, as in our model. However, the addition of prebiotics led to an increase primarily in *Bifidobacterium* populations, which does

not necessarily mean other species were also carbon limited. In addition, ATP pro-
duction may not be a good proxy for biomass production in some species. Species-
specific information on the relation between ATP and biomass production may aid
future modelling. Finally, gaps in the GEMs we use may cause ATP production itself to
be underestimated compared to what occurs *in vivo*.

Outside of our modelling of metabolism, our handling of diffusion could also cause
discrepancies between our data and the *in vivo* situation. The sensitivity analysis dis-
played an unrealistic metabolic output when metabolic diffusion was raised to $2.2 \cdot 10^{-4}$
cm^2/s (S12 Figure). Modelling of the passage of spheres in the adult gut has suggested
even higher values, around $1 \cdot 10^{-2}$ cm^2/s for both bacteria and metabolites ([60]). How-
ever, motor patterns strongly influence the amount of mixing in the gut ([61]) and it is
unclear to what extent motor patterns in the adult gut translate to the infant gut ([62]).
Motor activity which transports colonic contents over long distances is more common
in infants than in older children or adults ([63]), while motor activity which mixes the
colon contents and slows net distal movement is rarer in infants than in older chil-
dren or adults ([64], [63]). This corresponds with the much faster colonic transit in in-
fants ([32], [33]) compared to that found in older children and adolescents ([65]). Taken
together, these results suggest that the low diffusion that our model requires for a
plausible fecal composition may be a feature of the infant colon.

Our modeling approach is similar to several previously published frameworks,
such as Steadycom, BacArena, and COMETS ([16], [66], [17], [67]). While these previous mod-
els are useful as frameworks for many applications, we chose to continue developing
our own line of gut models based on the previous work by Van Hoek and Merks ([18]).
This previous model of a dynamic, evolving microbiota was a good fit to the infant gut
microbiota, which is similarly dynamic. Nonetheless, further integration of methods
would allow for better comparison between models, and our model could potentially
be implemented in the new COMETS 2 framework ([19]).

We have used our modeling approach to generate testable hypotheses on the
causes and mechanics of succession in the infant gut microbiota, potentially laying a
base for nutritional interventions that could improve the health of infants. It should
be emphasized that the current and future work on this model represents a tool for
generating hypotheses, and for testing potential mechanical explanations. We cannot
fully represent the complexities of the infant gut microbiota, and any generated hy-
potheses must necessarily be validated *in vitro* and *in vivo*. There is work to be done
to further study the relevant factors, and to bring these results into a more realistic
model context. We aim to do so by integrating the prebiotic oligosaccharide and pro-
tein content of nutrition into the model. This may lead to the creation of more niches
in the model, and so a diversity closer to that of the *in vivo* system. We will also con-
tinue to improve the selection and curation of metabolic models, such as by disabling
lactate uptake in certain species or including a wider diversity of GEMs. This may lead
to novel future insights on the interactions between differences in infant nutrition and
succession in the infant microbiota.

METHODS

We use a spatially explicit model to represent the newborn infant microbiota (Fig. 1B).
Our model is based on an earlier model of the adult gut microbiota ([18]). The model
consists of a regular square lattice of 225 by 8 lattice sites, where each lattice site
represents 2mm by 2mm of space, resulting in an infant colon of 450mm by 16mm.
Each lattice site can contain any number of metabolites of the 723 types represented
in the model in any concentration and a single simulated bacterial population. The

metabolism of these bacterial populations is based on genome-scale metabolic mod-
els (GEMs) from the AGORA collection of species-specific GEMs (23). From this set,
we have chosen 15 GEMs based on a consortium of known infant microbiota species
(30, 11). Their metabolic inputs and outputs are calculated using dynamic flux balance
analysis (FBA) (68) with an enzymatic constraint functioning as a limit on the total flux
through the network (20). The effects of the FBA solution are applied to the spatial
environment and then recalculated each timestep, creating a spatial dynamic FBA.

We identify the two narrow ends of the rectangle with the proximal and distal ends
of the colon. In each timestep metabolites both mix and flow from the proximal to the
distal end. To mimic adhesion to the gut wall and mixing through contractions (69),
bacterial populations are mixed, but do not flow distally as metabolites do.

At the start of each run, we initialize the system with a large number of small popu-
lations. We let these perform their metabolism each timestep. They take metabolites
from the environment, and deposit the resulting products. We let the populations
grow and divide according to their energy output. Both the initial placement locations
and movement of the populations is random, introducing stochasticity in the model.
The system develops differently depending on initial conditions, into a diverse and
complex ecosystem.

Species composition Each population is represented by a GEM of a species. 15
different metabolic models are used in our spatially explicit model (Table 1), selected
based on previous research (30). From this list of genera, the most prevalent species
within that genus in the vaginally born newborn data set from Bäckhed *et al.* was
used (11). One group from the list could only be determined at the family level - Lach-
nospiraceae. *Ruminococcus gnavus* was chosen to represent this group, based on its
high prevalence among this group and the prior inclusion of species from the *Blautia*
and *Dorea* genera (70). Because the genus *Bifidobacterium* is known to be particularly
diverse (71), we represented it with models of three different strains. All models are
based on the GEMs created by Magnúsdóttir et al. in the AGORA collection (23).

Checking the validity of the GEMs All GEMs used in the model were tested indi-
vidually to ensure that they could grow on a substrate of lactose. Only *Veillonella* did
not pass this test, which is consistent with *in vitro* observations (72). *Veillonella* did pass
when lactate, a common infant gut metabolite, was used instead. We also tested all
GEMs individually for spurious growth in absence of substrates. To this end, all uptake
bounds except water were set to 0. None of the GEMs grew under these conditions.
All models were tested for having a net neutral exchange of hydrogen, carbon, oxygen
and nitrogen at each timestep of the model during the simulations. There should be
no net change in the number of atoms in the medium due to the calculated fluxes,
because the reaction we set as an objective function does not remove any atoms, and
there are no other sinks in the simulations. In the present simulation, only water, oxy-
gen, and lactose are introduced into the simulation, so no other atoms than these
three are considered. The tests revealed two errors in glycogen metabolism in sev-
eral GEMs in AGORA that resulted in a energetically-favorable removal of intracellular
protons from the system. We corrected these GEMs by replacing the reactions respon-
sible for this erroneous energy source (S2 Table). With the corrections we applied to
the model, these tests always passed - allowing for rounding errors of less than $1 \cdot 10^{-8}$
μmol per FBA solution.

The thermodynamic correctness of all reactions was checked by calculating the
net difference in Gibbs free energy between input and output metabolites, using a
pre-existing dataset of thermodynamic source data (73, 74). The conditions assumed
for the calculation of the Gibbs free energy were a pH of 7 and an ionic strength of

0.1M ([73](#)). This was recorded for each population at each timestep over the course of a full run of 10080 timesteps. 99.999% of all FBA solutions in the simulations of Fig. [3](#) passed the test by containing less free energy than the associated inputs (Fig. [1](#)E). The remaining 0.0001% all had an amount of energy in the outputs equal to that of the inputs. All of these solutions had very low growth rates, less than 0.001% of the average growth rate. The sum of these growth rates over 30 simulations was less than 0.05 of an initial population, totalling $2 \cdot 10^6$ bacteria.

Changes from AGORA We use updated versions of the AGORA GEMs ([75](#)), to which we have applied checks and modifications. Firstly, the objective function was changed from the biomass reaction included in the models to a reaction only requiring ATP production. As this reaction yields only ADP and P_i, it is is mass-neutral. This allows us to focus on carbon dissimilation within the GEMs, and the differences in underlying metabolism. ATP yield has been a good proxy for biomass production in previous studies ([76](#)). Focusing on carbon dissimilation means we can leave all unknown uptake bounds at 0, instead of using an arbitrary or randomized level, as in some other studies ([17](#), [16](#)).

We also checked the metabolic networks for reactions that allow for the occurrence of unrealistic FBA solutions, and we have added additional reactions. In the *Bifidobacterium* models the ATP synthase reaction was made reversible. This allows it to function as a proton pump, which *Bifidobacterium* species use to maintain its internal pH ([77](#)). Lactose permease reactions were added to all *Bifidobacterium* species in the model ([78](#)) based on those available in other models in the AGORA collection ([23](#)). Combined with the existing reactions in the model and the metabolic constraints this leads to a set of reactions simulating a realistic bifid shunt yield, when substrate is abundant, of 5 mol ATP, 2 mol lactate and 3 mol acetate per mol of lactose([38](#), [79](#)). Lactose permease was also added to *Streptococcus salivarius* and *Ruminococcus gnavus* to bring them in line with existing literature on their *in vitro* behavior ([80](#), [81](#)). *Veillonella dispar* is the only species in the model that does not have any lactose uptake ([72](#)). A complete list of changes is presented in [S2 Table](#).

FBA with enzymatic constraint Each timestep of the model, a modified version of flux balance analysis with an enzymatic constraint is used by each population to determine what reactions it should use to achieve the most biomass production from the metabolites available to it ([82](#), [20](#)). First, each GEM is converted to a stoichiometric matrix S. All reversible reactions are converted to two irreversible reactions, so that flux is always greater than or equal to 0. All reactions identified as exchange, sink, or demand in the metabolic reconstruction are marked as exchange in the matrix. These reactions exchange nutrients or metabolites with the environment. Each timestep, all reactions are assumed to be in internal steady state ([82](#)):

$$S \cdot \vec{f} = 0 \tag{1}$$

Where $\vec{f}$ is a vector of the metabolic fluxes through each reaction in the network (in mol per time unit per population unit).

Each exchange reaction (i.e. each reaction that takes up metabolites from the environment or deposits them) is constrained by an upper bound $F_{ub}(m)$ which represents the limited availability of most nutrients from the environment:

$$\vec{F}(\vec{m}) \leq \vec{F}_{ub}(\vec{m}) \tag{2}$$

Where $\vec{F}$ is a vector of fluxes between the environment and the bacterial population (in mol per time unit per population unit) and $\vec{m}$ is a vector of all metabolites.

$\vec{F}_{ub}(\vec{m})$ is set dynamically at each timestep t by the spatial environment (discussed in detail below) at each lattice site x:

$$\vec{F}_{ub}(\vec{x}, \vec{m}, t) = \frac{\vec{c}(\vec{x}, \vec{m}, t)}{B(\vec{x}, t)} \tag{3}$$

Where $\vec{c}$ is a vector of all metabolite concentrations in mol per lattice site, $\vec{x}$ is a vector of all lattice sites and $B(\vec{x})$ is the size of the local bacterial population in population units. Population units are continuous, the size of B can range from $5 \cdot 10^7$ to $2 \cdot 10^{10}$, or $4 \cdot 10^{10}$ when division is not possible.

There is an additional constraint on the total flux. This constraint represents the limited amount of metabolism that can be performed per cell in each timestep. Each cell can only contain a limited number of enzymes, and each enzyme can only perform a limited number of reactions in limited time. Not limiting flux would lead to an unrealistic situation. The enzymatic constraint uses the sum of all fluxes $\vec{f}$, not just those that exchange with the environment:

$$\sum \vec{f} \leq a \tag{4}$$

The enzymatic constraint a is in mol per time unit per population unit (Table 2). As both $\vec{f}$ and a are per population unit, this limit scales with population size, allowing each bacterium to contribute equally and independently to the metabolic flux attained in a lattice site. The enzymatic constraint is included as a constraint on each local FBA solution separately at each timestep. The enzymatic constraint was first proposed in a metabolic modelling context in 1990 (20), adapted from the study of other capacitated flow networks. In our context the enzymatic constraint represents the limited availability of physical space for enzymes. Our method is similar to FBA with molecular crowding in that we use an additional constraint to model metabolic capacity (21). However, we do not utilize the necessarily reaction-specific crowding coefficients.

FBA then identifies the solution space that adheres to these constraints and from this spaces identifies the solution that optimizes the objective function. The set of solutions consists of a set of exchange fluxes $\vec{F}(\vec{x}, \vec{m}, t)$, and a growth rate $\vec{g}(\vec{x}, t)$. These exchange fluxes are taken as the derivatives of a set of partial-differential equations to model the transport of intermediary metabolites (see below). The size of the population increases in proportion to the growth rate produced by the solution.

Environmental metabolites We modeled a set of 723 extracellular metabolites - the union of all metabolites that can be exchanged with the environment by at least one GEM in the model. In practice, only a fraction of the metabolites occur in any exchange reaction that has flux over it, and only 17 metabolites are ever present in the medium in more than micromolar amounts in our simulations outside of the sensitivity analysis (S1 Table). Though the model distinguishes between L-lactate and D-lactate, we display them together in our figures. Nearly all lactate produced and consumed in our model is L-lactate.

To mimic advection, the complete metabolic contents of the lattice except oxygen are moved towards the distal end by 2mm (usually one lattice site) per timestep, i.e. once every 3 simulated minutes. This leads to an average transit time of approximately 11 hours, in agreement with the observed cecum to rectum transit time in newborns (32, 33). Metabolites moving out of the distal end are removed entirely and analysed separately (see Section 'Analysis' for detail).

Every timestep all metabolites diffuse to the four nearest neighbours $NB(\vec{x})$ at an equal rate for all metabolites (Table 2). We used a baseline of $4.4 \cdot 10^{-5}$ to represent mixing of metabolites, which is an order of magnitude higher than normal diffusion

712 for common metabolites (83) to represent active mixing due to colonic contractions.
713 Metabolites are also added and removed by bacterial populations as a result of the
714 FBA solutions, yielding

$$\frac{d\vec{c}(\vec{x}, \vec{m}, t)}{dt} = \vec{F}(\vec{x}, \vec{m}, t)B(\vec{x}, t) + \frac{D}{L^2} \sum_{\vec{i} \in NB(\vec{x})} \left(\vec{c}(\vec{i}, \vec{m}, t) - \vec{c}(\vec{x}, \vec{m}, t) \right). \tag{5}$$

All lattice sites initially contain water as their only metabolite, except for the con-
ditions where oxygen is added.

Metabolites representing the food intake are inserted into the first six columns
of lattice sites every three hours (60 timesteps) to approximate a realistic interval for
neonates (84). It consists solely of lactose, in a concentration in line with human milk
(85), assuming 98% host uptake of carbohydrates before reaching the colon, a com-
monly used assumption (16). Water is provided as a metabolite in unlimited quantities.
Oxygen is placed in some simulations. No other metabolites are available, other than
those produced as a result of bacterial metabolism within the model.

Population dynamics Each local population solves the FBA problem based on its
own GEM, a set enzymatic constraint a, its current population size $B(\vec{x}, t)$ and the local
concentrations of metabolites $\vec{c}(\vec{m}, t)$ at each timestep, and applies the outcome to
the environment (see above) and its own population size, as follows:

$$\frac{d\vec{B}(\vec{x}, t)}{dt} = \vec{B}(\vec{x}, t) \cdot \vec{g}(\vec{x}, t) \tag{6}$$

Populations at least 200 times the initial size (Table 2) will create a new population
in one empty adjacent lattice site, if possible. Half of the old population size is trans-
ferred to the new population, in such a way that the total size is preserved. To mimic
colonization events new populations are introduced at random into empty lattice sites
during the simulation, representing new populations that emerge from sources such
as ingestion or emergence from crypts acting as reservoirs (34). Each empty lattice has
a probability of 0.00005 (Table 2) each step to acquire a new population of a randomly
selected species. This is scaled along with the total number of lattice sites for lattices
of different dimensions. There is an equal probability for any species in the model to
be selected. As we consider these new populations to be new colonizers we initialize
them at the same population size B as the initial populations in the model (Table 2).
Each population dies out at a probability of 0.0075 per timestep, creating a turnover
within the range of estimated microbial turnover rates in the mouse microbiota (86).

To mimic the spread of and mixing of bacterial populations, the lattice sites swap
population contents each timestep. At the default lattice size, one swap is attempted
per lattice site, in a random order, with a random direction in either dimension. This
can cause the lattice site to pick itself. The swap is performed if the origin and destina-
tion lattice site have not been previously been swapped this timestep. The swap only
affects any present population or populations, not the metabolites. Any populations
on the most distal column, i.e. at the exit of the colon, are removed from the system.
To maintain the same diffusion in lattices with different dimensions than the default,
the number of swaps is limited or the swap algorithm is performed multiple times.

This approach to diffusion allows us to diffuse our population units without break-
ing them into very small units. Having many even smaller units would require the pres-
ence of multiple populations in the same space and greatly increase computation time,
because each FBA solution is calculated per population. The mixing method we use for
populations leads to a diffusion constant of $5.7 \cdot 10^{-5}$ cm^2/s (Table 2), as derived from
the mean square displacement calculated over two steps from a point source in the

center of a 20 by 20 field. The diffusion constant for bacterial populations is 1.3 times higher than that of the metabolites, which diffuse in a continuous manner. We have maintained the diffusion rates at these two slightly different levels. Decreasing the diffusion of bacterial populations would no longer allow some bacterial populations to even attempt movement in a timestep. Increasing metabolic diffusion to exactly match bacterial diffusion would necessitate performing our diffusion algorithm twice to prevent instability, which would also cause a portion of metabolites to move further than any population can move in a single step, and increase computational load in all simulations.

The swapping procedure we use is based on Kawasaki-Ising dynamics originally used for modelling the diffusion of spins in ferromagnets ([87]). Methods based on Kawasaki-Ising dynamics have been used to model the mixing of bacteria ([88]). Exchange of neighbours like in Kawasaki-Ising dynamics were previously used to model the mixing of bacterial populations in the model by van Hoek and Merks ([18]).

Initial conditions The simulation is initialized by placing a number of very small populations (B) of the various species randomly across different lattice sites of the environment (Table [2]). There is a probability of 0.3 for each lattice site to acquire a population, an average of 540 for our lattice. The size of initial populations is scaled to be roughly equivalent to a plausible initial total load of approximately $3 \cdot 10^{10}$([89]), assuming a total colon volume of approximately 90 ml. As there is little information on the relative abundance of species in the very early infant gut, we place all species at equal probability. In initially oxygenated conditions, oxygen is also placed as a metabolite now. Water is always considered to be present everywhere. No other metabolites are initially present except for the first feeding.

The ratio of population to bacterial abundance was chosen as a balance between a high bacterial resolution and computational complexity, as the FBA solution of each population is calculated independently per timestep. This resolution splits the infant microbiota into approximately 1-600 units (depending on conditions), allowing for the presence of species at relatively low abundances and spatial variation. There is no hard limit on the number of populations in the model - it is limited only by the balance between growth (through consumption) and death.

Analysis At each timestep we record the location and exchange fluxes $\vec{F}(\vec{x}, t)$ of major metabolites as well as the size $B(\vec{x}, t)$ and species of all populations. This is used to analyze both population composition and metabolic fluxes over time and space. In addition, each timestep we record the location and quantity $\vec{c}(\vec{x}, t)$ of all metabolites present in more than micromolar quantities. We also record this metabolite data for all metabolites exiting the system at the distal side. This is considered the closest equivalent to a fecal composition in the model, and these results are compared to data from *in vivo* fecal samples.

To detect any irregularities, we also record the net carbon, hydrogen, and oxygen flux of every population and the system as a whole. The difference in Gibbs free energy per timestep is also recorded per for each population per FBA solution, and seperately over the whole system. Estimated Gibbs free energy is derived from the Equillibrator database ([74]). Energy loss I in joules per timestep per population unit is recorded as follows, where m are metabolites, F is the exchange fluxes in mol per population unit, B is the population size and E contains the Gibbs free energy in joules per mol for each metabolite,

$$I = \sum_m F(m) \cdot E(m) \qquad (7)$$

For specific simulations, reactions are removed from the models. This is per-
formed by deleting some reactions from the GEM before the conversion to the sto-
ichiometric matrix. To remove Fructose-6-phosphate phosphoketolase, we removed
the reactions R_PKL and R_F6PE4PL. For other simulations, the uptake of certain metabo-
lites is disabled. This is done by placing the upper flux bound $\vec{F}_{ub}$ of the relevant ex-
change reaction at 0 for the relevant populations.

Parameters Relevant parameters are listed in Table 2. Based on measurements
of the typical length and diameter of the infant colon (26, 27) we estimated a volume
of 90 ml. Combined with the average abundance per ml of around 10^{10} after the first
days (89), this leads to a very rough estimate of 10^{12} bacteria in the young infant colon.
To remain computationally feasible, while still modelling at a high resolution, we divide
this population into units of at most $1 \cdot 10^{10}$. Local populations at or above this limit will
divide into two equally sized populations when space is available. This prevents the
local population from becoming unrealistically high. Local populations of more than
$2 \cdot 10^{10}$, which can only form if no space is available to divide for a longer time period,
cease metabolism.

TABLE 2 Default parameters of the model

Parameter	Value	Unit
Timestep	180	seconds
Lattice site side length	2	millimetre
Width of the lattice	225	lattice sites
Height of the lattice	8	lattice sites
Colon transit time	11	hours
Average number of initial populations	540	-
Initial size per population	$5 \cdot 10^{7}$	number of bacteria
Population size required for division	$1 \cdot 10^{10}$	number of bacteria
Death probability	0.0075	per timestep per population
Enzymatic constraint	2	µmol flux per timestep per $1 \cdot 10^{10}$ population
New species placement probability	0.00005	per timestep per empty lattice site
Diffusion of metabolites	$4.4 \cdot 10^{-5}$	cm^2/s
Diffusion of populations	$5.7 \cdot 10^{-5}$	cm^2/s
Growth per µmol ATP	$1 \cdot 10^{9}$	number of bacteria
Lactose input	211	µmol per three hours

The lactose input is estimated from the known intake of milk, its lactose concen-
tration, and an estimate of pre-colonic lactose absorption of 98% (90, 16). Little data
is available on the growth rate of bacteria within the human colon. Growth rates are
expected to be much lower than those found in *in vitro* cultures of individual species
(91). In absence of precise data for infants, here we use a death probability that places
the replacement rate within the range of estimated doubling times of the whole gut
microbiota in mice (86). The colonic transit time is based on data for total transit time
gathered with carmine red dye (32), adjusted for the mouth to cecum transit time (33).
The timestep interval was set at three minutes, to be able to capture individual feed-
ings at a high resolution. Other parameters were determined experimentally to reach
plausible outcomes on a metabolic and species level while maintaining computational
feasibility.

Implementation The model was implemented in C++11. This code was based on
one used earlier to model the gut microbiota (18). Random numbers are generated
with Knuth's subtractive random number generator algorithm. The GEMs are loaded
using libSBML 5.18.0 for C++. We use the GNU Linear Programming Kit 4.65 (GLPK) as a
linear programming tool to solve each FBA with enzymatic constraint problem. We use

the May 2019 update of AGORA, the latest at time of writing, from the Virtual Metabolic
Human Project website (vmh.life). Python 3.6 was used to extract thermodynamic data
extracted from the eQuilibrator API (December 2018 update) (74) and determine mean
square displacement of our bacterial diffusion. Model screenshots are made using
the libpng16 and pngwriter libraries. Other visualizations and statistical analysis were
performed with R 4.1.2 and Google Sheets.

SUPPLEMENTAL MATERIAL

S1 Table. List of present metabolites.csv
A table of the extracellular metabolites present in the system in more than micromolar
amounts with common names and BiGG ID as a .csv file.

S2 Table. Table of changed or deleted reactions and annotations.csv
A table of changes made to the AGORA models as a .csv file.

S1 Video. Video of a run without initially oxygenated conditions

S2 Video. Visualisation of the run from S1 Video, displaying population and
metabolite pool sizes, and fluxes between populations and metabolites

S3 Video. Visualisation of a run from Fig. 4 A, followed by a run from Fig. 4
B, and a run from Fig. 4 F, displaying population and metabolite pool sizes, and
fluxes between populations and metabolites

S4 Video. Video of a run with an initial 0.1 µmol of oxygen per lattice site

S5 Video. Visualisation of the run from S4 Video, displaying population and
metabolite pool sizes, and fluxes between populations and metabolites

S6 Video. Visualisation of a run from Fig. 5 D, followed by a run from 5 H, dis-
playing population and metabolite pool sizes, and fluxes between populations
and metabolites

S7 Video. Visualisation of a run from Fig. 6 A without the enzymatic con-
straint, displaying population and metabolite pool sizes, and fluxes between
populations and metabolites

S1 Figure. Histogram displaying the distribution of change in Gibbs free en-
ergy for each non-zero FBA solution over 30 simulations of the model as dis-
played in Fig. 3A

S2 Figure. (A) Growth per timestep by lactose concentration for populations
$5 \cdot 10^9$ **with access to one lattice site (0.05ml) of some major bacterial species with**
the enzymatic constraint disabled (B)Production of metabolites per timestep by
a *E. coli* population of $5 \cdot 10^9$ **with access to one lattice site (0.05ml) (C) Produc-**
tion of metabolites by *B. vulgatus* population of $5 \cdot 10^9$ **with access to one lattice**
site (0.05ml). Formate and propionate formation are always equal, and so are
hydrogen and succinate. (D)Growth per timestep by lactose concentration for
populations $5 \cdot 10^9$ **with access to one lattice site (0.05ml) of some major bacterial**
species with the enzymatic constraint disabled

S3 Figure. Populations over time with bacteria in the "others" category of
fig. 5 left out. Abundances for grouped species over 21 days. One day is 480
timesteps. Curve shows mean value and shaded area one standard deviation
over n=30 simulations.

S4 Figure. (A,B) Spatial distribution of (A) lactose and (B) acetate over the last
two days (960 timesteps) of the simulation. Curve shows mean value and shaded
area one standard deviation over n=30 simulations.

S5 Figure. (A) Populations over time with lactate uptake disabled for *Bifi-*
***dobacterium* species. Abundances for grouped species over 21 days. One day**
is 480 timesteps. Curve shows mean value and shaded area one standard devi-

ation over n=30 simulations. **(B) Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation in (A). n=30.**

S6 Figure. Populations over time with lactate production disabled for *Bifidobacterium* species. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

S7 Figure. (A)Production of metabolites by lactose concentration in the presence of abundant oxygen for an *E. coli* population of $5 \cdot 10^9$ with access to one lattice site (0.05ml) (B) Production of metabolites by lactose concentration in the presence of abundant oxygen for an *E. cloacea* population of $5 \cdot 10^9$ with access to one lattice site (0.05ml) (C) Production of metabolites by glucose concentration in the presence of abundant oxygen for an *E. coli* population of $5 \cdot 10^9$ with access to one lattice site (0.05ml)

S8 Figure. Populations over time with (A) 0.001 (B) 0.01 (C) 0.1 (D) 1 μmol oxygen released in total per step divided over the top and bottom rows. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations per condition.

S9 Figure. (A) Populations over time with 1 μmol oxygen per step initially (as in S8 Figure D), decreasing to 0 at step 5040. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations per condition. (B) Distribution of total *Bifidobacterium* abundance at 21 days (10080 timesteps) corresponding to the simulations of (A)

S10 Figure. Populations over time with new populations after the initial populations only placed in the first column, with the placement probability (Table 2) increased to 0.01125. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

S11 Figure. (A,B,C) Spatial distribution of lactose, lactate, and acetate, respectively, over the last two days (960 timesteps) of the simulation with a smaller lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (D) Abundance per location for grouped species over the last two days (960 timesteps) of the simulations with a smaller lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (E,F,G) Spatial distribution of lactose, lactate, and acetate, respectively, over the last two days (960 timesteps) of the simulation with a larger lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (H) Abundance per location for grouped species over the last two days (960 timesteps) of the simulations with a larger lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

S12 Figure. Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation with diffusion of metabolites increased by a factor of 5. n=30.

CONTRIBUTIONS

P.V.J, M.P., J.M.W.G., and R.M.H.M acquired funding. D.M.V., P.V.J, M.P., J.M.W.G., and R.M.H.M. conceived and planned the simulations. D.M.V. and D.M. wrote software used for the simulations. D.M.V. performed the simulations and analyzed the data. R.S, E.L., J.M.W.G., and R.M.H.M contributed to the interpretation of the results. J.M.W.G.,

and R.M.H.M. supervised the project. D.M.V. drafted the manuscript. D.M.V., R.S., E.L.,
J.M.W.G. and R.M.H.M. revised and edited the manuscript.

ACKNOWLEDGMENTS
This study was financially supported by FrieslandCampina. R.S, E.L., P.V.J., M.P. and
J.M.W.G. are currently or were previously employed by FrieslandCampina. The work
was carried out in part on the Dutch national e-infrastructure with the support of SURF
Cooperative. This work was performed in part using the ALICE compute resources
provided by Leiden University.

REFERENCES

1. **Dogra S, Sakwinska O, Soh SE, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Karnani N, Lee YS, Yap F, Chong YS, Godfrey KM, Holbrook JD**. 2015. Rate of establishing the gut microbiota in infancy has consequences for future health. Gut Microbes 6 (5):321–325. doi: 10.1080/19490976.2015.1078051.

2. **Dogra S, Sakwinska O, Soh SEE, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Lee YS, Yap F, Chong YSS, Godfrey KM, Holbrook JD, GUSTO Study Group, Sakwinska O, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Soh SEE, Yap F, Godfrey KM, Holbrook JD**. 2015. Dynamics of infant gut microbiota are influenced by delivery mode and gestational duration and are associated with subsequent adiposity. mBio 6 (1):1–9. doi:10.1128/mBio.02419-14.

3. **Scholtens PAMJ, Oozeer R, Martin R, Amor KB, Knol J**. 2012. The Early Settlers: Intestinal Microbiology in Early Life. Annu Rev Food Sci Technol 3 (1):425–447. doi:10.1146/annurev-food-022811-101120.

4. **Stark PL, Lee A**. 1982. The microbial ecology of the large bowel of breast-fed and formula-fed infants during the first year of life. J Med Microbiol 15 (2):189–203. doi:10.1099/00222615-15-2-189.

5. **Matamoros S, Gras-Leguen C, Le Vacon F, Potel G, De La Cochetiere MF**. 2013. Development of intestinal microbiota in infants and its impact on health. Trends Microbiol 21 (4):167–173. doi: 10.1016/j.tim.2012.12.001.

6. **Dotinga BM, Mintzer JP, Moore JE, Hulscher JBF, Bos AF, Kooi EMW**. 2020. Maturation of Intestinal Oxygenation: A Review of Mechanisms and Clinical Implications for Preterm Neonates. Front Pediatr 8 (July). doi:10.3389/fped.2020.00354.

7. **Friedman ES, Bittinger K, Esipova TV, Hou L, Chau L, Jiang J, Mesaros C, Lund PJ, Liang X, FitzGerald GA, Goulian M, Lee D, Garcia BA, Blair IA, Vinogradov SA, Wu GD**. 2018. Microbes vs. chemistry in the origin of the anaerobic gut lumen. Proc Natl Acad Sci United States Am 115 (16):4170–4175. doi: 10.1073/pnas.1718635115.

8. **Lay C, Chu CW, Purbojati RW, Acerbi E, Drautz-Moses DI, de Sessions PF, Jie S, Ho E, Kok YJ, Bi X, Chen S, Mak SY, Chua MC, Goh AEN, Chiang WC, Rao R, Chaithongwongwatthana S, Khemapech N, Chongsrisawat V, Martin R, Koh Y, Lohar SR, Tan ICH, Anng WA, Jie C, Bartke N, Ben-Amor K, Renes IB, Wong F, Roeselers G, Ho YS, Hibberd ML, Schuster SC, Knol J**. 2021. A synbiotic intervention modulates meta-omics signatures of gut redox potential and acidity in elective caesarean born infants. BMC Microbiol 21 (1):1–17. doi: 10.1186/s12866-021-02230-1.

9. **Johnson JM, Adams ED**. mar 2022. The Gastrointestinal Microbiome in Infant Colic. MCN: The Am J Matern Nurs Publish Ah:1–12. doi: 10.1097/NMC.0000000000000832.

10. **Duar RM, Kyle D, Casaburi G**. 2020. Colonization Resistance in the Infant Gut: The Role of B. infantis in Reducing pH and Preventing Pathogen Growth. High-Throughput 9 (2):7. doi:10.3390/ht9020007.

11. **Bäckhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, Li Y, Xia Y, Xie H, Zhong H, Khan MT, Zhang J, Li J, Xiao L, Al-Aama J, Zhang D, Lee YS, Kotowska D, Colding C, Tremaroli V, Yin Y, Bergman S, Xu X, Madsen L, Kristiansen K, Dahlgren J, Jun W**. 2015. Dynamics and stabilization of the human gut microbiome during the first year of life. Cell Host Microbe 17 (5):690–703. doi: 10.1016/j.chom.2015.04.004.

12. **van Best N, Hornef MW, Savelkoul PHM, Penders J**. 2015. On the origin of species: Factors shaping the establishment of infant's gut microbiota. Birth Defects Res Part C - Embryo Today: Rev 105 (4):240–251. doi:10.1002/bdrc.21113.

13. **Berger B, Porta N, Foata F, Grathwohl D, Delley M, Moine D, Charpagne A, Siegwald L, Descombes P, Alliet P, Puccio G, Steenhout P, Mercenier A, Sprenger N**. 2020. Linking human milk oligosaccharides, infant fecal community types, and later risk to require antibiotics. mBio 11 (2):1–18. doi:10.1128/mBio.03196-19.

14. **Francavilla R, Calasso M, Calace L, Siragusa S, Ndagijimana M, Vernocchi P, Brunetti L, Mancino G, Tedeschi G, Guerzoni E, Indrio F, Laghi L, Miniello VL, Gobbetti M, De Angelis M**. 2012. Effect of lactose on gut microbiota and metabolome of infants with cow's milk allergy. Pediatr Allergy Immunol 23 (5):420–427. doi:10.1111/j.1399-3038.2012.01286.x.

15. **Magnúsdóttir S, Thiele I**. 2018. Modeling metabolism of the human gut microbiome. Curr Opin Biotechnol 51:90–96. doi: 10.1016/j.copbio.2017.12.005.

16. **Chan SHJ, Simons MN, Maranas CD**. 2017. SteadyCom: Predicting microbial abundances while ensuring community stability. PLoS Comput Biol 13 (5):1–25. doi:10.1371/journal.pcbi.1005539.

17. **Bauer E, Zimmermann J, Baldini F, Thiele I, Kaleta C**. 2017. BacArena: Individual-based metabolic modeling of heterogeneous microbes in complex communities. PLoS Comput Biol 13 (5):1–22. doi:10.1371/journal.pcbi.1005544.

18. **Hoek MJAv, Merks RMH**. 2017. Emergence of microbial diversity due to cross-feeding interactions in a spatial model of gut microbial metabolism. BMC Syst Biol 11 (1):1–18. doi:10.1186/s12918-017-0430-4.

19. **Dukovski I, Bajić D, Chacón JM, Quintin M, Vila JCC, Sulheim S, Pacheco AR, Bernstein DB, Riehl WJ, Korolev KS, Sanchez A, Harcombe WR, Segrè D**. 2021. A metabolic modeling platform for the computation of microbial ecosystems in time and space (COMETS). Nat Protoc doi:10.1038/s41596-021-00593-3.

20. **Majewski RA, Domach MM**. 1990. Simple constrained-optimization view of acetate overflow in E. coli. Biotechnol Bioeng 35 (7):732–738. doi:10.1002/bit.260350711.

21. **Beg QK, Vazquez A, Ernst J, De Menezes MA, Bar-Joseph Z, Barabási AL, Oltvai ZN**. 2007. Intracellular crowding defines the mode and sequence of substrate uptake by Escherichia coli and constrains its metabolic activity. Proc Natl Acad Sci United States

Am 104 (31):12663–12668. doi:10.1073/pnas.0609845104.

22. **de Groot DH, Lischke J, Muolo R, Planqué R, Bruggeman FJ, Teusink B**. 2020. The common message of constraint-based optimization approaches: overflow metabolism is caused by two growth-limiting constraints. Cell Mol Life Sci 77 (3):441–453. doi: 10.1007/s00018-019-03380-2.

23. **Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, Greenhalgh K, Jäger C, Baginska J, Wilmes P, Fleming RMT, Thiele I**. 2017. Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota. Nat Biotechnol 35 (1):81–89. doi:10.1038/nbt.3703.

24. **Medina DA, Pinto F, Ovalle A, Thomson P, Garrido D**. 2017. Prebiotics mediate microbial interactions in a consortium of the infant gut microbiome. Int J Mol Sci 18 (10):1–16. doi:10.3390/ijms18102095.

25. **Monteagudo-Mera A, Arthur JC, Jobin C, Keku T, Bruno-Barcena JM, Azcarate-Peril MA**. 2016. High purity galacto-oligosaccharides enhance specific Bifidobacterium species and their metabolic activity in the mouse gut microbiome. Benef Microbes 7 (2):247–264. doi: 10.3920/BM2015.0114.

26. **Valentin J, Streffer C**. 2002. Basic anatomical and physiological data for use in radiological protection: Reference values - ICRP Publication 89. Ann ICRP 32 (3-4):1–277. doi:10.1016/S0146-6453(03)00002-2.

27. **Davis WS, Parker Allen R, Favara BE**. 1974. Neonatal small left colon syndrome. Am journal Roentgenol 120 (2):322–329.

28. **Swidsinski A, Loening-Baucke V, Verstraelen H, Osowska S, Doerffel Y**. 2008. Biostructure of Fecal Microbiota in Healthy Subjects and Patients With Chronic Idiopathic Diarrhea. Gastroenterology 135 (2):568–579. doi:10.1053/j.gastro.2008.04.017.

29. **Hong PY, Croix JA, Greenberg E, Gaskins HR, Mackie RI**. 2011. Pyrosequencing-based analysis of the mucosal microbiota in healthy individuals reveals ubiquitous bacterial groups and microheterogeneity. PLoS ONE 6 (9). doi:10.1371/journal.pone.0025042.

30. **Delsing D, Lindner C, van Dijck H, Looijesteijn E, Lima ES, Heerikhuizen M, Agamennone V, Triantis V, Bovee-Oudenhoven I**. 2019. Gut microbiota modulating effects of GOS and 2'-FL studied by in vitro infant and toddler faecal fermentations. J Pediatr Gastroenterol Nutr 68 (Supplement 1):1037. http://ovidsp.ovid.com/ovidweb.cgi?T=JS{&}PAGE=reference{&}D=emexa{&}NEWS=N{&}AN=628093791.

31. **Schuetz R, Kuepfer L, Sauer U**. 2007. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. Mol Syst Biol 3 (119). doi:10.1038/msb4100162.

32. **Rubaltelli FF, Largajolli G**. 1973. Effect of Light Exposure on Gut Transit Time in Jaundiced Newborns. Acta Pædiatrica 62 (2):146–148. doi: 10.1111/j.1651-2227.1973.tb08082.x.

33. **Hyams JS, Geertsma MA, Etienne NL, Treem WR**. 1989. Colonic hydrogen production in infants with colic. The J Pediatr 115 (4):592–594. doi:10.1016/S0022-3476(89)80289-6.

34. **Lee SM, Donaldson GP, Mikulski Z, Boyajian S, Ley K, Mazmanian SK**. 2013. Bacterial colonization factors control specificity and stability of the gut microbiota. Nature 501 (7467):426–429. doi: 10.1038/nature12447.

35. **van Hoek MJA, Merks RMH**. 2012. Redox balance is key to explaining full vs. partial switching to low-yield metabolism. BMC Syst Biol 6. doi:10.1186/1752-0509-6-22.

36. **Vazquez A, Oltvai ZN**. 2016. Macromolecular crowding explains overflow metabolism in cells. Sci Reports 6:1–7. doi:10.1038/srep31007.

37. **de Vries W, Stouthamer AH**. 1968. Fermentation of glucose, lactose, galactose, mannitol, and xylose by bifidobacteria. J Bacteriol 96 (2):472–478.

38. **De Vuyst L, Moens F, Selak M, Rivière A, Leroy F**. 2014. Summer

Meeting 2013: Growth and physiology of bifidobacteria. J Appl Microbiol 116 (3):477–491. doi:10.1111/jam.12415.

39. **Kotarski SF, Salyers AA**. 1981. Effect of long generation times on growth of Bacteroides thetaiotaomicron in carbohydrate-limited continuous culture. J Bacteriol 146 (3):853–860. doi: 10.1128/jb.146.3.853-860.1981.

40. **Martin FPJ, Moco S, Montoliu I, Collino S, Da Silva L, Rezzi S, Prieto R, Kussmann M, Inostroza J, Steenhout P**. 2014. Impact of breast-feeding and high-and low-protein formula on the metabolism and growth of infants from overweight and obese mothers. Pediatr Res 75 (4):535–543. doi:10.1038/pr.2013.250.

41. **Tsukuda N, Yahagi K, Hara T, Watanabe Y, Matsumoto H, Mori H, Higashi K, Tsuji H, Matsumoto S, Kurokawa K, Matsuki T**. 2021. Key bacterial taxa and metabolic pathways affecting gut short-chain fatty acid profiles in early life. ISME J doi:10.1038/s41396-021-00937-7.

42. **Schwab C, Ruscheweyh HJ, Bunesova V, Pham VT, Beerenwinkel N, Lacroix C**. 2017. Trophic interactions of infant bifidobacteria and eubacterium hallii during L-fucose and fucosyllactose degradation. Front Microbiol 8 (JAN):1–14. doi:10.3389/fmicb.2017.00095.

43. **Wolin MJ, Yerry S, Miller TL, Zhang Y, Bank S**. 1998. Changes in Production of Ethanol, Acids and H2from Glucose by the Fecal Flora of a 16- to 158-d-Old Breast-Fed Infant. The J Nutr 128 (1):85–90. doi: 10.1093/jn/128.1.85.

44. **Bridgman SL, Azad MB, Field CJ, Haqq AM, Becker AB, Mandhane PJ, Subbarao P, Turvey SE, Sears MR, Scott JA, Wishart DS, Kozyrskyj AL**. 2017. Fecal Short-Chain Fatty Acid Variations by Breastfeeding Status in Infants at 4 Months: Differences in Relative versus Absolute Concentrations. Front Nutr 4 (April). doi: 10.3389/fnut.2017.00011.

45. **de Vries W, Stouthamer AH**. 1967. Pathway of glucose fermentation in relation to the taxonomy of bifidobacteria. J bacteriology 93 (2):574–576. doi:10.1128/jb.93.2.574-576.1967.

46. **Kennedy KM, Gerlach MJ, Adam T, Heimesaat MM, Rossi L, Surette MG, Sloboda DM, Braun T**. 2021. Fetal meconium does not have a detectable microbiota before birth. Nat Microbiol 6 (7):865–873. doi:10.1038/s41564-021-00904-0.

47. **Albenberg L, Esipova TV, Judge CP, Bittinger K, Chen J, Laughlin A, Grunberg S, Baldassano RN, Lewis JD, Li H, Thom SR, Bushman FD, Vinogradov SA, Wu GD**. 2014. Correlation between intraluminal oxygen gradient and radial partitioning of intestinal microbiota. Gastroenterology 147 (5):1055–1063.e8. doi: 10.1053/j.gastro.2014.07.020.

48. **Zheng L, Kelly CJ, Colgan SP**. 2015. Physiologic hypoxia and oxygen homeostasis in the healthy intestine. A review in the theme: Cellular responses to hypoxia. Am J Physiol - Cell Physiol 309 (6):C350–C360. doi:10.1152/ajpcell.00191.2015.

49. **Matsuki T, Yahagi K, Mori H, Matsumoto H, Hara T, Tajima S, Ogawa E, Kodama H, Yamamoto K, Yamada T, Matsumoto S, Kurokawa K**. 2016. A key genetic factor for fucosyllactose utilization affects infant gut microbiota development. Nat Commun 7 (May):1–12. doi:10.1038/ncomms11939.

50. **Pham VT, Lacroix C, Braegger CP, Chassard C**. 2016. Early colonization of functional groups of microbes in the infant gut. Environ microbiology 18 (7):2246–2258. doi:10.1111/1462-2920.13316.

51. **Sim K, Shaw AG, Randell P, Cox MJ, McClure ZE, Li MS, Haddad M, Langford PR, Cookson WOCM, Moffatt MF, Kroll JS**. 2015. Dysbiosis anticipating necrotizing enterocolitis in very premature infants. Clin Infect Dis 60 (3):389–397. doi:10.1093/cid/ciu822.

52. **Mori M, Hwa T, Martin OC, De Martino A, Marinari E**. 2016. Constrained Allocation Flux Balance Analysis. PLoS Comput Biol 12 (6):1–24. doi:10.1371/journal.pcbi.1004913.

53. **Balakrishnan R, Silva RT, Hwa T, Cremer J**. 2021. Suboptimal resource allocation in changing environments constrains response and growth in bacteria. Mol Syst Biol 17 (12):1–13. doi: 10.15252/msb.202110597.

54. **Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, Lovley DR**. 2011. Genome-scale dynamic modeling of the competition between Rhodoferax and Geobacter in anoxic subsurface environments. ISME J 5 (2):305–316. doi:10.1038/ismej.2010.117.

55. **Stansbridge EM, Walker V, Hall MA, Smith SL, Millar MR, Bacon C, Chen S**. 1993. Effects of feeding premature infants with Lactobacillus GG on gut fermentation. Arch Dis Child 69 (5 SPEC NO):488–492. doi:10.1136/adc.69.5_Spec_No.488.

56. **Falony G, Lazidou K, Verschaeren A, Weckx S, Maes D, De Vuyst L**. 2009. In vitro kinetic analysis of fermentation of prebiotic inulin-type fructans by Bifidobacterium species reveals four different phenotypes. Appl Environ Microbiol 75 (2):454–461. doi: 10.1128/AEM.01488-08.

57. **Cocks GT, Aguilar J, Lin ECC**. 1974. Evolution of L-1, 2-Propanediol Catabolism in Escherichia coli by Recruitment of Enzymes for L-Fucose and L-Lactate Metabolism.

58. **Pham VT, Lacroix C, Braegger CP, Chassard C**. 2017. Lactate-utilizing community is associated with gut microbiota dysbiosis in colicky infants. Sci Reports 7 (1):1–13. doi:10.1038/s41598-017-11509-1.

59. **Schwab C, Ruscheweyh HJ, Bunesova V, Pham VT, Beerenwinkel N, Lacroix C**. 2017. Trophic interactions of infant bifidobacteria and eubacterium hallii during L-fucose and fucosyllactose degradation. Front Microbiol 8 (JAN):1–14. doi:10.3389/fmicb.2017.00095.

60. **Cremer J, Arnoldini M, Hwa T**. 2017. Effect of water flow and chemical environment on microbiota growth and composition in the human colon. Proc Natl Acad Sci United States Am 114 (25):6438–6443. doi:10.1073/pnas.1619598114.

61. **Cremer J, Segota I, Yang CY, Arnoldini M, Sauls JT, Zhang Z, Gutier-reza E, Groisman A, Hwa T**. 2016. Effect of flow and peristaltic mixing on bacterial growth in a gut-like channel. Proc Natl Acad Sci United States Am 113 (41):11414–11419. doi:10.1073/pnas.1601306113.

62. **Corsetti M, Costa M, Bassotti G, Bharucha AE, Borrelli O, Din-ning P, Di Lorenzo C, Huizinga JD, Jimenez M, Rao S, Spiller R, Spencer NJ, Lentle R, Pannemans J, Thys A, Benninga M, Tack J**. 2019. First translational consensus on terminology and definitions of colonic motility in animals and humans studied by manometric and other techniques. Nat Rev Gastroenterol Hepatol 16 (9):559–579. doi:10.1038/s41575-019-0167-1.

63. **Nurko S**. 2006. Motility of the Colon and Anorectum. NeoReviews 7 (1):e34–e48. doi:10.1542/neo.7-1-e34.

64. **Sarna SK**. 2010. Colonic Motility From Bench Side to Bed-side. Morgan&Claypool Life Sciences, San Rafael(CA). doi: 10.4199/C00020ED1V01Y201011ISP011.

65. **Velde SV, Notebaert A, Meersschaut V, Herregods N, Van Winckel M, Van Biervliet S**. 2013. Colon transit time in healthy children and adolescents. Int J Colorectal Dis 28 (12):1721–1724. doi: 10.1007/s00384-013-1750-5.

66. **Chan, Friedman, Wu, Maranas**. 2019. Predicting the Longitudinally and Radially Varying Microbial Metabolic Modeling. processes .

67. **Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, Bonilla G, Kar A, Leiby N, Mehta P, Marx CJ, Segrè D**. 2014. Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. Cell Reports 7 (4):1104–1115. doi:10.1016/j.celrep.2014.03.070.

68. **Varma A, Palsson BØ**. 1994. Metabolic Flux Balancing: Basic Concepts, Scientific and Practical Use. Biotechnology 12 (October):994–998.

69. **Arnoldini M, Cremer J, Hwa T**. 2018. Bacterial growth, flow, and mixing shape human gut microbiota density and composition. Gut microbes 9 (6):559–566. doi:10.1080/19490976.2018.1448741.

70. **Sagheddu V, Patrone V, Miragoli F, Puglisi E, Morelli L**. 2016. Infant early gut colonization by Lachnospiraceae: High frequency of Ruminococcus gnavus. Front Pediatr 4 (JUN):7–12. doi: 10.3389/fped.2016.00057.

71. **Turroni F, Peano C, Pass DA, Foroni E, Severgnini M, Claesson MJ, Kerr C, Hourihane J, Murray D, Fuligni F, Gueimonde M, Margolles A, de Bellis G, O'Toole PW, van Sinderen D, Marchesi JR, Ventura M**. 2012. Diversity of bifidobacteria within the infant gut microbiota. PLoS ONE 7 (5):20–24. doi:10.1371/journal.pone.0036957.

72. **Rogosa M**. 1964. the Genus Veillonella. I. General Cultural, Ecological, and Biochemical Considerations. J Bacteriol 87:162–170. doi: 10.1128/JB.87.1.162-170.1964.

73. **Flamholz A, Noor E, Bar-Even A, Milo R**. 2012. EQuilibrator - The biochemical thermodynamics calculator. Nucleic Acids Res 40 (D1):770–775. doi:10.1093/nar/gkr874.

74. **Noor E, Haraldsdóttir HS, Milo R, Fleming RMT**. 2013. Consistent Estimation of Gibbs Energy Using Component Contributions. PLoS Comput Biol 9 (7). doi:10.1371/journal.pcbi.1003098.

75. **Magnúsdóttir S, Heinken A, Fleming RMT, Thiele I**. 2018. Reply to "Challenges in modeling the human gut microbiome". Nat Biotechnol 36 (8):686–691. doi:10.1038/nbt.4212.

76. **Schuetz R, Kuepfer L, Sauer U**. 2007. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. Mol Syst Biol 3 (119). doi:10.1038/msb4100162.

77. **Sánchez B, De Los Reyes-Gavilán CG, Margolles A**. 2006. The F1F0-ATPase of Bifidobacterium animalis is involved in bile tolerance. Environ Microbiol 8 (10):1825–1833. doi:10.1111/j.1462-2920.2006.01067.x.

78. **Krzewinski F, Brassart C, Gavini F, Bouquelet S**. 1996. Characterization of the lactose transport system in the strain Bifidobacterium bifidum DSM 20082. Curr Microbiol 32 (6):301–307. doi: 10.1007/s002849900054.

79. **Sela DA, Adeuya A, Kim JH, Whitehead TR, Chapman J, Rokhsar DS, Richardson PM, Lapidus A, German JB, Price NP, Chen F, Lebrilla CB, Mills DA**. 2008. The genome sequence of Bifidobacterium longum subsp. infantis reveals adaptations for milk utilization within the infant microbiome. Proc Natl Acad Sci 105 (48):18964–18969. doi:10.1073/pnas.0809584105.

80. **Lessard C, Cochu A, Lemay JD, Roy D, Vaillancourt K, Frenette M, Moineau S, Vadeboncoeur C**. 2003. Phosphorylation of Streptococcus salivarius Lactose Permease (LacS) by HPr(His~P) and HPr(Ser-P)(His~P) and Effects on Growth. J Bacteriol 185 (23):6764–6772. doi: 10.1128/JB.185.23.6764-6772.2003.

81. **Holdeman LV**. 1976. Emendation of Bacteroidaceae and B utyrivib-rio and Descriptions of Desulfornonas gen. nov. and Ten New Species in the Genera Desulfomonas, Butyrivibrio, Eubacterium, Clostridium, and Ruminococcus. Int J Syst Bacteriol 26 (2):238–252.

82. **Orth JD, Thiele I, Palsson BO**. 2010. What is flux balance analysis? Nat Biotechnol 28 (3):245–248. doi:10.1038/nbt.1614.

83. **Stewart PS**. 2003. Diffusion in biofilms. J Bacteriol 185 (5):1485–1491. doi:10.1128/JB.185.5.1485-1491.2003.

84. **Grummer-Strawn LM, Scanlon KS, Fein SB**. 2008. Infant feeding and feeding transitions during the first year of life. Pediatrics 122 (SUPPL. 2). doi:10.1542/peds.2008-1315d.

85. **Ballard O, Morrow AL**. 2013. Human Milk Composition. Nutrients and Bioactive Factors. Pediatr Clin North Am 60 (1):49–74. doi: 10.1016/j.pcl.2012.10.002.

86. **Gibbons RJ, Kapsimalis B**. 1967. Estimates of the overall rate of

growth of the intestinal microflora of hamsters, guinea pigs, and mice. J Bacteriol 93 (1):510–512.

87. **Kawasaki K**. 1966. Diffusion constants near the critical point for time-dependent ising models. I. Phys Rev 145 (1):224–230. doi: 10.1103/PhysRev.145.224.

88. **Koumakis N, Devailly C, Poon WCK**. jun 2018. Motile bacteria in a critical fluid mixture. Phys Rev E 97 (6). doi: 10.1103/PhysRevE.97.062604.

89. **Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO**. 2007. De-velopment of the human infant intestinal microbiota. PLoS Biol 5 (7):1556–1573. doi:10.1371/journal.pbio.0050177.

90. **Anderson PO, Valdés V**. 2015. Variation of milk intake over time: Clin-ical and pharmacokinetic implications. Breastfeed Med 10 (3):142–144. doi:10.1089/bfm.2014.0170.

91. **Gibson B, Wilson DJ, Feil E, Eyre-Walker A**. 2018. The distribution of bacterial doubling times in the wild. Proc Royal Soc B: Biol Sci 285 (1880). doi:10.1098/rspb.2018.0789.