A multiscale spatiotemporal model including a switch from aerobic to anaerobic metabolism reproduces succession in the early infant gut microbiota

David M. Versluis,[a] Ruud Schoemaker,[b] Ellen Looijesteijn,[b] Daniël Muysken,[d],†
Prescilla V. Jeurink,[b] Marcel Paques,[b] Jan M. W. Geurts,[b] Roeland M. H.
Merks,[a,c,*]

Leiden University, Institute of Biology, Leiden, The Netherlands[a]; FrieslandCampina, Amersfoort, the Netherlands[b]
Leiden University, Mathematical Institute, Leiden, The Netherlands[c]; Centrum Wiskunde & Informatica (CWI),
Amsterdam, The Netherlands[d];

ABSTRACT   The human intestinal microbiota starts to form immediately after birth, and is important for the health of the host. During the first days facultatively anaerobic bacterial species generally dominate, such as Enterobacteriaceae. These are succeeded by strictly anaerobic species, particularly *Bifidobacterium* species. An early transition to *Bifidobacterium* species is associated with health benefits: for example, *Bifidobacterium* species repress growth of pathogenic competitors and modulate the immune response. Succession to *Bifidobacterium* is thought to be due to consumption of intracolonic oxygen present in newborns by facultative anaerobes, including Enterobacteriaceae. To study if oxygen depletion suffices for the transition to *Bifidobacterium* species, here we introduce a multiscale mathematical model that considers metabolism, spatial bacterial population dynamics and cross-feeding. Using publicly available metabolic network data from the AGORA collection, the model simulates *ab initio* the competition of strictly and facultatively anaerobic species in a gut-like environment under the influence of lactose and oxygen. The model predicts that individual differences in intracolonic oxygen in newborn infants can explain the observed individual variation in succession to anaerobic species, in particular *Bifidobacterium* species. *Bifidobacterium* species become dominant in the model by using the bifid shunt, which allows *Bifidobacterium* to switch to sub-optimal yield metabolism with fast growth at high lactose concentrations as predicted here using flux-balance analysis. The computational model thus allows us to test the internal plausibility of hypotheses for bacterial colonization and succession in the infant colon.

IMPORTANCE   The composition of the infant microbiota has a great impact on infant health, but its controlling factors are still incompletely understood. The frequently dominant anaerobic *Bifidobacterium* species benefit health, e.g., because they can keep harmful competitors under control and modulate the intestinal immune response. Controlling factors could include nutritional composition and intestinal mucus composition, as well as environmental factors, such as antibiotics. We introduce a modeling framework of a metabolically realistic intestinal microbial ecology in which hypothetical scenarios can be tested and compared. We present simulations that suggest that greater levels of intra-intestinal oxygenation more strongly delay the dominance of

Bifidobacterium species, explaining the observed variety of microbial composition and
demonstrating the use of the model for hypothesis generation. The framework al-
lows us to test a variety of controlling factors, including intestinal mixing and transit
time. Future versions will also include detailed modeling of oligosaccharide and mucin
metabolism.

KEYWORDS: Infant microbiota, Microbial ecology, Flux balance analysis.

INTRODUCTION
The human infant microbiota starts forming directly after birth, and differs greatly
from the adult microbiota ([1]). Three main community types are observed in the infant:
(a) a Proteobacteria-dominated microbiota; (b) an Actinobacteria-dominated micro-
biota; and, less frequently (12-14% in one study ([2])), (c) a Bacilli-dominated microbiota
([1]). The three main community types are established after an ecological succession of
early communities. This ecological succession is thought to be controlled by nutrition
and early oxygen in the colon. To develop hypotheses on the potential mechanisms
and controlling factors of the initial development of the human microbiota, in particu-
lar the role of early oxygen, we introduce a computational model.
In the first 24 to 48 hours after birth, Proteobacteria, including *Escherichia coli* and
Enterobacter cloacae, and Bacilli, including *Streptococcus*, *Lactobacillus*, and *Staphylo-*
coccus, are the most common ([3], [1]). In the following weeks Proteobacteria are often
replaced by anaerobic Actinobacteria, mainly *Bifidobacterium* species, whereas Bacilli
are succeeded by either Proteobacteria or Actinobacteria. Other anaerobic species
are also found, but typically do not dominate. Actinobacteria generally persist as the
dominant group until weaning ([1]). A possible trigger for the replacement of the Enter-
obacteriacea by *Bifidobacterium* spp., is depletion of a hypothesized initial amount of
oxygen ([4], [5]). Oxygen diffuses into the gut lumen from the body and is taken up by
bacteria, by colonocytes, and by non-biological chemical processes in the cecal con-
tents ([6], [7], [8]) leading to oxygen depletion. The relative importance of these processes
is still under debate ([7], [8]).
Bifidobacterium species ([1]) are associated with a range of health benefits. For ex-
ample, early succession to a *Bifidobacterium* dominated microbiota is correlated with
reduced probabilities to be underweight at 18 months of age or to experience colic
([2], [9]). Acetate and lactate produced by *Bifidobacterium* also contribute to the acidifica-
tion of the infant gut, which suppresses many pathogens ([10]). A key question, there-
fore, is if and how *Bifidobacterium* can be promoted. The composition and develop-
ment of the infant microbiota is thought to be determined by many factors, including
nutrition ([11], [12], [13]), i.e. human milk or infant formula. The presence of lactose, the
most abundant carbohydrate in human milk and most infant formulas, is essential for
acquiring high levels of *Bifidobacterium* ([14]). This suggests that nutrition can play a
crucial role in promoting *Bifidobacterium*.
To obtain more insight into the potential ecological and metabolic mechanisms
underlying bacterial colonization and succession in the infant colon, here we propose
a multiscale, spatial computational model of the infant gut microbiota. Earlier work
has introduced similar models of the adult gut microbiota ([15], [16], [17], [18]). These
make use of flux balance analysis (FBA) to model metabolism. Given a number of
constraints including substrate availability and enzyme availability, FBA uses genome-
scale metabolic network models (GEMs) to predict optimal biomass or energy produc-
tion and the exchange of metabolites with the environment. In spatial FBA, FBA models

representing small subpopulations of bacteria are coupled together to model a micro-
bial, metabolizing ecosystem. While some approaches assume the whole system is
in steady-state, allowing them to use the optimization approach for the whole ecosys-
tem ([16]), here we build upon dynamic approaches such as BacArena ([17]), COMETS
([19]), and the approach by Van Hoek and Merks ([18]). In these models, the metabolite
exchange rates given by FBA are used to dynamically change metabolite concentra-
tions in the environment, and predict the resulting effects on the bacterial populations.
These properties make these dynamic, spatial FBA approaches highly suitable for our
purpose of modeling the formation of the infant gut microbiota.

To analyze succession of bacterial species in a dynamic ecosystem such as the in-
fant gut, it is key to accurately describe the dependency of bacterial growth rate and
metabolite exchange on environmental metabolite concentrations. Previous work has
shown that these are well described by constraints on the metabolic capacity of bac-
terial cells. These are given by an 'enzymatic constraint' which poses a limit on the
summed metabolic flux through the whole cell ([20]), or through more advanced meth-
ods such as flux balance analysis with molecular crowding (FBAwMC) ([21]) which weighs
the flux constraint according to the efficiency and volume of the individual enzymes
([22]). Our model is, therefore, based upon the previous approach by Van Hoek and
Merks ([18]) that uses FBAwMC. However, because such detailed data on enzyme ef-
ficiency and volume is unavailable for the majority of GEMs, here we have replaced
FBAwMC with the simpler 'enzymatic constraint' approach ([20]). This approach allows
us to model metabolic limitations and substrate-dependent switches in metabolism.

The model represents the first 21 days of development of the infant microbiota
in a dynamic, spatially extended, gut-like environment, and considers three scales. At
the microscopic scale, the model simulates individual metabolites. At the mesoscopic
scale, metabolism and growth of bacteria occurs as a function of the local availability of
metabolites. The local environment is depleted by the bacterial models, and receives
the metabolites they deposit. This will influence the availability of metabolites in the
next timesteps. The macroscopic scale represents the whole ecosystem. At this level
a diffusion model simulates the spread of metabolites between adjacent lattice sites,
and a local mixing model simulates mobility of bacterial populations. The metabolites
also advect distally to represent luminal movements, while to represent adherence
to the mucus the bacterial populations do not advect. Bacterial populations are also
randomly deleted to represent local extinction, and they can form new populations
in their immediate environment. The system currently considers 15 bacterial species
based on 15 GEMs taken from the AGORA collection ([23]), a database of 773 semi-
automatically reconstructed GEMs of human gut bacteria that includes all major infant
species ([23]).

The infant gut microbiota is partially limited by carbohydrates. Prebiotic carbohy-
drates increase population sizes *in vitro* and in mice inoculated with infant gut bacteria
([24], [25]). The primary carbohydrate in human milk and nearly all infant formula is lac-
tose, and infant formulas without lactose lead to a different infant microbiota with
greatly reduced *Bifidobacterium* abundance ([14]). Therefore to a first approximation,
we ignore key nutrient sources such as human milk oligosaccharides, fats, protein
residues, or intestinal mucus and focus on carbohydrate metabolism, here taking lac-
tose as the only input carbon source in the model.

In comparison with the previous approach ([18]) upon which the present work builds,
the technological advance is that the present framework simulates a dynamic ecology
of a large diversity of bacterial species. To this end, the modeling framework can, with
minor modifications, import any GEM written in SBML, such as those in the AGORA

FIG 1   The multi-scale metabolic model
(A) Schematic of the model at a system level. Circles represent bacterial populations, color represents species. Flow through the tube is from left (proximal) to right (distal). Lactose is placed at the proximal side. Output metabolites are examples, and depend on bacterial metabolism. Lattice dimensions and ratio are schematic. (B) Screenshot of the bacterial layer of the model at a single time point. Color indicates species, brightness indicates growth rate on the current timestep. (C,D) Screenshots showing, respectively, lactose and lactate concentration of the model corresponding to B. Intensity indicates concentration.

database that we use in this project (23). The previous system (18) could only describe
a 'superbacterial' metabolic network representative of the adult microbiota, while bac-
terial diversity was simulated by activating or repressing individual pathways. In the
present work we study a selection of species representative of the infant microbiota.
The new system also allows us to analyze and visualise the flow of fluxes through the
emergent network of dynamically interacting spatially distributed populations. The
conceptual advance following from this technological advance is that we can now pre-
dict the effect of environmental conditions (oxygen, food, etc.) on the relative abun-
dance of individual bacterial species. This advance has allowed us to use the system to
demonstrate the mechanistic consistency of the hypothesis (4, 5) that a transition from
an Enterobacteriaceae-dominated microbiota to a *Bifidobacterium* spp.-dominated mi-
crobiota is explained through gradual depletion of oxygen leading to anaerobiosis.
The model suggests that high levels of initial oxygen can lead to prolonged dominance
by Enterobacteriaceae, even after oxygen has been depleted, possibly explaining ob-
served variation in the composition of the infant microbiota (1, 13), and reveals pos-
sible cross-feeding interactions between the species. Altogether the model shows to
be a useful tool for qualitatively evaluating hypotheses on the dynamics of the infant
gut microbiota.

RESULTS
Model outline The model of the infant colon is discussed in detail in Section Meth-
ods. Briefly, given a typical infant colon length of 45 cm (26) and diameter of approxi-
mately 1.5 cm (27) the colon is described on a regular square lattice of 225×8 boxes

of 2 $mm \times 2\ mm$ each, thus reaching a good balance between computational efficiency
and accuracy (Fig. 1A). Because bacterial populations can vary at a much smaller scale
(28, 29), the boxes are purely used as a computational discretization. Each lattice site
can contain a population of one species, representing only the locally dominant bacte-
rial species. It is described by a GEM from a list of 15 species prevalent in the earliest
infant gut microbiota ((30, 11) selected from the AGORA collection (23)) (Table 1, Meth-
ods section Species composition).

A simulation proceeds as follows. We initialize the system by placing approxi-
mately 540 small populations of the 15 species randomly across the lattice (Fig. 1B). We
run the model in timesteps representing three minutes of real time, these timesteps
proceed as follows: To mimic carbohydrates entering the colon from the small intes-
tine, a pulse of lactose is introduced every 60 timesteps to the lattice sites of the six
most proximal columns (Fig. 1A-1). Then we predict the metabolism of local bacte-
rial populations using FBA. To mimic metabolic trade-offs, we use an enzymatic con-
straint that sets the maximum summed flux for each FBA solution. The uptake bounds
for each population are set according to the local concentrations of metabolites (Fig.
1C&D). FBA returns a solution that maximizes the rate of ATP production, which is a
suitable proxy for the rate of biomass production (31) given the focus on carbohydrate
metabolism. This biomass production rate (indicated by brightness in Fig. 1B) is used
to update the local population size. Populations spread into an available adjacent
lattice site once the population size has exceeded a threshold. The FBA solution also
returns a set of metabolite exchange rates, which are used to update the local metabo-
lite concentrations (Fig. 1D). In total, 723 extracellular metabolites are described in the
model, but only 17 of these are typically found in more than micromolar amounts in
our simulations (Methods, Table 2). Bacterial populations diffuse through swaps of
the content of adjacent lattice sites (Fig. 1A-2). Metabolites diffuse between adjacent
lattice sites (Fig. 1A-3) and advect to more distal sites (Fig. 1A-4) over one lattice per
timestep, reaching transit times of effectively 11 hours (32, 33). Metabolites and pop-
ulations that reach the distal side of the system (Fig. 1A-5) are removed. To mimic the
appearance of new bacterial populations (e.g. from ingested spores or introduction
from intestinal crypts (34)) each empty lattice site now has a small probability to be
filled with a new bacterial population selected at random from all available species
(Table 1). This completes the timestep, after which we record the metabolite concen-
trations and bacterial population in the whole model. All parameters are given in Sec-
tion Parameters. We have also performed a sensitivity analysis for several parameters
around the default parameter set shown in Table 2 (Section Sensitivity analysis).

We performed several checks to ensure the biochemical and thermodynamic plau-
sibility of the model (Methods section Checking the validity of the GEMs). After the
resulting corrections to the GEMs (S1 Table), no atoms were generated or lost by the
FBA solutions, there was no flux through the objective function without a substrate,
and 99.999% of all fluxes of Fig. 3A contained less free energy in the output than in the
input. The remaining 0.001% all had no energy increase and growth rates less than
0.001% of the average growth rate. We therefore concluded that the system conserved
mass and energy.

Enzymatic constraint reproduces metabolic switching in *Bifidobacterium* Many
microorganisms switch between metabolic pathways of higher and lower yield de-
pending on substrate availability. These so called metabolic switches can be repro-
duced with FBA through an enzymatic constraint (20, 35, 36, 22) that limits the summed
metabolic flux of a bacterium. *Bifidobacterium* also displays a metabolic switch through
a pathway known as the bifid shunt. At low extracellular sugar concentrations, *Bifi-*

TABLE 1 Species included in our model, all from the AGORA collection([23]). Color indicates color used in figures

Species	Family	Anaerobic? ([23])	Reactions
Bacteroides vulgatus	Bacteroidaceae	Yes	2474
Bifidobacterium breve	Bifidobacteriaceae	Yes	1987
Bifidobacterium longum ssp. *infantis*	Bifidobacteriaceae	Yes	1005
Bifidobacterium longum ssp. *longum*	Bifidobacteriaceae	Yes	2043
Blautia hansenii	Lachnospiraceae	Yes	2055
Collinsella aerofaciens	Coriobacteriaceae	Yes	920
Dorea formicigenerans	Lachnospiraceae	Yes	2072
Parabacteroides distasonis	Porphyromonadaceae	Yes	2519
Ruminococcus gnavus	Lachnospiraceae	Yes	2225
Veillonella dispar	Veillonellaceae	Yes	1124
Enterococcus faecalis	Enterococcaceae	Facultative	1174
Enterobacter cloacae	Enterobacteriaceae	Facultative	1800
Escherichia coli SE11	Enterobacteriaceae	Facultative	2356
Lactobacillus gasseri	Lactobacillaceae	Facultative	1163
Streptococcus salivarius	Streptococcaceae	Facultative	1190

dobacterium produces a mixture of acetate, ethanol, and formate in a 1:1:2 ratio from
pyruvate. At increased sugar concentrations the bifid shunt instead reroutes pyruvate
to lactate. In addition acetate is produced from acetyl-phosphate independently, such
that at low concentration of sugars *Bifidobacterium* produces acetate, ethanol and for-
mate in a 4:1:2 ratio, and at increased sugar concentrations *Bifidobacterium* switches
to production of acetate and lactate in a 3:2 ratio ([37], [38]).

To test if the enzymatic constraint suffices to explain the metabolic switch in *Bifi-*
dobacterium, we performed simulations of a single timestep in increasing lactose con-
centrations (Fig. 2A). These correctly predicted the metabolic switch. The ratios be-
tween the metabolites before and after the metabolic switch, as well as the transient
between the two regimes and the pathways used (S2 Table) match the experimental
observations. Simulations without the constraint only reproduced the situation at low
concentrations of lactose (S1 Figure A), showing that the enzymatic constraint is re-
quired. The model also predicts an unobserved metabolic switch on lactose in *E. coli*
(S1 Figure B) and reproduced the metabolic switch in *Bacteroides vulgatus* ([39]) (S1 Fig-
ure C). Another effect of the enzymatic constraint is that the growth rate saturates at a
species-dependent concentration (Fig. 2B, S1 Figure D). As a consequence the relative
growth rates of the species depend on the lactose concentration.

Model predicts *Bifidobacterium* dominance through metabolism under anaer-
obic conditions After having established the effect of the enzymatic constraint in
models of individual bacterial populations, we studied the behavior of the full mul-
tiscale model for anaerobic conditions. After 21 days (10080 timesteps) *Bifidobac-*
terium spp. had become the most abundant in 27 of 30 simulations (Fig. 3A, S1 Video), a
typical situation found in infants of a corresponding age ([11], [2]). Also in agreement with
in vivo data, the model predicted the presence of populations of *E. coli*. Our model also
predicted populations of *Blautia hansenii*, which is only occasionally present in the in-
fant colon ([11]). We will investigate the niche of this species further in the section 'Non-
Bifidobacterium species benefit from lactate consumption'. All species other than the
aforementioned had abundances either lower or not significantly higher than their
initial abundance after 21 simulated days (p>0.05, two-sample t-test). To test if these
species were required for the formation of the community we ran an additional 30 sim-
ulations in which we only included species that had an abundance significantly larger

FIG 2 **Enzymatic constraint reproduces metabolic switching in *Bifidobacterium***
(A) Production of metabolites per timestep by a *B. longum infantis* population of $5 \cdot 10^9$
bacteria with access to one lattice site (0.05ml) using our FBA with enzymatic constraint
method. (B) Growth per timestep by lactose concentration for populations of $5 \cdot 10^9$ **bac-**
teria with access to one lattice site (0.05ml) of some major bacterial species using our
FBA with enzymatic constraint method. *B. longum infantis* is used as the *Bifidobacterium*
representative.

than their initial abundance at the end of other simulations. This led to a very similar
set of outcomes (S2 Figure A). In these simulations none of the species abundances
differed from the original simulations with all species included (p>0.05, two-sample
t-test). In the remaining simulations all species are included in order to prevent an ini-
tial bias, thus leaving in the possibility that any of the rare species become abundant
under one of the conditions considered. In practice this did not occur.

We next analysed the metabolites diffusing or advecting out of the distal end of
the gut over the last two days of the simulations of Fig. 3A. We found considerable
amounts of acetate, formate, lactate, hydrogen, ethanol, and succinate (Fig. 3B). All
other metabolites were present in much smaller quantities. The high abundance of ac-
etate and lower abundance of lactate agree with the fecal composition of three month
old formula-fed infants (40), and is characteristic of microbiota dominated by *Bifidobac-*
terium. Formate, hydrogen, ethanol and succinate are also all reported in infant feces
(41, 42, 43, 44).

To examine how this metabolic pattern is formed we analysed the metabolic pro-
duction and uptake in a sample run (Fig. 3C). *Bifidobacterium* converts lactose into a
mix of acetate, lactate, ethanol, and formate, as predicted (Fig. 2A), with a higher lac-
tate production than ethanol or formate. Some of this lactate is reabsorbed in later
timesteps and converted into the three other metabolites by *Bifidobacterium*. A mix-
ture of lactose and crossfeeding products is converted by *E. coli* and *B. hansenii* into
the remaining metabolites of Fig. 3B. The main crossfeeding nutrient absorbed is lac-
tate, produced by *Bifidobacterium*, but there is also uptake of ethanol, formate, acetate,
and succinate. These are also produced by the crossfeeders. The network has become
simpler at the end of this run, as *B. hansenii* is at a very low abundance and *E. coli* now
only consumes lactose, lactate, and ethanol, which it does not produce (Fig. 3D).

FIG 3 *Bifidobacterium* spp become dominant anaerobically and mostly proximally with a unique metabolic profile

(A) Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (B) Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation. n=30. (C,D) Visualisation of metabolic interactions in the sample run of **S1 Video**. Green lines represent production, yellow lines represent consumption. Line width and intensity is proportional to the amount exchanged with the environment, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels. Data from hour 90 to 93 (step 1800 to 1860) for C and last 3 hours (step 10020 to 10080) for D (E) Spatial distribution of lactate over the last two days (960 timesteps) of the simulation. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (F) Abundance per location for grouped species over the last two days (960 timesteps). Curve shows mean value and shaded area one standard deviation over n=30 simulations. (G) Metabolism of lactose by grouped species. All data from last two days. n=30. (H) Metabolism of lactate by grouped species. All data from last two days. n=30.

Crossfeeding emerges in simulations The observed crossfeeding in the sample run (Fig. 3C&D) suggested that production of metabolites by *Bifidobacterium* spp. may be crucial for *E. coli* and *B. hansenii*. To further analyse such crossfeeding interactions we next analysed the spatial distribution of lactose and the Bifidobacterial products lactate and acetate. From day 19 to 21, lactose was present mostly proximally (S2 Figure B), lactate was most abundant in the middle of the colon (Fig. 3E), and acetate was most abundant distally (S2 Figure C). To analyse the role of *Bifidobacterium* spp. in shaping this metabolic pattern, we analysed the location of populations from day 19 to 21 (Fig. 3F). *Bifidobacterium* spp. was located proximally, whereas the other species were located more distally. These results suggested that separated metabolic niches had formed in the simulated colon. To quantify crossfeeding interactions we summed the flux of lactose and lactate for *Bifidobacterium* and for all non-*Bifidobacterium* species grouped together. *Bifidobacterium* spp. was the largest consumer of lactose (Fig. 3G) and both the largest producer and consumer of lactate (Fig. 3H). An additional consumption of lactate towards acetate, formate and ethanol takes place at lower lactose concentrations, when the enzymatic constraint is not saturated by lactose uptake. However, the non-*Bifidobacterium* groups consumed more lactate relative to their lactose consumption. To test the effect of lactate uptake by *Bifidobacterium*, additional simulations were run in which the conversion from lactate to pyruvate was blocked, preventing *Bifidobacterium* from consuming lactate. In these simulations *Bifidobacterium* was the most abundant group in 20 of the 30 simulations, with the largest average abundance (S2 Figure D). Thus lactate consumption played a small role but was not the primary cause of *Bifidobacterium*'s dominance. We hypothesized that *Bifidobacterium*'s primary niche is as primary consumer, consuming lactose, while for other species niches exist as primary and/or secondary consumer around consuming lactose or lactate. Combined with the observation (Fig. 3E) that lactate concentration is lower in the distal colon and the recorded crossfeeding fluxes in our network visualisation (S1 Video), we can conclude that the population at the distal end of the simulated colon consists at least partially of such secondary consumers, i.e. crossfeeders.

Lactose uptake through the bifid shunt is essential for *Bifidobacterium* dominance Next we determined whether the bifid shunt is essential to *Bifidobacterium* dominance in the model. We disabled the bifid shunt by blocking all flux through the reactions associated with fructose-6-phosphate phosphoketolase and ran 30 simulations. Fructose-6-phosphate phosphoketolase is unique to and necessary for the bifid shunt, and characteristic of *Bifidobacterium* as a genus (45). *Bifidobacterium* spp. were dominant in none of these simulations (Fig. 4A, S2 Video). Instead, *E. coli*, *B. vulgatus*, and *B. hansenii* were dominant. This indicates that the bifid shunt is crucial to the dominance of *Bifidobacterium* spp. in our model. We also ran 30 simulations in which only the production of lactate was disabled in *Bifidobacterium* (S2 Figure F). 27 of these simulations still led to *Bifidobacterium* dominance. This indicates that the metabolic switch to lactate production was not essential to *Bifidobacterium* dominance in our model. *B. vulgatus* also had a higher average abundance, but this was not significant (p=0.27, two-sided t-test).

B. hansenii but not _E. coli_ can sustain itself on crossfeeding We observed that non-*Bifidobacterium* species appeared in the simulation behind a proximal population of *Bifidobacterium* spp. (Fig. 3F). These non-*Bifidobacterium* species consumed lactose (Fig. 3D&G), despite the fact that lactose concentrations have dropped compared to the proximal part (S2 Figure B). We therefore wondered how these non-*Bifidobacterium* species could persist in the model. At reduced lactose concentrations, *E. coli* produces more ATP per timestep than *Bifidobacterium* spp. (Fig. 2B). We hypothe-

FIG 4 *Bifidobacterium* and *E. coli* require lactose, but *B. hansenii* requires lactate
(A) Abundances for grouped species over 21 days. The bifid shunt was disabled for all *Bifidobacterium* species. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (B) Abundances for grouped species over 21 days, but with lactose disabled for non-*Bifidobacterium* species from 5040 timesteps (10.5 days) onwards. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (C,D) Visualisation of metabolic interactions in a sample run. Green lines represent production, yellow lines represent consumption. Line width and intensity is proportional to the amount exchanged with the environment, with a threshold of 0.5 μmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels. Data from hour 90 to 93 (step 1800 to 1860) for C and last 3 hours (step 10020 to 10080) for D. (E) Growth per timestep by lactate concentration for populations $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml) of some major bacterial species (F) Abundances for grouped species, with lactate consumption disabled for all species. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

sized that non-*Bifidobacterium* species outcompete *Bifidobacterium* at reduced lactose concentrations in the model. To test this hypothesis, we blocked lactose consumption by non-*Bifidobacterium* species after 10.5 days, the midway point of the simulation, and ran 30 simulations. In this way, we ensured that sufficient secondary resources were produced by *Bifidobacterium* in case non-*Bifidobacterium* species made use of them. We observed a sharp population decrease of non-*Bifidobacterium* species starting directly after blocking lactose uptake (Fig. 4B). *E. coli* had a near-zero abundance at the end of these simulations. *B. hansenii* also had a significantly lower abundance compared to the baseline (p<0.01, two-sample t-test), but still had a presence (S2 Video). Thus, in our simulations, some amount of primary consumption was essential for *E. coli*, but not for *B. hansenii*.

Non-*Bifidobacterium* species benefit from lactate consumption To analyse how *B. hansenii* could sustain a population as secondary consumer, we investigated the substrates used by non-*Bifidobacterium* species in the model. Analysis of the flow of metabolites between species in a new run showed that the uptake of lactose by *B. hansenii* decreases over time, while the uptake of lactate increased (Fig. 4C&D). *E. coli* and *B. hansenii* also produce more ATP from lactate than *Bifidobacterium* at any concentration (Fig. 4E). Combined with our earlier observations of crossfeeding on lactate (Fig. 3D&H) we hypothesized that *Bifidobacterium* species cannot compete with *E. coli* and *B. hansenii* on pure lactate. Consumption of lactate by *B. hansenii* could be essential to its ability to sustain populations without lactose uptake.

To investigate if lactate is essential in feeding secondary consumer populations, we blocked the lactate uptake reaction for all species and ran 30 simulations. (Fig. 4F, S2 Video). At 21 days *B. hansenii* populations were much smaller than in simulations with functional lactate uptake, not significantly larger than the initial populations (p>0.05, two-sample t-test). *E. coli* populations were also significantly smaller than those in simulations with lactate uptake, but also significantly larger than their initial population at the start of the run (p<0.05, two-sample t-test). *Bifidobacterium* populations were similar to those in the baseline simulations (Fig. 3A). Together with the results in Fig. 4B this indicates that *E. coli* does not require lactate as a crossfeeding substrate like *B. hansenii* does.

E. coli outcompetes *Bifidobacterium* by taking up early oxygen During the first days Enterobacteriaceae or Bacilli such as *Streptococcus* are dominant in the *in vivo* infant gut microbiota, after which *Bifidobacterium* spp. are often dominant (1, 11). This pattern may be explained by the presence of oxygen in the gut shortly after birth (5, 3), e.g. via diffusion through the gut wall (7) or accumulation in the newborn infant gut *in utero* in the absence of microbes (46). Intracolonic oxygen diffusion depends on host tissue oxygenation and colonic blood flow (47, 48). To mimic the early presence of oxygen we introduced 0.1 to 10 µmol initial oxygen per lattice site. These values were chosen arbitrarily in absence of precise data, but cover a wide range of outcomes in our model. The oxygen diffused, but did not advect in the model.

0.1 µmol initial oxygen sufficed to reproduce the pattern observed *in vivo*: *E. coli* initially dominated and *Bifidobacterium* became dominant after a few days or weeks of simulated time (Fig. 5A, S3 Video). However, we did not see dominance of *Streptococcus* in any of the simulations, in contrast to observations *in vivo* (2). The initial presence of oxygen strongly affected the other species in the model: *B. hansenii* remained nearly entirely absent, while *B. vulgatus* had a higher abundance than it did without initial oxygen. In the network analysis of a sample run a similar pattern occurs as in the networks of Fig. 3C, but with different early species (Fig. 5B). Here the Enterobacteriaceae *E. coli* and *Enterobacter cloacea* consume more of the early lactose,

FIG 5 Initial oxygen leads to initial *E. coli* dominance and succession by *Bifidobacterium* (A,D,E) Abundances for grouped species. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. Initial oxygen per lattice site in µmol is (A) 0.1 (D) 1 (E) 10 (B,C) Visualisation of metabolic interactions in a sample run. Green lines represent production, yellow lines represent consumption. Line width and intensity is proportional to the amount exchanged with the environment, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels. Data from hour 30 to 33 (step 600 to 660) for B and last 3 hours (step 10020 to 10080) for C (F) Distribution of total *Bifidobacterium* abundance at 21 days (10080 timesteps) performed with 10 µmol initial oxygen per lattice site n=30. (G) Growth per timestep by lactose concentration in the presence of abundant oxygen for populations of $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml) of some major bacterial species (H) Abundances for grouped species with oxygen uptake disabled for *E. coli*. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

along with oxygen. By the end of the run *Bifidobacterium* spp. again took up the role of primary lactose consumer and lactate producer, whereas *E. coli* had fed largely on lactate (Fig. 5C), thus assuming a role of secondary consumer.

At 1 or 10 μmol initial oxygen per lattice site *E. coli* and *E. cloacea* remained dominant and the main primary consumers for longer (Fig. 5D, S4 Video), and succession to *Bifidobacterium* took much longer, or did not occur at all within the simulated timeframe (Fig. 5E). At 10 μmol oxygen per lattice site *Bifidobacterium* either dominated the population or remained much smaller, with few intermediate cases, leading to a bimodal distribution (Fig. 5F). This prediction matches *in vivo* observations (1, 13). The proportion of infants dominated by Enterobacteriaceae at 21 days of age varies depending on the study population, but can be 22% (2) or 25% (49) after 21 days, an intermediate value between the 10% (Fig. 5D) and the 60% (Fig. 5E) that we observe with different oxygen levels. We also initialised four sets of simulations in which oxygen was released from the upper and lower boundaries of the system every timestep (S3 Figure A-D). This led to an increasingly stronger stimulation of *E. coli* with higher levels of oxygen, but did not lead to any of the temporal effects observed *in vivo*. In a separate set of simulations we stopped oxygen release at the midpoint of our simulations. We observed an increase of *Bifidobacterium* spp. and a decrease of *E. coli* after that point (S3 Figure E), leading to a bimodal outcome similar to the simulations with initial oxygen (S3 Figure F).

We further examined the causes of *E. coli* dominance over *Bifidobacterium* in the presence of oxygen in our model. Fig. 5G shows that when oxygen is available *E. coli* has a much higher growth per timestep in our model than *Bifidobacterium* spp. for all concentrations of lactose, even though *Bifidobacterium* spp. also produce more ATP than under anaerobic conditions. *B. hansenii* and *Streptococcus salivarius*, in contrast, grew slower than *Bifidobacterium* for all concentrations in the presence of oxygen. This effect depends on the local oxygen concentration, but *E. coli* grows faster than other species even at concentrations much lower than our initial value (S3 Figure G). *E. coli* does not use overflow metabolism in our model, though *E. cloacea* does, and *E. coli* does when in the presence of glucose (S3 Figure H,I,J).

To test if the direct uptake of oxygen was indeed responsible for *E. coli*'s dominance of the microbiota, we disabled the oxygen uptake reaction of *E. coli* for 30 simulations with 1 μmol initial oxygen per lattice site. Under these conditions, *E. coli* failed to dominate the population (Fig. 5H). Other species, primarily *E. cloacae*, became dominant early in the simulations as a primary consumer (S4 Video). These species are replaced in all simulations by a population composition similar to that from the simulations without oxygen. In some studies, *Klebsiella* species were the dominant early members of the Enterobacteriaceae instead of *E. coli* (50). To examine whether our model could also reproduce early dominance by a *Klebsiella* species we initialized the simulations with a set of species containing *K. pneumoniae* instead of *E. coli*. At 0.1 μmol initial oxygen per lattice site, these simulations behaved the same as the original simulations initialized with *E. coli* (S3 Figure K): *Bifidobacterium* became the most abundant group after 21 days in 26 out of 30 simulations. In simulations initialized with both *K. pneumoniae* and *E. coli*, *K. pneumoniae* became the most abundant in 15 of 30 simulations, and *Bifidobacterium* in the other 15 (S3 Figure L). Altogether, the model predicts that *E. coli* relies on direct consumption of oxygen to dominate the microbiota under initially oxygenated conditions in the model. It also indicates that *E. cloacea* does not have an anaerobic metabolism competitive enough to sometimes become dominant over *Bifidobacterium* spp. as *E. coli* does, and that other Enterobacteriaceae such as *K. pneumoniae* can perform similarly to *E. coli*.

Sensitivity analysis A number of key parameters in our model were based on reasonable estimates. To test the effects of these parameter values on the simulation results, we performed a sensitivity analysis for these parameters under anaerobic conditions, which best represents the situation at the end of the simulations (Fig. 6). As shown in Fig. 2B and S1 Figure D, after relaxing the enzymatic constraint *Bifidobacterium* spp. loses its competitive advantage at high lactose concentrations (S4 Video). This leads to reduced dominance or extinction in whole-gut simulations (Fig. 6A). Tightening the enzymatic constraint led to *Bifidobacterium* spp. remaining dominant, but with a much smaller population size. Thus the presence of an enzymatic constraint sufficiently large for metabolic shifts, but not its exact level, is crucial for the prediction of *Bifidobacterium* dominance. We set the enzyme constraint at a level strong enough to predict the metabolic shifts observed *in vitro*, but not so strong as to prevent reasonable growth. The ATP required for a unit of population growth had a linear effect on population size, as expected, but had little effect on the relative abundance of *Bifidobacterium* (Fig. 6B). The large number of populations with a low ATP requirement also made the model computationally unfeasible. The rate of colonization had little effect on the simulation outcomes: at increased colonization rates, *Bifidobacterium* spp. still dominated the microbiota and its relative abundance remained unchanged (Fig. 6C). Even eliminating colonization altogether hardly affected *Bifidobacterium* spp. dominance in most simulations, but led to complete extinction of all *Bifidobacterium* species in 5/30 simulations and of all 12 non-*Bifidobacterium* species in 9/30 simulations. Thus we kept a moderate amount of colonization in the model. We also examined the effect of placing new populations after initialization only in the first column of the lattice, with a probability increased to match the lower number of eligible squares (S4 Figure A). This also led to a slightly larger but still comparatively small population for the non-*Bifidobacterium* species. To determine the optimal spatial discretization we ran the model on a more coarse and on a more refined lattice, scaling the diffusion, advection, and initialization and colonization parameters accordingly (Fig. 6D; see methods section Population dynamics). Lattice refinement did not change the dominance of *Bifidobacterium* nor the spatial distribution of metabolites and populations (S4 Figure B-E), but on a more coarse lattice the smaller number of lattice sites seemed to increase the rate of extinctions and variability of metabolite distributions (S4 Figure F-I). The diffusion of metabolites had a larger effect on the simulation results (Fig. 6E). At low diffusion rates, the populations tended to die out, because they had less opportunity to take up lactose as they were exposed only as it 'passed by' due to advection. At high diffusion rates cross-feeders and *Bifidobacterium* consumed all lactate (S4 Figure J); in infant feces substantial amounts of lactate are found (40, 51). We therefore decided to choose a value for which the system produced the observed concentration of lactate. Finally the amount of mixing of bacteria was varied (Fig. 6F). *Bifidobacterium* remained dominant regardless of the diffusion parameter, even when the bacteria were fully mixed (i.e. placed at random locations) each timestep. Without mixing the populations could not spread, for lack of empty space. The bacterial diffusion constant is 1.3 times higher than the metabolic diffusion constant. As shown in Fig. 6F the exact setting of the bacterial diffusion has little effect on the outcomes of the simulations, so we have maintained bacterial diffusion at this baseline.

DISCUSSION

To develop new hypotheses for possible external effects on the initial phases of the infant gut microbiota, we have developed a dynamic multiscale model. Our simulations reproduce the initial dominance of Enterobacteriaceae, particularly *E. coli*, over *Bifi-*

FIG 6   Sensitivity analysis for estimated parameters
Abundances for *Bifidobacterium* spp. and other species at the end of day 21 with different parameters. n=30 for each condition. (A) Effect of enzymatic constraint around the default value of 2 µmol flux per timestep per $1 \cdot 10^{10}$ population (B) Effect of the amount of ATP required for growth of $1 \cdot 10^{10}$ bacteria around the default value of 10 µmol (C) Effect of variation in continued influx of populations per timestep per empty square around the default of 0.00005 (D) Effect of variation in width and height of lattice, including adjustment of associated parameters, around the default of 8x225 lattice sites (E) Effect of variation in diffusion for metabolites away from the baseline of $4.4 \cdot 10^{-5}$ cm²/s (F) Effect of variation in diffusion of bacterial populations away from the baseline of $5.7 \cdot 10^{-5}$ cm²/s

dobacterium species, and the subsequent dominance of *Bifidobacterium* species, out
of a broad consortium of infant species. Moreover, our simulations suggest show the
consistency of the classical hypothesis that this succession could be due to an initial
presence of oxygen, followed by microbial depletion ([4](), [5]()). These predictions on oxy-
gen suggest that higher levels of initial oxygen may explain the absence or strong delay
of *Bifidobacterium* development that is observed in some infants ([2]()). A continuous in-
put of oxygen did not explain the pattern observed *in vivo* (S3 Figure A-D), but stopping
the input of oxygen at the midway point of the simulation did lead to a similar pattern
(S3 Figure E-F).

While it is known that there is variation in colonic oxygenation ([47](), [48]()), the range
of this variation in newborns is not known. In premature infants a positive association
exists between the number of days of supplemental oxygen and *Klebsiella* sp. abun-
dance ([50]()). *Klebsiella* was the primary representative of the Enterobacteriaceae in this
study population. However, only very few of these infants developed a *Bifidobacterium*
dominated microbiota, regardless of the duration of supplemental oxygen. Nonethe-
less, it may be desirable to reduce oxygen concentration faster to achieve the positive
health effects of *Bifidobacterium* more consistently. As the direct oxidation of lipids
and other organic substrates may contribute to creating anaerobic conditions in the
colon ([7]()), altered nutrition may allow for a decrease in oxygen in the colon without
stimulating facultative anaerobic bacteria such as *E. coli*. There are also indications
that short-chain fatty acids produced by *Bifidobacterium* may decrease oxygen levels
by stimulating the oxygen use of colonocytes ([8]()). Future versions of the model could
represent these oxygen-depleting mechanisms by localized (e.g., near the intestinal
wall) and distributed oxygen consumption terms. At present the model predicts that
distributed oxygen usage by such additional processes will speed up the succession
to an anaerobic, *Bifidobacterium*-dominated microbiota. The limit case, i.e. extremely
fast oxygen consumption, is represented by the anaerobic case (Fig. [3]()). The effect of
localized oxygen consumption processes, e.g., near the intestinal wall is a topic of our
ongoing research.

Our simulations predict that the dominance of *Bifidobacterium* spp. in the anaero-
bic infant system could be explained from its consumption of lactose through its bifid
shunt metabolism. Our model can correctly represent bifid shunt metabolism due
to the implementation of an enzymatic constraint on the maximum total metabolic
flux in the bacteria (Fig. [2]()). Both the low and high yield metabolism, and the switch
between them based on nutrient availability, have been extensively described *in vitro*
([38]()). The FBA with enzymatic constraint method we use is similar to flux balance analy-
sis with molecular crowding (FBAwMC) ([21]()). As in FBAwMC we consider the enzymatic
constraint to represent the limited availability of space and production capacity for en-
zymes within a bacterial cell. However, in the absence of good reaction-specific ther-
modynamic information for our current model, we do not utilize the reaction-specific
technique of FBAwMC. We instead base an enzymatic constraint only on the size of the
local population, which is sufficient for modeling a metabolic switch. This is equivalent
to using FBAwMC and placing all crowding coefficients at the same positive number,
which has been done previously to model metabolic switches ([21]()). More broadly, our
method is similar to other existing methods that model metabolic switches by includ-
ing limits on fluxes that represent some physical limit on metabolism, such as pro-
teome constraints or membrane occupancy ([52](), [35](), [53](), [54]()). It has been shown that
metabolic switches in microbes can be modelled in many ways, provided that two si-
multaneous constraints are in place ([22]()). In our case, these are the concentration of
substrate and the enzymatic constraint.

The metabolic switch of *Bifidobacterium* in our model determines the production
of its metabolites. The two most abundant, acetate and lactate, are present in the
simulated feces at a median ratio of 3.5:1 (Fig. 3B), similar to that found in formula-
fed infant feces (40, 51). We can derive a concentration in mM if we assume each of
the eight lattice sites whose metabolites are advected out of the system each step
contains 0.05ml of water. This is based on an estimated total volume of 90ml, divided
over 1800 lattice sites (Table 3). This results in an average concentration of 29 mM
acetate and 8 mM lactate in the simulated fecal output analyzed in Fig. 3B. These
values are close to, and well within the variability, of the average values for acetate and
lactate of 31 and 12 mM respectively in the feces of two-week old infants in (51). The
median ethanol:acetate ratio of 1:21 is similar to the ratios of 1:18 and 1:10 reported
in term infants (43) and the 1:50 reported in pre-term infants (55). We use ratios here
because the measures per gram dry weight are not available in our model. We observe
a lower quantity of ethanol in the simulated fecal output of the simulations where
lactate uptake by *Bifidobacterium* was disabled (S2 Figure E).

Bifidobacterium takes up lactate in our model and converts it into pyruvate, cat-
alyzed by lactate dehydrogenase. In our model this pyruvate is then further converted
to formate, ethanol, and acetate through the high-yield pathway also used in lactose
metabolism. However, strong lactate uptake by *Bifidobacterium* spp. has not been ob-
served *in vitro* (in e.g. (56)). Thus, while the enzymes used for lactate consumption
exist in *Bifidobacterium*, it is not known if the pathway is used *in vivo* to the extent
predicted by the model. Lactate uptake by *Bifidobacterium* has only little impact on
the simulation results (S2 Figure D). Similarly, though growth on lactate by *E. coli* in
the presence of oxygen is well documented, this is not the case for the anaerobic con-
dition (57). Lactate uptake is not essential for *E. coli*, as it can exist in a more purely
primary consumer role (Fig. 4F). The third consumer of lactate, *B. hansenii*, is present
only marginally *in vivo* (11). This matches with our results for initially oxygenated con-
ditions, in which *B. hansenii* is also largely absent. However, other species that could
fill the same lactate-consuming secondary consumer niche as *B. hansenii* in our model
are common *in vivo* (58). Our selection of species may be further improved to allow the
correct species to emerge in this secondary consumer niche. For example, it might be
appropriate to add *Eubacterium hallii*, a common infant gut bacterium known to con-
sume *Bifidobacterium*-produced lactate and acetate *in vitro* (59).

More generally, our model predicts a lower diversity overall in both bacterial species
and metabolites compared to what is observed *in vivo* (11). While the abundant species
in our model largely match with the abundant species *in vivo*, many other species such
as those of the genera *Streptococcus* and *Lactobacillus* have a lower abundance in our
model compared to the *in vivo* conditions(11). We have decided to keep all less abun-
dant species in our simulations, to show how in most cases the correct species out
of a broad consortium become dominant. There were no large differences when we
ran the simulation without these smaller species (S2 Figure A). Particularly the lack of
representation of the Bacilli species in our model outcomes is notable. While some
sources report a portion (e.g. 12-14% (2)) of their subjects to be dominated by Bacilli,
mainly *Streptococcus*, the *Streptococcus* species in our model never became dominant.
We showed that *S. salivarius* has an inferior metabolism on lactose and lactate com-
pared to other species in our model (Fig. 2A, Fig. 4E, Fig. 5G).

The discrepancy between our model and the *in vivo* data in the abundance of less
common species such as *Streptococcus* may partially be explained through the focus
of our model on carbon dissimilation. Though it would be preferable to represent the
whole metabolism of the infant microbiota, including factors such as consumption of

amino acids, oligosaccharides and intestinal mucus, the initial version presented here
only considers carbon metabolism from lactose. While a more extensive metabolism
might have allowed more mechanisms and niches to be discerned, it would also intro-
duce additional free parameters, as there is no clear data on the concentration and
uptake of these nutrients. The uptake bounds would have to be set arbitrarily, with
uniform values or random sampling. Many substances such as fatty acids or protein
residues are not even included as metabolites in the database we use for bacterial
metabolism ([23]), further complicating their introduction to the model. By focusing on
carbon metabolism, using ATP production as a proxy for growth rate, and only using
lactose as an input nutrient, we can circumvent these problems. ATP production has
been shown to be a good proxy for biomass production in *E. coli* ([31]). Supplementation
of the *in vitro* and model organism infant gut microbiota with prebiotic carbohydrates
led to a larger microbiota, primarily due to larger *Bifidobacterium* populations([24], [25]).
This indicates that carbon metabolism is a limiting factor, especially in the absence of
prebiotics, as in our model. However, as the increased population in these *in vitro* and
in vivo studies consisted largely of *Bifidobacterium*, other species were not necessarily
also carbon limited. In addition, ATP production may not be a good proxy for biomass
production in some species. Species-specific information on the relation between ATP
and biomass production may aid future modelling. Finally, gaps in the GEMs we use
may cause ATP production itself to be underestimated compared to what occurs *in*
vivo.
Besides details on the modelling of metabolism, the way that diffusion and ad-
vection of bacteria and metabolites is handled in the simulations may impact the pre-
dictions of our model. The sensitivity analysis demonstrated an unrealistic metabolic
output when the metabolic diffusion coefficient was raised to $2.2 \cdot 10^{-4}$ cm^2/s ([S4 Fig-
ure] J). Although this value is much lower than diffusion coefficients measured in the
adult intestinal lumen ($1 \cdot 10^{-2}$ cm^2/s) ([60]), in the infant colon mixing may be reduced
compared to the adult colon: motor activity that mixes the colon contents ([68]) is rarer
in infants than in older children or adults ([61], [62]). Exact measurements of mixing in
the infant colon will be hard to obtain. Related to this, while we did not examine the
effect of bacterial advection in our current work, our previous work ([18]) suggested
that increased bacterial advection greatly reduces diversity and spatial patterning in
the model. In fact, compared to the adult gut, in the infant gut there is increased mo-
tor activity driving advection ([62]) and infants display faster colonic transit than adults
([32], [33]). In light of these data, which seem to indicate much faster bacterial advection
and stronger luminal mixing than what we have assumed in our model, a potential
interpretation of the present set-up of our model is that it represents the dynamics
of bacterial populations adhering to the intestinal wall in interaction with metabolites
advecting through the lumen. Our future work will explore in more detail how a bal-
ance between adhesion of microbiota to the mucus ([63]), versus advection of bacteria
and metabolites in the lumen affects the colonization of the infant colon.
Our modeling approach relates to alternative simulation frameworks, such as Steady-
com, BacArena, and COMETS ([16], [64], [17], [65]). These frameworks, in particular the new
COMETS 2 framework ([19]) would certainly be suitable for answering questions similar
to those asked in the present work. In absence of suitable frameworks at the initial
phases of this project, and given the flexibility that comes with using an in-house code
base, we preferred to continue developing our own line of gut models ([18]). A future
implementation of our model in one of the available simulation frameworks would be
useful exercise, facilitating future development and comparison of models.
We have used our modeling approach to generate testable hypotheses on the

causes and mechanics of succession in the infant gut microbiota, potentially laying the foundation for nutritional interventions that could improve the health of infants. It should be emphasized that the current and future work on this model represent a tool for generating hypotheses, and for testing potential mechanical explanations. We cannot fully represent the complexities of the infant gut microbiota, and any generated hypotheses must necessarily be validated *in vitro* and *in vivo*. There is work to be done to further study the relevant factors, and to bring these results into a more realistic model context. We aim to do so by integrating the prebiotic oligosaccharide and protein content of nutrition into the model. This may lead to the creation of more niches in the model, and so a diversity closer to that of the *in vivo* system. We will also continue to improve the selection and curation of metabolic models, such as by disabling lactate uptake in certain species or including a wider diversity of GEMs. This may lead to novel future insights on the interactions between differences in infant nutrition and succession in the infant microbiota.

METHODS

We use a spatially explicit model to represent the newborn infant microbiota (Fig. 1A). Our model is based on an earlier model of the adult gut microbiota ([18]). The model consists of a regular square lattice of 225×8 lattice sites, where each lattice site represents $2\ mm \times 2\ mm$ of space, resulting in an infant colon of 450mm by 16mm. Each lattice site can contain any number of metabolites of the 723 types represented in the model in any concentration and a single simulated bacterial population. The metabolism of these bacterial populations is based on genome-scale metabolic models (GEMs) from the AGORA collection of species-specific GEMs ([23]). From this set, we have chosen 15 GEMs based on a consortium of known infant microbiota species ([30], [11]). Their metabolic inputs and outputs are calculated using dynamic flux balance analysis (FBA) ([66]) with an enzymatic constraint functioning as a limit on the total flux through the network ([20]). The effects of the FBA solution are applied to the spatial environment and then recalculated each timestep, creating a spatial dynamic FBA.

We identify the two narrow ends of the rectangle with the proximal and distal ends of the colon. In each timestep metabolites both mix and flow from the proximal to the distal end. Bacterial populations are mixed, but do not flow distally as metabolites do.

At the start of each run, we initialize the system with a large number of small populations. We let these perform their metabolism each timestep. They take metabolites from the environment, and deposit the resulting products. We let the populations grow and divide according to their energy output. Both the initial placement locations and movement of the populations is random, introducing stochasticity in the model. The system develops differently depending on initial conditions, into a diverse and complex ecosystem.

Species composition Each population is represented by a GEM of a species. 15 different metabolic models are used in our spatially explicit model (Table 1), selected based on previous research ([30]). From this list of genera, the most prevalent species within that genus in the vaginally born newborn data set from Bäckhed *et al.* was used ([11]). One group from the list could only be determined at the family level - Lachnospiraceae. *Ruminococcus gnavus* was chosen to represent this group, based on its high prevalence among this group and the prior inclusion of species from the *Blautia* and *Dorea* genera ([69]). Because the genus *Bifidobacterium* is known to be particularly diverse ([70]), we represented it with models of three different strains. All models are based on the GEMs created by Magnúsdóttir et al. in the AGORA collection ([23]).

Changes from AGORA We use updated versions of the AGORA GEMs ([74]), to which we have applied checks and modifications. Firstly, the objective function was changed from the biomass reaction included in the models to a reaction only requiring ATP production. As this reaction yields only ADP and P_i, it is mass-neutral. This allows us to focus on carbon dissimilation within the GEMs, and the differences in underlying metabolism. ATP yield has been a good proxy for biomass production in previous studies ([75]). Focusing on carbon dissimilation means we can leave all unknown uptake bounds at 0, instead of using an arbitrary or randomized level, as in some other studies ([17], [16]).

We also checked the metabolic networks for reactions that allow for the occurrence of unrealistic FBA solutions, and we have added additional reactions. In the *Bifidobacterium* models the ATP synthase reaction was made reversible. This allows it to function as a proton pump, which *Bifidobacterium* species use to maintain their internal pH ([76]). Lactose permease reactions were added to all *Bifidobacterium* species in the model ([77]) based on those available in other models in the AGORA collection ([23]). Combined with the existing reactions in the model and the metabolic constraints this leads to a set of reactions simulating a realistic bifid shunt yield, when substrate is abundant, of 5 mol ATP, 2 mol lactate and 3 mol acetate per mol of lactose([38], [78]). Lactose permease was also added to *Streptococcus salivarius* and *Ruminococcus gnavus* to bring them in line with existing literature on their *in vitro* behavior ([79], [80]). *Veillonella dispar* is the only species in the model that does not have any lactose uptake ([71]). A complete list of changes is presented in S1 Table.

Checking the validity of the GEMs After the changes in S1 Table were applied all GEMs used in the model were tested individually to ensure that they could grow on a substrate of lactose. Only *Veillonella* did not pass this test, which is consistent with *in vitro* observations ([71]). *Veillonella* did pass when lactate, a common infant gut metabolite, was used instead. We also tested all GEMs individually for spurious growth in absence of substrates. To this end, all uptake bounds except water were set to 0. None of the GEMs grew under these conditions. All models were tested for having a net neutral exchange of hydrogen, carbon, oxygen and nitrogen at each timestep of the model during the simulations. There should be no net change in the number of atoms in the medium due to the calculated fluxes, because the reaction we set as an objective function does not remove any atoms, and there are no other sinks in the simulations. In the present simulation, only water, oxygen, and lactose are introduced into the simulation, so no other atoms than these three are considered. The tests revealed two errors in glycogen metabolism in several GEMs in AGORA that resulted in a energetically-favorable removal of intracellular protons from the system. We corrected these GEMs by replacing the reactions responsible for this erroneous energy source (S1 Table). With the corrections we applied to the model, these tests always passed - allowing for rounding errors of less than $1 \cdot 10^{-8}$ μmol per FBA solution.

The thermodynamic correctness of all reactions was checked by calculating the net difference in Gibbs free energy between input and output metabolites, using a pre-existing dataset of thermodynamic source data ([72], [73]). The conditions assumed for the calculation of the Gibbs free energy were a pH of 7 and an ionic strength of 0.1M ([72]). This was recorded for each population at each timestep over the course of a full run of 10080 timesteps. 99.999% of all FBA solutions in the simulations of Fig. 3 passed the test by containing less free energy than the associated inputs. The remaining 0.0001% all had an amount of energy in the outputs equal to that of the inputs. All of these solutions had very low growth rates, less than 0.001% of the average growth rate. The sum of these growth rates over 30 simulations was less than 0.05 of an initial

population, totalling $2 \cdot 10^6$ bacteria.

FBA with enzymatic constraint Each timestep of the model, a modified version of flux balance analysis with an enzymatic constraint is used by each population to determine what reactions it should use to achieve the most biomass production from the metabolites available to it ([81], [20]). First, each GEM is converted to a stoichiometric matrix S. All reversible reactions are converted to two irreversible reactions, so that flux is always greater than or equal to 0. All reactions identified as exchange, sink, or demand in the metabolic reconstruction are marked as exchange in the matrix. These reactions exchange nutrients or metabolites with the environment. Each timestep, all reactions are assumed to be in internal steady state ([81]):

$$S \cdot \vec{f} = 0 \tag{1}$$

Where $\vec{f}$ is a vector of the metabolic fluxes through each reaction in the network (in mol per time unit per population unit).

Each exchange reaction that takes up metabolites from the environment ($F_{in}(m)$) is constrained by an upper bound $F_{ub}(m)$ which represents the limited availability of most nutrients from the environment:

$$\vec{F_{in}}(\vec{m}) \leq \vec{F_{ub}}(\vec{m}) \tag{2}$$

Where $\vec{F_{in}}$ is a vector of fluxes between the environment and the bacterial population (in mol per time unit per population unit) and $\vec{m}$ is a vector of all metabolites. $\vec{F_{ub}}(\vec{m})$ is set dynamically at each timestep t by the spatial environment (discussed in detail below) at each lattice site x:

$$\vec{F_{ub}}(\vec{x}, \vec{m}, t) = \frac{\vec{c}(\vec{x}, \vec{m}, t)}{B(\vec{x}, t)} \tag{3}$$

Where $\vec{c}$ is a vector of all metabolite concentrations in mol per lattice site, $\vec{x}$ is a vector of all lattice sites and $B(\vec{x})$ is the size of the local bacterial population in population units. Population units are continuous, the size of B can range from $5 \cdot 10^7$ to $2 \cdot 10^{10}$, or $4 \cdot 10^{10}$ when division is not possible.

There is an additional constraint on the total flux. This constraint represents the limited amount of metabolism that can be performed per cell in each timestep. Each cell can only contain a limited number of enzymes, and each enzyme can only perform a limited number of reactions in limited time. Not limiting flux would lead to an unrealistic situation. The enzymatic constraint uses the sum of all fluxes $\vec{f}$, not just those that exchange with the environment:

$$\sum \vec{f} \leq a \tag{4}$$

The enzymatic constraint a is in mol per time unit per population unit (Table [3]). As both $\vec{f}$ and a are per population unit, this limit scales with population size, allowing each bacterium to contribute equally and independently to the metabolic flux attained in a lattice site. The enzymatic constraint is included as a constraint on each local FBA solution separately at each timestep. The enzymatic constraint was first proposed in a metabolic modelling context in 1990 ([20]), adapted from the study of other capacitated flow networks. In our context the enzymatic constraint represents the limited availability of physical space for enzymes. Our method is similar to FBA with molecular crowding in that we use an additional constraint to model metabolic capacity ([21]). However, we do not utilize the necessarily reaction-specific crowding coefficients.

FBA then identifies the solution space that adheres to these constraints and from
this spaces identifies the solution that optimizes the objective function. The set of so-
lutions consists of a set of exchange fluxes $\vec{F}(\vec{x}, \vec{m}, t)$, and a growth rate $\vec{g}(\vec{x}, t)$. These
exchange fluxes are taken as the derivatives of a set of partial-differential equations
to model the transport of intermediary metabolites (see below). The size of the popu-
lation increases in proportion to the growth rate produced by the solution.

Environmental metabolites We modeled a set of 723 extracellular metabolites
- the union of all metabolites that can be exchanged with the environment by at least
one GEM in the model. In practice, only a fraction of the metabolites occur in any ex-
change reaction that has flux over it, and only 17 metabolites are ever present in the
medium in more than micromolar amounts in our simulations outside of the sensitiv-
ity analysis (Table 2). Though the model distinguishes between L-lactate and D-lactate,
we display them together in our figures. Nearly all lactate produced and consumed in
our model is L-lactate.

TABLE 2 Extracellular metabolites present in the model in more than micromolar amounts

Common name	BiGG ID
Acetate	ac
Alpha-Ketoglutaric acid	akg
Carbon dioxide	co2
Ethanol	etoh
Formate	for
Hexadecanoate	hdca
Hydrogen	h2
L-Lactate	lac_L
D-Lactate	lac_D
Lactose	lcts
Octadecanoate	ocdca
Oxygen	o2
Propionate	ppa
Proton	h
Succinate	succ
Tetradecanoate	ttdca
Water	h2o

To mimic advection, the complete metabolic contents of the lattice except oxygen
are moved towards the distal end by 2mm (one lattice site with the default param-
eters) per timestep, i.e. once every 3 simulated minutes. This leads to an average
transit time of approximately 11 hours, in agreement with the observed cecum to rec-
tum transit time in newborns (32, 33). Metabolites moving out of the distal end are
removed entirely and analysed separately (see Section ' Analysis' for detail).
Every timestep all metabolites diffuse to the four nearest neighbours $NB(\vec{x})$ at
an equal rate for all metabolites (Table 3). We used a baseline of $4.4 \cdot 10^{-5}$ cm^2/s to
represent mixing of metabolites, which is an order of magnitude higher than normal
diffusion for common metabolites (82) to represent active mixing due to colonic con-
tractions. Metabolites are also added and removed by bacterial populations as a result
of the FBA solutions, yielding

$$\frac{d\vec{c}(\vec{x}, \vec{m}, t)}{dt} = \vec{F_{out}}(\vec{x}, \vec{m}, t)B(\vec{x}, t) - \vec{F_{in}}(\vec{x}, \vec{m}, t)B(\vec{x}, t) + \frac{D}{L^2} \sum_{\vec{i} \in NB(\vec{x})} \left(\vec{c}(\vec{i}, \vec{m}, t) - \vec{c}(\vec{x}, \vec{m}, t) \right).$$

$\hspace{12cm}$ (5)

Where F_{out} is a vector of fluxes from the bacterial populations to the environment,
in mol per time unit per population unit.

All lattice sites initially contain water as their only metabolite, except for the con-
ditions where oxygen is added. Metabolites representing the food intake are inserted
into the first six columns of lattice sites every three hours (60 timesteps) to approxi-
mate a realistic interval for neonates (83). In the current model this food intake con-
sists solely of lactose, in a concentration in line with human milk (84), assuming 98%
host uptake of carbohydrates before reaching the colon, a commonly used assump-
tion (16). Water is provided as a metabolite in unlimited quantities. Oxygen is placed
evenly distributed or at the upper and lower boundaries in some simulations. No
other metabolites are available, other than those produced as a result of bacterial
metabolism within the model.

Population dynamics Each local population solves the FBA problem based on its
own GEM, an enzymatic constraint a, its current population size $B(\vec{x}, t)$ and the local
concentrations of metabolites $\vec{c}(\vec{m}, t)$ at each timestep, and applies the outcome to
the environment (see above) and its own population size, as follows:

$$\frac{d\vec{B}(\vec{x}, t)}{dt} = \vec{B}(\vec{x}, t) \cdot \vec{g}(\vec{x}, t) \tag{6}$$

Populations at least 200 times the initial size (Table 3) will create a new population
in one empty adjacent lattice site, if possible. Half of the old population size is trans-
ferred to the new population, in such a way that the total size is preserved. To mimic
colonization events new populations are introduced at random into empty lattice sites
during the simulation, representing both dormant bacteria from intestinal crypts (34)
and small bacterial populations that are formed from ingested bacteria, which may
only become active after having diffused far into the gut. Each empty lattice has a
probability of 0.00005 (Table 3) each step to acquire a new population of a randomly
selected species, as follows:

for $site \in lattice sites$ **do**
if $site == empty$ **then**
813   **if** $random number \leq placement probability$ **then**
816   Place metabolic network of random species at $site$

The probability is scaled by $\frac{1}{n^2}$, with n the scaling factor of Δx, the side length of a lattice
site. There is an equal probability for any species in the model to be selected. As we
consider these new populations to be new colonizers we initialize them at the same
population size B as the initial populations in the model (Table 3). Each population
dies out at a probability of 0.0075 per timestep, creating a turnover within the range
of estimated microbial turnover rates in the mouse microbiota (85).

To mimic the mixing of bacterial populations, the lattice sites swap population con-
tents each timestep. We use the following algorithm, inspired by Kawasaki dynamics
(86), as also used previously for bacterial mixing (18, 87): In random order, the content
of each site, i.e., the bacterial population represented by its size and the GEM but not
the metabolites, are swapped with a site randomly selected from the set consisting
of the site itself and the first and second order neighbours. This swap only occurs if
both the origin and destination site have not already swapped. Bacterial populations
at the most distal column, i.e. at the exit of the colon, are deleted from the system.
With this mixing method the diffusion constant of the bacterial populations becomes
$5.7 \cdot 10^{-5}$ cm^2/s (Table 3). In the simulations that use a finer or a coarser grid (Fig. 6D)
the number of swaps is scaled as $\frac{1}{n^2}$, with n the scaling factor of Δx, the side length of
a lattice site, thus maintaining the same diffusion rate. For $n < 1$ all sites are marked

as unswapped once all sites have attempted a swap. This allows for sites to swap mul-
tiple times. The same approach was used to change the bacterial diffusion rate in the
sensitivity analysis (Fig. 6F). To achieve full mixing all bacteria were assigned random
non-overlapping locations at every timestep. Note that the diffusion rate of bacterial
populations with the default parameters (Table 3) is 1.3 times higher than that of the
metabolites. As shown by the sensitivity analysis (Fig. 6F) this has little effect on the
results.

Initial conditions The simulation is initialized by placing a number of very small
populations (B) of the various species randomly across different lattice sites of the
environment (Table 3). There is a probability of 0.3 for each lattice site to acquire a
population, an average of 540 for our lattice. The size of initial populations is scaled to
be roughly equivalent to a plausible initial total load of approximately $3 \cdot 10^{10}$(88), as-
suming a total colon volume of approximately 90 ml. As there is little information on
the relative abundance of species in the very early infant gut, we place all species with
equal probability. In initially oxygenated conditions, oxygen is also placed as a metabo-
lite now. Water is always considered to be present everywhere. No other metabolites
are initially present except for the first feeding.

The ratio of population to bacterial abundance was chosen as a balance between
a high bacterial resolution and computational complexity, as the FBA solution of each
population is calculated independently per timestep. This resolution splits the infant
microbiota into approximately 1-600 units (depending on conditions), allowing for the
presence of species at relatively low abundances, and spatial variation. There is no
hard limit on the number of populations in the model - it is limited only by the balance
between growth (through consumption) and death.

Analysis At each timestep we record the location and exchange fluxes $\vec{F}(\vec{x}, t)$ of
major metabolites as well as the size $B(\vec{x}, t)$ and species of all populations. This is used
to analyze both population composition and metabolic fluxes over time and space. In
addition, each timestep we record the location and quantity $\vec{c}(\vec{x}, t)$ of all metabolites
present in more than micromolar quantities. We also record this metabolite data for
all metabolites exiting the system at the distal side. This is considered the closest
equivalent to a fecal composition in the model, and these results are compared to
data from *in vivo* fecal samples.

To detect any irregularities, we also record the net carbon, hydrogen, and oxygen
flux of every population and the system as a whole. The difference in Gibbs free energy
per timestep is also recorded per for each population per FBA solution, and seperately
over the whole system. Estimated Gibbs free energy is derived from the Equillibrator
database (73). Energy loss I in joules per timestep per population unit is recorded as
follows, where m are metabolites, F is the exchange fluxes in mol per population unit,
B is the population size and E contains the Gibbs free energy in joules per mol for
each metabolite,

$$I = \sum_m F(m) \cdot E(m) \tag{7}$$

For specific simulations, reactions are removed from the models. This is per-
formed by deleting some reactions from the GEM before the conversion to the sto-
ichiometric matrix. To remove Fructose-6-phosphate phosphoketolase, we removed
the reactions R_PKL and R_F6PE4PL. For other simulations, the uptake of certain metabo-
lites is disabled. This is done by placing the upper flux bound $\vec{F}_{ub}$ of the relevant ex-
change reaction at 0 for the relevant populations.

Parameters Relevant parameters are listed in Table 3. Based on measurements of the typical length and diameter of the infant colon (26, 27) we estimated a volume of 90 ml. Combined with the average abundance per ml of around 10^{10} after the first days (88), this leads to a very rough estimate of 10^{12} bacteria in the young infant colon. To remain computationally feasible, while still modelling at a high resolution, we divide this population into units of at most $1 \cdot 10^{10}$. Local populations at or above this limit will divide into two equally sized populations when space is available. This prevents the local population from becoming unrealistically high. Local populations of more than $2 \cdot 10^{10}$, which can only form if no space is available to divide for a longer time period, cease metabolism.

TABLE 3 Default parameters of the model

Parameter	Value	Unit
Timestep	180	seconds
Lattice site side length (Δx)	2	millimetre
Width of the lattice	225	lattice sites
Height of the lattice	8	lattice sites
Colon transit time	11	hours
Average number of initial populations	540	-
Initial size per population	$5 \cdot 10^7$	number of bacteria
Population size required for division	$1 \cdot 10^{10}$	number of bacteria
Death probability	0.0075	per timestep per population
Enzymatic constraint	2	µmol flux per timestep per $1 \cdot 10^{10}$ population
New species placement probability	0.00005	per timestep per empty lattice site
Diffusion of metabolites	$4.4 \cdot 10^{-5}$	cm^2/s
Diffusion of populations	$5.7 \cdot 10^{-5}$	cm^2/s
Growth per µmol ATP	$1 \cdot 10^9$	number of bacteria
Lactose input	211	µmol per three hours

The lactose input is estimated from the known intake of milk, its lactose concentration, and an estimate of pre-colonic lactose absorption of 98% (89, 16). Little data is available on the growth rate of bacteria within the human colon. Growth rates are expected to be much lower than those found in *in vitro* cultures of individual species (90). In absence of precise data for infants, here we use a death probability that places the replacement rate within the range of estimated doubling times of the whole gut microbiota in mice (85). The colonic transit time is based on data for total transit time gathered with carmine red dye (32), adjusted for the mouth to cecum transit time (33). The timestep interval was set at three minutes, to be able to capture individual feedings at a high resolution. Other parameters were selected to reach plausible outcomes on a metabolic and species level while maintaining computational feasibility. We do not expect the parameters to match precisely with *in vivo* values, if these were measured.

Implementation The model was implemented in C++11. Our code was based on code used earlier to model the gut microbiota (18). Random numbers are generated with Knuth's subtractive random number generator algorithm. Diffusion of metabolites was implemented using the Forward Euler method. The GEMs are loaded using libSBML 5.18.0 for C++. We used the GNU Linear Programming Kit 4.65 (GLPK) as a linear programming tool to solve each FBA with enzymatic constraint problem. We used the May 2019 update of AGORA, the latest at time of writing, from the Virtual Metabolic Human Project website (vmh.life). Python 3.6 was used to extract thermo-dynamic data from the eQuilibrator API (December 2018 update) (73) and determine mean square displacement of our bacterial diffusion. Model screenshots are made us-

ing the libpng16 and pngwriter libraries. Other visualisations and statistical analysis
were performed with R 4.1.2 and Google Sheets.

Data availability The code used for the model is available from GitHub at
https://github.com/DMvers/IGMOST. The data files used for the model are available
from GitHub at https://github.com/DMvers/IGMOSTdatafiles.

SUPPLEMENTAL MATERIAL

S1 Figure. (A) Growth per timestep by lactose concentration for populations
of $5 \cdot 10^9$ **bacteria with access to one lattice site (0.05ml) of some major bacterial**
species with the enzymatic constraint disabled (B) Production of metabolites per
timestep by an *E. coli* population of $5 \cdot 10^9$ **bacteria with access to one lattice site**
(0.05ml) (C) Production of metabolites by a *B. vulgatus* population of $5 \cdot 10^9$ **bacteria**
with access to one lattice site (0.05ml). Formate and propionate formation are
always equal, and so are hydrogen and succinate. (D) Growth per timestep by lac-
tose concentration for populations of $5 \cdot 10^9$ **bacteria with access to one lattice site**
(0.05ml) of some major bacterial species with the enzymatic constraint disabled

S2 Figure. (A) Populations over time with bacteria in the "others" category of Fig. 5 left out. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (B,C) Spatial distribution of (B) lactose and (C) acetate over the last two days (960 timesteps) of the simulation. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (D) Populations over time with lactate uptake disabled for *Bifidobacterium* species. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (E) Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation in D. n=30. (F) Populations over time with lactate production disabled for *Bifidobacterium* species. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations.

S3 Figure. (A,B,C,D) Populations over time with (A) 0.001 (B) 0.01 (C) 0.1 (D) 1 μmol oxygen released in total per step divided over the top and bottom rows. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations per condition. (E) Populations over time with 1 μmol oxygen released per step initially (as in D), decreasing to 0 at step 5040. Abundances for grouped species over 21 days. One day is 480 timesteps. Curve shows mean value and shaded area one standard deviation over n=30 simulations per condition. (F) Distribution of total *Bifidobacterium* abundance at 21 days (10080 timesteps) corresponding to the simulations of E (G) Growth on unlimited lactose per timestep by oxygen concentration for populations of $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml) of some major bacterial species (H) Production of metabolites by lactose concentration in the presence of abundant oxygen for an *E. coli* population of $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml) (I) Production of metabolites by lactose concentration in the presence of abundant oxygen for an *E. cloacea* population of $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml) (J) Production of metabolites by glucose concentration in the presence of abun-

dant oxygen for an *E. coli* population of $5 \cdot 10^9$ bacteria with access to one lattice
site (0.05ml) (K) Populations over time with *K. pneumoniae* instead of *E. coli* and
0.1 µmol initial oxygen per lattice site. Abundances for grouped species. One
day is 480 timesteps. Curve shows mean value and shaded area one standard
deviation over n=30 simulations. (L) Populations over time with *K. pneumoniae*
in addition to the 15 species from table 1, and 0.1 µmol initial oxygen per lattice
site. Abundances for grouped species. One day is 480 timesteps. Curve shows
mean value and shaded area one standard deviation over n=30 simulations.

**S4 Figure. (A) Populations over time with new populations after the initial
populations only placed in the first column, with the placement probability (Ta-
ble 3) increased to 0.01125. Abundances for grouped species over 21 days. One
day is 480 timesteps. Curve shows mean value and shaded area one standard
deviation over n=30 simulations. (B,C,D) Spatial distribution of lactose, lactate,
and acetate, respectively, over the last two days (960 timesteps) of the simula-
tion with a larger lattice. Curve shows mean value and shaded area one stan-
dard deviation over n=30 simulations. (E) Abundance per location for grouped**

species over the last two days (960 timesteps) of the simulations with a larger lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (F,G,H) Spatial distribution of lactose, lactate, and acetate, respectively, over the last two days (960 timesteps) of the simulation with a smaller lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (I) Abundance per location for grouped species over the last two days (960 timesteps) of the simulations with a smaller lattice. Curve shows mean value and shaded area one standard deviation over n=30 simulations. (J) Distribution of metabolites exiting the system distally over the last two days (960 timesteps) of the simulation with diffusion of metabolites increased by a factor of 5. n=30.

S1 Table. Table of changed or deleted reactions and annotations.csv
A table of changes made to the AGORA models as a .csv file.

S2 Table Table of reactions used by *Bifidobacterium longum infantis* in model depending on lactose concentration
A table containing the flux through each reaction with more than nanomolar flux for *Bifidobacterium longum infantis* populations of $5 \cdot 10^9$ bacteria with access to one lattice site (0.05ml). Water was unlimited, no other metabolites were present. Lactose was present at 0.05, 0.35, or 0.5 µmol per ml.

S1 Video. Video of a run with no initial oxygen, consisting of a visualisation of the distribution of bacterial species and major metabolites, and a visualisation displaying fluxes between population and metabolite pools. Line width and intensity is proportional to the amount exchanged with the environment over the last 60 timesteps, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels.

S2 Video. Visualisations, from left to right, of a run from Fig. 4A, a run from Fig. 4B, and a run from Fig. 4F, displaying population and metabolite pool sizes, and fluxes between populations and metabolites. Line width and intensity is proportional to the amount exchanged with the environment over the last 60 timesteps, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels.

S3 Video. Video of a run with an initial 0.1 µmol of oxygen per lattice site, consisting of a visualisation of the distribution of bacterial species and major metabolites, and a visualisation displaying fluxes between population and metabolite pools. Line width and intensity is proportional to the amount exchanged with the environment over the last 60 timesteps, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels.

S4 Video. Visualisations, from left to right, of a run from Fig. 5 D, a run from Fig. 5 H, and a run from Fig. 6 A without the enzymatic constraint, displaying population and metabolite pool sizes, and fluxes between populations and metabolites. Line width and intensity is proportional to the amount exchanged with the environment over the last 60 timesteps, with a threshold of 0.5 µmol, with no normalization. Metabolite circle size is relative to the most abundant metabolite, with a minimum displayed size of 26 pixels.

CONTRIBUTIONS

P.V.J, M.P., J.M.W.G., and R.M.H.M acquired funding. D.M.V., P.V.J, M.P., J.M.W.G., and R.M.H.M. conceived and planned the simulations. D.M.V. and D.M. wrote software used for the simulations. D.M.V. performed the simulations and analyzed the data. R.S, E.L., J.M.W.G., and R.M.H.M contributed to the interpretation of the results. J.M.W.G., and R.M.H.M. supervised the project. D.M.V. drafted the manuscript. D.M.V., R.S., E.L., J.M.W.G. and R.M.H.M. revised and edited the manuscript.

ACKNOWLEDGMENTS

This study was financially supported by FrieslandCampina. R.S, E.L., P.V.J., M.P. and J.M.W.G. are currently or were previously employed by FrieslandCampina. The work was carried out in part on the Dutch national e-infrastructure with the support of SURF Cooperative. This work was performed in part using the ALICE compute resources provided by Leiden University.

REFERENCES

1. **Dogra S, Sakwinska O, Soh SE, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Karnani N, Lee YS, Yap F, Chong YS, Godfrey KM, Holbrook JD**. 2015. Rate of establishing the gut microbiota in infancy has consequences for future health. Gut Microbes 6 (5):321–325. doi:10.1080/19490976.2015.1078051.

2. **Dogra S, Sakwinska O, Soh SEE, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Lee YS, Yap F, Chong YSS, Godfrey KM, Holbrook JD, GUSTO Study Group, Sakwinska O, Ngom-Bru C, Brück WM, Berger B, Brüssow H, Soh SEE, Yap F, Godfrey KM, Holbrook JD**. 2015. Dynamics of infant gut microbiota are influenced by delivery mode and gestational duration and are associated with subsequent adiposity. mBio 6 (1):1–9. doi:10.1128/mBio.02419-14.

3. **Scholtens PAMJ, Oozeer R, Martin R, Amor KB, Knol J**. 2012. The Early Settlers: Intestinal Microbiology in Early Life. Annu Rev Food Sci Technol 3 (1):425–447. doi:10.1146/annurev-food-022811-101120.

4. **Stark PL, Lee A**. 1982. The microbial ecology of the large bowel of breast-fed and formula-fed infants during the first year of life. J Med Microbiol 15 (2):189–203. doi:10.1099/00222615-15-2-189.

5. **Matamoros S, Gras-Leguen C, Le Vacon F, Potel G, De La Cochetiere MF**. 2013. Development of intestinal microbiota in infants and its impact on health. Trends Microbiol 21 (4):167–173. doi:10.1016/j.tim.2012.12.001.

6. **Dotinga BM, Mintzer JP, Moore JE, Hulscher JBF, Bos AF, Kooi EMW**. 2020. Maturation of Intestinal Oxygenation: A Review of Mechanisms and Clinical Implications for Preterm Neonates. Front Pediatr 8:354. doi:10.3389/fped.2020.00354.

7. **Friedman ES, Bittinger K, Esipova TV, Hou L, Chau L, Jiang J, Mesaros C, Lund PJ, Liang X, FitzGerald GA, Goulian M, Lee D, Garcia BA, Blair IA, Vinogradov SA, Wu GD**. 2018. Microbes vs. chemistry in the origin of the anaerobic gut lumen. Proc Natl Acad Sci United States Am 115 (16):4170–4175. doi:10.1073/pnas.1718635115.

8. **Lay C, Chu CW, Purbojati RW, Acerbi E, Drautz-Moses DI, de Sessions PF, Jie S, Ho E, Kok YJ, Bi X, Chen S, Mak SY, Chua MC, Goh AEN, Chiang WC, Rao R, Chaithongwongwatthana S, Khemapech N, Chongsrisawat V, Martin R, Koh Y, Lohar SR, Tan ICH, Anng WA, Jie C, Bartke N, Ben-Amor K, Renes IB, Wong F, Roeselers G, Ho YS, Hibberd ML, Schuster SC, Knol J**. 2021. A synbiotic intervention modulates meta-omics signatures of gut redox potential and acidity in elective caesarean born infants. BMC Microbiol 21 (1):1–17. doi:10.1186/s12866-021-02230-1.

9. **Johnson JM, Adams ED**. 2022. The Gastrointestinal Microbiome in Infant Colic. MCN: The Am J Matern Nurs p 1–12. doi:10.1097/NMC.0000000000000832.

10. **Duar RM, Kyle D, Casaburi G**. 2020. Colonization Resistance in the Infant Gut: The Role of B. infantis in Reducing pH and Preventing Pathogen Growth. High-Throughput 9 (2):7. doi:10.3390/ht9020007.

11. **Bäckhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, Li Y, Xia Y, Xie H, Zhong H, Khan MT, Zhang J, Li J, Xiao L, Al-Aama J, Zhang D, Lee YS, Kotowska D, Colding C, Tremaroli V, Yin Y, Bergman S, Xu X, Madsen L, Kristiansen K, Dahlgren J, Jun W**. 2015. Dynamics and stabilization of the human gut microbiome during the first year of life. Cell Host Microbe 17 (5):690–703. doi:10.1016/j.chom.2015.04.004.

12. **van Best N, Hornef MW, Savelkoul PHM, Penders J**. 2015. On the origin of species: Factors shaping the establishment of infant's gut microbiota. Birth Defects Res Part C - Embryo Today: Rev 105 (4):240–251. doi:10.1002/bdrc.21113.

13. **Berger B, Porta N, Foata F, Grathwohl D, Delley M, Moine D, Charpagne A, Siegwald L, Descombes P, Alliet P, Puccio G, Steenhout P, Mercenier A, Sprenger N**. 2020. Linking human milk oligosaccharides, infant fecal community types, and later risk to require antibiotics. mBio 11 (2):1–18. doi:10.1128/mBio.03196-19.

14. **Francavilla R, Calasso M, Calace L, Siragusa S, Ndagijimana M, Vernocchi P, Brunetti L, Mancino G, Tedeschi G, Guerzoni E, Indrio F, Laghi L, Miniello VL, Gobbetti M, De Angelis M**. 2012. Effect of lactose on gut microbiota and metabolome of infants with cow's milk allergy. Pediatr Allergy Immunol 23 (5):420–427. doi:10.1111/j.1399-3038.2012.01286.x.

15. **Magnúsdóttir S, Thiele I**. 2018. Modeling metabolism of the human gut microbiome. Curr Opin Biotechnol 51:90–96. doi:10.1016/j.copbio.2017.12.005.

16. **Chan SHJ, Simons MN, Maranas CD**. 2017. SteadyCom: Predicting microbial abundances while ensuring community stability. PLoS Comput Biol 13 (5):1–25. doi:10.1371/journal.pcbi.1005539.

17. **Bauer E, Zimmermann J, Baldini F, Thiele I, Kaleta C**. 2017. BacArena: Individual-based metabolic modeling of heterogeneous microbes in complex communities. PLoS Comput Biol 13 (5):1–22. doi:10.1371/journal.pcbi.1005544.

18. **van Hoek MJA, Merks RMH**. 2017. Emergence of microbial diversity due to cross-feeding interactions in a spatial model of gut microbial

metabolism. BMC Syst Biol 11 (1):1–18. doi:10.1186/s12918-017-0430-4.

19. **Dukovski I, Bajić D, Chacón JM, Quintin M, Vila JCC, Sulheim S, Pacheco AR, Bernstein DB, Riehl WJ, Korolev KS, Sanchez A, Harcombe WR, Segrè D**. 2021. A metabolic modeling platform for the computation of microbial ecosystems in time and space (COMETS). Nat Protoc doi:10.1038/s41596-021-00593-3.

20. **Majewski RA, Domach MM**. 1990. Simple constrained-optimization view of acetate overflow in E. coli. Biotechnol Bioeng 35 (7):732–738. doi:10.1002/bit.260350711.

21. **Beg QK, Vazquez A, Ernst J, De Menezes MA, Bar-Joseph Z, Barabási AL, Oltvai ZN**. 2007. Intracellular crowding defines the mode and sequence of substrate uptake by Escherichia coli and constrains its metabolic activity. Proc Natl Acad Sci United States Am 104 (31):12663–12668. doi:10.1073/pnas.0609845104.

22. **de Groot DH, Lischke J, Muolo R, Planqué R, Bruggeman FJ, Teusink B**. 2020. The common message of constraint-based optimization approaches: overflow metabolism is caused by two growth-limiting constraints. Cell Mol Life Sci 77 (3):441–453. doi:10.1007/s00018-019-03380-2.

23. **Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, Greenhalgh K, Jäger C, Baginska J, Wilmes P, Fleming RMT, Thiele I**. 2017. Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota. Nat Biotechnol 35 (1):81–89. doi:10.1038/nbt.3703.

24. **Medina DA, Pinto F, Ovalle A, Thomson P, Garrido D**. 2017. Prebiotics mediate microbial interactions in a consortium of the infant gut microbiome. Int J Mol Sci 18 (10):1–16. doi:10.3390/ijms18102095.

25. **Monteagudo-Mera A, Arthur JC, Jobin C, Keku T, Bruno-Barcena JM, Azcarate-Peril MA**. 2016. High purity galacto-oligosaccharides enhance specific Bifidobacterium species and their metabolic activity in the mouse gut microbiome. Benef Microbes 7 (2):247–264. doi:10.3920/BM2015.0114.

26. **Valentin J, Streffer C**. 2002. Basic anatomical and physiological data for use in radiological protection: Reference values - ICRP Publication 89. Ann ICRP 32 (3-4):1–277. doi:10.1016/S0146-6453(03)00002-2.

27. **Davis WS, Parker Allen R, Favara BE, Slovis TL**. 1974. Neonatal small left colon syndrome. Am journal Roentgenol 120 (2):322–329. doi:10.2214/ajr.120.2.322.

28. **Swidsinski A, Loening-Baucke V, Verstraelen H, Osowska S, Doerffel Y**. 2008. Biostructure of Fecal Microbiota in Healthy Subjects and Patients With Chronic Idiopathic Diarrhea. Gastroenterology 135 (2):568–579. doi:10.1053/j.gastro.2008.04.017.

29. **Hong PY, Croix JA, Greenberg E, Gaskins HR, Mackie RI**. 2011. Pyrosequencing-based analysis of the mucosal microbiota in healthy individuals reveals ubiquitous bacterial groups and micro-heterogeneity. PLoS ONE 6 (9). doi:10.1371/journal.pone.0025042.

30. **Delsing D, Lindner C, van Dijck H, Looijesteijn E, Lima ES, Heerikhuizen M, Agamennone V, Triantis V, Bovee-Oudenhoven I**. 2019. Gut microbiota modulating effects of GOS and 2'-FL studied by in vitro infant and toddler faecal fermentations. ESPGHAN Conf. Abstr. N-P-006 .

31. **Schuetz R, Kuepfer L, Sauer U**. 2007. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. Mol Syst Biol 3 (119). doi:10.1038/msb4100162.

32. **Rubaltelli FF, Largajolli G**. 1973. Effect of Light Exposure on Gut Transit Time in Jaundiced Newborns. Acta Pædiatrica 62 (2):146–148. doi:10.1111/j.1651-2227.1973.tb08082.x.

33. **Hyams JS, Geertsma MA, Etienne NL, Treem WR**. 1989. Colonic hydrogen production in infants with colic. The J Pediatr 115 (4):592–594. doi:10.1016/S0022-3476(89)80289-6.

34. **Lee SM, Donaldson GP, Mikulski Z, Boyajian S, Ley K, Mazmanian SK**. 2013. Bacterial colonization factors control specificity and stability of the gut microbiota. Nature 501 (7467):426–429. doi:10.1038/nature12447.

35. **van Hoek MJA, Merks RMH**. 2012. Redox balance is key to explaining full vs. partial switching to low-yield metabolism. BMC Syst Biol 6. doi:10.1186/1752-0509-6-22.

36. **Vazquez A, Oltvai ZN**. 2016. Macromolecular crowding explains overflow metabolism in cells. Sci Reports 6:1–7. doi:10.1038/srep31007.

37. **de Vries W, Stouthamer AH**. 1968. Fermentation of glucose, lactose, galactose, mannitol, and xylose by bifidobacteria. J Bacteriol 96 (2):472–478. doi:10.1128/jb.96.2.472-478.1968.

38. **De Vuyst L, Moens F, Selak M, Rivière A, Leroy F**. 2014. Summer Meeting 2013: Growth and physiology of bifidobacteria. J Appl Microbiol 116 (3):477–491. doi:10.1111/jam.12415.

39. **Kotarski SF, Salyers AA**. 1981. Effect of long generation times on growth of Bacteroides thetaiotaomicron in carbohydrate-limited continuous culture. J Bacteriol 146 (3):853–860. doi:10.1128/jb.146.3.853-860.1981.

40. **Martin FPJ, Moco S, Montoliu I, Collino S, Da Silva L, Rezzi S, Prieto R, Kussmann M, Inostroza J, Steenhout P**. 2014. Impact of breast-feeding and high-and low-protein formula on the metabolism and growth of infants from overweight and obese mothers. Pediatr Res 75 (4):535–543. doi:10.1038/pr.2013.250.

41. **Tsukuda N, Yahagi K, Hara T, Watanabe Y, Matsumoto H, Mori H, Higashi K, Tsuji H, Matsumoto S, Kurokawa K, Matsuki T**. 2021. Key bacterial taxa and metabolic pathways affecting gut short-chain fatty acid profiles in early life. ISME J doi:10.1038/s41396-021-00937-7.

42. **Schwab C, Ruscheweyh HJ, Bunesova V, Pham VT, Beerenwinkel N, Lacroix C**. 2017. Trophic interactions of infant bifidobacteria and eubacterium hallii during L-fucose and fucosyllactose degradation. Front Microbiol 8 (JAN):1–14. doi:10.3389/fmicb.2017.00095.

43. **Wolin MJ, Yerry S, Miller TL, Zhang Y, Bank S**. 1998. Changes in Production of Ethanol, Acids and H2 from Glucose by the Fecal Flora of a 16- to 158-d-Old Breast-Fed Infant. The J Nutr 128 (1):85–90. doi:10.1093/jn/128.1.85.

44. **Bridgman SL, Azad MB, Field CJ, Haqq AM, Becker AB, Mandhane PJ, Subbarao P, Turvey SE, Sears MR, Scott JA, Wishart DS, Kozyrskyj AL**. 2017. Fecal Short-Chain Fatty Acid Variations by Breastfeeding Status in Infants at 4 Months: Differences in Relative versus Absolute Concentrations. Front Nutr 4 (April). doi:10.3389/fnut.2017.00011.

45. **de Vries W, Stouthamer AH**. 1967. Pathway of glucose fermentation in relation to the taxonomy of bifidobacteria. J bacteriology 93 (2):574–576. doi:10.1128/jb.93.2.574-576.1967.

46. **Kennedy KM, Gerlach MJ, Adam T, Heimesaat MM, Rossi L, Surette MG, Sloboda DM, Braun T**. 2021. Fetal meconium does not have a detectable microbiota before birth. Nat Microbiol 6 (7):865–873. doi:10.1038/s41564-021-00904-0.

47. **Albenberg L, Esipova TV, Judge CP, Bittinger K, Chen J, Laughlin A, Grunberg S, Baldassano RN, Lewis JD, Li H, Thom SR, Bushman FD, Vinogradov SA, Wu GD**. 2014. Correlation between intraluminal oxygen gradient and radial partitioning of intestinal microbiota. Gastroenterology 147 (5):1055–1063.e8. doi:10.1053/j.gastro.2014.07.020.

48. **Zheng L, Kelly CJ, Colgan SP**. 2015. Physiologic hypoxia and oxygen homeostasis in the healthy intestine. A review in the theme: Cellular responses to hypoxia. Am J Physiol - Cell Physiol 309 (6):C350–C360. doi:10.1152/ajpcell.00191.2015.

49. **Matsuki T, Yahagi K, Mori H, Matsumoto H, Hara T, Tajima S, Ogawa E, Kodama H, Yamamoto K, Yamada T, Matsumoto S,**

Kurokawa K. 2016. A key genetic factor for fucosyllactose utilization affects infant gut microbiota development. Nat Commun 7 (May):1–12. doi:10.1038/ncomms11939.

50. **Sim K, Shaw AG, Randell P, Cox MJ, McClure ZE, Li MS, Haddad M, Langford PR, Cookson WOCM, Moffatt MF, Kroll JS**. 2015. Dysbiosis anticipating necrotizing enterocolitis in very premature infants. Clin Infect Dis 60 (3):389–397. doi:10.1093/cid/ciu822.

51. **Pham VT, Lacroix C, Braegger CP, Chassard C**. 2016. Early colonization of functional groups of microbes in the infant gut. Environ microbiology 18 (7):2246–2258. doi:10.1111/1462-2920.13316.

52. **Mori M, Hwa T, Martin OC, De Martino A, Marinari E**. 2016. Constrained Allocation Flux Balance Analysis. PLoS Comput Biol 12 (6):1–24. doi:10.1371/journal.pcbi.1004913.

53. **Balakrishnan R, Silva RT, Hwa T, Cremer J**. 2021. Suboptimal resource allocation in changing environments constrains response and growth in bacteria. Mol Syst Biol 17 (12):1–13. doi:10.15252/msb.202110597.

54. **Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, Lovley DR**. 2011. Genome-scale dynamic modeling of the competition between Rhodoferax and Geobacter in anoxic subsurface environments. ISME J 5 (2):305–316. doi:10.1038/ismej.2010.117.

55. **Stansbridge EM, Walker V, Hall MA, Smith SL, Millar MR, Bacon C, Chen S**. 1993. Effects of feeding premature infants with Lactobacillus GG on gut fermentation. Arch Dis Child 69 (5 SPEC NO):488–492. doi:10.1136/adc.69.5_Spec_No.488.

56. **Falony G, Lazidou K, Verschaeren A, Weckx S, Maes D, De Vuyst L**. 2009. In vitro kinetic analysis of fermentation of prebiotic inulin-type fructans by Bifidobacterium species reveals four different phenotypes. Appl Environ Microbiol 75 (2):454–461. doi:10.1128/AEM.01488-08.

57. **Cocks GT, Aguilar J, Lin ECC**. 1974. Evolution of L-1,2 propanediol catabolism in Escherichia coli by recruitment of enzymes for L-fucose and L-lactate metabolism. J Bacteriol 118 (1):83–88. doi:10.1128/jb.118.1.83-88.1974.

58. **Pham VT, Lacroix C, Braegger CP, Chassard C**. 2017. Lactate-utilizing community is associated with gut microbiota dysbiosis in colicky infants. Sci Reports 7 (1):1–13. doi:10.1038/s41598-017-11509-1.

59. **Schwab C, Ruscheweyh HJ, Bunesova V, Pham VT, Beerenwinkel N, Lacroix C**. 2017. Trophic interactions of infant bifidobacteria and eubacterium hallii during L-fucose and fucosyllactose degradation. Front Microbiol 8:1–14. doi:10.3389/fmicb.2017.00095.

60. **Cremer J, Arnoldini M, Hwa T**. 2017. Effect of water flow and chemical environment on microbiota growth and composition in the human colon. Proc Natl Acad Sci United States Am 114 (25):6438–6443. doi:10.1073/pnas.1619598114.

61. **Sarna SK**. 2010. Colonic Motility From Bench Side to Bedside. Morgan&Claypool Life Sciences, San Rafael(CA). doi:10.4199/C00020ED1V01Y201011ISP011.

62. **Nurko S**. 2006. Motility of the Colon and Anorectum. NeoReviews 7 (1):e34–e48. doi:10.1542/neo.7-1-e34.

63. **Juntunen M, Kirjavainen PV, Ouwehand AC, Salminen SJ, Isolauri E**. 2001. Adherence of probiotic bacteria to human intestinal mucus in healthy infants and during rotavirus infection. Clin Diagn Lab Immunol 8 (2):293–296. doi:10.1128/CDLI.8.2.293-296.2001.

64. **Chan, Friedman, Wu, Maranas**. 2019. Predicting the Longitudinally and Radially Varying Microbial Metabolic Modeling. Processes 7:394. doi:10.3390/pr7070394.

65. **Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, Bonilla G, Kar A, Leiby N, Mehta P, Marx CJ, Segrè D**. 2014. Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. Cell Reports 7 (4):1104–1115. doi:10.1016/j.celrep.2014.03.070.

66. **Varma A, Palsson BØ**. 1994. Metabolic Flux Balancing: Basic Concepts, Scientific and Practical Use. Biotechnology 12 (October):994–998. doi:10.1038/nbt1094-994.

67. **Arnoldini M, Cremer J, Hwa T**. 2018. Bacterial growth, flow, and mixing shape human gut microbiota density and composition. Gut microbes 9 (6):559–566. doi:10.1080/19490976.2018.1448741.

68. **Cremer J, Segota I, Yang CY, Arnoldini M, Sauls JT, Zhang Z, Gutierreza E, Groisman A, Hwa T**. 2016. Effect of flow and peristaltic mixing on bacterial growth in a gut-like channel. Proc Natl Acad Sci United States Am 113 (41):11414–11419. doi:10.1073/pnas.1601306113.

69. **Sagheddu V, Patrone V, Miragoli F, Puglisi E, Morelli L**. 2016. Infant early gut colonization by Lachnospiraceae: High frequency of Ruminococcus gnavus. Front Pediatr 4:7–12. doi:10.3389/fped.2016.00057.

70. **Turroni F, Peano C, Pass DA, Foroni E, Severgnini M, Claesson MJ, Kerr C, Hourihane J, Murray D, Fuligni F, Gueimonde M, Margolles A, de Bellis G, O'Toole PW, van Sinderen D, Marchesi JR, Ventura M**. 2012. Diversity of bifidobacteria within the infant gut microbiota. PLoS ONE 7 (5):20–24. doi:10.1371/journal.pone.0036957.

71. **Rogosa M**. 1964. the Genus Veillonella. I. General Cultural, Ecological, and Biochemical Considerations. J Bacteriol 87:162–170. doi:10.1128/JB.87.1.162-170.1964.

72. **Flamholz A, Noor E, Bar-Even A, Milo R**. 2012. EQuilibrator - The biochemical thermodynamics calculator. Nucleic Acids Res 40 (D1):770–775. doi:10.1093/nar/gkr874.

73. **Noor E, Haraldsdóttir HS, Milo R, Fleming RMT**. 2013. Consistent Estimation of Gibbs Energy Using Component Contributions. PLoS Comput Biol 9 (7). doi:10.1371/journal.pcbi.1003098.

74. **Magnúsdóttir S, Heinken A, Fleming RMT, Thiele I**. 2018. Reply to "Challenges in modeling the human gut microbiome". Nat Biotechnol 36 (8):686–691. doi:10.1038/nbt.4212.

75. **Schuetz R, Kuepfer L, Sauer U**. 2007. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. Mol Syst Biol 3 (119). doi:10.1038/msb4100162.

76. **Sánchez B, De Los Reyes-Gavilán CG, Margolles A**. 2006. The F1F0-ATPase of Bifidobacterium animalis is involved in bile tolerance. Environ Microbiol 8 (10):1825–1833. doi:10.1111/j.1462-2920.2006.01067.x.

77. **Krzewinski F, Brassart C, Gavini F, Bouquelet S**. 1996. Characterization of the lactose transport system in the strain Bifidobacterium bifidum DSM 20082. Curr Microbiol 32 (6):301–307. doi:10.1007/s002849900054.

78. **Sela DA, Adeuya A, Kim JH, Whitehead TR, Chapman J, Rokhsar DS, Richardson PM, Lapidus A, German JB, Price NP, Chen F, Lebrilla CB, Mills DA**. 2008. The genome sequence of Bifidobacterium longum subsp. infantis reveals adaptations for milk utilization within the infant microbiome. Proc Natl Acad Sci 105 (48):18964–18969. doi:10.1073/pnas.0809584105.

79. **Lessard C, Cochu A, Lemay JD, Roy D, Vaillancourt K, Frenette M, Moineau S, Vadeboncoeur C**. 2003. Phosphorylation of Streptococcus salivarius Lactose Permease (LacS) by HPr(His~P) and HPr(Ser-P)(His~P) and Effects on Growth. J Bacteriol 185 (23):6764–6772. doi:10.1128/JB.185.23.6764-6772.2003.

80. **Holdeman LV**. 1976. Emendation of Bacteroidaceae and Butyrivibrio and Descriptions of Desulfornonas gen. nov. and Ten New Species in the Genera Desulfomonas, Butyrivibrio, Eubacterium, Clostridium, and Ruminococcus. Int J Syst Bacteriol 26 (2):238–252. doi:10.1099/00207713-26-2-238.

81. **Orth JD, Thiele I, Palsson BO**. 2010. What is flux balance analysis? Nat Biotechnol 28 (3):245–248. doi:10.1038/nbt.1614.

82. **Stewart PS**. 2003. Diffusion in biofilms. J Bacteriol 185 (5):1485–1491. doi:10.1128/JB.185.5.1485-1491.2003.

83. **Grummer-Strawn LM, Scanlon KS, Fein SB**. 2008. Infant feeding and feeding transitions during the first year of life. Pediatrics 122 (SUPPL. 2). doi:10.1542/peds.2008-1315d.

84. **Ballard O, Morrow AL**. 2013. Human Milk Composition. Nutrients and Bioactive Factors. Pediatr Clin North Am 60 (1):49–74. doi:10.1016/j.pcl.2012.10.002.

85. **Gibbons RJ, Kapsimalis B**. 1967. Estimates of the overall rate of growth of the intestinal microflora of hamsters, guinea pigs, and mice. J Bacteriol 93 (1):510–512. doi:10.1128/jb.93.1.510-512.1967.

86. **Kawasaki K**. 1966. Diffusion constants near the critical point for time-dependent ising models. I. Phys Rev 145 (1):224–230. doi:10.1103/PhysRev.145.224.

87. **Koumakis N, Devailly C, Poon WCK**. jun 2018. Motile bacteria in a critical fluid mixture. Phys Rev E 97 (6). doi:10.1103/PhysRevE.97.062604.

88. **Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO**. 2007. Development of the human infant intestinal microbiota. PLoS Biol 5 (7):1556–1573. doi:10.1371/journal.pbio.0050177.

89. **Anderson PO, Valdés V**. 2015. Variation of milk intake over time: Clinical and pharmacokinetic implications. Breastfeed Med 10 (3):142–144. doi:10.1089/bfm.2014.0170.

90. **Gibson B, Wilson DJ, Feil E, Eyre-Walker A**. 2018. The distribution of bacterial doubling times in the wild. Proc Royal Soc B: Biol Sci 285 (1880). doi:10.1098/rspb.2018.0789.