1

Self-assembly of pericentriolar material in

interphase cells lacking centrioles

Fangrui Chen[1], Jingchao Wu[1], Malina K. Iwanski[1], Daphne Jurriens[1], Arianna Sandron[1],

Milena Pasolli[1], Gianmarco Puma[1], Jannes Z. Kromhout[1], Chao Yang[1], Wilco Nijenhuis[1,2],

Lukas C. Kapitein[1,2], Florian Berger[1] and Anna Akhmanova[1]

[1] Cell Biology, Neurobiology and Biophysics, Department of Biology, Faculty of Science, Utrecht
University, Utrecht, The Netherlands

[2] Center for Living Technologies, Eindhoven-Wageningen-Utrecht Alliance, The Netherlands

Correspondence: a.akhmanova@uu.nl

Abstract

The major microtubule-organizing center (MTOC) in animal cells, the centrosome, comprises a pair of centrioles surrounded by pericentriolar material (PCM), which nucleates and anchors microtubules. Centrosome assembly depends on PCM binding to centrioles, PCM self-association and dynein-mediated PCM transport, but the self-assembly properties of PCM in interphase cells are poorly understood. Here, we used experiments and modeling to study centriole-independent features of interphase PCM assembly. We showed that when centrioles are lost due to PLK4 depletion or inhibition, dynein-based PCM transport and PCM self-clustering are sufficient to form a single compact MTOC, which generates a dense radial microtubule array. Interphase PCM self-assembly depends on γ-tubulin, pericentrin, CDK5RAP2 and ninein, but not NEDD1, CEP152 or CEP192. Formation of a compact acentriolar MTOC is inhibited by AKAP450-dependent PCM recruitment to the Golgi or by randomly organized CAMSAP2-stabilized microtubules, which keep PCM mobile and prevent its coalescence. Linking of CAMSAP2 to a minus-end-directed motor leads to the formation of an MTOC, but MTOC compaction requires cooperation with pericentrin-containing self-clustering PCM. Our data reveal that interphase PCM contains a set of components that can self-assemble into a compact structure and organize microtubules, but PCM self-organization is sensitive to motor- and microtubule-based rearrangement.

Introduction

The centrosome is the major microtubule organizing center (MTOC) in animal cells. It consists of two centrioles surrounded by pericentriolar material (PCM) (reviewed in (Conduit et al., 2015; Paz and Luders, 2017)). Major PCM components are microtubule-nucleating and anchoring proteins, which can associate with centrioles and with each other. For a long time it was thought that the PCM is amorphous, but super-resolution microscopy studies have shown that it has a distinct organization, with some proteins likely attached to the centriole wall and others organized around them (Fu and Glover, 2012; Lawo et al., 2012; Mennella et al., 2014; Mennella et al., 2012). This distinct organization is more obvious in interphase than in mitosis, when the microtubule-organizing capacity of the centrosome increases due to enhanced PCM recruitment. Many PCM components are known to oligomerize and interact with each other, and recent work suggested that the phase separation of interacting PCM components might contribute to centrosome assembly during mitosis (Raff, 2019; Woodruff et al., 2017). This idea is underscored by data showing that various cell-type-specific assemblies of PCM components can form clusters that are able to nucleate and organize microtubules and serve as MTOCs in the absence of centrioles, particularly during the formation of mitotic spindle poles (Balestra et al., 2021; Chinen et al., 2021; Gartenmann et al., 2020; Meitinger et al., 2020; Watanabe et al., 2020; Yeow et al., 2020). Furthermore, an important centrosome component, cytoplasmic dynein, is a motor that can bind to different PCM proteins and transport them to the centrosome-anchored microtubule ends, where these PCM proteins can nucleate and anchor additional microtubules, thus generating a positive feedback loop in centrosome assembly (Balczon et al., 1999; Burakov et al., 2008; Purohit et al., 1999; Redwine et al., 2017). Dynein and its mitotic binding partner NuMA also strongly participate in the formation of mitotic and meiotic spindle poles (Chinen et al., 2020; Khodjakov et al., 2000; Kolano et al., 2012). The relative importance of different molecular pathways of PCM assembly varies between cell systems and phases of the cell cycle and has not been explored systematically in interphase cells.

Here, we set out to investigate the centriole-independent self-assembly properties of interphase PCM. These properties, such as the ability of PCM to cluster or form molecular condensates, nucleate and anchor microtubules and move with motor proteins, are relevant because in most differentiated cell types, centrosome function is suppressed, and PCM forms acentrosomal MTOCs that are still poorly understood (Muroyama and Lechler, 2017; Sallee and Feldman, 2021). To study centriole-independent PCM function and dynamics, we removed centrioles using the PLK4 kinase inhibitor centrinone B which blocks centriole duplication (Wong et al., 2015). Importantly, in most commonly studied cultured cell lines, such as fibroblasts, epithelial, endothelial or cancer cells, microtubule networks are dense, and the centrosome is not the only MTOC. In such cells, non-centrosomal microtubule minus

ends are often stabilized by the members of CAMSAP family (Jiang et al., 2014; Meng et al., 2008;

Tanaka et al., 2012), and the Golgi apparatus serves as a second MTOC which nucleates and anchors

a very significant proportion of microtubules (Efimov et al., 2007; Rios, 2014; Wu et al., 2016; Zhu and

Kaverina, 2013). If centrosomes are lost because centriole duplication is blocked by inhibiting PLK4 or

depleting another essential centriole component, Golgi-dependent microtubule organization

becomes predominant (Gavilan et al., 2018; Martin et al., 2018; Wu et al., 2016). The ability of the

Golgi complex to serve as an MTOC critically depends on the Golgi adaptor AKAP450, which recruits

several PCM components that nucleate microtubules including the γ-tubulin ring complex (γ-TuRC),

CDK5RAP2 and pericentrin (Gavilan et al., 2018; Rivero et al., 2009; Wu et al., 2016). Moreover,

AKAP450 also tethers microtubule minus ends stabilized by CAMSAP2 to the Golgi membranes (Wu et

al., 2016). In the absence of AKAP450, the Golgi ribbon is maintained, but neither PCM components

nor CAMSAP-stabilized microtubule minus ends can be attached to the Golgi membranes and they are

instead dispersed in cytoplasm, leading to a randomly organized microtubule network (Gavilan et al.,

2018; Rivero et al., 2009; Wu et al., 2016). These data seem to suggest that centrioles and/or Golgi

membranes are essential to assemble PCM into an MTOC in interphase. However, this notion appears

to be inaccurate: in our previous study in RPE1 cells, we observed that a single compact acentriolar

MTOC (caMTOC) can still form after centriole loss in AKAP450 knockout cells, if the stabilization of

free minus ends in these cells is disabled by knocking out CAMSAP2 (Wu et al., 2016).

We used this observation as a starting point to investigate which properties of PCM allow it to self-

assemble in interphase mammalian cells and how the presence of non-centrosomal microtubules

affects this process. AKAP450 knockout cells provided a system to study PCM assembly in the absence

of competition with the Golgi-associated MTOC. caMTOC formation in AKAP450 knockout cells

required microtubules and depended on dynein, which brought together small PCM clusters with

attached minus ends. Experiments and modeling showed CAMSAP2-mediated minus-end stabilization

strongly perturbed PCM coalescence, because in the absence of AKAP450, randomly oriented

CAMSAP2-stabilized microtubules supported PCM motility and prevented PCM clustering. In the

absence of CAMSAP2, caMTOCs did form, but were often cylindrical rather than spherical in shape

and contained a subset of the major centrosome components. γ-tubulin, pericentrin, CDK5RAP2 and

ninein were necessary for the formation of caMTOCs, whereas some other major PCM proteins,

namely CEP192, CEP152 and NEDD1, were neither enriched in these structures nor required for their

formation, indicating that not all PCM components associate with each other in the absence of

centrioles. A single caMTOC containing PCM components could also form in the presence of CAMSAP2

when this protein was directly linked to a microtubule minus-end-directed motor. Importantly, in the

absence of pericentrin, minus-end-directed transport of CAMSAP2-stabilized minus ends organized

these ends into a ring, indicating that self-associating PCM is required for the formation of a caMTOC.

This conclusion was supported by modeling. Taken together, our data show that a subset of interphase

PCM components can self-assemble and efficiently nucleate and tether microtubules, but PCM

clustering is sensitive to microtubule- and motor-dependent rearrangements. These properties of

interphase PCM may also be involved in the transition from a centrosomal to a non-centrosomal

microtubule network, as typically occurs during cell differentiation.

Results

Assembly of microtubule-dependent caMTOCs in AKAP450/CAMSAP2 KO cells lacking PLK4 activity

To study centriole-independent PCM organization and dynamics in interphase cells, we induced centriole loss in RPE1 cells by treating them for 11 days with the PLK4 inhibitor centrinone B (Wong et al., 2015) (Figure 1A,B). In wild type (WT) cells, PCM (detected with antibodies against pericentrin) relocalized to the Golgi apparatus, and the microtubule array reorganized around the Golgi membranes, as described previously (Gavilan et al., 2018; Wu et al., 2016) (Figure 1C,D). In AKAP450 knockout cells, centriole loss led to the appearance of strongly dispersed PCM clusters, which could no longer bind to the Golgi, and a highly disorganized microtubule system, consistent with published work (Gavilan et al., 2018; Wu et al., 2016) (Figure 1D,E). In contrast, in AKAP450/CAMSAP2 double knockout cells, a single caMTOC with microtubules focused around it was observed (Figure 1D-F). Formation of a single caMTOC was also observed in AKAP450 knockout cells transiently depleted of CAMSAP2 by siRNA transfection (Figure 1 – figure supplement 1A,B). As an alternative approach to block the ability of the Golgi apparatus to recruit PCM, we treated the cells with Brefeldin A, which disrupts the Golgi (Klausner et al., 1992). As expected, Brefeldin A by itself had no effect on centrosome integrity but led to the dispersal of the Golgi marker GM130 (Figure 1 – figure supplement 1C-E). In centrinone-treated wild type cells, PCM was dispersed after Golgi disruption, and the same was true for the majority of CAMSAP2 knockout cells (Figure 1 – figure supplement 1C-F). However, in 12 % of CAMSAP2 knockout cells treated with both centrinone B and Brefeldin A, we observed PCM compaction (Figure 1 – figure supplement 1D-F), similar to that seen in AKAP450/CAMSAP2 double knockout cells. The efficiency of caMTOC formation was low likely because dispersed AKAP450 could still recruit PCM to some extent or because the duration of the Brefeldin A treatment (2 hrs) was too short to allow for PCM compaction in most cells (longer Brefeldin A treatments were not performed due to potential cell toxicity). The acentriolar PCM can thus form a compact structure if both the Golgi MTOC and CAMSAP2-mediated minus-end stabilization are disabled.

Unlike centrosomes, which always have a spherical shape, caMTOCs in AKAP450/CAMSAP2 knockout cells were cylindrical in ~35% of the cells, whereas in the remaining cells that lacked centrioles based on staining for centrin, MTOCs had a round shape (~38% of the cells); the rest of the centrin-negative cells either had dispersed PCM clusters (~7%) or no detectable PCM clusters (~11%) (Figure 1 – figure supplement 2A, B). In contrast, ~72% of acentriolar AKAP450 knockout cells had dispersed PCM, while caMTOCs were very rare (Figure 1E, Figure 1 – figure supplement 2A, B). Analysis by Stimulated Emission Depletion (STED) microscopy revealed that cylindrical caMTOCs in AKAP450/CAMSAP2 knockout cells consisted of small clusters of PCM components, including pericentrin, CDK5RAP2, γ-

tubulin, ninein and dynein heavy chain (Figure 1F). caMTOCs with a clearly elongated shape had an average length of ~11 μm and an average width of ~1.5 μm (Figure 1G).

To study PCM dynamics in caMTOCs, we generated cell lines stably expressing the PCM component CDK5RAP2 tagged with GFP. In control, untreated cells, GFP-CDK5RAP2 was localized to the centrosome and the Golgi apparatus as expected (Figure 1 – figure supplement 3A). In centrinone-treated AKAP450/CAMSAP2 knockout cells, it was strongly enriched within caMTOCs and sometimes also present in small motile clusters around a caMTOC (Figure 1H, Figure 1 – figure supplement 3B). Fluorescence recovery after photobleaching (FRAP) assays showed that when the whole caMTOC was bleached, the recovery was very slow and incomplete (Figure 1H, I). If only a part of a caMTOC was photobleached, the dynamics of recovery showed cell-to-cell variability. Highly condensed caMTOCs displayed a slow redistribution of GFP-CDK5RAP2 signal, suggesting that their components are largely immobile and do not rearrange. In more loosely organized caMTOCs, some rearrangement of small PCM clusters was observed (Figure 1H,I); however, the recovery was still far from complete. These data indicate that caMTOCs display variable degrees of compaction and are composed of PCM clusters that display limited exchange of GFP-CDK5RAP2 with the cytoplasmic pool, possibly because most of the GFP-CDK5RAP2 is accumulated within the caMTOC.

Next, we investigated whether centriole loss induced by means other than pharmacological PLK4 inhibition could also cause the formation of a single caMTOC in AKAP450/CAMSAP2 knockout cells. To achieve efficient protein depletion in RPE1 cells, they were transfected with siRNAs twice (on day 0 and day 2), treated with thymidine starting from day 4 to block cell cycle progression and fixed and stained on day 5 or day 7 (Figure 1 – figure supplement 2C). Depletion of PLK4 using siRNAs caused the appearance of round or cylindrical caMTOCs, similar to those observed after PLK4 inhibition with centrinone B, indicating that catalytically inactive PLK4 had no scaffolding role within these structures (Figure 1 – figure supplement 2C-F). The percentage of cells with caMTOCs increased over time (Figure 1 – figure supplement 2D), possibly due to the gradual depletion of PLK4. In contrast, depletion of CPAP, an essential centriole biogenesis factor (Kohlmaier et al., 2009; Schmidt et al., 2009; Tang et al., 2009), which also led to centriole loss, was much less efficient in inducing caMTOCs, and cylindrical caMTOCs were never observed (Figure 1 – figure supplement 2C-E). After CPAP depletion, cells in which pericentrin formed dispersed clusters or no visible clusters predominated (~67%, Figure 1 – figure supplement 2D, E). Treatment of CPAP-depleted AKAP450/CAMSAP2 knockout cells with centrinone B for 1 day promoted the assembly of round or cylindrical caMTOCs, and the proportion of such cells increased to ~55% after 3 days of centrinone B treatment (Figure 1 – figure supplement 2C-G). We also tested whether the inhibition of PLK1, a kinase that is known to be a major regulator of PCM self-assembly in mitosis (Haren et al., 2009; Joukov et al., 2014; Lee and Rhee, 2011), has an

effect on the formation of caMTOCs by treating cells with BI2536 (a highly selective and potent
inhibitor of PLK1), but found this not to be the case (Figure 1 – figure supplement 2C, D). We conclude
that PCM can assemble into a single stable MTOC in a centriole-independent manner if PLK4 is either
inactivated or depleted and the two major pathways of microtubule nucleation and minus-end
stabilization dependent on the Golgi membranes and CAMSAP2 are disabled.

Composition of caMTOCs and their effect on microtubule organization

PCM is composed of numerous proteins that can bind to each other and interact with microtubules,
and we next set out to investigate which PCM components can self-assemble in the absence of
centrioles. We first stained centrinone-treated AKAP450/CAMSAP2 knockout cells with antibodies
against different centrosome and centriole markers and microtubule-associated proteins (MAPs). As
indicated above, the abundant PCM components pericentrin, CDK5RAP2, ninein and γ-tubulin
colocalized within caMTOCs (Figure 1F). In contrast, three other major PCM proteins, CEP152, CEP192
and NEDD1, could not be detected in these structures although they were present in centrosomes of
AKAP450/CAMSAP2 knockout RPE1 cells that were not treated with centrinone B (Figure 2A-C) and
were also expressed in centrinone-treated cells (Figure 2 – figure supplement 1A). We then
individually depleted all these proteins in centrinone-treated AKAP450/CAMSAP2 knockout cells using
siRNAs. After the depletion of pericentrin, no clusters of other PCM components could be detected
(Figure 2D,J,K, Figure 2 – figure supplement 1B). To confirm this result, we also attempted to knock
out pericentrin in AKAP450/CAMSAP2 knockout cells, but such cells were not viable, likely because
centrosome defects in these cells caused prolonged mitosis and p53-dependent G1 arrest (Fong et al.,
2016; Lambrus et al., 2016; Meitinger et al., 2016). However, we were able to knock out pericentrin in
cells lacking AKAP450, CAMSAP2 and p53 (Figure 2 – figure supplement 2), confirming that the loss of
p53 makes the cells more tolerant to centrosome absence and allows them to divide even in the
presence of centrinone B (Figure 2 – figure supplement 3, Video 1). Similar to pericentrin-depleted
cells, these quadruple knockout cells were unable to form a single caMTOC when treated with
centrinone B (Figure 2E, J). In these acentriolar quadruple knockout cells, CDK5RAP2, γ-tubulin and
cytoplasmic dynein displayed no clustering, while ninein and PCM1, a centriolar satellite protein that
localizes closely around the centrosome in normal cells (Prosser and Pelletier, 2020), formed small
clusters distributed throughout the cytoplasm (Figure 2 – figure supplement 4A). Loss of pericentrin
had no effect on the expression of CDK5RAP2, γ-tubulin and ninein (Figure 2 – figure supplement 4B),
indicating that clustering by pericentrin affects the organization but not the stability of these PCM
components.

The depletion of CDK5RAP2, γ-tubulin or ninein in centrinone-treated AKAP450/CAMSAP2 knockout
cells did not prevent the formation of small pericentrin clusters, but these failed to coalesce into a
single caMTOC (Figure 2F-H, J, Figure 2 – figure supplement 1B). In contrast, the depletion of CEP152,
CEP192 or NEDD1 had no effect on the formation of caMTOCs (Figure 2A-C, J, Figure 2 – figure
supplement 1B), in agreement with the fact these proteins could not be detected within these
structures. caMTOCs contained several centriole biogenesis factors, including CPAP, CP110 and
CEP120, but lacked centrin and CEP135; however, the depletion of different centriolar proteins did
not affect caMTOC formation (Figure 2K, Figure 2 – figure supplement 1C). Within caMTOCs, we also
detected a component of the HAUS complex (HAUS2), the centrosomal protein CEP170, dynein,
dynactin, CLASP1/2, CLIP-115, CLIP-170, chTOG, KIF2A and KIF1C (Figure 2K, Figure 2 – figure
supplement 1C). We tested the importance of some of these proteins for caMTOC formation by siRNA-
mediated depletion (see Figure 2K for an overview), but among the tested proteins, only cytoplasmic
dynein appeared essential for this process. In dynein-depleted cells, no clusters of pericentrin or other
PCM components could be detected after centrinone B treatment (Figure 2I-K). It is important to note,
however, that because we used siRNAs to reduce protein expression, we cannot exclude that the
residual levels of some of the investigated proteins were sufficient to support caMTOC formation.
Because we detected several microtubule plus-end tracking proteins (+TIPs) in the caMTOCs, such as
CLIP-170, CLASP1/2 and the large subunit of dynactin, p150Glued, we also tested for the presence of
the core components of the +TIP machinery, EB1 and EB3, but found that they were not enriched
within the caMTOCs (Figure 2K, Figure 2 – supplement 5A). Using the previously described cells that
lack EB3, CAMSAP2 and the C-terminal partner-binding half of EB1 (EB1/EB3/CAMSAP2mut cells; Yang
et al., 2017), we generated a knockout cell line that also lacks AKAP450 and found that caMTOCs could
still form in these cells (Figure 2K, Figure 2 – figure supplement 5B,C). We conclude that a subset of
PCM components binds to each other in the absence of centrioles, and in AKAP450/CAMSAP2
knockout cells, these proteins form caMTOCs that recruit a number of additional PCM proteins and
MAPs normally present in interphase centrosomes.

Despite containing only a subset of centrosomal proteins, caMTOCs strongly affected the organization
and density of the microtubule network in acentriolar cells: microtubules were focused around
caMTOCs if present and disorganized in cells lacking caMTOCs. The strongest loss of microtubule
density was observed in cells lacking pericentrin, dynein or γ-tubulin, while milder phenotypes were
observed in cells lacking CDK5RAP2 or ninein (Figure 3A,B,E). To further characterize microtubule
organization after the loss of these proteins, we analyzed the proportion of the radial and non-radial
microtubules. Whereas control cells (AKAP450/CAMSAP2 knockout cells treated with centrinone B
and control siRNA) formed radial microtubule networks with ~12% non-radial microtubules,

acentriolar cells lacking pericentrin or cytoplasmic dynein had ~46% non-radial microtubules, and the
depletion of CDK5RAP2, ninein and γ-tubulin led to an intermediate phenotype with 25-30% non-
radial microtubules (Figure 3C,D,E). An acentriolar PCM assembly containing pericentrin, CDK5RAP2,
ninein, γ-tubulin and dynein is thus sufficient to form a radial microtubule array, similar to a
centrosome, and PCM clustering promotes dense microtubule organization.

Dynamics of caMTOC disassembly

To test whether the formation and maintenance of caMTOCs depends on microtubules, we
depolymerized them by treating cells with nocodazole at 37°C and found that caMTOCs fragmented
into small clusters upon nocodazole treatment and reassembled into a single structure after
nocodazole washout (Figure 4A-C). Because we found that caMTOC formation is dynein-dependent,
we also included the dynein inhibitor dynapyrazole A in these experiments (Steinman et al., 2017). We
confirmed that treatment with dynapyrazole A for 3 hrs had no effect on dynein expression (Figure
4D) and found that the addition of this drug before nocodazole treatment prevented the disassembly
of caMTOCs, whereas the treatment of cells with dynapyrazole A during nocodazole washout strongly
inhibited caMTOC re-assembly (Figure 4A-C). These data indicate that both microtubule-dependent
dispersal and coalescence of PCM clusters into caMTOCs are driven by dynein activity.

We next studied PCM dynamics using stably expressed GFP-CDK5RAP2 as a marker in live cells where
microtubules were labeled with SiR-tubulin. GFP-CDK5RAP2 was mostly immobile within caMTOCs
before nocodazole treatment (Figure 1 – figure supplement 3B, Video 2). After a few minutes of
nocodazole treatment, when the microtubule density was significantly reduced, small PCM clusters
started to move out of the caMTOC and undergo rapid directional motility with speeds of up to 2
μm/sec, which is within the range characteristic for cytoplasmic dynein (Schlager et al., 2014) (Figure
4E,F, Figure 4 – figure supplement 1A, Video 2). Once microtubules were completely disassembled,
the movement of GFP-CDK5RAP2-positive clusters stopped, indicating that it is microtubule-
dependent but occurs only when the microtubule network is partially depolymerized. Since cluster
dispersal towards the cell periphery could be blocked by a dynein inhibitor, and since cytoplasmic
dynein is a minus-end-directed motor, these data indicate that during microtubule disassembly by
nocodazole at 37°C, there is a transient stage when PCM clusters interact with only a few
microtubules, some of which have their minus-ends facing outwards, and these microtubules serve as
tracks for PCM transport. To support this idea, we used motor-PAINT, a technique that employs
nanometric tracking of purified kinesin motors on the extracted cytoskeleton of fixed cells to super-
resolve microtubules and determine their orientation (Tas et al., 2017). Using this approach, we

determined microtubule orientations in centrinone-treated AKAP450/CAMSAP2 knockout cells and in

cells that were also treated with nocodazole for 15 min to induce partial microtubule disassembly

(Figure 4G, Figure 4 – figure supplement 1B). We found that the cells contained a significant number

of minus-end-out microtubules, and their proportion increased during early stages of nocodazole

treatment, possibly because minus-end-out microtubules are more stable (Figure 4G, e.g., minus-end-

out microtubules constituted ~23% of the total microtubule length determined from kinesin-1

trajectories in the untreated cell and ~46% in the nocodazole-treated cell). These microtubules could

serve as tracks for outward movement of PCM, causing the disassembly of caMTOC when the overall

microtubule density around the caMTOC was strongly reduced (Figure 4H). These data suggest that

the dense network of PCM-anchored microtubule minus-ends around a caMTOC allows for its

compaction via dynein-mediated forces, but that dynein can pull the caMTOC apart when

microtubules are disorganized.

To further confirm that caMTOC disassembly is an active microtubule-dependent process, we also

depolymerized microtubules by a combination of cold (4°C) and nocodazole treatment. When all

microtubules were depolymerized, caMTOCs did not fall apart, even when the cells were subsequently

warmed to 37°C in the presence of nocodazole, so that microtubules could not re-grow (Figure 4I).

However, we noticed that in these conditions, the continuity and cylindrical organization of the PCM

cluster were often perturbed. This raised the possibility that the elongated arrangement of PCM

components within caMTOCs is microtubule-driven. Indeed, when cells were subjected to cold (4°C)

treatment in the absence of nocodazole, most of microtubules depolymerized, but some short cold-

stable microtubules remained associated with the caMTOC (Figure 4 – figure supplement 1C). These

data indicate that PCM self-assembly in the absence of centrioles is microtubule-dependent, and

microtubules are involved in shaping the PCM cluster. Once assembled, the PCM cluster is quite stable,

unless microtubule organization is altered and dynein-driven microtubule-based transport pulls it

apart.

Dynamics of caMTOC assembly

When nocodazole-mediated microtubule disassembly was carried out at 4°C, the caMTOC remained

intact and after nocodazole washout it served as the major microtubule nucleation site, similar to the

centrosome in untreated wild-type cells (Figure 5 – figure supplement 1A). However, when

nocodazole-mediated disassembly of the caMTOC was carried out at 37°C, the cluster fell apart and

reassembled upon nocodazole washout (Figure 4A, B), providing a way to study the dynamics of PCM

self-assembly and the roles of different PCM components during this process. Small PCM clusters

positive for pericentrin, CDK5RAP2, γ-tubulin and the centriolar satellite protein PCM1 that co-
localized with the plus-ends of microtubules (labeled with EB1) could be detected 30 s after
nocodazole washout; these PCM clusters and nascent microtubules did not colocalize with the Golgi
membranes (Figure 5A, Figure 5 – figure supplement 1B). Ninein was not detected within the clusters
at this early stage of microtubule regrowth but could be found 2 min after nocodazole washout. In
contrast, no clusters of CEP192 or NEDD1 were observed even 10 min after nocodazole washout
(Figure 5A, Figure 5 – figure supplement 1B,C). The depletion of pericentrin, CDK5RAP2 and γ-tubulin
strongly inhibited microtubule nucleation in these conditions, whereas the depletion of dynein heavy
chain or ninein had a milder effect (Figure 5B,C). Live cell imaging with GFP-CDK5RAP2 and SiR-tubulin
showed that CDK5RAP2 clusters with attached microtubules coalesced by undergoing microtubule-
based movements (Figure 5D), and measurements in cells fixed at different time points after
nocodazole washout showed that a partly radial microtubule system emerged already 2 min after
nocodazole washout (Figure 5 – figure supplement 1D). Reassembly of a single caMTOC in the central
part of the cell occurred within ~15 min after nocodazole washout, though it was less compact than
in cells that were not treated with nocodazole (Figure 5D-G). Depletion of pericentrin, γ-tubulin and
dynein heavy chain strongly inhibited the reformation of a radial microtubule network during
nocodazole washout, whereas the effect of depleting CDK5RAP2 and ninein was less strong (Figure 5
– figure supplement 1D-F). Live cell imaging of acentriolar AKAP450/CAMSAP2 knockout RPE1 cells
stably expressing GFP-CDK5RAP2 showed that when pericentrin was depleted, CDK5RAP2 clusters
were not detectable, and the microtubule network, both before nocodazole treatment and after
nocodazole washout, was disorganized (Figure 5 – figure supplement 2, Video 3). Taken together, our
data show that pericentrin and γ-tubulin form microtubule-nucleating and anchoring units, which are
clustered by the self-association of pericentrin and assembled into larger structures by dynein-based
transport. CDK5RAP2 contributes to microtubule nucleation efficiency, whereas ninein appears to act
somewhat later and contributes to the formation of a compact PCM cluster and a radial microtubule
network. Importantly, all these proteins can cluster in the absence of centrioles, and together they
can efficiently nucleate and anchor microtubules.

The role of CAMSAP2-stabilized minus ends in defining microtubule network geometry

The results of nocodazole treatment and washout suggested that PCM can self-assemble into a
caMTOC which nucleates and anchors microtubules, but this structure is sensitive to microtubule
organization. This observation prompted us to investigate in more detail how the microtubules that
are not anchored at PCM clusters affect PCM organization in steady state conditions. An abundant

population of stable minus ends that do not attach to PCM is decorated by CAMSAP2. In centrinone-treated wild type cells, CAMSAP2-bound microtubule minus ends were anchored at the Golgi (Wu et al., 2016)(Figure 1D), whereas in centrinone-treated AKAP450 knockout cells they were distributed randomly (Figure 1D, 6A). Live imaging showed that CAMSAP2-decorated minus ends displayed only limited motility on the scale of hours and thus formed a relatively stationary, disorganized microtubule network (Video 4). Live imaging of GFP-CDK5RAP2 together with SiR-tubulin in these cells demonstrated that small PCM clusters were distributed throughout the cytoplasm (Figure 6B). These clusters moved along microtubules and encountered each other, but the direction of the movements was random and the clusters did not coalesce into a single structure (Figure 6B, Figure 6 – figure supplement 1A, Video 5). Treatment with nocodazole and subsequent nocodazole washout confirmed that the motility of GFP-CDK5RAP2 clusters in centrinone-treated AKAP450 knockout cells was microtubule-dependent, and that these clusters could nucleate microtubules and move together with microtubule minus ends, but did not converge into a single caMTOC (Figure 6 – figure supplement 1A-C, Video 5). Treatment with dynapyrazole A strongly inhibited the movements of small PCM clusters (Figure 6 – figure supplement 1D,E, Video 6), indicating that they are dynein-driven. After the depletion of pericentrin, GFP-CDK5RAP2 became completely diffuse, and it failed to form clusters during nocodazole treatment or washout (Figure 6 – figure supplement 2, Video 7), indicating that clustering of GFP-CDK5RAP2 in AKAP450 knockout cells is pericentrin-dependent. Based on these data, we conclude that in AKAP450 knockout cells, pericentrin still forms PCM clusters that can nucleate microtubules and can be moved by dynein along other microtubules, similar to what occurs in wild-type cells. However, in the absence of AKAP450, CAMSAP2-stabilized microtubules form a disorganized network, which imposes a random motility pattern on pericentrin-dependent PCM clusters and thus prevents their assembly into a single caMTOC, likely because PCM interactions are not sufficient to trigger their stable association (Figure 6C).

If the geometry of the CAMSAP2-stabilized microtubule network determines PCM distribution, focusing CAMSAP2-bound minus ends is expected to bring PCM together. To test this idea, we linked CAMSAP2-stabilized minus ends to a minus-end-directed motor. In order to avoid potential cell toxicity associated with manipulating cytoplasmic dynein, we used the motor-containing part of a moss kinesin-14, type VI kinesin-14b from the spreading earthmoss *Physcomitrella patens* (termed here ppKin14). The C-terminal motor-containing part of this protein can efficiently induce minus-end-directed motility of different cargoes in mammalian cells when it is tetramerized through a fusion with the leucine zipper domain of GCN4 (GCN4-ppKin14-VIb (861–1321)) and recruited to cargoes using inducible protein heterodimerization (Jonsson et al., 2015; Nijenhuis et al., 2020). We employed a chemical heterodimerization system that is based on inducible binding of two protein domains, FRB

and FKBP, upon the addition of a rapamycin analog (rapalog) (Pollock et al., 2000). To ensure that all

CAMSAP2-decorated microtubule minus ends were linked to kinesin-14, we rescued centrinone-

treated AKAP450/CAMSAP2 knockout cells by expressing CAMSAP2 fused to a tandemly repeated

FKBP domain (2FKBP-mCherry-CAMSAP2) (Figure 6D-F). This construct was co-expressed with the FRB-

GCN4-tagBFP-ppKin14 fusion, which by itself localized quite diffusely, with only a weak enrichment

along microtubules, as described previously (Nijenhuis et al., 2020) (Figure 6D,F). In the absence of

rapalog, CAMSAP2-decorated microtubule minus ends were distributed randomly, similar to

endogenous CAMSAP2 in AKAP450 knockout cells (Figure 6F). However, upon rapalog addition,

ppKin14 was rapidly recruited to CAMSAP2-decorated microtubule ends, and after 2 hrs, more than

90% of cells acquired a radial microtubule organization (Figure 6E-G). In rapalog-treated cells,

CAMSAP2-bound microtubule minus ends formed either a tight cluster or a "whirlpool-like" ring in the

cell center (Figure 6D-G, Figure 6 – figure supplement 1F). The whirlpool-like arrangement likely comes

about when CAMSAP2-stretches are a bit longer and continue to slide against each other, forming a

nematic circular bundle. The major caMTOC components, pericentrin, CDK5RAP2, γ-tubulin and ninein

were also concentrated within the CAMSAP2 cluster (Figure 6F, Figure 6 – figure supplement 1F).

These data indicate that the positioning of stabilized minus-ends is an important determinant of the

overall microtubule organization and PCM localization in interphase cells.

The role of PCM in CAMSAP2-driven microtubule organization

Presence of PCM in the caMTOC induced by minus-end-directed transport of CAMSAP2 might be a

passive consequence of microtubule reorganization, but might also play an active role in forming this

caMTOC. To distinguish between these possibilities, we attached CAMSAP2 to ppKin14 in centrinone

-treated   cells   where   both   AKAP450   and   pericentrin   were   knocked   out

(AKAP450/CAMSAP2/p53/pericentrin knockout). In the absence of rapalog, CAMSAP2-stabilized

minus ends and the whole microtubule network were disorganized, as expected, and the same was

true when the two constructs were expressed separately, with or without rapalog (Figure 7A,D, Figure

7- figure supplement 1). After rapalog addition, microtubules in cells expressing both constructs

acquired a radial organization, but their minus ends usually did not converge in a single spot but rather

accumulated in a ~30-70 μm-large ring-like structure (Figure 7A,D, Figure 7- figure supplement 2,

Video 8). Staining for PCM markers showed that CDK5RAP2 and γ-tubulin were enriched in the vicinity

of CAMSAP2-positive microtubule minus ends, whereas ninein appeared rather diffuse (Figure 7A). To

determine the nature of the structure "corralled" by the ring of CAMSAP2-decorated minus ends in

rapalog-treated cells, we stained for different membrane organelles and found that whereas there

was no strong correlation with the nucleus, Golgi membranes or lysosomes, the majority of mitochondria were found within the CAMSAP2 ring, and the endoplasmic reticulum (ER) displayed increased density overlapping with the CAMSAP2 ring (Figure 7B, Figure 7 - figure supplement 2B). It therefore appeared that in the absence of pericentrin, CAMSAP2-decorated minus ends were brought together by ppKin14, but their convergence was inefficient and possibly impeded by membrane organelles enriched in the central, thicker part of the cell before rapalog addition (see the upper panel of Figure 7 - figure supplement 2B). Transient transfection of centrinone-treated AKAP450/CAMSAP2/p53/pericentrin knockout cells with GFP-pericentrin rescued the formation of a tight CAMSAP2 cluster upon rapalog treatment (Figure 7C). Our data show that pericentrin-containing PCM contributes to the formation of a caMTOC driven by minus-end-directed transport of CAMSAP2-stabilized minus-ends.

To support this notion further, we also generated cells that were knockout for AKAP450, CAMSAP2, CDK5RAP2, myomegalin (MMG, homologue of CDK5RAP2), p53 and pericentrin. To achieve this, we used the previously described RPE1 cell line knockout for AKAP450, CAMSAP2, CDK5RAP2 and MMG (Wu et al., 2016), in which we sequentially knocked out p53 and pericentrin (Figure 7 – figure supplement 3A-G). While it was not possible to induce centriole loss by centrinone B treatment in AKAP450/CAMSAP2/CDK5RAP2/MMG knockout cells because the proliferation of these cells was arrested in the absence of centrioles (Wu et al., 2016), centriole removal was successful in AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/pericentrin knockout cells due to the absence of p53 and led to microtubule disorganization (Figure 7 – figure supplement 3H). Interestingly, when these cells were co-transfected with FKBP-linked CAMSAP2 and FRB-linked ppKin14 and treated with rapalog (Figure 6D), we observed that CAMSAP2 clustering was even less efficient than in AKAP450/CAMSAP2/p53/pericentrin knockout cells (Figure 7D,E). 49% of AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/pericentrin knockout cells had small bundles of CAMSAP2 stretches dispersed throughout the cytoplasm, and only 37% of these cells formed a ring of CAMSAP2-decorated minus ends, whereas 80% of AKAP450/CAMSAP2/p53/pericentrin knockout cells formed such a ring. We examined the ER and mitochondria in these cells and found that in cells that did form a CAMSAP2 ring, the ER displayed an overlapping ring-like density, although the mitochondria inside the CAMSAP2 ring were more scattered compared to those of AKAP450/CAMSAP2/p53/pericentrin knockout cells (Figure 7 – figure supplement 2B). In cells with dispersed CAMSAP2-positive bundles, no increased ER density or central accumulation of mitochondria were observed (Figure 7 – figure supplement 2B). These data further support the notion that minus-end-directed transport of stable minus ends alone is insufficient to generate a caMTOC, and that synergy with PCM is required.

Recapitulation of PCM self-organization by computer simulations

To rationalize the appearance of the different microtubule arrangements and to find a minimal set of interactions between filaments and motors that would lead to self-organization into structures that we observed experimentally, we set up agent-based computer simulations with Cytosim (Nedelec and Foethke, 2007). First, we sought to recapitulate the formation of a single PCM cluster. The numerical values for the parameters of our simulations have been taken from literature or reasonably chosen otherwise (Supplemental Table S1). For simplicity, we considered a two-dimensional circular cell with a radius of 10 μm. We described a mobile PCM complex as a bead with a radius of 50 nm from which one microtubule plus end could grow. Microtubule growth and shrinkage were simulated with the classical microtubule model from Cytosim. When the microtubule reached a maximal length of 7.5 μm, its growth was stopped. In this way, we limited the microtubule length to avoid long microtubules that push their minus end to the periphery of the cell. Additionally, one dynein molecule was attached to a PCM complex with its cargo-binding domain. With this configuration, dynein molecules could transport PCM complexes along microtubules growing from other PCM complexes. Once they were bound to a microtubule, they walked towards the minus end in a force-dependent manner. When a dynein motor reached a minus end, it detached. Furthermore, we implemented a reversible binding interaction between PCM complexes to make them adhere to each other. Because we were interested in the spatial arrangement of the PCM complexes, we introduced steric interactions between them. However, we neglected steric interactions between microtubules to effectively account for the three-dimensional space that we projected to two dimensions. Simulations of 300 of such PCM complexes reproducibly reached a steady state in which they formed a compact centrally located cluster (Figure 8A, Video 9A). Quantification of 10 simulations showed that the mean and standard deviation of the distance from each PCM complex to the average location of all complexes (the center of mass) were small compared to the radius of the cell (Figure 8E, F). However, if we disabled the adhesive interactions between PCM complexes, they formed a loose ring-like arrangement and not a compact cluster (Figure 8B, Video 9B), and the mean and standard deviation of the distance of PCM complexes to the center of mass were larger (Figure 8E, F). Taken together, our simulations support the idea that PCM components form a central cluster through positive feedback of dynein molecules carrying them towards the minus ends of microtubules attached to other PCM proteins, and that adhesive interactions between the PCM complexes promote cluster compaction.

In our experiments, we saw that the presence of CAMSAP2-stabilized microtubule network prevented PCM clustering (Figure 1D,E, Figure 6A-C), and that CAMSAP2-bound minus ends appeared almost stationary on the scale of minutes (Video 4) compared to the rapid dynein-driven movement of small PCM clusters (Video 5). To simulate such a situation, we assumed that 300 microtubules could grow

and shrink with random orientations from CAMSAP2-stabilized microtubule ends that were randomly
distributed and stationary within the cell. Furthermore, we assumed that some of the moving PCM
complexes were not efficient in nucleating microtubules, because the presence of a PCM-independent
microtubule population would reduce the concentration of soluble tubulin and thus nucleation
efficiency. To effectively account for such an effect, we allowed only 50 of the 300 PCM complexes to
nucleate microtubules. In this system, the PCM cluster formation was disrupted, and PCM complexes
randomly moved around the cell (Figure 8C, Video 9C). The distributions of the distances to the
average position of the PCM complexes were broad and had a large mean and standard deviation,
indicating the dispersed localization (Figure 8E, F). However, if we increased the strength of adhesive
interactions between the PCM complexes, the randomly oriented microtubule network no longer
suppressed the clustering of PCM complexes (Figure 8D, Video 9D). Therefore, the presence of a
stationary randomly organized microtubule network can be sufficient to block the formation of a
compact PCM cluster even though individual PCM complexes can still move with dynein and adhere
to each other, provided that the interaction between PCM complexes is not too tight.

Next, we examined how CAMSAP2-stabilized microtubules would organize when minus-end-directed
kinesin-14 motors could attach to them and carry the minus ends. CAMSAP2-stabilized microtubule
ends bound to kinesin-14 motors (termed "CAMSAP-kin14 complexes" in the text below) were
modelled as a bead with a radius of 50 nm from which a microtubule could grow and to which 6
kinesin-14 molecules were attached. Because the biophysical properties of single plant kinesin-14
molecules used in our assays are poorly understood, we assumed typical kinesin-1 values with an
opposite directionality. CAMSAP-kin14 complexes only interacted sterically but did not adhere to each
other. In a simulation of 300 CAMSAP-kin14 complexes, a loose ring-like arrangement appeared
(Figure 8G), similar to the one observed in our simulations of PCM complexes that could not bind to
each other (Figure 8B). Introducing adhesive interactions between CAMSAP-kin14 complexes was
sufficient to promote compact cluster formation (Figure 8H).

Finally, to investigate if a compact cluster of CAMSAP-kin14 complexes would emerge in the presence
of self-associating PCM complexes, we set up simulations with 150 CAMSAP-kin14 complexes together
with 150 adherent PCM complexes. The steady state of such a system displayed a compact central
cluster, in which both types of complexes were mixed (Figure 8I,K,L, Video 9I), while making PCM
complexes non-adhesive prevented cluster compaction (Figure 8J,K,L, Video 9J). Taken together, our
simulations suggest that the PCM complexes provide enough adhesive interactions to compact the
cluster of non-interacting CAMSAP-kin14 and PCM complexes, very similar to our experimental
observations.

Discussion

In this study, we explored the mechanisms of interphase PCM self-assembly in the absence of centrioles. Our experiments and simulations support the idea that PCM complexes can form a single cluster through a positive feedback mechanism, whereby dynein motors carry microtubule minus-ends to other microtubule minus ends (Cytrynbaum et al., 2004). However, for a compact cluster to emerge, the minus ends must not only be able to move towards each other but also to bind to each other, and this idea is fully supported by our simulations. Interphase PCM is thus capable both of dynein binding and self-association sufficient to organize a compact microtubule-nucleating and anchoring structure in the absence of centrioles. However, the formation of a compact MTOC in acentriolar cells is slow and less robust than in centriole-containing cells, and the resulting structure is sensitive to the overall organization of microtubules and to dynein function. This means that interphase PCM self-association is by itself reversible and not sufficiently tight to resist dynein-driven forces. Centrioles can thus be regarded as catalysts of PCM assembly and stabilizers of interphase centrosomes, preventing PCM movement on microtubules oriented with their minus ends away from the centrosome.

Our experimental system allowed us to examine which PCM components are capable of associating with each other and with dynein to promote microtubule nucleation and anchoring independently of centrioles. The major scaffold for interphase acentriolar PCM assembly is pericentrin, which can self-associate (Jiang et al., 2021) and form clusters that recruit CDK5RAP2 and ninein, two proteins known to bind to pericentrin (Chen et al., 2014; Delaval and Doxsey, 2010; Kim and Rhee, 2014; Lawo et al., 2012). CDK5RAP2 is important for efficient microtubule nucleation, consistent with its role as an activator of γ-TuRC (Choi et al., 2010). The same may be true for pericentrin itself (Takahashi et al., 2002) and, to a lesser extent, for ninein (Delgehyr et al., 2005; Mogensen et al., 2000), which was less important for microtubule nucleation in our assays. Ninein might be required for promoting minus-end anchoring, as proposed by previous studies (Abal et al., 2002; Chong et al., 2020; Delgehyr et al., 2005; Goldspink et al., 2017; Lechler and Fuchs, 2007; Mogensen et al., 2000; Shinohara et al., 2013; Zheng et al., 2020). The importance of ninein for the formation of caMTOCs highlights its function within the PCM independent of its role at the centriolar appendages, a major site of ninein localization at the centrosome (Chong et al., 2020; Delgehyr et al., 2005; Sonnen et al., 2012). Pericentrin can also directly interact with dynein through the dynein light intermediate chain (Purohit et al., 1999). Ninein was shown to form a triple complex with dynein and dynactin and activates dynein motility (Redwine et al., 2017), and an interaction between CDK5RAP2 and dynein has also been reported (Jia et al., 2013; Lee and Rhee, 2010). All these interactions likely contribute to dynein-dependent PCM coalescence in the absence of centrioles and also to the function of PCM in microtubule organization,

as in the absence of PCM clustering, microtubule density is strongly reduced even though all PCM

proteins are expressed.

Pericentrin-dependent MTOC assembly has also been observed in acentriolar mitotic cells (Chinen et

al., 2021; Watanabe et al., 2020), but the interphase pathway displays some interesting differences.

Most notably, CEP192 and its binding partners CEP152 and NEDD1 (Gomez-Ferreria et al., 2012;

Joukov et al., 2014; Kim et al., 2013; Sonnen et al., 2013) were not enriched at caMTOCs, and their

depletion appeared to have no impact on caMTOC formation. This is surprising, because CEP192 is

essential for cell division (Gomez-Ferreria et al., 2007; Joukov et al., 2014; Yang and Feldman, 2015;

Zhu et al., 2008), and in acentriolar cells that rely on pericentrin and CDK5RAP2 for spindle pole

formation, CEP192 is recruited to mitotic pericentrin clusters (Chinen et al., 2021; Watanabe et al.,

2020). Moreover, although NEDD1 is targeted to centrosomes by CEP192, pericentrin can also

contribute to the centrosomal targeting of NEDD1 independently of CEP192 during mitosis (Chi et al.,

2021). However, our data indicate that in interphase, pericentrin and CEP192/NEDD1 are brought

together by centrioles. This conclusion is important for understanding MTOC assembly in

differentiated cells, where centrosome function is suppressed. Strikingly, pericentrin, ninein,

CDK5RAP2 and γ-TuRC are also present in non-centrosomal PCM-based MTOCs at the nuclear

envelope and the Golgi membranes in muscle cells, where they are targeted by AKAP450 (Gimpel et

al., 2017; Oddoux et al., 2013; Vergarajauregui et al., 2020). Recent work also demonstrated that at

the ciliary base of certain types of worm neurons, there is an acentriolar PCM-dependent MTOC that

is formed by the functional counterparts of CDK5RAP2 (SPD-5), pericentrin (PCMD-1) and γ-tubulin,

but lacks CEP192/SPD-2 (Garbrecht et al., 2021; Magescas et al., 2021). The co-assembly of

pericentrin, ninein and CDK5RAP2 counterparts into structures that can promote γ-TuRC-dependent

microtubule nucleation and also anchor minus ends may thus be a general property of interphase

PCM-dependent acentriolar MTOCs. Additionally, the participation of dynein is likely to be a general

feature of acentrosomal microtubule organization, as exemplified by MTOCs formed by the Golgi

apparatus or endosomal membranes (Liang et al., 2020; Zhu and Kaverina, 2013).

Our simple cellular system allowed us to dissect the molecular details of acentriolar PCM assemblies.

We found that caMTOCs recruited multiple MAPs and a subset of centriolar proteins. For example,

caMTOCs accumulated several +TIPs, such as CLASPs, CLIP170 and chTOG, though the depletion of

these proteins had no effect on caMTOC formation or function. In contrast, the core components of

the +TIP complexes, EB1 and EB3, were neither enriched in caMTOCs nor required for their assembly.

While the negative results on other +TIPs might be due to their incomplete depletion, EB1 and EB3

function was tested using genetic knockouts, indicating that interphase PCM function is EB-

independent. Among the centriolar proteins, we detected CPAP, CP110 and CEP120 in caMTOCs, and

it will be interesting to test whether any of these microtubule-binding factors contribute to microtubule organization independently of their participation in centriole and centrosome assembly.

Furthermore, our work provided insight into the self-assembly properties of interphase PCM. It seems that while oligomerization and clustering of pericentrin molecules is important for MTOC formation both in interphase and in mitosis, during mitotic entry, pericentrin forms condensates (Jiang et al., 2021), while in interphase, pericentrin clusters show no hallmarks of liquid droplets or condensates. Moreover, the compaction and shape of these acentriolar PCM clusters depend on dynein and microtubules. While caMTOCs were stable when all microtubules were disassembled, indicating that attractive interactions between PCM components are sufficient to keep them together, cold treatment experiments suggested that the cylindrical shape of caMTOCs is likely due to their association with some stable microtubules. Furthermore, when the number of PCM-unattached stable minus-end-out microtubules was increased by either nocodazole treatment or by the presence of CAMSAP2, PCM clusters could no longer form a compact structure but continued to move along microtubules. Therefore, an important outcome of our study is the key role of stabilized minus ends in determining interphase PCM organization in the absence of the centrosome. This property may help to explain how acentrosomal MTOCs form in cells where both PCM proteins and PCM-independent microtubule stabilizers such as CAMSAPs are present. Two well studied examples of such systems are the MTOCs at the Golgi membranes and the apical cortex of epithelial cells, where CAMSAPs and PCM proteins are present simultaneously and may cooperate with each other or play redundant roles (Goldspink et al., 2017; Nashchekin et al., 2016; Noordstra et al., 2016; Sanchez et al., 2021; Toya et al., 2016; Wang et al., 2015; Wu et al., 2016; Zhu and Kaverina, 2013). We found that CAMSAP2-stabilized minus ends can exert a highly dominant effect on PCM organization, and this property likely explains how the Golgi, which anchors CAMSAP2-stabilized microtubules, recruits PCM and becomes the major MTOC in cells lacking centrosomes. Importantly, our experiments and simulations showed that coupling stable minus-ends to a minus-end directed motor is by itself insufficient to form a compact MTOC, but self-clustering PCM can contribute to this process. Altogether, self-association of interphase PCM appears to be strong enough to promote its clustering but is sufficiently dynamic to allow PCM reorganization dependent on other microtubule regulators present in the cell.

An interesting question that remains unanswered by our work is the inhibitory role of PLK4 in interphase caMTOC formation. We did observe caMTOCs in cells depleted of PLK4, indicating that, unlike cells lacking TRIM37, which form PCM clusters containing catalytically inactive PLK4 (Meitinger et al., 2020; Yeow et al., 2020), interphase cells studied here do not rely on enzymatically inactive PLK4 for PCM assembly. PLK4 is known to phosphorylate NEDD1 (Chi et al., 2021), and it is possible that the lack of phosphorylation prevents this γ-TuRC activator and its partners, such as CEP192, from

606   participating in interphase caMTOC assembly. It is, of course, also possible that PLK4 phosphorylation
607   inhibits the interactions or activities of some of the players driving caMTOC formation. The easy-to-
608   manipulate cellular model that we have described here will allow these questions to be addressed and
609   facilitate detailed studies of the interactions and functions of PCM components in nucleating and
610   stabilizing interphase microtubule minus ends.

Materials and Methods

DNA constructs and protein purification

To generate the lentiviral vector pLVX-GFP-CDK5RAP2-IRES-Puro, pLVX-IRES-Puro plasmid (Clontech) was digested with AgeI and NotI (FastDigest, Thermo Fisher), and then Gibson Assembly (NEB) was performed with gel-purified PCR product of GFP-CDK5RAP2 (Wu et al., 2016). To generate pB80-FRB-TagBFP-GCN4-ppKin14 and pB80-FRB-HA-GCN4-ppKin14, pB80-FRB-GFP-GCN4-ppKin14-VIb was digested with XbaI and BsrGI (FastDigest, Thermo Fisher), and then TagBFP and HA-tag encoding DNA fragments were subcloned into the linearized vector by Gibson Assembly.

To generate the PX459 with single guide RNA (sgRNA) sequences, pSpCas9(BB)-2A-Puro (PX459) V2.0 (Ran et al., 2013)(purchased from Addgene) was digested with FastDigest BbsI (Thermo Fisher), and the annealing product of single-strand sgRNA-encoding oligonucleotides was inserted into the linear PX459 linear vector by T4 ligation (Thermo Fisher). The sgRNA sequences that were used in this study are: sgRNA targeting AKAP450 5'- gAGGGTTACCTATGGGACTGA -3'; sgRNA targeting CAMSAP2 encoding gene 5'-gCATGATCGATACCCTCATGA-3'; sgRNA targeting p53-encoding gene exon 2 #1 5'-gCGTCGAGCCCCCTCTGAGTC-3'; sgRNA targeting p53 exon 4 #2 5'-gCCATTGTTCAATATCGTCCG-3'; sgRNA targeting PCNT exon 5 #1 5'-gAGACGGCATTGACGGAGCTG-3'; sgRNA targeting pericentrin-encoding gene exon 5 #2 5'-GCTCAACAGCCGGCGTGCCC-3'.

To generate the GST-DmKHC(1-421)-mNeonGreen construct used for protein purification for motor-PAINT, the fragment containing amino acids 1 to 421 of the Drosophila melanogaster Kinesin Heavy Chain (DmKHC) was amplified from donor construct DmKHC(1-421)-GFP-6x-His with a C-terminal mNeonGreen tag by PCR and then cloned into a pGEX vector. The plasmid was transformed into *E. coli* BL21 cells for purification. Bacteria were cultured until OD600 $\approx$0.7 and cultures were cooled prior to inducing protein expression with 0.15 mM IPTG at 18°C overnight. Cells were then pelleted by centrifugation, snap frozen in liquid nitrogen, and stored at -80°C until use. Cells were rapidly thawed at 37°C before being resuspended in chilled lysis buffer (phosphate buffered saline (PBS) supplemented with 5 mM MgCl$_2$, 0.5 mM ATP, 0.1% Tween 20, 250 mM NaCl, and 1x complete protease inhibitor; pH 7.4). Bacteria were lysed by sonication (5 rounds of 30 s) and supplemented with 5 mM DTT and 2mg/mL lysozyme and then incubated on ice for 45 min. The lysate was clarified by centrifuging at 26000 xg for 30 minutes before being incubated with equilibrated Glutathione Sepharose 4B resin for 1.75 hrs. Beads were then pelleted, resuspended in wash buffer (PBS supplemented with 5 mM MgCl$_2$, 0.1% Tween 20, 250 mM NaCl, 1 mM DTT, and 0.5 mM ATP; pH 7.4), and transferred to a BioRad column. Once settled, the resin was washed with 2 x 10 column volumes (CV) wash buffer, followed by 1 x 10CV PreScission buffer (50 mM Tris-HCl, 5 mM MgCl$_2$, 100mM NaCl,

1 mM DTT, 0.5 mM ATP; pH 8.0). The resin was then incubated overnight in 4CV PreScission buffer

with 80U PreScission protease to cleave off the GST tag. The following morning, after allowing the

resin to settle, the eluent was collected, concentrated by spinning through a 3000kDa MWCO filter,

supplemented with an additional 0.1mM ATP, 1mM DTT, and 20% w/v sucrose before flash freezing

in liquid nitrogen, and finally stored at -80°C. Concentration was determined using a Nanodrop. All

steps from lysis onwards were performed at 4°C.

Cell culture and drug treatment

hTERT immortalized RPE-1 (RPE1) cell lines were grown in an 1:1 mix of DMEM and F-10 (Lonza) and

Human Embryonic Kidney (HEK) 293T cells line were cultured in DMEM, both supplemented with 10%

fetal bovine serum (FBS, GE Healthcare) and 1% penicillin and streptomycin (Sigma-Aldrich). All cells

were grown in tissue culture polystyrene flasks (Corning) and were maintained in a humidified

incubator at 37°C with 5% CO_2. Mycoplasma contamination was routinely checked with LT07-518

Mycoalert assay (Lonza).

FuGENE 6 (Promega) was used to transfect RPE1 cells with plasmids for generating CRISPR/Cas9

knockouts, immunofluorescence staining and live cell imaging; RNAiMAX (Thermo Fisher Scientific)

was used to transfect RPE1 cells with siRNAs at 20 nM; MaxPEI was used to transfect HEK293T cells

for lentivirus packaging. Transfections were performed according to the manufacturer's instructions

within the recommended reagent/DNA or reagent/siRNA ratio range.

We used the following drugs: centrinone B (Tocris Bioscience), nocodazole (Sigma), rapalog (A/C

Heterodimerizer, Takara), dynapyrazole A (Sigma-Aldrich) , BI2536 (Selleckchem), Thymidine (Sigma-

Aldrich) , proTAME (Boston Biochem) and Brefeldin A (Peptrotech).

To remove centrioles, RPE1 cells were treated with 125 nM centrinone B containing complete medium

for ~10 days, and drug-containing medium was refreshed every 24 hrs; cell confluence was maintained

around ~50-80% during the treatment.

For the microtubule disassembly and regrowth assay, the acentriolar RPE1 cells were seeded onto

coverslips in 24-well plates and incubated for 24 hrs, then cells were treated with 10 μM nocodazole

for 1 hr in an incubator (37°C, 5% CO_2) and followed by another 1 hr treatment at 4°C to achieve

complete disassembly of stable microtubule fragments. Nocodazole washout was then carried out by

at least six washes on ice with ice-cold complete medium; subsequently, plates were moved to a 37°C

water bath and pre-warmed medium was added to each well to allow microtubule regrowth.

For cell cycle synchronization, centrinone-treated AKAP450/CAMSAP2/P53 knockout cells were

treated with 5 mM Thymidine (Sigma-Aldrich) overnight, released in centrinone containing medium

for 4 hrs and subsequently treated with 5µM proTAME (Boston Biochem, I-440) for 2 hrs before being

released in centrinone containing medium for 1-4 hrs followed by live imaging and fixation.

For the inducible ppKin14-CAMSAP2 heterodimerization experiment, acentriolar cells were seeded

onto coverslips in 24-well plates, cultured with centrinone B containing medium and co-transfected

with 2FKBP-mCherry-CAMSAP2 and FRB-TagBFP-GCN4-ppKin14 vectors. 24 hrs after transfection,

rapalog was added into the medium at a final concentration of 50 nM and incubated overnight for

preparation of fixed cells. For live imaging, rapalog was used at 100 nM.

Lentivirus packaging and generation of transgenic stable cell lines

Lentiviruses were produced by MaxPEI-based co-transfection of HEK293T cells with the transfer

vectors together with the packaging vector psPAX2 and envelope vector pMD2.G (psPAX2 and

pMD2.G were a gift from Didier Trono, Addgene plasmid #12259 and #12260; RRID:Addgene_12259

and RRID:Addgene_12260). Supernatant of packaging cells was harvested 48-72 hrs after transfection,

filtered through a 0.45 µm filter, incubated with a polyethylene glycol (PEG)-6000-based precipitation

solution overnight at 4°C and centrifuged for 30 minutes at 1500 rpm to concentrate the virus.

Lentiviral pellet was resuspended in PBS.

Wild type, AKAP450 and AKAP450/CAMSAP2 knockout RPE1 cells were infected with lentivirus and

incubated in complete medium supplemented with 8 µg/ml polybrene (Sigma-Aldrich). After 24 hrs,

the cell medium was replaced with fresh medium. Starting 72 hrs after viral transduction, cells were

subjected to selection with puromycin at a concentration of 25 µg/ml for wild-type, 20 µg/ml for

AKAP450 knockout and 15 µg/ml for AKAP450/CASMAP2 knockout for up to 3 days (until most of the

untransduced control cells, treated with the same concentration of antibiotic, were dead). After

selection, cells were grown in normal medium for 3 days and individual colonies expressing GFP were

isolated into 96-well plates by fluorescence-activated cell sorting (FACS). Sorted single transgenic

stable cell lines were further confirmed by immunofluorescence staining to check the expression level

of GFP-CDK5RAP2 and its colocalization with other centrosomal proteins.

Generation of CRIPSR/Cas9 knockout cell lines

The CRISPR/Cas9-mediated knockout of p53-, pericentrin-, AKAP450- and CAMSAP2-encoding genes

was performed as described previously (Ran et al., 2013). In brief, AKAP450/CAMSAP2 knockout RPE1

cells (Wu et al., 2016) were transfected with the vectors bearing the appropriate targeting sequences

using FuGENE 6. One day after transfection, the transfected AKAP450/CAMSAP2 knockout RPE1 cells

were subjected to selection with 15 μg/ml puromycin for up to 3 days. After selection, cells were

allowed to recover in normal medium for ~7 days, and knockout efficiency was checked by

immunofluorescence staining. Depending on the efficiency, 50–500 individual clones were isolated

and confirmed by immunofluorescence staining, and the resulted single colonies were characterized

by Western blotting, immunostaining and genome sequencing. AKAP450/CAMSAP2/p53 and

AKAP450/CAMSAP2/MMG/CDK5RAP2/p53 knockout cell lines were generated first and subsequently,

each of them was used to knock out the gene encoding pericentrin. The mutated portions of the p53-

and pericentrin-encoding genes were sequenced using gel-purified PCR products obtained with

primers located in the vicinity of the corresponding sgRNA targeting sites.

Antibodies, immunofluorescence staining and Western blotting

Antibodies used for immunostaining and Western blotting are listed in the Key Reagent or Resource

table. For immunofluorescence cell staining, cultured cells were fixed with –20°C methanol for 5 min

or with 4% paraformaldehyde (PFA) for 12 min at room temperature, rinsed in PBS for 5 min,

permeabilized with 0.15% Triton X-100 in PBS for 2 min, washed 3 times for 5 min with 0.05% Tween-

20 in PBS, sequentially incubated for 20 min in the blocking buffer (2% BSA and 0.05% Tween-20 in

PBS), 1 hr with primary antibodies in the blocking buffer, washed 3 times for 5 min with 0.05% Tween-

20 in PBS, then for 1 hr in secondary antibodies in the blocking buffer, washed 3 times for 5 min with

0.05% Tween-20 in PBS, and air-dried after a quick wash in 96% ethanol. Cells were mounted in

Vectashield mounting medium with or without DAPI (Vector laboratories, Burlingame, CA). Alexa Fluor

-405, -488, –594 and –647 conjugated goat antibodies against rabbit, rat and mouse IgG were used as

secondary antibodies (Molecular Probes, Eugene, OR).

For Western blotting, cells were harvested from six-well plates or 10 cm dishes at 90% confluence and

protein extracts were prepared using the lysis buffer containing 20 mM Tris-Cl, pH 7.5, 150 mM NaCl,

0.5 mM EDTA, 1 mM DTT, 1% Triton X-100 or RIPA buffer containing 50 mM Tris-HCl, pH 7.5, 150 mM

NaCl, 1% Triton X-100, 0.5% Sodium Deoxycholate supplemented with protease inhibitor and

phosphatase inhibitors (Roche). Samples were run on polyacrylamide gels, followed by transfer on

0.45 μm nitrocellulose membrane (Sigma-Aldrich). Blocking was performed in 2% BSA in PBS for 30

min at room temperature. The membrane was first incubated with the primary antibodies overnight

at 4°C and washed with 0.05% Tween-20 in PBS 3 times and subsequently incubated with secondary

antibodies for 1 hr at room temperature and washed 3 times with 0.05% Tween-20 in PBS. IRDye

800CW/680 LT Goat anti-rabbit and anti-mouse were used as secondary antibodies (Li-Cor Biosciences, Lincoln, LE) and membranes were imaged on Odyssey CLx infrared imaging system (Image Studio version 5.2.5, Li-Cor Biosciences).

Imaging and analysis of fixed cells

Images of fixed cells were collected with a Nikon Eclipse Ni upright fluorescence microscope equipped with a DS-Qi2 CMOS camera (Nikon), an Intensilight C-HGFI epi-fluorescence illuminator (Nikon), Plan Apo Lambda 100× NA 1.45 or Plan Apo Lambda 60x N.A. 1.40 oil objectives (Nikon) and driven by NIS-Elements Br software (Nikon).

Gated STED imaging was performed with Leica TCS SP8 STED 3X microscope driven by LAS X software using HC PL APO 100x/1.4 oil STED WHITE objective, white laser (633 nm) for excitation and 775 nm pulsed lased for depletion. Images were acquired in 2D STED mode with vortex phase mask. Depletion laser power was equal to 90% of maximum power and an internal Leica HyD hybrid detector with a time gate of $1 \leqslant tg \leqslant 8$ ns was used.

ImageJ was used for adjustments of intensity levels and contrast, quantification of the immunofluorescence signal intensity and maximum intensity projections. To analyze PCM clustering after nocodazole washout in AKAP450/CAMSAP2 knockout RPE1 cells, images were separated into concentric circular areas using Concentric Circles plugin of ImageJ. The biggest PCM cluster (which normally also had the highest fluorescence intensity) was selected as the center, around which 20 circles with 2 μm inner radius and 20 μm outer radius were drawn. Fluorescence intensity of PCM clusters in these concentric circles was measured automatically and normalized by the sum of the total PCM intensity in each cell per condition. To quantify the areas occupied by PCM clusters, immunofluorescence images of fixed cells and time lapse images of live cells were analyzed by drawing the smallest circle that covered visible PCM clusters to indicate the area occupied by the PCM clusters, and the diameters of the circles were used for the quantification.

Measurements of microtubule radiality

To analyze microtubule radiality, images of fluorescently labeled microtubules were separated into radial and non-radial components using a customized ImageJ macro (https://github.com/ekatrukha/radialitymap). First, a local orientation angle map was calculated for each pixel using the OrientationJ plugin. We used 'cubic spline gradient' method and tensor sigma

parameter of 6 pixels (0.4 μm). The new origin of coordinates was specified by selecting the

centrosome position in the corresponding channel, or the brightest spot in case of centrinone

treatment. Radial local orientation angle was calculated as a difference between the local orientation

angle and the angle of the vector drawn from the new origin of coordinates to the current pixel

position. A radial map image was calculated as an absolute value of the cosine of the radial local

orientation angle at each pixel providing values between zero and one. A non-radial map image was

calculated as one minus the radial map. Both maps were multiplied with the original image to account

for different signal intensities; the two maps illustrate separated radial and non-radial image

components.

Live cell imaging and analysis

Live fluorescent imaging was performed with spinning disk confocal microscopy on inverted research

microscope Nikon EclipseTi-E (Nikon), equipped with the Perfect Focus System (Nikon), Nikon Plan

Apo VC 60x NA 1.4 and Nikon Plan Apo VC 100x N.A. 1.40 oil objectives (Nikon) and spinning-disc

confocal scanner unit (CSU-X1-A1, Yokogawa). The system was also equipped with ASI motorized stage

with the piezo top plate MS-2000-XYZ (ASI), Photometrics Evolve 512 EMCCD camera (Photometrics)

and controlled by the MetaMorph 7.8 software (Molecular Devices). Vortran Stradus lasers (405 nm

100 mW, 488 nm 150 mW and 642 nm 165 mW) and Cobolt Jive 561 nm 110 mW laser were used as

the light sources. System was equipped with ET-DAPI (49000), ET-GFP (49002), ET-mCherry (49008)

and ET-Cy5 (49006) filter sets (Chroma). 16-bit images were projected onto the EMCCD chip with the

intermediate lens 2.0X (Edmund Optics) at a magnification of 110 nm per pixel (60x objective) and 67

nm per pixel (100x objective). To keep cells at 37°C and 5% CO_2 we used stage top incubator (INUBG2E-

ZILCS, Tokai Hit). Cells were plated on round 25 mm coverslips, which were mounted in Attofluor Cell

Chamber (Thermo fisher). Cells were imaged with a 2 s interval and 200 ms exposure for 1-3 hrs at

10% laser power.

Phase-contrast live cell imaging was performed on a Nikon Ti equipped with a perfect focus system

(Nikon), a super high pressure mercury lamp (C-SHG1, Nikon, Japan), a Plan Apo 60x NA 1.4 (Ph3), a

CoolSNAP HQ2 CCD camera (Photometrics, Tucson, AZ), a motorized stage MS-2000-XYZ with Piezo

Top Plate (ASI, Eugene, OR) and a stage top incubator (Tokai Hit, Japan) for 37°C/5% CO2 incubation.

The microscope setup was controlled by Micro-manager software. Cells were plated on round 25 mm

coverslips, which were mounted in Attofluor Cell Chamber (Thermo fisher), and imaged with a 1 min

interval for ~24 hrs.

For live imaging of nocodazole treatment and washout experiments, cells were incubated with the
medium containing 100 nM SiR-tubulin (Tebu-bio) overnight to image the microtubule network.
Centrinone-treated cells were imaged for a desired period of time prior to the nocodazole treatment,
and then nocodazole was added into the medium at a final concentration of 10 µM while imaging
simultaneously. Culture medium was carefully removed when microtubules were completely
depolymerized and washed with prewarmed medium six times to let microtubules regrow. GFP-
CDK5RAP2 and SiR-tubulin imaging was performed with a 2 s interval with 200 ms exposure for 1-3
hrs in total, and maximum intensity projections, contrast adjustment and further processing was
performed using ImageJ.

FRAP

FRAP experiments were performed on the spinning disc microscope describe above, equipped with
iLas platform and using Targeted Laser Action options of iLas and controlled with iLas software (Roper
Scientific, now Gataca Systems). Photobleaching in the GFP channel was performed with the 488 nm
laser. For the FRAP analysis, Polygon ROIs were set in photobleached and non-bleached regions as
well as in the background. The average fluorescence intensity was measured using ImageJ for each
frame, the background intensity was subtracted from the bleached and non-bleached areas and
normalized to the average of the frames acquired prior to the bleach. The mean fluorescence
intensities of the images before photobleaching were set as 100%, and the subsequent relative
recovery percentages were calculated. Time lapse acquisitions were corrected for drift with the
ImageJ plugins Template Matching.

motor-PAINT and analysis

For motor-PAINT, a protocol published previously (Tas et al., 2017) was used, with minor adjustments.
Cells were incubated with 50 nM SiR-tubulin and 500 nM verapamil overnight to allow fields of view
suitable for imaging to be located before the addition of purified GST-DmKHC(1-421)-mNeonGreen.
For nocodazole-treated samples, cells were first incubated with 10 µM nocodazole for 15 minutes at
37°C. A single nocodazole-treated or control sample was then transferred to an imaging chamber, and
cells were subjected to extraction for 1 minute in extraction buffer (BRB80: 80 mM K-Pipes, 1 mM
$MgCl_2$, 1 mM EGTA; pH 6.8, supplemented with 1M sucrose and 0.15% TritonX-100) pre-warmed to
37°C. Pre-warmed fixation buffer (BRB80 supplemented with 2% PFA) was added to this (i.e. final PFA
concentration of 1%) and the solutions were mixed by gentle pipetting for 1 minute. This buffer was

removed and the chamber was washed for 4 times for 1 minute in pre-warmed wash buffer (BRB80
supplemented with 1 µM Taxol) before adding imaging buffer (BRB80 supplemented with 583 µg/mL
catalase, 42 µg/mL glucose oxidase, 1.7% w/v glucose, 1 mM DTT, 1 µM Taxol, and 5mM ATP). An
aliquot of GST-DmKHC(1-421)-mNeonGreen motors was warmed, spun in the Airfuge at 20 psi for 5
minutes in a pre-chilled rotor to remove any aggregates, and then transferred to a clean tube prior to
use. Motors were kept on ice and added locally to cells in 0.3 µl increments.

Imaging was performed immediately after sample preparation at room temperature on a Nikon Ti-E
microscope equipped with a 100x Apo TIRF oil immersion objective (NA. 1.49) and Perfect Focus
System 3 (Nikon). Excitation was achieved with a Lighthub-6 laser combiner (Omicron) containing a
647 nm laser (LuxX 140 mW, Omicron), a 488 nm laser (LuxX 200 mW, Omicron), and optics allowing
for a tunable angle of incidence. Illumination was adjusted for (pseudo-) total internal reflection
fluorescence (TIRF) microscopy. Emission light was separated from excitation light using a quad-band
polychroic mirror (ZT405/488/561/640rpc, Chroma), a quad-band emission filter
(ZET405/488/561/640m, Chroma), and an additional single-band emission filter (ET525/50m for
mNeonGreen emission, Chroma). Detection was achieved using a Hamamatsu Flash 4.0v2 sCMOS
camera. Image stacks were acquired with a 60 ms exposure time, 7% laser power, and 15000-22000
images per field of view. Components were controlled using MicroManager (Edelstein et al., 2014).

Acquired stacks were pre-processed using the Faster Temporal Median ImageJ plugin
(https://github.com/HohlbeinLab/FTM2; (Jabermoradi et al., 2021)) with a window size of 100 frames.
These stacks were then analyzed using Detection of Molecules (DoM) plugin v.1.2.1 for ImageJ
(https://github.com/ekatrukha/DoM_Utrecht), as has been described previously (Chazeau et al.,
2016; Tas et al., 2017). Each image in an acquired stack is convoluted with a two-dimensional Mexican
hat kernel. The resulting intensity histogram is used to create a thresholded mask based on a cut-off
of three standard deviations above the mean. This mask is then subject to rounds of dilution and
erosion to create a filtered mask used to calculate the centroids on the original image. These centroids
are used as initial values to perform unweighted nonlinear least squares fitting with a Levenberg-
Marquardt algorithm to an asymmetric two-dimensional Gaussian point spread function (PSF),
allowing for the sub-pixel localization of particles.

Images were drift-corrected using DoM. The normalized cross-correlation between intermediate
reconstructions of consecutive sub-stacks is used to calculate the drift in x and y between sub-stacks,
which is then linearly interpolated to adjust each individual frame in the stack.

Detected particles were linked into tracks again using DoM, which performs a quicker variant of a
nearest neighbor search, with a maximum distance of 5 pixels (~320nm) between consecutive frames

and no permitted frame gap. Tracks were later filtered to remove those shorter than 4 frames or

longer than 200 frames, those in which an angle between parts of the trajectory exceeded 90 degrees,

and those in which the speed of the motor was less than 100nm/s or more than 1500nm/s.

The particle table was then split into four particle tables corresponding to the four quadrants of the

image with tracks sorted based on their net displacement (i.e., $\triangle$x>0 $\wedge$ $\triangle$y>0; $\triangle$x>0 $\wedge$ $\triangle$y<0; $\triangle$

x<0 $\wedge$ $\triangle$y>0; $\triangle$x<0 $\wedge$ $\triangle$y<0), as described previously (Tas et al., 2017). These directionality-filtered

particle tables were reconstructed using DoM, creating four super-resolved images of microtubule

segments pointing in a similar direction. These were merged with the reconstructed image of all

localizations to determine the direction of each microtubule segment. Each microtubule was manually

assessed to assign it as being plus-end-in or plus-end-out. Microtubules were manually traced with

lines 4 pixels (80 nm) wide, assigned a color based on their orientation, flattened onto the image,

filtered with a Gaussian Blur of radius 2, and finally merged with the reconstructed image of all

localizations.

To quantify the percentage of minus-end-out microtubule length to total microtubule length before

and after nocodazole treatment, the length of each microtubule (determined from kinesin-1

trajectories) in the cell was measured by calculating the Euclidean distance between all subsequent

pairs of points along the microtubule and summed. The ratio was calculated as the total minus-end-

out microtubule length divided by the total microtubule length.

Analysis of PCM cluster dynamics

To represent the motion of PCM clusters during nocodazole treatment, ImageJ plugin

KymoResliceWide v.0.4 (https://github.com/ekatrukha/KymoResliceWide) was used for generating

kymographs from the time lapse images. The velocity of PCM clusters was measured manually using

kymographs starting from the time point when a small PCM cluster moved out of a caMTOC.

Microtubule density around each PCM cluster was determined by measuring the mean fluorescence

intensity of SiR-tubulin in a circular area with a 2 μm radius centered on the PCM cluster and

normalizing it to the mean fluorescence intensity of 20 images prior to nocodazole addition (set as

100%). The moment when a PCM cluster started to move out of the caMTOC was set as the initial time

point (0 min), and the subsequent PCM cluster motion velocity and the relative local microtubule

density at 43 time points were calculated and averaged.

The movement trajectories of PCM clusters were generated using ImageJ plugin TrackMate (version

is 6.0.2). The parameters and the settings used were as following: LoG (Laplacian of Gaussian) detector

with estimated blob diameter: 14.9 µm; thresholding value 12.25; sub-pixel localization was selected.

HyperStack Displayer was selected to overlay the spots and tracks on the current hyperstack window.

Simple LAP tracker was selected to track the distance and time with the linking max distance: 32.0 µm,

gap-closing max distance: 55.0 µm and gap-closing max frame gap: 2. All other parameters and

settings were used as the default.

Computer simulations and analysis

Simulations were performed with Cytosim (version June 2019). Cytosim solves a set of Langevin-

equations that describe the movement of flexible cytoskeletal filaments and associated proteins, such

as molecular motors (Nedelec and Foethke, 2007). The numerical values for the parameters are given

in Table S1. The configuration file is provided as Supplemental file 1.

We defined the following components in the simulation:

Cell shape: We considered a two-dimensional system with a circular cell with a radius of 10 µm. As

commonly used in Cytosim simulations, we set the intracellular viscosity to 1 pN s/ μm^2.

Molecular motors: The binding process of a molecular motor to a microtubule was described by a

binding rate k_{on} and the unbinding from the microtubule by a force-dependent unbinding rate k_{off} =

$k^0_{off} \exp(F/F_d)$. When a motor was engaged with the microtubule it moved along the microtubule with

a linear force-dependent velocity, characterized by $v(F) = v_0 (1 - F/F_s)$. Dynein, as well as kinesin-14

motors moved to the minus end of microtubules.

Microtubule filaments: We used a classical model for microtubule dynamics which is described by a

catastrophe rate, a growth speed, and a shrinkage speed. The growth speed is force-dependent with

a characteristic growing force. For simplicity, we ignored rescue events. The catastrophe rate was set

as $k_{cat} = v_g/L_{MT}$, in which the mean microtubule length L_{MT} was 5 µm. To further restrict the microtubule

length, we set a maximum of 7.5 µm. This limitation was necessary to avoid that long microtubules

were pushing the minus ends to the periphery.

PCM complexes: We described a PCM complex as a bead with a radius of 50 nm. We randomly placed

one microtubule nucleation site and one dynein on the bead. To effectively account for an unspecific

adhesive interaction between PCM complexes, we introduced two molecules that can bind to each

other. One was implemented as a 10 nm Cytosim fiber and the binding partner as a Cytosim hand with

a binding range of 100 nm, binding rate of 10 s^{-1}, force-free unbinding rate of 0.01 s^{-1}, and

characteristic unbinding force of 3 pN. We randomly placed one of each molecule on a PCM complex.

In the simulations with strong adhesive interactions, we increased the number of adhesive binding

molecules on the beads and kept all the other parameters the same. We defined five random

attachment points on the beads and placed to each point five molecules. In this setup each PCM

complex was covered with 25 Cytosim binding filaments and 25 Cytosim binding hands. Therefore,

when two PCM complexes were close to each they formed multiple bonds between each other.

CAMSAP-kin14 complexes: We described a complex consisting of a CAMSAP-stabilized microtubule

end with kinesin 14 motors attached as a Cytosim bead with a radius of 50 nm. We attached five kin14

motors and one microtubule nucleation site randomly on the bead. When we implemented adhesive

interaction between CAMSAP-kin14 complexes, we used exactly the same binding molecules and

arrangements as used for the PCM complexes.

Steric interactions: In all simulations, we considered steric interaction between the beads which either

describe the PCM complex or the CAMSAP-kin14 complexes. All other steric interactions, except with

the cell boundary were ignored.

Simulations and data analysis: We set the time step for the simulation to 0.01 s and simulated for a

total of 30 min, after which a definite steady state was reached. For each configuration, we run 10

simulations and analyzed them afterward to obtain statistics on the emerging structures. From the

last frame of the simulation, we obtained all positions of the complexes and calculated the mean

position, which defines the center of mass. We subtracted the center of mass from all positions and

derived all distances of the complexes to the center of mass. For a few examples, we determined the

empirical cumulative distribution of the distances to the center of mass. We used the standard NumPy

functions to determine the mean and standard deviation of the distances to the center of mass for

each simulation.

Statistical analysis

955   All statistical analyses were performed using GraphPad Prism 9. Statistical details for each experiment

956   can be found in the corresponding figure legends and supporting files.

Data and software availability

All mentioned ImageJ plugins have source code available and are licensed under open-source GNU

GPL v3 license. The source data for the original Western blots are available within the paper.

Acknowledgements

We thank Lynne Cassimeris (Lehigh University, USA), Pierre Gönczy and Didier Trono (EPFL,

Switzerland), Dr. Duane Compton (Geisel School of Medicine at Dartmouth, USA) and Dr. Laurence

Pelletier (Lunenfeld-Tanenbaum Research Institute, Canada) for the gift of materials and Ilya Grigoriev

and Eugene Katrukha (Biology Imaging Center, Utrecht University) for the help with imaging and image

analysis. This work was supported by China Scholarship Council scholarships to Fangrui Chen, Jingchao

Wu and Chao Yang, the Netherlands Organization for Scientific Research Spinoza prize to A.A, as well

as the European Research Council Consolidator Grant 819219 to L.C.K. and the Eindhoven-

Wageningen-Utrecht Alliance (www.ewuu.nl) that supports the Center for Living Technologies.

Competing financial interests

The authors declare no competing financial interests.

References

Abal, M., M. Piel, V. Bouckson-Castaing, M. Mogensen, J.B. Sibarita, and M. Bornens. 2002.
Microtubule release from the centrosome in migrating cells. *J Cell Biol*. 159:731-737.
Akhmanova, A., C.C. Hoogenraad, K. Drabek, T. Stepanova, B. Dortland, T. Verkerk, W. Vermeulen,
B.M. Burgering, C.I. De Zeeuw, F. Grosveld, and N. Galjart. 2001. Clasps are CLIP-115 and -
170 associating proteins involved in the regional regulation of microtubule dynamics in
motile fibroblasts. *Cell*. 104:923-935.
Balczon, R., C.E. Varden, and T.A. Schroer. 1999. Role for microtubules in centrosome doubling in
Chinese hamster ovary cells. *Cell Motil Cytoskeleton*. 42:60-72.
Balestra, F.R., A. Dominguez-Calvo, B. Wolf, C. Busso, A. Buff, T. Averink, M. Lipsanen-Nyman, P.
Huertas, R.M. Rios, and P. Gonczy. 2021. TRIM37 prevents formation of centriolar protein
assemblies by regulating Centrobin. *Elife*. 10:e62640.
Belyy, V., M.A. Schlager, H. Foster, A.E. Reimer, A.P. Carter, and A. Yildiz. 2016. The mammalian
dynein-dynactin complex is a strong opponent to kinesin in a tug-of-war competition. *Nat
Cell Biol*. 18:1018-1024.
Berger, F., S. Klumpp, and R. Lipowsky. 2019. Force-Dependent Unbinding Rate of Molecular Motors
from Stationary Optical Trap Data. *Nano Lett*. 19:2598-2602.
Brenner, S., F. Berger, L. Rao, M.P. Nicholas, and A. Gennerich. 2020. Force production of human
cytoplasmic dynein is limited by its processivity. *Sci Adv*. 6:eaaz4295.
Burakov, A., O. Kovalenko, I. Semenova, O. Zhapparova, E. Nadezhdina, and V. Rodionov. 2008.
Cytoplasmic dynein is involved in the retention of microtubules at the centrosome in
interphase cells. *Traffic*. 9:472-480.
Burakov, A., E. Nadezhdina, B. Slepchenko, and V. Rodionov. 2003. Centrosome positioning in
interphase cells. *J Cell Biol*. 162:963-969.
Carter, N.J., and R.A. Cross. 2005. Mechanics of the kinesin step. *Nature*. 435:308-312.
Cassimeris, L., and J. Morabito. 2004. TOGp, the human homolog of XMAP215/Dis1, is required for
centrosome integrity, spindle pole organization, and bipolar spindle assembly. *Mol Biol Cell*.
15:1580-1590.
Charrasse, S., M. Schroeder, C. Gauthier-Rouviere, F. Ango, L. Cassimeris, D.L. Gard, and C. Larroque.
1998. The TOGp protein is a new human microtubule-associated protein homologous to the
Xenopus XMAP215. *J Cell Sci*. 111 (Pt 10):1371-1383.
Chazeau, A., E.A. Katrukha, C.C. Hoogenraad, and L.C. Kapitein. 2016. Studying neuronal microtubule
organization and microtubule-associated proteins using single molecule localization
microscopy. *Methods Cell Biol*. 131:127-149.
Chen, C.T., H. Hehnly, Q. Yu, D. Farkas, G. Zheng, S.D. Redick, H.F. Hung, R. Samtani, A. Jurczyk, S.
Akbarian, C. Wise, A. Jackson, M. Bober, Y. Guo, C. Lo, and S. Doxsey. 2014. A unique set of
centrosome proteins requires pericentrin for spindle-pole localization and spindle
orientation. *Curr Biol*. 24:2327-2334.
Chi, W., G. Wang, G. Xin, Q. Jiang, and C. Zhang. 2021. PLK4-phosphorylated NEDD1 facilitates
cartwheel assembly and centriole biogenesis initiations. *J Cell Biol*. 220:e202002151.
Chinen, T., S. Yamamoto, Y. Takeda, K. Watanabe, K. Kuroki, K. Hashimoto, D. Takao, and D.
Kitagawa. 2020. NuMA assemblies organize microtubule asters to establish spindle bipolarity
in acentrosomal human cells. *EMBO J*. 39:e102378.
Chinen, T., K. Yamazaki, K. Hashimoto, K. Fujii, K. Watanabe, Y. Takeda, S. Yamamoto, Y. Nozaki, Y.
Tsuchiya, D. Takao, and D. Kitagawa. 2021. Centriole and PCM cooperatively recruit CEP192
to spindle poles to promote bipolar spindle assembly. *J Cell Biol*. 220:e202006085.
Choi, Y.K., P. Liu, S.K. Sze, C. Dai, and R.Z. Qi. 2010. CDK5RAP2 stimulates microtubule nucleation by
the gamma-tubulin ring complex. *J Cell Biol*. 191:1089-1095.

Chong, W.M., W.J. Wang, C.H. Lo, T.Y. Chiu, T.J. Chang, Y.P. Liu, B. Tanos, G. Mazo, M.B. Tsou, W.N. Jane, T.T. Yang, and J.C. Liao. 2020. Super-resolution microscopy reveals coupling between mammalian centriole subdistal appendages and distal appendages. *Elife*. 9:e53580.

Cizmecioglu, O., M. Arnold, R. Bahtz, F. Settele, L. Ehret, U. Haselmann-Weiss, C. Antony, and I. Hoffmann. 2010. Cep152 acts as a scaffold for recruitment of Plk4 and CPAP to the centrosome. *Journal of Cell Biology*. 191:731-739.

Conduit, P.T., A. Wainman, and J.W. Raff. 2015. Centrosome function and assembly in animal cells. *Nat Rev Mol Cell Biol*. 16:611-624.

Cytrynbaum, E.N., V. Rodionov, and A. Mogilner. 2004. Computational model of dynein-dependent self-organization of microtubule asters. *J Cell Sci*. 117:1381-1397.

Delaval, B., and S.J. Doxsey. 2010. Pericentrin in cellular function and disease. *J Cell Biol*. 188:181-190.

Delgehyr, N., J. Sillibourne, and M. Bornens. 2005. Microtubule nucleation and anchoring at the centrosome are independent processes linked by ninein function. *J Cell Sci*. 118:1565-1575.

Dogterom, M., and B. Yurke. 1997. Measurement of the force-velocity relation for growing microtubules. *Science*. 278:856-860.

Edelstein, A.D., M.A. Tsuchida, N. Amodaj, H. Pinkard, R.D. Vale, and N. Stuurman. 2014. Advanced methods of microscope control using muManager software. *J Biol Methods*. 1:e10.

Efimov, A., A. Kharitonov, N. Efimova, J. Loncarek, P.M. Miller, N. Andreyeva, P. Gleeson, N. Galjart, A.R. Maia, I.X. McLeod, J.R. Yates, 3rd, H. Maiato, A. Khodjakov, A. Akhmanova, and I. Kaverina. 2007. Asymmetric CLASP-dependent nucleation of noncentrosomal microtubules at the trans-Golgi network. *Dev Cell*. 12:917-930.

Fong, C.S., G. Mazo, T. Das, J. Goodman, M. Kim, B.P. O'Rourke, D. Izquierdo, and M.F. Tsou. 2016. 53BP1 and USP28 mediate p53-dependent cell cycle arrest in response to centrosome loss and prolonged mitosis. *Elife*. 5:e16270.

Fong, K.W., Y.K. Choi, J.B. Rattner, and R.Z. Qi. 2008. CDK5RAP2 is a pericentriolar protein that functions in centrosomal attachment of the gamma-tubulin ring complex. *Mol Biol Cell*. 19:115-125.

Fu, J.Y., and D.M. Glover. 2012. Structured illumination of the interface between centriole and peri-centriolar material. *Open Biology*. 2:120104.

Ganem, N.J., and D.A. Compton. 2004. The KinI kinesin Kif2a is required for bipolar spindle assembly through a functional relationship with MCAK. *J Cell Biol*. 166:473-478.

Ganem, N.J., K. Upton, and D.A. Compton. 2005. Efficient mitosis in human cells lacking poleward microtubule flux. *Curr Biol*. 15:1827-1832.

Garbrecht, J., T. Laos, E. Holzer, M. Dillinger, and A. Dammermann. 2021. An acentriolar centrosome at the C. elegans ciliary base. *Curr Biol*. 31:2418-2428.

Gartenmann, L., C.C. Vicente, A. Wainman, Z.A. Novak, B. Sieber, J.H. Richens, and J.W. Raff. 2020. Drosophila Sas-6, Ana2 and Sas-4 self-organise into macromolecular structures that can be used to probe centriole and centrosome assembly. *J Cell Sci*. 133:jcs244574.

Gavilan, M.P., P. Gandolfo, F.R. Balestra, F. Arias, M. Bornens, and R.M. Rios. 2018. The dual role of the centrosome in organizing the microtubule network in interphase. *EMBO Rep*. 19:e45942.

Gibeaux, R., A.Z. Politi, P. Philippsen, and F. Nedelec. 2017. Mechanism of nuclear movements in a multinucleated cell. *Mol Biol Cell*. 28:645-660.

Gimpel, P., Y.L. Lee, R.M. Sobota, A. Calvi, V. Koullourou, R. Patel, K. Mamchaoui, F. Nedelec, S. Shackleton, J. Schmoranzer, B. Burke, B. Cadot, and E.R. Gomes. 2017. Nesprin-1alpha-Dependent Microtubule Nucleation from the Nuclear Envelope via Akap450 Is Necessary for Nuclear Positioning in Muscle Cells. *Curr Biol*. 27:2999-3009.

Goldspink, D.A., C. Rookyard, B.J. Tyrrell, J. Gadsby, J. Perkins, E.K. Lund, N. Galjart, P. Thomas, T. Wileman, and M.M. Mogensen. 2017. Ninein is essential for apico-basal microtubule formation and CLIP-170 facilitates its redeployment to non-centrosomal microtubule organizing centres. *Open Biol*. 7:160274.

Gomez-Ferreria, M.A., M. Bashkurov, A.O. Helbig, B. Larsen, T. Pawson, A.C. Gingras, and L. Pelletier.
2012. Novel NEDD1 phosphorylation sites regulate gamma-tubulin binding and mitotic
spindle assembly. *J Cell Sci*. 125:3745-3751.
Gomez-Ferreria, M.A., U. Rath, D.W. Buster, S.K. Chanda, J.S. Caldwell, D.R. Rines, and D.J. Sharp.
2007. Human Cep192 is required for mitotic centrosome and spindle assembly. *Curr Biol*.
17:1960-1966.
Grigoriev, I., S.M. Gouveia, B. van der Vaart, J. Demmers, J.T. Smyth, S. Honnappa, D. Splinter, M.O.
Steinmetz, J.W. Putney, Jr., C.C. Hoogenraad, and A. Akhmanova. 2008. STIM1 is a MT-plus-
end-tracking protein involved in remodeling of the ER. *Curr Biol*. 18:177-182.
Gros, O.J., H.G.J. Damstra, L.C. Kapitein, A. Akhmanova, and F. Berger. 2021. Dynein self-organizes
while translocating the centrosome in T-cells. *Mol Biol Cell*. 32:855-868.
Haren, L., M.H. Remy, I. Bazin, I. Callebaut, M. Wright, and A. Merdes. 2006. NEDD1-dependent
recruitment of the gamma-tubulin ring complex to the centrosome is necessary for centriole
duplication and spindle assembly. *J Cell Biol*. 172:505-515.
Haren, L., T. Stearns, and J. Luders. 2009. Plk1-dependent recruitment of gamma-tubulin complexes
to mitotic centrosomes involves multiple PCM components. *PLoS One*. 4:e5976.
Hoogenraad, C.C., P. Wulf, N. Schiefermeier, T. Stepanova, N. Galjart, J.V. Small, F. Grosveld, C.I. de
Zeeuw, and A. Akhmanova. 2003. Bicaudal D induces selective dynein-mediated microtubule
minus end-directed transport. *Embo J*. 22:6004-6015.
Jabermoradi, A., S. Yang, M. Gobes, J.P.M. van Duynhoven, and H. J. 2021. Enabling single-molecule
localization microscopy in turbid food emulsions. *bioRxiv*:2021.2003.2003.433739.
Jia, Y., K.W. Fong, Y.K. Choi, S.S. See, and R.Z. Qi. 2013. Dynamic recruitment of CDK5RAP2 to
centrosomes requires its association with dynein. *PLoS One*. 8:e68523.
Jiang, K., S. Hua, R. Mohan, I. Grigoriev, K.W. Yau, Q. Liu, E.A. Katrukha, A.F. Altelaar, A.J. Heck, C.C.
Hoogenraad, and A. Akhmanova. 2014. Microtubule minus-end stabilization by
polymerization-driven CAMSAP deposition. *Dev Cell*. 28:295-309.
Jiang, X., D.B.T. Ho, K. Mahe, J. Mia, G. Sepulveda, M. Antkowiak, L. Jiang, S. Yamada, and L.E. Jao.
2021. Condensation of pericentrin proteins in human cells illuminates phase separation in
centrosome assembly. *J Cell Sci*. 134.
Jonsson, E., M. Yamada, R.D. Vale, and G. Goshima. 2015. Clustering of a kinesin-14 motor enables
processive retrograde microtubule-based transport in plants. *Nat Plants*. 1:15087.
Joukov, V., J.C. Walter, and A. De Nicolo. 2014. The Cep192-organized aurora A-Plk1 cascade is
essential for centrosome cycle and bipolar spindle assembly. *Mol Cell*. 55:578-591.
Khodjakov, A., R.W. Cole, B.R. Oakley, and C.L. Rieder. 2000. Centrosome-independent mitotic
spindle formation in vertebrates. *Curr Biol*. 10:59-67.
Kim, S., and K. Rhee. 2014. Importance of the CEP215-pericentrin interaction for centrosome
maturation during mitosis. *PLoS One*. 9:e87016.
Kim, T.S., J.E. Park, A. Shukla, S. Choi, R.N. Murugan, J.H. Lee, M. Ahn, K. Rhee, J.K. Bang, B.Y. Kim, J.
Loncarek, R.L. Erikson, and K.S. Lee. 2013. Hierarchical recruitment of Plk4 and regulation of
centriole biogenesis by two centrosomal scaffolds, Cep192 and Cep152. *Proc Natl Acad Sci U*
S A. 110:E4849-4857.
Klausner, R.D., J.G. Donaldson, and J. Lippincott-Schwartz. 1992. Brefeldin A: insights into the control
of membrane traffic and organelle structure. *J Cell Biol*. 116:1071-1080.
Klumpp, S., C. Keller, F. Berger, and R. Lipowsky. 2015. Molecular Motors: Cooperative Phenomena
of Multiple Molecular Motors. *In* Multiscale Modeling in Biomechanics and Mechanobiology.
S. De, W. Hwang, and E. Kuhl, editors. Springer London, London. 27-61.
Kohlmaier, G., J. Loncarek, X. Meng, B.F. McEwen, M.M. Mogensen, A. Spektor, B.D. Dynlacht, A.
Khodjakov, and P. Gonczy. 2009. Overly Lona Centrioles and Defective Cell Division upon
Excess of the SAS-4-Related Protein CPAP. *Current Biology*. 19:1012-1018.

Kolano, A., S. Brunet, A.D. Silk, D.W. Cleveland, and M.H. Verlhac. 2012. Error-prone mammalian female meiosis from silencing the spindle assembly checkpoint without normal interkinetochore tension. *Proc Natl Acad Sci U S A*. 109:E1858-1867.

Kole, T.P., Y. Tseng, I. Jiang, J.L. Katz, and D. Wirtz. 2005. Intracellular mechanics of migrating fibroblasts. *Mol Biol Cell*. 16:328-338.

Komarova, Y., G. Lansbergen, N. Galjart, F. Grosveld, G.G. Borisy, and A. Akhmanova. 2005. EB1 and EB3 control CLIP dissociation from the ends of growing microtubules. *Mol Biol Cell*. 16:5334-5345.

Lambrus, B.G., V. Daggubati, Y. Uetake, P.M. Scott, K.M. Clutario, G. Sluder, and A.J. Holland. 2016. A USP28-53BP1-p53-p21 signaling axis arrests growth after centrosome loss or prolonged mitosis. *J Cell Biol*. 214:143-153.

Lansbergen, G., Y. Komarova, M. Modesti, C. Wyman, C.C. Hoogenraad, H.V. Goodson, R.P. Lemaitre, D.N. Drechsel, E. van Munster, T.W.J. Gadella Jr., F. Grosveld, N. Galjart, G.G. Borisy, and A. Akhmanova. 2004. Conformational changes in CLIP-170 regulate its binding to microtubules and dynactin localisation. *J Cell Biol*. 166:1003-1014.

Lawo, S., M. Bashkurov, M. Mullin, M.G. Ferreria, R. Kittler, B. Habermann, A. Tagliaferro, I. Poser, J.R. Hutchins, B. Hegemann, D. Pinchev, F. Buchholz, J.M. Peters, A.A. Hyman, A.C. Gingras, and L. Pelletier. 2009. HAUS, the 8-subunit human Augmin complex, regulates centrosome and spindle integrity. *Curr Biol*. 19:816-826.

Lawo, S., M. Hasegan, G.D. Gupta, and L. Pelletier. 2012. Subdiffraction imaging of centrosomes reveals higher-order organizational features of pericentriolar material. *Nat Cell Biol*. 14:1148-1158.

Lechler, T., and E. Fuchs. 2007. Desmoplakin: an unexpected regulator of microtubule organization in the epidermis. *J Cell Biol*. 176:147-154.

Lee, K., and K. Rhee. 2011. PLK1 phosphorylation of pericentrin initiates centrosome maturation at the onset of mitosis. *J Cell Biol*. 195:1093-1101.

Lee, S., and K. Rhee. 2010. CEP215 is involved in the dynein-dependent accumulation of pericentriolar matrix proteins for spindle pole formation. *Cell Cycle*. 9:774-783.

Letort, G., F. Nedelec, L. Blanchoin, and M. Thery. 2016. Centrosome centering and decentering by microtubule network rearrangement. *Mol Biol Cell*. 27:2833-2843.

Liang, X., M. Kokes, R.D. Fetter, M.D. Sallee, A.W. Moore, J.L. Feldman, and K. Shen. 2020. Growth cone-localized microtubule organizing center establishes microtubule orientation in dendrites. *Elife*. 9:e56547.

Lin, Y.N., C.T. Wu, Y.C. Lin, W.B. Hsu, C.J. Tang, C.W. Chang, and T.K. Tang. 2013. CEP120 interacts with CPAP and positively regulates centriole elongation. *J Cell Biol*. 202:211-219.

Luders, J., U.K. Patel, and T. Stearns. 2006. GCP-WD is a gamma-tubulin targeting factor required for centrosomal and chromatin-mediated microtubule nucleation. *Nat Cell Biol*. 8:137-147.

Magescas, J., S. Eskinazi, M.V. Tran, and J.L. Feldman. 2021. Centriole-less pericentriolar material serves as a microtubule organizing center at the base of C. elegans sensory cilia. *Curr Biol*. 31:2410-2417.

Martin, M., A. Veloso, J. Wu, E.A. Katrukha, and A. Akhmanova. 2018. Control of endothelial cell polarity and sprouting angiogenesis by non-centrosomal microtubules. *Elife*. 7:e33864.

Meitinger, F., J.V. Anzola, M. Kaulich, A. Richardson, J.D. Stender, C. Benner, C.K. Glass, S.F. Dowdy, A. Desai, A.K. Shiau, and K. Oegema. 2016. 53BP1 and USP28 mediate p53 activation and G1 arrest after centrosome loss or extended mitotic duration. *J Cell Biol*. 214:155-166.

Meitinger, F., M. Ohta, K.Y. Lee, S. Watanabe, R.L. Davis, J.V. Anzola, R. Kabeche, D.A. Jenkins, A.K. Shiau, A. Desai, and K. Oegema. 2020. TRIM37 controls cancer-specific vulnerability to PLK4 inhibition. *Nature*. 585:440-446.

Meng, W., Y. Mushika, T. Ichii, and M. Takeichi. 2008. Anchorage of microtubule minus ends to adherens junctions regulates epithelial cell-cell contacts. *Cell*. 135:948-959.

Mennella, V., D.A. Agard, B. Huang, and L. Pelletier. 2014. Amorphous no more: subdiffraction view
of the pericentriolar material architecture. *Trends Cell Biol*. 24:188-197.
Mennella, V., B. Keszthelyi, K.L. McDonald, B. Chhun, F. Kan, G.C. Rogers, B. Huang, and D.A. Agard.
2012. Subdiffraction-resolution fluorescence microscopy reveals a domain of the
centrosome critical for pericentriolar material organization. *Nat Cell Biol*. 14:1159-1168.
Mimori-Kiyosue, Y., I. Grigoriev, G. Lansbergen, H. Sasaki, C. Matsui, F. Severin, N. Galjart, F.
Grosveld, I. Vorobjev, S. Tsukita, and A. Akhmanova. 2005. CLASP1 and CLASP2 bind to EB1
and regulate microtubule plus-end dynamics at the cell cortex. *J Cell Biol*. 168:141-153.
Mogensen, M.M., A. Malik, M. Piel, V. Bouckson-Castaing, and M. Bornens. 2000. Microtubule
minus-end anchorage at centrosomal and non-centrosomal sites: the role of ninein. *J Cell Sci*.
113 (Pt 17):3013-3023.
Muroyama, A., and T. Lechler. 2017. Microtubule organization, dynamics and functions in
differentiated cells. *Development*. 144:3012-3021.
Nashchekin, D., A.R. Fernandes, and D. St Johnston. 2016. Patronin/Shot Cortical Foci Assemble the
Noncentrosomal Microtubule Array that Specifies the Drosophila Anterior-Posterior Axis.
Dev Cell. 38:61-72.
Nedelec, F., and D. Foethke. 2007. Collective Langevin dynamics of flexible cytoskeletal fibers. *New
Journal of Physics*. 9:427-427.
Nijenhuis, W., M.M.P. van Grinsven, and L.C. Kapitein. 2020. An optimized toolbox for the
optogenetic control of intracellular transport. *J Cell Biol*. 219:e201907149.
Noordstra, I., Q. Liu, W. Nijenhuis, S. Hua, K. Jiang, M. Baars, S. Remmelzwaal, M. Martin, L.C.
Kapitein, and A. Akhmanova. 2016. Control of apico-basal epithelial polarity by the
microtubule minus-end-binding protein CAMSAP3 and spectraplakin ACF7. *J Cell Sci*.
129:4278-4288.
Oddoux, S., K.J. Zaal, V. Tate, A. Kenea, S.A. Nandkeolyar, E. Reid, W. Liu, and E. Ralston. 2013.
Microtubules that form the stationary lattice of muscle fibers are dynamic and nucleated at
Golgi elements. *J Cell Biol*. 203:205-213.
Ohashi, K.G., L. Han, B. Mentley, J. Wang, J. Fricks, and W.O. Hancock. 2019. Load-dependent
detachment kinetics plays a key role in bidirectional cargo transport by kinesin and dynein.
Traffic. 20:284-294.
Paz, J., and J. Luders. 2017. Microtubule-Organizing Centers: Towards a Minimal Parts List. *Trends
Cell Biol*:S0962-8924(0917)30187-30183.
Pollock, R., R. Issner, K. Zoller, S. Natesan, V.M. Rivera, and T. Clackson. 2000. Delivery of a stringent
dimerizer-regulated gene expression system in a single retroviral vector. *Proc Natl Acad Sci U
S A*. 97:13221-13226.
Prosser, S.L., and L. Pelletier. 2020. Centriolar satellite biogenesis and function in vertebrate cells. *J
Cell Sci*. 133:jcs239566.
Purohit, A., S.H. Tynan, R. Vallee, and S.J. Doxsey. 1999. Direct interaction of pericentrin with
cytoplasmic dynein light intermediate chain contributes to mitotic spindle organization. *J
Cell Biol*. 147:481-492.
Pyrpassopoulos, S., H. Shuman, and E.M. Ostap. 2020. Modulation of Kinesin's Load-Bearing Capacity
by Force Geometry and the Microtubule Track. *Biophys J*. 118:243-253.
Raff, J.W. 2019. Phase Separation and the Centrosome: A Fait Accompli? *Trends Cell Biol*. 29:612-
622.
Ran, F.A., P.D. Hsu, J. Wright, V. Agarwala, D.A. Scott, and F. Zhang. 2013. Genome engineering using
the CRISPR-Cas9 system. *Nat Protoc*. 8:2281-2308.
Redwine, W.B., M.E. DeSantis, I. Hollyer, Z.M. Htet, P.T. Tran, S.K. Swanson, L. Florens, M.P.
Washburn, and S.L. Reck-Peterson. 2017. The human cytoplasmic dynein interactome
reveals novel activators of motility. *Elife*. 6:e28257.
Rios, R.M. 2014. The centrosome-Golgi apparatus nexus. *Philos Trans R Soc Lond B Biol Sci*.
369:20130462.

Rivero, S., J. Cardenas, M. Bornens, and R.M. Rios. 2009. Microtubule nucleation at the cis-side of
the Golgi apparatus requires AKAP450 and GM130. *EMBO J*. 28:1016-1028.
Sallee, M.D., and J.L. Feldman. 2021. Microtubule organization across cell types and states. *Curr Biol*.
31:R506-R511.
Sanchez, A.D., T.C. Branon, L.E. Cote, A. Papagiannakis, X. Liang, M.A. Pickett, K. Shen, C. Jacobs-
Wagner, A.Y. Ting, and J.L. Feldman. 2021. Proximity labeling reveals non-centrosomal
microtubule-organizing center components required for microtubule growth and
localization. *Curr Biol*. 31:3586-3600 e3511.
Schlager, M.A., A. Serra-Marques, I. Grigoriev, L.F. Gumy, M. Esteves da Silva, P.S. Wulf, A.
Akhmanova, and C.C. Hoogenraad. 2014. Bicaudal d family adaptor proteins control the
velocity of Dynein-based movements. *Cell Rep*. 8:1248-1256.
Schmidt, T.I., J. Kleylein-Sohn, J. Westendorf, M. Le Clech, S.B. Lavoie, Y.D. Stierhof, and E.A. Nigg.
2009. Control of centriole length by CPAP and CP110. *Curr Biol*. 19:1005-1011.
Shinohara, H., N. Sakayori, M. Takahashi, and N. Osumi. 2013. Ninein is essential for the
maintenance of the cortical progenitor character by anchoring the centrosome to
microtubules. *Biol Open*. 2:739-749.
Sonnen, K.F., A.M. Gabryjonczyk, E. Anselm, Y.D. Stierhof, and E.A. Nigg. 2013. Human Cep192 and
Cep152 cooperate in Plk4 recruitment and centriole duplication. *J Cell Sci*. 126:3223-3233.
Sonnen, K.F., L. Schermelleh, H. Leonhardt, and E.A. Nigg. 2012. 3D-structured illumination
microscopy provides novel insight into architecture of human centrosomes. *Biology Open*.
1:965-976.
Spektor, A., W.Y. Tsang, D. Khoo, and B.D. Dynlacht. 2007. Cep97 and CP110 suppress a cilia
assembly program. *Cell*. 130:678-690.
Splinter, D., M.E. Tanenbaum, A. Lindqvist, D. Jaarsma, A. Flotho, K.L. Yu, I. Grigoriev, D. Engelsma,
E.D. Haasdijk, N. Keijzer, J. Demmers, M. Fornerod, F. Melchior, C.C. Hoogenraad, R.H.
Medema, and A. Akhmanova. 2010. Bicaudal D2, dynein and kinesin-1 associate with nuclear
pore complexes and regulate centrosome and nuclear positioning during mitotic entry. *PLoS*
Biol. 8:e1000350.
Steinman, J.B., C.C. Santarossa, R.M. Miller, L.S. Yu, A.S. Serpinskaya, H. Furukawa, S. Morimoto, Y.
Tanaka, M. Nishitani, M. Asano, R. Zalyte, A.E. Ondrus, A.G. Johnson, F. Ye, M.V. Nachury, Y.
Fukase, K. Aso, M.A. Foley, V.I. Gelfand, J.K. Chen, A.P. Carter, and T.M. Kapoor. 2017.
Chemical structure-guided design of dynapyrazoles, cell-permeable dynein inhibitors with a
unique mode of action. *Elife*. 6:e25174.
Stolz, A., N. Ertych, and H. Bastians. 2015. A phenotypic screen identifies microtubule plus end
assembly regulators that can function in mitotic spindle orientation. *Cell Cycle*. 14:827-837.
Takahashi, M., A. Yamagiwa, T. Nishimura, H. Mukai, and Y. Ono. 2002. Centrosomal proteins CG-
NAP and kendrin provide microtubule nucleation sites by anchoring gamma-tubulin ring
complex. *Mol Biol Cell*. 13:3235-3245.
Tanaka, N., W. Meng, S. Nagae, and M. Takeichi. 2012. Nezha/CAMSAP3 and CAMSAP2 cooperate in
epithelial-specific organization of noncentrosomal microtubules. *Proc Natl Acad Sci U S A*.
109:20029-20034.
Tang, C.J.C., R.H. Fu, K.S. Wu, W.B. Hsu, and T.K. Tang. 2009. CPAP is a cell-cycle regulated protein
that controls centriole length. *Nat Cell Biol*. 11:825-831.
Tas, R.P., A. Chazeau, B.M.C. Cloin, M.L.A. Lambers, C.C. Hoogenraad, and L.C. Kapitein. 2017.
Differentiation between Oppositely Oriented Microtubules Controls Polarized Neuronal
Transport. *Neuron*. 96:1264-1271.
Tibelius, A., J. Marhold, H. Zentgraf, C.E. Heilig, H. Neitzel, B. Ducommun, A. Rauch, A.D. Ho, J.
Bartek, and A. Kramer. 2009. Microcephalin and pericentrin regulate mitotic entry via
centrosome-associated Chk1. *J Cell Biol*. 185:1149-1157.

Toya, M., S. Kobayashi, M. Kawasaki, G. Shioi, M. Kaneko, T. Ishiuchi, K. Misaki, W. Meng, and M.
Takeichi. 2016. CAMSAP3 orients the apical-to-basal polarity of microtubule arrays in
epithelial cells. *Proc Natl Acad Sci U S A*. 113:332-337.
Vergarajauregui, S., R. Becker, U. Steffen, M. Sharkova, T. Esser, J. Petzold, F. Billing, M.S. Kapiloff, G.
Schett, I. Thievessen, and F.B. Engel. 2020. AKAP6 orchestrates the nuclear envelope
microtubule-organizing center by linking golgi and nucleus via AKAP9. *Elife*. 9:e61669.
Vinopal, S., M. Cernohorska, V. Sulimenko, T. Sulimenko, V. Vosecka, M. Flemr, E. Draberova, and P.
Draber. 2012. gamma-Tubulin 2 nucleates microtubules and is downregulated in mouse
early embryogenesis. *PLoS One*. 7:e29919.
Wang, G., Q. Chen, X. Zhang, B. Zhang, X. Zhuo, J. Liu, Q. Jiang, and C. Zhang. 2013. PCM1 recruits
Plk1 to the pericentriolar matrix to promote primary cilia disassembly before mitotic entry. *J
Cell Sci*. 126:1355-1365.
Wang, S., D. Wu, S. Quintin, R.A. Green, D.K. Cheerambathur, S.D. Ochoa, A. Desai, and K. Oegema.
2015. NOCA-1 functions with gamma-tubulin and in parallel to Patronin to assemble non-
centrosomal microtubule arrays in C. elegans. *Elife*. 4:e08649.
Watanabe, S., F. Meitinger, A.K. Shiau, K. Oegema, and A. Desai. 2020. Centriole-independent
mitotic spindle assembly relies on the PCNT-CDK5RAP2 pericentriolar matrix. *J Cell Biol*.
219:e202006010.
Wong, Y.L., J.V. Anzola, R.L. Davis, M. Yoon, A. Motamedi, A. Kroll, C.P. Seo, J.E. Hsia, S.K. Kim, J.W.
Mitchell, B.J. Mitchell, A. Desai, T.C. Gahman, A.K. Shiau, and K. Oegema. 2015. Cell biology.
Reversible centriole depletion with an inhibitor of Polo-like kinase 4. *Science*. 348:1155-
1160.
Woodruff, J.B., B. Ferreira Gomes, P.O. Widlund, J. Mahamid, A. Honigmann, and A.A. Hyman. 2017.
The Centrosome Is a Selective Condensate that Nucleates Microtubules by Concentrating
Tubulin. *Cell*. 169:1066-1077.
Wu, J., C. de Heus, Q. Liu, B.P. Bouchet, I. Noordstra, K. Jiang, S. Hua, M. Martin, C. Yang, I. Grigoriev,
E.A. Katrukha, A.F. Altelaar, C.C. Hoogenraad, R.Z. Qi, J. Klumperman, and A. Akhmanova.
2016. Molecular Pathway of Microtubule Organization at the Golgi Apparatus. *Dev Cell*.
39:44-60.
Yang, C., J. Wu, C. de Heus, I. Grigoriev, N. Liv, Y. Yao, I. Smal, E. Meijering, J. Klumperman, R.Z. Qi,
and A. Akhmanova. 2017. EB1 and EB3 regulate microtubule minus end organization and
Golgi morphology. *J Cell Biol*. 216:3179-3198.
Yang, R., and J.L. Feldman. 2015. SPD-2/CEP192 and CDK Are Limiting for Microtubule-Organizing
Center Function at the Centrosome. *Curr Biol*. 25:1924-1931.
Yeow, Z.Y., B.G. Lambrus, R. Marlow, K.H. Zhan, M.A. Durin, L.T. Evans, P.M. Scott, T. Phan, E. Park,
L.A. Ruiz, D. Moralli, E.G. Knight, L.M. Badder, D. Novo, S. Haider, C.M. Green, A.N.J. Tutt, C.J.
Lord, J.R. Chapman, and A.J. Holland. 2020. Targeting TRIM37-driven centrosome
dysfunction in 17q23-amplified breast cancer. *Nature*. 585:447-452.
Zheng, Y., R.A. Buchwalter, C. Zheng, E.M. Wight, J.V. Chen, and T.L. Megraw. 2020. A perinuclear
microtubule-organizing centre controls nuclear positioning and basement membrane
secretion. *Nat Cell Biol*. 22:297-309.
Zhu, F., S. Lawo, A. Bird, D. Pinchev, A. Ralph, C. Richter, T. Muller-Reichert, R. Kittler, A.A. Hyman,
and L. Pelletier. 2008. The mammalian SPD-2 ortholog Cep192 regulates centrosome
biogenesis. *Curr Biol*. 18:136-141.
Zhu, X.D., and I. Kaverina. 2013. Golgi as an MTOC: making microtubules for its own good.
Histochemistry and Cell Biology. 140:361-367.

1319

 Figure legends.

Figure 1. Formation and characterization of caMTOCs in AKAP450/CAMSAP2 knockout cells

(A) Immunofluorescence images of control or centrinone-treated wild type (WT) RPE1 cells stained for

centrioles (CEP135, red; centrin, green). The zooms of the boxed area show the centrioles stained with

the indicated markers.

(B) Quantification shows the percentage of cells with centrioles before and after the centrinone

treatment. 350 cells (n=7 fields of view) analyzed for each measurement in three independent

experiments. The statistical significance was determined by unpaired two-tailed Mann-Whitney test

in Prism 9.1 (***p<0.001). Values represent mean±SD.

(C) Immunofluorescence images of centrinone-treated WT RPE1 cells stained for pericentrin (PCNT,

green) and the Golgi marker GM130 (red). Inset shows the merged image of the boxed area.

(D) Diagrams of the microtubule organization in WT and knockout (KO) cells used.

(E) Immunofluorescence images of control and centrinone-treated WT and knockout RPE1 cell lines

stained for pericentrin (green) and microtubules (α-tubulin, red). Enlargements on the right show the

boxed areas.

(F) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells

stained for different PCM components as indicated and imaged by STED microscopy.

(G) Quantification of the length and width of cylindrical PCM clusters. n=65 cells analyzed in three

independent experiments. Values represent mean±SD.

(H) (Top left) Two frames of time-lapse images of centrinone-treated AKAP450/CAMSAP2 knockout

RPE1 cells stably expressing GFP-CDK5RAP2 prior to FRAP experiments. (Top right) Schemes show

regions of caMTOC where photobleaching was performed. (Middle) Kymographs illustrating

fluorescence of unbleached caMTOC (No FRAP), fully photobleached caMTOC (Whole FRAP) and

partially photobleached caMTOC (Partial FRAP). (Bottom) Time-lapse images illustrating partial FRAP

of a caMTOC. Time is min:sec.

(I) Normalized fluorescence intensity as a function of time. The blue line shows averaged intensity

traces of unbleached caMTOCs (No FRAP), the black line shows averaged intensity traces of fully

photobleached caMTOCs (Whole FRAP), the red line shows averaged intensity traces of whole

caMTOC that were partially photobleached (whole caMTOC intensity, Partial FRAP) and the green line

shows averaged intensity traces of the photobleached region of the partially photobleached caMTOC

(FRAP region intensity, Partial FRAP). n=3 for No FRAP, 3 for Whole FRAP, 5 for Partial FRAP (whole

caMTOC intensity) and 5 for Partial FRAP (FRAP region intensity); time-lapse images of ~1600

timepoints with 2 s interval were analyzed for each measurement. Values are mean±SD.

1353

Figure 2. Molecular composition of caMTOCs in AKAP450/CAMSAP2 knockout cells

All the cells used in this figure are AKAP450/CAMSAP2 knockout cells, except for panel E as indicated.

(A-C) Immunofluorescence images of control or centrinone-treated AKAP450/CAMSAP2 knockout

RPE1 cells stained for and depleted of the indicated proteins.

(D, F-I) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells

stained for and depleted of the indicated proteins.

(E) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2/p53 knockout and

AKAP450/CAMSAP2/p53/pericentrin knockout RPE1 cells stained as indicated.

In panels A-I, insets show enlargements of the merged channels of the boxed areas, and dashed lines

indicate cell edges.

(J) Quantification of the main PCM organization types, as described in Figure 1- figure supplement 2A,

for cells prepared as described in panel A-C, E-I. Numbers on the histogram show the percentages.

1293 (-CentB), 1547(+CentB), 2021(siLuci), 1822(siCEP152), 1345(siCEP192), 1161(siNEDD1), 2302

(AKAP450/CAMSAP2/p53/PCNT knockout (PCNT KO)), 2264(siCDK5RAP2), 2510(siγ-tubulin),

2408(siNIN) and 2526(siDHC) cells were analyzed for each measurement in three independent

experiments (n=3). Values represent mean±SD.

(K) Summarizing table of PCM localization and the depletion effects on caMTOC formation in

AKAP450/CAMSAP2 knockout RPE1 cells. NT, not tested.

1372

Figure 3. Microtubule organization in acentriolar cells missing different caMTOC components

(A) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells

depleted of the indicated proteins and stained for microtubules (α-tubulin, red) and different PCM

proteins (green). Insets show enlargements of the merged channels of the boxed areas and dashed

lines show cell boundaries.

(B) Quantification of the normalized overall microtubule intensity for the indicated conditions. The

number of cells analyzed in three independent experiments: n=56 (siLuci), 45 (siPCNT), 33

(siCDK5RAP2), 36 (siNinein), 43 (siγ-tubulin) and 28 (siDHC). The statistical significance was

determined by unpaired two-tailed Mann-Whitney test in Prism 9.1 (***p<0.001). Values represent

mean±SD.

(C) Microtubule images were split into a radial and non-radial components (heat maps) based on

microtubule orientation in relation to the PCM clusters or the brightest point, as described in

Methods.

(D) Quantification of the proportion of the non-radial microtubules shown in panel C (see Methods

for details). The number of cells analyzed for each measurement in three independent experiments:

n=25 (siLuci), 43 (siPCNT), 32 (siCDK5RAP2), 34 (siNinein), 37 (siγ-tubulin) and 25 (siDHC). The

statistical significance was determined by unpaired two-tailed Mann-Whitney test in Prism 9.1

(***p<0.001). Values represent mean±SD.

(E) Diagram illustrating the distribution of PCM clusters and microtubule organization upon the

depletion of the indicated proteins in centrinone-treated AKAP450/CAMSAP2 knockout cells.

1393

Figure 4. Microtubule- and dynein-dependent disassembly of caMTOCs

(A) Diagram illustrating different order of cell treatments with nocodazole (Noco) and/or dynapyrazole

A (Dyna, 5 µM) and the time points when the cells were fixed.

(B) Immunofluorescence staining of centrinone-treated AKAP450/CAMSAP2 knockout cells treated as

shown in panel A. Dashed red circles represent the areas occupied by PCM clusters in each condition.

(C) Quantification of the area occupied by PCM clusters in each condition, as shown in panels A and B.

n=35-53 cells analyzed for each measurement in three independent experiments. The statistical

significance was determined by unpaired two-tailed Mann-Whitney test in Prism 9.1 (not significant

(NS), *P*<0.12; **P*<0.033; ***P*<0.001). Values are represented as mean±SD.

(D) Western blot showing that 3 hrs treatment with dynaprazole A does not affect the expression of

the endogenous dynein heavy chain and the dynactin large subunit p150Glued.

(E) Time-lapse images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stably

expressing GFP-CDK5RAP2. Microtubules were visualized by treating cells with 100 nM SiR-tubulin

overnight. Red arrows show the immobilized PCM clusters at indicated timepoints. Time is min:sec.

Time-lapse images of the same cell prior to the nocodazole treatment were shown in Figure 1 - figure

supplement 3B.

(F) (Top) Kymograph illustrating the motility of PCM clusters during microtubule disassembly with

nocodazole. (Bottom) Measurements of the normalized microtubule (SiR-tubulin) fluorescence

intensity (red plot, left Y-axis) and the instantaneous velocity of GFP-CDK5RAP2 clusters (green plot,

right Y-axis) during the movement of GFP-CDK5RAP2 clusters away from caMTOC. Microtubule density

around each PCM cluster was determined by measuring mean fluorescence intensity of SiR-tubulin in

a circular area with a 2 μm radius centered on the PCM cluster and normalizing it to the mean

fluorescence intensity of 20 images prior to nocodazole addition (set as 100%). The moment when a

PCM cluster started to move out of the caMTOC was set as the initial time point (0 min) for this cluster,

and the subsequent PCM cluster motion velocity and the relative local microtubule density of 43 time

points were calculated and averaged. n=12 clusters were analyzed in each condition. Values are

represented as mean±SD.

(G) Motor-PAINT images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells before and

after nocodazole treatment. Plus-end-out microtubules are shown in white whereas minus-end-out

microtubules are shown in magenta. Asterisks represent the putative position of caMTOC.

(H) Summarizing diagram illustrating microtubule organization and the motility of GFP-CDK5RAP2-

positive PCM clusters during nocodazole treatment and dynapyrazole A (treat first) and nocodazole

co-treatment.

(I) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells

stained for microtubules (α-tubulin, red) and PCM (PCNT, green). Enlargements show the merged and

single channels of the boxed areas.

1430

Figure 5. Dynamics of microtubule nucleation and caMTOC re-assembly in acentriolar cells

(A) Immunofluorescence images of microtubule regrowth after nocodazole washout at the indicated

timepoints in centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stained for PCM

components (green) and newly nucleated microtubules (EB1, red). A Golgi marker, GM130 (blue), is

included in the left row, and zooms of the boxed regions (numbered 1 and 2) show that microtubules

nucleate from PCM clusters but not from the Golgi membranes. Dashed lines show cell boundaries.

(B) (Top) Timeline shows the time course of protein depletion (siRNA transfection), nocodazole

treatment, nocodazole washout and microtubule regrowth. (Bottom) Immunofluorescence images of

microtubule regrowth experiments after depletion of the indicated proteins in centrinone-treated

AKAP450/CAMSAP2 knockout RPE1 cells stained for the indicated PCM markers (green) and EB1 as a

marker of nascent microtubules (red). Cell outlines are indicated with dashed lines and enlargements

of the merged channels of the boxed areas are shown on the right.

(C) Quantification of normalized microtubule intensity at 30 s after nocodazole washout in control

cells and cells depleted of the indicated PCM proteins. n=40 (siLuci, siPCNT), 57 (siCDK5RAP2), 48

(siNIN), 45 (siγ-tubulin) and 50 (siDHC) cells analyzed for each measurement in three independent

experiments. The statistical significance was determined by unpaired two-tailed Mann-Whitney test

in Prism 9.1 (**P<0.002; ***P<0.001). Values are represented as mean±SD.

(D) Time-lapse images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stably

expressing GFP-CDK5RAP2 before and after nocodazole washout. Dispersed GFP-CDK5RAP2-positive

PCM clusters (GFP, green) serve as microtubule nucleation sites (SiR-tubulin, red) and coalesce into a

big cluster after nocodazole washout. Time is min:sec.

(E) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells

stained for pericentrin (green) and microtubules (α-tubulin, red) at the indicated timepoints after

nocodazole washout.

(F) Measurements of normalized fluorescence intensity of PCM clusters at the indicated distances in

relation to the brightest point, as described in Methods. The biggest PCM cluster (which normally also

had the highest fluorescence intensity) was selected as the center, around which 10 concentric circles

with 2 μm width were drawn. Fluorescence intensity of PCM clusters in these concentric circles was

measured automatically and normalized by the sum of the total PCM intensity in each cell per

condition. n=12 cells per plot per timepoint. Values represent mean±SD.

(G) Summarizing diagram illustrating microtubule organization and motility of GFP-CDK5RAP2-positive

PCM clusters during nocodazole washout.

1463

Figure 6. caMTOC formation in the presence of CAMSAP2 using inducible motor recruitment

(A) Immunofluorescence images of control or centrinone treated AKAP450 knockout RPE1 cells

stained for CAMSAP2 (green), PCM protein (γ-tubulin, cyan) and microtubules (α-tubulin, red).

Enlargements show the boxed regions of the merged images.

(B) Time lapse images of centrinone treated AKAP450 knockout RPE1 cells stably expressing GFP-

CDK5RAP2 (green). Microtubules were labeled with 100 nM SiR-tubulin overnight (red, top row). The

maximum intensity projection includes 200 frames, 200 ms/frame. Red arrows show the motion

directions of GFP-CDK5RAP2-positive PCM clusters. Time is min:sec.

(C) A diagram of microtubule organization and PCM motility in AKAP450 knockout cells.

(D, E) Diagram of the inducible heterodimerization assay with ppKin14 and CAMSAP2. (D) CAMSAP2
was tagged with mCherry and fused to a tandemly repeated FKBP domain; tetramerized ppKin14 was
tagged with TagBFP and fused to FRB. Rapalog induces the binding of CAMSAP2 and ppKin14 by linking
FKBP to FRB. (E) Rapalog treatment induces the binding of CAMSAP2 (red) and ppKin14 (blue) and the
formation of radial microtubule network. In this scheme, PCM-anchored microtubules are not shown.

(F) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells co-
transfected with 2FKBP-mCherry-CAMSAP2 and FRB-TagBFP-GCN4-ppKin14 and stained for the
indicated proteins in cells treated with DMSO or rapalog. Zooms show the magnifications of boxed
areas.

(G) Quantification of the proportion of cells with a radial, whirlpool-like or non-radial microtubule
organization with and without rapalog treatment. Numbers on the histogram show the percentages.
414 cells treated with DMSO (-Rapa) and 385 cells treated with rapalog (+Rapa) analyzed for each
measurement in three independent experiments (n=3). Values represent mean±SD.

1486

Figure 7. The role of the PCM in CAMSAP2-driven formation of caMTOCs

(A,B) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2/p53/pericentrin
knockout RPE1 cells transfected with 2FKBP-mCherry-CAMSAP2 and FRB-TagBFP-GCN4-ppKi14 and
stained for the indicated components before (top) or after an overnight rapalog treatment. Zooms
show magnifications of boxed areas. Black dashed lines show the position of the nucleus.

(C) Cells treated as described for panel A were co-transfected with GFP-pericentrin and stained for
mitochondria (cytochrome C, red) and CAMSAP2 (red) in same channel overnight after rapalog
addition. Zooms show magnifications of boxed areas.

(D) Quantification of the proportion of cells with different types of microtubule minus end
organization before and after overnight rapalog treatment. Numbers on the histogram show the
percentages. 334 (-Rapa), 424(+Rapa) cells of AKAP450/CAMSAP2/p53/pericentrin knockout RPE1
cells, 206(-Rapa) and 239(+Rapa) of AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/pericentrin knockout
RPE1 cells analyzed for each measurement in three independent experiments (n=3). Values represent
mean±SD.

(E)   Immunofluorescence   images   of   centrinone-treated
AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/pericentrin knockout RPE1 cells transfected with 2FKBP-

mCherry-CAMSAP2 and FRB-TagBFP-GCN4-ppKi14 and stained for microtubules (α-tubulin, green)

after an overnight rapalog treatment. Zooms show magnifications of boxed areas.

1505

Figure 8. Cytosim simulations of PCM and microtubule self-organization.

In all simulations, we considered a circular cell with a radius of 10 μm (gray line), containing

microtubules (red), PCM complexes (green), and CAMSAP-kin14 complexes (blue).

(A) Simulation of 300 PCM complexes; each PCM complex is attached to a dynein motor, and from

each complex, a MT can grow. A weak adhesive interaction is introduced between the PCM complexes.

A compact PCM cluster is observed.

(B) The same simulation as in (A), but with the adhesive interaction disabled; PCM is organized in a

ring-like structure.

(C) The simulation in (A) modified in two ways: a randomly organized, dynamic microtubule network

with 300 filaments is added, and only 50 of the 300 PCM complexes can nucleate microtubules. PCM

cluster formation is disrupted.

(D) The same simulation as in (C), but the strength of the adhesive interaction is increased. The

formation of a compact cluster is recovered.

(E) Cumulative distributions of distances of the PCM complexes to the average position (the center of

mass), determined from the last frame of the simulations.

(F) The mean and standard deviation (SD) of the distances of the PCM complexes to the center of mass

from different repetitions of the simulations. Each system was simulated 10 times. Small values for

the mean and for the standard deviation indicate a compact cluster. For a ring-like structure, the mean

values are is larger and describe the radius of the ring. A dispersed localization of PCM complexes is

characterized by large values for the mean and the standard deviation.

(G) Simulation of 300 CAMSAP-kin14 complexes, which consist of a microtubule nucleation site and

five kinesin-14 motors, but do not adhere to each other. The complexes form a ring-like structure.

(H) The same simulation as in (G), but with complexes that have the same weak adhesive interaction

as for the PCM complexes in (A). The complexes form a central compact cluster.

(I) Simulation of a mixed system with 150 CAMSAP-kin14 complexes that do not adhere to each other

and 150 weakly adhesive PCM complexes. All complexes associate in a central compact cluster.

(J) The same simulation as in (I), but with non-adhesive PCM complexes. A ring-like structure is formed.

(K) Cumulative distributions of the distances from all complexes to the center of mass of the last frame

of the simulation.

(L) The mean and standard deviation of the distances from all complexes to the center of mass from

different repetitions of the simulations. Each system was simulated 10 times. A compact cluster is

characterized by a small value for the mean and for the standard deviation. A ring-like structure has a

larger mean value.

 Legends to Supplemental Figures

 Figure 1 – figure supplement 1. Characterization of caMTOCs in wild type, AKAP450 knockout and

 CAMSAP2 knockout cells

(A) Immunofluorescence images of centrinone B treated AKAP450 knockout RPE1 cells stained for

CAMSAP2 (green) and γ-tubulin (red), transfected either with control (Luciferase) or CAMSAP2 siRNAs.

Insets show enlargements of the merged images of the boxed areas.

(B) Quantification of CAMSAP2 depletion and caMTOC formation related to panel A. For normalized

CAMSAP2 intensity, n=76 (siLuciferase) and 62 (siCAMSAP2) cells from three independent experiment

were analyzed; to calculate the percentage of cells showing dispersed or clustered PCM, n=300

(siLuciferase) and 299 (siCAMSAP2) cells analyzed for each measurement in three independent

experiments. The statistical significance was determined by unpaired two-tailed Mann-Whitney test

in Prism 9.1 (***$p < 0.001$). Values represent mean±SD.

(C-E) Immunofluorescence images of wild type and CAMSAP2 knockout cells treated with DMSO,

CentB, Brefeldin A (BFA) (5 μg/ml (17.8 μM) for 2 hrs), and stained for centrioles (Centrin, red), PCM

(PCNT, green), microtubules (α-tubulin, red) and Golgi apparatus (GM130, blue). Insets show

enlargements of the merged channels of the boxed areas.

(F) Quantification of the proportion of cells with different types of microtubule minus end organization

in wild type (WT) and CAMSAP2 knockout RPE1 cell lines with indicated treatment. 319 (+DMSO),

413(+CentB), 412(+BFA), 353(+CentB+BFA) WT cells and 340(+DMSO), 445(+CentB), 384(+BFA),

553(+CentB+BFA) CAMSAP2 knockout RPE1 cells analyzed for each measurement in three

independent experiments (n=3). Values represent mean±SD.

1560

Figure 1 – figure supplement 2. Characterization of caMTOCs in AKAP450/CAMSAP2 knockout cells

(A) Immunofluorescence images of control (-CentB) and centrinone-treated AKAP450/CAMSAP2

knockout RPE1 cells stained for pericentrin and centrin to show the main PCM organization types

(centrin-negative cells with cylindrical or round PCM cluster, dispersed PCM or no cluster, and centrin-

positive centrosomes) in each condition. Enlargements of the merged channels of the boxed areas are

shown on the right.

(B) Quantification of the main PCM organization types in centrinone-treated AKAP450 knockout and

AKAP450/CAMSAP2 knockout RPE1 cells. Numbers on the histogram show the percentages. 465

(AKAP450 KO) and 495 (AKAP450/CAMSAP2 KO) cells analyzed for each measurement in three

independent experiments (n=3). Values represent mean±SD.

(C) Diagram showing the generation of CPAP or PLK4 depleted AKAP450/CAMSAP2 knockout RPE1

cells with different drug treatments. Cells were transfected with siRNAs to depleted CPAP and PLK4

respectively. After 2 days, the transfection was performed again to increase the depletion efficiency.

After 2 more days, cells were treated with thymidine to block cell proliferation or with a combination

of thymidine and centrinone B, or thymidine, centrinone B and the PLK1 inhibitor BI 2536 for one or

three days. Cells were fixed for the first time (Fix1) after a 24 hr drug treatment, and for the second

time (Fix2) after a 72 hr drug treatment.

(D) Quantification of the main PCM organization types, as described in panel A, for cells prepared as

described in panel E-G. Numbers on the histogram show the percentages and numbers in brackets

show cells analyzed for each measurement in three independent experiments (n=3; 247-612 cells

analyzed per condition). Values represent mean±SD.

(E-G) Immunofluorescence images of Fix2 (as described in panel C) showing control (transfected with

siRNA against Luciferase), or depleted of PLK4 or CPAP and treated as indicated. Cells were stained for

centrioles (CEP135, centrin), PCM proteins (PCNT) and microtubules (α-tubulin). Images illustrate the

PCM organization types in each condition. Insets show the merged channels of the boxed areas.

1586

Figure 1 – figure supplement 3. PCM dynamics visualized with GFP-CDK5RAP2

(A) Immunofluorescence images of control (-CentB) and centrinone B treated WT RPE1 cell stably

expressing GFP-CDK5RAP2 (green) stained for α-tubulin (red) and pericentrin (PCNT, blue). Zooms

show enlargements of the boxed regions.

(B) Time-lapse images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stably

expressing GFP-CDK5RAP2. Microtubules are visualized by treating the cells with 100 nM SiR-tubulin

overnight. PCM labeled with GFP-CDK5RAP2 (green) forms a stable cluster that functions as the MTOC.

Zooms show the enlarged GFP channel of the boxed regions. Time is min:sec.

1595

1596 **Figure 2 – figure supplement 1. Characterization of PCM components localizing to caMTOCs in**

1597 **AKAP450/CAMSAP2 knockout cells**

1598 (A) Western blot showing that endogenous NEDD1 and CEP192 are present in centrinone-treated

1599 AKAP450/CAMSAP2 knockout cells.

(B) Western blot results showing the depletion of the indicated PCM proteins.

(C) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells showing the localization and the effects of depletion of the indicated proteins on the caMTOCs. siRNA against luciferase was used as a control. Zooms show enlargements of the merged channels of the boxed areas.

Figure 2 – figure supplement 2. Generation of the AKAP450/CAMSAP2/p53/pericentrin knockout RPE1 cell line

(A) Diagram illustrating two sgRNA sites targeting p53 exon 2 and exon 4. The green and red boxes indicate the position of the PAM sites, and the predicted Cas9 cut sites are indicated by scissors.

(B,C) Genotyping results of the genomic mutation using gel-purified PCR product which covers exon 2 and exon 4 of the p53-encoding gene in AKAP450/CAMSAP2/p53 knockout cell line and a diagram illustrating the induced inversion.

(D) Diagram illustrating the binding sites of p53 antibodies and the position of the mutation induced in p53-encoding gene.

(E,F) Immunofluorescence and Western blot images showing p53 expression in control and AKAP450/CAMSAP2/p53 knockout cells.

(G-H) Genotyping results of the genomic mutation using gel-purified PCR product which covers the two sgRNAs targeting sites within pericentrin-encoding gene in AKAP450/CAMSAP2/p53/pericentrin knockout cell line and a diagram illustrating the induced genomic mutations within exon 5 of the pericentrin-encoding gene. The green and red boxes indicate the position of the PAM sites, and the predicted Cas9 cut sites are indicated by scissors.

(I) Diagram illustrating the binding domain of the pericentrin antibody and the positions of the mutations induced downstream of the Cas9 cutting sites.

(J) Immunofluorescence images confirming the loss of pericentrin in AKAP450/CAMSAP2/p53/pericentrin knockout cells.

(K) Western blot confirming the loss of p53 and pericentrin in AKAP450/CAMSAP2/p53/pericentrin knockout cells.

Figure 2 – figure supplement 3. Characterization of cell division in centrinone-treated

AKAP450/CAMSAP2/p53 knockout cells

(A) Immunofluorescence images of metaphase AKAP450/CAMSAP2/p53 knockout cells treated with

DMSO and centrinone (CentB) and stained for h microtubules (α-tubulin, magenta), PCM (PCNT, red),

centrioles (Centrin, green) and DNA (DAPI, blue).

(B-C) Phase-contrast time-lapse images of centrinone-treated AKAP450/CAMSAP2/p53 knockout

RPE1 cells. Time is hh:mm. (B) The red and blue arrows show two cells with different mitosis duration.

(C) Mitotic phases in a representative cell.

(D) Quantification of the mitosis duration of centrinone-treated AKAP450/CAMSAP2/p53 knockout

RPE1 cells. 36 cells from 14 videos were analyzed. Values represent mean±SD.

1639

Figure 2 – figure supplement 4. Characterization of PCM organization in AKAP450/CAMSAP2/p53/

pericentrin knockout cells

(A) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2/p53 knockout (Control)

and AKAP450/CAMSAP2/p53/pericentrin knockout (PCNT KO) RPE1 cells showing staining for the

indicated proteins. Zooms of the boxed areas show merged channels.

(B) Western blots showing the expression levels of CDK5RAP2, γ-tub, and ninein in control (+DMSO)

and centrinone-treated (+CentB) AKAP450/CAMSAP2/p53 knockout and

AKAP450/CAMSAP2/p53/PCNT knockout RPE1 cells.

1648

Figure 2 – figure supplement 5. Effects of the depletion or knockout of EB1 and EB3 on caMTOC

formation in AKAP450/CAMSAP2 knockout cells

(A) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout cells stained for

EB1 or EB3 and pericentrin showing effects of EB1 or EB3 depletion on caMTOCs.

(B) Western blot confirming the loss of AKAP450 and CAMSAP2 in AKAP450/CAMSAP2/EB1/EB3

mutant cells.

(C) Immunofluorescence images of control or centrinone-treated AKAP450/CASMSAP2/EB1/EB3

mutant RPE1 cells stained for the indicated proteins. Zooms of the boxed areas show merged

channels.

1658

Figure 4 – Figure supplement 1. PCM dynamics during nocodazole treatment in centrinone-treated AKAP450/CAMSAP2 knockout cells

(A) Time-lapse images illustrating the dynamics of GFP-CDK5RAP2 clusters (green, white arrows) and microtubules (visualized with an overnight treatment with 100 nM SiR-tubulin, red) during nocodazole treatment. A single frame of the whole cell is on the bottom left, with the boxed area indicating the location of the enlarged images. Time is min:sec.

(B) (Top) Motor-PAINT images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells before and after treatment with 10 µM nocodazole for 15 min (the same cells, colorized differently, are displayed in Figure 4G). Microtubule segments were assigned a color based on their absolute orientation, with their plus-ends pointing in the directions indicated with the compass. Asterisks indicate approximate assumed positions of caMTOCs. (Bottom) Directionality-filtered reconstruction of motor tracks used for generating the Motor-PAINT images.

(C) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stained for microtubules (α-tubulin, red) and PCM (PCNT, green) after cold treatment for 30 min at 4°C. Enlargements show the merged and single channels of the boxed areas.

1674

Figure 5 – figure supplement 1. PCM dynamics and microtubule regrowth during nocodazole washout in AKAP450/CAMSAP2 knockout cells

(A) Immunofluorescence images of control (+DMSO) and centrinone-treated AKAP450/CAMSAP2 knockout cells treated with cold alone or cold and nocodazole, and stained for nascent microtubules (EB1, red), PCM (PCNT, green) and the Golgi apparatus (GM130, blue) 30 s after nocodazole washout.

(B) Immunofluorescence images of acentriolar AKAP450/CAMSAP2 knockout RPE1 cells fixed at 30 s and 10 min after nocodazole washout and stained for the indicated markers. Zooms show the enlargements of boxed areas. Dashed lines show the cell boundaries.

(C) Summarizing table showing the presence of PCM components in clusters (including centrosome and caMTOC) before and after nocodazole washout in indicated conditions.

(D) Immunofluorescence images of control acentriolar AKAP450/CAMSAP2 knockout RPE1 cells and cells lacking the indicated PCM proteins fixed 2 min after nocodazole washout and stained for EB1 (a marker of growing microtubules, red) and different PCM proteins (γ-tubulin, ninein (NIN) and dynein heavy chain (DHC), green; CDK5RAP2 and PCNT, blue). Insets show the enlargements of the boxed

areas and the heat maps on the right show the radial and non-radial part of growing microtubules.

Dashed lines show the cell boundaries.

(E) Quantification of the proportion of non-radial microtubules 2 min after nocodazole washout. The

radial and the non-radial microtubule intensities were measured and the non-radial microtubule

intensity was normalized by dividing it by the sum of the radial and the non-radial microtubule

intensities for each group. n=20-25 cells analyzed for each measurement in three independent

experiments. The statistical significance was determined by unpaired two-tailed Mann-Whitney test

in Prism 9.1 (***p<0.001). Values represent mean±SD.

(F) Diagram showing microtubule regrowth and PCM assembly in control acentriolar

AKAP450/CAMSAP2 knockout RPE1 cells and cells depleted of the indicated PCM proteins at 30 s and

2 min after nocodazole washout.

1700

Figure 5 – figure supplement 2. Pericentrin is required for PCM clustering in acentriolar

AKAP450/CAMSAP2 knockout RPE1 cells

Time-lapse images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells stably expressing

GFP-CDK5RAP2 (green), which is diffusely distributed and does not form any clear clusters.

Microtubule regrowth experiments show random nucleation of microtubules (labeled with 100nM

SiR-tubulin overnight) that remain disorganized. Summarizing diagram shows PCM clusters and

microtubule network in the indicated conditions.

1708

Figure 6 – figure supplement 1. PCM dynamics and the effects of CAMSAP2 clustering on PCM

organization in centrinone-treated RPE1 cells lacking AKAP450

(A) Time-lapse images illustrating the dynamics of PCM clusters (green, white arrows) and

microtubules (SiR-tubulin, red) after nocodazole washout in centrinone-treated AKAP450 knockout

RPE1 cells stably expressing GFP-CDK5RAP2. A single frame of the whole cell with numbered boxed

areas indicating the location of enlarged time-lapse images is shown on the left. White arrows indicate

moving PCM clusters after microtubules re-grow. Time is min:sec.

(B) Time-lapse images of centrinone-treated AKAP450 knockout RPE1 cells stably expressing GFP-

CDK5RAP2. GFP-CDK5RAP2 forms small clusters that are dynamic before microtubules (SiR-tubulin,

red) are disassembled by nocodazole treatment. Red arrows show that these clusters become

immobile when microtubules are completely depolymerized (enlargements of the boxed areas shown

in the bottom rows). GFP-CDK5RAP2-positive clusters are sites of microtubule nucleation after nocodazole washout and become dynamic again once microtubules re-grow. The maximum intensity projections include 100 or 400 frames, 200 ms/frame. Time is min:sec.

(C) Summarizing diagram illustrating microtubule organization and the motility of PCM clusters in acentriolar AKAP450 knockout RPE1 cells.

(D) Movement trajectories of PCM clusters in acentriolar AKAP450 knockout cells before and after 1 hr treatment with 5µM dynapyrazole A (Dyna). 350 frames with a 2 s interval from the same cell were analyzed. Shaded areas represent the cell body.

(E) Quantification of the length of the trajectories of moving PCM clusters. n=132 (without dynapyrazole A, from 5 cells) and 111 (after 1 hr dynapyrazole A treatment, from 4 cells) trajectories were analyzed for each condition. The statistical significance was determined by unpaired two-tailed Mann-Whitney test in Prism 9.1 (***$p < 0.001$). Values represent mean±SD.

(F) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2 knockout RPE1 cells transfected with 2FKBP-mCherry-CAMSAP2 and FRB-TagBFP-GCN4-ppKin14 and stained for the indicated proteins after overnight rapalog treatment. Zooms show the magnification of boxed area.

Figure 6 – figure supplement 2. Pericentrin is required for PCM clustering in acentriolar AKAP450 knockout RPE1 cells

Time-lapse images of centrinone-treated AKAP450 knockout RPE1 cells stably expressing GFP-CDK5RAP2 (green) depleted of pericentrin. GFP-CDK5RAP2 is diffusely distributed and does not form any clear clusters. Microtubule regrowth experiments show random nucleation of microtubules (labeled with 100nM SiR-tubulin overnight) that remain disorganized. Summarizing diagram shows PCM clusters and microtubule network in the indicated conditions.

Figure 7 – figure supplement 1. Inducible CAMSAP2-driven microtubule rearrangement in AKAP450/CAMSAP2/p53/pericentrin knockout cell

Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2/p53/pericentrin knockout cells expressing either FKBP-mCh-CAMSAP2 alone, FRB-GFP-ppKin14 alone or both, with or without rapalog treatment, and stained for microtubules (α-tubulin). Enlargements show the merged channels of the boxed areas.

1750

Figure 7 – figure supplement 2. Inducible CAMSAP2-driven radial microtubule rearrangement in
AKAP450/CAMSAP2/p53/pericentrin knockout cell

(A) Time-lapse images of a centrinone-treated AKAP450/CAMSAP2/p53/pericentrin knockout cell
transiently expressing 2FKBP-mCherry-CAMSAP2 and FRB-GFP-GCN4-ppKin14 imaged for 10 min (100
frames, 6 s interval) prior to treatment with 100 nM rapalog and ~90 min after. Time is hr: min: sec.

(B) Immunofluorescence images of centrinone-treated AKAP450/CAMSAP2/p53/pericentrin knockout
and AKAP450/CAMSAP2/MMG/CDK5RAP2/p53/pericentrin knockout RPE1 cells, stained for the
indicated markers. In the upper row, the cells were neither transfected nor treated with rapalog,
whereas in the other panels, cells were transfected with 2FKBP-mCherry-CAMSAP2 and FRB-HA-
GCN4-ppKin14 and treated with rapalog overnight. Zooms show the magnification of boxed areas.

1761

Figure 7 – figure supplement 3. Generation of the AKAP450/CAMSAP2/CDK5RAP2/MMG/
p53/pericentrin knockout RPE1 cell line

(A) Diagram illustrating two sgRNA sites targeting p53 exon 4. The green and red boxes indicate the
position of the PAM sites, and the predicted Cas9 cut sites are indicated by scissors. Genotyping results
of the genomic mutation using gel-purified PCR product which covers exon 4 of the p53-encoding gene
in AKAP450/CAMSAP2/MMG/CDK5RAP2/p53 knockout cell line are shown below.

(B) Diagram illustrating the binding sites of p53 antibodies and the position of the mutation induced
in p53-encoding gene.

(C) Immunofluorescence images showing p53 staining in control and
AKAP450/CAMSAP2/MMG/CDK5RAP2/p53 knockout cells.

(D) Diagram illustrating two sgRNA sites targeting pericentrin exon 5. The green and red boxes indicate
the position of the PAM sites, and the predicted Cas9 cut sites are indicated by scissors. Genotyping
results of the genomic mutation using gel-purified PCR product which covers exon 5 of the pericentrin-
encoding gene in AKAP450/CAMSAP2/MMG/CDK5RAP2/p53/pericentrin knockout cell line are shown
below.

(E) Diagram illustrating the binding domain of the pericentrin antibody and the positions of the
mutations induced downstream of the Cas9 cut sites.

(F) Immunofluorescence images showing the confirmation of pericentrin loss in control and
AKAP450/CAMSAP2/MMG/CDK5RAP2/p53/pericentrin knockout cells.

(G) Western blot illustrating the p53 and pericentrin loss in control and
AKAP450/CAMSAP2/MMG/CDK5RAP2/p53/pericentrin knockout cells.

(H) Immunofluorescence images of control and centrinone-treated AKAP450/CAMSAP2/
CDK5RAP2/MMG/p53/pericentrin knockout cells staining microtubules (α-tubulin, red) and
centrosomes (CEP192, green). Zooms show magnification of the boxed area.

1786

 Legends to Supplemental Videos

Video 1. Cell cycle progression in acentriolar AKAP450/CAMSAP2/p53 knockout cells

Cell cycle progression visualized by phase-contrast imaging of centrinone-treated
AKAP450/CAMSAP2/p53 knockout RPE1 cells. The cells were imaged for ~15 hrs with 1min interval.
Time is hh:mm.

Video 2. caMTOC disassembly during nocodazole treatment of acentriolar AKAP450/CAMSAP2
knockout cells

PCM dynamics visualized by stable expression of GFP-CDK5RAP2 (green) in centrinone-treated
AKAP450/CAMSAP2 knockout RPE1 cells. Microtubules were labeled overnight with 100 nM SiR-
tubulin (red). The cell was imaged for ~3.5min (100 frames, 2 s interval) prior to the addition of 10 µM
nocodazole. Time is min: sec.

Video 3. Depletion of pericentrin inhibits PCM clustering in acentriolar AKAP450/CAMSAP2
knockout cells

A pericentrin-depleted acentriolar AKAP450/CAMSAP2 knockout cell stably expressing GFP-CDK5RAP2
(green) and labeled overnight with 100 nM SiR-tubulin (red) was imaged for ~4.5 min (140 frames, 2 s
time interval) prior to treatment with 10 µM nocodazole. Nocodazole was washed out at ~20 min
(frame 591), when all microtubules were depolymerized. Time is hr: min: sec.

Video 4. CAMSAP2 distribution and dynamics in acentriolar AKAP450 knockout cells

An acentriolar AKAP450 knockout RPE1 cell transiently expressing mCherry-CAMSAP2 was imaged for
15 hrs (900 frames, 1min interval). Time is hr:mm.

Video 5. PCM dynamics during nocodazole treatment and washout in acentriolar AKAP450 knockout
cells

An acentriolar AKAP450 knockout RPE1 cell stably expressing GFP-CDK5RAP2 (green) and labeled
overnight with 100 nM SiR-tubulin (red) was imaged for ~7min (200 frames, 2 s interval) prior to
treatment with 10 µM nocodazole. Nocodazole was washed out at ~27 min (frame 801) when all
microtubules were depolymerized. Time is min: sec.

Video 6. PCM dynamics in acentriolar AKAP450 knockout cells are inhibited by dynapyrazole

An acentriolar AKAP450 knockout RPE1 cell stably expressing GFP-CDK5RAP2 (green) and labeled
overnight with 100 nM SiR-tubulin (red) was imaged for ~12 min (350 frames, 2 s interval) prior to the

dynapyrazole treatment, treated with 5 µM Dynapyrazole A for 1h, and then the same cell was imaged

for ~12 min. Time is min: sec.

Video 7. Depletion of pericentrin inhibits PCM clustering in acentriolar AKAP450 knockout cells

A pericentrin-depleted acentriolar AKAP450 knockout cell stably expressing GFP-CDK5RAP2 (green)

and labeled overnight with 100 nM SiR-tubulin (red) was imaged for ~3 min (90 frames, 2 s time

interval) prior to treatment with 10 µM nocodazole. Nocodazole was washed out at ~23 min (frame

701) when all microtubules were depolymerized. Time is hr: min: sec.

Video 8. Inducible CAMSAP2-driven radial microtubule rearrangement in an

AKAP450/CAMSAP2/p53/pericentrin knockout cell

An acentriolar AKAP450/CAMSAP2/p53/pericentrin knockout cell transiently expressing 2FKBP-

mCherry-CAMSAP2 (red) and FRB-GFP-GCN4-ppKin14 (green) was imaged for 10 min (100 frames, 6 s

interval) prior to treatment with 100 nM rapalog. Subsequently, the cell was imaged for ~1 hr and 35

min after rapalog addition. Time is hr: min: sec.

Video 9. Cytosim simulations of PCM and microtubule self-organization

Example videos of our Cytosim simulations. Each simulation represents 1800 seconds of real time, and

the last frame is shown in the panels of Figure 8. In all simulations, we considered a circular cell with

a radius of 10 µm (gray line), containing microtubules (red), PCM complexes (green), and CAMSAP-

kin14 complexes (blue). The videos correspond to panels A,B,C, D, I and J of Figure 8, as indicated.

1834

1835

Source data 1

Uncropped Western blots shown in this manuscript

(A,B) Western blots showing that NEDD1 and CEP192 are present in centrinone-treated

AKAP450/CAMSAP2 knockout cells shown in Figure 2 – figure supplement 1A.

(C-J) Western blots showing the depletion of indicated proteins in centrinone-treated

AKAP450/CAMSAP2 knockout cells shown in Figure 2 – figure supplement 1B.

White dashed lines indicate where the blots were cut before incubation.

(K) Western blot showing that 3 hrs treatment with dynaprazole A does not affect the expression of

the endogenous dynein heavy chain and the dynactin large subunit p150Glued in centrinone-treated

AKAP450/CAMSAP2 knockout cells shown in Figure 4D.

1846

Source data 2

Uncropped Western blots shown in this manuscript

(A) Western blots showing the knockout of p53 from AKAP450/CAMSAP2 knockout cell line shown in

Figure 2 – figure supplement 2F.

(B) Western blots showing the knockout of pericentrin from AKAP450/CAMSAP2/p53 knockout cell

line shown in Figure 2 – figure supplement 2K.

(C-D) Western blots showing expression levels of CDK5RAP2, γ-tub, and ninein (NIN) in control (-CentB)

and centrinone-treated AKAP450/CAMSAP2/p53 knockout and AKAP450/CAMSAP2/p53/PCNT

knockout cell lines shown in Figure 2 – figure supplement 4B.

(E) Western blots showing the knockout of AKAP450 and CAMSAP2 from EB1/EB3 mutant RPE1 cell

line shown in Figure 2 – figure supplement 5B.

(F) Western blots showing the knockout of pericentrin in AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/

knockout cell line shown in Figure 7 – figure supplement 3G.

White dashed lines indicate where the blots were cut before incubation.

1861

1862 **Supplemental file 1. Cytosim configuration file for the simulations.**

1863 The configuration file was executed with a compiled Cytosim (www.cytosim.org) version June 2019**.**

 Supplemental Table S1: Parameters with numerical values used in the Cytosim simulations.

	Symbol	Value	Note	Reference
Cell				
Cell radius		10 µm		
Viscosity		1 pN s/ µm^2	Typical value for Cytosim simulations	(Kole et al., 2005; Letort et al., 2016)
Thermal energy	kT	4.2 pN nm		
Dynein				
Binding rate	k_{on}	5 s^{-1}	Typical order of magnitude	(Ohashi et al., 2019)
Force-free unbinding rate	k^0_{off}	0.1 s^{-1}		(Ohashi et al., 2019)
Detachment force	F_d	2		(Ohashi et al., 2019)
Binding range		75 nm	Estimated	
Force-free velocity	v_0	500 nm/s		(Brenner et al., 2020)
Stall force	F_s	4 pN		(Belyy et al., 2016)
Kin14				
Binding rate	k_{on}	1 s^{-1}	Typical order of magnitude for molecular motors	(Klumpp et al., 2015)
Force-free unbinding rate	k^0_{off}	1 s^{-1}	Assumption based on kinesin-1 value	(Berger et al., 2019)
Detachment force	F_d	3	Assumption based on kinesin-1 value	(Pyrpassopoulos et al., 2020)
Binding range		75 nm	Estimated	
Force-free velocity	v_0	800 nm/s	Assumption based on kinesin-1 value	(Carter and Cross, 2005)
Stall force	F_s	7 pN	Assumption based on kinesin-1 value	(Carter and Cross, 2005)
Microtubules				
Rigidity		20 pN µm^2		(Gibeaux et al., 2017)
Catastrophe rate	k_{cat}	0.026 s^{-1}	A similar value has been used in the reference	(Gibeaux et al., 2017)
Rescue rate		0	We ignore rescues	
Growing force		1.7 pN		(Dogterom and Yurke, 1997)

Growing speed	v_g	0.13 µm/s		(Burakov et al., 2003; Letort et al., 2016)
Shrinkage speed		0.272 µm/s		(Burakov et al., 2003; Letort et al., 2016)
Maximum length		7.5 µm	To avoid boundary effects	
Adhesive interactions of PCM complexes				
Binding rate		10 s^{-1}	Estimated	
Binding range		100 nm	Estimated	
Unbinding rate		0.01 s^{-1}	Estimated	
Detachment force		3 pN	Typical value, estimated	
General parameters				
Stiffness of all linking elements		100 pN/ µm	Typical value for molecular motors	(Gros et al., 2021; Letort et al., 2016)

1865

1866

 Key reagent or resource table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	anti-Pericentrin (mouse monoclonal)	Abcam	Abcam Cat# ab28144, RRID:AB_2160664	(1:500) for IF
Antibody	anti-Pericentrin (rabbit polyclonal)	Abcam	Abcam Cat# ab4448, RRID:AB_304461	(1:500) for IF; (1:1000) for WB
Antibody	anti-CDK5RAP2 (rabbit polyclonal)	Bethyl Laboratories	Bethyl Cat# A300-554A, RRID:AB_477974	(1:300) for IF; (1:1000) for WB
Antibody	anti-γ-tubulin (mouse monoclonal)	Sigma-Aldrich	Sigma-Aldrich: T6557; RRID:AB_477584	(1:300) for IF; (1:2000) for WB
Antibody	anti-γ-tubulin (rabbit polyclonal)	Sigma-Aldrich	Sigma-Aldrich:T3559, RRID:AB_477575	(1:300) for IF
Antibody	anti-NEDD1 (mouse monoclonal)	Abnova	Abnova Corporation Cat# H00121441-M05, RRID:AB_534956	(1:300) for IF
Antibody	anti-NEDD1 (rabbit polyclonal)	Rockland	Rockland Cat# 109-401-C38S, RRID:AB_10893219	(1:1000) for WB
Antibody	anti-Ninein (rabbit polyclonal)	BETHYL	Bethyl Cat# A301-504A, RRID:AB_999627	(1:300) for IF; (1:2000) for WB
Antibody	anti-Ninein (mouse monoclonal)	Santa Cruz Biotechnology	Santa Cruz Biotechnology Cat# sc-376420, RRID:AB_11151570	(1:300) for IF
Antibody	anti-Dynein HC (rabbit polyclonal)	Santa Cruz Biotechnology	Santa Cruz Biotechnology Cat# sc-9115, RRID:AB_2093483	(1:300) for IF; (1:500) for WB
Antibody	anti-p150Glued (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 610473, RRID:AB_397845	(1:100) for IF; (1:500) for WB
Antibody	anti-PCM1 (mouse monoclonal)	Santa Cruz Biotechnology	Santa Cruz Biotechnology Cat# sc-398365, RRID:AB_2827155	(1:300) for IF
Antibody	anti-PCM1 (rabbit polyclonal)	Bethyl Laboratories	Bethyl Cat# A301-150A, RRID:AB_873100	(1:300) for IF
Antibody	anti-AKAP450 (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 611518, RRID:AB_398978	(1:500) for WB
Antibody	anti-CAMSAP2 (rabbit polyclonal)	Novus	Novus:NBP1-21402; RRID:AB_1659977	(1:200) for IF; (1:1000) for WB
Antibody	anti-p53 (mouse monoclonal)	Santa Cruz Biotechnology	Santa Cruz Biotechnology Cat# sc-126, RRID:AB_628082	(1:300) for IF; (1:1000) for WB
Antibody	anti-p53 (rabbit polyclonal)	BETHYL	Bethyl Cat# A300-248A, RRID:AB_263349	(1:300) for IF
Antibody	anti-EB1 (mouse monoclonal)	BD Biosciences	BD Biosciences:610535; RRID:AB_397892	(1:400) for IF

Antibody	anti-EB3 (rabbit polyclonal)	Martin, et al., 2018;		(1:300) for IF
Antibody	anti-Centrin (mouse monoclonal)	Millipore	Millipore Cat# 04-1624, RRID:AB_10563501	(1:500) for IF
Antibody	anti-CEP120 (rabbit polyclonal)	Thermo Fisher Scientific	Thermo Fisher Scientific Cat# PA5-55985, RRID:AB_2639665	(1:300) for IF
Antibody	anti-CEP135 (rabbit polyclonal)	Sigma-Aldrich	Sigma-Aldrich:SAB4503685; RRID:AB_10746232	(1:300) for IF
Antibody	anti-CEP152 (rabbit polyclonal)	Abcam	Abcam, Cat # ab183911	(1:300) for IF; (1:1000) for WB
Antibody	anti-CEP170 (mouse monoclonal)	Thermo Fisher Scientific	Thermo Fisher Scientific Cat# 41-3200, RRID:AB_2533502	(1:200) for IF
Antibody	anti-CEP192 (rabbit polyclonal)	Bethyl Laboratories	Bethyl Cat# A302-324A, RRID:AB_1850234	(1:300) for IF; (1:1000) for WB
Antibody	anti-GM130 (mouse monoclonal)	BD Biosciences	BD Biosciences:610823; RRID:AB_398142	(1:300) for IF; (1:2000) for WB
Antibody	anti-α-tubulin YL1/2 (rat monoclonal)	Pierce	Pierce: MA1-80017; RRID:AB_2210201	(1:300) for IF
Antibody	anti-α-tubulin (mouse monoclonal)	Sigma-Aldrich	Sigma-Aldrich:T5168; RRID:AB_477579	(1:400) for IF
Antibody	anti-α-tubulin (rabbit monoclonal antibody)	Abcam	Abcam Cat# ab52866, RRID:AB_869989	(1:800) for IF
Antibody	Anti-β-tubulin (mouse monoclonal)	Sigma-Aldrich	Sigma-Aldrich Cat# T8660, RRID:AB_477590	(1:2000) for WB
Antibody	anti-CLASP1 (rabbit polyclonal)	(Akhmanova et al., 2001)		(1:400) for IF
Antibody	anti-CLASP2 (rabbit polyclonal)	(Akhmanova et al., 2001)		(1:400) for IF
Antibody	anti-CLIP-115 #2238 (rabbit polyclonal)	(Akhmanova et al., 2001)		(1:300) for IF
Antibody	anti-CLIP-170 #2360 (rabbit polyclonal)	(Akhmanova et al., 2001)		(1:300) for IF
Antibody	anti-ch-TOG (rabbit polyclonal)	(Charrasse et al., 1998)	Dr. Lynne Cassimeris (Lehigh University, USA)	(1:200) for IF
Antibody	anti-CPAP (rabbit polyclonal)	(Kohlmaier et al., 2009)	Dr. Pierre Gönczy (EPFL, Switzerland)	(1:200) for IF
Antibody	anti-CP110 (rabbit monoclonal)	Proteintech	Proteintech Cat# 12780-1-AP, RRID:AB_10638480	(1:300) for IF
Antibody	anti-KIF1C (rabbit polyclonal)	Cytoskeleton	Cytoskeleton Cat# AKIN11-A, RRID:AB_10708792	(1:300) for IF

Antibody	anti-KIF2A (rabbit polyclonal)	(Ganem and Compton, 2004)	Dr. Duane Compton (Geisel School of Medicine at Dartmouth, USA)	(1:300) for IF
Antibody	anti-HAUS2 (rabbit polyclonal)	(Lawo et al., 2009)	Dr. Laurence Pelletier (Lunenfeld-Tanenbaum Research Institute, Canada)	(1:200) for IF
Antibody	anti-BICD2 (rabbit polyclonal)	(Hoogenraad et al., 2003)		(1:2500) for WB
Antibody	anti-Actin (mouse monoclonal)	Millipore	Millipore Cat# MAB1501, RRID:AB_2223041	(1:4000) for WB
Antibody	anti-Ku80 (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 611360, RRID:AB_398882	(1:2000) for WB
Antibody	anti-LaminA/C (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 612162, RRID:AB_399533	(1:400) for IF
Antibody	anti-Cytochrome C (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 556432, RRID:AB_396416	(1:300) for IF
Antibody	anti-Calnexin (rabbit polyclonal)	Abcam	Abcam Cat# ab22595, RRID:AB_2069006	(1:300) for IF
Antibody	Anti-Lamtor4 (rabbit monoclonal)	Cell Signaling (CST)/Bioke	Cell Signaling Technology Cat# 12284, RRID:AB_2797870	(1:800) for IF
	Anti-Tom20 (mouse monoclonal)	BD Biosciences	BD Biosciences Cat# 612278, RRID:AB_399595	(1:200) for IF
Antibody	IRDye 800CW/680LT secondaries	Li-Cor Biosciences	LI-COR Biosciences Cat# 926-32219, RRID:AB_1850025' LI-COR Biosciences Cat# 926-68020, RRID:AB_10706161; LI-COR Biosciences Cat# 926-32211, RRID:AB_621843; LI-COR Biosciences Cat# 926-68021, RRID:AB_10706309	(1:5000) for WB
Antibody	Alexa Fluor 405–, 488–, and 594–secondaries	Molecular Probes/ Thermo Fisher Scientific	Molecular Probes Cat# A-11007, RRID:AB_141374; Cat# A-11034, RRID:AB_2576217; Cat# A32723, RRID:AB_2633275; Cat# A-31553, RRID:AB_221604; Cat# A-11029, RRID:AB_138404; Cat# A-11032, RRID:AB_2534091; Cat# A-11006, RRID:AB_141373; Thermo Fisher Scientific Cat# A-11012, RRID:AB_2534079	(1:500) for IF
Sequence-based reagent	siRNA against PCNT #1	(Gavilan et al., 2018)	5'-AAAAGCUCUGAUUUAUCAAAAGAAG-3'	

Sequence-based reagent	siRNA against PCNT #2	(Gavilan et al., 2018)	5'-UGAUUGGACGUCAUCCAAUGAGAAA-3'	
Sequence-based reagent	siRNA against PCNT #3	(Tibelius et al., 2009)	5'-GCAGCUGAGCUGAAGGAGA-3'	
Sequence-based reagent	siRNA against CDK5RAP2	(Fong et al., 2008)	5'-UGGAAGAUCUCCUAACUAA-3'	
Sequence-based reagent	siRNA against γ-tubulin #1	(Luders et al., 2006)	5'-GGAGGACAUGUUCAAGGAA-3'	
Sequence-based reagent	siRNA against γ-tubulin #2	(Vinopal et al., 2012)	5'-CGCAUCUCUUUCUCAUAU-3'	
Sequence-based reagent	siRNA against Ninein	(Goldspink et al., 2017)	5'-CGGUACAAUGAGUGUAGAAUU-3'	
Sequence-based reagent	siRNA against PCM1	(Wang et al., 2013)	5'-UCAGCUUCGUGAUUCUCAG-3'	
Sequence-based reagent	siRNA against CEP152	(Cizmecioglu et al., 2010; Komarova et al., 2005)	5'-GCGGAUCCAACUGGAAAUCUA-3'	
Sequence-based reagent	siRNA against CEP120	(Ganem et al., 2005; Lin et al., 2013)	5'-AAUAUAUCUUCUUGCAUCUCCUUCC-3'	
Sequence-based reagent	siRNA against CEP192	(Sonnen et al., 2013)	5'-CAGAGGAAUCAAUAAUAAA-3'	
Sequence-based reagent	siRNA against NEDD1 #1	(Luders et al., 2006)	5'-GCAGACAUGUGUCAAUUUA-3'	
Sequence-based reagent	siRNA against NEDD1 #2	(Haren et al., 2006)	5'-GGGCAAAAGCAGACAUGUG-3'	
Sequence-based reagent	siRNA against DHC #1	(Splinter et al., 2010)	5'-CGUACUCCCGUGAUUGAUG-3'	
Sequence-based reagent	siRNA against DHC #2	(Splinter et al., 2010)	5'-GCCAAAAGUUACAGACUUU-3'	
Sequence-based reagent	siRNA against CAMSAP2	(Jiang et al., 2014)	5'-GUACUGGAUAAAUAAGGUA-3'	
Sequence-based reagent	siRNA against CEP170	(Stolz et al., 2015)	5'-GAAGGAAUCCUCCAAGUCA-3'	
Sequence-based reagent	siRNA against CPAP	(Tang et al., 2009)	5'-AGAAUUAGCUCGAAUAGAA-3'	
Sequence-based reagent	siRNA against CLIP170 #1	(Lansbergen et al., 2004; Mimori-Kiyosue et al., 2005)	5'-GGAGAAGCAGCAGCACAUU-3'	

Sequence-based reagent	siRNA against CLIP170 #2	(Lansbergen et al., 2004; Mimori-Kiyosue et al., 2005)	5'-UGAAGAUGUCAGGAGAUAA-3'	
Sequence-based reagent	siRNA against CLIP115 #1	(Lansbergen et al., 2004)	5'-GGCACAGCAUGAGCAGUAU-3'	
Sequence-based reagent	siRNA against CLIP115 #2	(Lansbergen et al., 2004)	5'-CUGGAAAUCCAAGCUGGAC-3'	
Sequence-based reagent	siRNA against ch-TOG:	(Cassimeris and Morabito, 2004; Lansbergen et al., 2004)	5'-GAGCCCAGAGUGGUCCAAA-3'	
Sequence-based reagent	siRNA against EB1	(Grigoriev et al., 2008; Lansbergen et al., 2004)	5'-AUUCCAAGCUAAGCUAGAA-3'	
Sequence-based reagent	siRNA against EB3	(Cassimeris and Morabito, 2004; Komarova et al., 2005)	5'-CUAUGAUGGAAAGGAUUAC-3'	
Sequence-based reagent	siRNA against KIF2A	(Ganem et al., 2005; Grigoriev et al., 2008)	5'-GGCAAAGAGAUUGACCUGG-3'	
Sequence-based reagent	siRNA against CP110	(Cizmecioglu et al., 2010; Spektor et al., 2007)	5'-AAGCAGCAUGAGUAUGCCAGU-3'	
Sequence-based reagent	siRNA against Luciferase	(Lansbergen et al., 2004; Lin et al., 2013)	5'-CGUACGCGGAAUACUUCGA-3'	
Sequence-based reagent	sgRNA target CAMSAP2	(Lansbergen et al., 2004; Wu et al., 2016)	5'-gCATGATCGATACCCTCATGA-3	
Sequence-based reagent	sgRNA target p53 e2 #1	This study	5'-gCGTCGAGCCCCCTCTGAGTC-3';	
Sequence-based reagent	sgRNA target p53 e4 #2	This study	5'-gCCATTGTTCAATATCGTCCG-3';	
Sequence-based reagent	sgRNA target PCNT e5-1 #1	This study	5'-gAGACGGCATTGACGGAGCTG-3';	

Sequence-based reagent	sgRNA target PCNT e5-2 #2	This study	5′-GCTCAACAGCCGGCGTGCCC-3′;	
Sequence-based reagent	p53 KO sequencing primer F	This study	5′-TCAGACACTGGCATGGTGTT-3′;	
Sequence-based reagent	p53 KO sequencing primer R	This study	5′-AGAAATGCAGGGGGATACGG-3′;	
Sequence-based reagent	PCNT KO sequencing primer F	This study	5′-ATACAGCGAGGGAATTCGGG-3′;	
Sequence-based reagent	PCNT KO sequencing primer R	This study	5′-TAGAATGCCCACACCGAGC-3′;	
Chemical compound	Centrinone B	Tocris Bioscience	Tocris Bioscience Cat # 5690	125 nM
Chemical compound	Nocodazole	Sigma-Aldrich	Sigma-Aldrich, Cat # M1404-10MG	10 µM
Chemical compound	Rapalog (A/C Heterodimerizer)	Takara	Takara, Cat # 635056	50 nM (fixation), 100 nM (live imaging).
Chemical compound	Dynapyrazole A	Sigma-Aldrich	Sigma-Aldrich, Cat # SML2127-25MG	5 µM
Chemical compound	BI2536	Selleckchem	Selleckchem, Cat # S1109	500 nM
Chemical compound	Thymidine	Sigma-Aldrich	Sigma-Aldrich, Cat # T9250-25G	5 mM
Chemical compound	proTAME	Boston Biochem	Boston Biochem, Cat # I-440	5 µM
Chemical compound	Brefeldin A	Peptrotech	Peptrotech, Cat # 2031560	5 µg/ml
Chemical compound	SiR-tubulin	Tebu-bio	Tebu-bio, Cat # SC002	100 nM
Software, algorithm	ImageJ radiality plugin	https://github.com/ekatrukha/radialitymap	Katrukha, 2017. radialitymap. Github. https://github.com/ekatrukha/radialitymap cf1e78f	
Recombinant DNA reagent	pLVX-IRES-puro (plasmid)	Clontech		
Recombinant DNA reagent	pB80-FRB-TagBFP-GCN4-ppKin14 (plasmid)	This work		

Recombinant DNA reagent	pB80-FRB-GFP-GCN4-ppKin14 (plasmid)	This work		
Recombinant DNA reagent	pB80-FRB-HA-GCN4-ppKin14 (plasmid)	This work		
Recombinant DNA reagent	2FKBP-mCherry-CAMSAP2 (plasmid)	This work		
Recombinant DNA reagent	GFP-PCNT (plasmid)	This work		
Recombinant DNA reagent	GST-DmKHC(1-421)-mNeonGreen (plasmid)	This work		
Cell line (Homo sapiens)	hTERT-RPE-1	ATCC	CRL-4000	
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450 knockout	(Wu et al., 2016)		
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450/CAMSAP2 knockout	(Wu et al., 2016)		
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450/CAMSAP2/p53 knockout	This work		
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450/CAMSAP2/p53/Pericentrin knockout	This work		
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450/CAMSAP2/EB1/EB3 mutant	This work		
Cell line (Homo sapiens)	hTERT-RPE-1 AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/Pericentrin knockout	This work		

Cell line (Homo sapiens)	HEK 293T	ATCC	CRL-11268	

1868

Figure 2

Figure 3

Figure 4

A Noco only

| No drug | +Noco 1h | Noco washout 15min |
Fix 1 (red) | Fix 2 (green) | Fix 3 (blue)

Noco+Dyna (Noco first)

| No drug | +Noco 1h | +Noco / +Dyna 1h | Noco washout / +Dyna 15min |
Fix 1 | Fix 2 | Fix 3

Noco+Dyna (Dyna first)

| +Dyna 1h | +Noco / +Dyna 1h | Noco washout / +Dyna 15min |
Fix 1 | Fix 2 | Fix 3

B AKAP450/CAMSAP2 KO + CentB

Noco only: PCNT Fix 1, PCNT Fix 2, PCNT Fix 3
Noco first: Fix 1, Fix 2, Fix 3
Dyna first: Fix 1, Fix 2, Fix 3

10µm

C Area occupied by PCM clusters

Diameter (µm)

Noco only, Noco first, Dyna first

Fix 1, Fix 2, Fix 3

* *** *** *** ***

D

kDa   -Dyna +Dyna
530- DHC
142-

E AKAP450/CAMSAP2 KO stable GFP-CDK5RAP2 + CentB

Noco treatment

GFP CDK5RAP2   SiR-tubulin

05:00 mm:ss, 11:00, 11:16, 15:10, 16:50, 22:30, 30:28, 36:26   5µm

GFP   5µm

F

5min
5µm

Norm. MT inten. (%) 2 µm around PCM clusters (red)
PCM cluster velocity (µm/s) (green)

PCM clusters transport time (s)

G AKAP450/CAMSAP2 KO + CentB

-Noco   +Noco (10µM, 15min)

10µm   10µm

Magenta MT= minus-end-out MT
White MT= plus-end-out MT

H AKAP450/CAMSAP2 KO

+CentB

+Noco → Early stage → Late stage

+Dyna(treat first) +Noco

Centrosome
MT
Nucleus
Compact PCM cluster
Small PCM clusters
Mobile PCM
Immobile PCM

I AKAP450/CAMSAP2 KO + CentB

Noco treatment (10min 4°C)

α-tubulin PCNT   α-tubulin PCNT   PCNT

Cold (10min 4°C) → Noco treatment (1h 4°C → 1h 37°C)

α-tubulin PCNT   α-tubulin PCNT   PCNT

5µm   5µm

Figure 5

Figure 6

Figure 7

Figure 8

A 300 adhesive PCM complexes (all MT-nucleating)

B 300 non-adhesive PCM complexes (all MT-nucleating)

C 300 adhesive PCM complexes (50 MT-nucleating) 300 immobile MTs

D 300 strongly adhesive PCM complexes (50 MT-nucleating) 300 immobile MTs

E

F

G 300 non-adhesive CAMSAP-kin14 complexes (all MT-nucleating)

H 300 adhesive CAMSAP-kin14 complexes (all MT-nucleating)

I 150 non-adhesive CAMSAP-kin14 complexes (all MT-nucleating) 150 adhesive PCM complexes (all MT-nucleating)

J 150 non-adhesive CAMSAP-kin14 complexes (all MT-nucleating) 150 non-adhesive PCM complexes (all MT-nucleating)

K

L

Figure 1 - figure supplement 2

Figure 1 - figure supplement 3

Figure 2- figure supplement 1

Figure 2 - figure supplement 2

Figure 2 - figure supplement 3

A AKAP450/CAMSAP2/p53 KO
α-tubulin   PCNT   Centrin   Merge

+DMSO

+CentB

10µm

B AKAP450/CAMSAP2/p53 KO + CentB

00:00 hh:mm   01:00   02:00   03:00   04:00   05:00

06:00   06:20   06:30   06:35   06:40   06:50

07:00   07:40   07:45   07:50   07:55   08:00

08:20   08:30   08:50   09:00   09:30   10:30

10:35   10:50   11:00   11:20   11:50   12:20

20µm

C AKAP450/CAMSAP2/p53 KO + CentB

Prophase   Metaphase   Anaphase   Telophase

00:00 hh:mm   00:40   01:04   01:08

10µm

D

Figure 2 - figure supplement 4

A Control=AKAP450/CAMSAP2/p53 KO + CentB

PCNT KO=AKAP450/CAMSAP2/p53/PCNT KO + CentB

Figure 2 - figure supplement 5

Figure 4 - figure supplement 1

A AKAP450/CAMSAP2 KO stable GFP-CDK5RAP2 + CentB

Noco treatment

00:00 mm:ss 00:02 00:18 00:24 00:32 00:34 00:38 00:44 00:52 00:56

5µm 5µm

B AKAP450/CAMSAP2 KO + CentB

-Noco +Noco (10µM, 15min)

10µm 10µm

Directionality-filtered reconstruction of motor tracks

10µm 10µm

C AKAP450/CAMSAP2 KO + CentB, Cold treatment (30min 4°C)

α-tubulin PCNT α-tubulin PCNT α-tubulin PCNT

5µm 5µm

C Presence of PCM proteins in clusters before and after Noco washout

CentB	-	+	+	+	+	+
Nocodazole	-	-	+	+	+	+
Washout			30s	2min	5min	10min
PCNT	+	+	+	+	+	+
CDK5RAP2	+	+	+	+	+	+
γ-tubulin	+	+	+	+	+	+
NEDD1	+	-	-	-	-	-
Ninein	+	+	-	+	+	+
CEP192	+	-	-	-	-	-
PCM1	+	+	+	+	+	+

E Nocodazole washout 2 min non-radial part (%)

F AKAP450/CAMSAP2 KO + CentB

Figure 5 - figure supplement 2

Figure 6 - figure supplement 1

Figure 6 - figure supplement 2

Figure 7 - figure supplement 1

AKAP450/CAMSAP2/p53/PCNT KO + CentB

Overexpression FKBP-mCh-CAMSAP2

FKBP-mCh-CAMSAP2 | α-tubulin | Merge

-Rapalog / +Rapalog

Overexpression FRB-GFP-ppKin14

FRB-GFP-ppKin14 | α-tubulin | Merge

-Rapalog / +Rapalog

5µm | 5µm

AKAP450/CAMSAP2/p53/PCNT KO + CentB

Overexpression FKBP-mCh-CAMSAP2, FRB-GFP-ppKin14

FKBP-mCh-CAMSAP2 | FRB-GFP-ppKin14 | α-tubulin | Merge

-Rapalog / +Rapalog

5µm

Figure 7 - figure supplement 2

A AKAP450/CAMSAP2/p53/PCNT KO, +CentB

-Rapalog | +Rapalog

FKBP-CAMSAP2
00:00:00 hh:mm:ss | 00:10:00 | 00:13:00 | 00:17:24 | 00:27:48 | 00:50:18 | 01:39:06

FRB-ppKin14
00:00:00 | 00:10:00 | 00:13:00 | 00:17:24 | 00:27:48 | 00:50:18 | 01:39:06

Merge
00:00:00 | 00:10:00 | 00:13:00 | 00:17:24 | 00:27:48 | 00:50:18 | 01:39:06

10µm

B AKAP450/CAMSAP2/p53/PCNT KO + CentB

Untransfected -Rapalog
α-tubulin | Calnexin | Merge | Merge | α-tubulin | Calnexin

+Rapalog
α-tubulin | Calnexin | FKBP-CAMSAP2 | Merge | Merge | FKBP-CAMSAP2 | Calnexin

AKAP450/CAMSAP2/MMG/CDK5RAP2/p53/PCNT KO + CentB

+Rapalog

Ring-like
α-tubulin | Calnexin | FKBP-CAMSAP2 | Merge | Merge | FKBP-CAMSAP2 | Calnexin

Bundled

Ring-like
α-tubulin | Tom20 | FKBP-CAMSAP2 | Merge | Merge | FKBP-CAMSAP2 | Tom20

Bundled

5µm | 5µm

A AKAP450/CAMSAP2/MMG/CDK5RAP2 KO

p53 sgRNA targets

sg 1   sg 2

exon 4

PAM1   PAM2

...CCCATCTACAGTCCCCCCTTGCCGTCCCAAGCAATGGATGATTTGATGCTGTCCCCGGACGATATTGAACAATGGTTCACTGAAGA...

37 bp

TGCTCTTTTCACCCATCTACAGTCCCCCTTGACGATATTGAACAATGGTTCACTGAAGACCCA

allele 1+2

PAM1

B Mouse anti p53 (N)
(target 11-25aa)

Rabbit anti p53(C)
(target 325-375aa)

p53 protein (393aa)

sg1 sg2

* =mutated from 36th aa
* =1st stop codon at 110th aa
■ =sg1 covers the 35-41 aa
■ =sg2 covers the 47-53 aa

C

| | p53 (N) | p53(C) | DAPI | Merge |

AKAP450/CAMSAP2/
MMG/CDK5RAP2 KO

AKAP450/CAMSAP2/
MMG/CDK5RAP2/
p53 KO

10μm

D AKAP450/CAMSAP2/MMG/CDK5RAP2/p53 KO

sg e5-1   sg e5-2

PCNT sgRNA targets

exon 5

PAM1   PAM2

...GGAGAAGGAGACGGCATTGACGGAGCTGCGGGAGATGCTCAACAGCCGGCGTGCCCAGGAGCTGGCCCTGCTACA...

28 bp

AAGGAGAAGGAGACGGCATTGACGGAGCCCAGGCAACTGGCCCTGCTACCGACC

allele 1+2

28 bp deletion CCCAGGAGCTGGCCCTGCTACAGAGCAGGCA

E Rabbit anti PCNT
(target 100-600aa)

PCNT protein(3336aa)

* =mutated from 280th aa
* =1st stop codon at 319th aa

F

| | NIN | PCNT | Merge |

AKAP450/CAMSAP2/
CDK5RAP2/MMG/
p53 KO

AKAP450/CAMSAP2
/CDK5RAP2/MMG/
p53/PCNT KO

10μm

H AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/PCNT KO

| α-tubulin | CEP192 | Merge | CEP192 | α-tubulin |

-CentB

+CentB

5μm

5μm

G

| | AKAP450/CAMSAP2 KO | AKAP450/CAMSAP2/p53/PCNT KO | AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/PCNT KO |

kDa

PCNT

250-

p53

50-

100-

Ku80

Source data 1
Uncropped Western blots shown in this manuscript

Source data 2
Uncropped Western blots shown in this manuscript

A

Ctl AKAP450/CAMSAP2/p53 KO   Ctl AKAP450/CAMSAP2/p53 KO

anti p53

anti Actin

B

Ctl AKAP450/CAMSAP2/p53/PCNT KO

anti PCNT

anti Ku80

C

AKAP450/CAMSAP2/p53 KO
AKAP450/CAMSAP2/p53/PCNT KO
AKAP450/CAMSAP2/p53 KO
AKAP450/CAMSAP2/p53/PCNT KO

CentB - - + +

anti CDK5RAP2

anti Ku80

D

AKAP450/CAMSAP2/p53 KO
AKAP450/CAMSAP2/p53/PCNT KO
AKAP450/CAMSAP2/p53 KO
AKAP450/CAMSAP2/p53/PCNT KO

CentB - - + +

anti Ku80

anti γ-tubulin

anti NIN

E

EB1/EB3 mut   AKAP450/CAMSAP2/EB1/EB3 mut
EB1/EB3 mut   AKAP450/CAMSAP2/EB1/EB3 mut
EB1/EB3 mut   AKAP450/CAMSAP2/EB1/EB3 mut

anti AKAP450

anti CAMSAP2

anti Ku80

(Short Exposure)   (Long Exposure)

F

Ctl
AKAP450/CAMSAP2/p53/PCNT KO
AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/PCNT KO
Ctl
AKAP450/CAMSAP2/p53/PCNT KO
AKAP450/CAMSAP2/CDK5RAP2/MMG/p53/PCNT KO

anti PCNT

anti Ku80

anti p53