Machine learning prediction of antiviral-HPV protein interactions for anti-HPV pharmacotherapy

Hui-Heng Lin [a*] (ORCID 0000-0003-4060-7336), Qian-Ru Zhang [b] (ORCID 0000-0001-8986-5219), Xiangjun Kong [c], Liuping Zhang [d] (ORCID 0000-0001-7696-0161), Yong Zhang [e] (ORCID 0000-0003-1820-0927), Yanyan Tang [f] (ORCID 0000-0002-1189-4498) and Hongyan Xu [d*] (ORCID 0000-0003-2402-0973)

[a]   Yuebei People's Hospital, Shantou University Medical College, No. 133 of Huimin South road, Wujiang District, Shaoguan City, Guangdong Province, 512025, China.

[b]   School of Pharmacy, Key Lab of the Basic Pharmacology of the Ministry of Education, Zunyi Medical University, 6 West Xue-Fu Road, Zunyi city, Guizhou Province, 563000, China

[c]   State Key Laboratory of Quality Research in Chinese Medicine, Institute of Chinese Medical Sciences, University of Macau, Avenida de Universidade, Taipa, Macau, 999078, China.

[d]   Department of Gynecology, Yuebei People's Hospital, Shantou University Medical College, No. 133 of Huimin South road, Wujiang District, Shaoguan City, Guangdong Province, 512025, China.

[e]   Interdisciplinary Research Center for Agriculture Green Development in Yangtze River Basin, Southwest University, No.1-2-1 Tiansheng Road, Beibei District,

Chongqing, 400715 China.

f   Department of Neurology, the First Affiliated Hospital of Guangxi Medical

University, No.6 Shuangyong Road, Nanning, Guangxi, 530021 China.

*   Correspondence to Hui-Heng Lin scienceds@outlook.com and Hongyan Xu:

medxh@outlook.com . Corresponding Postal Address: Yuebei People's hospital, No.

133 of Huimin South road, Wujiang District, Shaoguan City, Guangdong Province,

512025 China.

Email of coauthors:

- Qian-Ru Zhang: qrzhang_iris@foxmail.com

- Xiangjun Kong: xiangjunkong@um.edu.mo

- Liuping Zhang: lpzhang0912@163.com

- Yangyan Tang: yuji0301@126.com

- Yong Zhang: bioyongzhang@swu.edu.cn

Abstract

Background: Persistent infection with high-risk types Human Papillomavirus could

cause diseases including cervical cancers and oropharyngeal cancers. Nonetheless, so

far there is no effective pharmacotherapy for treating the infection from high-risk HPV

types, and hence it remains to be a severe threat to health of female.

Methods: In light of drug repositioning strategy, we trained and benchmarked multiple machine learning predictive models so as to predict potential effective antiviral drugs for HPV infection in this work. Based on antiviral-target interaction dataset, we generated high dimension feature set of drug-target interaction pairs and used the dataset to train and construct machine learning predictive models.

Results: Through optimizing models, measuring models' predictive performance using 182 pairs of antiviral-target interaction dataset which were all approved by United States Food and Drug Administration, and benchmarking different models' predictive performance, we identified the optimized Support Vector Machine and K-Nearest Neighbor classifier with high precision score were the best two predictors (0.80 and 0.85 respectively) amongst classifiers of Support Vector Machine, Random forest, Adaboost, Naïve Bayes, K-Nearest Neighbors, and Logistic regression classifier. We applied these two predictors together and successfully predicted 58 pairs of antiviral-HPV protein interactions from 846 pairs of antiviral-HPV protein associations.

Conclusions: Our work provided good drug candidates for anti-HPV drug discovery. So far as we know, we are the first one to conduct such HPV-oriented computational drug repositioning study.

Keywords

Machine Learning Prediction; Human Papillomavirus; HPV infection; Antivirals; Drug repositioning; Drug-Protein interaction prediction; Support Vector Machine; K-Nearest

64   Neighbors

Introduction

Human Papillomavirus (HPV) can infect human body and cause different types of

phenotypes. Specifically, HPVs can infect females' reproductive system and causes

different types of gynecological diseases. For instance, a variety of warts and the genital

cancers [1]. What's more, it is reported that HPV infection is one of the risk factors of

oropharyngeal cancer [2]. Researchers have classified the subtypes of HPVs into the

low-risk types, and high-risk types according to their virulence and relevant risk levels

of infections. For low-risk types [3], e.g., the type 6, 11, 40, etc, they might be

disappeared after several periods of infection, and the infected hosts might generally be

fine. While for those high-risk subtypes of HPV, e.g., the HPV-16 and HPV-18, their

persistent infection on hosts could finally result to severe disease like cervical cancer [4],

which could results to the death of hosts. According to report, it is estimated that

569,000 cases of cervical cancer newly occurred in 2018 globally and 311,000 deaths

were found [5]. Therefore, the HPV infection remains a large threat to female's health

especially in developing countries, and treating HPV infection remains an urgent task

and difficult challenge due to there lacks effective pharmacotherapy. Though HPV

vaccines are available, they are ineffective for those who has already been infected by

HPVs [6].

Scientists have been trying hard to combat against HPV infection. For instance, several

researchers have identified HPV's E6 and E7 proteins to be the virulent tumorigenesis

risk factors [7, 8], and parts of their molecular 3D structural conformations have

revealed through approaches of *in silico* simulation [9] and structural biology [10]. Other

studies have tested, discussed and reviewed the *in vitro* effects of existed drug, i.e., the

Human Immunodeficiency Virus (HIV) protease inhibitor, on HPV proteins and cells

infected with HPVs [11, 12, 13, 14, 15]. These reports targeting existed drugs for HPV

treatments showed that, compared with *de novo* drug discovery, repositioning exited

drugs is indeed the better and quicker strategy.

Nonetheless, drug efficacies from above evidences were moderate and no further

progress is seen in later stages, .e.g., in clinical contexts. And hence, above research

progresses are yet far from being able to identify drug candidates with good therapeutic

and anti-HPV potential. Limitation of them could be due to two reasons. One is that

inappropriate compound or drug candidates have been chosen for testing. The other

reason could be the number of drug candidates to be tested is too small. Testing only

limited numbers of compound or drug candidates surely restricts the probability of

identifying those appropriate ones.

In order to meet the urgent needs for effective anti-HPV drug discovery, based on

target-oriented drug repositioning strategy, we collected and analyzed 96 antiviral drugs

to do the relatively large-scale *in silico* screening for 9 HPV-16 proteins, so as to

computationally and effectively identify effectively antivirals with good potential for

targeting HPV proteins. Briefly, in this work, we constructed, benchmarked and selected

machine learning predictive models (also known as predictors) to predict antivirals that

could have potential interactions with HPV proteins. This is because Drug-target

interactions are vital prerequisite of molecular therapeutic mechanisms. Through

benchmarking, we selected the high-precision K-Nearest Neighbor (KNN) [16] and

Support vector machine (SVM) [17] predictors to detect those confidence interaction

pairs of antiviral-HPV protein.

To the best of our knowledge, no prior study similar to our work has been done. Lots of

researchers predicted targets of drugs, co   mpound-protein   interactions   or

protein-protein interactions using machine learning or other computational methods [18,

19, 20, 21]. However, so far as we know, no study has focused on studying relationships

between antiviral drugs and HPV proteins.

Methods

Research question formulation

Theoretically, a therapeutic target and its drug molecule have interactive binding

relation to each other. Therefore, trying to identify potential HPV protein targets of

antivirals could be considered as a binary classification task, i.e., to predictively classify

proteome of HPV into two classes of proteins. One class is HPV proteins which have

potential interaction with drug molecules, and the other class is HPV proteins do not

have potential interactions with drugs. Machine learning is a good and state-of-art

methods to solve such binary classifications **(Figure 1)**. Considered that known antiviral

drug-target interaction pairs were available, which could serve as the known-label

validation dataset, we thus chose supervised (machine) learning methods for this study.

Figure 1: Research framework of this study. Predicting antiviral drug-HPV protein interaction could be

considered a binary classification task, and machine learning is a good method for such task. In this work,

antiviral drug-target pairs' features were transformed into vectors for constructing machine learning

predictors. Through benchmarking, the best predictors were selected to predict antiviral-HPV protein

interactions.

Data collection and preprocessing

We collected antiviral drugs and their associated target data from DrugBank [22],

Drugs@FDA [23], PubChem [24], Uniprot [25] and Therapeutic Targets [26] databases

(As of 19th July 2020). Drug-target interaction pairs which contained the United States

Food and Drug Administration (FDA)-approved antiviral drugs were treated as the

validation dataset for machine learning, while the rest drug-target interaction pairs were

treated as the training dataset for machine learning. In this work's machine learning

classification task, an interaction pair of an antiviral drug and a protein was defined to

be a positive instance, while negative instance indicated a non-interactive pair of

antiviral and protein. In order to balance data ratio for binary machine learning

classification task. We randomly generated non-interactive drug-target pairs so as to

assure the 1:1 ratio of positive instances to negative instances for machine learning.

Next, we integrated the proteome (9 proteins in total) of high-risk HPV-16 subtype and

all the antiviral drugs to form drug-protein interaction prediction dataset. See

Supplementary Table 1 for machine learning training dataset of antiviral drug-target

interaction pairs, **Supplementary Table 2** for drug-target interaction pair dataset used in

machine learning validation process, and **Supplementary Table 3** for Uniprot's HPV-16

proteome, i.e., 9 proteins.

Next, all antivirals' molecular structures were analyzed using ChemmineR [27] and

1024-dimension chemical fingerprint datasets were generated through R scripting [28].

All proteins were analyzed using ProtR [29] and 10784 high dimension protein

descriptor feature datasets were generated. As seen in **Table 1**. Descriptors used were

protein structural and physicochemical properties. These descriptors have been widely

used in and performed well for analyzing protein-protein interactions and protein-ligand

interactions [29].

All datasets were integrated, scaled and normalized using R computing environment

[28].

Table 1: Molecular descriptors used for machine learning analysis

ID	Molecular descriptor	Vector length
1	Drug molecule fingerprint	1024
2	Amino acid composition	20
3	Dipeptide composition	400
4	Tripeptide composition	8000
5	Normalized Moreau-Broto Autocorrelation	240
6	Moran Autocorrelation	240
7	Geary Autocorrelation	240

8	Composition descriptor	21
9	Transition descriptor	21
10	Distribution descriptor	105
11	Pseudo amino acid composition	50
12	Amphiphilic pseudo amino acid composition	80
13	Conjoint Triad	343
	Total	10784

Machine learning and prediction

Briefly, the machine learning processes of this research work followed such order and general steps. Using training dataset, 5-fold cross validation and grid searching were applied to different machine learning algorithms for obtaining identifying the best parameters of the best predictive power. Later, predictors with good performances were further applied to classify the validation dataset with known labels. Lastly, the verified best predictor was used to predict antiviral-HPV protein interaction pairs.

Diverse sorts of supervised learning algorithms with different purposes exist. Amongst, the Support Vector Machine, Random Forest [30], Logistic Regression [31], etc, are classic algorithms for tackling the binary classification questions. Initially, 6 types of machine learning classifiers friendly for binary classification were chosen for building

predictive models. The chosen predictors were Support Vector Machine, Random Forest,

AdaBoost [32], Logistic Regression, Naïve Bayes [33] and K-Nearest Neighbor

classifier. Next, with default parameters, 6 predictors went over simple checking

through quick training and performance measurement. At this early stage, as expected,

all predictors did not perform well. Subsequently, in order to identify better parameters

for predictors, grid search 5-fold cross validations and performance benchmarking were

conducted. The predictive performance of 6 different predictors with better parameters

were tested using known-label validation dataset. Upon checking performance of

different predictors, we selected the optimized K-Nearest Neighbors classifier, which

had the highest precision score and was the most appropriate predictor to identify high

confidence drug-protein interaction pairs from 864 pairs of antiviral-HPV protein

associations.

Aforementioned data processing and machine learning computations were done via

in-house scripts of Python [34] and R [28]. Libraries and modules used were Sci-kit

learn [35], Pandas [36], Numpy [37, 38] Scipy [39], and also Bioconductor [40, 41] and

Biomart [42].We acknowledge authors and developers of these computational tools.

204 Results

Dataset overview

The antiviral drugs and their associated targets were retrieved and analyzed as described

in method section. **Table 2** provides a summary of our dataset. We had totally 61

antiviral drugs, which formed totally 284 antiviral-target interaction pairs with their

targets. For the purpose of measuring machine learning predictors' performance,

antiviral-target interactions were split in to two classes where 102 pairs were used as the

dataset for training or fitting machine learning predictors, and the rest 182 pairs were

treated as dataset for validating the predictive performances of machine learning

predictors. And we also compiled 9 proteins of HPV (its complete proteome) with 96

antiviral drugs to form 864 pairs of antiviral-HPV protein association pairs (**Table 2**).

Table 2: Summary of antiviral-target and antiviral-HPV protein interaction dataset used in machine

learning processing of this study

Dataset	Antiviral drug	Target protein	Antiviral-protein interaction pair
Training set	35	34	102 [c]
Validation set [a]	61	47	182 [c]
Prediction set	96	9 [b]	864

218   [a] Validation set consisted of U.S.FDA-approved antiviral drugs and these drugs' binding target proteins

219   [b] 9 proteins of HPV-16.

220   [c] Ratio of positive instance to negative instance was 1:1.

Performances of machine learning models

Initially, we chose 6 type of machine learning models and applied 5-fold cross validation strategy to fit the antiviral-target interaction training dataset. A primary benchmarking of the predictive performance of 6 chosen predictors were as seen in **Table 3**.

Table 3: Performance of 6 machine learning predictors with default parameters.

ID	Predictor	Precision	Recall	F1-measure	Accuracy	AUC [a]
1	SVM	0.36	0.48	0.37	0.48	0.44
2	Logistic Regression	0.52	0.57	0.54	0.59	0.56
3	KNN	0.61	0.62	0.60	0.59	0.61
4	Naïve Bayes	0.46	0.65	0.52	0.49	0.52
5	Random Forest	0.61	0.61	0.63	0.66	0.68
6	AdaBoost	0.73	0.48	0.55	0.62	0.48

[a] AUC indicates the metric of Area Under Curve of Receiver-Operating Characteristic Curve.

Briefly, all predictors' predictive performances were less satisfying, as expected. SVM with default parameter (RBF kernel) performed the worst in all sorts of metrics among 6 predictors. AdaBoost classifier scored the best in terms of precision score but had the lowest recall value in recall. F1-measure is the harmonic average of precision and recall. The highest F1-measure was found from Random forest classifier, which was 0.63.

While we also found other metrics of Random forest were not high. All its metrics were

round 0.65 though the values were close to each other. The highest accuracy score and

AUC (Area Under Curve of Receiver-Operating Characteristic Curve) of 6 predictors'

were 0.66 and 0.68, respectively. And both of them were also found in Random forest's

performance. Metrics of default parameters' KNN were all around 0.6, indicating its

unsatisfying performances in 5-fold cross validation, too. Similar to KNN, Naïve bayes

classifier did not perform well, and one common point of KNN and Naïve bayes

classifier was that, the value of their recall score was higher than those of other metrics

(**Table 3**).

Next we tuned parameters of predictors through grid searching 5-fold cross validation,

and tested how combination set of parameters affected predictors' predictive

performances on known-label validation dataset. At beginning, we focused on optimize

predictors for obtaining better values of comprehensive metrics, such as the F1-measure,

accuracy or AUC value. Despite a great number of times' trying, no high sores of

aforementioned F1-measure, accuracy or AUC metric value was seen.

Given that high precision score indicates the low number of predictive false positive

instances, and high recall score indicates the low number of predictive false negative

instances, we changed our strategy and decided to do high precision-oriented

optimization. This was because the purpose of this work was to identify antivirals which

interact with HPV proteins. To this end, using high-precision predictor, predictive

positive instances could had lower false positive instance mixed inside. Therefore, in

this work, we preferred precision metric over recall metric for selecting appropriate

predictors to predict antiviral-HPV protein interactions (positive instances). Through

benchmarking the performances of predictors, we found optimized SVM and KNN

predictors had better precision score than others. SVM's was 0.8 and the KNN

classifier's was 0.85 **(Table 4)**. We hence used them for prediction task and we chose

the intersection of their prediction results as the final results.

Table 4: Precision scores of optimized machine learning predictors on the validation dataset of

antiviral-HPV protein interaction pairs

Predictor	**SVM**	**Logistic Regression**	**KNN**	**Naïve Bayes**	**Random Forest**	**AdaBoost**
Precision Score	**0.80**	0.50	**0.85**	0.65	0.68	0.75

Predicted antiviral-HPV protein interaction pairs

Upon selection of high-precision predictors, we applied them to predict the

antiviral-HPV protein interactions. We selected two predictors' result intersection as the

final prediction result, i.e., we only consider an antiviral-HPV protein association pair

has potential interaction if both predictors predicted this pair to be position (interactive).

As a result, within 864 antiviral-HPV association pairs, most antiviral-HPV protein

pairs were predicted to be negative, i.e., the antiviral drug does not interact with the

HPV protein. Only a small portion, i.e., 58, of antiviral-HPV protein pairs were

predicted to have interaction. Prediction results were summarized in **Table 5** in HPV

protein-oriented form**.** Full prediction results could be found in **Supplementary Table 3**.

Here we took the Docosanol as an example for analysis. The drug Docosanol was

predicted to interact with HPV-16's protein E7 using our high-precision machine

learning predictors. Docosanol is a U.S. FDA-approved antiviral drug targeting

Envelope glycoprotein GP350 and GP340 of Epstein-Barr Virus (EBV, also known as

Human Herpesvirus or HHV-4) and it is used to treat fever blisters, etc. Interestingly,

through literature survey, a recently published clinical case report was found to claim

that, the mixture usage of Docosanol, curcumin, and other drugs together treated HPV

infection and vaginal warts of a patient well [43]. This could be an evidence supporting

our predictive result about Docosanol and HPV protein. HPV protein-oriented antiviral

prediction results were summarized in **Table 5** and briefly description of the example

antiviral drugs, protein targets of the antiviral drug and relevant therapeutic indications

were also listed in **Table 5.**

Table 5: Summary of prediction result of antivirals targeting each protein of HPV-16.

HPV-16 protein	Number of antivirals [a]	Example
Protein E7	7	Docosanol targeting GP340 or GP350 protein of Epstein-Barr Virus has been approved to treat herpes labialis, fever blisters, etc.
Regular Protein E2	5	Voxilaprevir targeting NS3/4A protein of Hepatitis C Virus has been approved to treat chronic Hepatitis C caused by Hepatitis C Virus infection.
Protein E6	6	Telaprevir is an NS3/4A viral protease inhibitor. It has been approved to treat chronic Hepatitis C Virus infection in combination with other drugs.
Minor capsid protein L2	4	Grazoprevir targeting NS3/4A protein of Hepatitis C Virus has been approved to treat Hepatitis C viral infection.
Protein E4	8	Nelfinavir is a potent viral protease inhibitor for treating infections of Human Immunodeficiency Virus (HIV), and it targets the protease of HIV -1.
Probable protein E5	7	Maraviroc is a chemokine receptor antagonist drug targeting C-C chemokine receptor type 5. It has been approved to treat HIV-1 infection

Replication protein E1	7	Pirodavir (investigational drug) targets the genome polyprotein of Polioviruses and it seems to have broad-spectrum antiviral effects on multiple kinds of Human Rhinoviruses.
Major capsid protein L1	5	Docosanol targeting GP340 or GP350 protein of Epstein-Barr Virus has been approved to treat herpes labialis, fever blisters, etc.
Protein E8^E2C	7	TMC-310911 (investigational drug) is a protease inhibitor targeting HIV-1 protease and it seems to have effect on treating HIV-1 infection.

295   [a] Indicating the number of antivirals which was predicted to have potential interaction with specific

296   HPV-16 protein.

Discussion

While our results are to be validated by *in vitro* assays, in this work, we constructed

machine learning models, and predicted antiviral-HPV protein interactions so as to

identify potential drug candidates targeting HPV proteins. The high-risk types of HPV

are not limited to HPV-16. There are other types such as HPV-18. Indeed, we are not

only able to apply the research framework of this study to predict the potential drug

candidates for the proteome of other HPV subtypes, but also to other types of

pathogenic and infectious microbes, as well.

Limitation

Reviewing this current study, we found several significant points that could help us do better preparation for further works. Initially, in this work, though we tried our best to collect more antiviral drugs, due to the availability of antiviral drugs, we had limited size of dataset for machine learning. This could be one of factors why we did not obtain predictors with high score of F1-measure, accuracy, or AUC. Compared with antivirals, the number of other types of drugs, e.g., the cancer drugs or antibiotics, are higher. Thus, in future study, we would consider using other types of drugs for repositioning purpose.

Also, the final predictors selected did not have high F1-measure, accuracy, or AUC. Because current machine learning processes are black box which is difficult to interpret. Alternatively, in this study, considered the tradeoff between precision and recall, we chose to select the intersected prediction results from two high-precision predictors in order to get higher confidence antiviral-HPV protein interactions. For future study, we would learn and try to apply the state-of-art explainable machine learning methods which may be interpretable. In such case, we may be able to find out reasons causing low performances and obtaining guidance for model optimizations and obtaining more powerful machine learning predictors.

Conclusions

Inspired by the needs of anti-HPV drug discovery, drug repositioning and computational

analytics, we designed this research project and constructed machine learning models to

predict possible antiviral-HPV protein interactions so as to identify potential

pharmacotherapy for HPV infection. As a result, we optimized the predictors and

identified 58 antivirals which have potential interaction with proteins of HPV-16's.

To the best of our knowledge, we are the first pioneer to conduct this HPV-oriented

computational antiviral repositioning study. No similar study has been found so far.

Therefore, our work provides good insights to virologists, medicinal chemists,

gynecologist, clinical microbiologists, etc, those who are interested in the treatment and

therapy of HPV infections. Also, drug candidates pre-selected via computational

analytic screening could have lower probability of ineffectiveness than those that did

not go through computational analyses. It thus could save resources, and antivirals

identified by us could be good candidates for further *in vitro* and *in vivo* tests. In such

way, this work contributes to drug development for HPV infections. What is more, our

predicted antiviral-HPV protein interaction pairs also offer insights for fundamental

biomedical research on drug-protein interactions or molecular interaction mechanisms.

The last but not the least, the research framework of this study, i.e., the machine

learning-based compound-protein interaction prediction, could also be applied to

primary drug repositioning or drug discovery for those diseases or infectious microbial

pathogens lacking effective pharmacotherapy. E.g., the Noroviurs.

Supplementary Materials

Supplementary Materials are available in this study.

- Supplementary Table 1 is the antiviral-target interaction dataset for machine learning training processing.
- Supplementary Table 2 included drug-target interaction pair dataset used in machine learning validation process.
- Supplementary Table 3 included dataset of HPV-16 proteome.

Data Availability Statement

Data of this study were included in the supplementary materials

Author Contributions

- Conception of this work: Hui-Heng Lin and Hongyan Xu;
- Acquisition of datasets: Hui-Heng Lin, Xiangjun Kong and Qian-Ru Zhang;
- Analysis and interpretation of data: Hui-Heng Lin, Liuping Zhang
- Drafting or reviewing manuscript: Hui-Heng Lin, Yanyan Tang, and Yong Zhang.
- Approval of submission: All authors approved submission.

Funding Statement

This research did not received any specific grant from funding agencies in the public, commercial, or not-for-profit sections. Funding sources have no role in conducting, writing and publishing this research work.

References

[1]. Ljubojevic S, Skerlev M. HPV-associated diseases. Clinics in dermatology. 2014

Mar 1;32(2):227-34.

[2]. Fakhry C, Zhang Q, Nguyen-Tan PF, Rosenthal D, El-Naggar A, Garden AS,

Soulieres D, Trotti A, Avizonis V, Ridge JA, Harris J. Human papillomavirus and

overall survival after progression of oropharyngeal squamous cell carcinoma.

Journal of clinical oncology. 2014 Oct 20;32(30):3365.

[3]. Muñoz N, Bosch FX, De Sanjosé S, Herrero R, Castellsagué X, Shah KV, Snijders

PJ, Meijer CJ. Epidemiologic classification of human papillomavirus types

associated with cervical cancer. New England journal of medicine. 2003 Feb

6;348(6):518-27. doi:10.1056/NEJMoa021641. hdl:2445/122831.

[4]. Wardak S. Human Papillomavirus (HPV) and cervical cancer. Medycyna

doswiadczalna i mikrobiologia. 2016 Jan 1;68(1):73-84.

[5]. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer

statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for

36 cancers in 185 countries. CA: a cancer journal for clinicians. 2018

Nov;68(6):394-424.

[6]. Markowitz LE, Dunne EF, Saraiya M, Lawson HW, Chesson H, Unger ER.

Quadrivalent human papillomavirus vaccine: recommendations of the Advisory

Committee on Immunization Practices. MMWR Morb Mortal Wkly Rep.

2007;56(RR-02):1-24.

[7]. Ganguly N, Parihar SP. Human papillomavirus E6 and E7 oncoproteins as risk

factors for tumorigenesis. Journal of biosciences. 2009 Mar 1;34(1):113-23.

doi*:*10.1007/s12038-009-0013-7.

[8]. Tang S, Tao M, McCoy Jr JP, Zheng ZM. The E7 oncoprotein is translated from

spliced E6* I transcripts in high-risk human papillomavirus type 16-or type

18-positive cervical cancer cell lines via translation reinitiation. Journal of virology.

2006 May 1;80(9):4249-63. doi:10.1128/JVI.80.9.4249-4263.2006.

[9]. Ricci-López J, Vidal-Limon A, Zunñiga M, Jimènez VA, Alderete JB, Brizuela CA,

Aguila S. Molecular modeling simulation studies reveal new potential inhibitors

against HPV E6 protein. PloS one. 2019 Mar 15;14(3):e0213028.

[10].   Zanier K, Charbonnier S, Sidi AO, McEwen AG, Ferrario MG,

Poussin-Courmontagne P, Cura V, Brimer N, Babah KO, Ansari T, Muller I.

Structural basis for hijacking of cellular LxxLL motifs by papillomavirus E6

oncoproteins. Science. 2013 Feb 8;339(6120):694-8. doi*:*10.1126/science.1229934.

[11].   Bernstein WB, Dennis PA. Repositioning HIV protease inhibitors as cancer

therapeutics. Current Opinion in HIV and AIDS. 2008 Nov;3(6):666.

[12].   Hampson L, Oliver AW, Hampson IN. Using HIV drugs to target human

papilloma virus. Expert review of anti-infective therapy. 2014 Sep 1;12(9):1021-3.

[13].   Hampson L, Kitchener HC, Hampson IN. Specific HIV protease inhibitors

inhibit the ability of HPV16 E6 to degrade p53 and selectively kill E6-dependent

cervical carcinoma cells in vitro. Antiviral therapy. 2006;11(6):813–25.

[14].   Kim DH, Jarvis RM, Allwood JW, Batman G, Moore RE, Marsden-Edwards E,

Hampson L, Hampson IN, Goodacre R. Raman chemical mapping reveals site of

action of HIV protease inhibitors in HPV16 E6 expressing cervical carcinoma cells.

Analytical and bioanalytical chemistry. 2010 Dec;398(7):3051-61.

[15].   Kim DH, Allwood JW, Moore RE, Marsden-Edwards E, Dunn WB, Xu Y,

Hampson L, Hampson IN, Goodacre R. A metabolomics investigation into the

effects of HIV protease inhibitors on HPV16 E6 expressing cervical carcinoma

cells. Molecular BioSystems. 2014;10(3):398-411.

[16].   Guo G, Wang H, Bell D, Bi Y, Greer K. KNN model-based approach in

classification. In OTM Confederated International Conferences. On the Move to

Meaningful Internet Systems" 2003 Nov 3 (pp. 986-996). Springer, Berlin,

Heidelberg.

[17].   Noble WS. What is a support vector machine?. Nature biotechnology. 2006

Dec;24(12):1565-7.

[18].   Chen R, Liu X, Jin S, Lin J, Liu J. Machine learning for drug-target interaction

prediction. Molecules. 2018 Sep;23(9):2208.

[19].   Zhang W, Lin W, Zhang D, Wang S, Shi J, Niu Y. Recent advances in the

machine learning-based drug-target interaction prediction. Current drug metabolism.

2019 Mar 1;20(3):194-202.

[20].   Liu S, Liu C, Deng L. Machine learning approaches for protein–protein

interaction hot spot prediction: Progress and comparative assessment. Molecules.

2018 Oct;23(10):2535.

[21]. Das S, Chakrabarti S. Classification and prediction of protein–protein

interaction interface using machine learning algorithm. Scientific reports. 2021 Jan

19;11(1):1-2.

[22]. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, Sajed T,

Johnson D, Li C, Sayeeda Z, Assempour N. DrugBank 5.0: a major update to the

DrugBank database for 2018. Nucleic acids research. 2018 Jan 4;46(D1):D1074-82.

[23]. Schwartz LM, Woloshin S, Zheng E, Tse T, Zarin DA. ClinicalTrials. gov and

Drugs@ FDA: a comparison of results reporting for new drug approval trials.

Annals of internal medicine. 2016 Sep 20;165(6):421-30.

[24]. Kim S, Chen J, Cheng T, Gindulyte A, He J, He S, Li Q, Shoemaker BA,

Thiessen PA, Yu B, Zaslavsky L. PubChem 2019 update: improved access to

chemical data. Nucleic acids research. 2019 Jan 8;47(D1):D1102-9.

[25]. UniProt Consortium. UniProt: a hub for protein information. Nucleic acids

research. 2015 Jan 28;43(D1):D204-12.

[26]. Zhu F, Han B, Kumar P, Liu X, Ma X, Wei X, Huang L, Guo Y, Han L, Zheng

C, Chen Y. Update of TTD: therapeutic target database. Nucleic acids research.

2010 Jan 1;38(suppl_1):D787-91.

[27]. Cao Y, Charisi A, Cheng LC, Jiang T, Girke T. ChemmineR: a compound

mining framework for R. Bioinformatics. 2008 Aug 1;24(15):1733-4.

[28]. Ihaka R, Gentleman R. R: a language for data analysis and graphics. Journal of

computational and graphical statistics. 1996 Sep 1;5(3):299-314.

[29].   Xiao N, Cao DS, Zhu MF, Xu QS. protr/ProtrWeb: R package and web server

for generating various numerical representation schemes of protein sequences.

Bioinformatics. 2015 Jun 1;31(11):1857-9.

[30].   Pal M. Random forest classifier for remote sensing classification. International

journal of remote sensing. 2005 Jan 1;26(1):217-22.

[31].   Pregibon D. Logistic regression diagnostics. The annals of statistics. 1981

Jul;9(4):705-24.

[32].   Rätsch G, Onoda T, Müller KR. Soft margins for AdaBoost. Machine learning.

2001 Mar;42(3):287-320.

[33].   Soria D, Garibaldi JM, Ambrogi F, Biganzoli EM, Ellis IO. A

'non-parametric'version of the naive Bayes classifier. Knowledge-Based Systems.

2011 Aug 1;24(6):775-84.

[34].   Oliphant TE. Python for scientific computing. Computing in science &

engineering. 2007 Jun 18;9(3):10-20.

[35].   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O,

Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J. Scikit-learn:

Machine learning in Python. the Journal of machine Learning research. 2011 Nov

1;12:2825-30.

[36].   McKinney W. pandas: a foundational Python library for data analysis and

statistics. Python for high performance and scientific computing. 2011 Nov

18;14(9):1-9.

[37].   Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau

D, Wieser E, Taylor J, Berg S, Smith NJ, Kern R. Array programming with NumPy.

Nature. 2020 Sep;585(7825):357-62.

[38].   Van Der Walt S, Colbert SC, Varoquaux G. The NumPy array: a structure for

efficient numerical computation. Computing in science & engineering. 2011 Mar

7;13(2):22-30.

[39].   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D,

Burovski E, Peterson P, Weckesser W, Bright J, Van Der Walt SJ. SciPy 1.0:

fundamental algorithms for scientific computing in Python. Nature methods. 2020

Mar;17(3):261-72.

[40].   Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B,

Gautier L, Ge Y, Gentry J, Hornik K. Bioconductor: open software development for

computational biology and bioinformatics. Genome biology. 2004 Sep;5(10):1-6.

[41].   Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W.

BioMart and Bioconductor: a powerful link between biological databases and

microarray data analysis. Bioinformatics. 2005 Aug 15;21(16):3439-40.

[42].   Smedley D, Haider S, Ballester B, Holland R, London D, Thorisson G,

Kasprzyk A. BioMart–biological queries made easy. BMC genomics. 2009

Dec;10(1):1-2.

[43].   Psomiadou V, Iavazzo C, Douligeris A, Fotiou A, Prodromidou A, Blontzos N,

Karavioti E, Vorgias G. An Alternative Treatment for Vaginal Cuff Wart: a Case

Report. Acta Medica (Hradec Kralove). 2020;63(1):49-51. doi:

10.14712/18059694.2020.15.