1

Title

Quantitative live-cell imaging of GPCR downstream signaling dynamics.

Authors

Ryosuke Tany[1,2,3], Yuhei Goto[1,2,3,*], Yohei Kondo[1,2,3], and Kazuhiro Aoki[1,2,3,4,*]

Affiliations

[1]Quantitative Biology Research Group, Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

[2]Division of Quantitative Biology, National Institute for Basic Biology, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

[3]Department of Basic Biology, School of Life Science, SOKENDAI (The Graduate University for Advanced Studies), 5-1 Higashiyama, Myodaiji-cho, Okazaki, Aichi 444-8787, Japan.

[4]Lead Contact

*Correspondence: Yuhei Goto (y-goto@nibb.ac.jp), and Kazuhiro Aoki (k-aoki@nibb.ac.jp)

Keywords

GPCR, signaling dynamics, fluorescence imaging, dopamine, serotonin

Abstract

G-protein-coupled receptors (GPCRs) play an important role in sensing various extracellular stimuli, such as neurotransmitters, hormones, and tastants, and transducing the input information into the cell. While the human genome encodes more than 800 GPCR genes, only four Gα-proteins (Gα$_s$, Gα$_{i/o}$, Gα$_{q/11}$, and Gα$_{12/13}$) are known to couple with GPCRs. It remains unclear how such divergent GPCR information is translated into the downstream G-protein signaling dynamics. To answer this question, we report a live-cell fluorescence imaging system for monitoring GPCR downstream signaling dynamics at the single-cell level. Genetically encoded biosensors for cAMP, Ca^{2+}, RhoA, and ERK were selected as markers for GPCR downstream signaling, and were stably expressed in HeLa cells. GPCR was further transiently overexpressed in the cells. As a proof-of-concept, we visualized GPCR signaling dynamics of 5 dopamine receptors and 12 serotonin receptors, and found heterogeneity between GPCRs and between cells. Even when the same Gα proteins were known to be coupled, the patterns of dynamics in GPCR downstream signaling, including the signal strength and duration, were substantially distinct among GPCRs. These results suggest the importance of dynamical encoding in GPCR signaling.

INTRODUCTION

G-protein-coupled receptors (GPCRs), the largest family of membrane proteins, are triggered by various types of ligands, such as neurotransmitters (e.g., dopamine, serotonin), peptide hormones (e.g., angiotensin II), tastants, and light, to name a few. More than 800 GPCR genes are encoded in the human genome, and play indispensable roles in a vast variety of biological processes [1]. Moreover, GPCRs are one of the most important targets for drug discovery, as demonstrated by the fact that GPCRs constitute targets of approximately one-third of all Food and Drug Administration (FDA)-approved drugs [2].

      Upon ligand stimulation, GPCRs activate heterotrimeric G proteins composed of alpha, beta, and gamma subunits (Gα, Gβ, and Gγ) [3]. The activated GPCRs promote guanine nucleotide exchange of Gα proteins from a GDP-bound inactive state to a GTP-bound active state [4]. The different types of GTP-loaded Gα proteins bind to and activate/inactivate different downstream effectors (**Figure 1A**). Gα$_s$ positively regulates and Gα$_{i/o}$ negatively regulates adenylate cyclases, which are enzymes that catalyze cyclic AMP (cAMP) production [5,6]. Gα$_{q/11}$ induces intracellular Ca^{2+} increase through

phospholipase C beta (PLCβ), while $G\alpha_{12/13}$ activates the Rho family of small GTPases [7–9]. It has

also been reported that $G\alpha_s$, $G\alpha_{i/o}$, and $G\alpha_{q/11}$ regulate the ERK MAP kinase pathway [10–13]. In

addition to Gα proteins, accumulating evidence supports the independent roles of the Gβγ complex in

activating Rho family GTPases [14]. Other well-known mediators of GPCR signaling, including

regulators of G protein signaling (RGSs), G-protein-coupled receptor kinases (GRKs), and β-arrestin,

have also been shown to modulate the ERK MAP kinase pathway [10].

Much effort has been devoted to the development of imaging systems for quantifying GPCR

activation upon ligand stimulation. The imaging systems reported so far can be roughly classified into

two types: systems that utilize the principle of ligand-induced association/dissociation between GPCRs,

effectors, and heterotrimeric G proteins, and systems that employ downstream signaling molecules of

GPCRs as a marker. In the former imaging methods, protein-protein interactions such as GPCR-G

proteins, GPCR-β-arrestin, and Gα-Gβγ are detected by techniques such as Förster (or fluorescence)

resonance energy transfer (FRET) and bioluminescence resonance energy transfer (BRET) [15–17].

Other methods for measuring these protein-protein interactions by converting them into gene

expression or by cleavage of membrane molecules by metalloproteinases such as ADAM17/TACE

have been proposed to enable the large-scale GPCR screening [18,19]. These techniques directly

evaluate ligand-induced GPCR activation, but they do not allow estimation of the extent to which the

GPCR activation evokes downstream signaling. The latter type of imaging methods detect downstream

cell signaling induced by GPCRs such as cAMP and Ca^{2+} by small chemical compounds and

genetically encoded biosensors [20–23]. However, a drawback of this method is that it is challenging to

visualize two or more types of signal transduction systems in an efficient manner, and the throughput is

practically limited.

Many patterns of GPCR-Gα protein coupling have been investigated, revealing that some

GPCRs can promiscuously couple to multiple G proteins [19,24]. Yet, in most cases, only a limited

amount of quantitative information is available. Thus, it is still unknown to what extent the GPCRs

activate Gα proteins and what types of downstream signaling dynamics an activated GPCR generates.

This study reports a quantitative live-cell imaging system for quantifying cAMP, Ca^{2+}, RhoA, and ERK

activation dynamics induced by GPCR activation at the single-cell level. For this purpose, the FRET-

based biosensors for cAMP and RhoA activity, the red fluorescent Ca^{2+} biosensor, and the infra-red

fluorescent ERK biosensors were selected. We first established two types of HeLa cell lines stably

expressing cAMP/Ca^{2+} biosensors and RhoA/ERK biosensors. These two cell lines were then co-

cultured, transfected with GPCR genes, and stimulated with the ligand to monitor the changes in cAMP, Ca^{2+}, RhoA, and ERK activity. Taking advantage of this method, we quantified the GPCR signaling dynamics of 5 dopamine receptors and 12 serotonin receptors, demonstrating a substantial difference even between GPCRs that bind to the same Gα proteins and cellular heterogeneity of downstream signals.

Experimental

Plasmids

The cAMP biosensor (CFP-Epac-YFP) was developed based on the previous work [20], and it contained monomeric teal fluorescent protein (mTFP), the human RAPGEF3 (EPAC) gene (corresponding to amino acids 149-881) cloned from HeLa cells by RT-PCR, and mVenus. The cDNA of the cAMP biosensor was inserted into the pCX4neo vector [25], a retroviral vector with IRES-*neo* (the G418-resistance gene), providing pCX4neo-CFP-Epac-YFP. The cDNA of the red fluorescent genetically encoded Ca^{2+} indicators for optical imaging (R-GECO1.0) [26] was subcloned into the pPBbsr vector [27,28], a PiggyBac transposon vector with IRES-*bsr* (the blasticidin S-resistance gene), generating pPBbsr-R-GECO1.0. The dimerization-optimized reporter for activation (DORA) RhoA biosensor (DORA-RhoA) was used to visualize RhoA activity, and the DORA-RhoA was a gift from Dr. Yi Wu (University of Connecticut Health Center, CT) [22]; DORA-RhoA consists of the Rho-binding domain of PKN1, circularly permuted Venus (cpVenus), a linker, Cerulean3, and RhoA. The cDNA of DORA-RhoA was subcloned into pT2Absr [29], which was generated by inserting IRES-*bsr* into the pT2AL200R175 vector, generating pT2Absr-DORA-RhoA. The cDNA of the ERK activity biosensor, ERK-KTR [23], was fused with the cDNA of monomeric near-infrared fluorescent protein (miRFP703)[30], and subcloned into the pCSIIbleo vector, a lentivirus vector with IRES-*bleo* (the zeocin-resistance gene) [31], generating pCSIIbleo-ERK-KTR-miRFP703. The plasmid encoding the nuclear marker, Linker Histone-H1 (H1)-mCherry, was subcloned into a pCSIIneo vector [32], generating pCSIIneo-H1-mCherry. The GPCR plasmids were derived from the PRESTO-Tango kit originated from Dr. Bryan Roth (Addgene kit # 1000000068) [18]. pCMV-VSV-G-RSV-Rev was a gift of Dr. Miyoshi (RIKEN, Japan) [33], pGP and pCSIIpuro-MCS were a gift from Dr. Matsuda (Kyoto University, Japan) [34,35], psPAX2 was a gift from Dr. Trono (Addgene plasmid #12260) [36], pCAGGS-T2TP was a gift from Dr. Kawakami (National Institute for Genetics, Japan) [29], and pCMV-mPBase (neo-) was a gift from Dr. Bradley (Wellcome Trust Sanger Institute, UK) [27].

Reagents

Epidermal growth factor (EGF) was purchased from Sigma-Aldrich (St. Louis, MO). Blasticidin S, zeocin, G418, and puromycin were obtained from InvivoGen (Carlsbad, CA). Forskolin (FSK) and 3-isobutyl-1-methyl-xanthine (IBMX) were purchased from Wako (Osaka, Japan). Dopamine was obtained from Sigma-Aldrich (H8602) and solubilized into 10 mM HCl as a 1 M stock solution. Serotonin was purchased from Cayman (14332) and solubilized into 10 mM HCl as a 50 mM stock solution. Adenosine 5'-triphosphate disodium salt hydrate (ATP) was obtained from TCI (A0157) and solubilized into DDW as a 100 mM stock solution. Isoproterenol (ISO) was obtained from Sigma-Aldrich (I6504) and solubilized into DDW as a 50 mM stock solution.

Cell culture

HeLa cells and HEK-293T cells were gifted from Dr. Matsuda (Kyoto University, Japan), and cultured in Dulbecco's Modified Eagle's Medium (DMEM) high glucose (Wako; nacalai tesque) supplemented with 10% fetal bovine serum (Sigma-Aldrich) at 37°C in 5% CO_2. For the live-cell imaging, HeLa cells were plated on CELLview cell culture dishes (glass bottom, 35 mm diameter, 4 components: The Greiner Bio-One) one day before transfection. Transfection was performed with a mixture containing 230 ng plasmid encoding TANGO-GPCR, 20 ng plasmid carrying the puromycin-resistance gene (pCSIIpuro-MCS), and 0.25 uL of 293fectin transfection reagent (Thermo Fisher Scientific) in each well. In the dopamine receptor co-expression experiment in Figure 4, TANGO GPCR plasmids were used in a 1: 1 mixture. One day after the transfection, puromycin (final concentration 2 μg/ml) was administered, and transfected cells were selected for 2 days. After the selection of transfected cells, the medium was replaced with starvation medium (FluoroBrite (nacalai tesque)/1x GlutaMAX (GIBCO)/0.1% BSA) 18 ~ 25 h before the imaging was started.

Stable cell line construction

To establish a stable cell line expressing R-GECO1.0 and CFP-Epac-YFP (HeLa/cAMP/Ca^{2+}), the PiggyBac transposon system and the retroviral vector system were used, respectively. The pPBbsr-R-GECO1.0 was co-transfected with pCMV-mPBase (neo-) encoding the PiggyBac transposase by using 293fectin. One day after the transfection, cells were selected with 20 μg/ml blasticidin S for at least one week. For retroviral production, the pCX4neo-CFP-Epac-YFP was transfected into HEK-293T cells together with pGP and pCSV-VSV-G-RSV-Rev by using Polyethylenimine "MAX" MW 40,000 (Polyscience Inc., Warrington, PA). Virus-containing media were collected at 48 h after the

transfection, filtered, and used to infect the cell lines expressing R-GECO1.0 with 10 μg/mL polybrene.

After the infection, cells were selected with 20 μg/ml blasticidin S and 1 mg/mL G418 for at least one

week, followed by single-cell cloning. Clonal cells were used in the experiments Figure 2 and Figure 4.

To establish a stable cell line expressing ERK-KTR-miRFP703, DORA-RhoA, and H1-

mCherry (HeLa/RhoA/ERK), a lentiviral vector system (for ERK-KTR-miRFP703 and H1-mCherry)

and a Tol2 transposon-based system (for DORA-RhoA) were applied, respectively. For lentiviral

production, HEK-293T cells were co-transfected with the pCSIIbleo-ERK-KTR-miRFP703, psPAX2,

and pCSV-VSV-G-RSV-Rev by using Polyethylenimine "MAX" MW 40,000. Virus-containing media

were collected at 48 h after the transfection, filtered, and used to infect HeLa cells with 10 μg/mL

polybrene. After the infection, cells were selected with 100 μg/ml zeocin for at least one week. Next,

the pT2Absr-DORA-RhoA was co-transfected into HeLa cells stably expressing ERK-KTR-miRFP703

together with the pCAGGS-T2TP plasmid encoding transposase by using 293fectin. One day after the

transfection, cells were selected by at least one week of treatment with 100 μg/ml zeocin and 20 μg/ml

blasticidin S. The pCSIIneo-H1-mCherry was introduced into this cell line by using a lentiviral viral

vector system as well as ERK-KTR-miRFP703. One day after the infection, cells were selected by at

least one week of treatment with 100 μg/ml zeocin, 20 μg/ml blasticidin S, and 1 mg/mL G418,

followed by single-cell cloning.

Single-cell clones were used in the experiments in Figure 2, Supplementary Figures S2, S3, and

S5. In other experiments, bulk cells were used.

Time-lapse imaging

Images were acquired on an IX81 inverted microscope (Olympus) equipped with a Retiga 4000R

cooled Mono CCD camera (QImaging), a Spectra-X light engine illumination system (Lumencor), an

IX2-ZDC laser-based autofocusing system (Olympus), a UAPO/340 40x/1.35 oil iris objective lens

(Olympus), a MAC5000 controller for filter wheels and XY stage (Ludl Electronic Products), an

incubation chamber (Tokai Hit), and a GM-4000 CO_2 supplier (Tokai Hit). The following filters and

dichroic mirrors were used: for R-GECO1.0 and mCherry, an FF01-580/20 excitation filter (Semrock),

a 20/80 beamsplitter dichroic mirror (Chroma), and an FF01-641/75 emission filter (Semrock); for

iRFP, an FF01-632/22 excitation filter (Semrock), an FF408/504/581/667/762-Di01 dichroic mirror

(Semrock), and an FF01-692/LP emission filter (Semrock); for FRET, an FF01-438/24 excitation filter

(Semrock), an XF2034 455DRLP dichroic mirror (Omega Optical), an FF01-542/27 emission filter

(Semrock); for CFP, an FF01-438/24 excitation filter (Semrock), an XF2034 455DRLP dichroic mirror

(Omega Optical), and an FF01-483/32 emission filter (Semrock). The microscopes were controlled by

MetaMorph software (Molecular Devices). For live-cell imaging, cells were stimulated with ligands

(EGF, FSK+IBMX, ATP, dopamine, or serotonin) 5 min after the start of imaging, and followed by 25

min imaging with every 1 min interval. To measure the $G\alpha_{i/o}$ activity, cells were first treated with 100

nM ISO 5 min after the start of imaging to increase the intracellular cAMP level. Next, cells were

subsequently treated with vehicle or ligands (dopamine or serotonin) 10 min after the ISO treatment,

followed by 15 min imaging.

Immunofluorescence

Cells were fixed with the final 3.7% formaldehyde for 30 min in a culture medium. After washing with

PBS four times, cells were permeabilized with 0.2% TritonX-100 in PBS for 15 min for the

immunofluorescence of total GPCR expression, followed by soaked with 3% BSA and 0.02% TritonX-

100 in PBS (buffer 1). For the immunofluorescence of GPCRs expressing cell-surface, cells were

soaked with buffer 1 without permeabilization. Then, the cells were incubated with primary antibodies,

monoclonal anti-FLAG M2 antibody (1:500 dilution; Sigma-Aldrich, F1804) diluted in buffer 1 for

overnight at 4°C. Next, the cells were washed three times with PBS, and then incubated for 1 h at room

temperature with Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody Alexa

Fluor Plus 555 (1:1000 dilution; Invitrogen, A32727) in buffer 1. Finally, the cells were washed three

times with PBS and subjected to fluorescence imaging.

Image and data analysis

Fiji, a distribution of ImageJ [37], was used for image processing and analysis. For all images,

background signals were subtracted by the rolling ball method, and then the images were registered by

StackReg, a Fiji plugin to correct misregistration. Note that the median filter was used for the time-

lapse images before registration to remove camera noise to prevent the registration error. For FRET

images of CFP-Epac-YFP or DORA-RhoA, the ratio images of the CFP/FRET or FRET/CFP ratio

were generated, respectively, and then ratio images were visualized by the intensity-modulated display

(IMD) mode, where eight colors from red to blue are used to represent the CFP/FRET ratio.

To measure the intracellular cAMP and Ca^{2+} levels, we manually set a circular region of interest

(ROI) with a diameter of 20 pixels at the center of each cell. Note that we avoid selecting the cells that

overlapped with dying cells during the time course. To measure the ERK activity from the translocation

of ERK-KTR, we used CellTK [38]. In brief, pre-processing, segmentation, tracking, and post-

processing was performed using functions in CellTK. In pre-processing, nuclear images were processed

by a gaussian blur filter. In the segmentation process, an adaptive threshold algorithm was basically

applied except for some difficult images, which were segmented by Stardist [39,40]. Segmented

nuclear labels were tracked by the Linear Assignment Problem (LAP) algorithm [41], followed by a

track_neck_cut function in CellTK. As for post-processing, short tracking data were discarded, and the

cytoplasmic region was defined as a dilating ring around nuclear segmentation. Quantified feature

values (i.e., mean intensity, x-y positions, etc.) of each cell were exported and cleaned in Python.

Invalid cell tracks were discarded according to the following criteria: a threshold of CFP signal

intensity to remove non-fluorescent cells and dead cells; a threshold of coefficient of variation (CV) of

the CFP signal intensity along time course to remove the cells crossed by floating dead cells or debris

during time course; DORA-RhoA cells adjacent to CFP-Epac-YFP cells were removed because the

much stronger fluorescence from CFP-Epac-YFP cells affects the quantification of FRET signals of

DORA-RhoA. The quantified time traces were further normalized by the average values before

stimulation (0-5 min), except in Figure 6E and Figure 7F.

To classify signaling dynamics, K-means-based time-series clustering was used. Euclidean

distance was used as a measure of proximity. At first, all quantified time-course data of each reporter

were normalized as described above, and divided into three groups by the K-means algorithm using

scikit-learn ver. 1.0.2 and tslearn ver. 0.5.2 [42,43] modules for Python. At the next step, the fractions

of clusters were calculated in each receptor. Note that the number of resulting clusters was

conservatively determined as three for all reporters because the cluster number prediction methods,

such as the Elbow Method, did not provide reliable prediction in our case.

RESULTS

Development of live-cell fluorescence imaging for the quantification of GPCR signaling.

To cover a wide range of GPCR signals mediated by individual Gα subtypes, we chose cAMP, Ca^{2+},

RhoA, and ERK as markers (**Figure 1A**). We screened genetically encoded biosensors for these

signaling molecules, and finally selected the following biosensors: CFP-Epac-YFP (a Förster resonance

energy transfer (FRET)-based cAMP biosensor) [20], R-GECO1.0 (a red fluorescent Ca^{2+} biosensor)

[21], DORA-RhoA (a FRET-based RhoA biosensor) [22], and ERK-KTR fused with miRFP703 (an

infra-red fluorescent ERK biosensor) [23] (**Figure 1A**). For efficient imaging, two types of stable HeLa

cell lines were established: a cell line expressing the cAMP biosensor and R-GECO1.0

(HeLa/cAMP/Ca^{2+}), and a cell line expressing DORA-RhoA and ERK-KTR (HeLa/RhoA/ERK)

(**Figure 1B**) together with linker histone H1-mCherry as a nuclear marker to quantify the

cytoplasm/nucleus (C/N) ratio automatically. The two cell lines were co-cultured and transfected with

the plasmids for GPCR expression and puromycin-resistance gene expression. One day after

transfection, the cells were treated with puromycin to select GPCR-expressing cells. After the selection,

the cells were further serum-starved for 18~25 h before imaging to reduce the ERK activity at a basal

level. The plasmids from the PRESTO-Tango GPCR kit were used for the expression of each GPCR,

and therefore the exogenous GPCR protein was fused with the HA signal peptide and FLAG-tag at the

N-terminus and V2 tail, the TEV cleaved site, and the rtTA moieties at the C-terminus [18]. We suspect

that these parts have no considerable effects on GPCR signaling.

265
266

267
Figure 1. Development of the live-cell fluorescence imaging system for quantifying the GPCR
signaling dynamics.
(A) (Upper panel) Schematic model of Gα coupling and typical GPCR-mediated intracellular signaling
pathways. (Lower panel) The mode of action of the indicated biosensors is shown.
(B) Experimental procedure of the live-cell fluorescence imaging.

Validation of cell lines for monitoring GPCR signaling.

As a validation of the cell lines mentioned above, we stimulated the cells with chemical compounds,

ligands, and epidermal growth factor (EGF), which are well-known to upregulate cAMP, Ca^{2+}, RhoA,

and ERK. HeLa/cAMP/Ca^{2+} cells were stimulated with Forskolin and IBMX (**Figure 2A**) and ATP

(**Figure 2B**). The cells showed a sustained increase in the CFP/FRET ratio of the CFP-Epac-YFP

biosensor and pulsatile increase in red fluorescence of R-GECO1.0, indicating a sustained cAMP

increase and pulsatile Ca^{2+} increase, respectively. Similarly, HeLa/RhoA/ERK cells were stimulated

with EGF, resulting in a transient rise in the FRET/CFP ratio of DORA-RhoA and gradual

translocation of ERK-KTR-miRFP703 from the nucleus to the cytoplasm (**Figure 2C, 2D**). The data

indicate that EGF induced transient RhoA and gradual ERK activation. We represent ERK activity as

the cytoplasm/nucleus (C/N) ratio of ERK-KTR-miRFP703 fluorescence intensity [23,30]. To assess

the sensitivity and dynamic range of biosensors, we evaluated the dose-response of each biosensor to

the ligands (**Figure 2E-2H**). Further, no clear correlation was observed between the expression level of

the biosensors and the response to the ligands in the cell lines we used in this study (**Supplementary**

Figure S1A-D). It is noted that our microscope setting did not show any spectral cross-talk between

CFP-Epac-YFP and R-GECO1.0 or between DORA-RhoA and ERK-KTR-miRFP703. Taken together,

these results demonstrated that the cell lines expressing biosensors allowed monitoring of cAMP, Ca^{2+},

RhoA, and ERK activity at the single-cell level.

Figure 2. Validation of cell lines for monitoring GPCR signaling.
(A-D) (Left panels) HeLa/cAMP/Ca^{2+} cells (A, B) and HeLa/RhoA/ERK cells (C, D) were treated with
50 μM Forskolin and 100 μM IBMX (A), 100 μM ATP (B), and 10 ng/mL EGF (C and D).
Representative CFP/FRET ratio images (A) and FRET/CFP ratio images (C) are shown in the IMD
mode, where eight colors from red to blue are used to represent the CFP/FRET ratio (A) or FRET/CFP
ratio (C) with the intensity of each color indicating the denominator channels. (Right panels) The
response of the biosensors for cAMP levels, Ca^{2+} levels, ERK activity, and RhoA activity is normalized
by dividing by the averaged value before stimulation and plotted as a function of elapsed time after
stimulation. The red and gray lines represent the time-course for the average and individual cells,
respectively. N = 32, 45, 13, 16 cells, respectively. (E-H) The dose response curve of cAMP (E), Ca^{2+}
(F), RhoA (G), and ERK (H) to the indicated ligands. The mean $\Delta F/F_0$ or $\Delta R/R_0$ values (dots) are
plotted as a function of ligand concentrations with the fitted curve (gray line) and parameters for EC50
and Hill coefficient (nH). The experiments were repeated at least two times. Scale bar, 10 μm.

Signaling dynamics of dopamine receptors.

A neuromodulator, dopamine, controls many physiological functions, including locomotion, reward,

cognitive function, and learning [44]. Because dopamine is involved in a wide range of cellular and

behavioral processes, multiple human disorders have been linked to dopaminergic dysfunctions such as

Parkinson's disease, schizophrenia, ADHD, and Tourette's syndrome [45]. The five subtypes of

dopamine receptors, D1, D2, D3, D4, and D5 receptors, are encoded by the genes *DRD1*, *DRD2*,

DRD3, *DRD4*, and *DRD5*, respectively, in the human genome, and collectively these subtypes are

known to mediate all physiological functions of dopamine [45,46]. The dopamine receptors are GPCRs

and are divided into two groups: the D1-class dopamine receptors (D1 and D5) and the D2-class

dopamine receptors (D2, D3, and D4) [47]. The D1-class dopamine receptors are generally coupled to

$G\alpha_s$, while the D2-class dopamine receptors are coupled to $G\alpha_{i/o}$ (**Figure 3A**). Thus, the D1 and D2

receptors positively and negatively regulate adenylate cyclases (ACs), which catalyze the production of

cAMP, respectively. Furthermore, it has been reported that D1, D5, and a putative D1-D2 heterodimer

couple and activate $G\alpha_{q/11}$ and Ca^{2+} signaling [48–51]. However, this idea remains controversial

because D1and D2 are not coexpressed in most striatal neurons in mice [45], and further studies will be

needed to quantitatively investigate to extent to which these dopamine receptors contribute to the $G\alpha_{q/11}$

activation and Ca^{2+} increase.

First, the surface expression of D1-5 receptors was confirmed by immunofluorescence assay

(**Figure 3B and 3C**). DRD1 was mainly localized at the cell surface, while DRD3 and DRD4 were

partially expressed in the cell surface and the rest was localized intracellularly (**Figure 3B**). Cell

surface expression of DRD2 and DRD5 were weaker, but detectable, than the others, and most of these

receptors retained in the intracellular regions (**Figure 3B**). We quantified the surface expression levels

of DRDs by correcting cell-to-cell heterogeneity of transfection efficiency, showing the distinct

differences of surface expression levels among the DRDs (**Figure 3C**). Next, we quantified the cAMP,

Ca^{2+}, RhoA, and ERK activation dynamics evoked by dopamine receptors with the live-cell

fluorescence imaging system (**Figure 1B**). In HeLa cells lacking expression of dopamine receptors,

dopamine stimulation did not elicit any change in cAMP, Ca^{2+}, RhoA, or ERK activity (**Figure 3D**, left

column). This could be simply because HeLa cells do not express any dopamine receptors. As

expected, the expression of D1-type receptors, i.e., DRD1 and DRD5, increased cAMP levels upon

dopamine stimulation, while the expression of D2-type receptors, i.e., DRD2, DRD3, and DRD4, did

not change any cAMP levels (**Figure 3D**, first row). This was probably because the basal cAMP level

was low, and thus the cAMP biosensor could not detect the decrease in cAMP levels by D2-type

receptors. Weak, measurable Ca^{2+} pulses were observed in HeLa cells expressing DRD1 and DRD2,

suggesting that DRD1 and DRD2 receptors are weakly coupled to $G\alpha_{q/11}$ receptors (**Figure 3D**, second

row). RhoA was also activated in DRD1 or DRD5 expressing HeLa cells for an unknown reason

(**Figure 3D**, third row). ERK was transiently decreased in HeLa cells expressing DRD1 or DRD5, but

it is challenging to evaluate dopamine-induced ERK signaling due to significant cell heterogeneity

(**Figure 3D**, fourth row). Furthermore, to confirm the activation of $G\alpha_{i/o}$-coupled receptors, cells

expressing DRD2, DRD3, or DRD4 were pretreated with 100 nM isoproterenol to upregulate cAMP

production through endogenous β adrenoceptor activation, followed by dopamine stimulation. As we

expected, the expression of DRD2, 3, or 4 increased the proportion of cells that reduced cAMP levels

after dopamine stimulation (**Supplementary Figure S2A**), indicating the dopamine-induced activation

of $G\alpha_{i/o}$ signaling through DRD2, 3, and 4.

Figure 3. Signaling dynamics induced by each dopamine receptor.
(A) Schematic illustration of dopamine, dopamine receptors (DRDs) coupling to Gα, and downstream signaling. (B) Expression of DRDs at the cell surface and whole cell. HeLa cells were transfected with the plasmids expressing the indicated receptors, followed by immunofluorescence (see methods). The values in the lower right indicate the lower and upper limits of the display range. Scale bar, 20 μm. (C) The quantification of the surface expression levels of DRDs. To correct for cell-to-cell heterogeneity of transfection efficiency, the fluorescence intensity of cotransfected EGFP was used to normalize the fluorescence intensity of cell surface receptors in each cell. The values for each cell were plotted as a scatter plot with a box-whisker plot, in which the box shows the quartiles of data with the whiskers denoting the minimum and maximum except for the outliers detected by 1.5 times the interquartile range (n = 46, 26, 41, 50, 27 cells). (D) Normalized responses of each reporter in HeLa cells expressing the indicated dopamine receptors are shown. The response of the biosensors for cAMP levels, Ca^{2+} levels, ERK activity, and RhoA activity is normalized by dividing by the averaged value before stimulation and plotted as a function of elapsed time after administration of 10 μM dopamine. The red and gray lines represent the time-course for the average and individual cells (n > 10 cells), respectively.

Next, we examined whether heterodimerized dopamine receptors synergistically increase intracellular Ca^{2+} levels upon dopamine stimulation, because it has been reported that heterodimers of dopamine receptors such as the D1-D2 dimer activate $Gα_{q/11}$ signaling [48–50]. HeLa cells were co-transfected with ten possible combinations of dopamine receptors and stimulated with dopamine. Unexpectedly, we found no enhancement of the dopamine-induced Ca^{2+} increase by co-expression of dopamine receptors, but rather the co-expression inhibited the pulsatile Ca^{2+} increase observed in HeLa cells expressing DRD1 or DRD5 (**Supplementary Figure S3**).

Signaling dynamics of serotonin receptors.
Serotonin (5-hydroxytryptamine, 5-HT) plays essential roles in a wide range of physiological phenomena, including sensory perception and behaviors [52]. The serotonin receptors are the most prominent among the GPCR families for neurotransmitters; 13 distinct genes encode GPCRs for the serotonin, and there is one ligand-gated ion channel, the 5-HT3 receptor [53]. The G-protein coupled serotonin receptors are grouped into three families depending on which Gα proteins they primarily couple: the $Gα_{q/11}$-coupled receptors (HTR2A, HTR2B, and HTR2C), $Gα_{i/o}$-coupled receptors (HTR1A, HTR1B, HTR1D, HTR1E, HTR1F, HTR5), and $Gα_s$-coupled receptors (HTR4, HTR6, and HTR7) (**Figure 5A**). Although these receptors are divided based primarily on their coupling with Gα proteins, recent works have shown that GPCRs can couple to more than one type of Gα proteins [19]. In addition, the constitutive activity of serotonin receptors has been reported, though the basal- and ligand-induced activities of serotonin receptors have not yet been thoroughly evaluated.

We therefore analyzed the G-protein-coupled serotonin receptor-induced cAMP, Ca^{2+}, RhoA,

and ERK activation with the live-cell fluorescence imaging system (**Figure 1B**). First, $G\alpha_{q/11}$-coupled

serotonin receptors of HTR2A, HTR2B, and HTR2C, were investigated as did in DRDs. The surface

and total expression of HTR2A and HTR2B were detectable, while those of HTR2C were quite low

(**Figure 4B and 4C**). Parental HeLa cells did not show any response of these signaling molecules to

serotonin stimulation (**Figure 4D**, left column). The expression of $G\alpha_{q/11}$-coupled HTR2A and HTR2C

induced a sustained or pulsatile Ca^{2+} increase upon 5-HT stimulation, while slight or no response of

Ca^{2+} was observed in HTR2B-expressing cells (**Figure 4D**, second row). 5-HT stimulation slightly

activated RhoA in HTR2A- or HTR2B-expressing cells (**Figure 4D**, third row). Some cells showed a

substantial change in ERK activity, but on average, many cells did not show a shift in ERK activity

(**Figure 4D**, fourth row).

Figure 4. Signaling dynamics of $G\alpha_{q/11}$-coupled serotonin receptors.
(A) Schematic illustration of serotonin receptors coupling to $G\alpha$ and downstream signaling.
(B) Expression of $G\alpha_{q/11}$-coupled serotonin receptors at the cell surface and whole cell. The data were
obtained and are represented as in Figure 3B. Scale bar, 20 μm. (C) The quantification of the surface
expression levels of $G\alpha_q$-coupled serotonin receptors. The data were quantified and are represented as
in Figure 3C. (n = 50, 21, 32 cells). (D) Normalized response of each reporter in HeLa cells
overexpressing $G\alpha_{q/11}$-coupled serotonin receptors was plotted as a function of time after 10 μM 5-HT
administration. The response of the biosensors for cAMP levels, Ca^{2+} levels, ERK activity, and RhoA
activity was normalized by dividing by the averaged value before stimulation and plotted as a function
of elapsed time after stimulation. The red and gray lines represent the time-course for the average and
individual cells (n > 10 cells), respectively.

We next analyzed the signaling dynamics of $G\alpha_{i/o}$-coupled serotonin receptors, i.e., HTR1A,

HTR1B, HTR1D, HTR1E, HTR1F, and HTR5. The surface and total expression of these receptors

were confirmed by immunofluorescence (**Figure 5A and 5B**). As in D2-type dopamine receptors, the

expression of these receptors did not alter the cAMP levels with or without 5-HT (**Figure 5C**, first

row). Some of the receptors induced an increase in the Ca^{2+} level (HTR1A, 1B, 1D)(**Figure 5C**, second

row) and RhoA activity (HTR1B)(**Figure 5C**, third row), indicating the possibility of $G\alpha_{q/11}$-coupling

signaling in these receptors. We could not observe the characteristic dynamics of ERK activation with

high reproducibility (**Figure 5C**, fourth row). To confirm the activation of $G\alpha_{i/o}$, isoproterenol was

pretreated with the cells expressing the $G\alpha_{i/o}$-coupled serotonin receptors, followed by the treatment of

5-HT. To our surprise, we found that a part of the HeLa cell population responded to 5-HT without

overexpression of $G\alpha_{i/o}$-coupled serotonin receptors (**Supplementary Figure 2B**). This could be due to

the expression of HTR1D in HeLa cells [54]. The expression of the $G\alpha_{i/o}$-coupled serotonin receptors

further increased the population of responded cells to 5-HT except for HTR1E-expressing cells

(**Supplementary Figure 2B**).

Figure 5. Signaling dynamics of $G\alpha_{i/o}$-coupled serotonin receptors.

(A) Expression of $G\alpha_{i/o1}$-coupled serotonin receptors at the cell surface and whole cell. The data were obtained and are represented as in Figure 3B. Scale bar, 20 μm. (B) The quantification of the surface expression levels of $G\alpha_{i/o}$-coupled serotonin receptors. The data were quantified and are represented as in Figure 3C. (n = 50, 50, 41, 32, 28, 50 cells). (C) Normalized response of each reporter in HeLa cells overexpressing $G\alpha_{i/o}$-coupled serotonin receptors was plotted as a function of time after 10 μM 5-HT administration. The response of the biosensors for cAMP levels, Ca^{2+} levels, ERK activity, and RhoA activity was normalized by dividing by the averaged value before stimulation and plotted as a function of elapsed time after stimulation. The red and gray lines represent the time-course for the average and individual cells (n > 10 cells), respectively.

(E) FSK → Ligand

Constitutive activity of $G\alpha_s$-coupled serotonin receptors and their response to ligand

Finally, we analyzed $G\alpha_s$-coupled serotonin receptors, HTR4, HTR6, and HTR7. These receptors were expressed well at the cell surface and whole-cell (**Figure 6A and 6B**). HeLa cells expressing HTR4, HTR6, or HTR7 showed a sustained increase in cAMP level upon 5-HT stimulation with substantial

cell-to-cell heterogeneity (**Figure 6C**, first row). In most cells, Ca^{2+} levels did not vary by 5-HT

treatment (**Figure 6C**, second row), whereas RhoA activity slightly but reproducibly increased in cells

expressing HTR4, HTR6, or HTR7 (**Figure 6C**, third row). HTR4-expressing cells showed an increase

in ERK activity from the basal level on 5-HT stimulation, while did not in HTR6- or HTR7-expressing

cells (**Figure 6C**, fourth row).

We investigated the cause of the large cell-to-cell heterogeneity in the increase in cAMP, and

found that the basal cAMP levels varied between cells. HTR4 or HTR6 expression enhanced the basal

cAMP level in some cells despite the absence of 5-HT stimulation (**Figure 6D**, upper panels). The cells

with lower basal cAMP levels increased the cAMP upon 5-HT stimulation, whereas the cells with

higher basal cAMP levels did not (**Figure 6D**, lower panels). As a result, 5-HT stimulation caused

constant cAMP levels in all cells (**Figure 6D**, lower panels). Raw values of the CFP/YFP ratio of the

cAMP biosensor, which correlated with intracellular cAMP concentration, also demonstrated that 5-HT

stimulation kept cAMP levels constant (**Figure 6E**). These results suggest the constitutive activity of

HTR4 and HTR6 for $G\alpha_s$ signaling in the basal state and homeostatic mechanisms that control cAMP

levels in the activating state.

Figure 6. Constitutive activity of Gα$_s$-coupled serotonin receptors and their response to ligand.
(A) Expression of Gα$_s$-coupled serotonin receptors at the cell surface and whole cell. The data were
obtained and are represented as in Figure 3B. Scale bar, 20 μm. (B) The quantification of the surface
expression levels of Gα$_{i/o}$-coupled serotonin receptors. The data were quantified and are represented as
in Figure 3C. (n = 50, 50, 50 cells). (C) Normalized response of each reporter in HeLa cells
overexpressing Gα$_s$-coupled serotonin receptors was plotted as a function of time after 10 μM 5-HT
administration. Gray lines indicate single-cell responses (n > 10 cells), and the red line represents the
mean response. (D) Representative CFP/YFP ratio images of cAMP in HeLa cells overexpressing Gα$_s$-
coupled serotonin receptors before and after the stimulation of 10 μM 5-HT. CFP/FRET ratio images
are shown in intensity-modulated display mode as in Figure 2. Scale bar 10 μm. (E) Time course of

CFP/FRET ratio values without normalization. Gray lines indicate single cell (n > 10 cells) responses,
and the red line represents the mean response.
483

Systems analysis of signaling dynamics induced by activating dopamine or serotonin receptors.
To take full advantage, we analyzed the time-series data of GPCR signaling in more detail. We
performed hierarchical clustering with all data and demonstrated three clusters in the single-cell
imaging data of cAMP, Ca^{2+}, RhoA, and ERK (**Figure 7A-D**, left heatmaps and middle plots). Next,
we quantified the fraction of these three clusters in the signaling dynamics of each GPCR (**Figure 7A-**
7D, **Supplementary Figure S4A-D**). All $G\alpha_s$-coupled receptors including DRD1, DRD5, HTR4,
HTR6, and HTR7 demonstrated distinct cAMP dynamics in comparison to the other $G\alpha$-coupled
receptors, though the fraction of the clusters in the cAMP dynamics varied among these $G\alpha_s$-coupled
receptors (**Figure 7A**, **Supplementary Figure S4A**). Some cells showing characteristic Ca^{2+} dynamics
were found in DRD2, HTR1A, HTR1B, HTR2A, and HTR2C coupled to $G\alpha_{i/o}$ or $G\alpha_{q/11}$ (**Figure 7B**,
Supplementary Figure S4B). Many dopamine or serotonin receptors affected the activity of RhoA,
and among them, HTR2A, HTR4 and HTR7 demonstrated higher proportion of cells in cluster 3
showing strong activation of RhoA (**Figure 7C**, **Supplementary Figure S4C**). Interestingly, a fraction
of cells changing RhoA activity tended to be higher in activation of $G\alpha_s$-coupled GPCRs than the other
$G\alpha$-coupled GPCRs. Although in control cells the proportion of cluster 2 showing ERK inactivation
was observed, the distinct patterns of clustering in ERK activity dynamics were induced by the
activation of DRD1, DRD5, HTR4, HTR7, and HTR2A (**Figure 7D**, **Supplementary Figure S4D**).
These results suggest that GPCRs induce various downstream signaling dynamics, and that they
contain some parts that can be explained by the $G\alpha$ protein to which they are coupled and some that are
not.
505

Finally, we analyzed the interrelationships between GPCR downstream signaling dynamics. In
our imaging system, two of four GPCR signaling are simultaneously observed in one cell, providing an
opportunity to analyze crosstalk and feedback of two types of signal transduction at the single cell
level. In addition to HeLa/cAMP/Ca^{2+} and HeLa/RhoA/ERK cell lines, we further established a stable
cell line HeLa/cAMP/ERK expressing cAMP and ERK biosensors, thereby allowing to investigate the
relations in cAMP-Ca^{2+}, cAMP-ERK, and RhoA-ERK pairs. The dynamics of each signaling induced
by activation of DRD1, HTR4, or HTR7 were visualized by heatmaps with sorting basal cAMP level or
ERK activity (**Supplementary Figure S5**). Obviously, there is no correlation between cAMP and Ca^{2+}

dynamics (**Supplementary Figure S5A**), consistent with the result that the activation of $G\alpha_s$-coupled

GPCR did not induce any changes in Ca^{2+} levels (**Figure 7A** and **7B**). ERK activation was often

observed in HTR4 expressing cells with higher basal cAMP levels (**Supplementary Figure S5B**),

whereas ERK activation seemed not to be correlated with RhoA activation (**Supplementary Figure**

S5C). The findings provide a potential link between ERK and cAMP signaling, such as cross-talk

regulations.

524
525
Figure 7. Systems analysis of signaling dynamics induced by activating dopamine or serotonin
receptors.
(A-D) Left heatmaps represent the result of time-series *K*-means clustering analysis. Vertical axis of
heatmap represents each single cell and horizontal axis represents time (-5 min to 25 min since the
ligand addition). Middle plots are the average time-course of each cluster. The proportion of clusters in
each GPCR expressing cell are visualized in the right bar graphs. cAMP (A), Ca^{2+} (B), RhoA (C), and
ERK (D), respectively.
533
534
DISCUSSION

In this study, we demonstrate the effectiveness of the live-cell fluorescence imaging system to

simultaneously visualize multiple GPCR-induced downstream signal dynamics. Targeting dopamine

receptors and serotonin receptors, we found that GPCR downstream signaling dynamics are highly

variable. Although it is known which Gα protein the dopamine and serotonin receptors couple to, our

experimental results indicate that the strength and temporal patterns of the downstream signaling vary

among GPCRs. This is probably because the degree of Gα activation by GPCRs and/or inactivation of

GPCRs via GRKs or β-arrestins differs among GPCRs. The crosstalk of downstream signaling of

GPCRs may also affect their dynamics. In fact, the negative correlation between cAMP and ERK

activity was observed at the single-cell level (**Supplementary Figure S5B**). This could be explained

by the suppression of RAF by protein kinase A (PKA)[55,56]. In addition, the phosphodiesterases

(PDEs) are regulated by Ca^{2+}, PKA, and ERK [57,58], suggesting the potential contribution to the

dynamics of these signaling molecules.

The single-cell imaging data further clarify the striking cellular heterogeneity of GPCR

downstream signaling, showing diverse signaling dynamics such as responding vs. non-responding

cells, sustained vs. transient responses, and pulsatile responses. The observed diversity of dynamics

suggests that the GPCR signaling is enhanced by "dynamical encoding," i.e., utilization of time courses

of signaling activities as biological information. Despite dozens to hundreds of GPCRs being expressed

in individual cells [59], GPCRs transmit signals primarily through only four types of Gα proteins,

namely, $Gα_s$, $Gα_{i/o}$, $Gα_{q/11}$, and $Gα_{12/13}$. Thus, it is inevitable that multiple GPCRs in a single cell

activate the same type of Gα proteins, making it difficult for cells to interpret the input information.

The dynamical-encoding property has been found in many signaling components, including Ca^{2+} and

ERK [60–62]. Pioneering works have revealed rich signaling dynamics downstream of GPCRs [63].

However, due to the limited number of GPCRs characterized in terms of dynamics, we are still far from

unveiling the full diversity of GPCR signaling dynamics. For example, the cross-talk between the

dynamics of cAMP and Ca^{2+} is of critical importance in neuromodulator signaling [64]. We thus

considered that it would be of interest to apply our live-cell fluorescence imaging system to other

neuromodulator GPCRs.

One unexpected finding was that heterodimerization of dopamine receptors did not enhance the

intracellular Ca^{2+} increase induced by dopamine stimulation (**Supplementary Figure S2**). The most

probable reason for this disparity in results is that we used different cell lines from the previous study

(HeLa vs. HEK or neurons). A second possible reason is that the GPCR constructs we used in this

study contained an HA signal peptide and FLAG-tag fused to the N-terminus and a V2-tail, TEV

cleavage site, and rtTA fused to the C-terminus for the TANGO assay [18]. These moieties connected

to the GPCRs might prevent the dopamine receptors from forming heterodimers. The third possibility

is that the stoichiometry of the dopamine receptors was not appropriate. More careful analysis for

GPCR heterodimers will be required in the future. Another unexpected finding was the constitutive

activity of $G\alpha_s$-coupled serotonin receptors, HTR4 and HTR6 (**Figure 6**). Constitutive activity has been

reported in many types of GPCRs [65]. Indeed, it has been shown that HTR4 and HTR6 exhibit

constitutive $G\alpha_s$ activity [66–68]. The cellular heterogeneity of basal cAMP levels in cells expressing

HTR4 or HTR6 is simply explained by the variability in the expression levels of those receptors and

the constitutive activity of the receptors for $G\alpha_s$. It would be interesting to investigate whether the

expression levels of endogenous HTR4 and HTR6 are heterogenous, and if so, whether the cellular

heterogeneity plays a physiological role.

We established a live-cell fluorescence imaging system for quantifying four GPCR downstream

signals at the single-cell level. We would like to make two additional points in regard to technical

matters. The first concern is endogenous GPCRs. In this study, we targeted dopamine and serotonin

receptors, most of which are not expressed in HeLa cells, and therefore the data analysis was

straightforward. However, in cases in which the endogenous GPCR responds to the ligand, the data

should be analyzed with great care. The second concerns the biosensors for multiplexed fluorescence

imaging. Because of the spectral overlap of biosensors, we established two types of reporter cell lines

and co-cultured them, limiting multiplexed imaging at the single-cell level because of just two of the

permutations for biosensor pairs. Ideally, it is desirable to establish one type of cell line expressing all

reporters from the viewpoint of high-throughput screening. This issue could be solved by using the

recently developed single fluorophore-based biosensors [71,72], making it easier to understand cross-

talk and feedback control between signaling molecules at the single-cell level. In addition, recent
advances in fluorescence imaging allow us to visualize multiple cell signaling at the single-cell level
[73,74]. Further studies will be needed to visualize the downstream signaling dynamics of many types
of GPCRs and reveal the information processing mechanism.
597

598 **Data availability**
599 Data and reagents are available upon request to the corresponding authors.
600

601 **Competing Interests**
602 The authors declare that there are no competing interests associated with the manuscript.
603

604 **Funding**
605 K.A. was supported by the CREST program, a JST Grant (JPMJCR1654), and JSPS KAKENHI Grants
606 (nos. 18H04754 "Resonance Bio", 18H02444, and 19H05798). Y.G. was supported by a JSPS
607 KAKENHI Grant (no.19K16050) and a Jigami Yoshifumi Memorial Research Grant. Y.K. was
608 supported by JSPS KAKENHI Grants (nos. 19K16207 and 19H05675).
609

610 # Author contributions
611 R.T., Y.G., and K.A. designed the research. R.T., Y.K., and Y.G. performed all experiments and data
612 analysis. R.T., Y.G., Y.K., and K.A. wrote the manuscript.
613

Acknowledgments

We thank all members of the Aoki Laboratory for their helpful discussions and assistance.

References

Fredriksson, R., Lagerström, M. C., Lundin, L.-G. and Schiöth, H. B. (2003) The G-protein-coupled receptors in the human genome form five main families. Phylogenetic analysis, paralogon groups, and fingerprints. Mol. Pharmacol. **63**, 1256–1272.

Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schiöth, H. B. and Gloriam, D. E. (2017) Trends in GPCR drug discovery: new agents, targets and indications. Nat. Rev. Drug Discov. **16**, 829–842.

Gilman, A. G. (1987) G proteins: transducers of receptor-generated signals. Annu. Rev. Biochem. **56**, 615–649.

Sprang, S. R. (1997) G protein mechanisms: insights from structural analysis. Annu. Rev. Biochem. **66**, 639–678.

Ross, E. M., Howlett, A. C., Ferguson, K. M. and Gilman, A. G. (1978) Reconstitution of hormone-sensitive adenylate cyclase activity with resolved components of the enzyme. J. Biol. Chem., Elsevier **253**, 6401–6412.

Hazeki, O. and Ui, M. (1981) Modification by islet-activating protein of receptor-mediated regulation of cyclic AMP accumulation in isolated rat heart cells. J. Biol. Chem. **256**, 2856–2862.

Williamson, J. R., Cooper, R. H., Joseph, S. K. and Thomas, A. P. (1985) Inositol trisphosphate and diacylglycerol as intracellular second messengers in liver. Am. J. Physiol. **248**, C203–16.

Berridge, M. J. (1984) Inositol trisphosphate and diacylglycerol as second messengers. Biochem. J **220**, 345–360.

Kozasa, T., Jiang, X., Hart, M. J., Sternweis, P. M., Singer, W. D., Gilman, A. G., Bollag, G. and Sternweis, P. C. (1998) p115 RhoGEF, a GTPase activating protein for Galpha12 and Galpha13. Science **280**, 2109–2111.

Shenoy, S. K., Drake, M. T., Nelson, C. D., Houtz, D. A., Xiao, K., Madabushi, S., Reiter, E., Premont, R. T., Lichtarge, O. and Lefkowitz, R. J. (2006) β-Arrestin-dependent, G Protein-independent ERK1/2 Activation by the β2 Adrenergic Receptor. J. Biol. Chem. **281**, 1261–1273.

Copik, A. J., Baldys, A., Nguyen, K., Sahdeo, S., Ho, H., Kosaka, A., Dietrich, P. J., Fitch, B., Raymond, J. R., Ford, A. P. D. W., et al. (2015) Isoproterenol acts as a biased agonist of the alpha-1A-adrenoceptor that selectively activates the MAPK/ERK pathway. PLoS One **10**, e0115701.

Mochizuki, N., Ohba, Y., Kiyokawa, E., Kurata, T., Murakami, T., Ozaki, T., Kitabatake, A., Nagashima, K. and Matsuda, M. (1999) Activation of the ERK/MAPK pathway by an isoform of rap1GAP associated with Gαi. Nature, Macmillan Magazines Ltd. **400**, 891.

Smedler, E. and Uhlén, P. (2014) Frequency decoding of calcium oscillations. Biochim. Biophys. Acta **1840**, 964–969.

Sugiyama, K., Tago, K., Matsushita, S., Nishikawa, M., Sato, K., Muto, Y., Nagase, T. and Ueda, H. (2017) Heterotrimeric G protein Gαs subunit attenuates PLEKHG2, a Rho family-specific guanine nucleotide exchange factor, by direct interaction. Cell. Signal. **32**, 115–123.

Maziarz, M., Park, J.-C., Leyme, A., Marivin, A., Garcia-Lopez, A., Patel, P. P. and Garcia-Marcos, M. (2020) Revealing the Activity of Trimeric G-proteins in Live Cells with a Versatile

Biosensor Design. Cell **182**, 770–785.e16.

16 Olsen, R. H. J., DiBerto, J. F., English, J. G., Glaudin, A. M., Krumm, B. E., Slocum, S. T., Che,
T., Gavin, A. C., McCorvy, J. D., Roth, B. L., et al. (2020) TRUPATH, an open-source biosensor
platform for interrogating the GPCR transducerome. Nat. Chem. Biol. **16**, 841–849.

17 Wright, S. C. and Bouvier, M. (2021) Illuminating the complexity of GPCR pathway selectivity -
advances in biosensor development. Curr. Opin. Struct. Biol. **69**, 142–149.

18 Kroeze, W. K., Sassano, M. F., Huang, X.-P., Lansu, K., McCorvy, J. D., Giguère, P. M., Sciaky,
N. and Roth, B. L. (2015) PRESTO-Tango as an open-source resource for interrogation of the
druggable human GPCRome. Nat. Struct. Mol. Biol. **22**, 362–369.

19 Inoue, A., Raimondi, F., Kadji, F. M. N., Singh, G., Kishi, T., Uwamizu, A., Ono, Y., Shinjo, Y.,
Ishida, S., Arang, N., et al. (2019) Illuminating G-Protein-Coupling Selectivity of GPCRs. Cell,
Elsevier **0**.

20 Ponsioen, B., Zhao, J., Riedl, J., Zwartkruis, F., van der Krogt, G., Zaccolo, M., Moolenaar, W.
H., Bos, J. L. and Jalink, K. (2004) Detecting cAMP-induced Epac activation by fluorescence
resonance energy transfer: Epac as a novel cAMP indicator. EMBO Rep. **5**, 1176–1180.

21 Wu, J., Liu, L., Matsuda, T., Zhao, Y., Rebane, A., Drobizhev, M., Chang, Y.-F., Araki, S., Arai,
Y., March, K., et al. (2013) Improved orange and red $Ca^2\pm$ indicators and photophysical
considerations for optogenetic applications. ACS Chem. Neurosci. **4**, 963–972.

22 van Unen, J., Reinhard, N. R., Yin, T., Wu, Y. I., Postma, M., Gadella, T. W. J. and Goedhart, J.
(2015) Plasma membrane restricted RhoGEF activity is sufficient for RhoA-mediated actin
polymerization. Sci. Rep., The Author(s) **5**, 14693.

23 Regot, S., Hughey, J. J., Bajar, B. T., Carrasco, S. and Covert, M. W. (2014) High-sensitivity
measurements of multiple kinase activities in live single cells. Cell, Journal Article, Elsevier **157**,
1724–1734.

24 Okashah, N., Wan, Q., Ghosh, S., Sandhu, M., Inoue, A., Vaidehi, N. and Lambert, N. A. (2019)
Variable G protein determinants of GPCR coupling selectivity. Proc. Natl. Acad. Sci. U. S. A.
116, 12054–12059.

25 Akagi, T., Shishido, T., Murata, K. and Hanafusa, H. (2000) v-Crk activates the phosphoinositide
3-kinase/AKT pathway in transformation. Proc. Natl. Acad. Sci. U. S. A. **97**, 7290–7295.

26 Zhao, Y., Araki, S., Wu, J., Teramoto, T., Chang, Y.-F., Nakano, M., Abdelfattah, A. S., Fujiwara,
M., Ishihara, T., Nagai, T., et al. (2011) An Expanded Palette of Genetically Encoded Ca2+
Indicators. Science, American Association for the Advancement of Science **333**, 1888–1891.

27 Yusa, K., Rad, R., Takeda, J. and Bradley, A. (2009) Generation of transgene-free induced
pluripotent mouse stem cells by the piggyBac transposon. Nat. Methods **6**, 363–369.

28 Komatsu, N., Aoki, K., Yamada, M., Yukinaga, H., Fujita, Y., Kamioka, Y. and Matsuda, M.
(2011) Development of an optimized backbone of FRET biosensors for kinases and GTPases.
Mol. Biol. Cell, Journal Article **22**, 4647–4656.

29 Kawakami, K. and Noda, T. (2004) Transposition of the Tol2 element, an Ac-like element from
the Japanese medaka fish Oryzias latipes, in mouse embryonic stem cells. Genetics **166**, 895–899.

30 Shcherbakova, D. M., Baloban, M., Emelyanov, A. V., Brenowitz, M., Guo, P. and Verkhusha, V.
V. (2016) Bright monomeric near-infrared fluorescent proteins as tags and biosensors for
multiscale imaging. Nat. Commun. **7**, 12405.

31 Aoki, K., Kondo, Y., Naoki, H., Hiratsuka, T., Itoh, R. E. and Matsuda, M. (2017) Propagating
Wave of ERK Activation Orients Collective Cell Migration. Dev. Cell, Elsevier Inc. **43**, 305–
317.e5.

32 Maryu, G., Matsuda, M. and Aoki, K. (2016) Multiplexed Fluorescence Imaging of ERK and Akt
Activities and Cell-cycle Progression. Cell Struct. Funct. **41**, 81–92.

33 Miyoshi, H., Blömer, U., Takahashi, M., Gage, F. H. and Verma, I. M. (1998) Development of a
self-inactivating lentivirus vector. J. Virol. **72**, 8150–8157.

34 Sakurai, A., Matsuda, M. and Kiyokawa, E. (2012) Activated Ras Protein Accelerates Cell Cycle
Progression to Perturb Madin-Darby Canine Kidney Cystogenesis*. J. Biol. Chem. **287**, 31703–
31711.
35 Hino, N., Rossetti, L., Marín-Llauradó, A., Aoki, K., Trepat, X., Matsuda, M. and Hirashima, T.
(2020) ERK-Mediated Mechanochemical Waves Direct Collective Cell Polarization. Dev. Cell,
Elsevier **0**.
36 Moffat, J., Grueneberg, D. A., Yang, X., Kim, S. Y., Kloepfer, A. M., Hinkle, G., Piqani, B.,
Eisenhaure, T. M., Luo, B., Grenier, J. K., et al. (2006) A lentiviral RNAi library for human and
mouse genes applied to an arrayed viral high-content screen. Cell **124**, 1283–1298.
37 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012) Fiji: an open-source platform for biological-
image analysis. Nat. Methods **9**, 676–682.
38 Kudo, T., Jeknić, S., Macklin, D. N., Akhter, S., Hughey, J. J., Regot, S. and Covert, M. W.
(2018) Live-cell measurements of kinase activity in single cells using translocation reporters. Nat.
Protoc. **13**, 155–169.
39 Schmidt, U., Weigert, M., Broaddus, C. and Myers, G. (2018) Cell Detection with Star-Convex
Polygons. In Medical Image Computing and Computer Assisted Intervention – MICCAI 2018, pp
265–273, Springer International Publishing.
40 Weigert, M., Schmidt, U., Haase, R., Sugawara, K. and Myers, G. (2020) Star-convex Polyhedra
for 3D Object Detection and Segmentation in Microscopy. In 2020 IEEE Winter Conference on
Applications of Computer Vision (WACV), pp 3655–3662.
41 Jaqaman, K., Loerke, D., Mettlen, M., Kuwata, H., Grinstein, S., Schmid, S. L. and Danuser, G.
(2008) Robust single-particle tracking in live-cell time-lapse sequences. Nat. Methods **5**, 695–702.
42 Tavenard, R., Faouzi, J., Vandewiele, G., Divo, F., Androz, G., Holtz, C., Payne, M., Yurchak, R.,
Rußwurm, M., Kolar, K., et al. (2020) Tslearn, A Machine Learning Toolkit for Time Series Data.
J. Mach. Learn. Res. **21**, 1–6.
43 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011) Scikit-learn: Machine Learning in Python. J.
Mach. Learn. Res. **12**, 2825–2830.
44 Wise, R. A. (2004) Dopamine, learning and motivation. Nat. Rev. Neurosci. **5**, 483–494.
45 Beaulieu, J.-M. and Gainetdinov, R. R. (2011) The physiology, signaling, and pharmacology of
dopamine receptors. Pharmacol. Rev. **63**, 182–217.
46 Beaulieu, J.-M., Espinoza, S. and Gainetdinov, R. R. (2015) Dopamine receptors - IUPHAR
Review 13. Br. J. Pharmacol. **172**, 1–23.
47 Andersen, P. H., Gingrich, J. A., Bates, M. D., Dearry, A., Falardeau, P., Senogles, S. E. and
Caron, M. G. (1990) Dopamine receptor subtypes: beyond the D1/D2 classification. Trends
Pharmacol. Sci. **11**, 231–236.
48 Lee, S. P., So, C. H., Rashid, A. J., Varghese, G., Cheng, R., Lança, A. J., O'Dowd, B. F. and
George, S. R. (2004) Dopamine D1 and D2 receptor Co-activation generates a novel
phospholipase C-mediated calcium signal. J. Biol. Chem. **279**, 35671–35678.
49 Rashid, A. J., So, C. H., Kong, M. M. C., Furtak, T., El-Ghundi, M., Cheng, R., O'Dowd, B. F.
and George, S. R. (2007) D1-D2 dopamine receptor heterooligomers with unique pharmacology
are coupled to rapid activation of Gq/11 in the striatum. Proc. Natl. Acad. Sci. U. S. A. **104**, 654–
659.
50 Hasbi, A., Fan, T., Alijaniaram, M., Nguyen, T., Perreault, M. L., O'Dowd, B. F. and George, S.
R. (2009) Calcium signaling cascade links dopamine D1-D2 receptor heteromer to striatal BDNF
production and neuronal growth. Proc. Natl. Acad. Sci. U. S. A. **106**, 21377–21382.
51 Hasbi, A., O'Dowd, B. F. and George, S. R. (2010) Heteromerization of dopamine D2 receptors
with dopamine D1 or D5 receptors generates intracellular calcium signaling by different

mechanisms. Curr. Opin. Pharmacol. **10**, 93–99.

52 Davis, K. L., Dennis, C., Joseph, C. and Charles, N. (2002) Neuropsychopharmacology: The Fifth
Generation of Progress, Lippincott Williams & Wilkins.

53 Nichols, D. E. and Nichols, C. D. (2008) Serotonin receptors. Chem. Rev. **108**, 1614–1641.

54 Uhlén, M., Fagerberg, L., Hallström, B. M., Lindskog, C., Oksvold, P., Mardinoglu, A.,
Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., et al. (2015) Proteomics. Tissue-based map
of the human proteome. Science **347**, 1260419.

55 Cook, S. J. and McCormick, F. (1993) Inhibition by cAMP of Ras-dependent activation of Raf.
Science **262**, 1069–1072.

56 Mischak, H., Seitz, T., Janosch, P., Eulitz, M., Steen, H., Schellerer, M., Philipp, A. and Kolch,
W. (1996) Negative regulation of Raf-1 by phosphorylation of serine 621. Mol. Cell. Biol. **16**,
5409–5418.

57 Hoffmann, R., Baillie, G. S., MacKenzie, S. J., Yarwood, S. J. and Houslay, M. D. (1999) The
MAP kinase ERK2 inhibits the cyclic AMP-specific phosphodiesterase HSPDE4D3 by
phosphorylating it at Ser579. EMBO J. **18**, 893–903.

58 Baillie, G. S., MacKenzie, S. J., McPhee, I. and Houslay, M. D. (2000) Sub-family selective
actions in the ability of Erk2 MAP kinase to phosphorylate and regulate the activity of PDE4
cyclic AMP-specific phosphodiesterases. Br. J. Pharmacol. **131**, 811–819.

59 Insel, P. A., Wilderman, A., Zambon, A. C., Snead, A. N., Murray, F., Aroonsakool, N.,
McDonald, D. S., Zhou, S., McCann, T., Zhang, L., et al. (2015) G Protein-Coupled Receptor
(GPCR) Expression in Native Cells: "Novel" endoGPCRs as Physiologic Regulators and
Therapeutic Targets. Mol. Pharmacol. **88**, 181–187.

60 Purvis, J. E. and Lahav, G. (2013) Encoding and decoding cellular information through signaling
dynamics. Cell **152**, 945–956.

61 Albeck, J. G., Mills, G. B. and Brugge, J. S. (2013) Frequency-modulated pulses of ERK activity
transmit quantitative proliferation signals. Mol. Cell **49**, 249–261.

62 Aoki, K., Kumagai, Y., Sakurai, A., Komatsu, N., Fujita, Y., Shionyu, C. and Matsuda, M. (2013)
Stochastic ERK activation induced by noise and cell-to-cell propagation regulates cell density-
dependent proliferation. Mol. Cell **52**, 529–540.

63 Grundmann, M. and Kostenis, E. (2017) Temporal Bias: Time-Encoded Dynamic GPCR
Signaling. Trends Pharmacol. Sci. **38**, 1110–1124.

64 Averaimo, S. and Nicol, X. (2014) Intermingled cAMP, cGMP and calcium spatiotemporal
dynamics in developing neuronal circuits. Front. Cell. Neurosci. **8**, 376.

65 Seifert, R. and Wenzel-Seifert, K. (2002) Constitutive activity of G-protein-coupled receptors:
cause of disease and common property of wild-type receptors. Naunyn. Schmiedebergs. Arch.
Pharmacol. **366**, 381–416.

66 Blondel, O., Gastineau, M., Langlois, M. and Fischmeister, R. (1998) The 5-HT4 receptor
antagonist ML10375 inhibits the constitutive activity of human 5-HT4(c) receptor. Br. J.
Pharmacol. **125**, 595–597.

67 Claeysen, S., Sebben, M., Becamel, C., Bockaert, J. and Dumuis, A. (1999) Novel brain-specific
5-HT4 receptor splice variants show marked constitutive activity: role of the C-terminal
intracellular domain. Mol. Pharmacol. **55**, 910–920.

68 Deraredj Nadim, W., Chaumont-Dubel, S., Madouri, F., Cobret, L., De Tauzia, M.-L., Zajdel, P.,
Bénédetti, H., Marin, P. and Morisset-Lopez, S. (2016) Physical interaction between
neurofibromin and serotonin 5-HT6 receptor promotes receptor constitutive activity. Proc. Natl.
Acad. Sci. U. S. A. **113**, 12310–12315.

69 Bush, A., Vasen, G., Constantinou, A., Dunayevich, P., Patop, I. L., Blaustein, M. and Colman-
Lerner, A. (2016) Yeast GPCR signaling reflects the fraction of occupied receptors, not the
number. Mol. Syst. Biol. **12**, 898.

70   Cai, X., Flores-Hernandez, J., Feng, J. and Yan, Z. (2002) Activity-dependent bidirectional
regulation of GABA(A) receptor channels by the 5-HT(4) receptor-mediated signalling in rat
prefrontal cortical pyramidal neurons. J. Physiol. **540**, 743–759.
71   Mehta, S., Zhang, Y., Roth, R. H., Zhang, J.-F., Mo, A., Tenner, B., Huganir, R. L. and Zhang, J.
(2018) Single-fluorophore biosensors for sensitive and multiplexed detection of signalling
activities. Nat. Cell Biol.
72   Mahlandt, E. K., Arts, J. J. G., van der Meer, W. J., van der Linden, F. H., Tol, S., van Buul, J. D.,
Gadella, T. W. J. and Goedhart, J. (2021, February 8) Visualizing endogenous RhoA activity with
an improved localization-based, genetically encoded biosensor. bioRxiv.
73   Scott, B. L. and Hoppe, A. D. (2015) Optimizing fluorescent protein trios for 3-Way FRET
imaging of protein interactions in living cells. Sci. Rep. **5**, 10270.
74   Linghu, C., Johnson, S. L., Valdes, P. A., Shemesh, O. A., Park, W. M., Park, D., Piatkevich, K.
D., Wassie, A. T., Liu, Y., An, B., et al. (2020) Spatial Multiplexing of Fluorescent Reporters for
Imaging Signaling Network Dynamics. Cell **183**, 1682–1698.e24.

819