Title:

Development of affinity beads-based *in vitro* metal-ligand binding assay reveals

dominant cadmium affinity of thiol-rich small peptides phytochelatins beyond

glutathione

Author names and addresses:

Shimpei Uraguchi[1], Kenichiro Nagai[2], Fumii Naruse[1], Yuto Otsuka[1], Yuka Ohshiro[1],

Ryosuke Nakamura[1], Yasukazu Takanezawa[1], Masako Kiyono[1]

[1] Department of Public Health, School of Pharmacy, Kitasato University, 5-9-1

Shirokane, Minato-ku, Tokyo 108-8641, Japan

[2] Medicinal Research Laboratories, School of Pharmacy, Kitasato University, 5-9-1

Shirokane, Minato-ku, Tokyo, 108-8641, Japan

*Corresponding author:

Prof. Dr. Masako Kiyono, telephone: +81-3-5791-6264, email address:

kiyonom@pharm.kitasato-u.ac.jp

Running title:

Metal-thiol binding affinity by beads-based assay

Abbreviations:

DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene; DIPCI, *N,N′*-diisopropylcarbodiimide; DMF, *N,N*-dimethylformamide; GSH, glutathione; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; HFIP, 1,1,1,3,3,3-hexafluoro-2-propanol; HOBt, hydroxybenzotriazole; ICP-OES, inductively coupled plasma-optical emission spectrometry, ICT, isothermal titration calorimetry; LC-ESI-MS, liquid chromatography-electrospray ionization mass-spectrometry, LC-ICP-MS, liquid chromatography-inductively coupled plasma mass-spectrometry; NA, nicotianamine; NMR, nuclear magnetic resonance; PC, phytochelatin; PCS, phytochelatin synthase; TFA, trifluoroacetic acid; TIPS, triisopropylsilane.

Keywords:

glutathione; metal; ligands; phytochelatins; small peptides

Conflicts of interest:

No conflicts of interest to be declared.

Abstract:

For a better understanding of metal-ligand interaction and its function in cells, we developed an easy, sensitive, and high-throughput method to quantify ligand-metal(loid) binding affinity under physiological conditions, by combining ligand-attached affinity beads and inductively coupled plasma-optical emission spectrometry (ICP-OES). Glutathione (GSH) and two phytochelatins (PC2 and PC3, small peptides with different thiol numbers) were employed as model ligands and attached to affinity beads. The principal of the assay resembles that of affinity purification of proteins in biochemistry: metals binding to the ligand on the beads and the rest in the buffer are separated by a spin-column and quantified by ICP-OES. The binding assay using the GSH-attached beads and various metal(loid)s suggested that the assay can detect the different affinity of the metal-ligand interactions, in accordance with the order of the Irving–Williams series and the reported stability constants. The binding assay using PC2 or PC3-attached beads suggested positive binding between PCs and nickel, copper, zinc, cadmium, and arsenite ions in a thiol-dependent manner. We then conducted the competition assay using cadmium, manganese, iron, copper, and zinc and the results suggested a better binding affinity of PC2 with cadmium than with the essential metals. Another competition assay using PC2 and GSH suggested a robust binding affinity between PCs and cadmium compared to GSH and cadmium. These results suggested the dominance of PC-Cd complex formation *in vitro*, supporting the physiological importance of PCs for the detoxification of cadmium *in vivo*. We also discuss the further potential application of the assay.

Introduction

The essential metals are utilized by cells as cofactors for many proteins [1]. A bioinformatics study estimated that approximately 40% of the enzymes with known three-dimensional structures required metals at their catalytic centers [2]. The proportions of essential metals associated with enzymes are 6–9% of the analyzed enzymes for zinc (Zn), iron (Fe), and manganese (Mn), respectively, and 1% for copper (Cu). These metals are preferentially utilized in respective classes of Enzyme Commission (EC): for instance, Fe is mostly utilized by oxidoreductases (EC1) and Zn-containing enzymes are prominently enriched in hydrolases (EC3)[2]. Zn also plays structural roles in the Zn-binding proteins containing zinc-finger or RING domains, suggesting the importance of Zn in transcriptional and post-translational regulations, especially in eukaryotes [3,4].

Meanwhile, the essential heavy metals in the free ionic forms are toxic due to the high reactivity. Thus, intracellular metal-buffering systems tightly maintain labile metal concentrations in the cell at very low levels to avoid their toxicity. The studies on bacteria and yeast suggest that there is practically no cytosolic pool of free Cu and Zn ions under steady-state conditions [5–7]. The essential heavy metals imported to the cells are preferentially delivered to the target proteins requiring respective metals, and the rest are "buffered", bound to various small-molecular ligands as well as proteins like metallothionein in mammals [8,9]. Under conditions with the excess metals, the biological chelators then function in "muffling", chelating the excess metal ions in the cell and avoiding the cytotoxicity [8,9].

Among such metabolites, GSH is believed as a major metal-chelating small

molecule especially in mammalian cells [8,10]. In addition, many diverse

small-molecular ligands have been discovered as physiologically significant metal

chelators in bacteria, yeast, worm, and higher plants [11–13]. (Phyto)siderophores and

its precursor nicotianamine (NA) are typical examples that function as Fe and Zn

binding ligands, respectively. Another established example of the small molecular

chelator is phytochelatins (PCs), cysteine-rich small peptides. The general structure of

PCs is (γ-Glu-Cys)n-Gly with *n* usually between 2 and 7. PCs are non-ribosomally

synthesized from GSH (γ-Glu-Cys-Gly) by phytochelatin synthases (PCSs). Compared

to (phyto)siderophores and NA that function as ligands for essential heavy metals [12],

PCs can be characterized as high-affinity chelators for a wide range of toxic

metal(loid)s such as cadmium (Cd) and arsenic (As). In plants, PCs play a central role in

the detoxification of various toxic metal(loid)s [14–18]. Interestingly, the PC-mediated

detoxification of toxic metal(loid)s is also demonstrated in the fission yeast

Schizosaccharomyces pombe [19], in which PCs were first discovered [20], and in the

nematode worm *Caenorhabditis elegans* [21]. These suggest the conserved functions of

PC/PCS for toxic metal(loid) detoxification in taxonomically distant organisms.

The physiological significance of GSH and PCs in metal(loid) detoxification

is mostly demonstrated by phenotypes induced by GSH synthesis inhibitors or PC/GSH

synthesis genes disruption, which lead to metal(loid) hypersensitivity [14–16,21,22].

These results indicate that ligand-metal complex formation is a key process for the

PC/GSH-mediated   toxic   metal(loid)   detoxification,   however,   there   is   limited

information on complex formation between PCs/GSH and metal(loid)s, especially under

physiologically relevant conditions. It appears first due to the technical difficulties

overall [23], and secondly, due to unfamiliarity with required analytical techniques

and/or limited access to the analytical devices especially for many molecular

metal-biologists. For example, the PCs/GSH-metal(loid) complex formation *in vitro* is

often examined by potentiometry, voltammetry, and nuclear magnetic resonance (NMR).

A problem with such an assay is that it is often carried out under physiologically

irrelevant conditions (e.g. extremely low pH conditions)[24]. Isothermal titration

calorimetry (ITC) would be a better approach, providing detailed *in vitro* data of

binding chemistry under physiological conditions, but the device is expensive and not

so widely available. Another disadvantage of these analytical chemistry approaches is

that it is technically difficult to take account of a cellular matrix containing various

metals and metabolites other than the targets, although they are also potential

interactants. Liquid-chromatography-inductively coupled plasma mass-spectrometry

(LC-ICP-MS) and liquid-chromatography-electrospray ionization mass-spectrometry

(LC-ESI-MS) enable direct detection of multiple metal-ligand complexes *in vitro* and *in*

vivo, however, the technical difficulty of analyzing the intact complex form remains the

bottleneck [25].

For these reasons, an alternative assay especially for metal biologists would

be a useful option to advance understanding of metal-ligand interaction in the cells. The

present study aimed to develop an ICP-based simple but quantitative method to evaluate

ligand-metal(loid) interaction under various physiological conditions. ICP-optical

emission spectrometry (OES) was used as an analytical tool. ICP-OES as well as

ICP-MS is a common device among metal-biologists, compared to the other analytical

techniques, and more importantly, ICP-based element analysis provides highly sensitive

and high-throughput quantification. We employed GSH and PCs as the model ligands

and prepared the ligand-attached affinity beads for the assay. The GSH-metal(loid)s

binding ratios derived from our method coincided with the reported values. With the

established assay, we examined PC and Cd interactions under physiological conditions

containing GSH and several essential metals. The competitive assays demonstrated

preferential interaction of PCs and Cd, which supports the *in vivo* observations that PCs

play a central role in Cd detoxification in plants, fission yeast, and *C. elegans*.

Results

Preparation of phytochelatin- and glutathione-attached beads

PCs are commercially available only on a 1 mg scale (Eurogentec, Seraing,

Belgium) as far as we searched, and it was a considerable cost disadvantage to our large

demand of PCs for the affinity beads preparation. Thus, we first synthesized a shorter

PC [(γ-Glu-Cys)$_2$-Gly, PC2] and a longer PC [(γ-Glu-Cys)$_3$-Gly, PC3] by a standard

Fmoc solid-phase peptide synthesis protocol (Fig. 1). The two different length PCs are

common PC isoforms synthesized in *S. pombe*, *C. elegans,* and various plant cells

exposed to toxic metal(loid) ions such as Cd(II) and As(III) [16,19,21,26,27].

Fmoc-Gly-OH was loaded onto 2-chlorotrityl chloride resin (1.38 mmol/g, 200 mg).

Fmoc protecting group was removed using a solution of 2%

1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) in *N,N*-dimethylformamide (DMF).

Condensation of Fmoc-Cys(Trt)-OH and Fmoc-Glu-O*t*Bu was performed using

hydroxybenzotriazole (HOBt) and *N,N′*-diisopropylcarbodiimide (DIPCI). The resulting

peptide was cleaved from the resin with 20% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP)

in CH_2Cl_2, and then deprotected with trifluoroacetic acid (TFA)/triisopropylsilane

(TIPS)/H_2O (95:2.5:2.5). After purification by preparative HPLC, PC2 and PC3 were

obtained in 64% yield (95.0 mg) and 64% yield (126 mg), respectively. Then we

analyzed the synthetic PC2 and PC3 by total ion-chromatography (Fig. 2A, B),

UV-spectroscopy (Fig. 2C, D), and mass-spectroscopy (Fig. 2E, F), and confirmed the

adequate purity and expected molecular size of the products.

Using the synthetic PC2, PC3, and commercially obtained GSH, we

attempted to prepare PC- or GSH-attached affinity beads (Fig. 2G). Hydrophilic beads

which are only modified with formyl groups were used as a carrier resin. GSH, PC2, or

PC3 was loaded onto the beads via reductive amination using $NaBH_3CN$. This reaction

was conducted twice independently to obtain two different preparations of the

respective ligand-attached beads. The yield was quantified by Ellman's reagent [28] and

was overall within a similar range (64.1 to 78.4 nmol thiol per mg beads) irrespective of

the ligands and preparations (Fig. 2G). The beads treated only with the reaction buffer

not containing GSH, PC2, or PC3 were prepared as control beads in the subsequent

binding assay.

Establishment of the assay protocol for metal(loid)-ligand interaction

We then designed and tested protocols for evaluating the ligand-metal(loid)

interactions using the prepared ligand-attached beads (Fig. 3). A reaction mixture for the

standard protocol was prepared in a 5 ml plastic tube containing 3.0 mg of the beads and

200 nmol of metal(loid)-of-interest. 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

(HEPES) buffer (pH 7.2) was used as a matrix instead of a phosphate buffer to avoid the

potential interaction of the metal(loid) and phosphate ions. The prepared tubes were

incubated at room temperature for 5 min or overnight with continuous rotation to avoid

sedimentation of the beads. Sample collection after the incubation was performed by

two different protocols: collecting a filtrate without beads using a spin column (Protocol

A) and collecting beads without a solution by centrifugation (Protocol B). Element

contents in the collected samples were quantified by ICP-OES.

For calculation of $[M]_{binding}$ [the metal(loid) binding to the ligand on the

beads], in Protocol A, $[M]_{ligand\ filtrate}$ [the metal(loid) amount in the filtrate from the

ligand-attached beads reaction] was subtracted from $[M]_{control\ filtrate}$ [the metal(loid)

amount in the filtrate from the control beads reaction] (Fig. 3). In Protocol B, $[M]_{control}$

$_{beads}$ [the metal(loid) amount in the control beads] was subtracted from $[M]_{ligand\ beads}$ [the

metal(loid) amount in the ligand-attached beads] to obtain $[M]_{binding}$ (Fig. 3).

The two protocols were tested with Cd as a model metal. To compare the

accuracy of Protocol A and B, we first quantified total Cd adsorption to the control and

ligand-attached beads (Fig. 4). For Protocol A, $[Cd]_{ligand\ filtrate}$ and $[Cd]_{control\ filtrate}$ were

respectively subtracted from Cd input to each reaction (200 nmol). For Protocol B,

[Cd]$_{control\ beads}$ and [Cd]$_{ligand\ beads}$ were directly used to assess total Cd adsorption to the

beads. Regardless of the types of the beads, the Cd adsorption to the beads did not

significantly differ between Protocol A and B (Fig. 4), suggesting that both protocols

can equally evaluate the metal binding to the beads as designed. Protocol A is more

convenient and practical because the separation of the beads and the buffer can be easily

and quickly achieved by a spin column centrifugation, whereas Protocol B requires

careful but quick removal of the buffer by pipetting after centrifugation. Probably

reflecting these differences, the standard deviation was a little smaller for Protocol A.

Taken together, we decided to choose Protocol A for further experiments.

Affinity variation of GSH, PC2, and PC3 with various metal(loid)s

206   The experiment to test the two protocols using Cd as a model metal indicated

the substantial chelation between the ligands and Cd because all the ligand-attached

beads (GSH, PC2, and PC3) adsorbed nearly 3-times more Cd than the control beads

(Fig. 4). The results motivated us to further conduct a full-scale experiment and we

quantified GSH and PCs-mediated chelation of Mn, Fe, Ni, Cu, Zn, As, and Cd (Fig. 5).

Divalent cation forms were used for Mn, Fe, Ni, Cu, Zn, and Cd, and trivalent arsenite

anion was used for As. 5-min incubation was applied for the initial experiment, and

overnight incubation was also examined for Mn, Fe, Cu, and As. The experiment for Fig.

5 was carried out with the first lot beads, except for Cd for which the independently

prepared second lot beads were also applied to examine possible lot-specific effects.

We first checked the metal(loid) adsorption to the controls beads as

non-specific background adsorption of the assay (Fig. 5A). The presented values are

percentages of the metal(loid) binding to the beads to the total input to the reaction (200

nmol). Non-specific adsorption to the hydrophilic beads ranged up to 30% of the input,

regardless of the tested metal(loid) species and incubation time, except for overnight

incubation with Fe (nearly 100% of the input to the reaction was adsorbed to the carrier

beads). The result of the non-specific background adsorption indicates that the

experimental condition provides enough metal(loid)s left in the reaction buffer which

can interact with the ligand on the ligand-attached beads, except for overnight Fe

incubation.

Then we calculated the metal(loid) binding ratio of GSH (Fig. 5B), PC2 (Fig.

5C), and PC3 (Fig. 5D), dividing $[M]_{binding}$ (Fig. 3) by the molar number of the ligand in

the reaction. For GSH which harbors one cysteine, the metal-binding ratio of 0.5 was

referred to as a standard to indicate a binding stoichiometry of one metal to two GSH.

The results of 5-min incubation for Ni, Zn, and Cd were close to 0.5 (Fig. 5B). The ratio

of 5-min incubation with Cu showed a somewhat intermediate value (0.36). Those of

Mn and Fe were significantly lower even than this value of Cu, suggesting a lower

affinity of Mn and Fe with GSH. For As which was spiked as arsenite, the binding ratio

of 0.33 was referred to as a standard indicating a binding stoichiometry of one As to

three GSH [29]. However, the ratio of As/GSH after 5-min incubation was 0.02, which

was far below the standard value of 0.33. To further examine the lower binding ratio of

(Cu), Mn, Fe, and As, the incubation time was prolonged from 5 min to overnight. The

binding ratios of Mn and As with GSH after overnight incubation was still quite lower

than the standards 0.5 and 0.33, respectively (Fig. 5B). The overnight incubation

increased the binding ratio of Cu and GSH from 0.36 (5-min) to 0.62, both of which

were within the range of the standard value of 0.5. The binding of Fe with GSH after the

overnight incubation was not detected due to the huge background Fe adsorption to the

carrier beads (Fig. 5A).

The metal(loid) binding ratio with PC2 was shown in Fig. 5C. For PC2, which

harbors two cysteines, the binding ratios of 0.5 and 1.0 as standards were referred to

indicate 1:2 and 1:1 binding of metal(loid) and PC2, respectively. The binding ratios of

Ni, Cu (overnight), Zn, and Cd were statistically similar and were close to 1.0 (Fig. 5C).

The ratio obtained from 5-min incubation with Cu was again a bit lower than that from

overnight incubation, but the difference was smaller than the case of GSH (Fig. 5B).

Binding ratios with PC2 for Mn (5-min and overnight) and Fe (5-min) were far below

1.0. The ratio for As/PC2 was close to 0.5 after overnight incubation, suggesting 1:2

binding of As(III) and PC2, whereas the result of 5-min incubation was much lower.

The metal(loid) binding ratio with PC3 was shown in Fig. 5D. PC3 contains

three cysteines, and the binding ratios of 1.0 and 1.5 as standards were referred to

indicate 1:1 and 3:2 binding of metal(loid) and PC3, respectively. The binding ratios of

Ni, Cu (5-min and overnight), Zn, and Cd were around 1.5. It should be noted that there

was no statistical difference between the incubation periods for Cu. The ratios for Mn

(5-min) and Fe (5-min) were 0.67 and 1.04, respectively, which were significantly lower

than the ratios for other divalent metals tested. The binding ratio of As and PC3

(overnight) was 0.98, indicating the 1:1 binding relationship, however, the 5-min

incubation again resulted in a lower value (0.22).

Effects of different beads preparation

264   Conducting the experiments to produce the most results for Fig. 4 and Fig. 5,

we used a large portion of the first preparation of the ligand-attached beads. Thus, we

prepared a second lot of the respective beads. The second preparation resulted in similar

yields however, the yield was overall a bit higher (Fig. 3G). We examined the possible

effects of the different lots on the metal(loid)-ligand-binding assay, using Cd as a model

again. Background non-specific Cd adsorption (Fig. 5A), and binding ratios with GSH

(Fig. 5B), PC2 (Fig. 5C), and PC3 (Fig. 5D) were not statistically different between the

first and second preparations by Tukey's HSD. In addition, the results of the Student's

T-test were also not significant ($P > 0.05$). These results suggest that differently

prepared ligand-attached beads can be equally used in the binding assay and we decided

to use the second lot for further experiments.

Binding competition between Cd and essential metals

277   The results of the single metal(loid)-ligand assay (Fig. 5) suggested different

affinity of the small thiol compounds to the various essential metals and toxic

metal(loid)s. We next planned to explore functional differences of the different length

thiols, GSH, PC2, and PC3, using the developed ligand-attached beads system. PCs are

believed to function in chelating free toxic metal(loid)s in the cytosol. However,

competition over PCs' binding sites can be expected in the cytosol between the toxic

metals and co-existing essential heavy metals. Thus, we conducted a competition assay

using Cd as a model. Cd, not As, was selected because Cd binding was stably detected

by the 5-min incubation, whereas As required the overnight incubation (Fig. 5). In this

competition assay, Cd (200 nmol) as well as essential metals [Mn (20 nmol), Fe (40

nmol), Cu (2 nmol), and Zn (20 nmol)] were added to the reaction. The ratio of the

supplemented metals was derived from a rough calculation based on the total element

concentrations in leaves of the plant model *Arabidopsis thaliana* [30–32].

As a result of the competition assay, Mn and Fe exhibited inferior affinity to

GSH, PC2, and PC3 compared to Cu, Zn, and Cd (Fig. 6A, B, C). Smaller fractions of

the Mn and Fe input (~30%) were associated with the ligands, and approximately 60%

or more of the Cu, Zn, and Cd input were associated with the ligands. We then

calculated profiles of the metals binding to GSH, PC2, and PC3, and compared them

with the profile of the input (Fig. 6D). Reflecting the lower affinity with the ligands, the

fractions of Mn and Fe binding to the ligands were smaller than those of the input. In

contrast, the Cd binding to the ligands was more than 80% of the total metals captured

by the ligands, which was larger than the input supplemented Cd fraction (73%). These

results further support the results of the single-metal binding assay (Fig. 5), suggesting a

preference of Cd in binding to the thiol ligands, compared to the tested essential metals.

Cd-binding competition between GSH and PCs

Next, we conducted another competition assay to examine a possible

preference of PC and Cd interaction (Fig. 7). First, the binding of 100 nmol Cd and

GSH beads competed with or without 100 nmol PC2 added as a free form competitor in

the reaction mixture. The Cd binding to GSH beads under the condition without the

competitor PC2 was 55% of the total Cd input (Fig. 7A), but the addition of PC2 as the

competitor to the reaction significantly decreased the Cd binding to GSH beads to 16%

of the total Cd input (Fig. 7A). Then, the inverted assay conditions were also tested (Fig.

7B): Binding of 100 nmol Cd by PC2 beads competed with 25 nmol GSH or 200 nmol

GSH supplemented to the reaction mixture in a free form. In the absence of GSH, 55%

of the input Cd was bound with PC2 on the beads, and the addition of 25 nmol nor 200

nmol GSH to the mixture as a competitor did not change the Cd binding profile to PC2

on the beads. The results overall suggest the superior binding affinity of Cd and PCs

over compared to GSH.

Discussion

Advantages of our metal-ligand binding assay

Physiological roles of metal-ligand interaction have been attracting attention

because of their significance in the essential metal homeostasis and toxic metal(loid)

detoxification in bacteria, plants, and humans [25,33–35]. The first goal of this study is

to establish a new assay system practically feasible for biologists interested in

evaluating metal-biological ligand interactions under cellular-mimicking conditions.

The idea of our assay resembles that of protein affinity purification protocols such as

immunoprecipitation in biochemistry. Using ICP as a readout tool is the first unique

advantage of our assay. ICP-OES and ICP-MS are widely used by metal biologists.

Moreover, the application of ICP enables rapid, highly sensitive, and stable

quantification of a large sample set (Figs. 3-4). Note that a standard ICP protocol takes

only 3-4 min for a single sample measurement of even more than 10 elements at ppb

(ICP-OES) or sub-ppb levels (ICP-MS). In addition, a spin column-based efficient

separation of the beads by centrifugation is also a key advantage of the assay. The

spin-column procedure is very familiar to molecular biologists and enables immediate

termination of the reaction of multiple samples. Besides technical advantages, it should

be noted that our assay system allows variable assay conditions which mimic cellular

environments. Thus, we can easily examine the effects of various co-existing metals and

metabolites (Figs. 5-7). These features are unique to our assay and imply the broad

utility of the protocol.

A few limitations exist in our assay: the assay cannot provide direct complex

form information and stability constants unlike the other conventional methods such as

potentiometry, voltammetry, NMR, ITC, and ESI-MS. However, the proposed assay has

significant advantages in terms of flexibility of assay conditions, high-throughput, and

technical feasibility especially for molecular biologists who are interested in

metal-ligand interactions under cellular environments.

GSH and metals interaction as a test model of the assay

In the present study, we employed GSH and two different length PCs (PC2

and PC3) as the model ligands for our assay (Figs. 1-3). First, the reliability of the

proposed assay was verified by assessing GSH-metal interactions (Fig. 5B). The affinity

between GSH and various metals is well characterized by conventional analytical

methods. GSH is believed as one of the most possible metabolites in the eukaryotic

cells to form metal-complexes through its sulfhydryl group [36]. The abundance in the

cells is another advantage of GSH as a metal-ligand: GSH concentrations in eukaryotic

cells (ranging from 0.1 mM to 8 mM [37–39]) are likely substantial enough to bind free

metals which would exist at much lower in the cells.

The results of binding experiments between GSH and well-studied cations

(Mn, Fe, Ni, Cu, Zn, and Cd) coincide with the reported results and the order of the

Irving–Williams series (Mn<Fe<Co<Ni<Cu, Zn) [40] which describes the complex

stability between divalent metal ions and organic ligands (note that Cd is not included in

the series, and Co is not tested in this study). For example, our assay suggests a weak

interaction of Mn and Fe with GSH (Fig. 5B), supporting the models suggested in the

previous studies: Mn(II)-GSH and Fe(II)-GSH interactions are 1:1 with markedly lower

stability constant (log K_1 values) of 2.7 and 5.1, respectively [41,42]. In contrast,

stronger affinity with GSH is suggested for highly reactive metals up in the

Irving–Williams series (Ni, Cu, and Zn) as well as Cd (Fig. 5B). The reported stability

constants (log β values) for these metals are around 10 or much higher even under

physiological pH conditions [43,44]. In our study, the metal/GSH ratio of around 0.5

was obtained for Ni, Cu, Zn, and Cd (Fig. 5B), suggesting 1:2 metal-GSH complex

formation as previously reported [43,44]. For Cu, there would be other possibilities. The

Cu/GSH ratios were slightly yet significantly lower than 0.5 under 5 min-incubating

conditions, and overnight incubation increased the ratio (Fig. 5B). This variation would

be explained by the controversial nature of Cu-GSH complex formation. Previous

studies imply Cu and GSH affinity with 1:2 and 1:1 stoichiometry [44] and in addition,

the formation of $Cu(I)_4(GS)_6$ is suggested [45]. Another note for the Cu assay is that we

applied Cu(II) to the assay, not Cu(I), because of its low solubility and extremely high

background adsorption to the beads in the preliminary experiment using CuCl. Cu(II) is

converted to Cu(I) under the reducing cellular conditions [45], which may explain the

increased interaction of Cu and GSH by the longer reaction period (Fig. 5B). Therefore,

taken together with these points, our results for Cu cannot suggest an exact

stoichiometry of Cu-GSH interaction, but at least suggest a stronger affinity with GSH

under physiological (cellular) conditions compared to Mn and Fe (Fig. 5B).

Nevertheless, the results of GSH beads (Fig. 5B) overall demonstrate that

under physiological pH conditions, the assay can differentiate the affinity of metals,

corresponding to the order of the Irving–Williams series and the established stability

constants.

Preferential interaction between phytochelatin and Cd

Several metabolites other than GSH are also suggested as biological metal

ligands. Among phytochelatin (PC) is an established example that mainly functions in

Cd detoxification through Cd-PC complex formation in higher plants, *S. pombe*, and *C.*

elegans [15,16,19,21]. Due to its exceeding significance in Cd detoxification, the PCS

activation mechanism has been extensively examined using recombinant proteins,

heterologous expression systems in *S. pombe*, plant cell cultures, and plant seedlings. In

brief, PCS activity is triggered substantially by various environmentally relevant ions

[e.g. As(III), As(V), Ag(I), Cd(II), Hg(II), and Pb(II)] and the excess of a few

physiologically essential metal ions [Cu(II) and Zn(II)] as well, irrespective of their

valency and positions in the periodic table [16–18,26,27,46–48]. On the other hand,

several elements essential for plant nutrition [K(I), Mg(II), Ca(II), Mn(II), and Ni(II)]

even in their excess levels do not activate PCSs [47,48]. Whether Fe activates PCSs is

still controversial [48]. Such metal-selectivity of PCSs raises a significant question of

how PCSs distinguish the activating ions and others. Recent studies from our group and

others suggest that the different regions of the C-terminal regulatory domain of PCSs

are involved in the metal(loid)-specific activation [16,49–52].

In contrast to the activation mechanism, much less is known about the subsequent

metal(loid)-PC complex formation. Only a few studies reported the detection of PC-Cd

and PC-As complexes in plant extracts or *in vitro* samples using LC-ESI-MS or

LC-ICP-MS [53–58]. Thus, the second purpose of the present study is to examine

preferences of metal(loid)-PC complex formation using our binding assay, mainly

focusing on Cd-PC interaction. The assay using PC2 beads and a single metal species

(Fig. 5C) suggests a 1:1 interaction between PC2 and Cd as well as Ni, Cu, and Zn, but

a much weaker interaction is suggested for Mn-PC2 and Fe-PC2. The 3:2 interaction

between Cd and PC3 is suggested in accordance with the three SH residues in PC3 (Fig.

5D). A similar 3:2 interaction was also suggested for Ni, Cu, and Zn. It should be noted

that unlike the case of PC2, Fe (5-min) is likely to interact with PC3 at the ratio of 1:1

(Fig. 5D). These results suggest a potential of Fe, Ni, Cu, and Zn interacting with PCs,

aside from their PCS activation ability. This implies that Cd and these metals may

compete for the PCs' binding sites. However, the Cd-PCs complexes are the major form

detected in plants exposed to Cd stress [55–57], suggesting that Cd-PC interaction is

dominant over the competitive essential metals.

A series of competitive assays were conducted to examine the hypothesis (Figs.

6-7). The competition assay between Cd and several plant essential metals suggests the

preferential binding affinity of PCs with Cd over the tested essential metals (Fig. 6). The

result supports the hypothesis and *in planta* observation of the dominance of Cd-PCs

formation [55–57]. In plant cellular conditions, sub-millimolar levels of GSH exit as

another potent Cd chelator other than PCs. Thus, we examined the Cd binding

preference between PC2 and GSH (Fig. 7). The assays clearly demonstrate the marked

preference of Cd-PC2 interaction over Cd-GSH (probably Cd-GS$_2$). The Cd-PC2

interaction was not interfered with even by the excess level of free GSH (Fig. 7B),

which supports the reported higher stability constant of Cd-PC2 than Cd-GS$_2$ [44,59].

Overall, our competition assays suggest the robust and preferential interaction of PC

and Cd over other co-existing essential metals and GSH. It should be noted that the

actual mechanism of PC-metal(loid) complex formation is still unclear. A previous

study suggested a possibility that the pre-formed Cd-GS$_2$ complex interacts with the

C-terminal regulatory domain of the PCS and activates the transpeptidase activity to

induce PC synthesis [60]. The previous study also proposed that a Cd ion is transferred

from the Cd-GS$_2$ complex to the PCS protein and activates PC synthesis [60]. Thus, it

can be postulated that Cd-PC complex formation (partly) occurs during the process of

PC synthesis [60]. Nevertheless, our competition assays (Figs. 6-7) suggest that the

formed Cd-PC complex would be stable enough in the cytosol, not interfered by GSH

and other essential metals until subsequent vacuolar compartmentation of the Cd-PC

complex.

Interaction between thiols and As(III)

Another promising partner of PC other than Cd(II) is As(III). In plants, As(III)

stress also induces PC synthesis, and the subsequent As(III)-PC complex formation and

vacuolar sequestration by ABC transporters are suggested to be crucial for As(III)

detoxification [16,50,61,62]. Formation of As(III)-(PC2)$_2$ and As(III)-PC3 is reported *in*

vitro and *in planta* [53,54,58]. The binding stoichiometry is in accordance with the

number of the thiol groups in PCs [note that three SH residues are required to catch one

As(III)]. Our results of As(III)-PC beads interaction assay under the overnight condition

clearly support these observations, with the As/PC2 ratio of 0.5 (As:PC2=1:2) and

As/PC3 ratio of 1 (As:PC3=1:1) (Fig. 5C and 5D). In addition to PCs, As(III) would

also interact with GSH and form As(GS)$_3$ *in vitro* [63]. However, the complex *in vivo* is

only observed under extreme biological conditions: the bile of rats injected with a very

high dose of arsenic [64] and the roots of Arabidopsis PCS1 mutant *cad1-3* [58]. The

instability of As(GS)$_3$ at physiological pH and temperature [24] would partly explain the

rareness of As(GS)$_3$ *in vivo*. Supporting this hypothesis, the As/GSH ratio obtained

under physiological conditions was much lower than 0.33 (Fig. 5B), the index for

As(GS)$_3$ formation. Taken together with the previous observations *in vitro* and *in vivo*,

our results suggest that As(GS)$_3$ complex is not a major form in the cells challenged by

As(III) except for the extreme conditions, and As-PCs complex is much more stable and

thus crucial for the As detoxification in plants.

Conclusion and future perspectives

The importance of toxic metals and essential metals homeostasis in plants as food

resources is increasing to achieve better human health, and the small molecular ligands

are major (potential) players in regulating these biological processes [25,65]. *In vitro*

evaluation of metal-ligand interactions is a crucial initial step for characterizing

ligand-based metal homeostasis in the cells. The proposed assay system in this study

has advantages regarding throughput, sensitivity, robustness, and specificity. Another

strong point is that the assay is flexible in terms of reaction conditions. Thus, it is

suitable for testing various metal-ligand combinations. Taking all these advantages, the

present study demonstrated dominant natures of PC-Cd interactions by a series of

binding assays.

Beyond PCs, the assay can be applied to test the potent metal affinity of other

ligands (candidates). For example, characterizing the metal interactions of the

Fe/Zn-chelator NA or other siderophores under physiological matrix conditions would

advance our understanding of the biological chelators. Other interesting ligand

candidates include isoforms of GSH and PCs in which the C-terminal Gly is replaced by

β-Ala, Ser, Gln, or Glu [66–68]. These iso-thiols are found in various plant species but

are little characterized. From the viewpoint of environmental science/ecotoxicology, the

assay can be used for the assessment of environmentally important metal(loid)s and

biological ligands interaction. Recently, in addition to the "classical metals", a number

of elements are newly utilized by high-tech industries and their levels in the

environment are accordingly increasing [69,70]. For these further applications,

detection by ICP-MS instead of ICP-OES is an option that would increase the

sensitivity of the assay and target elements to be tested.

Materials and methods

Synthesis of PC2 and PC3

PC2 and PC3 were synthesized by the Fmoc solid-phase peptide synthesis strategy. The

resulting peptides were purified by a preparative HPLC and then analyzed by an

ESI-MS to verify the mass of the purified peptides. A full method was provided as

supporting information.

Loading of GSH, PC2, or PC3 onto the affinity beads

TOYOPEARL AF-Formyl-650 (TOSOH, Tokyo, Japan) was selected as the basal beads.

TOYOPEARL AF-Formyl-650 (1 mL, 60 μmol/mL) was placed in a polypropylene

tube and washed with H_2O (1 mL, 1 min × 3). To the bead was added a solution of GSH

(18 mg, 60.0 µmol) in 0.1 M phosphate buffer (pH 7.0, 1 mL), and then sodium

cyanoborohydride (11 mg, 180 µmol) was added. The resulting mixture was agitated at

room temperature for 22 h. The solvents and soluble reagents were removed by

filtration, and the bead was washed with H_2O (1 mL, 1 min $\times$ 3), brine (1 mL, 1 min $\times$

3), H_2O (1 mL, 1 min $\times$ 3), and MeOH (1 mL, 1 min $\times$ 3). PC2 or PC3 were loaded onto

the affinity bead according to the loading procedure described above. The loading of the

resulting bead was determined by the amount of 2-nitro-5-thiobenzoate anion formed

from the reaction of 5,5'-dithiobis(2-nitrobenzoic acid), Ellman's reagent, with

sulfhydryl groups on the bead. The affinity resin (10 mg) was treated with 2.5 mM

Ellman's reagent in MeOH (2 mL) and N,N-diisopropylethylamine (5 µL) at room

temperature for 30 min. The total volume was adjusted to 5 mL by the addition of

MeOH and measured spectrophotometrically at 412nm. The yields were as shown in

Fig. 2G.

Single metal(loid)-beads assay

The interaction assay was conducted in a 5 mL Eppendorf tube containing 3.0 mg of the

beads (control, GSH, PC2, or PC3). The assay was repeated with at least three

replications. To evaluate the interaction between the ligand beads and various

metal(loid)s, 5 mL of 40 mM HEPES (pH 7.2) containing 200 nmol of $MnCl_2$, $FeCl_2$,

$(NH_4)_2Ni(SO_4)_2$, $CuSO_4$, $ZnSO_4$, $NaAsO_2$, or $CdCl_2$, was added to the 5 mL tube. The

tube was immediately set to a rotator and incubated at room temperature for 5 min. For

Mn, Fe, Cu, and As, overnight incubation was also tested.

Multiple metals competition assay

The competition assay using multiple metals was conducted as described for the single

metal(loid)-beads assay, except for the metal composition in the reaction buffer. The

reaction buffer [5 mL of 40 mM HEPES (pH 7.2)] contained 20 nmol of $MnCl_2$, 40

nmol of $FeCl_2$, 2 nmol of $CuSO_4$, 20 nmol of $ZnSO_4$, and 200 nmol of $CdCl_2$ and was

added to 3.0 mg of the beads (control, GSH, PC2, or PC3) in the 5 mL tube. The tube

was immediately set to a rotator and incubated at room temperature for 5 min. The

assay was repeated with three independent reactions.

PC2 and GSH competition assay

Cd-binding competition between PC2 and GSH was examined by modifying the single

metal(loid)-beads assay. For the assay using GSH-attached beads (3.0 mg), the reaction

buffer containing 40 mM HEPES (pH 7.2), 100 nmol of $CdCl_2$, and 100 nmol of PC2

was used. For the assay using PC2-attached beads (3.0 mg), the reaction buffer

containing 40 mM HEPES (pH 7.2), 100 nmol of $CdCl_2$, and 25 or 200 nmol of GSH

was used. The prepared tube was immediately set to a rotator and incubated at room

temperature for 5 min. The assay was repeated with three independent reactions.

Quantification of metal(loid) binding to the ligand

For Protocol A (Fig. 3), which was used for most of the assay in this study, an aliquot

(720 μL) of the reaction mixture was immediately applied to an empty spin column

(EconoSpin, Ajinomoto Bio-Pharma, Osaka, Japan) which harbors a polyethylene

membrane. The affinity beads were removed from the mixture by immediately

centrifuging the column at 15,300 g for 10 sec. An aliquot (500 µL) of the flow-through

was added to 500 µL of 60% nitric acid (for poisonous metals analysis grade,

FUJIFILM Wako Pure Chemical, Osaka, Japan) and then diluted to 5 mL with MilliQ

water. Elemental concentrations in the diluted samples were determined by ICP-OES

(iCAP7400Duo, Thermo-Fisher Scientific, Waltham, MA), and then the residual (free)

metal(loid) quantity in the reaction mixture after the reaction were calculated for control

beads ($M_{[control\ filtrate]}$) and ligand-attached beads ($M_{[ligand\ filtrate]}$), respectively (Fig. 3).

The amount of the metal(loid) that interacted with the ligand attached to the beads

($M_{[binding]}$) was obtained by subtracting $M_{[ligand\ filtrate]}$ from $M_{[control\ filtrate]}$ (Fig. 3). The

metal (loid)/ligand binding ratio was calculated by dividing $M_{[binding]}$ by the ligand

amount in the reaction mixture.

For Protocol B (Fig. 3), the reaction between Cd and the beads was examined. The

whole reaction mixture (5 mL) was centrifuged at 4,255 g for 1 min immediately after

the reaction. The supernatant was quickly but carefully removed by a vacuum aspirator

not to disturb the bead pellet. The beads were resuspended with 1 mL of 60% nitric acid.

The resuspension and another 1 ml of 60% nitric acid for rinse were transferred to a

10mL PTFE tube for acid digestion at 140 °C for 3 h. The digested samples were filled

up to 5 ml with 2% nitric acid, and their Cd concentrations were determined by

ICP-OES. The quantity of Cd binding to control beads ($M_{[control\ beads]}$) and to

ligand-attached beads ($M_{[ligand\ beads]}$) after the reaction was calculated, respectively (Fig.

3). Then, the amount of Cd interacted with the ligand attached to the beads ($M_{[binding]}$)

was obtained by subtracting $M_{[control\ beads]}$ from $M_{[ligand\ beads]}$ (Fig. 3).

Author contribution statement

SU, KN, and MK designed the experiments and prepared the manuscript. SU, KN, FN,

and YOt conducted the experiments. SU, KN, YOh, RN, YT, and MK analyzed the data.

All authors discussed the results and approved the manuscript.

Acknowledgments

We thank Prof. Tohru Nagamitsu (Kitasato University) and Dr. Mitsuru Abo (Meiji

University) for helpful advice and discussion. This work was partly supported by the

Japan Society for the Promotion of Science (Grant nos. 18K05377, 21K05330 to SU).

References

1 Waldron KJ, Rutherford JC, Ford D & Robinson NJ (2009) Metalloproteins and metal

sensing. *Nature* **460**, 823–30.

2 Andreini C, Bertini I, Cavallaro G, Holliday GL & Thornton JM (2008) Metal ions in

biological catalysis: from enzyme databases to general principles. *J Biol Inorg Chem*

13, 1205–18.

3 Dupont CL, Butcher A, Valas RE, Bourne PE & Caetano-Anollés G (2010) History of

biological metal utilization inferred through phylogenomic analysis of protein

structures. *Proc Natl Acad Sci USA* **107**, 10567–10572.

4 Dupont CL, Yang S, Palenik B & Bourne PE (2006) Modern proteomes contain

putative imprints of ancient shifts in trace metal geochemistry. *Proc Natl Acad Sci*

USA **103**, 17822–17827.

5 Hitomi Y, Outten CE & O'Halloran TV (2001) Extreme zinc-binding thermodynamics

of the metal sensor/regulator protein, ZntR. *J Am Chem Soc* **123**, 8614–8615.

6 Outten CE & O'Halloran TV (2001) Femtomolar sensitivity of metalloregulatory

proteins controlling zinc homeostasis. *Science* **292**, 2488–2492.

7 Rae T, Schmidt P, Pufahl R, Culotta V & O'Halloran TV (1999) Undetectable

intracellular free copper: The requirement of a copper chaperone for superoxide

dismutase. *Science* **284**, 805–808.

8 Colvin RA, Holmes WR, Fontaine CP & Maret W (2010) Cytosolic zinc buffering and

muffling: their role in intracellular zinc homeostasis. *Metallomics* **2**, 306–317.

9 Finney LA & O'Halloran TV (2003) Transition metal speciation in the cell: insights

from the chemistry of metal ion receptors. *Science* **300**, 931–936.

10 Hogstrand C, Kille P, Nicholson RI & Taylor KM (2009) Zinc transporters and

cancer: a potential role for ZIP7 as a hub for tyrosine kinase activation. *Trends Mol*

Med **15**, 101–111.

11 Bashir K, Rasheed S, Kobayashi T, Seki M & Nishizawa NK (2016) Regulating

subcellular metal Homeostasis: The key to crop improvement. *Front Plant Sci* **7**,

1192.

12 Clemens S, Deinlein U, Ahmadi H, Höreth S & Uraguchi S (2013) Nicotianamine is

a major player in plant Zn homeostasis. *BioMetals* **26**, 623–632.

13 Ghssein G, Brutesco C, Ouerdane L, Fojcik C, Izaute A, Wang S, Hajjar C, Lobinski

R, Lemaire D, Richaud P, Voulhoux R, Espaillat A, Cava F, Pignol D,

Borezee-Durant E & Arnoux P (2016) Biosynthesis of a broad-spectrum

nicotianamine-like metallophore in *Staphylococcus aureus*. *Science* **352**, 1105–1109.

14 Howden R, Goldsbrough PB, Andersen CR & Cobbett CS (1995)

Cadmium-sensitive, *cad1* mutants of *Arabidopsis thaliana* are phytochelatin

deficient. *Plant Physiol* **107**, 1059–1066.

15 Ha SB, Smith AP, Howden R, Dietrich WM, Bugg S, O'Connell MJ, Goldsbrough

PB & Cobbett CS (1999) Phytochelatin synthase genes from Arabidopsis and the

yeast *Schizosaccharomyces pombe*. *Plant Cell* **11**, 1153–1164.

16 Uraguchi S, Tanaka N, Hofmann C, Abiko K, Ohkama-Ohtsu N, Weber M, Kamiya

T, Sone Y, Nakamura R, Takanezawa Y, Kiyono M, Fujiwara T & Clemens S (2017)

Phytochelatin synthase has contrasting effects on cadmium and arsenic accumulation

in rice grains. *Plant Cell Physiol* **58**, 1730–1742.

17 Fischer S, Kühnlenz T, Thieme M, Schmidt H & Clemens S (2014) Analysis of plant

Pb tolerance at realistic submicromolar concentrations demonstrates the role of

phytochelatin synthesis for Pb detoxification. *Environ Sci Technol* **48**, 7552–7559.

18 Tennstedt P, Peisker D, Böttcher C, Trampczynska A & Clemens S (2009)

Phytochelatin synthesis is essential for the detoxification of excess zinc and

contributes significantly to the accumulation of zinc. *Plant Physiol* **149**, 938–948.

19 Clemens S, Kim EJ, Neumann D & Schroeder JI (1999) Tolerance to toxic metals by

a gene family of phytochelatin synthases from plants and yeast. *EMBO J* **18**,

3325–3333.

20 Kondo N, Imai K, Isobe M, Goto T, Murasugi A, Wada-Nakagawa C & Hayashi Y

(1984) Cadystin a and b, major unit peptides comprising cadmium binding peptides

induced in a fission yeast ----- separation, revision of structures and synthesis.

Tetrahedron Lett **25**, 3869–3872.

21 Vatamaniuk OK, Bucher EA, Ward JT & Rea PA (2001) A new pathway for heavy

metal detoxification in animals. Phytochelatin synthase is required for cadmium

tolerance in *Caenorhabditis elegans*. *J Biol Chem* **276**, 20817–20820.

22 Meister A (1991) Glutathione deficiency produced by inhibition of its synthesis, and

its reversal; Applications in research and therapy. *Pharmacol Therapeut* **51**,

155–194.

23 Wood BA & Feldmann J (2012) Quantification of phytochelatins and their

metal(loid) complexes: critical assessment of current analytical methodology. *Anal*

Bioanal Chem **402**, 3299–3309.

24 Raab A, Meharg AA, Jaspars M, Genney DR & Feldmann J (2003)

Arsenic–glutathione complexes—their stability in solution and during separation by

different HPLC modes. *J Anal Atom Spectrom* **19**, 183–190.

25 Clemens S (2019) Metal ligands in micronutrient acquisition and homeostasis. *Plant*

Cell Environ **42**, 2902–2912.

26 Cazalé AC & Clemens S (2001) *Arabidopsis thaliana* expresses a second functional

phytochelatin synthase. *FEBS Lett* **507**, 215–219.

27 Grill E, Winnacker EL & Zenk MH (1987) Phytochelatins, a class of

heavy-metal-binding peptides from plants, are functionally analogous to

metallothioneins. *Proc Natl Acad Sci USA* **84**, 439–443.

28 Badyal JP, Cameron AM, Cameron NR, Coe DM, Cox R, Davis BG, Oates LJ, Oye

G & Steel PG (2001) A simple method for the quantitative analysis of resin bound

thiol groups. *Tetrahedron Lett* **42**, 8531–8533.

29 Schmöger ME, Oven M & Grill E (2000) Detoxification of arsenic by phytochelatins

in plants. *Plant Physiol* **122**, 793–801.

30 Uraguchi S, Ohshiro Y, Otsuka Y, Tsukioka H, Yoneyama N, Sato H, Hirakawa M,

Nakamura R, Takanezawa Y & Kiyono M (2020) Selection of agar reagents for

medium solidification is a critical factor for metal(loid) sensitivity and ionomic

profiles of *Arabidopsis thaliana*. *Front Plant Sci* **11**, 503.

31 Verret F, Gravot A, Auroy P, Leonhardt N, David P, Nussaume L, Vavasseur A &

Richaud P (2004) Overexpression of *AtHMA4* enhances root-to-shoot translocation

669 of zinc and cadmium and plant metal tolerance. *FEBS Lett* **576**, 306–312.

670 32 Semane B, Cuypers A, Smeets K, Belleghem FV, Horemans N, Schat H &

671 Vangronsveld J (2007) Cadmium responses in *Arabidopsis thaliana*: glutathione

672 metabolism and antioxidative defence system. *Physiol Plant* **129**, 519–528.

673 33 Li Y, Nguyen M, Baudoin M, Vendier L, Liu Y, Robert A & Meunier B (2019) Why

674 is tetradentate coordination essential for potential copper homeostasis regulators in

675 Alzheimer's Disease? *Eur J Inorg Chem* **2019**, 4712–4718.

676 34 Fasae KD, Abolaji AO, Faloye TR, Odunsi AY, Oyetayo BO, Enya JI, Rotimi JA,

677 Akinyemi RO, Whitworth AJ & Aschner M (2021) Metallobiology and therapeutic

678 chelation of biometals (copper, zinc and iron) in Alzheimer's disease: Limitations,

679 and current and future perspectives. *J Trace Elem Med Bio* **67**, 126779.

680 35 Kenney GE, Dassama LMK, Pandelia ME, Gizzi AS, Martinie RJ, Gao P, DeHart CJ,

681 Schachner LF, Skinner OS, Ro SY, Zhu X, Sadek M, Thomas PM, Almo SC,

682 Bollinger JM, Krebs C, Kelleher NL & Rosenzweig AC (2018) The biosynthesis of

683 methanobactin. *Science* **359**, 1411–1416.

684 36 Fahey RC (2013) Glutathione analogs in prokaryotes. *Biochim Biophys Acta Gen*

685 *Subj* **1830**, 3182–3198.

686 37 Soboll S, Gründel S, Harris J, Kolb-Bachofen V, Ketterer B & Sies H (1995) The

687 content of glutathione and glutathione S-transferases and the glutathione peroxidase

688 activity in rat liver nuclei determined by a non-aqueous technique of cell

689 fractionation. *Biochem J* **311**, 889–894.

690 38 Sun X, Shih AY, Johannssen HC, Erb H, Li P & Murphy TH (2006) Two-photon

imaging of glutathione levels in intact brain indicates enhanced redox buffering in

developing neurons and cells at the cerebrospinal fluid and blood-brain interface. *J*

Biol Chem **281**, 17420–17431.

39 Gutiérrez-Alcalá G, Gotor C, Meyer AJ, Fricker M, Vega JM & Romero LC (2000)

Glutathione biosynthesis in Arabidopsis trichome cells. *Proc Natl Acad Sci* **97**,

11108–11113.

40 Irving H & Williams RJP (1953) The stability of transition-metal complexes. *J Chem*

Soc **3**, 3192–3210.

41 Martin RB & Edsall JT (1959) The association of divalent cations with glutathione 1.

J Am Chem Soc **81**, 4044–4047.

42 Hider RC & Kong XL (2011) Glutathione: a key component of the cytoplasmic

labile iron pool. *Biometals* **24**, 1179–1187.

43 Krężel A & Bal W (2004) Studies of zinc(II) and nickel(II) complexes of GSH,

GSSG and their analogs shed more light on their biological relevance. *Bioinorg*

Chem Appl **2**, 293–305.

44 Walsh MJ & Ahner BA (2013) Determination of stability constants of Cu(I), Cd(II)

& Zn(II) complexes with thiols using fluorescent probes. *J Inorg Biochem* **128**,

112–123.

45 Morgan MT, Nguyen LAH, Hancock HL & Fahrni CJ (2017) Glutathione limits

aquacopper(I) to sub-femtomolar concentrations through cooperative assembly of a

tetranuclear cluster. *J Biol Chem* **292**, 21558–21567.

46 Maitani T, Kubota H, Sato K & Yamada T (1996) The composition of metals bound

to class III metallothionein (phytochelatin and its desglycyl peptide) induced by

various metals in root cultures of *Rubia tinctorum*. *Plant Physiol* **110**, 1145–1150.

47 Oven M, Page JE, Zenk MH & Kutchan TM (2002) Molecular characterization of

the homo-phytochelatin synthase of soybean *Glycine max*: relation to phytochelatin

synthase. *J Biol Chem* **277**, 4747–4754.

48 Loscos J, Naya L, Ramos J, Clemente MR, Matamoros MA & Becana M (2006) A

reassessment of substrate specificity and activation of phytochelatin synthases from

model plants by physiologically relevant metals. *Plant Physiol* **140**, 1213–1221.

49 Li M, Barbaro E, Bellini E, Saba A, Toppi LS di & Varotto C (2020) Ancestral

function of the phytochelatin synthase C-terminal domain in inhibition of heavy

metal-mediated enzyme overactivation. *J Exp Bot* **71**, 6655–6669.

50 Uraguchi S, Sone Y, Ohta Y, Ohkama-Ohtsu N, Hofmann C, Hess N, Nakamura R,

Takanezawa Y, Clemens S & Kiyono M (2018) Identification of C-terminal regions

in *Arabidopsis thaliana* phytochelatin synthase 1 specifically involved in activation

by Arsenite. *Plant Cell Physiol* **59**, 500–509.

51 Kühnlenz T, Hofmann C, Uraguchi S, Schmidt H, Schempp S, Weber M, Lahner B,

Salt DE & Clemens S (2016) Phytochelatin synthesis promotes leaf Zn accumulation

of *Arabidopsis thaliana* plants grown in soil with adequate Zn supply and is essential

for survival on Zn-contaminated soil. *Plant Cell Physiol* **57**, 2342–2352.

52 Ruotolo R, Peracchi A, Bolchi A, Infusini G, Amoresano A & Ottonello S (2004)

Domain organization of phytochelatin synthase: functional properties of truncated

enzyme species identified by limited proteolysis. *J Biol Chem* **279**, 14686–14693.

735 53 Raab A, Feldmann J & Meharg AA (2004) The nature of arsenic-phytochelatin

736 complexes in *Holcus lanatus* and *Pteris cretica*. *Plant Physiol* **134**, 1113–1122.

737 54 Bluemlein K, Raab A, Meharg AA, Charnock JM & Feldmann J (2008) Can we trust

738 mass spectrometry for determination of arsenic peptides in plants: comparison of

739 LC–ICP–MS and LC–ES-MS/ICP–MS with XANES/EXAFS in analysis of

740 *Thunbergia alata*. *Anal Bioanal Chem* **390**, 1739–1751.

741 55 Sadi BBM, Vonderheide AP, Gong J-M, Schroeder JI, Shann JR & Caruso JA (2008)

742 An HPLC-ICP-MS technique for determination of cadmium–phytochelatins in

743 genetically modified *Arabidopsis thaliana*. *J Chromatogr B* **861**, 123–129.

744 56 Yen T, Villa JA & DeWitt JG (1999) Analysis of phytochelatin–cadmium complexes

745 from plant tissue culture using nano electrospray ionization tandem mass

746 spectrometry and capillary liquid chromatography/electrospray ionization tandem

747 mass spectrometry. *J Mass Spectrom* **34**, 930–941.

748 57 Chen L, Guo Y, Yang L & Wang Q (2007) SEC-ICP-MS and ESI-MS/MS for

749 analyzing *in vitro* and *in vivo* Cd-phytochelatin complexes in a Cd-hyperaccumulator

750 *Brassica chinensis*. *J Anal At Spectrom* **22**, 1403–1408.

751 58 Liu W-JJ, Wood BA, Raab A, McGrath SP, Zhao F-JJ & Feldmann J (2010)

752 Complexation of arsenite with phytochelatins reduces arsenite efflux and

753 translocation from roots to shoots in Arabidopsis. *Plant Physiol* **152**, 2211–2221.

754 59 Dorčák V & Krężel A (2003) Correlation of acid–base chemistry of phytochelatin

755 PC2 with its coordination properties towards the toxic metal ion Cd(II). *Dalton*

756 *Trans*, 2253–2259.

60 Vatamaniuk OK, Mari S, Lang A, Chalasani S, Demkiv LO & Rea PA (2004)

Phytochelatin synthase, a dipeptidyltransferase that undergoes multisite acylation

with gamma-glutamylcysteine during catalysis: stoichiometric and site-directed

mutagenic analysis of *Arabidopsis thaliana* PCS1-catalyzed phytochelatin synthesis.

J Biol Chem **279**, 22449–22460.

61 Song W-YY, Yamaki T, Yamaji N, Ko D, Jung K-HH, Fujii-Kashino M, An G,

Martinoia E, Lee Y & Ma JF (2014) A rice ABC transporter, OsABCC1, reduces

arsenic accumulation in the grain. *Proc Natl Acad Sci* **111**, 15699–15704.

62 Song W-YY, Park J, Mendoza-Cózatl DG, Suter-Grotemeyer M, Shim D,

Hörtensteiner S, Geisler M, Weder B, Rea PA, Rentsch D, Schroeder JI, Lee Y &

Martinoia E (2010) Arsenic tolerance in Arabidopsis is mediated by two ABCC-type

phytochelatin transporters. *Proc Natl Acad Sci* **107**, 21187–21192.

63 Rey NA, Howarth OW & Pereira-Maia EC (2004) Equilibrium characterization of

the As(III)–cysteine and the As(III)–glutathione systems in aqueous solution. *J Inorg*

Biochem **98**, 1151–1159.

64 Kala SV, Neely MW, Kala G, Prater CI, Atwood DW, Rice JS & Lieberman MW

(2000) The MRP2/cMOAT transporter and arsenic-glutathione complex formation

are required for biliary excretion of arsenic. *J Biol Chem* **275**, 33404–33408.

65 Clemens S (2019) Safer food through plant science: reducing toxic element

accumulation in crops. *J Exp Bot* **70**, 5537–5557.

66 Klapheck S, Fliegner W & Zimmer I (1994) Hydroxymethyl-phytochelatins

[γ-glutamylcysteine)n-serine] are metal-induced peptides of the *Poaceae*. *Plant*

Physiol **104**, 1325–1332.

67 Klapheck S, Schlunz S & Bergmann L (1995) Synthesis of phytochelatins and

homo-phytochelatins in *Pisum sativum* L. *Plant Physiol* **107**, 515–521.

68 Meuwly P, Thibault P, Schwan AL & Rauser WE (1995) Three families of thiol

peptides are induced by cadmium in maize. *Plant J* **7**, 391–400.

69 Merschel G & Bau M (2015) Rare earth elements in the aragonitic shell of

freshwater mussel *Corbicula fluminea* and the bioavailability of anthropogenic

lanthanum, samarium and gadolinium in river water. *Sci Total Environ* **533**, 91–101.

70 Tokumaru T, Ozaki H, Onwona-Agyeman S, Ofosu-Anim J & Watanabe I (2017)

Determination of the extent of trace metals pollution in soils, sediments and human

hair at e-waste recycling site in Ghana. *Arch Environ Contam Toxicol* **73**, 377–390.

 Figure legends

Fig. 1 Synthesis of phytochelatins 2 and 3 (PC2 and PC3) by Fmoc solid-phase peptide

synthesis using a 2-chlorotrityl chloride resin.

Fig. 2 Verification of the synthesized phytochelatins (A-F) and loading to the beads (G).

(A-F) Synthesized PC2 (A, C, E) and PC3 (B, D, F) were analyzed by total ion

chromatography (A, B), UV chromatography (C, D), and mass spectroscopy (E, F). (G)

Loadings of GSH, PC2, and PC3 onto TOYOPEARL AF-Fornyl-650 beads were carried

out under reductive amination conditions. The loadings were estimated by detecting the

amount of 2-nitro-5-thiobenzoate anion released from the reaction of Ellman's reagent

and thiol groups on the affinity beads.

Fig. 3 Schematic diagram describing the two protocols of the metal-ligand binding

assay using the ligand-attached beads and ICP. A reaction mixture for the standard

protocol was prepared in a 5 ml plastic tube containing the beads (normally 3.0 mg) and

a metal(loid)-of-interest (normally 200 nmol). The HEPES buffer (pH 7.2) was used as

a matrix instead of a phosphate buffer to avoid the potential interaction of the

metal(loid) and phosphate ions. The prepared tubes were incubated at room temperature

for 5 min or overnight with continuous rotation to avoid sedimentation of the beads.

After the incubation sample was immediately collected by Protocol A or Protocol B. In

Protocol A, a filtrate not containing beads was collected using a spin column and used

for the ICP quantification. In Protocol B, the beads were collected by centrifugation and

subjected to the ICP analysis.

Fig. 4 Comparison of Protocol A and Protocol B using Cd as a metal model. Total Cd

adsorption to the respective types of affinity beads was analyzed by ICP-OES after the

5-min incubation with 200 nmol Cd. Data represent means with SD from three

independent reactions for each combination. There is no significant difference between

the two protocols for all affinity beads tested ($P > 0.05$, Student's T-test). n.s., not

significant.

Fig. 5 Metal(loid)-ligand interaction assay using the ligand-attached affinity beads. A

reaction mixture containing 3.0 mg of each type of the beads and 200 nmol of each

metal(loid) species was incubated for 5 min or overnight at room temperature with

contentious rotation. After the incubation, the reaction mixture was immediately

subjected to spin column filtration (Protocol A). The metal(loid) content in the filtrate

was quantified by ICP-OES. Data represent means with SD from three independent

reactions for each combination. Means sharing the same letter are not significantly

different ($P < 0.05$, Tukey's HSD). (A) Non-specific metal(loid) adsorption to the

control (blank) beads. (B-D) Metal(loid) binding ratios of GSH-attached beads (B),

PC2-attached beads (C), and PC3-attached beads (D). n.d., not detected. n.a., not

analyzed.

Fig. 6 Metals competition assay using the ligand-attached beads. The reaction buffer

containing 20 nmol of $MnCl_2$, 40 nmol of $FeCl_2$, 2 nmol of $CuSO_4$, 20 nmol of $ZnSO_4$,

and 200 nmol of $CdCl_2$ was added to 3.0 mg of each type of the beads. The tube was

immediately set to a rotator and incubated at room temperature for 5 min. After the

incubation, the reaction mixture was immediately subjected to spin column filtration

(Protocol A). The metals content in the filtrate was quantified by ICP-OES. (A-C)

Proportions of metals binding to GSH (A), PC2 (B), and PC3 (C) on the beads,

calculated from total metal input to the reaction and the metals associated with each

ligand on the beads. Data represent means with SD from three independent reactions.

Means sharing the same letter are not significantly different ($P < 0.05$, Tukey's HSD).

(D) Profiles of the metals in the reaction mixture (input) and on the respective beads.

Data represent means from three independent reactions.

Fig. 7 PC2 and GSH competition assay using the ligand-attached beads. (A) The

reaction buffer containing 100 nmol of $CdCl_2$ and 100 nmol of PC2 was added to 3.0

mg of the GSH-attached beads. (B) The reaction buffer containing 100 nmol of $CdCl_2$,

and 25 or 200 nmol of GSH was added to 3.0 mg of the PC2-attached beads. (A, B) The

tube was immediately set to a rotator after adding the buffer and incubated at room

temperature for 5 min. After the incubation, the reaction mixture was immediately

subjected to spin column filtration (Protocol A). The Cd content in the filtrate was

quantified by ICP-OES. Data represent means with SD from three independent

reactions. *** $P < 0.001$, (Student's T-test). n.s. $P > 0.05$ (Dunnett's test).

2-Chrolotrityl chloride resin

Loading
Fmoc-Gly-OH
DIPEA, DCM

Method A
1) 2%DBU/DMF
2) Fmoc-Cys(Trt)-OH
DIPCI, HOBt,
DMF

Method B
1) 2%DBU/DMF
2) Fmoc-Glu-OtBu
DIPCI, HOBt,
DMF

1) Repeat A & B
2) 2%DBU/DMF
3) 20%HFIP/DCM
4) TFA/TIPS/H₂O
5) Preparative HPLC
Yield: 64%

PC2: (γ-Glu-Cys)₂-Gly, MW=539.58

1) Repeat A & B
2 times
2) 2%DBU/DMF
3) 20%HFIP/DCM
4) TFA/TIPS/H₂O
5) Preparative HPLC
Yield: 64%

PC3: (γ-Glu-Cys)₃-Gly, MW=771.84

A phytochelatin2 (PC2): (γ-Glu-Cys)₂-Gly

B phytochelatin3 (PC3): (γ-Glu-Cys)₃-Gly

G

	yield (nmol thiol mg⁻¹)	
	1st lot	2nd lot
GSH (n=1)	64.1	70.2
PC2 (n=2)	67.5	78.4
PC3 (n=3)	64.8	69.6

Mixture Preparation

beads · metal(loid)

40 mM HEPES (pH 7.2)

Incubation

5 min or O/N at RT with rotation

Sample Collection

Binding reaction

Protocol A · Collecting filtrate

Immediate filtration with a spin column

Protocol B · Collecting beads

Immediate centrifugation

ICP Quantification & Calculation

Dilution & ICP

Protocol A

$[M]_{binding} = [M]_{control\ filtrate} - [M]_{ligand\ filtrate}$

Acid extraction (140°C, 3h) & ICP

Protocol B

$[M]_{binding} = [M]_{ligand\ beads} - [M]_{control\ beads}$

A GSH beads

Metal binding to GSH (% of input)

B PC2 beads

Metal binding to PC2 (% of input)

C PC3 beads

Metal binding to PC3 (% of input)

D

Metal profiles (%)

Mn
Fe
Cu
Zn
Cd

Input   GSH beads   PC2 beads   PC3 beads

Net metals binding to