Title:

Development of affinity beads-based *in vitro* metal-ligand binding assay reveals

dominant cadmium affinity of thiol-rich small peptides phytochelatins beyond

glutathione

Author names and affiliations:

Shimpei Uraguchi[1], Kenichiro Nagai[2], Fumii Naruse[1], Yuto Otsuka[1], Yuka Ohshiro[1],

Ryosuke Nakamura[1], Yasukazu Takanezawa[1], Masako Kiyono[1]

[1] Department of Public Health, School of Pharmacy, Kitasato University, 5-9-1

Shirokane, Minato-ku, Tokyo 108-8641, Japan

[2] Medicinal Research Laboratories, School of Pharmacy, Kitasato University, 5-9-1

Shirokane, Minato-ku, Tokyo, 108-8641, Japan

*Corresponding author:

Prof. Dr. Masako Kiyono, telephone: +81-3-5791-6264, email address:

kiyonom@pharm.kitasato-u.ac.jp

Short running title:

Metal-thiol interaction by affinity beads assay

Abstract:

For a better understanding of metal-ligand interaction and its function in cells, we developed an easy, sensitive, and high-throughput method to quantify ligand-metal(loid) binding affinity under physiological conditions by combining ligand-attached affinity beads and inductively coupled plasma-optical emission spectrometry (ICP-OES). Glutathione (GSH) and two phytochelatins (PC2 and PC3, small peptides with different thiol numbers) were employed as model ligands and attached to hydrophilic beads. The principle of the assay resembles that of affinity purification of proteins in biochemistry: metals binding to the ligand on the beads and the rest in the buffer are separated by a spin-column and quantified by ICP-OES. The binding assay using the GSH-attached beads and various metal(loid)s suggested the different affinity of the metal-GSH interactions, in accordance with the order of the Irving–Williams series and the reported stability constants. The binding assay using PC2 or PC3-attached beads suggested positive binding between PCs and Ni(II), Cu(II), Zn(II), Cd(II), and As(III) in accordance with the number of thiols in PC2 and PC3. We then conducted the competition assay using Cd(II), Mn(II), Fe(II), Cu(II), and Zn(II) and the results suggested a better binding affinity of PC2 with Cd(II) than with the essential metals. Another competition assay using PC2 and GSH suggested a robust binding affinity between PCs and Cd(II) compared to GSH and Cd(II). These results suggested the dominance of PC-Cd complex formation *in vitro*, supporting the physiological importance of PCs for the detoxification of cadmium *in vivo*. We also discuss the further potential application of the assay.

Abbreviations:

DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene; DIPCI, *N,N'*-diisopropylcarbodiimide; DMF,

N,N-dimethylformamide;   GSH,   glutathione;   HEPES,

4-(2-hydroxyethyl)-1-piperazineethanesulfonic   acid;   HFIP,

1,1,1,3,3,3-hexafluoro-2-propanol; HOBt, hydroxybenzotriazole; ICP-OES, inductively

coupled plasma-optical emission spectrometry, ICT, isothermal titration calorimetry;

LC-ESI-MS,   liquid   chromatography-electrospray   ionization   mass-spectrometry,

LC-ICP-MS, liquid chromatography-inductively coupled plasma mass-spectrometry;

NA, nicotianamine; NMR, nuclear magnetic resonance; PC, phytochelatin; PCS,

phytochelatin synthase; TFA, trifluoroacetic acid; TIPS, triisopropylsilane.

Introduction

The essential metals are utilized by cells as cofactors for many proteins [1]. A bioinformatics study estimated that approximately 40% of the enzymes with known three-dimensional structures required metals at their catalytic centers [2]. The proportions of essential metals associated with enzymes are 6–9% of the analyzed enzymes for zinc (Zn), iron (Fe), and manganese (Mn), respectively, and 1% for copper (Cu). These metals are preferentially utilized in respective classes of Enzyme Commission (EC): for instance, Fe is mostly utilized by oxidoreductases (EC1) and Zn-containing enzymes are prominently enriched in hydrolases (EC3)[2]. Zn also plays structural roles in the Zn-binding proteins containing zinc-finger domains, suggesting the importance of Zn in transcriptional and post-translational regulations, especially in eukaryotes [3,4].

Meanwhile, the essential heavy metals in the cells are tightly regulated to avoid their potential toxicity. The studies on bacteria and yeast suggest that there is practically no cytosolic pool of free Cu and Zn ions under steady-state conditions [5–7]. The essential heavy metals imported to the cells are preferentially delivered to the target proteins requiring respective metals, and the rest are "buffered", bound to various small-molecular ligands as well as proteins like metallothionein in mammals [8,9]. Under conditions with excess metals, the biological chelators then function in "muffling", chelating the excess metal ions in the cell and avoiding cytotoxicity [8,9].

Among such metabolites, GSH is believed as a major metal-chelating small molecule especially in mammalian cells [8,10]. In addition, many diverse

small-molecular ligands have been discovered as physiologically significant metal

chelators in bacteria, yeast, worm, and higher plants [11–13]. (Phyto)siderophores and

its precursor nicotianamine (NA) are typical examples that function as Fe and Zn

binding ligands, respectively. Other established examples of the small molecular

chelators are phytochelatins (PCs), cysteine-rich small peptides. The general structure

of PCs is (γ-Glu-Cys)n-Gly with n usually between 2 and 7 [14]. PCs are

non-ribosomally synthesized from GSH (γ-Glu-Cys-Gly) by phytochelatin synthases

(PCSs). Compared to (phyto)siderophores and NA that function as ligands for essential

heavy metals [12], PCs can be characterized as high-affinity chelators for a wide range

of toxic metal(loid)s such as cadmium (Cd) and arsenic (As). In plants, PCs play a

central role in the detoxification of various toxic metal(loid)s [15–19]. Interestingly, the

PC-mediated detoxification of toxic metal(loid)s is also demonstrated in the fission

yeast *Schizosaccharomyces pombe* [20], in which PCs were first discovered [21], and in

the nematode worm *Caenorhabditis elegans* [22]. These suggest the conserved

functions of PC/PCS for toxic metal(loid) detoxification in taxonomically distant

organisms.

The physiological significance of GSH and PCs in metal(loid) detoxification

is mostly demonstrated by phenotypes induced by GSH synthesis inhibitors or PC/GSH

synthesis genes disruption, which lead to metal(loid) hypersensitivity [15–17,22,23].

These results indicate that ligand-metal complex formation is a key process for the

PC/GSH-mediated toxic metal(loid) detoxification, however, there is limited

information on complex formation between PCs/GSH and metal(loid)s, especially under

physiologically relevant conditions. It appears first due to the technical difficulties

overall [24], and secondly, due to unfamiliarity with required analytical techniques

and/or limited access to the analytical devices especially for many molecular

metal-biologists. For example, the PCs/GSH-metal(loid) complex formation *in vitro* is

often examined by potentiometry, voltammetry, and nuclear magnetic resonance (NMR)

spectrometry [25]. Isothermal titration calorimetry (ITC) is also a promising approach,

providing detailed *in vitro* data of binding chemistry under physiological conditions, but

the device is expensive and not so widely available. Another disadvantage of these

analytical chemistry approaches for molecular metal-biologists is that it is technically

difficult to mimic a cellular matrix containing various metals and metabolites other than

the   targets,   which   may   also   be   potential   interactants.

Liquid-chromatography-inductively coupled plasma mass-spectrometry (LC-ICP-MS)

and liquid-chromatography-electrospray ionization mass-spectrometry (LC-ESI-MS)

enable direct detection of multiple metal-ligand complexes *in vitro* and *in vivo*, however,

the technical difficulty of analyzing the intact complex form remains the bottleneck

[26].

For these reasons, an alternative assay especially for metal biologists would

be a useful option to advance understanding of metal-ligand interaction in the cells. The

present study aimed to develop an ICP-based simple but quantitative method to evaluate

ligand-metal(loid) interaction under various physiological conditions. ICP-optical

emission spectrometry (OES) was used as an analytical tool. ICP-OES as well as

ICP-MS is a common device among metal-biologists, compared to the other analytical

techniques, and more importantly, ICP-based element analysis provides highly sensitive

and high-throughput quantification. We employed GSH and PCs as model ligands and

prepared ligand-attached affinity beads for the assay. The GSH-metal(loid)s binding

ratios derived from our method coincided with the reported values. With the established

assay, we examined PC and Cd(II) interactions under physiological conditions

containing GSH and several essential metals. The competitive assays demonstrated

preferential interaction of PCs and Cd(II), which supports the *in vivo* observations that

PCs play a central role in Cd detoxification in plants, fission yeast, and *C. elegans*.

Materials and methods

Synthesis of PC2 and PC3

PC2 and PC3 were synthesized by the Fmoc solid-phase peptide synthesis strategy [27]

and the purity and mass sizes were verified by total ion current-chromatography,

UV-spectroscopy, and mass-spectrometry. A full method was provided as supplemental

materials and methods (see Supplementary information). LC-MS was performed on an

Agilent1200 system (G1379B degasser, G1312A binary pump, G1329A auto liquid

sampler, G1316A thermostat column compartment, and G1315B diode array detector)

and a JEOL AccuTOF LC-plus T100LP mass spectrometer equipped with an ESI

interface. The acquisition mass range was *m/z* 100–1000. Mobile phase A was 0.1%

HCO_2H in H_2O, and mobile phase B was 0.1% HCO_2H in CH_3CN. Separations on the

LC-MS were performed on a CAPCELL CORE C18 (2.7 μm, 2.1 × 50 mm, SHISEIDO

Loading of GSH, PC2, or PC3 onto the affinity beads

TOYOPEARL AF-Formyl-650 (TOSOH, Tokyo, Japan) was selected as the basal beads.

TOYOPEARL AF-Formyl-650 (1 mL, 60 μmol/mL) was placed in a polypropylene

tube and washed with H_2O (1 mL, 1 min $\times$ 3). To the bead was added a solution of GSH

(18 mg, 60.0 μmol) in 0.1 M phosphate buffer (pH 7.0, 1 mL), and then sodium

cyanoborohydride (11 mg, 180 μmol) was added. The resulting mixture was agitated at

room temperature for 22 h. The solvents and soluble reagents were removed by

filtration, and the bead was washed with H_2O (1 mL, 1 min $\times$ 3), saturated sodium

chloride (1 mL, 1 min $\times$ 3), H_2O (1 mL, 1 min $\times$ 3), and MeOH (1 mL, 1 min $\times$ 3). PC2

or PC3 were loaded onto the affinity bead according to the loading procedure described

above. The loading of the resulting bead was determined by the amount of

2-nitro-5-thiobenzoate anion formed from the reaction of 5,5'-dithiobis(2-nitrobenzoic

acid), Ellman's reagent, with sulfhydryl groups on the bead. The affinity resin (10 mg)

was treated with 2.5 mM Ellman's reagent in MeOH (2 mL) and

N,N-diisopropylethylamine (5 µL) at room temperature for 30 min. The total volume

was adjusted to 5 mL by the addition of MeOH and measured spectrophotometrically at

412nm.

Single metal(loid)-beads assay

The interaction assay was conducted in a 5 mL Eppendorf tube containing 3.0 mg of the

beads (control, GSH, PC2, or PC3). The assay was repeated with at least three

replications. To evaluate the interaction between the ligand beads and various

metal(loid)s, 5 mL of 40 mM HEPES (pH 7.2) containing 200 nmol of $MnCl_2$, $FeCl_2$,

$(NH_4)_2Ni(SO_4)_2$, $CuSO_4$, $ZnSO_4$, $NaAsO_2$, or $CdCl_2$, was added to the 5 mL tube. The

tube was immediately set to a rotator (RT-50, TAITEC, Saitama, Japan) and incubated at

room temperature for 5 min at approximately 4 rpm. For Mn(II), Fe(II), Cu(II), and

As(III), overnight incubation was also tested with rotation. The metal reagents used in

this study were all Guaranteed Reagents grade purchased from Nacalai Tesque (Kyoto,

Japan) or FUJIFILM Wako Pure Chemical (Osaka, Japan). The reagent for $NaAsO_2$ was

USP grade, purchased from Merck (Darmstadt, Germany).

Multiple metals competition assay

The competition assay using multiple metals was conducted as described for the single

metal(loid)-beads assay, except for the metal composition in the reaction buffer. The

reaction buffer [5 mL of 40 mM HEPES (pH 7.2)] contained 20 nmol of $MnCl_2$, 40

nmol of $FeCl_2$, 2 nmol of $CuSO_4$, 20 nmol of $ZnSO_4$, and 200 nmol of $CdCl_2$ and was

added to 3.0 mg of the beads (control, GSH, PC2, or PC3) in the 5 mL tube. The tube

was immediately set to a rotator and incubated at room temperature for 5 min. The

assay was repeated with three independent reactions.

PC2 and GSH competition assay

Cd-binding competition between PC2 and GSH was examined by modifying the single

metal(loid)-beads assay. For the assay using GSH-attached beads (3.0 mg), the reaction

buffer containing 40 mM HEPES (pH 7.2), 100 nmol of $CdCl_2$, and 100 nmol of PC2

was used. For the assay using PC2-attached beads (3.0 mg), the reaction buffer

containing 40 mM HEPES (pH 7.2), 100 nmol of $CdCl_2$, and 25 or 200 nmol of GSH

was used. The prepared tube was immediately set to a rotator and incubated at room

temperature for 5 min. The assay was repeated with three independent reactions.

Quantification of metal(loid) binding to the ligand

For Protocol A (Fig. 2), which was used for most of the assay in this study, an aliquot

(720 µL) of the reaction mixture was immediately applied to an empty spin column

(EconoSpin, Ajinomoto Bio-Pharma, Osaka, Japan) which harbors a polyethylene

membrane. The affinity beads were removed from the mixture by immediately

centrifuging the column at 15,300 g for 10 sec. An aliquot (500 µL) of the flow-through

was added to 500 µL of 60% nitric acid (for poisonous metals analysis, FUJIFILM

Wako Pure Chemical, Osaka, Japan) and then diluted to 5 mL with MilliQ water.

Elemental concentrations in the diluted samples were determined by ICP-OES

(iCAP7400Duo, Thermo-Fisher Scientific, Waltham, MA). Emission wavelengths used

for the detection were 189.04, 206.200, 224.700, 226.502, 239.562, and 260.569 nm for

As, Zn, Cu, Cd, Fe, and Mn, respectively. Then the residual (free) metal(loid) quantity

in the reaction mixture after the reaction was calculated for control beads ($M_{[control\ filtrate]}$)

and ligand-attached beads ($M_{[ligand\ filtrate]}$), respectively (Fig. 2). The amount of the

metal(loid) that interacted with the ligand attached to the beads ($M_{[binding]}$) was obtained

by subtracting $M_{[ligand\ filtrate]}$ from $M_{[control\ filtrate]}$ (Fig. 2). The metal (loid)/ligand binding

ratio was calculated by dividing $M_{[binding]}$ by the ligand amount in the reaction mixture.

For Protocol B (Fig. 2), the reaction between Cd and the beads was examined. The

whole reaction mixture (5 mL) was centrifuged at 4,255 g for 1 min immediately after

the reaction. The supernatant was quickly but carefully removed by a vacuum aspirator

not to disturb the bead pellet. The beads were resuspended with 1 mL of 60% nitric acid.

The resuspension and another 1 ml of 60% nitric acid for rinse were transferred to a

10mL PTFE tube. The sample was digested in an open vessel at 140 °C for 3 h. The

digested samples were filled up to 5 ml with 2% nitric acid, and their Cd concentrations

were determined by ICP-OES. The quantity of Cd binding to control beads ($M_{[control}$

$_{beads]}$) and to ligand-attached beads ($M_{[ligand\ beads]}$) after the reaction was calculated,

respectively (Fig. 2). Then, the amount of Cd interacted with the ligand attached to the

beads ($M_{[binding]}$) was obtained by subtracting $M_{[control\ beads]}$ from $M_{[ligand\ beads]}$ (Fig. 2).

Results

Preparation of phytochelatin- and glutathione-attached beads

PCs are commercially available only on a 1 mg scale (Eurogentec, Seraing,

Belgium) as far as we searched, and it was a considerable cost disadvantage to our large

demand of PCs for the affinity beads preparation. Thus, we first synthesized a shorter

PC [(γ-Glu-Cys)$_2$-Gly, PC2] and a longer PC [(γ-Glu-Cys)$_3$-Gly, PC3] by a standard

Fmoc solid-phase peptide synthesis protocol (Supplementary Fig. S1). The two different

length PCs are common PC isoforms synthesized in *S. pombe*, *C. elegans,* and various

plant cells exposed to toxic metal(loid) ions such as Cd(II) and As(III) [14,17,20,22,28].

Fmoc-Gly-OH was loaded onto 2-chlorotrityl chloride resin (1.38 mmol/g, 200 mg).

Fmoc   protecting   group   was   removed   using   a   solution   of   2%

1,8-diazabicyclo[5.4.0]undec-7-ene   (DBU)   in   *N,N*-dimethylformamide   (DMF).

Condensation of Fmoc-Cys(Trt)-OH and Fmoc-Glu-O*t*Bu was performed using

hydroxybenzotriazole (HOBt) and *N,N′*-diisopropylcarbodiimide (DIPCI). The resulting

peptide was cleaved from the resin with 20% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP)

in CH$_2$Cl$_2$, and then deprotected with trifluoroacetic acid (TFA)/triisopropylsilane

(TIPS)/H$_2$O (95:2.5:2.5). After purification by preparative HPLC, PC2 and PC3 were

obtained in 64% yield (95.0 mg) and 64% yield (126 mg), respectively. Then we

analyzed the synthesized PC2 and PC3 by HPLC/MS to confirm the purity and exact

mass. The synthesized PC2 was eluted as a peak with a retention time of 0.75 min in

total ion current chromatogram (Fig. 1A) and 0.68 min in PDA chromatogram (Fig. 1C)

and was detected as a protonated ion [M + H]+ at *m/z* 540.14 (Fig. 1E). The synthesized

PC3 was eluted in a peak with a retention time of 1.68 min in total ion current

chromatogram (Fig. 1B) and 1.60 min in PDA chromatogram (Fig. 1D) and was

detected as a protonated ion [M + H]+ at *m/z* 772.12 (Fig. 1F).

Using the synthetic PC2, PC3, and commercially obtained GSH, we

attempted to prepare PC- or GSH-attached affinity beads (Fig. 1G). Hydrophilic beads

which are only modified with formyl groups were used as a carrier resin. GSH, PC2, or

PC3 was loaded onto the beads via reductive amination using $NaBH_3CN$. This reaction

was conducted twice independently to obtain two different preparations of the

respective ligand-attached beads. The yield was quantified by Ellman's reagent [29] and

was overall within a similar range (64.1 to 78.4 nmol thiol per mg beads) irrespective of

the ligands and preparations (Fig. 1G). The beads treated only with the reaction buffer

not containing GSH, PC2, or PC3 were prepared as control beads in the subsequent

binding assay.

Establishment of the assay protocol for metal(loid)-ligand interaction

We then designed and tested protocols for evaluating the ligand-metal(loid)

interactions using the prepared ligand-attached beads (Fig. 2). A reaction mixture for the

standard protocol was prepared in a 5 ml plastic tube containing 3.0 mg of the beads and

200 nmol of metal(loid)-of-interest. 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

(HEPES) buffer (pH 7.2) was used as a matrix instead of a phosphate buffer to avoid the

potential interaction of the metal(loid) and phosphate ions. The prepared tubes were

incubated at room temperature for 5 min or overnight with continuous rotation to avoid

sedimentation of the beads. Sample collection after the incubation was performed by

two different protocols: collecting a filtrate without beads using a spin column (Protocol

A) and collecting beads without a solution by centrifugation (Protocol B). Element

contents in the collected samples were quantified by ICP-OES.

For calculation of $[M]_{binding}$ [the metal(loid) binding to the ligand on the

beads], in Protocol A, $[M]_{ligand\ filtrate}$ [the metal(loid) amount in the filtrate from the

ligand-attached beads reaction] was subtracted from $[M]_{control\ filtrate}$ [the metal(loid)

amount in the filtrate from the control beads reaction] (Fig. 2). In Protocol B, $[M]_{control}$

$_{beads}$ [the metal(loid) amount in the control beads] was subtracted from $[M]_{ligand\ beads}$ [the

metal(loid) amount in the ligand-attached beads] to obtain $[M]_{binding}$ (Fig. 2).

The two protocols were tested with Cd(II) as a model metal. To compare the

accuracy of Protocol A and B, we first quantified total Cd adsorption to the control and

ligand-attached beads (Fig. 3). For Protocol A, $[Cd]_{ligand\ filtrate}$ and $[Cd]_{control\ filtrate}$ were

respectively subtracted from Cd(II) input to each reaction (200 nmol). For Protocol B,

$[Cd]_{control\ beads}$ and $[Cd]_{ligand\ beads}$ were directly used to assess total Cd adsorption to the

beads. Regardless of the types of the beads, the Cd adsorption to the beads did not

significantly differ between Protocol A and B (Fig. 3), suggesting that both protocols

can equally evaluate the metal binding to the beads as designed. Protocol A is more

convenient and practical because the separation of the beads and the buffer can be easily

and quickly achieved by a spin column centrifugation, whereas Protocol B requires

careful but quick removal of the buffer by pipetting after centrifugation. Taken together,

we decided to choose Protocol A for further experiments.

Affinity variation of GSH, PC2, and PC3 with various metal(loid)s

The experiment to test the two protocols using Cd(II) as a model metal

indicated the substantial chelation between the ligands and Cd(II) because all the

ligand-attached beads (GSH, PC2, and PC3) adsorbed nearly 3-times more Cd than the

control beads (Fig. 3). The results motivated us to further conduct a full-scale

experiment and we quantified GSH and PCs-mediated chelation of Mn(II), Fe(II), Ni(II),

Cu(II), Zn(II), As(III), and Cd(II) (Fig. 4). Note that As(III) does not have a charge at

physiological pH. 5-min incubation was applied for the initial experiment, and

overnight incubation was also examined for Mn(II), Fe(II), Cu(II), and As(III). The

experiment for Fig. 4 was carried out with the first lot beads, except for Cd(II) for

which the independently prepared second lot beads were also applied to examine

possible lot-specific effects.

We first checked the metal(loid) adsorption to the controls beads as

non-specific background adsorption of the assay (Fig. 4A). The presented values are

percentages of the metal(loid) binding to the beads to the total input to the reaction (200

nmol). Non-specific adsorption to the hydrophilic beads reached up to 30% of the input,

regardless of the tested metal(loid) species and incubation time, except for overnight

incubation with Fe(II) (nearly 100% of the input to the reaction was adsorbed to the

carrier beads). The result of the non-specific background adsorption indicates that the

experimental condition provides enough metal(loid)s left in the reaction buffer which

can interact with the ligand on the ligand-attached beads, except for overnight Fe

incubation.

Then we calculated the metal(loid) binding ratio of GSH (Fig. 4B), PC2 (Fig.

4C), and PC3 (Fig. 4D), dividing $[M]_{binding}$ (Fig. 2) by the molar number of the ligand in

the reaction. For GSH which harbors one cysteine, the metal-binding ratio of 0.5 was

referred to as a standard to indicate a binding stoichiometry of one metal to two GSH.

The results of 5-min incubation for Ni(II), Zn(II), and Cd(II) were close to 0.5 (Fig. 4B).

The ratio of 5-min incubation with Cu(II) showed a somewhat intermediate value (0.36).

Those of Mn(II) and Fe(II) were significantly lower even than this value of Cu(II),

suggesting a lower affinity of Mn(II) and Fe(II) with GSH. For As(III), the binding ratio

of 0.33 was referred to as a standard indicating a binding stoichiometry of one As(III) to

three GSH [30]. However, the ratio of As/GSH after 5-min incubation was 0.02, which

was far below the standard value of 0.33. To further examine the lower binding ratio of

Cu(II), Mn(II), Fe(II), and As(III), the incubation time was prolonged from 5 min to

overnight. The binding ratios of Mn(II) and As(II) with GSH after overnight incubation

was still quite lower than the standards 0.5 and 0.33, respectively (Fig. 4B). The

overnight incubation increased the binding ratio of Cu(II) and GSH from 0.36 (5-min)

to 0.62, both of which were within the range of the standard value of 0.5. The binding of

Fe(II) with GSH after the overnight incubation was not detected due to the huge

background Fe adsorption to the carrier beads (Fig. 4A).

The metal(loid) binding ratio with PC2 is shown in Fig. 4C. For PC2, which

harbors two cysteines, the binding ratios of 0.5 and 1.0 as standards were referred to

indicate 1:2 and 1:1 binding of metal(loid) and PC2, respectively. The binding ratios of

Ni(II), Cu(II) (overnight), Zn(II), and Cd(II) were statistically similar and were close to

1.0 (Fig. 4C). The ratio obtained from 5-min incubation with Cu(II) was again a bit

lower than that from overnight incubation, but the difference was smaller than the case

of GSH (Fig. 4B). Binding ratios with PC2 for Mn(II) (5-min and overnight) and Fe(II)

(5-min) were far below 1.0. The ratio for As/PC2 was close to 0.5 after overnight

incubation, suggesting 1:2 binding of As(III) and PC2, whereas the result of 5-min

incubation was much lower.

347   The metal(loid) binding ratio with PC3 was shown in Fig. 4D. PC3 contains

three cysteines, and the binding ratios of 1.0 and 1.5 as standards were referred to

indicate 1:1 and 3:2 binding of metal(loid) and PC3, respectively. The binding ratios of

Ni(II), Cu(II) (5-min and overnight), Zn(II), and Cd(II) were around 1.5. It should be

noted that there was no statistical difference between the incubation periods for Cu(II).

The ratios for Mn(II) (5-min) and Fe(II) (5-min) were 0.67 and 1.04, respectively, which

were significantly lower than the ratios for other divalent metals tested. The binding

ratio of As(III) and PC3 (overnight) was 0.98, indicating the 1:1 binding relationship,

however, the 5-min incubation again resulted in a lower value (0.22).

Effects of different beads preparation

358   Conducting the experiments to produce the most results for Fig. 3 and Fig. 4,

we used a large portion of the first preparation of the ligand-attached beads. Thus, we

prepared a second lot of the respective beads. The second preparation resulted in similar

yields however, the yield was overall a bit higher (Fig. 1G). We examined the possible

effects of the different lots on the metal(loid)-ligand-binding assay, using Cd(II) as a

model again. Background non-specific Cd adsorption (Fig. 4A), and binding ratios with

GSH (Fig. 4B), PC2 (Fig. 4C), and PC3 (Fig. 4D) were not statistically different

between the first and second preparations by Tukey's HSD. In addition, the results of

the Student's T-test were also not significant ($P > 0.05$). These results suggest that

differently prepared ligand-attached beads can be equally used in the binding assay and

we decided to use the second lot for further experiments.

Binding competition between Cd and essential metals

The results of the single metal(loid)-ligand assay (Fig. 4) suggested different

affinity of the small thiol compounds to the various essential metals and toxic

metal(loid)s. We next planned to explore functional differences of the different length

thiols, GSH, PC2, and PC3, using the developed ligand-attached beads system. PCs are

believed to function in chelating free toxic metal(loid)s in the cytosol. However,

competition over PCs' binding sites can be expected in the cytosol between the toxic

metals and co-existing essential heavy metals. Thus, we conducted a competition assay

using Cd(II) as a model. Cd(II), not As(III), was selected because Cd(II) binding was

stably detected by the 5-min incubation, whereas As(III) required the overnight

incubation (Fig. 4). In this competition assay, Cd(II) (200 nmol) as well as essential

metals [Mn(II) (20 nmol), Fe(II) (40 nmol), Cu(II) (2 nmol), and Zn(II) (20 nmol)] were

added to the reaction. The ratio of the supplemented metals was derived from a rough

calculation based on the total element concentrations in leaves of the plant model

Arabidopsis thaliana [31–33].

As a result of the competition assay, Mn(II) and Fe(II) exhibited inferior

affinity to GSH, PC2, and PC3 compared to Cu(II), Zn(II), and Cd (II) (Fig. 5A, B, C).

Smaller fractions of the Mn(II) and Fe(II) input (~30%) were associated with the

ligands, and approximately 60% or more of the Cu(II), Zn(II), and Cd(II) input were

associated with the ligands. We then calculated profiles of the metals binding to GSH,

PC2, and PC3, and compared them with the profile of the input (Fig. 5D). Reflecting

the lower affinity with the ligands, the fractions of Mn(II) and Fe(II) binding to the

ligands were smaller than those of the input. In contrast, the Cd(II) binding to the

ligands was more than 80% of the total metals captured by the ligands, which was larger

than the input supplemented Cd(II) fraction (73%). These results further support the

results of the single-metal binding assay (Fig. 4), suggesting a preference of Cd(II) in

binding to the thiol ligands, compared to the tested essential metals.

Cd-binding competition between GSH and PCs

Next, we conducted another competition assay to examine a possible

preference of PC and Cd(II) interaction (Fig. 6). First, the binding of 100 nmol Cd(II)

and GSH beads competed with or without 100 nmol PC2 added as a free form

competitor in the reaction mixture. The Cd(II) binding to GSH beads under the

condition without the competitor PC2 was 55% of the total Cd(II) input (Fig. 6A), but

the addition of PC2 as the competitor to the reaction significantly decreased the Cd(II)

binding to GSH beads to 16% of the total Cd(II) input (Fig. 6A). Then, the inverted

assay conditions were also tested (Fig. 6B): Binding of 100 nmol Cd(II) by PC2 beads

competed with 25 nmol GSH or 200 nmol GSH supplemented to the reaction mixture in

a free form. In the absence of GSH, 55% of the input Cd(II) was bound with PC2 on the

beads, and the addition of 25 nmol nor 200 nmol GSH to the mixture as a competitor

did not change the Cd(II) binding profile to PC2 on the beads. The results overall

suggest the superior binding affinity of Cd(II) and PCs compared to GSH.

Discussion

Advantages of our metal-ligand binding assay

Physiological roles of metal-ligand interactions have been attracting attention

because of their significance in the essential metal homeostasis and toxic metal(loid)

detoxification in bacteria, plants, and humans [26,34–36]. The first goal of this study

was to establish a new assay system practically feasible for biologists interested in

evaluating metal-biological ligand interactions under cellular-mimicking conditions.

The idea of our assay resembles that of protein affinity purification protocols such as

immunoprecipitation in biochemistry. Using ICP as a readout tool is the first unique

advantage of our assay. ICP-OES and ICP-MS are widely used by metal biologists.

Moreover, the application of ICP enables rapid, highly sensitive, and stable

quantification of a large sample set. Note that a standard ICP protocol takes only 3-4

min for a single sample measurement of even more than 10 elements at ppb (ICP-OES)

or sub-ppb levels (ICP-MS). In addition, a spin column-based efficient separation of the

beads by centrifugation is also a key advantage of the assay. The spin-column procedure

is very familiar to molecular biologists and enables immediate termination of the

reaction of multiple samples. Besides technical advantages, it should be noted that our

assay system allows variable assay conditions which mimic cellular environments. Thus,

we can easily examine the effects of various co-existing metals and endogenous

metal-interactants (Figs. 4-6). These features are unique to our assay and imply the

broad utility of the protocol.

A few limitations exist in our assay: the assay cannot provide direct complex

form information and stability constants unlike the other conventional methods such as

potentiometry, voltammetry, NMR, ITC, and ESI-MS. However, the proposed assay has

significant advantages in terms of flexibility of assay conditions, high-throughput, and

technical feasibility especially for molecular biologists who are interested in

metal-ligand interactions under cellular environments.

GSH and metals interaction as a test model of the assay

In the present study, we employed GSH and two different length PCs (PC2

and PC3) as the model ligands for our assay (Figs. 1-3). First, the reliability of the assay

was verified by assessing GSH-metal interactions (Fig. 4B). The affinity between GSH

and various metals is well characterized by conventional analytical methods. GSH is

believed as one of the most possible ligands in the eukaryotic cells to form

metal-complexes through its sulfhydryl group [37]. The abundance in the cells is

another advantage of GSH as a metal-ligand: GSH concentrations in eukaryotic cells

(ranging from 0.1 mM to 8 mM [38–40]) are likely substantial enough to bind free

metals which would exist at much lower in cells.

The results of binding experiments between GSH and well-studied cations

[Mn(II), Fe(II), Ni(II), Cu(II), Zn(II), and Cd(II)] coincide with the reported results and

the order of the Irving–Williams series (Mn<Fe<Co<Ni<Cu, Zn) [41] which describes

the complex stability between divalent metal ions and organic ligands (note that Cd is

not included in the series, and Co is not tested in this study). For example, our assay

suggests a weak interaction of Mn(II) and Fe(II) with GSH (Fig. 4B), supporting the

models suggested in the previous studies: Mn(II)-GSH and Fe(II)-GSH interactions are

1:1 with markedly lower stability constant (log K_1 values) of 2.7 and 5.1, respectively

[42,43]. In contrast, stronger affinity with GSH is suggested for highly reactive metals

up in the Irving–Williams series [Ni(II), Cu(II), and Zn(II)] as well as Cd(II) (Fig. 4B).

The reported stability constants (log β values) for these metals are around 10 or much

higher even under physiological pH conditions [44,45]. In our study, the metal/GSH

ratio of around 0.5 was obtained for Ni(II), Cu(II), Zn(II), and Cd(II) (Fig. 4B),

suggesting 1:2 metal-GSH complex formation as previously reported [44,45]. For

Cu(II), there would be other possibilities. The Cu/GSH ratios were slightly yet

significantly lower than 0.5 under 5 min-incubating conditions, and overnight

incubation increased the ratio (Fig. 4B). This variation would be explained by the

controversial nature of Cu-GSH complex formation. Previous studies imply Cu(II) and

GSH affinity with 1:2 and 1:1 stoichiometry [45] and in addition, the formation of

$Cu(I)_4(GS)_6$ has been suggested [46]. Another note for the Cu-assay is that we applied

Cu(II) to the assay, not Cu(I), because of its low solubility and extremely high

background adsorption to the beads in the preliminary experiment using CuCl (data not

shown). Cu(II) is converted to Cu(I) under the reducing cellular conditions [46], which

may explain the increased interaction of Cu and GSH by the longer reaction period (Fig.

4B). Therefore, taken together with these points, our results for Cu(II) cannot suggest an

exact stoichiometry of Cu(II)-GSH interaction, but at least suggest a stronger affinity

with GSH under physiological (cellular) conditions compared to Mn(II) and Fe(II) (Fig.

4B).

480   Nevertheless, the results of GSH beads (Fig. 4B) overall demonstrate that

481   under physiological pH conditions, the assay can differentiate the affinity of metals,

482   corresponding to the order of the Irving–Williams series and the established stability

483   constants.

485   *Preferential interaction between phytochelatin and Cd*

486   Several metabolites other than GSH are also suggested as biological metal

ligands. PC is an established example that mainly functions in Cd detoxification

through Cd-PC complex formation in higher plants, *S. pombe*, and *C. elegans*

[16,17,20,22]. Due to its exceeding significance in Cd detoxification, the PCS activation

mechanism has been extensively examined using recombinant proteins, heterologous

expression systems in *S. pombe*, plant cell cultures, and plant seedlings. In brief, PCS

activity is triggered substantially by various environmentally relevant ions [e.g. As(III),

As(V), Ag(I), Cd(II), Hg(II), and Pb(II)] and the excess of a few physiologically

essential metal ions [Cu(II) and Zn(II)] as well, irrespective of their valency and

positions in the periodic table [14,17–19,28,47–49]. On the other hand, several elements

essential for plant nutrition [K(I), Mg(II), Ca(II), Mn(II), and Ni(II)] even in their

excess levels do not activate PCSs [48,49]. Whether Fe activates PCSs is still

controversial [49]. Such metal-(loid) PCS activation mechanism has been a significant

question. Recent studies suggest that the different regions of the C-terminal regulatory

domain of PCSs are required for the metal(loid)-specific activation [17,50–53],

although the detailed enzymatic action of PCS remains elusive mainly due to the lack of

the crystal structure of the full-length PCS.

In contrast to the activation mechanism, much less is known about the subsequent

metal(loid)-PC complex formation. Only a few studies reported the detection of Cd-PC

and As-PC complexes in plant extracts or *in vitro* samples using LC-ESI-MS or

LC-ICP-MS [54–59]. Thus, the second purpose of the present study is to examine

preferences of metal(loid)-PC complex formation using our binding assay, mainly

focusing on Cd-PC interaction. The assay using PC2 beads and a single metal species

(Fig. 4C) suggests a 1:1 interaction between Cd(II) and PC2. The 3:2 interaction

between Cd(II) and PC3 is suggested in accordance with the three SH residues in PC3

(Fig. 4D). These interaction ratios are in line with the suggested values obtained by

ESI-MS analysis of the plant samples and *in vitro* ITC analysis [57,60]. Similarly, our

assay suggests 1:1 interaction with PC2 and 3:2 interaction with PC3 were also

suggested for Ni(II), Cu(II), and Zn(II). Fe(II)-PC2 interaction is suggested weaker (Fig.

4C), but Fe(II) is likely to interact with PC3 at the ratio of 1:1 (Fig. 4D). These results

suggest a potential of Fe(II), Ni(II), Cu(II), and Zn(II) interacting with PCs, aside from

their PCS activation ability. This implies that Cd(II) and these metals may compete for

the PCs' binding sites. However, the Cd-PC complexes are the major form detected in

plants exposed to Cd stress [56–58], suggesting that Cd(II)-PC interaction is dominant

over the competitive essential metals.

A series of competitive assays were conducted to examine the dominance of

Cd-PC interaction (Figs. 5-6). The competition assay between Cd(II) and several plant

essential metals suggests the preferential binding affinity of PCs with Cd(II) over the

tested essential metals (Fig. 5). The result supports the hypothesis and *in planta*

observation of the dominance of Cd-PCs formation [56–58]. In plant cellular conditions,

sub-millimolar levels of GSH exist as another potent Cd(II) chelator other than PCs.

Thus, we examined the Cd(II) binding preference between PC2 and GSH (Fig. 6). The

assays clearly demonstrate the marked preference of Cd-PC2 interaction over Cd-GSH

(probably Cd-GS$_2$). The Cd-PC2 interaction was not interfered with even by the excess

level of free GSH (Fig. 6B), which supports the reported higher stability constant of

Cd-PC2 than Cd-GS$_2$ [45,61]. Overall, our competition assays suggest the robust and

preferential interaction of PC and Cd(II) over other co-existing essential metals and

GSH. It should be noted that the actual mechanism of PC-metal(loid) complex

formation is still unclear. A previous study suggested a possibility that the pre-formed

Cd-GS$_2$ complex interacts with the C-terminal regulatory domain of the PCS and

activates the transpeptidase activity to induce PC synthesis [62]. The previous study

also proposed that a Cd ion is transferred from the Cd-GS$_2$ complex to the PCS protein

and activates PC synthesis [62]. Thus, it can be postulated that Cd-PC complex

formation (partly) occurs during the process of PC synthesis [62]. Nevertheless, our

competition assays (Figs. 5-6) suggest that the formed Cd-PC complex would be stable

enough in the cytosol, not interfered by GSH and other essential metals. This stability

would be significant to ensure subsequent transport of the Cd-PC complex from the

cytosol to the vacuole, which is mediated by the tonoplast ABC transporters [63,64].

Interaction between thiols and As(III)

Another promising partner of PC other than Cd(II) is As(III). In plants, As(III)

stress also induces PC synthesis, and the subsequent As(III)-PC complex formation and

vacuolar sequestration by ABC transporters are suggested to be crucial for As(III)

detoxification [17,51,65,66]. Formation of As(III)-(PC2)$_2$ and As(III)-PC3 is reported *in*

vitro and *in planta* [54,55,59]. The binding stoichiometry is in accordance with the

number of the thiol groups in PCs [note that three SH residues are required to catch one

As(III)]. Our results of As(III)-PC beads interaction assay under the overnight condition

clearly support these observations, with the As/PC2 ratio of 0.5 (As:PC2=1:2) and

As/PC3 ratio of 1 (As:PC3=1:1) (Fig. 5C and 5D). In addition to PCs, As(III) would

also interact with GSH and form As(GS)$_3$ *in vitro* [67]. However, the complex *in vivo* is

only observed under extreme biological conditions: the bile of rats injected with a very

high dose of arsenite [68] and the roots of Arabidopsis PCS1 mutant *cad1-3* [59]. The

instability of As(GS)$_3$ at physiological pH and temperature [25] would partly explain the

rareness of As(GS)$_3$ *in vivo*. Supporting this hypothesis, the As/GSH ratio obtained

under physiological conditions was much lower than 0.33 (Fig. 4B), the index for

As(GS)$_3$ formation. Taken together with the previous observations *in vitro* and *in vivo*,

our results suggest that As(GS)$_3$ complex is not a major form in the cells challenged by

As(III) except for the extreme conditions, and an As-PC complex is much more stable

[69] and thus crucial for the As detoxification in plants.

Conclusion

The importance of toxic metals and essential metals homeostasis in plants as food

resources is increasing to achieve better human health, and the small molecular weight

ligands are major (potential) players in regulating these biological processes [26,70]. *In*

vitro evaluation of metal-ligand interactions is a crucial initial step for characterizing

ligand-based metal homeostasis in the cells. The proposed assay system in this study

has advantages regarding throughput, sensitivity, robustness, and specificity. Another

strong point is that the assay is flexible in terms of reaction conditions. Thus, it is

suitable for testing various metal-ligand combinations. Taking all these advantages, the

present study demonstrated dominant natures of PC-Cd interactions by a series of

binding assays.

Beyond PCs, the assay can be applied to test the potent metal affinity of other

ligands. For example, characterizing the metal interactions of the Fe/Zn-chelator NA or

other siderophores under physiological matrix conditions would advance our

understanding of the biological chelators. Other interesting ligand candidates include

isoforms of GSH and PCs in which the C-terminal Gly is replaced by β-Ala, Ser, Gln,

or Glu [71–73]. These iso-thiols are found in various plant species but have not been

well characterized. From the viewpoint of environmental science/ecotoxicology, the

assay can be used for the assessment of environmentally important metal(loid)s and

biological ligands interaction. Recently, in addition to the "classical metals", a number

of elements are newly utilized by high-tech industries and their levels in the

environment are accordingly increasing [74,75]. For these further applications,

detection by ICP-MS instead of ICP-OES is an option that would increase the

sensitivity of the assay and target elements to be tested. Another possible modification

to the method includes the buffer (matrix) composition, depending on the application.

For example, certain metalloproteins are known to adversely interact with HEPES

buffer [76], suggesting that other buffer types should be considered for such cases.

Acknowledgments

We thank Prof. Tohru Nagamitsu (Kitasato University) and Dr. Mitsuru Abo (Meiji

University) for helpful advice and discussion. This work was partly supported by the

Japan Society for the Promotion of Science (Grant nos. 18K05377, 21K05330 to SU).

Author contributions

SU, KN, and MK designed the experiments and prepared the manuscript. SU, KN, FN,

and YOt conducted the experiments. SU, KN, YOh, RN, YT, and MK analyzed the data.

All authors discussed the results and approved the manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

Data availability

The data underlying this article are available in the article and in its online

supplementary material.

References

1 Waldron KJ, Rutherford JC, Ford D & Robinson NJ (2009) Metalloproteins and metal

sensing. *Nature* **460**, 823–30.

2 Andreini C, Bertini I, Cavallaro G, Holliday GL & Thornton JM (2008) Metal ions in

biological catalysis: from enzyme databases to general principles. *J Biol Inorg Chem*

13, 1205–18.

3 Dupont CL, Butcher A, Valas RE, Bourne PE & Caetano-Anollés G (2010) History of

biological metal utilization inferred through phylogenomic analysis of protein

structures. *Proc Natl Acad Sci USA* **107**, 10567–10572.

4 Dupont CL, Yang S, Palenik B & Bourne PE (2006) Modern proteomes contain

putative imprints of ancient shifts in trace metal geochemistry. *Proc Natl Acad Sci*

USA **103**, 17822–17827.

5 Hitomi Y, Outten CE & O'Halloran TV (2001) Extreme zinc-binding thermodynamics

of the metal sensor/regulator protein, ZntR. *J Am Chem Soc* **123**, 8614–8615.

6 Outten CE & O'Halloran TV (2001) Femtomolar sensitivity of metalloregulatory

proteins controlling zinc homeostasis. *Science* **292**, 2488–2492.

7 Rae T, Schmidt P, Pufahl R, Culotta V & O'Halloran TV (1999) Undetectable

intracellular free copper: The requirement of a copper chaperone for superoxide

dismutase. *Science* **284**, 805–808.

8 Colvin RA, Holmes WR, Fontaine CP & Maret W (2010) Cytosolic zinc buffering and

muffling: their role in intracellular zinc homeostasis. *Metallomics* **2**, 306–317.

9 Finney LA & O'Halloran TV (2003) Transition metal speciation in the cell: insights

from the chemistry of metal ion receptors. *Science* **300**, 931–936.

10 Hogstrand C, Kille P, Nicholson RI & Taylor KM (2009) Zinc transporters and

cancer: a potential role for ZIP7 as a hub for tyrosine kinase activation. *Trends Mol*

Med **15**, 101–111.

11 Bashir K, Rasheed S, Kobayashi T, Seki M & Nishizawa NK (2016) Regulating

subcellular metal Homeostasis: The key to crop improvement. *Front Plant Sci* **7**,

1192.

12 Clemens S, Deinlein U, Ahmadi H, Höreth S & Uraguchi S (2013) Nicotianamine is

a major player in plant Zn homeostasis. *BioMetals* **26**, 623–632.

13 Ghssein G, Brutesco C, Ouerdane L, Fojcik C, Izaute A, Wang S, Hajjar C, Lobinski

R, Lemaire D, Richaud P, Voulhoux R, Espaillat A, Cava F, Pignol D,

Borezee-Durant E & Arnoux P (2016) Biosynthesis of a broad-spectrum

nicotianamine-like metallophore in *Staphylococcus aureus*. *Science* **352**, 1105–1109.

14 Grill E, Winnacker EL & Zenk MH (1987) Phytochelatins, a class of

heavy-metal-binding peptides from plants, are functionally analogous to

metallothioneins. *Proc Natl Acad Sci USA* **84**, 439–443.

15 Howden R, Goldsbrough PB, Andersen CR & Cobbett CS (1995)

Cadmium-sensitive, *cad1* mutants of *Arabidopsis thaliana* are phytochelatin

deficient. *Plant Physiol* **107**, 1059–1066.

16 Ha SB, Smith AP, Howden R, Dietrich WM, Bugg S, O'Connell MJ, Goldsbrough

PB & Cobbett CS (1999) Phytochelatin synthase genes from Arabidopsis and the

yeast *Schizosaccharomyces pombe*. *Plant Cell* **11**, 1153–1164.

17 Uraguchi S, Tanaka N, Hofmann C, Abiko K, Ohkama-Ohtsu N, Weber M, Kamiya

T, Sone Y, Nakamura R, Takanezawa Y, Kiyono M, Fujiwara T & Clemens S (2017)

Phytochelatin synthase has contrasting effects on cadmium and arsenic accumulation

in rice grains. *Plant Cell Physiol* **58**, 1730–1742.

18 Fischer S, Kühnlenz T, Thieme M, Schmidt H & Clemens S (2014) Analysis of plant

Pb tolerance at realistic submicromolar concentrations demonstrates the role of

phytochelatin synthesis for Pb detoxification. *Environ Sci Technol* **48**, 7552–7559.

19 Tennstedt P, Peisker D, Böttcher C, Trampczynska A & Clemens S (2009)

Phytochelatin synthesis is essential for the detoxification of excess zinc and

contributes significantly to the accumulation of zinc. *Plant Physiol* **149**, 938–948.

20 Clemens S, Kim EJ, Neumann D & Schroeder JI (1999) Tolerance to toxic metals by

a gene family of phytochelatin synthases from plants and yeast. *EMBO J* **18**,

3325–3333.

21 Kondo N, Imai K, Isobe M, Goto T, Murasugi A, Wada-Nakagawa C & Hayashi Y

(1984) Cadystin a and b, major unit peptides comprising cadmium binding peptides

induced in a fission yeast ----- separation, revision of structures and synthesis.

Tetrahedron Lett **25**, 3869–3872.

22 Vatamaniuk OK, Bucher EA, Ward JT & Rea PA (2001) A new pathway for heavy

metal detoxification in animals. Phytochelatin synthase is required for cadmium

tolerance in *Caenorhabditis elegans*. *J Biol Chem* **276**, 20817–20820.

23 Meister A (1991) Glutathione deficiency produced by inhibition of its synthesis, and

its reversal; Applications in research and therapy. *Pharmacol Therapeut* **51**,

155–194.

Wood BA & Feldmann J (2012) Quantification of phytochelatins and their metal(loid) complexes: critical assessment of current analytical methodology. *Anal Bioanal Chem* **402**, 3299–3309.

Raab A, Meharg AA, Jaspars M, Genney DR & Feldmann J (2003) Arsenic–glutathione complexes—their stability in solution and during separation by different HPLC modes. *J Anal Atom Spectrom* **19**, 183–190.

Clemens S (2019) Metal ligands in micronutrient acquisition and homeostasis. *Plant Cell Environ* **42**, 2902–2912.

Fukuda T, Nagai K, Yagi A, Kobayashi K, Uchida R, Yasuhara T & Tomoda H (2019) Nectriatide, a Potentiator of Amphotericin B Activity from *Nectriaceae* sp. BF-0114. *J Nat Prod* **82**, 2673–2681.

Cazalé AC & Clemens S (2001) *Arabidopsis thaliana* expresses a second functional phytochelatin synthase. *FEBS Lett* **507**, 215–219.

Badyal JP, Cameron AM, Cameron NR, Coe DM, Cox R, Davis BG, Oates LJ, Oye G & Steel PG (2001) A simple method for the quantitative analysis of resin bound thiol groups. *Tetrahedron Lett* **42**, 8531–8533.

Schmöger ME, Oven M & Grill E (2000) Detoxification of arsenic by phytochelatins in plants. *Plant Physiol* **122**, 793–801.

Uraguchi S, Ohshiro Y, Otsuka Y, Tsukioka H, Yoneyama N, Sato H, Hirakawa M, Nakamura R, Takanezawa Y & Kiyono M (2020) Selection of agar reagents for medium solidification is a critical factor for metal(loid) sensitivity and ionomic

profiles of *Arabidopsis thaliana*. *Front Plant Sci* **11**, 503.

32 Verret F, Gravot A, Auroy P, Leonhardt N, David P, Nussaume L, Vavasseur A &

Richaud P (2004) Overexpression of *AtHMA4* enhances root-to-shoot translocation

of zinc and cadmium and plant metal tolerance. *FEBS Lett* **576**, 306–312.

33 Semane B, Cuypers A, Smeets K, Belleghem FV, Horemans N, Schat H &

Vangronsveld J (2007) Cadmium responses in *Arabidopsis thaliana*: glutathione

metabolism and antioxidative defence system. *Physiol Plant* **129**, 519–528.

34 Li Y, Nguyen M, Baudoin M, Vendier L, Liu Y, Robert A & Meunier B (2019) Why

is tetradentate coordination essential for potential copper homeostasis regulators in

Alzheimer's Disease? *Eur J Inorg Chem* **2019**, 4712–4718.

35 Fasae KD, Abolaji AO, Faloye TR, Odunsi AY, Oyetayo BO, Enya JI, Rotimi JA,

Akinyemi RO, Whitworth AJ & Aschner M (2021) Metallobiology and therapeutic

chelation of biometals (copper, zinc and iron) in Alzheimer's disease: Limitations,

and current and future perspectives. *J Trace Elem Med Bio* **67**, 126779.

36 Kenney GE, Dassama LMK, Pandelia ME, Gizzi AS, Martinie RJ, Gao P, DeHart CJ,

Schachner LF, Skinner OS, Ro SY, Zhu X, Sadek M, Thomas PM, Almo SC,

Bollinger JM, Krebs C, Kelleher NL & Rosenzweig AC (2018) The biosynthesis of

methanobactin. *Science* **359**, 1411–1416.

37 Fahey RC (2013) Glutathione analogs in prokaryotes. *Biochim Biophys Acta Gen*

Subj **1830**, 3182–3198.

38 Soboll S, Gründel S, Harris J, Kolb-Bachofen V, Ketterer B & Sies H (1995) The

content of glutathione and glutathione S-transferases and the glutathione peroxidase

activity in rat liver nuclei determined by a non-aqueous technique of cell

fractionation. *Biochem J* **311**, 889–894.

39 Sun X, Shih AY, Johannssen HC, Erb H, Li P & Murphy TH (2006) Two-photon

imaging of glutathione levels in intact brain indicates enhanced redox buffering in

developing neurons and cells at the cerebrospinal fluid and blood-brain interface. *J*

Biol Chem **281**, 17420–17431.

40 Gutiérrez-Alcalá G, Gotor C, Meyer AJ, Fricker M, Vega JM & Romero LC (2000)

Glutathione biosynthesis in Arabidopsis trichome cells. *Proc Natl Acad Sci* **97**,

11108–11113.

41 Irving H & Williams RJP (1953) The stability of transition-metal complexes. *J Chem*

Soc **3**, 3192–3210.

42 Martin RB & Edsall JT (1959) The association of divalent cations with glutathione 1.

J Am Chem Soc **81**, 4044–4047.

43 Hider RC & Kong XL (2011) Glutathione: a key component of the cytoplasmic

labile iron pool. *Biometals* **24**, 1179–1187.

44 Krężel A & Bal W (2004) Studies of zinc(II) and nickel(II) complexes of GSH,

GSSG and their analogs shed more light on their biological relevance. *Bioinorg*

Chem Appl **2**, 293–305.

45 Walsh MJ & Ahner BA (2013) Determination of stability constants of Cu(I), Cd(II)

& Zn(II) complexes with thiols using fluorescent probes. *J Inorg Biochem* **128**,

112–123.

46 Morgan MT, Nguyen LAH, Hancock HL & Fahrni CJ (2017) Glutathione limits

aquacopper(I) to sub-femtomolar concentrations through cooperative assembly of a tetranuclear cluster. *J Biol Chem* **292**, 21558–21567.

Maitani T, Kubota H, Sato K & Yamada T (1996) The composition of metals bound to class III metallothionein (phytochelatin and its desglycyl peptide) induced by various metals in root cultures of *Rubia tinctorum*. *Plant Physiol* **110**, 1145–1150.

Oven M, Page JE, Zenk MH & Kutchan TM (2002) Molecular characterization of the homo-phytochelatin synthase of soybean *Glycine max*: relation to phytochelatin synthase. *J Biol Chem* **277**, 4747–4754.

Loscos J, Naya L, Ramos J, Clemente MR, Matamoros MA & Becana M (2006) A reassessment of substrate specificity and activation of phytochelatin synthases from model plants by physiologically relevant metals. *Plant Physiol* **140**, 1213–1221.

Li M, Barbaro E, Bellini E, Saba A, Toppi LS di & Varotto C (2020) Ancestral function of the phytochelatin synthase C-terminal domain in inhibition of heavy metal-mediated enzyme overactivation. *J Exp Bot* **71**, 6655–6669.

Uraguchi S, Sone Y, Ohta Y, Ohkama-Ohtsu N, Hofmann C, Hess N, Nakamura R, Takanezawa Y, Clemens S & Kiyono M (2018) Identification of C-terminal regions in *Arabidopsis thaliana* phytochelatin synthase 1 specifically involved in activation by Arsenite. *Plant Cell Physiol* **59**, 500–509.

Kühnlenz T, Hofmann C, Uraguchi S, Schmidt H, Schempp S, Weber M, Lahner B, Salt DE & Clemens S (2016) Phytochelatin synthesis promotes leaf Zn accumulation of *Arabidopsis thaliana* plants grown in soil with adequate Zn supply and is essential for survival on Zn-contaminated soil. *Plant Cell Physiol* **57**, 2342–2352.

53 Ruotolo R, Peracchi A, Bolchi A, Infusini G, Amoresano A & Ottonello S (2004)

Domain organization of phytochelatin synthase: functional properties of truncated

enzyme species identified by limited proteolysis. *J Biol Chem* **279**, 14686–14693.

54 Raab A, Feldmann J & Meharg AA (2004) The nature of arsenic-phytochelatin

complexes in *Holcus lanatus* and *Pteris cretica*. *Plant Physiol* **134**, 1113–1122.

55 Bluemlein K, Raab A, Meharg AA, Charnock JM & Feldmann J (2008) Can we trust

mass spectrometry for determination of arsenic peptides in plants: comparison of

LC–ICP–MS and LC–ES-MS/ICP–MS with XANES/EXAFS in analysis of

Thunbergia alata. *Anal Bioanal Chem* **390**, 1739–1751.

56 Sadi BBM, Vonderheide AP, Gong J-M, Schroeder JI, Shann JR & Caruso JA (2008)

An HPLC-ICP-MS technique for determination of cadmium–phytochelatins in

genetically modified *Arabidopsis thaliana*. *J Chromatogr B* **861**, 123–129.

57 Yen T, Villa JA & DeWitt JG (1999) Analysis of phytochelatin–cadmium complexes

from plant tissue culture using nano□electrospray ionization tandem mass

spectrometry and capillary liquid chromatography/electrospray ionization tandem

mass spectrometry. *J Mass Spectrom* **34**, 930–941.

58 Chen L, Guo Y, Yang L & Wang Q (2007) SEC-ICP-MS and ESI-MS/MS for

analyzing *in vitro* and *in vivo* Cd-phytochelatin complexes in a Cd-hyperaccumulator

Brassica chinensis. *J Anal At Spectrom* **22**, 1403–1408.

59 Liu W-JJ, Wood BA, Raab A, McGrath SP, Zhao F-JJ & Feldmann J (2010)

Complexation of arsenite with phytochelatins reduces arsenite efflux and

translocation from roots to shoots in Arabidopsis. *Plant Physiol* **152**, 2211–2221.

60 Jacquart A, Brayner R, Chahine J-MEH & Ha-Duong N-T (2017) Cd^{2+} and Pb^{2+}

complexation by glutathione and the phytochelatins. *Chem-biol Interact* **267**, 2–10.

61 Dorčák V & Krężel A (2003) Correlation of acid–base chemistry of phytochelatin

PC2 with its coordination properties towards the toxic metal ion Cd(II). *Dalton*

Trans, 2253–2259.

62 Vatamaniuk OK, Mari S, Lang A, Chalasani S, Demkiv LO & Rea PA (2004)

Phytochelatin synthase, a dipeptidyltransferase that undergoes multisite acylation

with gamma-glutamylcysteine during catalysis: stoichiometric and site-directed

mutagenic analysis of *Arabidopsis thaliana* PCS1-catalyzed phytochelatin synthesis.

J Biol Chem **279**, 22449–22460.

63 Mendoza-Cózatl DG, Zhai Z, Jobe TO, Akmakjian GZ, Song W-YY, Limbo O,

Russell MR, Kozlovskyy VI, Martinoia E, Vatamaniuk OK, Russell P & Schroeder JI

(2010) Tonoplast-localized Abc2 Transporter Mediates Phytochelatin Accumulation

in Vacuoles and Confers Cadmium Tolerance. *J Biol Chem* **285**, 40416–40426.

64 Park J, Song W-YY, Ko D, Eom Y, Hansen TH, Schiller M, Lee TG, Martinoia E &

Lee Y (2012) The phytochelatin transporters AtABCC1 and AtABCC2 mediate

tolerance to cadmium and mercury. *Plant J* **69**, 278–288.

65 Song W-YY, Yamaki T, Yamaji N, Ko D, Jung K-HH, Fujii-Kashino M, An G,

Martinoia E, Lee Y & Ma JF (2014) A rice ABC transporter, OsABCC1, reduces

arsenic accumulation in the grain. *Proc Natl Acad Sci* **111**, 15699–15704.

66 Song W-YY, Park J, Mendoza-Cózatl DG, Suter-Grotemeyer M, Shim D,

Hörtensteiner S, Geisler M, Weder B, Rea PA, Rentsch D, Schroeder JI, Lee Y &

Martinoia E (2010) Arsenic tolerance in Arabidopsis is mediated by two ABCC-type phytochelatin transporters. *Proc Natl Acad Sci* **107**, 21187–21192.

Rey NA, Howarth OW & Pereira-Maia EC (2004) Equilibrium characterization of the As(III)–cysteine and the As(III)–glutathione systems in aqueous solution. *J Inorg Biochem* **98**, 1151–1159.

Kala SV, Neely MW, Kala G, Prater CI, Atwood DW, Rice JS & Lieberman MW (2000) The MRP2/cMOAT transporter and arsenic-glutathione complex formation are required for biliary excretion of arsenic. *J Biol Chem* **275**, 33404–33408.

Bluemlein K, Raab A & Feldmann J (2009) Stability of arsenic peptides in plant extracts: off-line versus on-line parallel elemental and molecular mass spectrometric detection for liquid chromatographic separation. *Anal Bioanal Chem* **393**, 357–366.

Clemens S (2019) Safer food through plant science: reducing toxic element accumulation in crops. *J Exp Bot* **70**, 5537–5557.

Klapheck S, Fliegner W & Zimmer I (1994) Hydroxymethyl-phytochelatins [γ-glutamylcysteine)n-serine] are metal-induced peptides of the *Poaceae*. *Plant Physiol* **104**, 1325–1332.

Klapheck S, Schlunz S & Bergmann L (1995) Synthesis of phytochelatins and homo-phytochelatins in *Pisum sativum* L. *Plant Physiol* **107**, 515–521.

Meuwly P, Thibault P, Schwan AL & Rauser WE (1995) Three families of thiol peptides are induced by cadmium in maize. *Plant J* **7**, 391–400.

Merschel G & Bau M (2015) Rare earth elements in the aragonitic shell of freshwater mussel *Corbicula fluminea* and the bioavailability of anthropogenic

lanthanum, samarium and gadolinium in river water. *Sci Total Environ* **533**, 91–101.

75 Tokumaru T, Ozaki H, Onwona-Agyeman S, Ofosu-Anim J & Watanabe I (2017)

Determination of the extent of trace metals pollution in soils, sediments and human

hair at e-waste recycling site in Ghana. *Arch Environ Contam Toxicol* **73**, 377–390.

76 Jahromi EZ, White W, Wu Q, Yamdagni R & Gailer J (2010) Remarkable effect of

mobile phase buffer on the SEC-ICP-AES derived Cu, Fe and Zn-metalloproteome

pattern of rabbit blood plasma. *Metallomics* 2, 460–468.

 Figure legends

Fig. 1 Verification of the synthesized phytochelatins. (A-F) and loading to the beads (G).

(A, B) Total ion current chromatograms of the synthesized PC2 and PC3. (C, D)

LC-PDA chromatograms of the synthesized PC2 and PC3 (at 210 nm). (E, F) Mass

spectrum of the synthesized PC2 and PC3 (acquisition range: *m/z* 100–1000). (G)

Loadings of GSH, PC2, and PC3 onto TOYOPEARL AF-Fornyl-650 beads were carried

out under reductive amination conditions. The loadings were estimated by detecting the

amount of 2-nitro-5-thiobenzoate anion released from the reaction of Ellman's reagent

and thiol groups on the affinity beads.

Fig. 2 Schematic diagram describing the two protocols of the metal-ligand binding

assay using the ligand-attached beads and ICP. A reaction mixture for the standard

protocol was prepared in a 5 ml plastic tube containing the beads (normally 3.0 mg) and

a metal(loid)-of-interest (normally 200 nmol). The HEPES buffer (pH 7.2) was used as

a matrix instead of a phosphate buffer to avoid the potential interaction of the

metal(loid) and phosphate ions. The prepared tubes were incubated at room temperature

for 5 min or overnight with continuous rotation to avoid sedimentation of the beads.

After the incubation sample was immediately collected by Protocol A or Protocol B. In

Protocol A, a filtrate not containing beads was collected using a spin column and used

for the ICP quantification. In Protocol B, the beads were collected by centrifugation and

subjected to the ICP analysis.

Fig. 3 Comparison of Protocol A and Protocol B using Cd as a metal model. Total Cd

adsorption to the respective types of affinity beads was analyzed by ICP-OES after the

5-min incubation with 200 nmol Cd. Data represent means with SD from three

independent reactions for each combination. There is no significant difference between

the two protocols for all affinity beads tested ($P > 0.05$, Student's T-test). n.s., not

significant.

Fig. 4 Metal(loid)-ligand interaction assay using the ligand-attached affinity beads. A

reaction mixture containing 3.0 mg of each type of the beads and 200 nmol of each

metal(loid) species was incubated for 5 min or overnight at room temperature with

contentious rotation. After the incubation, the reaction mixture was immediately

subjected to spin column filtration (Protocol A). The metal(loid) content in the filtrate

was quantified by ICP-OES. Data represent means with SD from three independent

reactions for each combination. Means sharing the same letter are not significantly

different ($P < 0.05$, Tukey's HSD). (A) Non-specific metal(loid) adsorption to the

control (blank) beads. (B-D) Metal(loid) binding ratios of GSH-attached beads (B),

PC2-attached beads (C), and PC3-attached beads (D). n.d., not detected. n.a., not

analyzed.

Fig. 5 Metal competition assay using the ligand-attached beads. The reaction buffer

containing 20 nmol of $MnCl_2$, 40 nmol of $FeCl_2$, 2 nmol of $CuSO_4$, 20 nmol of $ZnSO_4$,

and 200 nmol of $CdCl_2$ was added to 3.0 mg of each type of the beads. The tube was

immediately set to a rotator and incubated at room temperature for 5 min. After the

incubation, the reaction mixture was immediately subjected to spin column filtration

(Protocol A). The metals content in the filtrate was quantified by ICP-OES. (A-C)

Proportions of metals binding to GSH (A), PC2 (B), and PC3 (C) on the beads,

calculated from total metal input to the reaction and the metals associated with each

ligand on the beads. Data represent means with SD from three independent reactions.

Means sharing the same letter are not significantly different ($P < 0.05$, Tukey's HSD).

(D) Profiles of the metals in the reaction mixture (input) and on the respective beads.

Data represent means from three independent reactions.

Fig. 6 PC2 and GSH competition assay using the ligand-attached beads. (A) The

reaction buffer containing 100 nmol of $CdCl_2$ and 100 nmol of PC2 was added to 3.0

mg of the GSH-attached beads. (B) The reaction buffer containing 100 nmol of $CdCl_2$,

and 25 or 200 nmol of GSH was added to 3.0 mg of the PC2-attached beads. (A, B) The

tube was immediately set to a rotator after adding the buffer and incubated at room

temperature for 5 min. After the incubation, the reaction mixture was immediately

subjected to spin column filtration (Protocol A). The Cd content in the filtrate was

quantified by ICP-OES. Data represent means with SD from three independent

reactions. *** $P < 0.001$, (Student's T-test). n.s. $P > 0.05$ (Dunnett's test).

A phytochelatin2 (PC2): (γ-Glu-Cys)₂-Gly

B phytochelatin3 (PC3): (γ-Glu-Cys)₃-Gly

	yield (nmol thiol mg⁻¹)	
	1st lot	2nd lot
GSH (n=1)	64.1	70.2
PC2 (n=2)	67.5	78.4
PC3 (n=3)	64.8	69.6

Mixture Preparation

beads   metal(loid)

40 mM HEPES (pH 7.2)

Incubation

5 min or O/N at
RT with rotation

Binding reaction

Sample Collection

Protocol A   Collecting filtrate

Immediate filtration
with a spin column

Protocol B   Collecting beads

Immediate
centrifugation

Acid extraction
(140°C, 3h) & ICP

ICP Quantification & Calculation

Dilution & ICP

Protocol A

$[M]_{binding} = [M]_{control\ filtrate} - [M]_{ligand\ filtrate}$

Protocol B

$[M]_{binding} = [M]_{ligand\ beads} - [M]_{control\ beads}$

A

GSH beads

Cd binding to GSH (% of input)

***

without competitor | 100 nmol
PC2

B

PC2 beads

Cd binding to PC2 (% of input)

n.s.

without competitor | 25 nmol | 200 nmol
GSH