Short Communication

Genome-wide development of lncRNA-derived-SSR markers for Dongxiang wild

rice (*Oryza rufipogon* Griff.)

Wanling Yang[1#], Yuanwei Fan[2, 3#], Yong Chen[1], Gumu Ding[1], Hu Liu[4], Jiankun Xie[1*]

and Fantao Zhang[1*]

[1]*Jiangxi Provincial Key Lab of Protection and Utilization of Subtropical Plant*

Resources, College of Life Sciences, Jiangxi Normal University, Nanchang 330022,

China

[2]*College of Biological Sciences and Biotechnology, Beijing Forestry University,*

Beijing 100083, China

[3]*Current address: Department of Biology and Center for Engineering*

Mechanobiology, Washington University in St.Louis, St.Louis, MO 63130, USA

[4]*Huaian Zhuxiang Ecological Agriculture CO. LTD, Huaian 223299, Jiangsu*

Province, China

[#]Contributed equally to this work

[*]Corresponding authors. E-mail: zhang84004@163.com and xiejiankun11@163.com

Abstract Dongxiang wild rice (*Oryza rufipogon* Griff., DXWR) is the northernmost

distributed common wild rice found in the world. It contains a large number of

agronomically valuable genes, which makes it a natural gene pool for rice breeding.

Molecular markers, especially simple repeat sequence (SSR) markers, play important

roles in crop breeding. Although a large number of SSR markers have been developed,

most of them are derived from the genome coding sequences, rarely from non-coding

sequences. Meanwhile, long non-coding RNAs (lncRNAs), which are derived from

the transcription of non-coding sequences, play vital roles in plant growth,

development and stress responses. In this study, 1878 SSR loci were detected from the

lncRNA sequences of DXWR, and 1258 lncRNA-derived-SSR markers were

developed on the genome-wide scale. To verify the validity and applicability of these

markers, 72 pairs of primers were randomly selected to test 44 rice materials. The

results showed that 42 (58.33%) pairs of primers have abundant polymorphism among

these rice materials; the polymorphism information content (PIC) values ranged from

0.04 to 0.87 with an average of 0.50; the genetic diversity index of SSR loci varied

from 0.04 to 0.88 with an average of 0.56; and the number of alleles per marker

ranged from 2 to 11 with an average of 4.36. Thus, we concluded that these

lncRNA-derived-SSR markers are a very useful source for future basic and applied

research, including genetic diversity analysis, QTL mapping, and molecular breeding

programs, to make good use of the elite lncRNA genes from DXWR.

Keywords: common wild rice, genetic marker, germplasm resource, long non-coding

RNA, simple repeat sequence

Introduction

Dongxiang wild rice (*Oryza rufipogon* Griff., DXWR) is a common wild rice species found in Dongxiang County, Jiangxi Province, China. It is by far the northernmost ($28^{o}14'$ N) distributed common wild rice found in the world (Mao *et al.*, 2015). Previous studies have showed that DXWR possesses many stress tolerance and high yield-related genes, which have been lost in modern cultivated rice (Zhang *et al.*, 2016). Therefore, DXWR is an elite genetic resource to breed new rice varieties.

In eukaryotes, over 90% of the genome is transcribed to large amounts of non-coding RNAs (ncRNAs) that are not translated into proteins (Wierzbicki, 2012). Long non-coding RNAs (lncRNAs) are an important type of ncRNAs. In plants, lncRNAs act as key regulators of plant growth and development, as well as in response to various biotic and abiotic stresses (Xu *et al*., 2017). Molecular marker-assisted selection (MAS) can make great use of the valuable genetic resources in wild rice by precisely selecting genes for various traits, therefore producing superior germplasm (Collard and Mackill, 2008; Akhtamov *et al*., 2020). Simple repeat sequences (SSRs), also known as microsatellites, are DNA segments with 1-6 nucleotides short base-pair motif repeated several times in tandem (Kozlowski *et al*., 2010). They have been extensively used in genetic diversity analysis and MAS breeding because of their high reproducibility, abundant polymorphisms, co-dominant inheritance, high genome coverage and simple analysis methods (Xie *et al*., 2017; Singh *et al*., 2018). However, there are no reports on the development of

lncRNA-derived-molecular markers in common wild rice so far, which greatly limits

the discovery and utilization of the elite lncRNA genes from the important and

endangered germplasm resource. Therefore, the main objectives of this study were to:

(i) use the lncRNAs sequencing data to develop a set of lncRNA-derived-SSR

markers and (ii) test their stabilities and polymorphisms for DXWR.

Experimental

The information of rice materials used in this study was presented in Supplementary

Table S1. To develop more lncRNA-derived-SSR markers, SSR Hunter software was

used to identify all possible di-, tri-, tetra-, penta-, and hexa-nucleotide SSRs with a

minimum set of three repeats, respectively (Li and Wan, 2005). Subsequently, primers

were designed based on the flanking sequences of the SSRs using Primer3.0 software

(Untergasser *et al.*, 2012). The main parameters of the primer design were as follows:

the primer length was 18-25 bp, GC content was 40-60 %, melting temperature was

55-65 ^{o}C, the expected length of the amplification product was 100-250 bp.

Genomic DNA was extracted using the Plant Genomic DNA Rapid Extraction

Kit (Sangon Biotech Co., Ltd). The PCR reaction system was 10 µL, including 1 µL

(200 ng/µL) genomic DNA, 5 µL 2 × FastTaq Premix (Tolo Biotech Co., Ltd), 1 µL

(0.01 nmol/µL) primers and 3 µL ddH$_2$O. The PCR amplification reaction program

was as follows: pre-denaturation at 95^{o}C for 5 min; denaturation at 95^{o}C for 30 s,

annealing at 55^{o}C for 45 s, extension at 72^{o}C for 45 s for 30 cycles; and a final

extension at 72^{o}C for 10 min. The PCR products were run on 9 % denaturing

polyacrylamide gel with 0.5 × TBE buffer. After electrophoresis, the gels were

visualized using the silver staining method (Cook *et al.*, 2004). The genetic

parameters were analyzed using Powermarker software (Liu and Muse, 2005).

Discussion

In our previous study, a total of 1655 lncRNA transcripts were obtained from DXWR

using strand-specific RNA sequencing (Qi *et al.*, 2020). In this study, to develop a set

of lncRNA-derived-SSR markers for DXWR, SSR Hunter software was used to

search for SSRs present in these lncRNA transcripts. Totally, 1878 SSRs were

detected. Among them, the dinucleotide (1291, 68.74%) and trinucleotide (498,

26.52%) repeat motifs were the most abundant types (Table 1). Moreover, the types of

AG/CT (505, 39.12%) and CCG/CGG (77, 15.46%) were the most predominant in the

dinucleotide and trinucleotide repeat types, respectively (Fig. 1). These results

following the same pattern as the lncRNA-derived-SSR study in wheat (Bhandawat *et*

al., 2020).

Based on the detected lncRNA-derived-SSR loci, we successfully developed a

total of 1258 molecular markers, including 885 (70.35%) for dinucleotide repeat type,

316 (25.12%) for trinucleotide repeat type, 35 (2.78%) for tetranucleotide repeat type,

16 (1.27%) for pentanucleotide repeat type and 6 (0.48%) for hexanucleotide repeat

type (Supplementary Table S2). These lncRNA-derived-SSR markers were present

throughout all of the 12 chromosomes. Of these markers, 781 (62.08%) were found to

be present in the first six chromosomes (Chr. 1-6); while 477 (37.92%) were present

in the remaining six chromosomes (Chr. 7-12). This is largely due to the identified

SSR loci exhibit some preference for the lncRNA transcripts located in chromosomes

1-6 of DXWR. It was also observed that Chr. 2 has the highest number of

lncRNA-derived-SSR markers (169, 13.43%) while Chr.9 has the lowest number of

lncRNA-derived-SSR markers (66, 5.25%).

To test the stabilities and polymorphisms of the developed lncRNA-derived-SSR

markers, we randomly selected 72 pairs of primers for further analysis. As a result, 42

(58.33%) pairs of primers showed abundant polymorphisms among the 44 rice

materials (Fig. 2). The 42 polymorphic SSRs produced a total of 183 alleles among

the 44 rice materials, ranging from 2 to 11, with an average 4.36 alleles per locus. The

polymorphic information content (PIC) of these polymorphic SSR markers ranged

from 0.04 to 0.87 with an average of 0.50. The genetic diversity index was observed

from 0.04 to 0.88, with a mean value of 0.56 (online Supplementary Table S3). The

mean number of alleles and PIC value of the verified lncRNA-derived-SSR markers

in our study were more than capsicum which were 2.50 and 0.39 respectively (Jaiswal

et al., 2020), indicating that the developed lncRNA-derived-SSR markers were highly

polymorphic and can be wildly applied in DXWR and modern cultivated rice.

In summary, this is the first report of the development and characterization of

lncRNA-derived-markers in common wild rice, which lays the foundation for

discovery and utilization of the elite lncRNA genes to further make good use of the

valuable and endangered wild rice germplasm resource.

Acknowledgements

This research was partially supported by the National Natural Science Foundation of

China (32070374, 31960370, 31960085), the Natural Science Foundation of Jiangxi

Province, China (20202ACB205002), the Foundation of Jiangxi Provincial Key Lab

of Protection and Utilization of Subtropical Plant Resources (YRD201903).

References

Akhtamov M, Adeva C, Shim KC, Lee HS, Kim SH, Jeon YA, Luong NH, Kang JW,

Lee JY and Ahn SN (2020) Characterization of quantitative trait loci for

germination and coleoptile length under low-temperature condition using

introgression lines derived from an interspecific cross in rice. *Genes* 11: 1200.

Bhandawat A, Sharma H, Pundir N, Madhawan A and Roy J (2020) Genome-wide

identification and characterization of novel non-coding RNA-derived SSRs in

wheat. *Molecular Biology Reports* 47: 6111–6125.

Collard BC and Mackill DJ (2008) Marker-assisted selection: an approach for

precision plant breeding in the twenty-first century. *Philosophical Transactions*

of the Royal Society B: Biological Sciences 363: 557–572.

Cook GA, Prakash O, Zhang K, Shank LP, Takeguchi WA, Robbins A, Gong YX,

Iwamoto T, Schultz BD and Tomich JM (2004) Activity and structural

comparisons of solution associating and monomeric channel-forming peptides

derived from the glycine receptor m2 segment. *Biophysical Journal* 86:

1424–1435.

Jaiswal V, Rawoof A, Dubey M, Chhapekar SS, Sharma V and Ramchiary N (2020)

Development and characterization of non-coding RNA based simple sequence

repeat markers in *Capsicum* species. *Genomics* 112: 1554–1564.

Kozlowski P, de Mezer M and Krzyzosiak WJ (2010) Trinucleotide repeats in human

genome and exome. *Nucleic Acids Research* 38: 4027–4039.

Li Q and Wan JM (2005) SSR hunter: development of local searching software for

SSR sites. *Hereditas* 27: 808–810.

Liu K and Muse SV (2005) PowerMarker: an integrated analysis environment for

genetic marker analysis. *Bioinformatics* 21: 2128–2129.

Mao DH, Yu L, Chen DZ, Li L, Zhu YX, Xiao YQ, Zhang DC and Chen CY (2015)

Multiple cold resistance loci confer the high cold tolerance adaptation of

Dongxiang wild rice (*Oryza rufipogon*) to its high-latitude habitat. *Theoretical*

and Applied Genetics 128: 1359–1371.

Qi WD, Chen HP, Yang ZZ, Hu BL, Luo XD, Ai B, Luo Y, Huang Y, Xie JK and

Zhang FT (2020) Systematic Characterization of Long Non-Coding RNAs and

Their Responses to Drought Stress in Dongxiang Wild Rice. *Rice Science* 27:

21–31.

Singh AK, Chaurasia S, Kumar S, Singh R, Kumari J, Yadav MC, Singh N, Gaba S

and Jacob SR (2018) Identification, analysis and development of salt responsive

candidate gene based SSR markers in wheat. *BMC Plant Biology* 18: 249.

Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M and Rozen

SG (2012) Primer3 – new capabilities and interfaces. *Nucleic Acids Research* 40:

e115.

Wierzbicki AT (2012) The role of long non-coding RNA in transcriptional gene

silencing. *Current Opinion in Plant Biology* 15: 517–522.

Xie JK, Zhang M, Sun J and Zhang FT (2017) Genome-wide genic SSR marker

development for the endangered Dongxiang wild rice (*Oryza rufipogon*). *Plant*

Genetic Resources 15: 566–569.

Xu Q, Song ZH, Zhu CY, Tao CC, Kang LF, Liu W, He F, Yan J and Sang T (2017)

Systematic comparison of lncRNAs with protein coding mRNAs in population

expression and their response to environmental change. *BMC Plant Biology* 17:

42.

Zhang FT, Xu T, Mao LY, Yan SY, Chen XW, Wu ZF, Chen R, Luo XD, Xie JK and

Gao S (2016) Genome-wide analysis of Dongxiang wild rice (*Oryza rufipogon*

Griff.) to investigate lost/acquired genes during rice domestication. *BMC Plant*

Biology 16: 103.

Table 1. Percentage of the identified SSR motifs in different repeat types of DXWR

Fig. 1. Frequency of different SSR repeat motif types in DXWR.

Fig. 2. Representative results of polymorphic lncRNA-derived-SSR markers

amplified by the genome of DXWR and other worldwide rice cultivars. (a)

Lnc-SSR-566686; (b): Lnc-SSR-353993; M: DL2000 DNA Marker; 1-3: three

different populations of DXWR; 4-44: other worldwide cultivars. The information of

the 44 rice materials was presented in online Supplementary Table S1.

Table S1. Dongxiang wild rice and worldwide rice cultivars used for verification of

the developed lncRNA-derived-SSR markers in this study

Table S2. The information of developed lncRNA-derived-SSR markers for

Dongxiang wild rice

Table S3. The verified lncRNA-derived-SSR markers and their amplification results

with 44 rice materials in this study

Table 1. Percentage of the identified SSR motifs in different repeat types of DXWR

SSR motif	Repeat number					Percentage (%)
	3	4	5	≥ 6	Total	
Dinucleotide	1123	117	20	31	1291	68.74
Trinucleotide	424	47	23	4	498	26.52
Tetranucleotide	51	5	0	1	57	3.04
Pentanucleotide	18	0	1	0	19	1.01
Hexanucleotide	11	2	0	0	13	0.69
Total	1627	171	44	36	1878	100
Percentage (%)	86.63	9.11	2.34	1.92	100	

Fig. 1. Frequency of different SSR repeat motif types in DXWR.

Fig. 2. Representative results of polymorphic lncRNA-derived-SSR markers amplified by the genome of DXWR and other worldwide rice cultivars. (a) Lnc-SSR-566686; (b): Lnc-SSR-353993; M: DL2000 DNA Marker; 1-3: three different populations of DXWR; 4-44: other worldwide cultivars. The information of the 44 rice materials was presented in online Supplementary Table S1.