Genetic coupling of life-history and aerobic performance in Atlantic salmon

Short title: Genetic coupling of life-history and metabolism

Jenni M. Prokkola[1]*, Eirik R. Åsheim[1], Sergey Morozov[1], Paul Bangura[1], Jaakko Erkinaro[2], Annukka Ruokolainen[1], Craig R. Primmer[1,3], Tutku Aykanat[1]

[1]Organismal and Evolutionary Biology Research Programme, University of Helsinki, Finland
[2]Natural Resources Institute Finland
[3]Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Finland
*Corresponding author, PO Box 65, FI-00014 University of Helsinki, Finland, jenni.prokkola@helsinki.fi

Acknowledgements
We thank Nikolai Piavchenko, Katja Maamela, Markus Lauha, Petra Liljeström, Suvi Ikonen, Mikko Immonen, and Anna Toikkanen for fish care and technical help during the experiment, and Lammi Biological Station for a high-quality research environment. Dr. Markus Haapala and the Nordic University Hub NordicPOP (Nordforsk, project no. 85352) are acknowledged for the 3D printing service, and Heidrikur Bergsson for sharing the chamber cap model. The study was funded by Academy of Finland (T. Aykanat: 325964, 1328860, C. R. Primmer: 314254 and 314255, 327255), the European Research Council under the European Articles Union's Horizon 2020 research and innovation program (grant no. 742312), and the University of Helsinki.

Keywords: age-at-maturity, energetics, metabolic rate, life history evolution

Author contributions

JMP, ERÅ, SM, CRP and TA conceived the ideas and designed methodology; JMP, ERÅ,

SM, PB, JE, AR and TA collected the data; JMP, SM and TA analysed the data; JMP and TA

led the writing of the manuscript. All authors contributed to the drafts and gave final approval

for publication.

Abstract

A better understanding of the genetic and phenotypic architecture underlying life-history

variation is a longstanding aim in biology. Theories suggest energy metabolism determines

life-history variation by modulating resource acquisition and allocation trade-offs, but the

genetic underpinnings of the relationship and its dependence on ecological conditions have

rarely been demonstrated. The strong genetic determination of age-at-maturity by two

unlinked genomic regions (*vgll3* and *six6*) makes Atlantic salmon (*Salmo salar*) an ideal

model to address these questions. Using more than 250 juveniles in common garden

conditions, we quantified the covariation between metabolic phenotypes –standard and

maximum metabolic rates (SMR and MMR), and aerobic scope (AS) – and the life-history

genomic regions and tested if food availability modulates the relationships. We found that the

early maturation genotype in *vgll3* was associated with higher MMR and consequently AS.

Additionally, MMR exhibited physiological epistasis; it was decreased when late maturation

genotypes co-occurred in both genomic regions. Contrary to our expectation, the life-history

genotypes had no effects on SMR. Further, food availability had no effect on the genetic

covariation, suggesting a lack of genotype-by-environment interactions. Our results provide

insights on the key organismal processes that link energy use at the juvenile-stage to age-at-

maturity, indicating potential mechanisms by which metabolism and life-history can

coevolve.

Introduction

Physiological processes control how life-history diversity emerges from resource allocation and acquisition trade-offs[1]. The rate of aerobic energy metabolism is a pivotal mechanism contributing to life-history variation – it modulates resource acquisition, provides cells with ATP, and constrains energy allocation to different body components and functions. Theories, such as the metabolic theory of ecology and the pace-of-life syndrome theory[2,3], suggest metabolic rate covaries with life-history variation within and among species. This covariation may have a genetic basis, consequently constraining trait evolution[4], yet only a few studies have demonstrated intraspecific genetic covariation or co-evolution between metabolic rate and life-history traits[5-7]. Determining whether this relation is modified by different ecological contexts (e.g., food availability) is crucial to better understand the mechanisms shaping life history variation and demographic shifts in populations in response to environmental changes[8].

The quintessential components of energy metabolism at the organismal level (i.e., the metabolic phenotypes) are standard metabolic rate (SMR), maximum metabolic rate (MMR), and the absolute aerobic scope (AS) that is the difference between SMR and MMR[9-11]. SMR is the minimum metabolic rate of an ectothermic animal associated with self-maintenance, and therefore defines the minimal cost of living (excluding growth, digestion, and locomotion). MMR defines the upper limit of aerobic performance that is functionally linked to SMR and the capacity to increase oxygen uptake and delivery beyond SMR. SMR and MMR together are the integral components of AS, which is the measure of surplus energy that can be allocated into non-maintenance functions, such as locomotion and digestion[12]. Higher AS is predicted to increase fitness via facilitating energetically demanding behaviours (such as migration, aggression, predator avoidance, and prey capture) and tolerance to environmental stress[11,13-15]. However, high aerobic performance comes with costs, including maintaining a larger heart and gill surface area (associated with increased demand for osmoregulation)[12,16,17].

Allocation of energy to growth or improved condition can link metabolic phenotypes to life-history traits[18]. Life-history traits, such as the timing of maturation and migration, are determined by adaptive body-size thresholds[19-21], and metabolic phenotypes are often correlated with growth rate, albeit in a context-dependent manner[11]. Under high food

availability, a high SMR in combination with high AS can increase growth rate[22], as it often

correlates with traits that improve resource acquisition, such as dominance and digestive

capacity[23-25]. Under low food availability, the growth benefit of high SMR or AS can be

minimized (or even reversed for SMR) due to high self-maintenance costs[22,26,27]. In addition,

individuals can be forced to seek new habitats or take more risks to acquire resources,

exerting further fitness costs[28,29]. Hence, covariation between life-history and metabolism,

whether it is genetically or environmentally driven, could be modulated by resource

availability. A resource dependent change in genetic covariation, i.e., genotype-by-

environment interaction, could maintain genetic variation in these traits[30-32], but this has not

been demonstrated.

In anadromous (sea-migrating) salmonids, the number of years the fish spends at sea before

the first spawning, i.e., sea age-at-maturity, has a dramatic effect on its size-at-maturity[33].

Individuals spending one year at sea typically weigh 1–3 kg compared with 10–20 kg after

three or more years. Increased size in late maturing individuals also translates to marked

gains in reproductive investment in both sexes[33]. Earlier maturation, i.e., less time spent at

sea, provides a potential fitness advantage through a higher probability of survival prior to

reproduction and a shorter generation time, but comes at the expense of decreased fecundity

and mating success (due to smaller size at reproduction[34-36]. The probability of early

maturation at sea is also positively associated with faster growth and fat deposition in the

freshwater (juvenile) stage, since the size of salmon at the onset of sea-migration has a

significant influence on maturation timing at sea[19,37-40]. These relationships suggest that early

maturation in salmon may be associated with higher SMR or aerobic scope via resource

utilization already in early life-stages prior to sea migration[24,25]. In addition, metabolic

phenotypes in the juvenile stage may explain maturation at sea through genetic correlations

across life stages (e.g.,[41]) if metabolic rate or aerobic performance at sea is linked to earlier

maturation.

In Atlantic salmon (*Salmo salar* L. 1758), a large proportion of variation in age-at-maturity is

explained by a single genomic region that encompasses the *vgll3* gene on chromosome 25

[42,43]. In addition, variation in another locus on chromosome 9, *six6*, is a strong predictor of

mean age-at-maturity among populations [43] and associated with early maturation in

aquaculture salmon[44]. *Vgll3* and *six6* are also associated with size-at-maturity, with the alleles

conferring late maturation being associated with larger age-specific body size especially after

multiple years at sea[43]. Moreover, *vgll3* is associated with precocious maturation in male

salmon parr via body condition[45], emphasizing the causal effect of energy acquisition on

maturation. In the last few decades, many Atlantic salmon populations have been maturing,

on average, at younger ages[46], which is associated with a change in *vgll3* allele frequency in

some cases[47]. Recently, a link was found between the decrease in salmon age-at-maturity and

a change in prey species composition[48] (see also [49] for diet composition in relation to *six6*).

These observations further highlight that genotype dependent differences in SMR or aerobic

performance, which can contribute to foraging success or food assimilation (e.g.,[22,50,51]), may

be related to contemporary life-history evolution in Atlantic salmon.

The strong effects of the *six6* and *vgll3* genomic regions on life-history variation provide an

opportunity for the genetic covariation between age-at-maturity and energy metabolism to be

studied prior to maturation, i.e., at the juvenile stage, by genetic prediction. This approach

makes controlled, empirical settings more feasible, as salmon require several years to reach

maturation. In this study, we test if genetic covariation exists between life-history and

metabolic phenotypes in juveniles; we expect that early maturation genotypes show a higher

metabolic activity (SMR, MMR, and AS) than late maturation genotypes under high food

availability (e.g.,[7]). Further, because low resource availability weakens the relationship

between SMR and growth[27] and may constrain aerobic performance[8], we also explore if food

availability modulates the genetic covariation.

Material and Methods

145 The experiments were conducted under an animal experiment permit granted by the Finnish

146 Project Authorisation Board (permit nr. ESAVI/4511/2020). Additional experimental details

147 are provided in online supplemental material.

Fish rearing and genotyping

The parental Atlantic salmon (Table S1) were first-generation hatchery brood stock (from the

river Kymijoki in Finland) from Laukaa hatchery, managed by the Natural Resources

Institute Finland (LUKE). In October 2019, eggs and milt were transferred to the University

of Helsinki for fertilisation. Full-sib families were crossed using heterozygous parents in

vgll3 and *six6* loci, (*vgll3**E/L and *six6**E/L where E and L refer to the alleles associated

with early and late maturation, respectively). This provides offspring with all genotype

combinations within each full-sib family. Feed rations were calculated assuming feed

conversion efficiency of 0.8, using growth predictions by Elliott and Hurley[52]. In July 2020,

fish were tagged with passive integrated transponder (PIT) tags and genotyped from fin clips.

Experimental design

Food treatments were started in August 2020. Briefly, fish were fed twice week^{-1} in the low

food treatment (Fig. 1). This was preferred over constant low ration to minimize dominance

hierarchies in tanks[53]. Fish in the high food treatment were fed with the total ad-libitum daily

ration. After 28–31d in the treatments, 48 and 32 fish from each family in the low and high

food treatment, respectively (192 and 128 in total), were measured for their SMR and MMR

(Fig. 1), though only 290 homozygous individuals were used in the analysis (Table S2).

SMR and MMR measurements

Sixteen fish at once were moved into an acclimation tank two days before SMR measurement

(Fig. S3). Each batch was from the same family and tank and balanced for genotype-sex-

combinations. During acclimation, the fish were held individually, without feeding at 11°C ±

0.1°C in 20×20×10cm cages, to minimize the effects of temperature fluctuations, digestion,

growth, and social interactions on SMR[54]. We measured SMR using intermittent flow

respirometry[55-57]. The SMR measurements commenced after 42–47h acclimation, between

11:30 and 14:30, until 8:00 the following day. Afterwards, MMR was measured using the

chase method, similarly to, e.g.,[58], where MMR reflects increased aerobic respiration related

to exercise and the oxygen debt incurred by anaerobic respiration [59,60]. Fish were then

euthanized with an overdose of methanesulfonate, measured, and weighed. All fish were

confirmed to have immature gonads.

Analysis of respirometry data

For SMR, oxygen consumption rate (MO_2, mg O$_2$ h^{-1}) for each linear measurement phase

was derived from best-fit linear regression of dissolved oxygen concentration over time. The

mean of the lowest normal distribution (MLND) was used to estimate SMR for each

individual as mg O$_2$ h^{-1} from the extracted MO_2 slopes[10].

MMR was calculated for each individual as mg O$_2$ h^{-1} from the O$_2$ concentrations after

performing background correction in *FishResp*[61]. We used two methods to identify the slope

of the steepest decrease in O_2 saturation. First, we used *respR*[62] with the function *auto_rate*,

fitting one and two-minute windows[63] (example slope in Fig. S5A). Second, slopes for MMR

were extracted using a derivative of a polynomial curve fitted on each measurement (function

smooth.spline, df=10). This is the '*spline-MMR*' method (Fig. S5B). The slopes were then

used to calculate MMR in mg O_2 h^{-1} using *FishResp*-package function *calculate.MR*. MMR

values calculated by the 1-min *respR* and *spline-MMR* approaches were highly correlated

(Pearson-r = 0.98, 95% confidence interval 0.98 – 0.99). We selected the spline-MMR data

for further analysis.

Statistical analyses

Data were analysed in R v. 3.6.2[64]. To test for the effects of treatment and genotype on

metabolic variables, we ran separate linear mixed models using SMR, MMR, and AS as

response variables. The response variables and body mass were $\log_{10}$-transformed to account

for allometric scaling of metabolic rate, fixed effects were centred, and covariates were

centred and scaled[65] (Table S3). We included treatment, *vgll3* and *six6* genotypes, and sex as

fixed effects in all models. For SMR, family and measurement batch were used as random

terms (models including chamber as random term were singular, and no variance was

explained by chamber). For MMR and AS, family, person performing the chase test, and

chamber identity were included as random terms. The order in which pairs of fish were tested

for MMR each day (values 1–8) was included as a covariate for MMR and AS. To test if

genotype-specific metabolic rates were affected by sex and food availability, we fitted

pairwise interactions between *vgll3* and *six6* genotypes, between genotypes and treatment,

and between genotypes and sex. The interaction of $\log_{10}$ body mass with treatment was

included to test for potential treatment-specific allometric scaling of metabolic rate.

The full models were fitted using *lme4* v. 1.1-26 [66] with an alpha value 0.05. P- and F-test

values for fixed effects were computed using type III tests with Satterthwaite's method.

Pairwise differences between significant interaction terms were obtained by post hoc analysis

using *emmeans*[67]. Residuals of models were confirmed to be homoscedastic and normally

distributed (but the residuals of SMR between the high food and low food treatments were

slightly heteroscedastic). One outlier was identified and removed from each of SMR and

MMR (using *outlierTest* in package *car,* Bonferroni-corrected p < 0.05). The proportion of

variance explained by genotypes was calculated with *partR2*[68]. Predicted means were

obtained with *ggpredict* in package *ggeffects*[69]. The data were visualized using *ggplot2*

v.3.3.3[70] and *interactions*[71]. Pearson's correlation coefficient among mass- and family-

corrected SMR, MMR and AS were calculated using residuals from a mixed model.

We also evaluated alternative models, and assess parameter significance using the corrected

Akaike information criterion score (AICc), an AIC score with a stronger penalty for complex

models[72], using the *dredge* function from package *MuMIn*[73]. We employed model averaging

if more than one model was similarly parsimonious (Table S4), in which parameter estimates

were obtained from weighted averages of all best models[74] (i.e., models with ΔAICc <2

compared to the most parsimonious models) using *model.avg* function (subset option = full)

from package *MuMIn*[73].

Results

234 Low food treatment decreased both the specific growth rate and condition factor of the fish

235 compared to high food treatment (Fig. S6). The mean body length of fish was 70.6 ± 4.5 and

236 66.2 ± 4.9 mm (SD), and the mean body mass was 4.2 ± 0.8 and 3.3 ± 0.8g after the high and

237 low food treatment, respectively.

Standard metabolic rate

There was no significant genotype, food availability or sex effect on SMR (Table 1, Fig. 2A).

There was a marginally significant interaction effect of *six6* and food availability on SMR in

the full model (p = 0.045) but this was non-significant in the averaged model (Table S5), and

none of the pairwise contrasts were significant (the largest effect being: *six6* EE-genotype,

high food vs. low food, $t_{25.6} = -2.37$, p = 0.11). The metabolic scaling exponent, *b*, i.e., the

slope of log SMR with log body mass, was marginally higher in the high food treatment

(0.94, $R^2 = 0.73$) than in the low food treatment (0.87, $R^2 = 0.69$, Fig. S7a, p = 0.047 (Table

1), p = 0.08 (Table S5)).

Maximum metabolic rate

Fish with the *vgll3* early maturation genotype had a higher MMR than fish with the late

maturation genotype (Fig. 2A, Table 1). *Vgll3* genotype also interacted with *six6*, such that

MMR was decreased when late maturation genotypes of the two loci cooccurred compared to

other genotype combinations (Fig. 2, Table 1). The genotype effects together explained ~5%

of the variance in MMR (Table S6). None of the treatment-genotype or sex-genotype

interactions or the main effects of sex or food availability had a significant effect on MMR

(Table 1, Table S5). Unlike SMR, the metabolic scaling of MMR was not significantly

affected by food treatment ($b = 0.86$, $R^2 = 0.76$).

Aerobic scope

Fish with the *vgll3* early maturation genotype had a higher AS compared to the late

maturation genotype (Fig. 2A-B, Table 1, predicted means 460.2 and 440.9 mg O_2 kg^{-1} h^{-1} for

early and late maturation genotypes, respectively). *Vgll3* was estimated to explain 1.7% of the

variance in mass-corrected AS (Table S6). AS was marginally higher under low food

availability than high food availability, but only in smaller fish (interaction p = 0.037 (Table

1) and p = 0.18 (Table S6)); scaling exponent $b = 0.94$ ($R^2 = 0.57$) in the high food and 0.90

($R^2 = 0.68$) in the low food treatment (Fig. S7b). The *vgll3* and food treatment effects were

also significant when mass adjusted SMR was included as a covariate in the model (Table

S7), indicating that the genotype effect was independent of SMR. None of the other factors

had a significant effect on AS (Table 1).

Table 1. Linear mixed models for $\log_{10}$-transformed metabolic phenotypes. All variables were centred to a mean of 0 (the category with a positive value is shown in parentheses), $\log_{10}$ body mass was scaled and centred. Significant effects shown in bold.

	Fixed effect	Estimate	SE	SSq	Den DF	F	p	Random effect	Var (95% CI)
SMR	Intercept	-0.295	0.019						
	Treatment (LF)	0.019	0.012	0.0039	17.67	2.772	0.114		
	Sex (male)	0.002	0.005	0.0002	248.11	0.172	0.679		
	Vgll3 (LL)	-0.002	0.005	0.0003	248.98	0.235	0.628		
	Six6 (LL)	0.002	0.005	0.0002	248.11	0.172	0.679		
	Log10 BM	**0.102**	**0.003**	**1.4570**	**259.04**	**1025.739**	**<0.0001**		
	Treatment (LF):log10 BM	**0.013**	**0.006**	**0.0057**	**259.25**	**3.987**	**0.047**		
	Treatment (LF):*Vgll3* (LL)	0.006	0.010	0.0006	249.25	0.410	0.523		
	Treatment (LF):Six6 (LL)	**-0.020**	**0.010**	**0.0058**	**249.21**	**4.080**	**0.044**		
	Sex (male):*Vgll3* (LL)	0.001	0.010	0.00001	248.33	0.004	0.948	Batch	0.0005 (0.0002, 0.0012)
	Sex (male):*Six6* (LL)	0.001	0.009	0.00001	247.62	0.006	0.937	Family	0.0013 (0.0003, 0.0083)
	Vgll3 (LL):*Six6* (LL)	-0.016	0.009	0.0042	248.20	2.930	0.088	Residual	0.0014
MMR	Intercept	0.333	0.013						
	Treatment (LF)	0.008	0.006	0.003	249.48	1.632	0.203		
	Sex (male)	-0.003	0.005	0.001	254.25	0.295	0.588		
	***Vgll3* (LL)**	**-0.017**	**0.005**	**0.017**	**252.06**	**10.429**	**0.001**		
	Six6 (LL)	-0.006	0.005	0.002	252.45	1.371	0.243		
	Log10 BM	**0.088**	**0.003**	**1.231**	**256.37**	**737.850**	**<0.0001**		
	Test order	-0.003	0.004	0.001	13.81	0.769	0.396		
	Treatment (LF):log10 BM	-0.010	0.007	0.004	255.47	2.252	0.135		
	Treatment (LF):*Vgll3* (LL)	-0.011	0.011	0.002	253.11	1.057	0.305		
	Treatment (LF):*Six6* (LL)	-0.010	0.011	0.001	255.77	0.840	0.360	Chamber	0.0002 (0, 0.0005)
	Sex (male):*Vgll3* (LL)	0.016	0.010	0.004	249.17	2.474	0.117	Initial	0.0001 (0, 0.0012)
	Sex (male):*Six6* (LL)	0.008	0.010	0.001	251.83	0.654	0.420	Family	0.0004 (0.0001, 0.0027)
	***Vgll3* (LL) x *Six6* (LL)**	**-0.023**	**0.010**	**0.008**	**253.67**	**5.076**	**0.025**	Residual	0.0017
AS	Intercept	0.210	0.019						
	Treatment (LF)	0.012	0.008	0.006	238.60	2.168	0.142		
	Sex (male)	-0.004	0.007	0.001	243.05	0.443	0.507		
	***Vgll3* (LL)**	**-0.016**	**0.007**	**0.015**	**240.86**	**5.653**	**0.018**		
	Six6 (LL)	-0.008	0.007	0.004	242.39	1.468	0.227		
	Log10 BM	**0.089**	**0.004**	**1.183**	**245.15**	**437.482**	**<0.0001**		
	Test order	-0.003	0.005	0.001	14.796	0.387	0.543		
	Treatment (LF):log10 BM	**-0.018**	**0.009**	**0.012**	**245.12**	**4.409**	**0.037**		
	Treatment (LF):*Vgll3* (LL)	-0.016	0.014	0.003	241.59	1.231	0.268		
	Treatment (LF):*Six6* (LL)	0.001	0.014	0.000	244.70	0.002	0.964	Chamber	0.0002 (0, 0.0008)
	Sex (male):*Vgll3* (LL)	0.023	0.014	0.008	239.76	2.802	0.095	Initial	0.0002 (0, 0.0022)
	Sex (male):*Six6* (LL)	0.009	0.014	0.001	239.97	0.487	0.486	Family	0.001 (0.0003, 0.0064)
	Vgll3 (LL) x *Six6* (LL)	-0.020	0.014	0.006	243.02	2.086	0.150	Residual	0.0027

BM = Body mass, LF = Low food

Correlations among metabolic phenotypes

There was a positive correlation between mass- and family- corrected SMR and MMR in the high food, but not the low food treatment (Fig. S8), and a very strong correlation between MMR and AS in both treatments (Table 2).

Table 2. Pearson's correlation coefficients between metabolic phenotypes in high food (above diagonal) and low food (below diagonal) treatments. P-values given in parentheses.

	rSMR	rMMR	rAbsAS
rSMR		**0.28 (0.004)**	0.13 (0.09)
rMMR	0.11 (0.16)		**0.99 (<0.001)**
rAbsAS	**-0.17 (0.04)**	**0.96 (<0.001)**	

Discussion

The timing of maturation, just as many life-history traits, depends on reaching a certain body size threshold, i.e., the acquisition of sufficient energy that can be allocated for maturation processes[38,39,75,76]. In line with our hypothesis, we found that the *vgll3* early maturation genotype increased the aerobic scope (AS) of juvenile Atlantic salmon compared to the late maturation genotype. This effect was driven by a change in maximum metabolic rate (MMR), not standard metabolic rate (SMR) (nearly all variation in AS was explained by MMR in our study, Table 2). A previous study showed that higher condition factor, mediated by the *vgll3* early maturation genotype, positively affected the initiation of male maturation[45]. The results presented here suggest that superior resource acquisition or assimilation via higher AS, driven by a higher MMR, is a potential mechanism by which an increased condition factor in individuals with the *vgll3* early maturation genotype could be achieved compared conspecifics with the late maturation genotype. In addition to differences in mean performance between *vgll3* genotypes, the two loci in our study exhibited physiological epistasis[77], as the cooccurrence of the late maturing genotypes in both loci was associated with lower MMR than their additive effects. The epistasis may help to maintain genetic variation under rapid adaptive responses[78,79].

The functional pathways that may explain the epistasis and the main effect of *vgll3* are not well known. However, both *six6* and *vgll3* are expressed during development and have been

implicated in the control of cell fate commitment and the hypothalamus-pituitary-gonad axis

in salmon[80,81]. Hence, the epistatic interaction between two genomic regions may stem from,

e.g., developmental canalisation[82]. Addressing the causal physiological and morphological

mechanisms of the link between the genomic regions and aerobic performance can shed light

into the mechanisms of life history evolution in salmon; the *vgll3* genomic region is the major

genetic axis explaining variation in age-at-maturity in salmon[43], and variation in the locus is

spatially divergent among populations and under rapid adaptive evolution[43,47,83,84].

Because of context-dependent covariation between metabolism and growth rate, whereby

high SMR improves growth under high, but not low, resource availability[27] and because

aerobic performance can be reduced under food limitation (e.g.,[8]), we tested if food

availability modified the genetic covariation between metabolic phenotypes and age-at-

maturity. Against our predictions, there was no change in SMR or MMR due to feed

restriction, nor did we find genotype-by-environment interactions, despite a strong decrease

in growth rate in low food treatment. The results indicate that the different genotypes exhibit

no plastic responses to food availability. However, a stronger feed deprivation, e.g., similar to

those occurring in the winter[85], could have induced a more pronounced effect on SMR[86,87].

Our low food treatment included approximately 3 days of fasting in between feeding to

satiation, similar to a "feast and famine" feeding strategy[88]. A lack of metabolic response to

reduced food availability may be beneficial if it allows the individual to maximize acquisition

via food assimilation during "feasting".

Unlike MMR (and consequently AS), SMR did not exhibit *vgll3*-linked covariation with age-

at-maturity. A decoupling of SMR and MMR in relation to life-history variation was also

found by Archer et al.[89] in resident and migratory brown trout (*Salmo trutta*), and high MMR,

but not SMR, was positively selected for in Atlantic salmon under high food competition in a

study by Auer et al.[90]. Further, a lack of differences in SMR across the *vgll3* genotypes was

found in our parallel study, in which the fish were smaller (mean ~1g) compared to this study

(mean ~4g)[91]. The lack of association to the age-at-maturity loci is unexpected, as SMR, or

basal metabolic rate in endotherms, has been proposed to explain life-history variation along

the fast-slow axis[5,7], but see[92]. Our results suggest that the genetic control of maturation by

the *vgll3* genomic region via MMR mostly involves physiological pathways that do not alter

SMR simultaneously. Such pathways may be related to oxygen demand by tissues or its

supply during stress and/or exhaustive exercise. For example, structural and functional

variation in the heart (i.e., cardiac output) or muscle, and mechanisms that modulate oxygen

carrying capacity of the cardiovascular system might invoke changes in MMR without

altering SMR[93-95]. However, our study does not rule out the possibility of the metabolic

phenotypes affecting variation in age-at-maturity also phenotypically or via small-effect

loci[44,96].

The differences in aerobic performance between life-history genotypes may have arisen due

to correlated selection mediated by resource acquisition: higher aerobic scope, which was

associated with early maturation, could enable higher feeding capacity[50] and improve

foraging efficiency, for example, via a shorter searching time of prey[11,85]. Further, salmon in

the wild are increasingly experiencing higher than optimal temperatures due to climate

change[97], and MMR is typically less plastic than SMR in response to environmental

temperature[15]. Thus, our results indicate a potentially important advantage for individuals

carrying the early maturation genotype under global warming, which could be mediated by a

higher appetite[98]. This advantage may also extend to higher survival during spawning

migration if the genotype effect on AS persists across life-stages[13,99,100]. It can also be

relevant to survival of salmon after spawning, and thereby repeated spawning (iteroparity),

which is co-inherited with the same *vgll3* genotype as early maturation[101]. However, the lack

of sex differences in metabolic phenotypes in our study both across and within age-at-

maturity genotypes, suggests that sex-dependent life-history variation in salmon[33] is not

reflected in metabolic rates during the juvenile stage (see also [91]).

Although a causal relationship between metabolism at the juvenile stage and age at maturity

would not be surprising, pleiotropy or linkage provide alternatives for the basis of the

observed association. For example, the *vgll3* gene encodes a transcription cofactor associated

with cell fate commitment and is expressed in many tissues[80,102], thus it is likely pleiotropic

with multiple independent functions. Similarly, several polymorphic loci with putative

functional variation are co-localised (i.e., linked) in the genomic region[43]. Finally, stage-

dependent genetic correlations in metabolism may obscure the time point that the trait

influences the life-history variation. For instance, higher aerobic scope can induce maturation

via facilitating the size attained both in the freshwater and at sea (e.g.,[19,39]), but whether

aerobic scope genetically covaries across life stages is yet to be explored.

The presence of genetic covariation between aerobic scope at the juvenile stage and age-at-

maturity at the *vgll3* genomic region suggests potential for multi-trait evolution across life-

stages, whereby selection acting on either trait would alter the phenotypic variation of the

other[4,103]. For example, if natural selection towards later age-at-maturity increases the

frequency of late maturing allele, this would constrain the aerobic scope of juveniles in the

population, even if that may be a suboptimal phenotype. On the other hand, genetic

covariation may help to maintain optimal trait variation in age-at-maturity, i.e., by limiting

potentially maladaptive environmentally induced (i.e., plastic) variation in age-at-maturity

(e.g.,[104,105]). For example, river geophysical properties are important determinants of the

optimal age structure at maturity, whereby populations in smaller tributaries have a younger,

and populations in large, fast-flowing rivers have an older age structure[33]. Forecasting age-at-

maturity from aerobic performance at earlier stages (e.g., via improved growth[22,51]) would

result in maladaptive age structure if the covariation was explained entirely by environmental

effects. However, our study was aimed at measuring the statistical association, and does not

provide estimates using a classical quantitative genetic framework (i.e., we did not quantify

environmental sources of variation, or variation due to technical or other genetic effects) or

measure evolutionary change. Therefore, partitioning biologically meaningful covariation and

quantifying the correlated response to selection were beyond the scope of this study[106].

Understanding the physiological basis of life-history variation in different life-stages and

environmental conditions can provide insights into the factors driving life-history evolution,

and hence, better predictions of the responses of populations to environmental changes. Wild

salmon populations have declined in recent decades, with a concomitant decrease in the

frequency of late maturing individuals[47,97]. Our study used an eco-physiological approach to

identify a potentially adaptive phenotype relating genetic variation and age-at-maturity in

salmon and suggests that evolution towards an earlier age-at-maturity can cause correlated

selection towards increased MMR and aerobic scope. In conclusion, the integration of age-at-

maturity and aerobic performance in the early life-stages via simple genetic mechanisms, as

shown in this study, could contribute to the diversification of ecotypes within species.

Data availability

The data are available in https://doi.org/10.5281/zenodo.5667078. R codes for the analyses

are available in an archived repository (v.1.0.2 https://doi.org/10.5281/zenodo.5783978).

Conflicts of interest

The authors declare no conflicts of interest.

Supplemental Material

Pdf-file with Material and Methods, Figures, and Tables.

References

Ricklefs, R. E. & Wikelski, M. The physiology/life-history nexus. *Trends in Ecology & Evolution* **17**, 462-468, doi:10.1016/s0169-5347(02)02578-8 (2002).

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory of ecology. *Ecology* **85**, 1771-1789, doi:10.1890/03-9000 (2004).

Dammhahn, M., Dingemanse, N. J., Niemelä, P. T. & Reale, D. Pace-of-life syndromes: a framework for the adaptive integration of behaviour, physiology and life history. *Behavioral Ecology and Sociobiology* **72**, doi:10.1007/s00265-018-2473-y (2018).

Roff, D. A. Evolutionary quantitative genetics. (Chapman and Hall, New York., 1997).

Boratynski, Z., Koskela, E., Mappes, T. & Schroderus, E. Quantitative genetics and fitness effects of basal metabolism. *Evolutionary Ecology* **27**, 301-314, doi:10.1007/s10682-012-9590-2 (2013).

Krams, I. A. *et al.* Metabolic rate associates with, but does not generate covariation between, behaviours in western stutter-trilling crickets, Gryllus integer. *Proceedings of the Royal Society B-Biological Sciences* **284**, doi:10.1098/rspb.2016.2481 (2017).

Auer, S. K., Dick, C. A., Metcalfe, N. B. & Reznick, D. N. Metabolic rate evolves rapidly and in parallel with the pace of life history. *Nature Communications* **9**, doi:10.1038/s41467-017-02514-z (2018).

Lailvaux, S. P. & Husak, J. F. The life-history of whole-organism performance. *Quarterly Review of Biology* **89**, 285-318, doi:10.1086/678567 (2014).

Fry, F. E. J. Effects of the environment on animal activity. Ontario Fisheries Research Laboratory Publication, Biol. Ser. **55**, 1-62 (1947).

Chabot, D., Steffensen, J. F. & Farrell, A. P. The determination of standard metabolic rate in fishes. *Journal of Fish Biology* **88**, 81-121, doi:10.1111/jfb.12845 (2016).

Metcalfe, N. B., Van Leeuwen, T. E. & Killen, S. S. Does individual variation in metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**, 298-321, doi:10.1111/jfb.12699 (2016).

12   Farrell, A. P. Environment, antecedents and climate change: lessons from the study of

temperature physiology and river migration of salmonids. *Journal of Experimental Biology*

212, 3771-3780, doi:10.1242/jeb.023671 (2009).

13   Clark, T. D., Jeffries, K. M., Hinch, S. G. & Farrell, A. P. Exceptional aerobic scope and

cardiovascular performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie

resilience in a warming climate. *Journal of Experimental Biology* **214**, 3074-3081,

doi:10.1242/jeb.060517 (2011).

14   Eliason, E. J., Wilson, S. M., Farrell, A. P., Cooke, S. J. & Hinch, S. G. Low cardiac and

aerobic scope in a coastal population of sockeye salmon *Oncorhynchus nerka* with a short

upriver migration. *Journal of Fish Biology* **82**, 2104-2112, doi:10.1111/jfb.12120 (2013).

15   Sandblom, E. *et al.* Physiological constraints to climate warming in fish follow principles of

plastic floors and concrete ceilings. *Nature Communications* **7**, doi:10.1038/ncomms11447

(2016).

16   Killen, S. S. *et al.* Ecological Influences and Morphological Correlates of Resting and

Maximal Metabolic Rates across Teleost Fish Species. *American Naturalist* **187**, 592-606,

doi:10.1086/685893 (2016).

17   Priede, I. G. Metabolic scope in fishes. (In: Tytler P., Calow P. (eds) Fish Energetics.

Springer, Dordrecht. https://doi.org/10.1007/978-94-011-7918-8_2, 1985).

18   Thorpe, J. E. Maturation responses of salmonids to changing developmental opportunities.

Marine Ecology Progress Series **335**, 285-288, doi:10.3354/meps335285 (2007).

19   Salminen, M. Relationships between smolt size, postsmolt growth and sea age at maturity in

Atlantic salmon ranched in the Baltic Sea. *Journal of Applied Ichthyology-Zeitschrift Fur*

Angewandte Ichthyologie **13**, 121-130, doi:10.1111/j.1439-0426.1997.tb00111.x (1997).

20   Theriault, V., Garant, D., Bernatchez, L. & Dodson, J. J. Heritability of life-history tactics and

genetic correlation with body size in a natural population of brook charr (Salvelinus

fontinalis). *Journal of Evolutionary Biology* **20**, 2266-2277, doi:10.1111/j.1420-

9101.2007.01417.x (2007).

21   Roff, D. A. The evolution of dimorphic traits - predicting the genetic correlation between

environments. *Genetics* **136**, 395-401 (1994).

22   Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J. & Metcalfe, N. B. The optimal

combination of standard metabolic rate and aerobic scope for somatic growth depends on

food availability. *Functional Ecology* **29**, 479-486, doi:10.1111/1365-2435.12396 (2015).

23   Niemelä, P. T. & Dingemanse, N. J. Meta-analysis reveals weak associations between

intrinsic state and personality. *Proceedings of the Royal Society B-Biological Sciences* **285**,

doi:10.1098/rspb.2017.2823 (2018).

24 Mathot, K. J., Dingemanse, N. J. & Nakagawa, S. The covariance between metabolic rate and
behaviour varies across behaviours and thermal types: meta-analytic insights. *Biological*
Reviews **94**, 1056-1074, doi:10.1111/brv.12491 (2019).

25 Rosenfeld, J., Richards, J., Allen, D., Van Leeuwen, T. & Monnet, G. Adaptive trade-offs in
fish energetics and physiology: insights from adaptive differentiation among juvenile
salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* **77**, 1243-1255,
doi:10.1139/cjfas-2019-0350 (2020).

26 Zeng, L. Q. *et al.* Standard metabolic rate predicts growth trajectory of juvenile Chinese
crucian carp (*Carassius auratus*) under changing food availability. *Biology Open* **6**, 1305-
1309, doi:10.1242/bio.025452 (2017).

27 Auer, S. K., Solowey, J. R., Rajesh, S. & Rezende, E. L. Energetic mechanisms for coping
with changes in resource availability. *Biology Letters* **16**, doi:10.1098/rsbl.2020.0580 (2020).

28 Killen, S. S., Marras, S. & McKenzie, D. J. Fuel, fasting, fear: routine metabolic rate and food
deprivation exert synergistic effects on risk-taking in individual juvenile European sea bass.
Journal of Animal Ecology **80**, 1024-1033, doi:10.1111/j.1365-2656.2011.01844.x (2011).

29 Auer, S. K. *et al.* Metabolic Rate Interacts with Resource Availability to Determine Individual
Variation in Microhabitat Use in the Wild. *American Naturalist* **196**, 132-144,
doi:10.1086/709479 (2020).

30 Hughes, K. A., Rodd, F. H. & Reznick, D. N. Genetic and environmental effects on secondary
sex traits in guppies (Poecilia reticulata). *Journal of Evolutionary Biology* **18**, 35-45,
doi:10.1111/j.1420-9101.2004.00806.x (2005).

31 Gutteling, E. W. *et al.* Environmental influence on the genetic correlations between life-
history traits in Caenorhabditis elegans. *Heredity* **98**, 206-213, doi:10.1038/sj.hdy.6800929
(2007).

32 Gillespie, J. H. & Turelli, M. Genotype-environment interactions and the maintenance of
polygenic variation. *Genetics* **121**, 129-138 (1989).

33 Fleming, I. A. & Einum, S. Reproductive Ecology: A Tale of Two Sexes. in *Atlantic Salmon*
Ecology 33-65 (2011).

34 Mobley, K. B. *et al.* Maturation in Atlantic salmon (Salmo salar, Salmonidae): a synthesis of
ecological, genetic, and molecular processes. *Reviews in Fish Biology and Fisheries*,
doi:10.1007/s11160-021-09656-w (2021).

35 Hendry, A. P., Berg, O. K. & Quinn, T. P. Condition dependence and adaptation-by-time:
breeding date, life history, and energy allocation within a population of salmon. *Oikos* **85**,
499-514, doi:10.2307/3546699 (1999).

36 Stearns, S. C. Life history evolution: successes, limitations, and prospects.
Naturwissenschaften **87**, 476-486, doi:10.1007/s001140050763 (2000).

37 Friedland, K. D. & Haas, R. E. Marine post-smolt growth and age at maturity of Atlantic
salmon. *Journal of Fish Biology* **48**, 1-15, doi:10.1111/j.1095-8649.1996.tb01414.x (1996).

38 Skilbrei, O. T. Relationships between smolt length and growth and maturation in the sea of
individually tagged Atlantic salmon (*Salmo salar*). *Aquaculture* **83**, 95-108,
doi:10.1016/0044-8486(89)90064-1 (1989).

39 Hutchings, J. A. & Jones, M. E. B. Life history variation and growth rate thresholds for
maturity in Atlantic salmon, *Salmo salar*. *Canadian Journal of Fisheries and Aquatic*
Sciences **55**, 22-47, doi:10.1139/cjfas-55-S1-22 (1998).

40 Tréhin, C. *et al.* Growth during the first summer at sea modulates sex-specific maturation
schedule in Atlantic salmon. *Canadian Journal of Fisheries and Aquatic Sciences* **78**, 659-
669, doi:10.1139/cjfas-2020-0236 (2021).

41 Moore, M. P. & Martin, R. A. On the evolution of carry-over effects. *Journal of Animal*
Ecology **88**, 1832-1844, doi:10.1111/1365-2656.13081 (2019).

42 Ayllon, F. *et al.* The vgll3 Locus Controls Age at Maturity in Wild and Domesticated Atlantic
Salmon (*Salmo salar* L.) Males. *Plos Genetics* **11**, doi:10.1371/journal.pgen.1005628 (2015).

43 Barson, N. J. *et al.* Sex-dependent dominance at a single locus maintains variation in age at
maturity in salmon. *Nature* **528**, 405-408, doi:10.1038/nature16062 (2015).

44 Sinclair-Waters, M. *et al.* Beyond large-effect loci: large-scale GWAS reveals a mixed large-
effect and polygenic architecture for age at maturity of Atlantic salmon. *Genetics Selection*
Evolution **52**, doi:10.1186/s12711-020-0529-8 (2020).

45 Debes, P. V. *et al.* Polygenic and major-locus contributions to sexual maturation timing in
Atlantic salmon. *Molecular Ecology*, doi:10.1111/mec.16062 (2021).

46 Chaput, G. Overview of the status of Atlantic salmon (*Salmo salar*) in the North Atlantic and
trends in marine mortality. *Ices Journal of Marine Science* **69**, 1538-1548,
doi:10.1093/icesjms/fss013 (2012).

47 Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C. R. Rapid sex-specific
evolution of age at maturity is shaped by genetic architecture in Atlantic salmon. *Nature*
Ecology & Evolution **2**, 1800-1807, doi:10.1038/s41559-018-0681-5 (2018).

48 Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C. Evolution in salmon life-
history induced by direct and indirect effects of fishing. *bioRxiv*,
doi:10.1101/2021.01.08.425869 (2021).

49 Aykanat, T. *et al.* Life-history genomic regions explain differences in Atlantic salmon marine
diet specialization. *Journal of Animal Ecology* **89**, 2677-2691, doi:10.1111/1365-2656.13324
(2020).

50 Auer, S. K., Salin, K., Anderson, G. J. & Metcalfe, N. B. Aerobic scope explains individual
variation in feeding capacity. *Biology Letters* **11**, doi:10.1098/rsbl.2015.0793 (2015).

51 Norin, T. & Clark, T. D. Fish face a trade-off between 'eating big' for growth efficiency and
'eating small' to retain aerobic capacity. *Biology Letters* **13**, doi:10.1098/rsbl.2017.0298
(2017).
52 Elliott, J. M. & Hurley, M. A. A functional model for maximum growth of Atlantic Salmon
parr, Salmo salar, from two populations in northwest England. *Functional Ecology* **11**, 592-
603, doi:10.1046/j.1365-2435.1997.00130.x (1997).
53 Ward, A. J. W., Webster, M. M. & Hart, P. J. B. Intraspecific food competition in fishes. *Fish*
and Fisheries **7**, 231-261, doi:10.1111/j.1467-2979.2006.00224.x (2006).
54 Rosenfeld, J., Van Leeuwen, T., Richards, J. & Allen, D. Relationship between growth and
standard metabolic rate: measurement artefacts and implications for habitat use and life-
history adaptation in salmonids. *Journal of Animal Ecology* **84**, 4-20, doi:10.1111/1365-
2656.12260 (2015).
55 Forstner, H. An automated multiple-chamber intermittent-flow respirometer. (Springer Berlin
Heidelberg), 111-126 (1983).
56 Svendsen, M. B. S., Bushnell, P. G. & Steffensen, J. F. Design and setup of intermittent-flow
respirometry system for aquatic organisms. *Journal of Fish Biology* **88**, 26-50,
doi:10.1111/jfb.12797 (2016).
57 Killen, S. S. *et al.* Guidelines for reporting methods to estimate metabolic rates by aquatic
intermittent-flow respirometry. *Journal of Experimental Biology* **224**, doi:10.1242/jeb.242522
(2021).
58 Raby, G. D., Doherty, C. L. J., Mokdad, A., Pitcher, T. E. & Fisk, A. T. Post-exercise
respirometry underestimates maximum metabolic rate in juvenile salmon. *Conservation*
Physiology **8**, doi:10.1093/conphys/coaa063 (2020).
59 Brett, J. R. The respiratory metabolism and swimming performance of young Sockeye
salmon. *Journal of the Fisheries Research Board of Canada* **21**, 1183-1226 (1964).
60 Norin, T. & Clark, T. D. Measurement and relevance of maximum metabolic rate in fishes.
Journal of Fish Biology **88**, 122-151, doi:10.1111/jfb.12796 (2016).
61 Morozov, S., McCairns, R. J. S. & Merilä, J. FishResp: R package and GUI application for
analysis of aquatic respirometry data. *Conservation Physiology* **7**,
doi:10.1093/conphys/coz003 (2019).
62 Harianto, J., Carey, N. & Byrne, M. respR-An R package for the manipulation and analysis of
respirometry data. *Methods in Ecology and Evolution* **10**, 912-920, doi:10.1111/2041-
210x.13162 (2019).
63 Little, A. G. *et al.* Maxed Out: Optimizing Accuracy, Precision, and Power for Field
Measures of Maximum Metabolic Rate in Fishes. *Physiological and Biochemical Zoology* **93**,
243-254, doi:10.1086/708673 (2020).
64 R Core Team.   (R Foundation for Statistical Computing, Vienna, Austria, 2019).

65   Schielzeth, H. Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution **1**, 103-113, doi:10.1111/j.2041-210X.2010.00012.x
(2010).

66   Bates, D., Machler, M., Bolker, B. M. & Walker, S. C. Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* **67**, 1-48, doi:10.18637/jss.v067.i01 (2015).

67   Lenth, R. V.   (R package version 1.5.3, 2020).

68   Stoffel, M. A., Nakagawa, S. & Schielzeth, H. partR2: partitioning R-2 in generalized linear
mixed models. *Peerj* **9**, doi:10.7717/peerj.11414 (2021).

69   Lüdecke, D. *et al.* Vol. 3   (Journal of Open Source Software, 2018).

70   Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Ggplot2: Elegant Graphics for*
Data Analysis, 1-212, doi:10.1007/978-0-387-98141-3 (2009).

71   Long, J. A. interactions: Comprehensive, User-Friendly Toolkit for Probing Interactions.
(2019).

72   Hurvich, C. M. & Tsai, C. L. Regression and time-series model selection in small samples.
Biometrika **76**, 297-307, doi:10.1093/biomet/76.2.297 (1989).

73   Bartoń, K. MuMIn: Multi-Model Inference. R package version 1.43.17. URL:
https://CRAN.R-project.org/package=MuMIn (2020).

74   Johnson, J. B. & Omland, K. S. Model selection in ecology and evolution. *Trends in Ecology*
& Evolution **19**, 101-108, doi:10.1016/j.tree.2003.10.013 (2004).

75   Weimerskirch, H. Reproductive effort in long-lived birds – Age-specific patterns of
condition, reproduction and survival in the Wandering albatross. *Oikos* **64**, 464-473,
doi:10.2307/3545162 (1992).

76   Cheung, C. C. *et al.* Leptin is a metabolic gate for the onset of puberty in the female rat.
Endocrinology **138**, 855-858, doi:10.1210/en.138.2.855 (1997).

77   Cheverud, J. M. & Routman, E. J. Epistasis and its contribution to genetic variance
components. *Genetics* **139**, 1455-1461 (1995).

78   Cheverud, J. M. & Routman, E. J. Epistasis as a source of increased additive genetic variance
at population bottlenecks. *Evolution* **50**, 1042-1051, doi:10.2307/2410645 (1996).

79   Merilä, J. & Sheldon, B. C. Genetic architecture of fitness and nonfitness traits: empirical
patterns and development of ideas. *Heredity* **83**, 103-109, doi:10.1046/j.1365-
2540.1999.00585.x (1999).

80   Kurko, J. *et al.* Transcription Profiles of Age-at-Maturity-Associated Genes Suggest Cell Fate
Commitment Regulation as a Key Factor in the Atlantic Salmon Maturation Process. *G3-*
Genes Genomes Genetics **10**, 235-246, doi:10.1534/g3.119.400882 (2020).

81   Moustakas-Verho, J. E. *et al.* Developmental expression patterns of six6: A gene linked with
spawning ecotypes in Atlantic salmon. *Gene Expression Patterns* **38**,
doi:10.1016/j.gep.2020.119149 (2020).

82 Debat, V. & David, P. Mapping phenotypes: canalization, plasticity and developmental
stability. *Trends in Ecology & Evolution* **16**, 555-561, doi:10.1016/s0169-5347(01)02266-2
(2001).

83 Pritchard, V. L. *et al.* Genomic signatures of fine-scale local selection in Atlantic salmon
suggest involvement of sexual maturation, energy homeostasis and immune defence-related
genes. *Molecular Ecology* **27**, 2560-2575, doi:10.1111/mec.14705 (2018).

84 Zueva, K. J., Lumme, J., Veselov, A. E., Primmer, C. R. & Pritchard, V. L. Population
genomics reveals repeated signals of adaptive divergence in the Atlantic salmon of north-
eastern Europe. *Journal of Evolutionary Biology* **34**, 866-878, doi:10.1111/jeb.13732 (2021).

85 Johansen, M., Erkinaro, J. & Amundsen, P. A. The When, What and Where of Freshwater
Feeding, in *Atlantic Salmon Ecology*, Wiley-Blackwell (2011).

86 O'Connor, K. I., Taylor, A. C. & Metcalfe, N. B. The stability of standard metabolic rate
during a period of food deprivation in juvenile Atlantic salmon. *Journal of Fish Biology* **57**,
41-51, doi:10.1006/jfbi.2000.1280 (2000).

87 Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J. & Metcalfe, N. B. Flexibility in
metabolic rate confers a growth advantage under changing food availability. *Journal of
Animal Ecology* **84**, 1405-1411, doi:10.1111/1365-2656.12384 (2015).

88 Armstrong, J. B. & Schindler, D. E. Excess digestive capacity in predators reflects a life of
feast and famine. *Nature* **476**, 84-87, doi:10.1038/nature10240 (2011).

89 Archer, L. C. *et al.* Metabolic traits in brown trout (Salmo trutta) vary in response to food
restriction and intrinsic factors. *Conservation Physiology* **8**, doi:10.1093/conphys/coaa096
(2020).

90 Auer, S. K. *et al.* Nutrients from salmon parents alter selection pressures on their offspring.
Ecology Letters **21**, 287-295, doi:10.1111/ele.12894 (2018).

91 Åsheim, E. R., Prokkola, J. M., Morozov, S., Aykanat, T. & Primmer, C. R. Standard
metabolic rate does not associate with age-at-maturity genotype in juvenile Atlantic salmon.
Ecology and Evolution (in press).

92 Polverino, G., Santostefano, F., Diaz-Gil, C. & Mehner, T. Ecological conditions drive pace-
of-life syndromes by shaping relationships between life history, physiology and behaviour in
two populations of Eastern mosquitofish. *Scientific Reports* **8**, doi:10.1038/s41598-018-
33047-0 (2018).

93 Harter, T. S., Zanuzzo, F. S., Supuran, C. T., Gamperl, A. K. & Brauner, C. J. Functional
support for a novel mechanism that enhances tissue oxygen extraction in a teleost fish.
Proceedings of the Royal Society B-Biological Sciences **286**, doi:10.1098/rspb.2019.0339
(2019).

94   Nikinmaa, M., Berenbrink, M. & Brauner, C. J. Regulation of erythrocyte function: Multiple

evolutionary solutions for respiratory gas transport and its regulation in fish. *Acta*

Physiologica **227**, doi:10.1111/apha.13299 (2019).

95   McArley, T. J., Morgenroth, D., Zena, L. A., Ekstrom, A. T. & Sandblom, E. Normoxic

limitation of maximal oxygen consumption rate, aerobic scope and cardiac performance in

exhaustively exercised rainbow trout (Oncorhynchus mykiss). *Journal of Experimental*

Biology **224**, doi:10.1242/jeb.242614 (2021).

96   Johnston, S. E. *et al.* Genome-wide SNP analysis reveals a genetic basis for sea-age variation

in a wild population of Atlantic salmon (Salmo salar). *Molecular Ecology* **23**, 3452-3468,

doi:10.1111/mec.12832 (2014).

97   Friedland, K. D. *et al.* The recruitment of Atlantic salmon in Europe. *Ices Journal of Marine*

Science **66**, 289-304, doi:10.1093/icesjms/fsn210 (2009).

98   Jutfelt, F. *et al.* 'Aerobic scope protection' reduces ectotherm growth under warming.

Functional Ecology **35**, 1397-1407, doi:10.1111/1365-2435.13811 (2021).

99   Eliason, E. J. *et al.* Differences in Thermal Tolerance Among Sockeye Salmon Populations.

Science **332**, 109-112, doi:10.1126/science.1199158 (2011).

100   Mottola, G., Kristensen, T. & Anttila, K. Compromised thermal tolerance of cardiovascular

capacity in upstream migrating Arctic char and brown trout-are hot summers threatening

migrating salmonids? *Conservation Physiology* **8**, doi:10.1093/conphys/coaa101 (2020).

101   Aykanat, T. *et al.* Co-inheritance of sea age at maturity and iteroparity in the Atlantic salmon

vgll3 genomic region. *Journal of Evolutionary Biology* **32**, 343-355, doi:10.1111/jeb.13418

(2019).

102   Verta, J. P. *et al.* Cis-regulatory differences in isoform expression associate with life history

strategy variation in Atlantic salmon. *Plos Genetics* **16**, doi:10.1371/journal.pgen.1009055

(2020).

103   Lande, R. Quantitative genetic analysis of multivariate evolution, applied to brain - body size

allometry. *Evolution* **33**, 402-416, doi:10.2307/2407630 (1979).

104   De Jong, G. Unpredictable selection in a structured population leads to local genetic

differentiation in evolved reaction norms. *Journal of Evolutionary Biology* **12**, 839-851,

doi:10.1046/j.1420-9101.1999.00118.x (1999).

105   Tufto, J. The evolution of plasticity and nonplastic spatial and temporal adaptations in the

presence of imperfect environmental cues. *American Naturalist* **156**, 121-130,

doi:10.1086/303381 (2000).

106   Sgro, C. M. & Hoffmann, A. A. Genetic correlations, tradeoffs and environmental variation.

Heredity **93**, 241-248, doi:10.1038/sj.hdy.6800532 (2004).

Figure captions

Fig. 1. Timeline of the experiment. The duration of low food and high food treatments was

four weeks. "W" indicates when fish density of high food tanks was reduced to the level of

low food tanks. Each horizontal line represents a separate tank. Blue blocks represent

procedures (W. & L. = weight and length, and/or SMR & MMR measurements).

Fig 2. A) Predicted means for standard metabolic rate (SMR), maximum metabolic rate

(MMR), and aerobic scope (AS) in *vgll3* and *six6* early- and late-maturation genotypes with

90% confidence intervals. The means are for average body mass, treatment, and sex effects,

and back transformed to linear scale. P-values show significant pairwise differences between

genotypes for MMR on top of the points, and for the *vgll3* main effects (Table 1) in MMR

and AS between the points. N = 60–71 in each genotype combination (same individuals used

for all traits). B) Residual aerobic scope from a linear mixed model including $\log_{10}$ aerobic

scope as response, $\log_{10}$ body mass as predictor and family as random term, showing

individuals (by points) in each genotype combination.

Week 1 | Week 2 | Week 3 | Week 4 | Week 5 | Week 6 | Week 7 | Week 8

W. & L.
Low food treatment
Family 1
SMR & MMR
Family 2
Family 3
Family 4

W. & L.
High food treatment
Family 1
SMR & MMR
Family 2
Family 3
Family 4

A) Predicted mean, mg O$_2$/kg/h (90% CI)

B) Residual aerobic scope