Genetic coupling of life-history and aerobic performance in Atlantic salmon indicates potential for coevolution across life-stages

Short title: Genetic coupling of life-history and metabolism

Jenni M. Prokkola[1]*, Eirik R. Åsheim[1], Sergey Morozov[1], Paul Bangura[1], Jaakko Erkinaro[2], Annukka Ruokolainen[1], Craig R. Primmer[1,3], Tutku Aykanat[1]

[1]Organismal and Evolutionary Biology Research Programme, University of Helsinki, Finland

[2]Natural Resources Institute Finland

[3]Institute of Biotechnology, Helsinki Institute of Life Science (HiLIFE), University of Helsinki, Finland

*Corresponding author, PO Box 65, FI-00014 University of Helsinki, Finland, jenni.prokkola@helsinki.fi

Acknowledgements

We thank Nikolai Piavchenko, Katja Maamela, Markus Lauha, Petra Liljeström, Suvi Ikonen, Mikko Immonen, and Anna Toikkanen for fish care and technical help during the experiment, and Lammi Biological Station for a high-quality research environment. Dr. Markus Haapala and the Nordic University Hub NordicPOP (Nordforsk, project no. 85352) are acknowledged for the 3D printing service, and Heidrikur Bergsson for sharing the chamber cap model. The study was funded by Academy of Finland (T. Aykanat: 325964, 1328860, C. R. Primmer: 314254 and 314255, 327255), the European Research Council under the European Articles Union's Horizon 2020 research and innovation program (grant no. 742312), and the University of Helsinki.

Keywords: age-at-maturity, energetics, metabolic rate, life history evolution

Author contributions

JMP, ERÅ, SM, CRP and TA conceived the ideas and designed methodology; JMP, ERÅ, SM, PB,

JE, AR and TA collected the data; JMP, SM and TA analysed the data; JMP and TA led the writing

of the manuscript. All authors contributed to the drafts and gave final approval for publication.

Abstract

A better understanding of the genetic and phenotypic architecture underlying life-history variation

is a longstanding aim in biology. Theories suggest energy metabolism determines life-history

variation by modulating resource acquisition and allocation trade-offs, but the genetic

underpinnings of the relationship and its dependence on ecological conditions have rarely been

demonstrated. The strong genetic determination of age-at-maturity by two unlinked genomic

regions (*vgll3* and *six6*) makes Atlantic salmon (*Salmo salar*) an ideal model to address these

questions. Using more than 250 juveniles in common garden conditions, we quantified the

covariation between metabolic phenotypes –standard and maximum metabolic rates (SMR and

MMR), and aerobic scope (AS) – and the life-history genomic regions and tested if food availability

modulates the relationships. We found that the early maturation genotype in *vgll3* was associated

with higher MMR and consequently AS. Additionally, MMR exhibited physiological epistasis; it

was decreased when late maturation genotypes co-occurred in both genomic regions. Contrary to

our expectation, the life-history genotypes had no effects on SMR. Further, food availability had no

effect on the genetic covariation, suggesting a lack of genotype-by-environment interactions. Our

results provide insights on the key organismal processes that link energy use at the juvenile-stage to

age-at-maturity, indicating potential mechanisms by which metabolism and life-history can

coevolve.

Introduction

Physiological processes control how life-history diversity emerges from resource allocation and acquisition trade-offs[1]. The rate of aerobic energy metabolism is a pivotal mechanism contributing to life-history variation – it modulates resource acquisition, provides cells with ATP, and constrains energy allocation to different body components and functions. Theories, such as the metabolic theory of ecology and the pace-of-life syndrome theory[2,3], suggest metabolic rate covaries with life-history variation within and among species. This covariation may have a genetic basis, consequently constraining trait evolution[4], yet only a few studies have demonstrated intraspecific genetic covariation or co-evolution between metabolic rate and life-history traits[5-7]. Determining whether this relation is modified by different ecological contexts (e.g., food availability) is crucial to better understand the mechanisms shaping life history variation and demographic shifts in populations in response to environmental changes[8].

The quintessential components of energy metabolism at the organismal level (i.e., the metabolic phenotypes) are standard metabolic rate (SMR), maximum metabolic rate (MMR), and the absolute aerobic scope (AS) that is the difference between SMR and MMR[9-11]. SMR is the minimum metabolic rate of an ectothermic animal associated with self-maintenance, and therefore defines the minimal cost of living (excluding growth, digestion, and locomotion). MMR defines the upper limit of aerobic performance that is functionally linked to SMR and the capacity to increase oxygen uptake and delivery beyond SMR. SMR and MMR together are the integral components of AS, which is the measure of surplus energy that can be allocated into non-maintenance functions, such as locomotion and digestion [12]. Higher AS is predicted to increase fitness via facilitating energetically demanding behaviours (such as migration, aggression, predator avoidance, and prey capture) and tolerance to environmental stress[11,13-15]. However, high aerobic performance comes with costs, including maintaining a larger heart and gill surface area (associated with increased demand for osmoregulation)[12,16,17].

Allocation of energy to growth or improved condition can link metabolic phenotypes to life-history traits[18]. Life-history traits, such as the timing of maturation and migration, are determined by adaptive body-size thresholds[19-21], and metabolic phenotypes are often correlated with growth rate, albeit in a context-dependent manner[11]. Under high food availability, a high SMR in combination with high AS can increase growth rate[22], as it often correlates with traits that improve resource acquisition, such as dominance and digestive capacity[23-25]. Under low food availability, the growth benefit of high SMR or AS can be minimized (or even reversed for SMR) due to high self-

maintenance costs[22,26,27]. In addition, individuals can be forced to seek new habitats or take more

risks to acquire resources, exerting further fitness costs[28,29]. Hence, covariation between life-history

and metabolism, whether it is genetically or environmentally driven, could be modulated by

resource availability. A resource dependent change in genetic covariation, i.e., genotype-by-

environment interaction, could maintain genetic variation in these traits[30-32], but this has not been

demonstrated.

In anadromous (sea-migrating) salmonids, the number of years the fish spends at sea before the first

spawning, i.e., sea age-at-maturity, has a dramatic effect on its size-at-maturity[33]. Individuals

spending one year at sea typically weigh 1–3 kg compared with 10–20 kg after three or more years.

Increased size in late maturing individuals also translates to marked gains in reproductive

investment in both sexes[33]. Earlier maturation, i.e., less time spent at sea, provides a potential

fitness advantage through a higher probability of survival prior to reproduction and a shorter

generation time, but comes at the expense of decreased fecundity and mating success (due to

smaller size at reproduction[34-36]. The probability of early maturation at sea is also positively

associated with faster growth and fat deposition in the freshwater (juvenile) stage, since the size of

salmon at the onset of sea-migration has a significant influence on maturation timing at sea[19,37-40].

These relationships suggest that early maturation in salmon may be associated with higher SMR or

aerobic scope via resource utilization already in early life-stages prior to sea migration[24,25]. In

addition, metabolic phenotypes in the juvenile stage may explain maturation at sea through genetic

correlations across life stages (e.g.,[41]) if metabolic rate or performance at sea is linked to earlier

maturation.

In Atlantic salmon (*Salmo salar* L. 1758), a large proportion of variation in age-at-maturity is

explained by a single genomic region that encompasses the *vgll3* gene on chromosome 25 [42,43]. In

addition, variation in another locus on chromosome 9, *six6*, is a strong predictor of mean age-at-

maturity among populations [43] and associated with early maturation in aquaculture salmon[44]. *Vgll3*

and *six6* are also associated with size-at-maturity, with the alleles conferring late maturation being

associated with larger age-specific body size especially after multiple years at sea[43]. Moreover,

vgll3 is associated with precocious maturation in male salmon parr via body condition[45],

emphasizing the causal effect of energy acquisition on maturation. In the last few decades, many

Atlantic salmon populations have been maturing, on average, at younger ages[46], which is associated

with a change in *vgll3* allele frequency in some cases[47]. Recently, a link was found between the

decrease in salmon age-at-maturity and a change in prey species composition[48] (see also [49] for diet

composition in relation to *six6*). These observations further highlight that genotype dependent

differences in SMR or aerobic performance, which can contribute to foraging success or food

assimilation (e.g.,[22,50,51]), may be related to contemporary life-history evolution in Atlantic salmon.

The strong effects of the *six6* and *vgll3* genomic regions on life-history variation provide an

opportunity for the genetic covariation between age-at-maturity and energy metabolism to be

studied prior to maturation, i.e., at the juvenile stage, by genetic prediction. This approach makes

controlled, empirical settings more feasible, as salmon require several years to reach maturation. In

this study, we test if genetic covariation exists between life-history and metabolic phenotypes in

juveniles; we hypothesize that early maturation genotypes show a higher metabolic activity (SMR,

MMR, and AS) than late maturation genotypes under high food availability (e.g.,[7]). Further,

because low resource availability weakens the relationship between SMR and growth[27] and may

constrain aerobic performance[8], we also explore if food availability modulates the genetic

covariation.

Material and Methods

137 The experiments were conducted under an animal experiment permit granted by the Finnish Project

138 Authorisation Board (permit nr. ESAVI/4511/2020). Additional experimental details are provided

139 in online supplemental material.

Fish rearing and genotyping

The parental Atlantic salmon (Table S1) were first-generation hatchery brood stock (from the river

Kymijoki in Finland) from Laukaa hatchery, managed by the Natural Resources Institute Finland

(LUKE). In October 2019, eggs and milt were transferred to the University of Helsinki for

fertilisation. Full-sib families were crossed using heterozygous parents in *vgll3* and *six6* loci,

(*vgll3**E/L and *six6**E/L where E and L refer to the alleles associated with early and late

maturation, respectively). This provides offspring with all genotype combinations within each full-

sib family. Feed rations were calculated assuming feed conversion efficiency of 0.8, using growth

predictions by Elliott and Hurley[52]. In July 2020, fish were tagged with passive integrated

transponder (PIT) tags and genotyped from fin clips.

Experimental design

Food treatments were started in August 2020. Briefly, fish were fed twice week^{-1} in the low food

treatment (Fig. 1). This was preferred over constant low ration to minimize dominance hierarchies

in tanks[53]. Fish in the high food treatment were fed with the total ad-libitum daily ration. After 28–

31d in the treatments, 48 and 32 fish from each family in the low and high food treatment,

respectively (192 and 128 in total), were measured for their SMR and MMR (Fig. 1), though only

290 homozygous individuals were used in the analysis (Table S2).

Fig. 1. Timeline of the experiment. The duration of low food and high food treatments was four

weeks. "W" indicates when fish density of high food tanks was reduced to the level of low food

tanks. Each horizontal line represents a separate tank. Blue blocks represent procedures (W. & L. =

weight and length, and/or SMR & MMR measurements).

SMR and MMR measurements

Sixteen fish at once were moved into an acclimation tank two days before SMR measurement (Fig.

S3). Each batch was from the same family and tank and balanced for genotype-sex-combinations.

During acclimation, the fish were held individually, without feeding at $11°C \pm 0.1°C$ in

$20×20×10cm$ cages, to minimize the effects of temperature fluctuations, digestion, growth, and

social interactions on SMR[54]. We measured SMR using intermittent flow respirometry[55-57]. The

SMR measurements commenced after 42–47h acclimation, between 11:30 and 14:30, until 8:00 the

following day. Afterwards, MMR was measured using the chase method, similarly to, e.g.,[58], where

MMR reflects increased aerobic respiration related to exercise and the oxygen debt incurred by

anaerobic respiration [59,60]. Fish were then euthanized with an overdose of methanesulfonate,

measured, and weighed. All fish were confirmed to have immature gonads.

Analysis of respirometry data

For SMR, oxygen consumption rate (MO_2, mg O_2 h^{-1}) for each linear measurement phase was

derived from best-fit linear regression of dissolved oxygen concentration over time. The mean of

the lowest normal distribution (MLND) was used to estimate SMR for each individual as mg O_2 h^{-1}

from the extracted MO_2 slopes[10].

MMR was calculated for each individual as mg O_2 h^{-1} from the O_2 concentrations after performing

background correction in *FishResp*[61]. We used two methods to identify the slope of the steepest

decrease in O_2 saturation. First, we used $respR$[62] with the function $auto\_rate$, fitting one and two-

minute windows[63] (example slope in Fig. S5A). Second, slopes for MMR were extracted using a

derivative of a polynomial curve fitted on each measurement (function $smooth.spline,$ df=10). This

is the '$spline$-MMR' method (Fig. S5B). The slopes were then used to calculate MMR in mg O_2 h^{-1}

using $FishResp$-package function $calculate.MR$. MMR values calculated by the 1-min $respR$ and

$spline$-MMR approaches were highly correlated (Pearson-r = 0.98, 95% confidence interval 0.98 –

0.99). We selected the spline-MMR data for further analysis.

Statistical analyses

To test for the effects of treatment and genotype on metabolic variables, we ran separate linear

mixed models using SMR, MMR, and AS as response variables. The response variables and body

mass were log_{10}-transformed to account for allometric scaling of metabolic rate, fixed effects were

centred, and covariates were centred and scaled[64] (Table S3). We included treatment, $vgll3$ and $six6$

genotypes, and sex as fixed effects in all models. For SMR, family and measurement batch were

used as random terms (models including chamber as random term were singular, and no variance

was explained by chamber). For MMR and AS, family, person performing the chase test, and

chamber identity were included as random terms. The order in which pairs of fish were tested for

MMR each day (values 1–8) was included as a covariate in the full MMR and AS models but

excluded from final models due to lack of significance. To test if genotype-specific metabolic rates

were affected by sex and food availability, we fitted pairwise interactions between $vgll3$ and $six6$

genotypes, between genotypes and treatment, and between genotypes and sex. The interaction of

log_{10} body mass with treatment was included to test for potential treatment-specific allometric

scaling of metabolic rate. We employed stepwise backward model selection by omitting non-

significant interactions based on Type III test p-values with Satterthwaite's method. Pairwise

differences between genotypes in the significant interaction terms were obtained by post hoc

analysis using $emmeans$[65].

All models were fitted using $lme4$ v. 1.1-26 [66] in R v. 3.6.2[67] with an alpha-value 0.05. P- and F-test

values for fixed effects were computed using type III tests with Satterthwaite's method. Residuals of

models were confirmed to be homoscedastic and normally distributed (but the residuals of SMR

between the high food and low food treatments were slightly heteroscedastic). One outlier was

identified and removed from each of SMR and MMR (using $outlierTest$ in package $car,$ Bonferroni-

corrected p < 0.05). The proportion of variance explained by genotypes was calculated with

$partR2$[68]. Predicted means were obtained with $ggpredict$ in package $ggeffects$[69]. The data were

visualized using *ggplot2* v.3.3.3[70] and *interactions*[71]. Pearson's correlation coefficient among mass-
and family-corrected SMR, MMR and AS were calculated using residuals from a mixed model.

Results

222 Low food treatment decreased both the specific growth rate and condition factor of the fish
223 compared to high food treatment (Fig. S6). The mean body length of fish was 70.6 ± 4.5 and $66.2 \pm$
224 4.9 mm (SD), and the mean body mass was 4.2 ± 0.8 and 3.3 ± 0.8g after the high and low food
225 treatment, respectively.

Standard metabolic rate

There was no significant genotype, food availability or sex effect on SMR (Table 1, Fig. 2A). There
was a marginally significant interaction effect of *six6* and food availability on SMR (p = 0.045 in
the full model, p = 0.055 in the simplified model, Table S4), but none of the pairwise contrasts were
significant (the largest effect being: *six6* EE-genotype, high food vs. low food, $t_{25.6} = -2.37$, p =
0.11). The metabolic scaling exponent, *b*, i.e., the slope of log SMR with log body mass, was
significantly higher in the high food treatment (0.94, $R^2 = 0.73$) than in the low food treatment
(0.87, $R^2 = 0.69$) (Fig. S7a, p = 0.025, Table 1).

Maximum metabolic rate

Fish with the *vgll3* early maturation genotype had a higher MMR than fish with the late maturation
genotype (Fig. 2A, Table 1). *Vgll3* genotype also interacted with *six6*, such that MMR was
decreased when late maturation genotypes of the two loci cooccurred compared to other genotype
combinations (Fig. 2, Table 1). The genotype effects together explained ~4.4% of the variance in
MMR (Table S8). None of the treatment-genotype or sex-genotype interactions or the main effects
of sex or food availability had a significant effect on MMR (Table S5). Unlike SMR, the metabolic
scaling of MMR was not significantly affected by food treatment ($b = 0.86$, $R^2 = 0.76$).

Aerobic scope

Fish with the *vgll3* early maturation genotype had a higher AS compared to the late maturation
genotype (Fig. 2A-B, Table 1, predicted means 460.7 and 441.7 mg O_2 kg^{-1} h^{-1} for early and late
maturation genotypes, respectively). *Vgll3* was estimated to explain 1.7% of the variance in mass-
corrected AS, with an upper confidence limit close to 8% (Table S8). AS was marginally higher
under low food availability than high food availability, but only in smaller fish (interaction p =
0.054, Table 1); scaling exponent $b = 0.94$ ($R^2 = 0.57$) in the high food and 0.90 ($R^2 = 0.68$) in the
low food treatment (Fig. S7b). The *vgll3* and food treatment effects were also significant when mass

adjusted SMR was included as a covariate in the model (Table S6), indicating that the genotype

effect was independent of SMR. None of the other factors had a significant effect on AS (Table S7).

Table 1. Simplified linear mixed models for log10-transformed metabolic phenotypes. Full models are shown in online supplemental material. All variables were centred to a mean of 0 (the category with a positive value is shown in parentheses), log10 body mass was scaled and centred. Significant effects shown in bold.

	Fixed effect	Estimate	SE	SSq	Den DF	F	P	Random effect	Var (95% CI)
SMR	Intercept	-0.295	0.019						
	Treatment (LF)	0.019	0.012	0.0040	17.52	2.75	0.115		
	Sex (male)	-0.007	0.005	0.0027	252.11	1.82	0.179		
	Vgll3 (LL)	-0.005	0.005	0.0014	248.00	0.98	0.324		
	Six6 (LL)	0.002	0.005	0.0002	248.04	0.14	0.706	Batch	0.0005 (0.0002, 0.0012)
	Log10 BM	**0.102**	**0.003**	**1.4983**	**258.76**	**1026.08**	**<0.0001**	Family	0.0013 (0.0003, 0.0083)
	Treatment x log10 BM	**0.014**	**0.006**	**0.0074**	**258.83**	**5.09**	**0.025**	Residual	0.0014
MMR	Intercept	0.335	0.013						
	Treatment (LF)	0.007	0.006	0.002	250.04	1.36	0.245		
	Sex (male)	-0.003	0.005	0.001	253.98	0.42	0.517		
	***Vgll3* (LL)**	**-0.018**	**0.005**	**0.020**	**251.86**	**11.62**	**0.0008**	Chamber	0.0002 (0, 0.0005)
	Six6 (LL)	-0.007	0.005	0.003	253.93	1.69	0.195	Initial	0.0001 (0, 0.0013)
	Log10 BM	**0.088**	**0.003**	**1.428**	**257.64**	**833.45**	**<0.0001**	Family	0.0004 (0.0001, 0.0028)
	Vgll3 (LL) x *Six6* (LL)	**-0.022**	**0.011**	**0.008**	**253.84**	**4.48**	**0.035**	Residual	0.0017
AS	Intercept	0.211	0.019						
	Treatment (LF)	0.014	0.008	0.009	241.47	3.22	0.074		
	Sex (male)	-0.005	0.007	0.002	242.58	0.58	0.447		
	***Vgll3* (LL)**	**-0.018**	**0.007**	**0.020**	**240.66**	**7.21**	**0.008**	Chamber	0.0002 (0.00003, 0.00079)
	Six6 (LL)	-0.007	0.007	0.003	243.74	1.05	0.308	Initial	0.0003 (0, 0.00260)
	Log10 BM	**0.091**	**0.004**	**1.273**	**245.56**	**459.20**	**<0.0001**	Family	0.0010 (0.00028, 0.00656)
	Treatment x log10 BM	-0.017	0.009	0.010	246.33	3.75	0.054	Residual	0.0028

BM = Body mass, LF = Low food

Fig 2. A) Predicted means for standard metabolic rate (SMR), maximum metabolic rate (MMR),

and aerobic scope (AS) in *vgll3* and *six6* early- and late-maturation genotypes with 90% confidence

intervals. The means are for average body mass, treatment, and sex effects, and back transformed to

linear scale. P-values show significant pairwise differences between genotypes for MMR on top of

the points, and for the *vgll3* main effects in MMR and AS between the points. N = 60–71 in each

genotype combination (same individuals used for all traits). B) Residual aerobic scope from a linear

mixed model including $\log_{10}$ aerobic scope as response, $\log_{10}$ body mass as predictor and family as

random term, showing individuals (by points) in each genotype combination.

Correlations among metabolic phenotypes

There was a positive correlation between mass- and family- corrected SMR and MMR in the high

food, but not the low food treatment (Fig. S8), and a very strong correlation between MMR and AS

in both treatments (Table 2).

Table 2. Pearson's correlation coefficients between metabolic phenotypes in high food (above diagonal) and low food (below diagonal) treatments. P-values given in parentheses.

	rSMR	rMMR	rAbsAS
rSMR		**0.28 (0.004)**	0.13 (0.09)
rMMR	0.11 (0.16)		**0.99 (<0.001)**
rAbsAS	**-0.17 (0.04)**	**0.96 (<0.001)**	

Discussion

The timing of maturation, just as many life-history traits, depends on reaching a certain body size threshold, i.e., the acquisition of sufficient energy that can be allocated for maturation processes[38,39,72,73]. In line with our hypothesis, we found that the *vgll3* early maturation genotype increased the aerobic scope (AS) of juvenile Atlantic salmon compared to the late maturation genotype. This effect was driven by a change in maximum metabolic rate (MMR), not standard metabolic rate (SMR) (nearly all variation in AS was explained by MMR in our study, Table 2). A previous study showed that higher condition factor, mediated by the *vgll3* early maturation genotype, positively affected the initiation of male maturation[45]. The results presented here suggest that superior resource acquisition or assimilation via higher AS, driven by a higher MMR, is a potential mechanism by which an increased condition factor in individuals with the *vgll3* early maturation genotype could be achieved compared conspecifics with the late maturation genotype. In addition to differences in mean performance between *vgll3* genotypes, the two loci in our study exhibited physiological epistasis[74], as the cooccurrence of the late maturing genotypes in both loci was associated with lower MMR than their additive effects. The epistasis may help to maintain genetic variation under rapid adaptive responses[75,76].

The functional pathways that may explain the epistasis and the main effect of *vgll3* are not well known. However, both *six6* and *vgll3* are expressed during development and have been implicated in the control of cell fate commitment and the hypothalamus-pituitary-gonad axis in salmon[77,78]. Hence, the epistatic interaction between two genomic regions may stem from, e.g., developmental canalisation[79]. Addressing the causal physiological and morphological mechanisms of the link between the genomic regions and aerobic performance can shed light into the mechanisms of life history evolution in salmon; the *vgll3* genomic region is the major genetic axis explaining variation in age-at-maturity in salmon[43], and variation in the locus is spatially divergent among populations and under rapid adaptative evolution[43,47,80,81].

Because of context-dependent covariation between metabolism and growth rate, whereby high SMR improves growth under high, but not low, resource availability[27] and because performance can be reduced under food limitation (e.g.,[8]), we tested if food availability modified the genetic covariation between metabolic phenotypes and age-at-maturity. Against our predictions, there was no change in SMR or MMR due to feed restriction, nor did we find genotype-by-environment interactions, despite a strong decrease in growth rate in low food treatment. The results indicate that the different genotypes exhibit no plastic responses to food availability. However, a stronger feed deprivation,

e.g., similar to those occurring in the winter[82], could have induced a more pronounced effect on

SMR[83,84]. Our low food treatment included approximately 3 days of fasting in between feeding to

satiation, similar to a "feast and famine" feeding strategy[85]. A lack of metabolic response to reduced

food availability may be beneficial if it allows the individual to maximize acquisition via food

assimilation during "feasting".

Unlike MMR (and consequently AS), SMR did not exhibit *vgll3*-linked covariation with age-at-

maturity. A decoupling of SMR and MMR in relation to life-history variation was also found by

Archer et al.[86] in resident and migratory brown trout (*Salmo trutta*), and high MMR, but not SMR,

was positively selected for in Atlantic salmon under high food competition[87]. Further, a lack of

differences in SMR across the *vgll3* genotypes was found in our parallel study, in which the fish

were smaller (mean ~1g) compared to this study (mean ~4g)[88]. The lack of association to the age-at-

maturity loci is unexpected, as SMR, or basal metabolic rate in endotherms, has been proposed to

explain life-history variation along the fast-slow axis[5,7], but see[89]. Our results suggest that the

genetic control of maturation by the *vgll3* genomic region via MMR mostly involves physiological

pathways that do not alter SMR simultaneously. Such pathways may be related to oxygen demand

by tissues or its supply during stress and/or exhaustive exercise. For example, structural and

functional variation in the heart (i.e., cardiac output) or muscle, and mechanisms that modulate

oxygen carrying capacity of the cardiovascular system might invoke changes in MMR without

altering SMR[90-92]. However, our study does not rule out the possibility of the metabolic phenotypes

affecting variation in age-at-maturity also phenotypically or via small-effect loci[44,93].

The differences in performance between life-history genotypes may have arisen due to correlated

selection mediated by resource acquisition: higher aerobic scope, which was associated with early

maturation, could enable higher feeding capacity[50] and improve foraging efficiency, for example,

via a shorter searching time of prey[11,82]. Further, salmon in the wild are increasingly experiencing

higher than optimal temperatures due to climate change[94], and MMR is typically less plastic than

SMR in response to environmental temperature[15]. Thus, our results indicate a potentially important

advantage for individuals carrying the early maturation genotype under global warming, which

could be mediated by a higher appetite[95]. This advantage may also extend to higher survival during

spawning migration if the genotype effect on AS persists across life-stages[13,96,97]. It can also be

relevant to survival of salmon after spawning, and thereby repeated spawning (iteroparity), which is

co-inherited with the same *vgll3* genotype as early maturation[98]. However, the lack of sex

differences in metabolic phenotypes in our study both across and within age-at-maturity genotypes,

suggests that sex-dependent life-history variation in salmon[33] is not reflected in metabolic rates

during the juvenile stage (see also [88]).

Although a causal relationship between metabolism at the juvenile stage and age at maturity would

not be surprising, pleiotropy or linkage provide alternatives for the basis of the observed

association. For example, the *vgll3* gene encodes a transcription cofactor associated with cell fate

commitment and is expressed in many tissues[77,99], thus it is likely pleiotropic with multiple

independent functions. Similarly, several polymorphic loci with putative functional variation are co-

localised (i.e., linked) in the genomic region[43]. Finally, stage-dependent genetic correlations in

metabolism may obscure the time point that the trait influences the life-history variation. For

instance, higher aerobic scope can induce maturation via facilitating the size attained both in the

freshwater and at sea (e.g.,[19,39]), but whether aerobic scope genetically covaries across life stages is

yet to be explored.

The presence of genetic covariation between aerobic scope at the juvenile stage and age-at-maturity

at the *vgll3* genomic region suggests potential for multi-trait evolution across life-stages, whereby

selection acting on either trait would alter the phenotypic variation of the other[4,100]. For example, if

natural selection towards later age-at-maturity induces the frequency of late maturing allele, this

would constrain the aerobic scope of juveniles in the population, even if that may be a suboptimal

phenotype. On the other hand, genetic covariation may help to maintain optimal trait variation in

age-at-maturity, i.e., by limiting potentially maladaptive environmentally induced (i.e., plastic)

variation in age-at-maturity (e.g.,[101,102]). For example, river geophysical properties are important

determinants of the optimal age structure at maturity, whereby populations in smaller tributaries

have a younger, and populations in large, fast-flowing rivers have an older age structure[33].

Forecasting age-at-maturity from aerobic performance at earlier stages (e.g., via improved

growth[22,51]) would result in maladaptive age structure if the covariation was explained entirely by

environmental effects. However, our study was aimed at measuring the statistical association, and

does not provide estimates using a classical quantitative genetic framework (i.e., we did not

quantify environmental sources of variation, or variation due to technical or other genetic effects) or

measure evolutionary change. Therefore, partitioning biologically meaningful covariation and

quantifying the correlated response to selection were beyond the scope of this study[103].

Understanding the physiological basis of life-history variation in different life-stages and

environmental conditions can provide insights into the factors driving life-history evolution, and

hence, better predictions of the responses of populations to environmental changes. Wild salmon

populations have declined in recent decades, with a concomitant decrease in the frequency of late

maturing individuals[47,94]. Our study used an eco-physiological approach to identify a potentially

adaptive phenotype relating genetic variation and age-at-maturity in salmon and suggests that

evolution towards an earlier age-at-maturity can cause correlated selection towards increased MMR

and aerobic scope. In conclusion, the integration of age-at-maturity and aerobic performance in the

early life-stages via simple genetic mechanisms, as shown in this study, could contribute to the

diversification of ecotypes within species.

Data availability

385   The data are available in https://doi.org/10.5281/zenodo.5667078. R codes for the analyses are

386   available in an archived repository (v.1.0.1, https://doi.org/10.5281/zenodo.5667133).

Conflicts of interest

389   The authors declare no conflicts of interest.

Supplemental Material

392   Pdf-file with Material and Methods, Figures, and Tables.

References

1   Ricklefs, R. E. & Wikelski, M. The physiology/life-history nexus. *Trends in Ecology & Evolution*
17, 462-468, doi:10.1016/s0169-5347(02)02578-8 (2002).

2   Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory
of ecology. *Ecology* **85**, 1771-1789, doi:10.1890/03-9000 (2004).

3   Dammhahn, M., Dingemanse, N. J., Niemela, P. T. & Reale, D. Pace-of-life syndromes: a
framework for the adaptive integration of behaviour, physiology and life history. *Behavioral*
Ecology and Sociobiology **72**, doi:10.1007/s00265-018-2473-y (2018).

4   Roff, D. A. Evolutionary quantitative genetics. (Chapman and Hall, New York., 1997).

5   Boratynski, Z., Koskela, E., Mappes, T. & Schroderus, E. Quantitative genetics and fitness effects of
basal metabolism. *Evolutionary Ecology* **27**, 301-314, doi:10.1007/s10682-012-9590-2 (2013).

6   Krams, I. A. *et al.* Metabolic rate associates with, but does not generate covariation between,
behaviours in western stutter-trilling crickets, Gryllus integer. *Proceedings of the Royal Society B-*
Biological Sciences **284**, doi:10.1098/rspb.2016.2481 (2017).

7   Auer, S. K., Dick, C. A., Metcalfe, N. B. & Reznick, D. N. Metabolic rate evolves rapidly and in
parallel with the pace of life history. *Nature Communications* **9**, doi:10.1038/s41467-017-02514-z
(2018).

8   Lailvaux, S. P. & Husak, J. F. The life-history of whole-organism performance. *Quarterly Review of*
412   *Biology* **89**, 285-318, doi:10.1086/678567 (2014).

413   9   Fry, F. E. J. Effects of the environment on animal activity. Ontario Fisheries Research Laboratory
414   Publication, Biol. Ser. **55**, 1-62 (1947).

415   10   Chabot, D., Steffensen, J. F. & Farrell, A. P. The determination of standard metabolic rate in fishes.
416   *Journal of Fish Biology* **88**, 81-121, doi:10.1111/jfb.12845 (2016).

417   11   Metcalfe, N. B., Van Leeuwen, T. E. & Killen, S. S. Does individual variation in metabolic
418   phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**, 298-321,
419   doi:10.1111/jfb.12699 (2016).

420   12   Farrell, A. P. Environment, antecedents and climate change: lessons from the study of temperature
421   physiology and river migration of salmonids. *Journal of Experimental Biology* **212**, 3771-3780,
422   doi:10.1242/jeb.023671 (2009).

423   13   Clark, T. D., Jeffries, K. M., Hinch, S. G. & Farrell, A. P. Exceptional aerobic scope and
424   cardiovascular performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a
425   warming climate. *Journal of Experimental Biology* **214**, 3074-3081, doi:10.1242/jeb.060517 (2011).

426   14   Eliason, E. J., Wilson, S. M., Farrell, A. P., Cooke, S. J. & Hinch, S. G. Low cardiac and aerobic
427   scope in a coastal population of sockeye salmon *Oncorhynchus nerka* with a short upriver migration.
428   *Journal of Fish Biology* **82**, 2104-2112, doi:10.1111/jfb.12120 (2013).

429   15   Sandblom, E. *et al.* Physiological constraints to climate warming in fish follow principles of plastic
430   floors and concrete ceilings. *Nature Communications* **7**, doi:10.1038/ncomms11447 (2016).

431   16   Killen, S. S. *et al.* Ecological Influences and Morphological Correlates of Resting and Maximal
432   Metabolic Rates across Teleost Fish Species. *American Naturalist* **187**, 592-606,
433   doi:10.1086/685893 (2016).

434   17   Priede, I. G. Metabolic scope in fishes. (In: Tytler P., Calow P. (eds) Fish Energetics. Springer,
435   Dordrecht. https://doi.org/10.1007/978-94-011-7918-8_2, 1985).

436   18   Thorpe, J. E. Maturation responses of salmonids to changing developmental opportunities. *Marine*
437   *Ecology Progress Series* **335**, 285-288, doi:10.3354/meps335285 (2007).

438   19   Salminen, M. Relationships between smolt size, postsmolt growth and sea age at maturity in Atlantic
439   salmon ranched in the Baltic Sea. *Journal of Applied Ichthyology-Zeitschrift Fur Angewandte*
440   *Ichthyologie* **13**, 121-130, doi:10.1111/j.1439-0426.1997.tb00111.x (1997).

441   20   Theriault, V., Garant, D., Bernatchez, L. & Dodson, J. J. Heritability of life-history tactics and
442   genetic correlation with body size in a natural population of brook charr (Salvelinus fontinalis).
443   *Journal of Evolutionary Biology* **20**, 2266-2277, doi:10.1111/j.1420-9101.2007.01417.x (2007).

444   21   Roff, D. A. The evolution of dimorphic traits - predicting the genetic correlation between
445   environments. *Genetics* **136**, 395-401 (1994).

446   22   Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J. & Metcalfe, N. B. The optimal combination of
447   standard metabolic rate and aerobic scope for somatic growth depends on food availability.
448   *Functional Ecology* **29**, 479-486, doi:10.1111/1365-2435.12396 (2015).

23   Niemela, P. T. & Dingemanse, N. J. Meta-analysis reveals weak associations between intrinsic state
and personality. *Proceedings of the Royal Society B-Biological Sciences* **285**,
doi:10.1098/rspb.2017.2823 (2018).

24   Mathot, K. J., Dingemanse, N. J. & Nakagawa, S. The covariance between metabolic rate and
behaviour varies across behaviours and thermal types: meta-analytic insights. *Biological Reviews* **94**,
1056-1074, doi:10.1111/brv.12491 (2019).

25   Rosenfeld, J., Richards, J., Allen, D., Van Leeuwen, T. & Monnet, G. Adaptive trade-offs in fish
energetics and physiology: insights from adaptive differentiation among juvenile salmonids.
Canadian Journal of Fisheries and Aquatic Sciences **77**, 1243-1255, doi:10.1139/cjfas-2019-0350
(2020).

26   Zeng, L. Q. *et al.* Standard metabolic rate predicts growth trajectory of juvenile Chinese crucian carp
(*Carassius auratus*) under changing food availability. *Biology Open* **6**, 1305-1309,
doi:10.1242/bio.025452 (2017).

27   Auer, S. K., Solowey, J. R., Rajesh, S. & Rezende, E. L. Energetic mechanisms for coping with
changes in resource availability. *Biology Letters* **16**, doi:10.1098/rsbl.2020.0580 (2020).

28   Killen, S. S., Marras, S. & McKenzie, D. J. Fuel, fasting, fear: routine metabolic rate and food
deprivation exert synergistic effects on risk-taking in individual juvenile European sea bass. *Journal*
of Animal Ecology **80**, 1024-1033, doi:10.1111/j.1365-2656.2011.01844.x (2011).

29   Auer, S. K. *et al.* Metabolic Rate Interacts with Resource Availability to Determine Individual
Variation in Microhabitat Use in the Wild. *American Naturalist* **196**, 132-144, doi:10.1086/709479
(2020).

30   Hughes, K. A., Rodd, F. H. & Reznick, D. N. Genetic and environmental effects on secondary sex
traits in guppies (Poecilia reticulata). *Journal of Evolutionary Biology* **18**, 35-45, doi:10.1111/j.1420-
9101.2004.00806.x (2005).

31   Gutteling, E. W. *et al.* Environmental influence on the genetic correlations between life-history traits
in Caenorhabditis elegans. *Heredity* **98**, 206-213, doi:10.1038/sj.hdy.6800929 (2007).

32   Gillespie, J. H. & Turelli, M. Genotype-environment interactions and the maintenance of polygenic
variation. *Genetics* **121**, 129-138 (1989).

33   Fleming, I. A. & Einum, S. Reproductive Ecology: A Tale of Two Sexes. in *Atlantic Salmon*
Ecology 33-65 (2011).

34   Mobley, K. B. *et al.* Maturation in Atlantic salmon (Salmo salar, Salmonidae): a synthesis of
ecological, genetic, and molecular processes. *Reviews in Fish Biology and Fisheries*,
doi:10.1007/s11160-021-09656-w (2021).

35   Hendry, A. P., Berg, O. K. & Quinn, T. P. Condition dependence and adaptation-by-time: breeding
date, life history, and energy allocation within a population of salmon. *Oikos* **85**, 499-514,
doi:10.2307/3546699 (1999).

36   Stearns, S. C. Life history evolution: successes, limitations, and prospects. *Naturwissenschaften* **87**,
476-486, doi:10.1007/s001140050763 (2000).

37 Friedland, K. D. & Haas, R. E. Marine post-smolt growth and age at maturity of Atlantic salmon.
Journal of Fish Biology **48**, 1-15, doi:10.1111/j.1095-8649.1996.tb01414.x (1996).

38 Skilbrei, O. T. Relationships between smolt length and growth and maturation in the sea of
individually tagged Atlantic salmon (*Salmo salar*). *Aquaculture* **83**, 95-108, doi:10.1016/0044-
8486(89)90064-1 (1989).

39 Hutchings, J. A. & Jones, M. E. B. Life history variation and growth rate thresholds for maturity in
Atlantic salmon, *Salmo salar*. *Canadian Journal of Fisheries and Aquatic Sciences* **55**, 22-47,
doi:10.1139/cjfas-55-S1-22 (1998).

40 Tréhin, C. *et al.* Growth during the first summer at sea modulates sex-specific maturation schedule
in Atlantic salmon. *Canadian Journal of Fisheries and Aquatic Sciences* **78**, 659-669,
doi:10.1139/cjfas-2020-0236 (2021).

41 Moore, M. P. & Martin, R. A. On the evolution of carry-over effects. *Journal of Animal Ecology* **88**,
1832-1844, doi:10.1111/1365-2656.13081 (2019).

42 Ayllon, F. *et al.* The vgll3 Locus Controls Age at Maturity in Wild and Domesticated Atlantic
Salmon (*Salmo salar* L.) Males. *Plos Genetics* **11**, doi:10.1371/journal.pgen.1005628 (2015).

43 Barson, N. J. *et al.* Sex-dependent dominance at a single locus maintains variation in age at maturity
in salmon. *Nature* **528**, 405-408, doi:10.1038/nature16062 (2015).

44 Sinclair-Waters, M. *et al.* Beyond large-effect loci: large-scale GWAS reveals a mixed large-effect
and polygenic architecture for age at maturity of Atlantic salmon. *Genetics Selection Evolution* **52**,
doi:10.1186/s12711-020-0529-8 (2020).

45 Debes, P. V. *et al.* Polygenic and major-locus contributions to sexual maturation timing in Atlantic
salmon. *Molecular Ecology*, doi:10.1111/mec.16062 (2021).

46 Chaput, G. Overview of the status of Atlantic salmon (Salmo salar) in the North Atlantic and trends
in marine mortality. *Ices Journal of Marine Science* **69**, 1538-1548, doi:10.1093/icesjms/fss013
(2012).

47 Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C. R. Rapid sex-specific evolution of
age at maturity is shaped by genetic architecture in Atlantic salmon. *Nature Ecology & Evolution* **2**,
1800-1807, doi:10.1038/s41559-018-0681-5 (2018).

48 Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C. Evolution in salmon life-history
induced by direct and indirect effects of fishing. *bioRxiv*, doi:10.1101/2021.01.08.425869 (2021).

49 Aykanat, T. *et al.* Life-history genomic regions explain differences in Atlantic salmon marine diet
specialization. *Journal of Animal Ecology* **89**, 2677-2691, doi:10.1111/1365-2656.13324 (2020).

50 Auer, S. K., Salin, K., Anderson, G. J. & Metcalfe, N. B. Aerobic scope explains individual variation
in feeding capacity. *Biology Letters* **11**, doi:10.1098/rsbl.2015.0793 (2015).

51 Norin, T. & Clark, T. D. Fish face a trade-off between 'eating big' for growth efficiency and 'eating
small' to retain aerobic capacity. *Biology Letters* **13**, doi:10.1098/rsbl.2017.0298 (2017).

52   Elliott, J. M. & Hurley, M. A. A functional model for maximum growth of Atlantic Salmon parr,
Salmo salar, from two populations in northwest England. *Functional Ecology* **11**, 592-603,
doi:10.1046/j.1365-2435.1997.00130.x (1997).

53   Ward, A. J. W., Webster, M. M. & Hart, P. J. B. Intraspecific food competition in fishes. *Fish and*
Fisheries **7**, 231-261, doi:10.1111/j.1467-2979.2006.00224.x (2006).

54   Rosenfeld, J., Van Leeuwen, T., Richards, J. & Allen, D. Relationship between growth and standard
metabolic rate: measurement artefacts and implications for habitat use and life-history adaptation in
salmonids. *Journal of Animal Ecology* **84**, 4-20, doi:10.1111/1365-2656.12260 (2015).

55   Forstner, H. An automated multiple-chamber intermittent-flow respirometer. (Springer Berlin
Heidelberg), 111-126 (1983).

56   Svendsen, M. B. S., Bushnell, P. G. & Steffensen, J. F. Design and setup of intermittent-flow
respirometry system for aquatic organisms. *Journal of Fish Biology* **88**, 26-50,
doi:10.1111/jfb.12797 (2016).

57   Killen, S. S. *et al.* Guidelines for reporting methods to estimate metabolic rates by aquatic
intermittent-flow respirometry. *Journal of Experimental Biology* **224**, doi:10.1242/jeb.242522
(2021).

58   Raby, G. D., Doherty, C. L. J., Mokdad, A., Pitcher, T. E. & Fisk, A. T. Post-exercise respirometry
underestimates maximum metabolic rate in juvenile salmon. *Conservation Physiology* **8**,
doi:10.1093/conphys/coaa063 (2020).

59   Brett, J. R. The respiratory metabolism and swimming performance of young Sockeye salmon.
Journal of the Fisheries Research Board of Canada **21**, 1183-1226 (1964).

60   Norin, T. & Clark, T. D. Measurement and relevance of maximum metabolic rate in fishes. *Journal*
of Fish Biology **88**, 122-151, doi:10.1111/jfb.12796 (2016).

61   Morozov, S., McCairns, R. J. S. & Merilä, J. FishResp: R package and GUI application for analysis
of aquatic respirometry data. *Conservation Physiology* **7**, doi:10.1093/conphys/coz003 (2019).

62   Harianto, J., Carey, N. & Byrne, M. respR-An R package for the manipulation and analysis of
respirometry data. *Methods in Ecology and Evolution* **10**, 912-920, doi:10.1111/2041-210x.13162
(2019).

63   Little, A. G. *et al.* Maxed Out: Optimizing Accuracy, Precision, and Power for Field Measures of
Maximum Metabolic Rate in Fishes. *Physiological and Biochemical Zoology* **93**, 243-254,
doi:10.1086/708673 (2020).

64   Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods in*
Ecology and Evolution **1**, 103-113, doi:10.1111/j.2041-210X.2010.00012.x (2010).

65   Lenth, R. V.   (R package version 1.5.3, 2020).

66   Bates, D., Machler, M., Bolker, B. M. & Walker, S. C. Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67**, 1-48, doi:10.18637/jss.v067.i01 (2015).

67   R Core Team.   (R Foundation for Statistical Computing, Vienna, Austria, 2019).

68 Stoffel, M. A., Nakagawa, S. & Schielzeth, H. partR2: partitioning R-2 in generalized linear mixed
models. *Peerj* **9**, doi:10.7717/peerj.11414 (2021).

69 Lüdecke, D. *et al.* Vol. 3   (Journal of Open Source Software, 2018).

70 Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Ggplot2: Elegant Graphics for Data*
Analysis, 1-212, doi:10.1007/978-0-387-98141-3 (2009).

71 Long, J. A. interactions: Comprehensive, User-Friendly Toolkit for Probing Interactions. (2019).

72 Weimerskirch, H. Reproductive effort in long-lived birds – Age-specific patterns of condition,
reproduction and survival in the Wandering albatross. *Oikos* **64**, 464-473, doi:10.2307/3545162
(1992).

73 Cheung, C. C. *et al.* Leptin is a metabolic gate for the onset of puberty in the female rat.
Endocrinology **138**, 855-858, doi:10.1210/en.138.2.855 (1997).

74 Cheverud, J. M. & Routman, E. J. EPISTASIS AND ITS CONTRIBUTION TO GENETIC
VARIANCE-COMPONENTS. *Genetics* **139**, 1455-1461 (1995).

75 Cheverud, J. M. & Routman, E. J. Epistasis as a source of increased additive genetic variance at
population bottlenecks. *Evolution* **50**, 1042-1051, doi:10.2307/2410645 (1996).

76 Merilä, J. & Sheldon, B. C. Genetic architecture of fitness and nonfitness traits: empirical patterns
and development of ideas. *Heredity* **83**, 103-109, doi:10.1046/j.1365-2540.1999.00585.x (1999).

77 Kurko, J. *et al.* Transcription Profiles of Age-at-Maturity-Associated Genes Suggest Cell Fate
Commitment Regulation as a Key Factor in the Atlantic Salmon Maturation Process. *G3-Genes*
Genomes Genetics **10**, 235-246, doi:10.1534/g3.119.400882 (2020).

78 Moustakas-Verho, J. E. *et al.* Developmental expression patterns of six6: A gene linked with
spawning ecotypes in Atlantic salmon. *Gene Expression Patterns* **38**, doi:10.1016/j.gep.2020.119149
(2020).

79 Debat, V. & David, P. Mapping phenotypes: canalization, plasticity and developmental stability.
Trends in Ecology & Evolution **16**, 555-561, doi:10.1016/s0169-5347(01)02266-2 (2001).

80 Pritchard, V. L. *et al.* Genomic signatures of fine-scale local selection in Atlantic salmon suggest
involvement of sexual maturation, energy homeostasis and immune defence-related genes.
Molecular Ecology **27**, 2560-2575, doi:10.1111/mec.14705 (2018).

81 Zueva, K. J., Lumme, J., Veselov, A. E., Primmer, C. R. & Pritchard, V. L. Population genomics
reveals repeated signals of adaptive divergence in the Atlantic salmon of north-eastern Europe.
Journal of Evolutionary Biology **34**, 866-878, doi:10.1111/jeb.13732 (2021).

82 Johansen, M., Erkinaro, J. & Amundsen, P. A. in *Atlantic Salmon Ecology*   Ch. 4, (Wiley-
Blackwell, 2011).

83 O'Connor, K. I., Taylor, A. C. & Metcalfe, N. B. The stability of standard metabolic rate during a
period of food deprivation in juvenile Atlantic salmon. *Journal of Fish Biology* **57**, 41-51,
doi:10.1006/jfbi.2000.1280 (2000).

84 Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J. & Metcalfe, N. B. Flexibility in metabolic rate
confers a growth advantage under changing food availability. *Journal of Animal Ecology* **84**, 1405-
1411, doi:10.1111/1365-2656.12384 (2015).

85 Armstrong, J. B. & Schindler, D. E. Excess digestive capacity in predators reflects a life of feast and
famine. *Nature* **476**, 84-87, doi:10.1038/nature10240 (2011).

86 Archer, L. C. *et al.* Metabolic traits in brown trout (Salmo trutta) vary in response to food restriction
and intrinsic factors. *Conservation Physiology* **8**, doi:10.1093/conphys/coaa096 (2020).

87 Auer, S. K. *et al.* Nutrients from salmon parents alter selection pressures on their offspring. *Ecology*
Letters **21**, 287-295, doi:10.1111/ele.12894 (2018).

88 Åsheim, E. R., Prokkola, J. M., Morozov, S., Aykanat, T. & Primmer, C. R. Standard metabolic rate
does not associate with age-at-maturity genotype in juvenile Atlantic salmon. *Ecology and Evolution*
(in press).

89 Polverino, G., Santostefano, F., Diaz-Gil, C. & Mehner, T. Ecological conditions drive pace-of-life
syndromes by shaping relationships between life history, physiology and behaviour in two
populations of Eastern mosquitofish. *Scientific Reports* **8**, doi:10.1038/s41598-018-33047-0 (2018).

90 Harter, T. S., Zanuzzo, F. S., Supuran, C. T., Gamperl, A. K. & Brauner, C. J. Functional support for
a novel mechanism that enhances tissue oxygen extraction in a teleost fish. *Proceedings of the Royal*
Society B-Biological Sciences **286**, doi:10.1098/rspb.2019.0339 (2019).

91 Nikinmaa, M., Berenbrink, M. & Brauner, C. J. Regulation of erythrocyte function: Multiple
evolutionary solutions for respiratory gas transport and its regulation in fish. *Acta Physiologica* **227**,
doi:10.1111/apha.13299 (2019).

92 McArley, T. J., Morgenroth, D., Zena, L. A., Ekstrom, A. T. & Sandblom, E. Normoxic limitation of
maximal oxygen consumption rate, aerobic scope and cardiac performance in exhaustively exercised
rainbow trout (Oncorhynchus mykiss). *Journal of Experimental Biology* **224**,
doi:10.1242/jeb.242614 (2021).

93 Johnston, S. E. *et al.* Genome-wide SNP analysis reveals a genetic basis for sea-age variation in a
wild population of Atlantic salmon (Salmo salar). *Molecular Ecology* **23**, 3452-3468,
doi:10.1111/mec.12832 (2014).

94 Friedland, K. D. *et al.* The recruitment of Atlantic salmon in Europe. *Ices Journal of Marine Science*
66, 289-304, doi:10.1093/icesjms/fsn210 (2009).

95 Jutfelt, F. *et al.* 'Aerobic scope protection' reduces ectotherm growth under warming. *Functional*
Ecology **35**, 1397-1407, doi:10.1111/1365-2435.13811 (2021).

96 Eliason, E. J. *et al.* Differences in Thermal Tolerance Among Sockeye Salmon Populations. *Science*
332, 109-112, doi:10.1126/science.1199158 (2011).

97 Mottola, G., Kristensen, T. & Anttila, K. Compromised thermal tolerance of cardiovascular capacity
in upstream migrating Arctic char and brown trout-are hot summers threatening migrating
salmonids? *Conservation Physiology* **8**, doi:10.1093/conphys/coaa101 (2020).

98   Aykanat, T. *et al.* Co-inheritance of sea age at maturity and iteroparity in the Atlantic salmon vgll3
genomic region. *Journal of Evolutionary Biology* **32**, 343-355, doi:10.1111/jeb.13418 (2019).

99   Verta, J. P. *et al.* Cis-regulatory differences in isoform expression associate with life history strategy
variation in Atlantic salmon. *Plos Genetics* **16**, doi:10.1371/journal.pgen.1009055 (2020).

100   Lande, R. Quantitative genetic analysis of multivariate evolution, applied to brain - body size
allometry. *Evolution* **33**, 402-416, doi:10.2307/2407630 (1979).

101   De Jong, G. Unpredictable selection in a structured population leads to local genetic differentiation
in evolved reaction norms. *Journal of Evolutionary Biology* **12**, 839-851, doi:10.1046/j.1420-
9101.1999.00118.x (1999).

102   Tufto, J. The evolution of plasticity and nonplastic spatial and temporal adaptations in the presence
of imperfect environmental cues. *American Naturalist* **156**, 121-130, doi:10.1086/303381 (2000).

103   Sgro, C. M. & Hoffmann, A. A. Genetic correlations, tradeoffs and environmental variation.
Heredity **93**, 241-248, doi:10.1038/sj.hdy.6800532 (2004).