Comparison of Single-Trait and Multi-Trait Genome-Wide Association Models and Inclusion of Correlated Traits in the Dissection of the Genetic Architecture of a Complex Trait in a Breeding Program

Lance F. Merrick[1], Adrienne B. Burke[1], Zhiwu Zhang[1], and Arron H. Carter[1]

[1]Department of Crop and Soil Sciences, Washington State University, Pullman, WA 99164, USA

*Correspondence: Dr. Arron H. Carter ahcarter@wsu.edu

Abstract

Traits with an unknown genetic architecture make it difficult to create a useful bi-parental mapping population to characterize the genetic basis of the trait due to a combination of complex and pleiotropic effects. Seedling emergence of wheat (*Triticum aestivum* L.) from deep planting is a vital factor affecting stand establishment and grain yield, has a poorly understood genetic architecture, and is historically correlated with coleoptile length. The creation of bi-parental mapping populations can be overcome by using genome-wide association studies (GWAS). This study aimed to dissect the genetic architecture of seedling emergence while accounting for correlated traits using one multi-trait GWAS model (MT-GWAS) and three single-trait GWAS models (ST-GWAS) with the inclusion of covariates for correlated traits. The ST-GWAS models included one single locus model (MLM), and two multiple loci models (FarmCPU and BLINK). We conducted the GWAS using two populations, the first consisting of 473 varieties from a diverse association mapping panel (DP) phenotyped from 2015-2019, and the other population used as a validation population consisting of 279 breeding lines (BL) phenotyped in 2015 in Lind, WA, with 40,368 markers. We also compared the inclusion of coleoptile length and markers associated with reduced height as covariates in our ST-GWAS models for the DP. ST-GWAS found 107 significant markers across 19 chromosomes, while MT-GWAS found 82 significant markers across 14 chromosomes. MT-GWAS models were able to identify large-effect markers on chromosome 5A. FarmCPU and BLINK models were able to identify many small effect markers, and the inclusion of covariates helped to identify the large effect markers on chromosome 5A. Therefore, by using multi-locus models combined with pleiotropic covariates, breeding programs can uncover the complex nature of traits to help identify candidate genes and the underlying architecture of a trait, such as seedling emergence of deep-sown winter wheat.

Keywords: Covariates; Single-Locus Model; Multi-Locus Model; Reduced Height Alleles; Pleiotropic Effects; Seedling Emergence; Coleoptile Length

Introduction

Complex traits are controlled by many quantitative trait loci (QTLs) and influenced by environmental conditions (Bernardo 2020). Challenges due to complexity, small effect QTLs, and difficulty in phenotyping, can make it difficult to create useful bi-parental mapping populations to characterize the genetic basis of traits using QTLs. These challenges for linkage mapping for complex traits can result in inconsistent estimated QTL effects (Bernardo 2020; Tibbs Cortes et al. 2021). The difficulty in creating bi-parental mapping populations creates a need for plant breeders and geneticists: to increase the knowledge of the inheritance and genetic architecture of the trait and to identify markers using genome-wide association studies (GWAS; Bernardo 2020).

GWAS enables the discovery of QTLs in a collection of diverse populations or diversity panels rather than using a mapping population (Lander and Schork 1994). GWAS can dissect the genetic architecture of a trait by exploiting all historical recombination events in the population and allow the ability to understand the genetic basis by identifying the associations between genetic markers and phenotypes (Lipka et al. 2015). Not only are complex traits influenced by the environment and multiple QTLs, but they also interact with correlated traits that result in a complex genetic architecture. Using covariates and correlated secondary traits can help account for confounding factors that bias marker effects and improve the GWAS model's power (Aschard et al. 2015). Commonly used covariates in GWAS account for population structure or genetic relatedness (Tibbs Cortes et al. 2021).

Statistical models have been developed for GWAS to distinguish real associations from false positives caused by population structure and linkage disequilibrium. Some of the first GWAS models, such as general linear models and mixed linear models (MLM), were single-locus, single-trait GWAS (ST-GWAS) models created to implement the covariates along with kinship matrices (Yu et al. 2006). However, these simple models resulted in false negatives caused by weakened associations in order to control inflation of p-values due to population structure (Liu et al. 2016). The MLM was improved upon by compressed mixed linear models (CMLM), which clusters individuals and uses them as random effects rather than individual genotypic effects (Zhang et al. 2010). CMLM was further improved by using pseudo quantitative trait nucleotides (QTNs) to derive kinship instead of all of the genetic markers. The settlement of MLM under progressively exclusive relationship (SUPER) model sorts markers by association and are then divided into bins with the most significant marker designated as pseudo QTNs and then used to derive a reduced kinship matrix (Wang et al. 2014). All of these methods improved computational efficiency over MLM while increasing statistical power. However, the single-locus models tested each marker one at a time, reducing power and increasing computational speed compared to multi-locus models (Liu et al. 2016).

Newer multi-locus models were then developed to test multiple markers simultaneously (Liu et al. 2016). These multi-locus GWAS models, such as the fixed and random model

circulating probability unification (FarmCPU) and Bayesian information and linkage-
disequilibrium iteratively nested keyway (BLINK) allow the evaluation of big datasets while also
reducing false positives and negatives (Huang et al. 2019). FarmCPU was developed in order to
control false positives and confounding effects between testing markers and cofactors iteratively.
Associated markers are fitted as cofactors to control false positives for testing the rest of the
markers in a fixed-effect model, and then a random effect model is used to select the associated
markers. BLINK was developed to increase statistical power and efficiency (Huang et al. 2019).
BLINK eliminates the FarmCPU assumption that causal genes are evenly distributed across the
genome, which improves speed, due to the optimization of bin size and number no longer being
required (Huang et al. 2019). However, due to the intricate and pleiotropic nature of quantitative
traits, there is no best model for all situations, and it is recommended to use both single locus and
multi-locus models to dissect the unique genetic architecture of complex traits (Tibbs Cortes et
al. 2021). This is due to differing biological complexities that may be identified by one method
and not another, and depends on the genetic architecture and population structure of the trait
(Tibbs Cortes et al. 2021).
Additionally, multi-trait GWAS (MT-GWAS) can be used to analyze multiple traits
simultaneously. MT-GWAS methods were developed to increase statistical power and identify
pleiotropic loci (Porter and O'Reilly 2017). Pleiotropy occurs when genetic loci affect more than
one trait (Solovieff et al. 2013). Therefore, correlations and pleiotropy are important, and taking
them into account can increase power compared to ST-GWAS (Galesloot et al. 2014). There are
various MT-GWAS methods ranging from direct to indirect based models. Direct multivariate
approaches consist as a multi-trait mixed model (MTMM), MultiPhen, and bayesian models
implemented in MV-SNPTEST and MV-BIMBAM which model the effects of genetic loci
directly from the phenotypes (Marchini et al. 2007; O'Reilly et al. 2012; Korte et al. 2012;
Stephens 2013; Galesloot et al. 2014). Indirect approaches are based on a reduction of the trait
dimensions using canonical correlation analysis in MV-PLINK, or based on principal
components using PCHAT (Klei et al. 2008). MT-GWAS methods display an increase in power
even when the traits display a negative correlation, when only one of the traits is associated with
the loci, or when genetic correlations between the traits are weak (Galesloot et al. 2014).
Therefore, MT-GWAS is a valuable approach to understand correlated traits.
Seedling emergence in deep-sown winter wheat exhibits a complex genetic architecture
with a significant environmental effect which allows us to compare GWAS methods in order to
identify marker-trait associations for selection. Seedling emergence is dependent on deep sowing
(sowing depths of 10-15 cm) when precipitation is below 150 to 300 mm annually (Schillinger et
al. 1998; Mohan et al. 2013). In deep-sowing practices, fast-emerging cultivars are the most
desirable to emerge before precipitation events create soil crusting that can dramatically decrease
emergence. Wheat seedlings have decreased emergence when they cannot penetrate the soil

surface due to crusting or due to the inability to germinate under dry soil conditions. Seedling emergence and therefore stand establishment is a vital factor affecting grain yield and can reduce grain yields by 35 to 40% (Schillinger et al. 1998).

Previous studies have shown a significant positive relationship between coleoptile length and seedling emergence (Allan et al. 1962; Sunderman 1964; Chastain et al. 1995; Schillinger et al. 1998; Botwright et al. 2001; Schillinger 2011). The reduced height (Rht) genes *Rht-B1b* and *Rht-D1b* are mutant alleles that cause the semi-dwarfing stature of wheat (Vogel et al. 1956; Allan et al. 1962). Dwarfing genes are responsible for short stature and have a pleiotropic effect that includes gibberellin insensitivity, coleoptile length, yield, protein content, and disease resistance (Gale and Youssefian 1983; Allan 1989). Semi-dwarf wheat cultivars have improved resistance to lodging and grain yield but reduced coleoptile length by one-half to three-fourths of the standard varieties at the time of their development (Allan et al. 1961; Allan 1989; Mohan et al. 2013). The reduced coleoptile length was due to decreased gibberellic acid response, which reduced cell size and elongation (Allan et al. 1959). Historically, when crusting was present, or other unfavorable conditions, the shorter coleoptiles of semi-dwarf cultivars resulted in poor stand establishment and yield potential (Rebetzke et al. 1999).

Recently, after 60 years of breeding, emergence in modern varieties was shown to have a reduced correlation between emergence and coleoptile length (Mohan et al. 2013). Coleoptile length only accounted for 28% of the variability for seedling emergence, and some lines with short coleoptiles had the best emergence rating. The remaining variability is attributed to many factors that affect seedling emergence, leading to a complex system resulting in stand establishment. As stated previously, the two main scenarios that affect seedling emergence and subsequently stand establishment include adequate seed-zone water potential for the first leaf to reach the surface and the occurrence of surface soil crust that prevents penetration (Schillinger 2011). Successful seedling emergence is also dependent on the force exerted by the first leaf. The first leaf protrudes through the coleoptile and emerges around 10 to 12 days after planted. During this time, the first leaf can be prone to buckling before it emerges through the soil surface, which can be affected by coleoptile diameter, speed of emergence, and the emergence force and lifting capacity of the first leave along with the previously reported coleoptile length (Arndt 1965; Schillinger et al. 2017; Lutcher et al. 2019). These studies showed that the genetic basis of seedling emergence is a complex trait not solely controlled by genes for any one factor and results in a poorly understood genetic architecture that is dependent on the environment to display variation (Schillinger et al. 2017; Lutcher et al. 2019).

This study presents research to assess the genetic architecture of a complex trait using GWAS. This study's objectives were to (1) compare ST-GWAS and MT-GWAS models for correlated traits and (2) assess the inclusion of fixed effects to explore the genetic architecture of seedling emergence using different types of populations within and across years to account for

differences in genetic variation between populations and environmental variation. This study

aided in determining the genetic architecture for a complex trait such as seedling emergence in

deep-sown winter wheat.

Material and Methods

<u>Phenotypic Data</u>

Seedling emergence notes were taken to select varieties under low annual precipitation

(~150 mm annual precipitation) at Washington State University in Lind, WA, planted using a

deep-furrow planting system. Emergence notes were taken on two populations within the

breeding program. The diverse association mapping panel (DP) represents a diverse panel of

lines not selected exclusively in deep-furrow trials. In contrast, the second population is

composed of $F_{3:5}$ breeding lines (BL) and represents a population of closely related lines from a

single breeding program composed of pedigrees that have been selected for emergence over

previous generations. The two populations were used to compare GWAS models. The DP was

used as the primary population for genetic dissection with the BL population used as the

validating population. The DP was evaluated in 2015, 2017, 2018, and 2019 in Lind, WA (Table

1). The second population consists of an unreplicated trial of the BL that was evaluated in 2015

(Table 1). In 2016, no data were collected for the DP due to significant soil crusting that was

severe enough to impede all lines' seedling emergence.

Seedling emergence was visually assessed and recorded as a percentage of the total plot

that emerged six weeks after planting for each trial. Table 1 summarizes location, population,

year, and the number of genotyped individuals. The emergence issue for each trial was attributed

to moisture stress. Coleoptile length was also measured for the DP in 2014 and in two replicates

in 2016 under greenhouse conditions. Coleoptile length was recorded to the nearest millimeter

according to Murphy et al. (2008).

Table 1. Study Populations for Seedling Emergence.

Populations screened in unreplicated trials under
moisture stress in Lind, WA from 2015 to 2019.

Location	Trial	Year	Individuals
Lind	DP*	2015	473
Lind	DP	2017	473
Lind	DP	2018	473
Lind	DP	2019	473
Lind	$F_{3:5}$	2015	276

*DP: Quality Association Mapping Diversity Panel

     Phenotypic Adjustments

Adjusted means from the emergence data collected in the unreplicated trials were

adjusted using residuals calculated for the unreplicated lines in individual environments and

across environments using the modified augmented complete block design model (ACBD;

Federer 1956; Goldman 2019). The adjustments were made following the method implemented

in Merrick and Carter (2021), with the full model across environments as follows:

$$Y_{ij} = Block_i + Check_j + Env_k + Block_i \; x \; Env_k + Check_j \; x \; Env_k + \varepsilon_{ijk}$$

where Y_{ij} is the trait of interest; $Block_i$ is the fixed effect of the ith block; $Check_j$ is the fixed

effect of the jth replicated check cultivar; Env_k is the fixed effect of the kth environment; and

ε_{ijk} are the residual errors used to adjust the mean of the lines.

Heritability on a genotype-difference basis for broad-sense heritability was calculated

using the variance components from the models implemented in Merrick and Carter (2021) and

using best linear unbiased predictors for both individual environments and across environments

using the formula:

$$H^2_{Cullis} = 1 - \frac{\bar{v}^{BLUP}_{\Delta..}}{2\sigma^{\wedge 2}_g} \; (2)$$

where σ^2_g, and $\bar{V}^{BLUP}$ is the genotype variance and mean-variance of the BLUPs (Cullis et al. 2006).

BLUPs for coleoptile length were calculated across trials using a mixed linear model

as follows:

$$Y_{klm} = Gen_k + Env_l + Rep_{m(l)} + Gen_k \; x \; Env_l + \varepsilon_{klm}$$

Where Y_{klm} is the trait of interest; $Gen_{k(j)}$ is the genotype k in the jth Check; Env_l is the random

effect of the lth environment; $Rep_{m(l)}$ is the replication m in the lth Env; and ε_{ijkl} are the residual

errors. Trial evaluation and significant differences were evaluated using the coefficient of

variation, and analysis of variance (ANOVA) in individual and across trials. Phenotypic

correlations were conducted between seedling emergence in the DP across years along with

coleoptile length.

Genetic correlations between seedling emergence in the DP and coleoptile length was

calculated using the R package "sommer" with the bivariate model, which is represented as:

$$\begin{bmatrix} Y_A \\ Y_B \end{bmatrix} = \begin{bmatrix} X_A & 0 \\ 0 & X_B \end{bmatrix}\begin{bmatrix} b_A \\ b_B \end{bmatrix} + \begin{bmatrix} Z_A & 0 \\ 0 & Z_B \end{bmatrix}\begin{bmatrix} g_A \\ g_B \end{bmatrix} + \begin{bmatrix} \varepsilon_A \\ \varepsilon_B \end{bmatrix}$$

Where Y_A and Y_B are adjusted means and BLUPs for emergence and coleoptile length,

respectively. X and Z are fixed and random design matrix, subscript A and B represent the primary

trait seedling emergence and secondary trait coleoptile length, respectively. a, g, e are the fixed,

random genetic effects, and residuals, respectively. Variance components were calculated in the

package "sommer", assuming $\begin{bmatrix} g_A \\ g_B \end{bmatrix} \sim N(0, H \otimes G)$, where H is the genetic variance-covariance

matrix and G is the additive relationship matrix calculated using "a.mat" function, and

$\begin{bmatrix} \varepsilon_A \\ \varepsilon_B \end{bmatrix} \sim N(0, I \otimes R)$, where I is the identity matrix, and R is the residual variance-covariance

matrix. Genetic correlations were then calculated using the function "cov2cor" as:

208
$$r_G = \frac{cov(A,B)}{\sqrt{var(A) * var(B)}}$$

Where cov(A,B) is the covariance between seedling emergence and coleoptile length, and

var(A), and var(B) is the genetic variance of seedling emergence and coleoptile length,

individually (Covarrubias-Pazaran 2018; R Core Team 2018).

<u>Genotypic Data</u>

Lines were genotyped using genotyping-by-sequencing (GBS; Elshire et al., 2011) through

the North Carolina State Genomics Sciences Laboratory in Raleigh, North Carolina, using the

restriction enzymes *Msp*I and *Pst*I (Poland et al. 2012). Genomic DNA was isolated from seedlings

at the one to three leaf stage using Qiagen BioSprint 96 Plant kits and the Qiagen BioSprint 96

workstation (Qiagen, Germantown, MD). DNA libraries were prepared following the protocol of

DNA digestion with *Pst*I and *Msp*I restriction enzymes (Poland et al, 2012). Genotyping by

sequencing (GBS; (Elshire et al. 2011) was conducted at North Carolina State University Genomic

Sciences Laboratory with either an Illumina HiSeq 2500 or a NovaSeq 6000. DNA library barcode

adapters, DNA library analysis, and sequence SNP calling were provided by the USDA Eastern

Regional Small Grains Genotyping Laboratory (Raleigh, NC). Sequences were aligned to the

Chinese Spring International Wheat Genome Sequencing Consortium (IWGSC) RefSeq v1.0

(Appels et al. 2018), using the Burrows-Wheeler Aligner (BWA) 0.7.17 (Li and Durbin 2009).

Genetic markers with more than 20% missing data, minor allele frequency of less than 5%, and

those that were monomorphic were removed. Markers were then imputed using Beagle version 5.0

and filtered once more for markers under a 5% minor allele frequency (Browning et al. 2018). A

total of 40,368 single-nucleotide polymorphism (SNP) markers remained.

All winter wheat lines in the DP were genotyped with Kompetitive Allele Specific PCR

(KASP®) assays in the WSU Winter Wheat Breeding Laboratory using allele-specific SNP

markers for semi-dwarf causing mutant alleles *Rht-B1b* and *Rht-D1b* previously reported in

Grogan et al. 2016 and Rasheed et al. 2016. The KASP assays were performed using PACE[TM]

Genotyping Master Mix (3CR Bioscience, Harlow, UK) following the manufacturer's instructions

and endpoint genotyping was conducted from fluorescence using a Lightcycler 480 Instrument II

(Roche, Indianpolis, IN). The Rht markers were coded as lines with *Rht-B1b* (1), *Rht-D1b* (2), *Rht-*

B1b heterozygous (3), *Rht-D1b* heterozygous (4), *Rht-D1b* with a heterozygous *Rht-B1b* (5), *Rht-*

B1b with a heterozygous *Rht-D1b* (6), and both *Rht-B1b* and *Rht-D1b* heterozygous (7).

Linkage disequilibrium (LD) between marker pairs was evaluated using JMP Genomics

v.9.0 (SAS Institute, Inc 2011). Significant marker pairs in the same chromosome were considered

in LD at a p-value <0.05. Population structure within both populations was analyzed using

principal component (PC) analysis biplots and kmeans clustering using the markers in the DP and
BL populations individually using the function "prcomp" and "cluster" in R, respectively (R Core
Team 2018).

Genome-Wide Association Models

To dissect the genetic architecture of a complex trait (seedling emergence), ST-GWAS
models were implemented using the Genome Association and Prediction Integrated Tool (GAPIT;
Liu et al. 2016; Tang et al. 2016; Huang et al. 2019). Both the ST-GWAS and MT-GWAS models
were implemented with three principal components fitted as fixed effects. Three principal
components were used based on BIC values using model selection in GAPIT (Tang et al. 2016).
The GWAS models were conducted on seedling emergence using the adjusted means mentioned
previously and on the BLUPs for coleoptile length. The DP was used as the primary population
for genetic dissection with the BL population used as the validating population. Three ST-GWAS
models were used for comparison. The single-locus ST-GWAS model used was MLM, and the
multi-locus models were BLINK and FarmCPU. Within the DP, we compared each model within
each year combination without covariates, then with the Rht markers as covariates, coleoptile
length BLUPs, and both Rht markers and coleoptile length as covariates. This resulted in 28
datasets for ST-GWAS for seedling emergence in the DP. The GWAS models were then conducted
within the BL without covariates to validate the significant markers. In addition, the ST-GWAS
models were used to dissect coleoptile length within the DP for further validation to determine
whether the significant markers affected coleoptile length.

Additionally, MTMM was implemented using the "sommer" package for MT-GWAS in
order to identify pleiotropic interactions between seedling emergence and coleoptile length within
the DP. The multivariate Newton-Raphson algorithm was used for multiple random effects and
covariance structures using the multivariate model in Covarrubias-Pazaran (2018):

$$y_1 = X_1\beta_1 + Z_1u_1 + \varepsilon_1$$
$$y_2 = X_2\beta_2 + Z_2u_2 + \varepsilon_2$$
$$...$$
$$y_i = X_i\beta_i + Z_iu_i + \varepsilon_i$$

Where y_i is a vector of trait phenotypes, β_i i is a vector of fixed effects, u_i is a vector
of random effects for individuals and ε_i are residuals for trait "i" (i=1,…,t). The random effects (u_1
… u_i and ε) are assumed to be normally distributed with mean zero. X and Z are incidence matrices
for fixed and random effects, respectively. The distribution of the multivariate response and the
phenotypic variance covariance V following the models in Covarrubias-Pazaran (2018) are:

$$Y = X\beta + ZU + \varepsilon_i$$
$$Y \sim MVN(X\beta, V)$$

$$Y = \begin{bmatrix} y_1 \\ y_2 \\ y_i \end{bmatrix}$$

$$X = \begin{bmatrix} X_1 & 0 & 0 \\ \vdots & \ddots & \vdots \\ 0 & 0 & X_t \end{bmatrix}$$

$$V = \begin{bmatrix} Z_1 K \sigma_{g_1}^2 Z_1' + H \sigma_{\varepsilon_1}^2 & \cdots & Z_1 K \sigma_{g_{1,t}} Z_t' + H \sigma_{\varepsilon_{1,t}}^2 \\ \vdots & \ddots & \vdots \\ Z_1 K \sigma_{g_{1,t}} Z_t' + H \sigma_{\varepsilon_{1,t}} & \cdots & Z_t K \sigma_{g_t}^2 Z_t' + H \sigma_{\varepsilon_t}^2 \end{bmatrix}$$

where K is the relationship matrix for the kth random effect (u=1,…,k), and H=I is an identity matrix for the residual term. The terms $\sigma_{g_i}^2$ And $\sigma_{\varepsilon_i}^2$ denote the genetic and residual variance of trait "i", respectively, and $\sigma_{g_{ij}}$ and $\sigma_{\varepsilon_{ij}}$ are the genetic and residual covariance between traits "i" and "j" (i-1,…,t, and j=1,…,t). Further, the GWAS model implemented in the package "sommer" to obtain marker effects uses the inverse of the phenotypic variance matrix (V) and is a generalized linear model of the form:

$$b = (X'V - X)X'V - y$$
$$with\ X = ZM_i$$

Where: b is the marker effect matrix, y is the multivariate response variable, V- is the inverse of the phenotypic variance matrix, Z is the incidence matrix for the random effect to perform the GWAS, and M_i is the ith column of the marker matrix. Further, we implemented three additional F-tests for joint analysis using the scripts proposed in (Korte et al. 2012). The full (FULL) model, which includes the effect of the marker genotype and its interaction, was tested against a null model and identified both loci with common and interaction effects. The interaction model (IE) to identify the interaction effects between the traits tested the full model against a genetic model. Finally, we identified common (COM) genetic effects and tested the genetic model against a null model.

Significant associations based on a Benjamini-Hochberg false discovery rate (FDR; Benjamini and Hochberg 1995). Phenotypic variation and effect explained by each significant marker were calculated by conducting stepwise linear regression between phenotypic and genotypic data and calculating the difference between the effects and variation when a single significant marker was added to the null model in R (R Core Team 2018; Lozada et al. 2019). Significant markers were deemed stable when they were significant in at least two years or in both populations. We then tested the additive effect of pyramiding the stable markers identified across populations and the stable markers identified in multiple years from the DP in each population individually. Manhattan plots were created using the "ggplot2" package, and QQ plots were plotted using the package "CMplots" in R (LiLin 2018; R Core Team 2018).

Results

<u>Phenotypic Data</u>

Heritability for seedling emergence was moderately high in the BL and a few years in the DP,

while the heritability decreased in combined years. The highest heritability in a single trial was

0.88 in the DP in 2018, and the BL had a heritability of 0.77 (Table 2). The trials with minimum

values of zero have a larger standard deviation, which indicates a wider range of seedling

emergence and increased environmental pressure and phenotypic variation for selection

purposes. Heritability for coleoptile length was very high in the DP (0.89) (Table S1).

Table 2. Deep-sowing seedling emergence trial statistics. Cullis heritability and trial statistics for deep-sowing seedling emergence in individual and combined trials for the diversity panel (DP) population and breeding line (BL) population phenotyped from 2015 to 2019 and 2015, respectively.

Population	Trial	Year	Heritability	CV[a]	Max[b] (%)	Mean (%)	Min[c] (%)	SD[d]
DP	DP	2015	0.75	78.91	139	48	-43	38
DP	DP	2017	0.70	24.18	118	87	-26	21
DP	DP	2018	0.88	45.46	118	62	-43	28
DP	DP	2019	0.68	106	135	43	-45	45
DP	DP	2015-2017	0.63	34.53	126	68	-17	23
DP	DP	2015-2018	0.61	28	110	66	2	18
DP	DP	2015-2019	0.64	31.58	101	60	1	19
BL	$F_{3:5}$	2015	0.77	53.55	121	61	-59	33

[a]CV: coefficient of variation

[b]Max: maximum

[c]Min: minimum

[d]SD: standard deviation

Since the variation for seedling emergence depends on the environmental effect, it is

important to examine the trials' variance components. Genetic variances were significant for the

DP in 2015, 2017, and for all of the combined trials (Table S2). The environmental effect was

not significant in any of the combined trials, with the genotype-by-environment interaction (GEI)

effect only significant in the DP over the combined trials of 2015-2018 and 2015-2019 (Table

S2). Phenotypic correlations allow us to compare the results in our GWAS models. The DP trials

are significantly positively correlated to each other with the exception of three scenarios: DP

2015 to DP 2018; DP 2017 to DP 2018; and DP 2017 to DP 2019 (Table S3). The BL $F_{3:5}$ trial in

2015 was significantly correlated to DP 2017. The genetic correlations between the DP seedling

emergence to coleoptile length showed moderate to large correlations in and across years (Table

3). The highest genetic correlation was found in DP 2018 with a correlation of 0.86. However,

this was not the case for the phenotypic correlations. The phenotypic correlations in all years was

near zero between seedling emergence and coleoptile length (Table 3 and Figure 1). The lack of
phenotypic correlation further displays the dependency on the environment for phenotypic
variation and the complexity of seedling emergence.
332

Table 3. Genetic and Phenotypic correlations between seedling emergence and coleoptile length. Genetic correlations between coleoptile length and deep-sowing seedling emergence in individual and combined trials for the diversity panel (DP) population phenotyped from 2015 to 2019.

Year	DP 2015	DP 2017	DP 2018	DP 2019	DP 2015-2017	DP 2015-2018	DP 2015-2019
Genetic Correlations (R^2)	0.41	-0.42	0.86	0.75	0.17	0.43	0.61
Phenotypic Correlations (R^2)	0.00	-0.01	-0.06	0.03	0.00	-0.04	-0.01

Figure 1. Effect of Coleoptile Length on Seedling Emergence. Effect of coleoptile length on
seedling emergence in the diversity panel varieties across the combined years of 2015-2019.

<u>Genotypic Data</u>
The principal component biplot using the SNP markers for the DP displayed four clusters
according to the elbow method with much overlap between clusters 1 and 2 and a slight
overlapping between clusters 3 and 4 (Figure 2A). PC1 explained 12.9% of the variation, and

PC2 explained 6.9% of the variation. However, most of the lines clustered within a single cluster

in the BL population even though the elbow method revealed four clusters using k-means, and

the biplot explained less variation with 9.0% and 5.2% for PC1 and PC2, respectively (Figure

2B). We can visually see the larger genetic variation in the DP than in the BL, and therefore it is

important to be accounted for in our GWAS models.

Figure 2. Principal component analysis. Principal Component (PC) biplot of SNP GBS

markers and k-means clustering from the (A) diversity panel and (B) breeding line population

consisting of the $F_{3:5}$ trial.

The frequency of the Rht alleles in the DP population can be seen in Table S3. The

majority of the lines in the DP had either *Rht-B1b (0.564)* or *Rht-D1b* (0.347), with *Rht-B1b*

conveying a higher mean seedling emergence than *Rht-D1b* (Figure 3). Further, Figure 3

displays outliers for seedling emergence for lines with certain Rht alleles such as *Rht-D1b*. They

were not removed due to the inability to determine the cause of the poor seedling emergence

caused by the complexity between phenotypic variation and genotypic effect.

Figure 3. Effect of Rht alleles on Seedling Emergence. Effect of *Rht* allele frequency on
seedling emergence in the diversity panel varieties across the combined years of 2015-2019. The
Rht markers were coded as lines with *Rht-B1b* (1), *Rht-D1b* (2), *Rht-B1b* heterozygous (3), *Rht-D1b* heterozygous (4), *Rht-D1b* with a heterozygous *Rht-B1b* (5), *Rht-B1b* with a heterozygous
Rht-D1b (6), and both *Rht-B1b* and *Rht-D1b* heterozygous (7).

Single-Trait Genome-Wide Association Studies

To confirm pleiotropic effects of significant markers, ST-GWAS was conducted on coleoptile
length within the DP. Genome-wide association for coleoptile length displayed three unique
markers using MLM, BLINK, and FarmCPU (Table 5). BLINK and MLM both identified a
marker, *S1A_14084576*, on chromosome 1A. Marker *S1A_14084576* conveyed the largest R^2
value of any significant marker with a value of 7%, and an effect of 14.52. The marker with the
next largest R^2 of 0.05 was *S6A_543395015*. It was identified by both BLINK and FarmCPU on
chromosome 6A. It also conveyed the largest effect with 34.44. A third marker was identified
only by FarmCPU, *S2A_765677087*, and is located on chromosome 2A. However,
S2A_765677087, had a rather small effect and R^2 of 3.22 and 0.04, respectively. The three
markers significantly associated with coleoptile length were not identified in any population,
year, nor model for seedling emergence (File S1).

Table 5. Significant markers for coleoptile length. Significant loci controlling deep-sowing seedling emergence in a Pacific Northwest
winter wheat diversity panel using BLINK, FarmCPU and MLM genome-wide association studies (GWAS) models using covariates for
coleoptile length and Rht alleles.

Marker	Position[a]	Alleles[b]	Chr[a]	Model	P-value	MAF[c]	Effect (mm)[d]	R^{2e}

S1A_14084576	14084576	T/A	1A	BLINK	1.82E-11	0.01	14.52	0.07
				MLM	4.78E-08	0.01	14.52	0.07
S2A_765677087	7.66E+08	G/T	2A	FarmCPU	4.22E-09	0.14	3.23	0.04
S6A_543395015	5.43E+08	T/G	6A	BLINK	4.70E-08	0.00	34.44	0.05
				FarmCPU	6.32E-14	0.00	34.44	0.05

[a]Chromosomes and positions of markers were determined according to IWGSC RefSeq v.1.1.
[b]Allele: favorable allele is underlined.
[c]MAF: minor allele frequency.
[d]Effect: Increase of seedling emergence (%) with the inclusion of favorable allele of associated marker.
[e]R^2: Phenotypic variance explained by associated marker.

There were 107 unique markers over all of the combinations of ST-GWAS for seedling
emergence (File S1). Of the 107 significant markers, 96 markers were significant in the DP, and
15 were significant in the BL, with four markers significant in both populations. Seventy-five of
the markers were significant in at least two combinations over both populations. Seventy-one and
five markers were significant in at least two combinations in the DP and BL, respectively.
Significant markers were found on 19 of the 21 chromosomes in the DP, with the majority of the
significant markers in the DP located on chromosome 5A with 34 markers (Figure 3). In the BL,
only nine chromosomes contained significant markers and more consistent with only 15 total
unique markers.

Figure 3. Significant markers for seedling emergence chromosomes. Distribution of
significant markers across both the diversity panel varieties and breeding line populations from
all combinations of models.

All three models had more significant markers in the DP population than in the BL
population. The MLM model identified 36 unique markers over all GWAS, three in the BL, and
23 in the DP. FarmCPU identified 65 unique markers overall, nine in the BL, and 57 in the DP.
BLINK identified 31 unique significant markers overall, eight in the BL, and 23 in the DP. The
MLM model identified significant markers mainly on chromosome 5A with many neighboring

significant markers. In contrast, FarmCPU and BLINK identified the same markers on
chromosome 5A and identified more markers on various other chromosomes (File S1)
In addition to the effect of the model, there was an indication of the effect in including
covariates in BLINK and FarmCPU. This is seen in the number of significant markers identified
within the DP (Figure 4). For FarmCPU, including both coleoptile length and Rht alleles
individually and combined, decreased the number of significant markers compared to FarmCPU
without covariates. For BLINK, the Rht alleles and the combined Rht alleles and coleoptile
length increased the number of significant markers, and coleoptile length alone decreased the
number of significant markers compared to BLINK without covariates. However, this was not
the case for the MLM. For example, including coleoptile length and Rht alleles as covariates
alone had similar number of significant markers (64 and 63) compared to the MLM without
covariates. Including both Rht alleles and coleoptile length did reduce the number of significant
markers to 55. Further, the large number of markers in LD increased the number of significant
markers identified with MLM compared to the other ST-GWAS models.

411
Figure 4. Comparison of significant markers across genome-wide association models and
covariates. Distribution of significant markers across both the (A) diversity panel and (B)
breeding line (BL) populations from all combinations of models. The models compared are
BLINK, FarmCPU, and Mixed-Linear Model (MLM) with and without Rht and coleoptile length
as covariates in the DP and without covariates in the BL.

Further evidence is seen examining QQ plots. For example, in the combined analysis of
2015-2018 and in the BL in 2015, the MLM was consistent regardless of the inclusion of
covariates (Figure S1). However, this was not the case with BLINK or FarmCPU. Within the

DP, the inclusion of Rht or Rht and coleoptile length increased the deviation from the quantile
line more than the models without covariates (Figures S2 and S3). This displays the advantage of
using covariates such as Rht alleles within the DP for an increase in power when the trait in
question is genetically correlated with another trait. In addition, the QQ plots reveal the
difference between the models in the different populations. For both BLINK and FarmCPU, the
BL population had the largest deviations compared to the DP GWAS, whereas the opposite was
seen in the MLM.

In addition, the environmental impact on the genetic dissection of seedling emergence
had a large effect on the identification of significant markers. The GEI for seedling emergence
was shown by the varying number of significant markers in different years for the DP. The
GWAS models were able to identify the most significant markers in the 2015 trial with 132
markers (Figure S4). However, only a few significant markers were identified in the other
individual years with two in 2017 and five in 2019. Further, the combining of years and
accounting for GEI in our phenotypic adjustments allowed the GWAS models to increase the
ability to dissect seedling emergence compared to individual years. All three year combinations
increased the number of significant markers consistently compared to individual years with 109
in year combinations 2015-2017 and 105 in 2015-2018. Additionally, there was a decrease in
significant markers identified with the increase in year combinations with the combination of all
years (2015-2019) identifying only 47 markers (Figure S4).

<u>Multi-Trait Genome-Wide Association Studies</u>

The MT-GWAS models were conducted to identify pleiotropic loci between seedling emergence
and coleoptile length. The MT-GWAS models displayed very similar results for the individual
traits compared to the ST-GWAS models, especially in comparison to the single trait results
from the MT-GWAS model. However, we used the Bonferroni cutoff with an alpha =0.05
instead of FDR due to the large deviations and inflation of p-values seen on the QQ plots for the
joint models (Figure S5). Using FDR resulted in 924 unique markers across the majority of
chromosomes. In comparison, the Bonferroni cutoff resulted in 82 unique significant markers
across 14 chromosomes with the majority of the large effect alleles on chromosome 5A (File S2).
In comparing the results for MT-GWAS FULL, IE, and COM F-tests to the single trait results
from the MT-GWAS, we see no significant markers for coleoptile length significant for any
other model (Figure 5). The significant markers for coleoptile length were the same two
significant markers on chromosomes 1A and 6A identified in the ST-GWAS models.
Additionally, the significant markers for seedling emergence were located on chromosomes 2B
(1) and 5A (15).

Further, there were no significant markers in the IE, indicating no contradictory effect
markers between seedling emergence and coleoptile length. The lack of significant IE markers is
why the FULL model and the COM models are very similar. The COM model was also able to
identify significant markers in every year in the DP compared to the single trait results for
seedling emergence, which did not identify significant markers in the 2017 through 2019

individual years (File S2). Additionally, there were 64 significant markers in the COM model,
with 16 significant markers also found significant for seedling emergence (Figure 5). Therefore,
indicating the potential for pleiotropic markers and the ability to identify them using a joint
analysis model compared to the single trait results implemented in the MT-GWAS models.

464
Figure 5. Comparison of significant markers across Multi-Trait GWAS models for
Seedling Emergence, Coleoptile Length, Full, Interaction Effect, and Common Models
using the diversity panel lines. Venn diagram for the number of unique significant markers
across the multi-trait genome-wide association studies (MT-GWAS) for seedling emergence
(EM), coleoptile length (CL), and the joint analysis for seedling emergence and coleoptile using
the full effect (FULL), interaction effect (IE), and common effect (COM) multi-trait mixed
models for identifying significant loci controlling deep-sowing seedling emergence and
coleoptile length in a Pacific Northwest winter wheat diversity panel (DP) phenotyped across
trials from 2015-2019 in Lind, WA.

Stable Significant Markers
To validate the significant markers, it is important to compare their significance across years,
populations, and in correlated traits to determine their stability as significantly associated with
seedling emergence. As mentioned previously, no markers were found significant in both

seedling emergence and coleoptile length in either ST-GWAS or MT-GWAS. In the ST-GWAS
models, only 74 of the 107 markers were found significant in more than one GWAS. Within the
DP, 23 unique markers were found significant in more than one year across models. BLINK only
identified one marker, *S5A_522153944*, in multiple years. FarmCPU identified six markers
across years on chromosomes 2A (2), 2B (1), and 5A (3). MLM identified 16 markers across
years, with all but one on chromosome 5A with the other marker located on chromosome 5B. For
these stable markers in the DP, only *S5A_522153944* was identified by all three models.

     Out of the 107 unique significant markers, only four markers were identified in both
populations. Further, only *S5A_522153944*, *S5A_522153953*, and *S5A_523025549* were
identified across both populations with the same model (Table 6). MLM identified all three of
these markers in both populations. whereas BLINK did not identify any markers in both
populations, and FarmCPU only identified *S5A_522153944* in both populations. The effect of
covariates on identifying the stable markers was inconsistent in the DP. MLM displayed no
response to the inclusion of covariates (Figure S6). However, FarmCPU identified the stable
marker *S5A_522153944* on chromosome 5A using the Rht markers, coleoptile length, and Rht
markers with coleoptile length combined as covariates (Figure S7). Further, BLINK was able to
detect *S5A_522153944* with and without covariates but was not able to identify it in the BL
population (Figure S8).

Table 6. Stable significant markers across both the Diversity Panel and Breeding Lines. Significant loci controlling deep-sowing seedling emergence in a Pacific Northwest winter wheat breeding trial using BLINK, FarmCPU, and MLM genome-wide association studies (GWAS) models using covariates for coleoptile length and Rht alleles.

Marker	Position[a]	Alleles[b]	Chr[a]	Pop[c]	Year	Model	Covariate[d]	P-value	MAF[e]	R^{2f}	Effect (%)[g]
S5A_514260464	5.14E+08	<u>C</u>/T	5A	BL	15	FarmCPU	N	7.86E-07	0.23	0.13	17
				DP	15	MLM	N, R, C, B	1.81E-09	0.11	0.08	19
				DP	15-17	MLM	N, R, C, B	6.69E-07	0.11	0.05	9
				DP	15-18	MLM	N, R, C, B	6.48E-06	0.11	0.02	5
S5A_522153944	5.22E+08	<u>C</u>/T	5A	DP	15	BLINK	N, C	1.58E-11	0.04	0.09	29
				DP	15-17	BLINK	N, C, B	3.27E-12	0.04	0.08	17
				DP	15-18	BLINK	N, R, C	1.83E-06	0.04	0.10	15
				DP	15-19	BLINK	N, R, C, B	1.82E-09	0.04	0.06	12
				BL	15	FarmCPU	N	1.69E-06	0.04	0.12	30
				DP	15	FarmCPU	N, C	2.44E-06	0.04	0.09	29
				DP	15-18	FarmCPU	N, R	5.13E-06	0.04	0.10	15
				DP	15-19	FarmCPU	R, C, B	6.15E-06	0.04	0.06	12
				BL	15	MLM	N	9.29E-07	0.04	0.12	30
				DP	15	MLM	N, R, C, B	3.62E-09	0.04	0.09	29
				DP	15-17	MLM	N, R, C, B	7.77E-09	0.04	0.08	17
				DP	15-18	MLM	N, R, C, B	2.67E-09	0.04	0.10	15
S5A_522153953	5.22E+08	<u>A</u>/G	5A	DP	15-17	BLINK	R	6.88E-13	0.04	0.08	17
				DP	15	FarmCPU	B	1.03E-07	0.04	0.09	29
				DP	15-18	FarmCPU	B	4.54E-07	0.04	0.10	15
				BL	15	MLM	N	9.29E-07	0.04	0.12	30
				DP	15	MLM	N, R, C, B	3.62E-09	0.04	0.09	29
				DP	15-17	MLM	N, R, C, B	7.77E-09	0.04	0.08	17
				DP	15-18	MLM	N, R, C, B	2.67E-09	0.04	0.10	15
S5A_523025549	5.23E+08	<u>A</u>/G	5A	BL	15	BLINK	N	4.56E-11	0.04	0.12	31
				BL	15	MLM	N	2.57E-06	0.04	0.12	31
				DP	15	MLM	N, R, C, B	1.03E-06	0.05	0.06	23
				DP	15-17	MLM	N, R, C, B	1.15E-05	0.05	0.05	13
				DP	15-18	MLM	N, R, C, B	1.33E-06	0.05	0.07	12

[a]Chromosomes and positions of markers were determined according to IWGSC RefSeq v.1.1.

[b]Allele: favorable allele is underlined.

[c]Pop: BL: Breeding lines; DP: Diversity Panel

[d]Covariate: N: None; R: Reduced height alleles; C: BLUPs for coleoptile length; B: both reduced height and coleoptile length

When taking into account the number of significant markers across chromosomes, QQ

plots, covariate effect, and across population identification, FarmCPU displayed the ability to

identify both the stable large effect and small effect markers with fewer false positives on

chromosome 5A (Figure 6). In comparison, MLM consistently identified markers regardless of

covariates and population but also identified a large number of linked markers. For example,

MLM consistently identified both *S5A_522153944* and *S5A_522153953* which have an R^2 LD

value of 1, and identified 11 other markers on 5A with R^2 LD values above 0.80. Further, BLINK

was able to identify other small effect loci on other chromosomes but did not identify a

significant marker in both populations.

Figure 6. Stacked manhattan plot of Mixed-Linear Model (MLM), FarmCPU, and BLINK using the diversity
panel and breeding lines. Stacked manhattan plots for genome-wide association studies (GWAS) using MLM,
FarmCPU, and BLINK for identifying significant loci controlling deep-sowing seedling emergence in a Pacific

Northwest winter wheat diversity panel (DP) phenotyped across trials from 2015-2018 and a breeding line (BL) trial
phenotyped in 2015 in Lind, WA. Red triangles display GWAS results in the DP, and blue circles identify GWAS
results in the BL. Significant markers using an FDR cutoff with an alpha=0.05 are placed above the solid black line
for the DP and the dashed black line for the BL. Significant markers across both populations are identified with a
vertical red solid line identifying their positions. Markers enclosed in a white text box display significant markers
identified in GWAS studies for coleoptile length.

The stable markers identified across years in the DP were located on chromosome 2A (2),

2B (3), 5A (15), 5B (1), and 7A (1). The two largest effect stable markers in the DP,

S5A_522153944 and *S5A_522153953*, were found on chromosome 5A and had the largest effect

of 30% for seedling emergence. These two markers are in complete linkage with an R^2 LD value

of 1. They are both considered relatively large effect alleles and had a maximum R^2 of 10% in

the DP. The other stable markers across years identified on chromosome 5A were mainly

identified by MLM and their effects ranged from 3% to 29%, with R^2 values ranging from to 2%

to 8%. FarmCPU was able to identify stable markers on the other chromosomes mentioned

previously. However, they were minor effect markers with effects ranging from 4 to 24% on

seedling emergence and R^2 values from 1 to 6%. MLM was only able to identify the smaller

effect markers on chromosomes 2B and 5B. The multi-locus models, FarmCPU and BLINK,

were able to identify the remaining small effect markers primarily in the combined year trials.

Overall, the stable markers across years accounted for 30% of the total variation in the combined

2015-2019 DP.

GWAS in the BL population displayed more consistent results than in the DP, and only

identified stable markers on chromosome 5A. There were four significant stable markers

identified in the BL (Table 6). The four stable markers were all found in the DP, and had effect

ranges of 17 to 30%, with R^2 values of 12 to 13%. The stable markers in the BL population

accounted for 18% of the total variation for seedling emergence, displaying a more consistent

dissection of the complex trait compared to the DP.

The MT-GWAS models displayed very similar results for identifying stable markers

across years for both the single trait and joint models as the ST-GWAS models. The stable

markers for seedling emergence and coleoptile length as the ST-GWAS models, with the

seedling emergence stable markers being located exclusively on chromosomes 2B and 5A,

whereas the stable markers for coleoptile length were the same markers identified in the ST-

GWAS models on chromosomes 1A and 6A. In addition to the stable singe trait markers, the

COM MT-GWAS model identified 19 stable markers on chromosome 5A, including the large

effect marker *S5A_522153944* (Figure 7). In addition, the COM model identified stable markers

on chromosomes 2B (1), 5B (1), and 7B (3). The 2B and 5B markers were *S2B_301162652* and

S5B_491273019. The 7B markers are all completely linked with R^2 values of 1, with the marker

S7B_663828309 having the largest effect. The 7B markers were the only stable markers in MT-

GWAS not found stable in the ST-GWAS models. The 7B markers were only found significant

in the 2015-2017 combined DP for the ST-GWAS models, and therefore, were not deemed
stable.

Figure 7. Stacked manhattan plot of Multi-Trait GWAS models for Seedling Emergence, Coleoptile Length,
Full, Interaction Effect, and Common Models using the diversity panel lines. Stacked manhattan plots for multi-
trait genome-wide association studies (MT-GWAS) using seedling and emergence overlapped (EM_CL), the full
effect (Full), interaction effect (IE), and common effect (COM) multi-trait mixed models for identifying significant
loci controlling deep-sowing seedling emergence and coleoptile length in a Pacific Northwest winter wheat diversity
panel (DP) phenotyped across trials from 2015-2018 in Lind, WA. For the overlapped plot, EM_CL, red triangles
display GWAS results for seedling emergence, and blue circles identify GWAS results for coleoptile length.
Significant markers using a Bonferonni cutoff with an alpha=0.05 are placed above the solid black line for seedling
emergence and the dashed black line for coleoptile length. Significant markers across three models are highlighted
with a solid red vertical line, and significant markers across two models are highlighted with a dashed green vertical
line identifying their positions.

<u>Effect of Combining Favorable Alleles</u>

The frequencies of the stable markers across populations and the stable markers across years for

seedling emergence are important for understanding the populations' selection and makeup. The

stable markers with LD above 0.80 R^2 were binned together with the marker with the largest

effect identified to represent the bin. This resulted in three out of the four markers identified
across populations remaining and 12 out of 23 markers identified across years remaining for
displaying their additive effect for seedling emergence (Table S5). In addition, the significant
marker on 7B identified in the COM MT-GWAS model was included with the markers stable
across years. The majority of stable markers had high frequencies in the 90% range in both the
DP and BL (Table S5). This shows that the stable markers are already consistent in both
populations.
Even though the stable markers across populations and years were frequent in the
populations, the differences was shown when we combined them and compared the favorable
alleles' cumulative effect on seedling emergence (Figure 8A-D). We compared the additive
ability for favorable alleles using stable markers across populations and stable markers identified
in the DP in multiple years. For the stable markers across populations, the majority of lines in
both populations had favorable alleles for all three of the markers, which were all located on
chromosome 5A. Both populations showed an additive effect, but in the DP, the lines with all
three markers had a lower mean compared to lines with just two of the markers (Figure 8B).
However, in the BL there was an increasing trend in emergence with the accumulation in
favorable alleles (Figure 8A). The stable markers across the populations only accounted for 8%
of the total variation in the DP and 19% in the BL.
This additive effect is seen to a point for the stable markers across years for seedling
emergence in the BL. The DP showed an additive effect for all 11 favorable alleles, whereas the
BL showed a diminishing return signifying some of the markers may not have a large effect
within the BL population (Figure 8C and 8D). For the DP, the majority of lines had either eight
or nine favorable alleles with 113 and 125 lines, respectively. This was also the case in the BL
population with 72 lines having eight alleles and 72 lines having 9 alleles. The stable markers
across years accounted for only 23% of the variation in the BL and 31% in the DP.

Figure 8 Effect of markers on seedling emergence. Effect of pyramiding favorable alleles for seedling emergence for the stable markers identified in both populations (A and B), and stable markers identified across years in the DP (C and D) across both the (A and C) diversity panel (DP) and (B and D) breeding line (BL) populations.

Discussion

Complex Traits and Seedling Emergence

Complex traits are quantitative in nature and are affected by many small-effect QTLs (Holland 2007). The challenges that impede the understanding of complex traits are the inability to statistically detect and map minor effect QTLs, accurately understand GEI, and account for pleiotropic effects (Luo et al. 2017). This study attempted to characterize one such trait, seedling emergence for deep-sown winter wheat, to be used as a model for other complex traits.

The reduced height semi-dwarf genes *Rht-B1b* and *Rht-D1b* are mutant alleles that cause the dwarfing stature of wheat due to the decrease in response to endogenous gibberellic acid (Vogel et al. 1956; Allan et al. 1962; Peng et al. 1999). These dwarfing genes are responsible for

short stature and have a pleiotropic effect that includes gibberellin insensitivity, coleoptile
length, grain yield, protein content, and disease resistance (Allan et al. 1961; Allan 1989; Mohan
et al. 2013). The introgression of these genes into 'Norin 10' and the subsequent use by N.E.
Borlaug was an essential driver of the green revolution (Vogel et al. 1956; Allan 1989; Mohan et
al. 2013). Semi-dwarf wheat cultivars were resistant to lodging, which allowed more fertilizer
input and much-improved grain yield (Allan 1989; Mohan et al. 2013). A side effect of the semi-
dwarf genes' introgression was the reduction in coleoptile length of one-half to three-fourths of
the standard varieties at the time (Allan et al. 1961). The reduced coleoptile length is due to a
decreased response of GA, which reduced cell size and elongation (Allan et al. 1959).
Historically, when crusting was present, or other unfavorable conditions, the shorter coleoptiles
of semi-dwarf cultivars resulted in poor stand establishment and yield potential (Rebetzke et al.
1999). These previous studies showed the relationship between coleoptile length and seedling
emergence (Allan et al. 1962; Sunderman 1964; Chastain et al. 1995; Schillinger et al. 1998;
Botwright et al. 2001; Schillinger 2011). However, the study by Mohan et al. (2013) was able to
refute the previous claims in modern varieties using a collection of 662 wheat cultivars from
around the world. The previous studies had claimed that coleoptile length explained more than
60% of the variation for seedling emergence but only explained 28% in modern varieties (Mohan
et al. 2013).

Major QTLs for coleoptile length have been reported on chromosomes 4BS, 4BL, and
5AL in wheat (Rebetzke et al. 2001). The QTLs on 4B were reported on either side of the *Rht-B1*
allele. This study was further enhanced by a subsequent QTL analysis that resolved the two 4B
QTLs directly to the *Rht-B1* locus (Rebetzke et al. 2007). In our study, no markers were located
on 4B with two small effect markers on 4D found only in the joint COM and FULL MT-GWAS
models, but were not found stable across years. The major stable markers validated across
populations for seedling emergence in our study were identified on chromosome 5A. However,
in the follow-up study in Rebetzke et al. (2007), the QTL for coleoptile length on 5A was not
found repeatable across two or more populations. There have been small-effect QTLs for
coleoptile length identified on wheat chromosomes 1A, 2B, 2D, 3A, 3B, 5A, 6A (Rebetzke et al.
2007). We found significant markers in our GWAS models on these chromosomes, with our
large effect markers also located on chromosome 5A. However, in the DP, significant markers
for coleoptile length were located on chromosomes 1A, 2A, and 6B, and none of these markers
were significant for seedling emergence. Therefore, we can conclude that our GWAS models are
not selecting major markers for coleoptile length. Further, we can conclude that the seedling
emergence we are dissecting is indeed due to other factors affecting seedling emergence or
another poorly understood mechanism similar to what was indicated in Mohan et al. (2013).

Our study demonstrated that chromosome 5A has a large effect on seedling emergence
with four large effect stable markers identified consistently across both years in the DP and

validated in the BL. Therefore, a major effect QTL for seedling emergence may be present in
that chromosomal location. In our GWAS models, the significant markers are possibly affecting
both fast emergence and the ability to deal with moisture stress. The variation for emergence in
the DP and BL was due to moisture stress, and therefore, the significant markers would affect
both fast-emergence and the ability to deal with moisture stress. However, as mentioned
previously, seedling emergence is a complex system of multiple factors and pleiotropic effects
that are not controlled by genes for any sole component.
The lack of consistency between GWAS models and populations in individual years in
the DP confirms the complexity of seedling emergence and displays the reason why seedling
emergence evaded genetic characterization for so long. The complex quantitative nature of
seedling emergence is indicated due to a few major QTLs as well as QTL-by-Environment
Interaction (QEI). According to Bernardo (2014), a major marker accounts for >10% of the
phenotypic variation. By this definition, a few large effect markers have a range of up to 30%
effect on emergence. The identification of a few large effect markers with mainly small effect
significant markers by GWAS confirms that seedling emergence is a complex trait controlled by
many small-effect loci that may be affected by multiple pleiotropic factors.

Genome-Wide Association Models
Genetic mapping through association studies has been used to dissect the genetic architecture of
various traits in wheat (Adhikari et al. 2012; Lozada et al. 2017, 2019; Bajgain et al. 2019;
Lozada and Carter 2020). However, GWAS has not been implemented for deep-sowing seedling
emergence in winter wheat but has been conducted in rice (Zhao et al. 2018). Even further, few
studies have compared ST-GWAS models, and fewer have compared FarmCPU and BLINK
(Liu et al. 2016; Huang et al. 2019). By comparing multiple covariates and GWAS models, we
attempted to characterize and dissect a trait with unknown genetic architecture. This allowed us
to identify stable markers that are large effect markers with up to 30% effect and 10% R^2 values.
Therefore, we can conclude that these markers are important for seedling emergence and their
additivity is noted by an increase in emergence with their accumulation.
The single-locus model used in our study was MLM. The MLM identified significant
markers mainly on the 5A chromosome. The MLM has been shown to have difficulties with low
power and false positives when identifying significant markers as compared to FarmCPU (Liu et
al. 2016). This problem occurs in our study within the DP, and therefore, MLM displays low
power in identifying both the major markers and other lower effect significant markers. In our
study this was due to high levels of LD between significant markers. The MLM was more
consistent in identifying the large effect markers on 5A without the inclusion of the covariates.
MLM has been shown to be comparable to FarmCPU for simple traits (Habier et al. 2011; Ward
et al. 2019). However, for complex traits such as seedling emergence, the multi-locus models
show higher power for small effect markers. The MLM-based models evaluate each genetic

marker's relationship and the overall variation independently; therefore, as the trait increases in

complexity and number of pleiotropic effects, the proportion of variation due to a locus

decreases (Miao et al. 2019).

FarmCPU and multi-locus models provide the best trade-off between power and false

positives (Liu et al. 2016; Miao et al. 2019). The multi-locus models increase the proportion of

genetic variance by using the major effect markers as fixed effects and identifying more

significant markers (Miao et al. 2019). This is why our multi-locus models were able to dissect

the complex trait of seedling emergence accurately and identified both major and minor effect

markers. The multi-locus models used in our study were FarmCPU and BLINK and had very

similar results in the BL population, but there were fewer significant markers discovered by

BLINK than by FarmCPU in the DP. The difference resulted in BLINK not identifying a

significant marker in both populations. Other than the markers on 5A that MLM also found, most

markers found significant by FarmCPU and BLINK had relatively small effects and R^2 values

indicating multi-locus models' ability to identify small-effect markers for complex traits.

The majority of markers identified by the MT-GWAS models were also identified in the

ST-GWAS models. Further, we had to use the Bonferroni cutoff because the MT-GWAS models

displayed inflation of p-values in the QQ plots for the joint analyses. Additionally, the MT-

GWAS model uses a single-locus model, which may additionally add to the inflated p-values.

Therefore, there was little to no increase in the power for identifying pleiotropic loci compared to

the ST-GWAS models. However, the COM model did identify stable markers on chromosome

7A that were not found stable in any ST-GWAS model but were identified in a single year

combination. The lack of increase in power for the MT-GWAS models may be due to the lack of

phenotypic correlation (Korte et al. 2012). Further, the lack of significant interaction effect loci

confirms the positive genetic correlation seen in the majority of years between seedling

emergence and coleoptile length. Even though we see a positive genetic correlation, the

phenotypic correlation is near zero, indicating a small genetic effect in comparison to the

environmental effect. In this scenario, it has been shown that the MT-GWAS models do not

outperform ST-GWAS models and confirm why our MT-GWAS models do not outperform the

ST-GWAS models (Korte et al. 2012). However, MTMM can still be used as a complement to

ST-GWAS models to identify interactions and pleiotropic loci such as the markers on

chromosome 7A.

<u>Covariates</u>

In our study, all of the ST-GWAS and MT-GWAS models used included covariates for

population structure as principal components (Price et al. 2006). In addition, the MLM used

kinship matrices calculated by VanRaden's genomic relationship matrix and implemented

through GAPIT (VanRaden 2008; Tang et al. 2016). These covariates are commonly used in

GWAS studies and allow the models to differentiate genetic and environmental effects (Tibbs

Cortes et al. 2021). However, for a complex trait, more confounding associations such as
correlated traits or pleiotropic effects may account for a portion of seedling emergence variation
(Saltz et al. 2017).

To account for these correlated traits, we compared the use of covariates for Rht markers
and coleoptile length. These correlated traits were used due to previous studies reporting large
associations between the traits (Allan et al. 1962; Sunderman 1964; Chastain et al. 1995;
Schillinger et al. 1998; Botwright et al. 2001; Schillinger 2011). The Rht and coleoptile length
covariates had an effect on the multi-locus models but not the MLM. The multi-locus models
required the inclusion of covariates in order to identify the major effect of significant markers on
chromosome 5A. However, this was not the case for identifying the small effect markers.
Therefore, the multi-locus models display a weakness in complex traits to identify both small
effect and large effect markers when faced with pleiotropic effects. This signifies that even
though modern varieties are no longer dependent on coleoptile length for improved emergence,
they still confer pleiotropic effects, confirmed by the results for our MT-GWAS models. We see
in the DP that coleoptile length still has a large genotypic correlation depending on the year.
Therefore, we recommend using both multi-locus models and covariates for correlated traits to
identify both small and large effect loci and increase the power to dissect the genetic architecture
of complex traits.

<u>Association Mapping Populations and Environments</u>
We evaluated the population composition of our populations by using PCs. The DP had a more
distinct population structure compared to the BL. This indicates and confirms that the BL
population is more genetically related. We used PCs as fixed effects in our GWAS to account for
and remove bias due to the presence of underlying population structure in our populations (Price
et al. 2006). The closer related the genotypes are, the fewer recombination events, thus
preserving marker linkage disequilibrium, and less genetic variation needed to be accounted for
by the models (Habier et al. 2007; Bernardo 2020). Due to the increased genetic relatedness and
less population structure in the BL, the resolution was increased, and fewer false positives were
identified as seen by the large difference in MLM's significant markers. This may be due to the
fact that the BL was purposely selected over generations for deep-sowing seedling emergence in
the Washington State University breeding program. The DP on the other hand is composed of
varieties from various breeding programs in the Pacific Northwest, with the majority of lines not
bred specifically for deep-sowing seedling emergence. Even with this discrepancy, we were still
able to identify the stable markers on chromosome 5A in both populations.

The number of environments examined between the two populations differed in addition
to the differences between the composition and genetic relatedness of the populations. In the BL,
we used a population in a single environment. This restricted our ability to account for GEI in
our phenotypic adjustments. However, with the close genetic relationship between the lines, we

were still able to identify major effect markers. Whereas in the DP, for both the ST-GWAS and

MT-GWAS models, some years had very little significant markers compared to 2015 or the

combined years. The year 2015 was important for stable marker identification, with no other

individual year identifying the stable markers across populations on chromosome 5A for seedling

emergence. Therefore, combining years to partition GEI enabled GWAS models to dissect

genetic architectures accurately.

Since seedling emergence is dependent on the environment to create phenotypic

variation, it displays GEI. If a trait displays GEI, it follows that so would the QTLs responsible

for the phenotypic expression (Bernardo 2020). A change in the ranking of QTLs across

environments indicates QEI, and detection of QTLs in some environments, and not others, have

been used as a criterion for declaring QEI (Bernardo 2020). Therefore, QEI can be seen with the

differing number of significant markers in the different years and the combination of years.

Furthermore, the difficulty in dissecting seedling emergence within and across years can

be seen in the discrepancy between the varying genetic and phenotypic correlations between

years and traits. The differences in genetic correlations for seedling emergence and coleoptile

length from year to year and the near-zero phenotypic correlations can be explained by the large

effect of the environment and the multitude of factors that affect seedling emergence. The

phenotypic effect can be partitioned into both genetic and non-genetic effects such as the

environmental effect (Searle 1961). Since genetic correlation only takes into account the genetic

effect, we still see moderate values. The negative value in 2017 can be explained by the different

factors that affect seedling emergence, such as coleoptile diameter, force, and speed of

emergence. This is further confirmed to be the result of another factor due to the lack of

significant markers for the interaction effect in the MT-GWAS models (Korte et al. 2012). As

mentioned previously, if the negative genetic correlation was due to the significant loci for

coleoptile length, we would see significant markers for the interaction effect (Korte et al. 2012).

Further, the near-zero phenotypic correlations display that coleoptile length, while having

moderate genetic correlations, has a much smaller genetic effect than the environmental effects

such as moisture stress and crusting, and therefore has a little overall phenotypic effect for

seedling emergence. In other terms, when genes controlling two traits, such as seedling

emergence and coleoptile length, are similar but where the environments pertaining to the

expression of the traits have a low correlation. The phenotypic correlation would then be lower

than the genetic correlation (Searle 1961).

The inconsistency between years displays the difficulty of dissecting a complex trait

affected by multiple factors and heavily dependent on the environment to display phenotypic

variation such as seedling emergence. However, the stable markers for seedling emergence on

5A were still identified in both populations across years as well as in both ST-GWAS and MT-

GWAS models. Therefore, we can conclude that the genetic architecture's dissection was
successful for both the unselected DP population and selected BL populations.

<u>Combining favorable alleles</u>
We see a clear increase and additive effect of accumulating favorable alleles in the DP for the
stable markers over years. Additionally, there was an additive effect of accumulating favorable
alleles for the stable markers over populations in the BL. The difference between the two sets of
markers were that the stable markers over populations consisted exclusively of markers on
chromosome 5A, whereas the stable markers over years consisted of markers on various
chromosomes. This discrepancy shows that the BL may be controlled mainly by the 5A
chromosomes with little effect of the other small effect markers on the other chromosomes.
Further, seedling emergence for the DP is controlled by both the large-effect and small effect
markers on various chromosomes.
The differences in additivity in the populations displays the potential differences between
populations purposely selected for complex traits such as seedling emergence and the success of
accumulating favorable alleles through traditional phenotypic selection. With the 60 plus years
of selection in the BLs, a larger number of favorable alleles may have been accumulated than the
DP, consisting of lines from various programs and not centered around deep-sowing emergence
selection. Therefore, we can see that through traditional phenotypic selection, the Washington
State University breeding program has been able to increase seedling emergence while
maintaining the inclusion of Rht alleles and semi-dwarf stature.
Evaluating the cumulative effect of favorable alleles allows us to make assumptions
about the types of genetic effects that control seedling emergence. By displaying an increase in
seedling emergence with favorable alleles' accumulation, we can assume that an additive effect
conveys seedling emergence. It also shows the ability to pyramid favorable alleles for further
improvement of cultivars. Pyramiding favorable alleles has shown to be successful for disease
resistance traits (Naruoka et al. 2015; Lewien et al. 2018; Lozada et al. 2019).

<u>Applications in Breeding</u>
This study compared GWAS models, populations, and covariates in the attempt to dissect the
genetic architecture of a complex trait with pleiotropic effects that's previously been dependent
on the environment for phenotypic screening and selection. For these complex traits, including
covariates and correlated traits created an advantage in the trait's dissection. The complex genetic
architecture with pleiotropic effects, both small and large effect markers, and the markers'
additivity allowed us to conclude that modern breeding programs no longer are dictated by the
large number of studies displaying the dependence of the emergence of dwarf-varieties on
coleoptile length. Therefore, this study has also shown empirical results for the ability to breed
for correlated traits and overcome perceived genetic disadvantages. In the dryland winter wheat
region in Washington, where deep-sowing of winter wheat occurs, plant breeding efforts of 60

years have been able to develop modern varieties that are fast emerging while having the GA

insensitive mutant alleles *Rht-B1b* and *Rht-D1b*.

We have also shown the importance of multi-locus models over single-locus models for

highly complex traits. The multi-locus models increase in power allows identifying small effect

markers located around the genome while accurately identifying the major effect markers as seen

in previous comparison (Li et al. 2015; Liu et al. 2016; Miao et al. 2019; Ward et al. 2019;

Huang et al. 2019). Further, the MT-GWAS models displayed a lack of power over ST-GWAS

models in dissecting correlated traits with low phenotypic correlation as shown in previous

studies (Korte et al. 2012).

Even though the large effect markers are advantageous for selecting and improving the

trait, identifying smaller effect markers indicates that marker-assisted selection may not be as

successful for seedling emergence compared to simpler traits. Therefore, by dissecting the

genetic architecture, we can make breeding decisions on the most accurate method for rapid

improvements, such as the inclusion of genomic selection or phenotypic selection that may be

more accurate for substantial genetic gain (Bernardo 2008). For instance, our GWAS models

conclude there are only a few large effect markers while identifying mainly small effect markers.

The use of genomic selection may be more successful while still having the capability of

including large effect markers as fixed effects in the genomic selection models. As in this case,

the markers convey effects of 30% for seedling emergence and display additive effects. These

high effects meet the criteria for inclusion set forward by Bernardo (2014). Even when GWAS

and dissections may not provide results in identifying candidate genes or large effect markers for

use in selection, it can allow us to conclude the most effective method of selection, which allows

breeders to be more efficient with their limited resources while meeting their program's goals

and targeted genetic gains.

Conclusion

This study displayed the ability for GWAS to dissect the genetic architecture of a

complex trait such as deep-sown seedling emergence. Both ST-GWAS and MT-GWAS models

identified a few large effect and many small effect markers for seedling emergence.

Additionally, neither the ST-GWAS or MT-GWAS models identified large pleiotropic effect

markers between seedling emergence and coleoptile length. The ST-GWAS and MT-GWAS

models did not identify the same significant markers for seedling emergence and coleoptile

length, and the MT-GWAS models did not identify any interaction effect markers. However, by

using multi-locus models in conjunction with covariates for correlated traits, we were able to

increase power over single-locus models to dissect a complex traits in both the DP and BL

populations. Additionally, the DP displayed the necessity for combining years for consistent

identification of significant markers for a trait dependent on the environment for phenotypic

variation. Further, the MT-GWAS models displayed a lower power over ST-GWAS models for

single-trait analysis and inflated p-values for joint analysis but still identified the large effect
markers on 5A. Further, the MT-GWAS model uses a single-locus model, which may add to the
inflated p-values. Therefore, the MT-GWAS models displayed a lack of power in dissecting
correlated traits with low phenotypic correlation. Therefore, by using multi-locus models
combined with pleiotropic covariates, breeding programs can uncover the complex nature of
traits to help identify candidate genes and the underlying architecture of a trait to make more
efficient breeding decisions and selection methods.

Declarations

Funding: This research was partially funded by the National Institute of Food and Agriculture
(NIFA) of the U.S. Department of Agriculture (Award number 2016-68004-24770), Hatch
project 1014919, and the O.A. Vogel Research Foundation at Washington State University.:

Conflicts of Interest: The authors declare that the research was conducted in the absence of any
commercial or financial relationships that could be construed as a potential conflict of interest.

Availability of data and material: The datasets generated for this study can be found at
https://github.com/lfmerrick21/GWAS-Complex-Traits

Author's Contributions: LM: conceptualized the idea, analyzed data, and drafted the
manuscript; AB: genotyped the KASP markers, reviewed, and edited the manuscript; ZZ:
reviewed and edited the manuscript; AC: supervised the study, conducted field trials, edited the
manuscript, and obtained the funding for the project.

Acknowledgments: The authors would like to acknowledge the Washington State University
Winter Wheat Breeding Program personnel Gary Shelton and Kyall Hagemeyer for plot
maintenance and data collection under field conditions and Kerry Barlow for coleoptile length
data collection. We would also like to thank Gina Brown-Guedira, Jared Smith, Brian Ward and
staff at the Eastern Regional Small Grains Genotyping Laboratory for their assistance with DNA
library prep and GBS sequencing and analysis.

References

Adhikari TB, Gurung S, Hansen JM, et al (2012) Association Mapping of Quantitative Trait Loci in Spring
Wheat Landraces Conferring Resistance to Bacterial Leaf Streak and Spot Blotch. Plant Genome J
5:1. https://doi.org/10.3835/plantgenome2011.12.0032

Allan RE (1989) Agronomic comparisons between Rht1 and Rht2 semidwarf genes in winter wheat. Crop
Sci 29:1103–1108

Allan RE, Vogel OA, Burleigh JR, Peterson CJ (1961) Inheritance of Coleoptile Length and Its Association
with Culm Length in Four Winter Wheat Crosses 1. Crop Sci 1:328–332

Allan RE, Vogel OA, Craddock Jr JC (1959) Comparative response to gibberellic acid of dwarf, semidwarf,
and standard short and tall winter wheat varieties 1. Agron J 51:737–740

Allan RE, Vogel OA, Peterson CJ (1962) Seedling Emergence Rate of Fall-sown Wheat and Its Association
with Plant Height and Coleoptile Length [1]. Agron J 54:347–350.
https://doi.org/10.2134/agronj1962.00021962005400040022x

Appels R, Eversole K, Feuillet C, et al (2018) Shifting the limits in wheat research and breeding using a
fully annotated reference genome. Science 361:eaar7191

Arndt W (1965) The impedance of soil seals and the forces of emerging seedlings. Soil Res 3:55–68

Aschard H, Vilhjálmsson BJ, Joshi AD, et al (2015) Adjusting for heritable covariates can bias effect
estimates in genome-wide association studies. Am J Hum Genet 96:329–339

Bajgain P, Zhang X, Anderson JA (2019) Genome-Wide Association Study of Yield Component Traits in
Intermediate Wheatgrass and Implications in Genomic Selection and Breeding. G3amp58
GenesGenomesGenetics 9:2429–2439. https://doi.org/10.1534/g3.119.400073

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach
to multiple testing. J R Stat Soc Ser B Methodol 57:289–300

Bernardo R (2020) Breeding for quantitative traits in plants, 3rd ed. Stemma Press, Woodbury,
Minnesota

Bernardo R (2008) Molecular Markers and Selection for Complex Traits in Plants: Learning from the Last
20 Years. Crop Sci 48:1649. https://doi.org/10.2135/cropsci2008.03.0131

Botwright T, Rebetzke G, Condon T, Richards R (2001) The effect of rht genotype and temperature on
coleoptile growth and dry matter partitioning in young wheat seedlings. Funct Plant Biol
28:417–423. https://doi.org/10.1071/pp01010

Browning BL, Zhou Y, Browning SR (2018) A one-penny imputed genome from next-generation reference
panels. Am J Hum Genet 103:338–348

Chastain TG, Ward KJ, Wysocki DJ (1995) Stand Establishment Responses of Soft White Winter Wheat to
Seedbed Residue and Seed Size. 6

Covarrubias-Pazaran G (2018) Software update: Moving the R package sommer to multivariate mixed
models for genome-assisted prediction. bioRxiv 354639

Cullis BR, Smith AB, Coombes NE (2006) On the design of early generation variety trials with correlated
data. J Agric Biol Environ Stat 11:381. https://doi.org/10.1198/108571106X154443

Elshire RJ, Glaubitz JC, Sun Q, et al (2011) A Robust, Simple Genotyping-by-Sequencing (GBS) Approach
for High Diversity Species. PLoS ONE 6:e19379. https://doi.org/10.1371/journal.pone.0019379

Federer WF (1956) Experimental design. LWW

Gale MD, Youssefian S (1983) Pleiotropic effects of the Norin 10 dwarfing genes, Rht1 and Rht2, and
interactions in response to chlormequat. In: Proceedings of the sixth International Wheat
Genetics Symposium/edited by Sadao Sakamoto. Kyoto: Plant Germ-Plasm Institute, Faculty of
Agriculture, Kyoto University …

Galesloot TE, Van Steen K, Kiemeney LA, et al (2014) A comparison of multivariate genome-wide
association methods. PloS One 9:e95923

Goldman I (2019) Plant Breeding Reviews. John Wiley & Sons

Habier D, Fernando RL, Dekkers JCM (2007) The Impact of Genetic Relationship Information on Genome-
Assisted Breeding Values. Genetics 177:2389–2397.
https://doi.org/10.1534/genetics.107.081190

Habier D, Fernando RL, Kizilkaya K, Garrick DJ (2011) Extension of the Bayesian alphabet for genomic
selection. BMC Bioinformatics 12:1–12

Holland JB (2007) Genetic architecture of complex traits in plants. Curr Opin Plant Biol 10:156–161

Huang M, Liu X, Zhou Y, et al (2019) BLINK: a package for the next level of genome-wide association
studies with both individuals and markers in the millions. GigaScience 8:.
https://doi.org/10.1093/gigascience/giy154

Klei L, Luca D, Devlin B, Roeder K (2008) Pleiotropy and principal components of heritability combine to
increase power for association analysis. Genet Epidemiol Off Publ Int Genet Epidemiol Soc 32:9–
19

Korte A, Vilhjálmsson BJ, Segura V, et al (2012) A mixed-model approach for genome-wide association
studies of correlated traits in structured populations. Nat Genet 44:1066–1071.
https://doi.org/10.1038/ng.2376

Lander ES, Schork NJ (1994) Genetic dissection of complex traits. Science 265:2037–2048

Lewien MJ, Murray TD, Jernigan KL, et al (2018) Genome-wide association mapping for eyespot disease
in US Pacific Northwest winter wheat. PloS One 13:e0194698

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler transform.
bioinformatics 25:1754–1760

Li X, Li X, Fridman E, et al (2015) Dissecting repulsion linkage in the dwarfing gene Dw3 region for
sorghum plant height provides insights into heterosis. Proc Natl Acad Sci 112:11823–11828

LiLin Y (2018) Package "CMplot." 7

Lipka AE, Kandianis CB, Hudson ME, et al (2015) From association to prediction: statistical methods for
the dissection and selection of complex traits in plants. Curr Opin Plant Biol 24:110–118.
https://doi.org/10.1016/j.pbi.2015.02.010

Liu X, Huang M, Fan B, et al (2016) Iterative usage of fixed and random effect models for powerful and efficient genome-wide association studies. PLoS Genet 12:

Lozada D, Godoy JV, Murray TD, et al (2019) Genetic Dissection of Snow Mold Tolerance in US Pacific Northwest Winter Wheat Through Genome-Wide Association Study and Genomic Selection. Front Plant Sci 10:1337. https://doi.org/10.3389/fpls.2019.01337

Lozada DN, Carter AH (2020) Insights into the Genetic Architecture of Phenotypic Stability Traits in Winter Wheat. Agronomy 10:368

Lozada DN, Mason RE, Babar MA, et al (2017) Association mapping reveals loci associated with multiple traits that affect grain yield and adaptation in soft winter wheat. Euphytica 213:222

Luo Z, Wang M, Long Y, et al (2017) Incorporating pleiotropic quantitative trait loci in dissection of complex traits: seed yield in rapeseed as an example. Theor Appl Genet 130:1569–1585

Lutcher LK, Wuest SB, Johlke TR (2019) First Leaf Emergence Force of Three Deep-Planted Winter Wheat Cultivars. Crop Sci 59:772–777. https://doi.org/10.2135/cropsci2018.08.0495

Marchini J, Howie B, Myers S, et al (2007) A new multipoint method for genome-wide association studies by imputation of genotypes. Nat Genet 39:906–913

Merrick LF, Carter AH (2021) Comparison of Genomic Selection Models for Exploring Predictive Ability of Complex Traits in Breeding Programs. bioRxiv

Miao C, Yang J, Schnable JC (2019) Optimising the identification of causal variants across varying genetic architectures in crops. Plant Biotechnol J 17:893–905

Mohan A, Schillinger WF, Gill KS (2013) Wheat Seedling Emergence from Deep Planting Depths and Its Relationship with Coleoptile Length. PLoS ONE 8:e73314. https://doi.org/10.1371/journal.pone.0073314

Naruoka Y, Garland-Campbell KA, Carter AH (2015) Genome-wide association mapping for stripe rust (Puccinia striiformis F. sp. tritici) in US Pacific Northwest winter wheat (Triticum aestivum L.). Theor Appl Genet 128:1083–1101. https://doi.org/10.1007/s00122-015-2492-2

O'Reilly PF, Hoggart CJ, Pomyen Y, et al (2012) MultiPhen: joint model of multiple phenotypes can increase discovery in GWAS. PloS One 7:e34861

Peng J, Richards DE, Hartley NM, et al (1999) 'Green revolution'genes encode mutant gibberellin response modulators. Nature 400:256–261

Poland J, Endelman J, Dawson J, et al (2012) Genomic selection in wheat breeding using genotyping-by-sequencing. Plant Genome 5:103–113

Porter HF, O'Reilly PF (2017) Multivariate simulation framework reveals performance of multi-trait GWAS methods. Sci Rep 7:1–12

Price AL, Patterson NJ, Plenge RM, et al (2006) Principal components analysis corrects for stratification in
genome-wide association studies. Nat Genet 38:904–909

R Core Team (2018) R: A Language and Environment for Statistical Computing. Version 3.5.1. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Rasheed A, Wen W, Gao F, et al (2016) Development and validation of KASP assays for genes
underpinning key economic traits in bread wheat. Theor Appl Genet 129:1843–1860

Rebetzke GJ, Appels R, Morrison AD, et al (2001) Quantitative trait loci on chromosome 4B for coleoptile
length and early vigour in wheat (Triticum aestivum L.). Aust J Agric Res 52:1221–1234

Rebetzke GJ, Ellis MH, Bonnett DG, Richards RA (2007) Molecular mapping of genes for coleoptile
growth in bread wheat (Triticum aestivum L.). Theor Appl Genet 114:1173–1183

Rebetzke GJ, Richards RA, Fischer VM, Mickelson BJ (1999) Breeding long coleoptile, reduced height
wheats. Euphytica 106:159–168

Saltz JB, Hessel FC, Kelly MW (2017) Trait correlations in the genomics era. Trends Ecol Evol 32:279–290

SAS Institute, Inc (2011) SAS® 9.3 system options: Reference. SAS Institute Inc Cary, NC

Schillinger WF (2011) Rainfall Impacts Winter Wheat Seedling Emergence from Deep Planting Depths.
Agron J 103:730–734. https://doi.org/10.2134/agronj2010.0442

Schillinger WF, Donaldson E, Allan RE, Jones SS (1998) Winter Wheat Seedling Emergence from Deep
Sowing Depths. Agron J 90:582–586.
https://doi.org/10.2134/agronj1998.00021962009000050002x

Schillinger WF, Schofstoll SE, Smith TA, Jacobsen JA (2017) Laboratory Method to Evaluate Wheat
Seedling Emergence from Deep Planting Depths. Agron J 109:2004–2010.
https://doi.org/10.2134/agronj2016.12.0715

Searle SR (1961) Phenotypic, genetic and environmental correlations. Biometrics 17:474–480

Solovieff N, Cotsapas C, Lee PH, et al (2013) Pleiotropy in complex traits: challenges and strategies. Nat
Rev Genet 14:483–495

Stephens M (2013) A unified framework for association analysis with multiple related phenotypes. PloS
One 8:e65245

Sunderman DW (1964) Seedling Emergence of Winter Wheats and Its Association With Depth of Sowing,
Coleoptile Length Under Various Conditions, and Plant Height [1]. Agron J 56:23–25.
https://doi.org/10.2134/agronj1964.00021962005600010008x

Tang Y, Liu X, Wang J, et al (2016) GAPIT Version 2: An Enhanced Integrated Tool for Genomic
Association and Prediction. Plant Genome 9:0.
https://doi.org/10.3835/plantgenome2015.11.0120

Tibbs Cortes L, Zhang Z, Yu J (2021) Status and prospects of genome-wide association studies in plants.
Plant Genome 14:e20077

Tibbs Cortes L, Zhang Z, Yu J Status and prospects of genome-wide association studies in plants. Plant
Genome e20077

VanRaden PM (2008) Efficient Methods to Compute Genomic Predictions. J Dairy Sci 91:4414–4423.
https://doi.org/10.3168/jds.2007-0980

Vogel OA, Craddock Jr JC, Muir CE, et al (1956) Semidwarf Growth Habit in Winter Wheat Improvement
for the Pacific Northwest 1. Agron J 48:76–78

Wang Q, Tian F, Pan Y, et al (2014) A SUPER powerful method for genome wide association study. PloS
One 9:e107684

Ward BP, Brown-Guedira G, Kolb FL, et al (2019) Genome-wide association studies for yield-related traits
in soft red winter wheat grown in Virginia. PloS One 14:e0208217

Yu J, Pressoir G, Briggs WH, et al (2006) A unified mixed-model method for association mapping that
accounts for multiple levels of relatedness. Nat Genet 38:203–208

Zhang Z, Ersoz E, Lai C-Q, et al (2010) Mixed linear model approach adapted for genome-wide
association studies. Nat Genet 42:355–360

Zhao Y, Zhao W, Jiang C, et al (2018) Genetic architecture and candidate genes for deep-sowing
tolerance in rice revealed by non-syn GWAS. Front Plant Sci 9:332

1054