Prospective cohort study reveals unexpected aetiologies of livestock abortion in northern Tanzania

Kate M. Thomas[1,2*$], Tito Kibona[3], John R. Claxton[4], William A. de Glanville[4], Felix Lankester[5,6], Nelson Amani[2], Joram J. Buza[3], Ryan W. Carter[4], Gail E. Chapman[7], John A. Crump[1,8], Mark P. Dagleish[9], Jo E. B. Halliday[4], Clare M. Hamilton[9], Elisabeth A. Innes[9], Frank Katzer[9], Morag Livingstone[9], David Longbottom[9], Caroline Millins[7#], Blandina T. Mmbaga[2,8], Victor Mosha[2], James Nyarobi[4], Obed M. Nyasebwa[10], George C. Russell[9], Paul N. Sanka[11], George Semango[3], Nick Wheelhouse[12], Brian J. Willett[13], Sarah Cleaveland[4], and Kathryn J. Allan[4,7]

[1]Centre for International Health, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand
[2]Kilimanjaro Clinical Research Institute, Good Samaritan Foundation, Moshi, United Republic of Tanzania
[3]Nelson Mandela African Institution of Science and Technology (NM-AIST), Tengeru, United Republic of Tanzania
[4]Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom
[5]Paul G. Allen School for Global Animal Health, Washington State University, WA, United States of America
[6]Global Animal Health Tanzania, Arusha, United Republic of Tanzania
[7]School of Veterinary Medicine, University of Glasgow, Glasgow, United Kingdom
[8]Kilimanjaro Christian Medical University College, Moshi, United Republic of Tanzania
[9]Moredun Research Institute, Midlothian, United Kingdom
[10]Zonal Veterinary Centre – Arusha, Ministry of Livestock and Fisheries, Arusha, United Republic of Tanzania
[11]Tanzania Veterinary Laboratory Agency, Arusha, United Republic of Tanzania
[12]School of Applied Sciences, Edinburgh Napier University, Edinburgh, United Kingdom
[13]Medical Research Council, University of Glasgow Centre for Virus Research, Glasgow, United Kingdom

* Corresponding author
E-mail: katethomas144@gmail.com OR Kate.Thomas@mpi.govt.nz (KMT)

$ Current address: New Zealand Food Safety, Ministry for Primary Industries, Wellington, New Zealand
# Current address: Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, United Kingdom

Abstract

Livestock abortion is an important cause of productivity losses worldwide and many infectious causes of abortion are zoonotic pathogens that impact on human health. Little is known about the relative importance of infectious causes of livestock abortion in Africa, including in subsistence farming communities that are critically dependent on livestock for food, income, and wellbeing. We conducted a prospective cohort study of livestock abortion, supported by cross-sectional serosurveillance, to determine aetiologies of livestock abortions in livestock in Tanzania. This approach generated several important findings including detection of a Rift Valley fever virus outbreak in cattle; high prevalence of *C. burnetii* infection in livestock; and the first report of *Neospora caninum*, *Toxoplasma gondii,* and pestiviruses associated with livestock abortion in Tanzania. Our approach provides a model for abortion surveillance in resource-limited settings. Our findings add substantially to current knowledge in sub-Saharan Africa, providing important evidence from which to prioritise disease interventions.

Introduction

Infectious causes of abortion, including early foetal loss and stillbirth, affect livestock production worldwide with the potential to cause major economic losses [1]. Many abortions are caused by zoonotic pathogens that also pose a risk of infection to people [2]. Although the economic losses can be substantial in high-production, intensive agricultural systems [1], abortion-associated livestock production losses have marked negative impacts for the poorest communities in the world, many millions of whom are critically dependent on livestock for economic livelihoods, food security, health, and wellbeing [3, 4].

Livestock abortions can be caused by a wide range of pathogens as well as non-infectious causes such as nutritional deficiencies and trauma [2]. While many infectious causes of abortions occur worldwide, aetiological data based on robust laboratory-confirmed diagnoses are dominated by studies from farming systems in high- and middle-income countries [5, 6, 7] with data lacking from low-income settings. Comparison of aetiologies across geographic areas is challenging as studies vary in methodologies and in the specific pathogens that are investigated. Furthermore, the ability to identify a specific aetiology of a livestock abortion is often relatively low, with less than 35% cases attributed to specific diagnoses [5, 7, 8].

Considerable surveillance and diagnostic infrastructure is required for robust case attribution of livestock abortions [5, 9], which typically limits the data available from low resource settings. For countries where data do exist, geographic patterns appear to vary quite widely. For example, *Coxiella burnetii* has been determined to be a common cause of abortion in sheep and goats in Canada [10], and in goats in the

Netherlands [11], but only rarely in sheep in the United States of America (USA) [8, 12] or

the United Kingdom (UK) [9], despite evidence of widespread seropositivity in both

countries [13, 14]. *Chlamydia abortus* is one of the most common causes of infectious

ovine abortion worldwide, including the UK, the USA and most of northern Europe

but has not been found to be an abortion issue in sheep in Australia or New Zealand

[15] and there is little available information from the African continent, other than in a

few select countries (e.g. Tunisia, Algeria, Namibia, Zimbabwe, and Morocco [16, 17, 18,

19]. In tropical areas, epidemic-prone pathogens, such as Rift Valley fever virus

(RVFV) which poses a severe health threat to both animals and people, are also a

concern [20]. No less important are the endemic zoonoses such as *Brucella abortus*

and *B. melitensis*, and zoonotic protozoal agents such as *Toxoplasma gondii,* which

are of substantial concern in subsistence farming settings where people live closely

with their animals [3, 21] or in communities where HIV predisposes the human

population to severe infection e.g., with toxoplasmosis [22].

In Africa, inferences about the potential role and relative importance of abortigenic

agents are often made from serological studies of livestock [23, 24, 25]. This is due to a

number of factors including a lack of national-level surveillance for syndromes such

as abortions in livestock; restricted access to veterinary care for many subsistence

farmers, who are therefore unable to investigate individual abortion events; and

limitations in diagnostic laboratory capacity (e.g., little to no access to histopathology

services or molecular diagnostic tests that could help to generate data on different

aetiologies of abortion). While serological data is valuable to identify the presence of

abortigenic agents circulating within a livestock population, there is also a need to

recognise the limitations of such data and the inferences that can be made based on

it [26, 27]. Livestock seropositivity indicates only that an animal has been exposed to a

pathogen and does not necessarily indicate current disease status or causality of an

abortion event. Additionally, chronic low-level carriage of some abortigenic agents

(e.g., *C. burnetii*) in healthy animals complicates attribution of abortion. To date, in

sub-Saharan Africa, there remains an almost complete absence of published data on

livestock abortion aetiologies determined through confirmatory diagnostic testing,

resulting in a very weak evidence base on which to prioritise animal health

interventions.

In Tanzania, livestock diseases are prioritised for prevention and control largely on

the basis of the potential for transboundary spread and zoonotic impacts [28]. A recent

exercise to prioritise zoonotic diseases ranked two abortigenic agents, *Brucella* and

RVFV amongst the top six zoonoses for prevention and control [29]. While there is

clear evidence that brucellosis and Rift Valley fever (RVF) are both diseases of

public health concern in Tanzania [30, 31, 32], no data currently exist on the relative

importance of these or other zoonotic pathogens, such as *C. burnetii*, *Leptospira*

spp., *C. abortus* or *T. gondii*, as causes of livestock abortion. Nor are there any data

available on the contribution of other abortigenic pathogens of global importance,

including *N. caninum*, bluetongue virus and pestiviruses such as bovine viral

diarrhoea virus (BVDV) and border disease virus (BDV). Without robust aetiological

data, it is unlikely that current approaches to livestock disease prioritisation will

reflect the true impact of abortigenic pathogens, with the risk that scarce resources

for disease control and prevention will be targeted ineffectively and achieve sub-

optimal productivity and public health outcomes.

We sought to address knowledge gaps around the aetiology of livestock abortion in

Tanzania through a prospective cohort study conducted in three regions spanning a

range of agro-ecological settings [33]. We used a combination of field investigation and

laboratory diagnosis of abortion events in cattle, goats, and sheep to determine

aetiologies of livestock abortions for a targeted selection of known abortigenic

agents. For these selected agents, we developed a set of case definitions to attribute

abortion events to a specific aetiological agent based on molecular diagnostic

assays in the absence of comprehensive histopathology data. In the same study

area, we also carried out serological analyses of a large sample of livestock sera

collected by cross-sectional studies in the same area to determine seroprevalence to

the same range of abortigenic pathogens at a regional level.

Results

Figure 1 shows the location of sampling for the serological cross-sectional and

abortion cohort studies.

Figure 1: Distribution of villages in cross-sectional studies for serologic

baseline testing (2013-2016; red dots) and abortion cohort sampling wards

(2017-2019; yellow shading) with number of abortion events investigated,

Arusha, Kilimanjaro, and Manyara Regions, northern Tanzania.

Cross-sectional sample serology

Samples were available from 3,629 cattle, 3,344 goats, and 2,628 sheep from 567

households in 43 villages. Figure 2 summarises the cross-sectional seroprevalence

of pathogens in ruminant livestock sampled in northern Tanzania. Seroprevalence at

individual level for cross sectional livestock with Clopper Pearson 95% confidence

intervals is detailed in supplementary materials (S1, A).

Figure 2: Seroprevalence of pathogens from cross-sectional studies (2013-2015 and 2016) in cattle, goats, and sheep, northern Tanzania.

Where x = not tested and error bars represent Clopper-Pearson 95% confidence intervals

Abortion cohort study

During the prospective cohort study period, a total of 215 abortion events were investigated, 71 (33%) in cattle, 100 (46.5%) in goats, and 44 (20.5%) in sheep. The numbers of each sample type collected for abortion events are detailed in Table 1.

Table 1: Number and type of samples collected from abortion events for each livestock animal species, northern Tanzania, 2017-2019

Sample types collected	Cattle (n=71)	Goat (n=100)	Sheep (n=44)
Vaginal swabs	71 (100%)	98 (98.0%)	44 (100%)
Exterior foetal swabs	37 (52.1%)	25 (25%)	5 (11.4%)
Placental tissue	34 (47.9%)	25 (25%)	1 (2.3%)
Acute sera	71 (100%)	98 (98.0%)	43 (97.7%)

Convalescent sera	49 (69.0%)	56 (56.0%)	29 (65.9%)

<u>Abortion cohort acute serology</u>

The acute seroprevalence results by pathogen in the abortion cohort was broadly

similar to the cross-sectional survey. A table showing these results is available in the

supplementary materials (S1; A).

<u>Diagnosis of abortion aetiology</u>

We attributed 42 (19.5%) of 215 abortion events to one or more of the abortigenic

pathogens included in our diagnostic panel. Table 2 summarises the number of

abortion events with positive diagnostic test results that met the study case

definitions. Histopathology and immunohistochemistry (IHC) results are shown in

supplementary materials S1, B.

Table 2: Number of abortion cases (cattle, goats and sheep) attributed to
abortigenic agents using study case definitions, northern Tanzania 2017-2019
186

Pathogen	Case definition	Number of cases attributed by animal species				Strength of evidence of abortion cause
		Cattle n/n (%)	Goats n/n (%)	Sheep n/n (%)	Total (n)	
***Brucella* spp.**	qPCR	1/71 (1.4)	0/100 (0.0)	0/44 (0.0)	1	Presumptive
Chlamydia abortus	qPCR	0/71 (0.0)	0/100 (0.0)	0/44 (0.0)	0	Definitive
	Sero-conversion	n/a	1/56[‡] (0.0)	0/29 (0.0)		Suggestive
Coxiella burnetii	qPCR	0/71 (0.0)	5/100 (5.0)	1/44 (2.3)	6	Presumptive
***Leptospira* spp.**	qPCR	0/71 (0.0)	0/100 (0.0)	0/44 (0.0)	0	Presumptive
	Sero-conversion	0/49 (0.0)	n/a	n/a		
Neospora caninum	PCR	9/71[#] (12.7)	1/100 (1.0)	0/44 (0.0)	10[#]	Presumptive
Toxoplasma gondii	PCR	0/49 (0.0)	0/100 (0.0)	1/44 (2.3)	1	Definitive
Bovine herpes virus	Sero-conversion	2/49[#] (4.1)	n/a	n/a	2[#]	Suggestive

(BHV-1)									
Bluetongue virus (BTV)	RT-qPCR	0/71	(0.0)	0/100	(0.0)	0/44	(0.0)	0	Definitive
Pestiviruses	RT-qPCR	2/71	(2.8)	1/100	(1.0)	6/44	(13.6)	9	Definitive
Rift Valley fever virus (RVFV)	RT-qPCR	14/71	(19.7)	0/100	(0.0)	0/44	(0.0)	14	Definitive
	Sero-conversion	1/49*	(1.4)	0/56	(0.0)	0/29	(0.0)		
None	Not detected	44/71	(62.0)	92/100	(92.0)	36/44	(81.8)	173	n/a
TOTAL		**27/71**	**(38.0)**	**8/100**	**(8.0)**	**8/44**	**(18.2)**	**215**	

One case in a cow met both BHV-1 (seroconversion) and *Neospora* (PCR) case definition and was
thus included in both
* One case in a cow met both RT-qPCR and seroconversion case definitions for RVFV and was thus
included in both
‡ Seroconversion was observed in one goat with a qPCR-negative swab sample for *C. abortus* and
therefore this case did not meet the case definition.
n/a = not applicable

For some bacterial pathogens, despite evidence of infection in the dam, laboratory

data was not considered sufficient to support a diagnosis of abortion associated with

the pathogen. This was particularly true for *C. burnetii*, which was detected by qPCR

with a threshold cycle (Ct) value <40 in 16 (22.5%) of 71 cattle, 24 (24.5%) of 98

goats, and 12 (27.3%) of 44 sheep, however only six (11.5%) of these 52 animals

met the study case definition (Ct ≤ 27) for *C. burnetii* as a presumptive aetiology of

abortion. For *C. abortus*, although seroconversion between acute and convalescent

samples to *C. abortus* was detected in one (1.8%) of 56 goats with paired sera

available for testing, this animal had a negative vaginal swab qPCR result and

therefore did not meet our case definition for *C. abortus*-associated abortion.

Of the protozoal pathogens, *Neospora* DNA was diagnosed by nested conventional

polymerase chain reaction (PCR) in nine (12.7%) of 71 cattle and one (1.0%) of 98

goat abortion events. In one cattle case, a co-infection with BHV-1 was also detected

suggesting a multi-factorial aetiology. *Toxoplasma* DNA was detected by nested

conventional PCR in one (2.3%) of 44 sheep abortion events. The single *T. gondii*

PCR positive sheep sample was genotyped at 7/10 marker and was found to have

type II alleles at all of them, indicative of a type II clonal genotype infection.

Of the viral pathogens, RVF was most commonly diagnosed, with RVFV RNA

detected by reverse transcriptase qPCR (RT-qPCR) in 14 (19.7%) of 71 cattle

abortion events. Evidence of seroconversion was detected in one animal that was

positive by RT-qPCR. The other 13 RT-qPCR positive animals were positive in both

acute and convalescent sera, meeting World Organisation for Animal Health (OIE)

guidelines for definitive diagnosis of infection. Pestiviruses were also commonly

detected with two (2.8%) of 71 cattle, six (13.6%) of 44 sheep, and one (1.0%) of

100 goat abortion events positive by RT-qPCR.

<u>Pestivirus speciation results</u>

Among the nine animals that tested positive for pestiviruses by RT-qPCR, N^{Pro}

coding sequence was obtained from six samples for phylogenetic analysis (Figure

3). BVDV infection was confirmed in two animals (one cow and one sheep) with a

distinct genotype detected in each case. The cattle sample (SF1-190) was most

similar to reference sequences from BVDV genotype 1d whereas the sheep sample

(SD1-147) was most similar to reference sequences from BVDV genotype 1b.

Sequences obtained from the remaining four samples (SD1-144, SD1-145, SD1-155,

SD1-160), all obtained from sheep, were similar to each other but appeared to be

distinct from all of the reference sequences. These sequences represent a

potentially novel, previously unreported pestivirus species and formed a separate

clade on phylogenetic analysis. The sequences were assigned accession numbers

OU548650-OU548655 by the European Nucleotide Archive

(https://www.ebi.ac.uk/ena/browser/home) and, for the purposes of that submission,

this novel clade was named Pestivirus abortion/sheep/SD1/2018/Tanzania.

Figure 3: Results of evolutionary analysis of the N^{Pro} coding region from pestivirus sequences obtained from cattle and sheep sampled in the abortion cohort study, northern Tanzania.
The evolutionary history was inferred using the Maximum Likelihood method and Tamura-Nei model of nucleotide substitution [34]. The tree with the highest log likelihood is shown. The percentage of trees from 500 Bootstrap repeats in which the associated taxa clustered together is shown next to the branches. Evolutionary analyses were conducted in MEGA X [35]. Reference sequences representing nine pestivirus species were also included [36] and are labelled with virus species (A-H, K), genotype and strain name and Genbank accession numbers. Pestivirus species are also shown by bracket labels on the right of the figure. Samples analysed in this

250 study are shown in bold and labelled with unique identifiers (SD/F1-XXX), where SD
251 denotes vaginal swab taken from aborting dam; SF denotes swab taken from
252 aborted foetus. Abbreviations: APPV = atypical porcine pestivirus; BDV = border
253 disease virus; BVDV = bovine viral diarrhoea virus; CSFV = classical swine fever
254 virus.
255

Discussion

To our knowledge, this is the first multi-pathogen study of the aetiologies of livestock

abortion in Tanzania. We developed and applied an approach designed to optimise

abortion investigations in resource limited settings, complemented by traditional

cross-sectional surveillance to generate a more comprehensive understanding of the

distribution and impact of several important abortigenic pathogens. This approach

generated several new insights for Tanzania including: a) detection of an RVF

outbreak in cattle during an inter-epidemic period; b) relatively high prevalence of *C.*

burnetii infection in cattle, goats and sheep; c) the first demonstration of *N. caninum,*

T. gondii, and pestiviruses, including novel pestivirus species, associated with

livestock abortion in Tanzania, including detection in less typical hosts (for example,

N. caninum in goats); but, contrary to prior expectations, d) limited or no detection of

Brucella spp., *C. abortus,* and *Leptospira* spp. in cases of livestock abortion. Our

findings provide important information on the health and production impacts of a

range of livestock pathogens in East Africa with relevance for prioritising disease

interventions in the region.

Our study demonstrates the importance of zoonotic pathogens as causes of abortion

in ruminants in Tanzania, with implications for human as well as animal health. Two

important zoonotic pathogens were diagnosed in multiple abortion events by our

study: RVFV, a virus that can cause devastating outbreaks of human and animal

disease [37, 38], and *C. burnetii*, an important bacterial cause of human febrile illness in

Tanzania [39]. RVFV is a high priority zoonosis in East Africa and outbreaks are

frequently associated with livestock abortion storms [20]. Direct animal to human

transmission of infection can occur following inhalation of aerosolized virus particles

or during close contact with tissues and bodily fluids of infected animals, for

example, when assisting with parturition or handling aborted foetuses [40, 41]. Although

RVF usually occurs as large outbreaks occurring every 10-20 years, there is

increasing serological evidence of transmission in the period between these

epidemics [42]. This study detected the occurrence of an apparent outbreak in cattle

that was previously unreported in Tanzania, despite RVF outbreaks being detected

in Kenya, Uganda, and Rwanda at the same time [40, 43]. Although an alert of RVF was

issued in Tanzania in June 2018 at the time of this study [44], to our knowledge, no

human cases were reported. Our finding highlights the value of livestock abortion

surveillance in early detection of this epidemic-prone pathogen and also

demonstrates the value of molecular diagnostic capacity for detection of RVF

livestock cases.

Coxiella burnetii is a pathogen of concern in Europe, where outbreaks of infection in

small ruminant livestock have also led to outbreaks of human disease [45]. *C. burnetii*

has also been detected in humans and in a wide range of animal species across

Africa [46, 47], and accounts for notable proportions of severe human febrile illness in

Africa (Vanderburg et al., 2014), including in northern Tanzania [39, 48]. Our results

provide important evidence of the impact of *C. burnetii* as a cause of livestock

reproductive losses, particularly in small ruminants, revealing a dual human and

animal health burden of this zoonotic pathogen. In our abortion cohort study, we

applied a stringent case definition that exceeded the minimum guidelines stated by

the OIE for diagnosis of *C. burnetii*-associated abortion. Using this case definition,

we diagnosed six cases of *C. burnetii*-associated abortion, but it is also worth noting

that *C. burnetii* DNA was detected in a total of 52 (24.4%) abortion events indicating

that shedding of this zoonotic bacterium is even more prevalent than the number of

attributed cases would suggest. Based on our findings, we suggest that the

development of preventive and control measures against *C. burnetii* should be a

priority for future work that could have substantial benefits for human and animal

health.

Our study confirmed *T. gondii,* another zoonotic pathogen, is associated with

livestock abortion in Tanzania. Toxoplasmosis is also an important, sometimes fatal,

zoonosis in Tanzania, particularly among immunocompromised people [22]. There are

limited data on genotypes present in Africa but the Type II clonal lineage found in the

single PCR positive sheep abortion event is known to be dominant in North and East

Africa so aligns with previous studies [49]. The cross-sectional serological data

showing widespread exposure of *T. gondii* in sheep and goats are consistent with

previous published findings of high seroprevalence in people and animals, including

livestock [50, 51] and wild felid species in East Africa [52]. Given that toxoplasmosis is one

of the most important abortigenic agents of sheep worldwide, it is perhaps surprising

that we did not detect more cases of *T. gondii* as a cause of sheep abortion in this

study. However, high flock seroprevalences are typically associated with low rates of

abortion due to the development of protective immunity in ewes infected with *T.*

gondii prior to mating [53].

Other zoonotic pathogens were detected infrequently in the abortion cohort study,

with only a single case attributed to *Brucella* spp. in a cow, a single detection of

seroconversion to *C. abortus* in a goat, and no evidence of abortions associated with

Leptospira spp. in any livestock species. Although *C. abortus* has been reported as a

cause of abortion in sheep elsewhere in northern and southern Africa [54, 55, 56], our

study generated little evidence that *C. abortus* was contributing to the burden of

production losses in sheep in northern Tanzania at the time of our study. Our cross-

sectional serological findings also support the conclusion that levels of exposure in

small ruminants are very low across northern Tanzania. However, these findings do

raise concerns about the potential impact should infection be introduced to this

predominantly naïve population in the future.

Both *Brucella* spp. and *Leptospira* spp. are recognised as important zoonoses in

northern Tanzania, causing large proportions of severe human febrile illness [21, 48, 57,

58] and are known to be circulating in livestock in these areas of Tanzania.

Seroprevalence of *Brucella* spp. was low in all three livestock species (Fig. 2), which

is consistent with the low frequency of detection in abortion cases. In contrast,

although *Leptospira* was not detected by qPCR in any abortion event investigated,

evidence of exposure to *Leptospira borgpetersenii* serovar Hardjo, and *Leptospira*

interrogans serovar Hardjo (subsequently referred to as *Leptospira* Hardjo serovars)

was relatively high in cattle in the cross-sectional study. Several possible

explanations may explain this discrepancy. Similar to *T. gondii,* prior exposure to

Leptospira serovars may lead to protective immunity prior to animals becoming

pregnant. Additionally, livestock abortions associated with *Leptospira* are often seen

in outbreaks or 'abortion storms' following either a new introduction into a herd or a

seasonal outbreak where flooding or wet weather conditions support transmission of

novel serovars [59]. The timespan of our project meant that we may have missed

intermittent or infrequent abortion storms associated with *Leptospira* in livestock.

However, even accounting for these limitations, our study suggests that in these

agricultural settings, the proportion of abortions attributable to *Brucella* spp. and

Leptospira spp. may be lower than is widely perceived. Alongside other constraints

to the uptake and use of livestock brucellosis vaccines these data raise questions

about the likely effectiveness and sustainability of livestock vaccination-based control

strategies in this context if based on a farmer pays model only. To effectively

mitigate both the animal and human impacts of brucellosis a diversity of strategies

may be needed to achieve the vaccination coverage required to reduce animal

production impacts and also to reduce the substantial human health burden from this

disease.

In addition to zoonotic pathogens, our study provides also provides valuable data on

several production-limiting infections of ruminants that are often over-looked in sub-

Saharan Africa. In particular, while *N. caninum* and pestiviruses are considered

important causes of abortions and productivity loss in Europe and the USA, there are

few data available describing their impact in African livestock. Our results provide the

first direct evidence of livestock abortion associated with *N. caninum* infection in

Tanzania, a conclusion which is supported by a positive relationship between *N*

caninum seroprevalence and in-herd abortion rates in cattle detected by earlier work

in our study area [60]. In this study, *N. caninum* was predominately diagnosed among

cattle abortions, but notably also in one goat abortion event. Serological evidence of

low-level exposure to *N. caninum* in goats (1.7%) and sheep (1.5%) was also

detected across the region. Although abortion and foetal pathology associated with

N. caninum has been described in goats and sheep [61, 62, 63], natural infection of *N.*

caninum in small ruminants is not widely reported. Given the importance of small

ruminants within livestock economies and for food security in Africa [64, 65], further

investigation of the patterns and impact of *N. caninum* infection in small ruminants

are warranted.

Pestiviruses were the most commonly diagnosed pathogen associated with abortion

in sheep in this study, but pestiviruses are rarely discussed as priority production-

limiting pathogens in livestock in Africa. Analysis of the six pestivirus sequences

obtained in this study identified at least three different viral lineages associated with

abortion in livestock in northern Tanzania including a novel lineage in sheep. Based

on our analysis of the N^{Pro} coding region, this novel lineage appears to be distinct

from other pestiviruses and is likely to be a new, previously unreported pestivirus

species. Notably, several of the most similar pestivirus species were detected in

sympatric wildlife species including giraffe in Kenya (Genbank accession numbers

U80907 and AY163647). As wildlife interactions with livestock are common in many

areas of Tanzania, we hypothesise that spill-over from wildlife may also be involved

in the epidemiology of this novel pestivirus lineage, which appears to be a

contributing cause of ovine reproductive loss in northern Tanzania.

Serological data from the cross-sectional study confirmed widespread exposure to

BHV-1 in the study area (Fig. 2), and previous studies in this area have

demonstrated an association between seropositivity and cattle abortion [66]. However,

the evidence for BHV-1 as a cause of abortion in this study is relatively weak. As

latent BHV-1 infection may lead to false positive results by molecular detection [67], we

relied on serology to diagnose infection. Two cattle showed evidence of BHV-1

seroconversion suggesting recent infection which may be implicated in the aetiology

of abortion in these cases but further data are needed to evaluate the impact of

BHV-1 pathogen on livestock reproduction in Tanzania and our findings highlight the

challenges of attribution in settings where endemic pathogens may co-exist in cattle

herds [68].

BTV, which causes production losses in livestock in Europe [69, 70], was not detected

by RT-qPCR in any samples from livestock abortion events, despite high BTV

seroprevalences detected in all species in the cross-sectional study population.

Serological evidence of infection is widespread East Africa [71, 72, 73] and several

studies have also demonstrated viraemia using RT-qPCR assays, reporting infection

prevalence as high as 88.9% in cattle in Kenya [73] and 56.0% in goats in Uganda [72].

Collectively, these studies suggest that BTV infection is endemic in livestock in East

Africa. However, research indicates that African breeds of livestock may be relatively

resistant to clinical disease associated with BTV. Studies from southern Africa have

indicated that indigenous African breeds of livestock are less susceptible to clinical

disease from BTV infection than European breeds [74]. Furthermore, transplacental

foetal infection is only associated with certain BTV strains, especially attenuated

vaccine strains [75]. Our findings supported by these trends in BTV epidemiology and

pathogenesis suggest that BTV is not a major cause of livestock abortions in

northern Tanzania and underscores the need for caution in making inferences about

livestock diseases and causes of abortion in particular based solely on

seroprevalence data.

The approach described in our abortion cohort study enabled us to attribute

approximately 20% of reported abortion events to a specific aetiology. Our success

rate was not dissimilar to the quoted rate of aetiological abortion diagnosis in high

income settings (35%) [5, 7, 8] despite working in a challenging and resource-limited

environment. Our approach was specifically designed to optimise abortion

investigations in a resource-limited setting, where veterinary surveillance may be

limited. Our focus on the use of molecular diagnostics to detect a range of selected

pathogens proved to be valuable, particularly for high priority pathogens such as

RVFV. While molecular diagnostic capability was established for several known

abortigenic agents, it was not possible for our diagnostic panel to be fully

comprehensive of all possible aetiologies of abortion and did not include agents such

as *Campylobacter spp.*, *Listeria spp.*, and *Salmonella enterica* serovars that are

known to be important causes of livestock abortion worldwide. Non-infectious causes

were also not considered for investigation and these two factors may account for

some of the undiagnosed cases.

Although histopathology is conventionally considered the most robust approach for

attributing causality in livestock abortion, histopathology was of limited value in this

study due to the limited availability and poor quality of tissues samples obtained

through field investigations that were often conducted in remote settings. Additional

challenges to inference of causality were encountered due to the nature of our study

site. In areas such as rural Tanzania, where remote locations, high ambient

temperatures and scavenging animals mean that the type and quality of diagnostic

samples is often sub-optimal, inference about the cause can be particularly

challenging. For example, placental tissues were not available for sampling from

many small ruminants, and even where placental tissues were available for

collection, most tissue samples showed evidence of significant autolysis, which

precluded a definitive diagnosis in some cases. However, despite these sampling

limitations, molecular diagnostics provided a wealth of information regarding the

presence of abortigenic agents in swab and tissue samples, and therefore we

recommend that molecular diagnostics should be included as a core element of

abortion surveillance studies to infer aetiology in resource-limited settings.

In considering appropriate criteria for attributing the aetiology of abortion events

reported to this study, we became aware of variations in case definitions and the

limitations of the evidence base underpinning attribution criteria for livestock

abortions. While methods for assigning attribution in situations of imperfect

diagnostic testing are well-developed and widely adopted in aetiological studies of

human diseases [76], these questions are rarely addressed in veterinary

investigations. Attribution was particularly problematic for *C. burnetii*, as the

pathogen is frequently detected in vaginal fluids at parturition in clinically healthy

animals [77] and environmental contamination of placental material or foetal surfaces

may also lead to false positives in high risk environments [78]. In this study, we

developed a case definition for *C. burnetii* based on OIE guidelines [79, 80] but also

driven by population-level data generated by this study. Further development of this

case definition would be beneficial to understand how clinical history, gross

pathology (where available), measured bacterial load and other adjunct diagnostics

could be used to more confidently attribute abortion cases both within and outside

the Tanzanian context.

Substantial differences in reporting rates between wards further limits the

representativeness of our sample. Important differences in livestock seroprevalence

between production systems have been observed in northern Tanzania [81] as well as

in other areas of East Africa [82]. Spatial heterogeneity in livestock exposure risk, with

clustering in seropositivity at the village-level, have also been described for

pathogens such as *Brucella* [83]. It is therefore possible that sample collection in more

administrative areas, with improved representation of pastoral and agropastoral

livestock keepers, would reveal different frequencies of pathogen detection than

described here. Our data should therefore be considered as only indicative of the

relative importance of different pathogens as causes of livestock abortion in northern

Tanzania as a whole.

A key insight from our study is that, apart from RVF, none of the abortigenic

pathogens identified by this study have been considered important pathogens in

Tanzania, or prioritised for interventions. Gaps remain in our understanding of the

causes of livestock abortion in Tanzania. However, even with testing a limited range

of pathogens, our study has been able to generate data that adds substantially to the

existing evidence base for the development of interventions to reduce the burden of

production losses associated with livestock abortions. Our study further highlights

the value and importance of establishing laboratory-based molecular diagnostics and

developing clear case definitions and diagnostic criteria for use in resource-limited

settings. With laboratory capacity and technical capability expanding rapidly in Africa,

there is enormous potential and an urgent need for the veterinary sector to shift

towards more evidence-based policy and decision-making, with the potential for this

transition to lead to substantial improvements in animal and human health.

Methods

Research ethics

Ethics approval for this research was granted by Kilimanjaro Christian Medical

Centre (KCMC) Ethics Committee (No. 535 & No. 832); National Institute of Medical

Research (NIMR), Tanzania (NIMR/HQ/R.8a/Vol.IX/1522 &

NIMR/HQ/R.8a/Vol.IX/2028); Research Ethics Coordinating Committee and the

Institutional Review Board for Clinical Investigations of Duke University Health

System in the United States (Pro00037356), University of Otago Ethics Committee

(H15/069 & H17/069); and University of Glasgow College of Medical Veterinary and

Life Sciences Ethics Committee (200140152 & 200170006). Sample shipments

between the UK and Tanzania were performed in accordance with the Nagoya

Protocol on Access to Genetic Resources under Material Transfer Agreements

between KCMC and University of Glasgow, and between KCMC and Moredun

Research Institute (MRI) with authorisation for export and import, respectively, from

the Tanzanian Ministry of Livestock and Fisheries (permit number VIC/ZIS/8582),

and The Scottish Government's Agriculture, Food and Rural Communities

Directorate (TARP(S) 2019/07).

Cross-sectional serological study

Two cross-sectional livestock serological studies were performed from 2013 through

2016, to provide a baseline of exposure to a range of abortigenic pathogens.

Livestock were selected and sampled as previously described [60, 84] from six districts

in Arusha Region, three districts in Kilimanjaro Region, and four districts in Manyara

Region. Serum was separated from coagulated whole blood samples by

centrifugation (≤1,300 x g for 10 mins at room temperature) and aliquoted to create

replicate sets. Samples were transported to the Zoonoses Laboratory at Kilimanjaro

Clinical Research Institute (KCRI), Moshi, Tanzania, for archiving at -80°C prior to

testing. Cattle, goat, and sheep sera were tested for exposure to *Brucella* spp., *C.*

burnetii, and RVFV using commercial ELISA, as per manufacturer's instructions

(Table 3). Cattle sera were additionally tested for exposure to *Leptospira* Hardjo

serovars, BHV-1, and BVDV (Table 3). Absorbance was measured using a

MultiSkan FC plate reader (Fisher Scientific, Leicestershire, UK). For each sample,

one of the duplicate aliquots was heat treated at 56°C for two hours and archived at -

80°C until it was shipped, on dry ice, to MRI or University of Glasgow, UK, for

testing. All or a random selection of available samples were tested by indirect ELISA

for *N. caninum* and *T. gondii*, and *C. abortus* (goats and sheep only) as described

previously (Helmick et al., 2002, Marques et al., 2012, Wilson et al., 2009). A random

selection of cross-sectional sera was also tested for exposure BTV (Table 3).

Clopper-Pearson 95% confidence intervals were calculated using the *binom* package

[85] in R statistical software (version 3.6.0).

Table 3: Cross-sectional and abortion cohort serological tests (ELISA) for

abortigenic pathogens used for various livestock species, northern Tanzania,

2013-2019.

Target pathogen	Livestock species tested			ELISA kit name (manufacturer or source)
	Cattle	Goat	Sheep	

Brucella spp.	✓	✓	✓	*Brucella* COMPELISA (Animal and Plant Health Agency, UK)
Chlamydia abortus	NA	✓	✓	In-house MRI ELISA [86]
Coxiella burnetii	✓	✓	✓	PrioCHECK Ruminant Q Fever Ab Plate § (Thermofisher®)
Leptospira Hardjo	✓	NA	NA	Bovine Leptospira Hardjo 5 Plate Solid ELISA kit (Linnodee®)
Neospora caninum	✓	✓	✓	In-house MRI ELISA [60] **OR** *Neospora caninum* Indirect ELISA (ID Screen®)
Toxoplasma gondii	✓	✓	✓	In-house MRI ELISA [87] **OR** Toxoplasmosis Indirect Multi-species ELISA (ID Screen®)
Bluetongue virus (BTV)	✓	✓	✓	Bluetongue competition kit (ID Screen®)
Bovine herpesvirus (BHV-1)	✓	NA	NA	IBR Individual Ab Test (IDEXX®)
Pestiviruses (BVDV/ BDV)	✓	NA	NA	BVDV p80 Ab ELISA (IDEXX®)
Rift Valley fever virus (RVFV)	✓	✓	✓	Rift Valley Fever Competition Multispecies ELISA kit (ID Screen®) **OR** In-house nucleocapsid-based ELISA pre-screen of goat and sheep CS sera (University of Glasgow) (Nyarobi, 2020)

§: Formerly known as LSIVet Ruminant Q fever Serum/Milk

KCRI = Kilimanjaro Clinical Research Institute, Moshi, Tanzania; MRI = Moredun Research
Institute, UK
NA = not applicable, test not done. For *Leptospira,* BHV-1 and BVDV, serological testing of
goat and sheep sera was not performed because the ELISA kits were not validated for small
ruminant sera.
554

<u>Abortion cohort study</u>

A prospective abortion cohort study was undertaken from October 2017 through

September 2019 in 13 wards in Arusha, Kilimanjaro, and Manyara Regions of

northern Tanzania (Fig. 1). Study wards were selected from those included in the

cross-sectional exposure studies. Two wards were selected purposively from among

those included in these cross-sectional studies because of existing relationships with

the livestock-keeping community that were expected to promote participation. The

remaining eleven wards were selected randomly from the wards targeted in the

cross-sectional studies.

Livestock field officers (LFOs), animal health professionals working for the

Tanzanian Ministry of Livestock and Fisheries, were recruited to participate in the

study with one LFO responsible for a single ward. At the start of the study the LFOs

attended a training course on causes of livestock abortion, as well as the practical

investigation and management of abortion events. Training included the safe

collection, storage, and transport of biological samples from the aborting dam,

foetus, and placental material, the use of personal protective equipment for the safe

handling and disposal of aborted material and methods for data collection. Following

the training course, LFOs were asked to disseminate information about the study to

the community living within their respective wards, requesting livestock owners to

report any instances of livestock abortion to them so that an investigation could be

carried out. Following receipt of a report, the LFO was asked to pass on the

information to the research coordinator, a veterinarian, for investigation. For the

purposes of this study, an abortion event was defined as: farmer reported visual

evidence of premature foetal loss or stillbirth in cattle, goats, or sheep. The abortion

was considered eligible for inclusion in this study if the research field team or LFO

could attend within 72 hours of the event occurring. Two investigation options were

available. Where LFOs carried out the investigation, vaginal swabs and 10 ml blood

samples were collected in red-top vacutainer tubes (Becton, Dickinson and

Company, New Jersey, USA) from the dam(s) that had aborted, as identified by the

livestock keeper. Where the veterinarian-led research team was able to join the

LFO's within 72 hours, additional samples were collected, including swabs from the

surface of the foetus exterior and placental (cotyledon) tissues, when available.

Duplicate swab samples (breakpoint shaft, viscose tip; Technical Service

Consultants Ltd, Lancashire, UK) and duplicate tissue samples were stored in 1 ml

DNA/RNA shield (Zymo Research, California, USA) to inactivate infectious agents

and preserve nucleic acid integrity prior to molecular testing. Due to practical

limitations, samples comprehensive sets of tissues including foetal fluids for bacterial

culture were not collected in this study. Tissue samples were also stored in duplicate

in 10% buffered formal saline to preserve tissue structure for histopathology and

IHC. Samples were transported to the Zoonoses Laboratory at KCRI, for archiving at

-80°C prior to testing against a panel of abortigenic agents.

Abortion cohort serological testing

Sera from dams that aborted were tested using the range of host and pathogen

specific assays described in Table 3. For each sample, one of these duplicate tubes

was heat treated at 56°C for two hours prior to export to the UK for testing. The

remaining tube was retained for testing at KCRI and stored at -80°C. Households

were re-visited four to six weeks later to collect a convalescent blood sample from

the same dam(s). Seroconversion was defined as a negative acute serum with a

positive convalescent serum.

<u>DNA and RNA extraction</u>

RNA was extracted using a RNeasy® Mini kit (QIAGEN, Hilden, Germany) following

manufacturer's instructions. For RNA extraction from the placental tissues, a sterile

scalpel was used to add approximately 25 mg of tissue to 600 µl RLT buffer in a

sterile bead beating tube containing ceramic beads (CK28, Precellys, Bertin

Instruments, France). Bead beating was performed for one minute at 3,450 rpm

using a MiniBeadBeater-16 (BioSpec Products, Inc., Oklahoma, USA). The liquid

was aliquoted into a microcentrifuge tube with 590 µl nuclease-free water and 10 µl

Proteinase K (QIAGEN, Hilden, Germany). For RNA extraction from swabs, 200 µl

sample liquid was added to 600 µl RLT buffer, and 10 µl Proteinase K. All tubes

were then incubated at 55°C for 10 min. Once cooled to room temperature, tubes

were centrifuged at 10,000 x g for three minutes. The supernatant was transferred to

a new microcentrifuge tube where half the volume of ethanol (96-99%) was added to

the tube (e.g., 450 µl ethanol to 900 µl supernatant) and mixed by pipetting up and

down gently. RNA was extracted from 700 µl of the lysate/ethanol mix using a final

elution volume of 50 µl.

DNA was extracted using a DNeasy® blood & tissue kit (QIAGEN), following

manufacturer's instructions. For DNA extraction from placental tissues, a sterile

scalpel was used to add approximately 25 mg of placenta into a microcentrifuge tube

with 180 µl ATL buffer and 20 µl Proteinase K. The tubes were incubated at 56°C,

vortexing occasionally, until the tissue was lysed, after which, 200 µl AL buffer was

added, and the tube was vortexed briefly. For DNA extraction from swabs, 200 µl

sample liquid was added to 200 µl AL buffer with 20 µl Proteinase K and vortexed.

All tubes were incubated at 56°C for 10 min. When the buffer/sample mix had cooled

to room temperature, 200 µl ethanol (96-99%) was added, and the tube was

vortexed. DNA was extracted from the entire volume of the lysate/ethanol mix. Final

elution volumes for swabs and placental tissue samples were 100 µl and 200 µl,

respectively.

All batches of nucleic acid extractions included a negative extraction control where

sample was replaced by nuclease-free water and the run alongside the nucleic acid

extraction process.

<u>Molecular testing</u>

The molecular targets for all pathogens are listed in Table 4. Primer and probe

sequences and cycling conditions for each PCR, qPCR, and RT-qPCR assay, are

detailed in supplementary materials (S1; Table C). DNA or RNA extracts from

positive controls listed in Table 4 were used for individual PCR, qPCR, and RT-

qPCR assays. All available vaginal swab, foetal swab, and placental tissue samples

for all three livestock species were for all pathogens included in this study. Samples

were tested in duplicate except for pathogens tested using commercial kits (BTV and

pestiviruses) where single samples were tested. Nuclease-free water and extraction

controls were included in each PCR, qPCR, and RT-PCR run as negative controls.

Samples were considered positive only if negative extraction controls and negative

controls showed no evidence of amplification and the internal and external positive

controls also amplified. Where single wells amplified for samples tested in duplicate,

the assay was repeated. Unless specified, testing was performed at KCRI.

Table 4: Abortion cohort study molecular assays (PCR, qPCR and RT-qPCR)

for cattle, goat and sheep samples for each target pathogen, northern

Tanzania, 2017-2019.

Target pathogen	PCR/qPCR/RT-PCR testing		
	Target (source)	Positive control strain (source)	Commercial kit (manufacturer)
Brucella spp.	*IS711* [88] **AND** *bcsp31* [89]	*Brucella abortus* strain 16M (Animal and Plant Health Agency, UK)	n/a
Chlamydia abortus	*Omp1* [90]	*C. abortus* strain S26/3 (Moredun Research Institute (MRI), UK)	n/a
Coxiella burnetii	*IS1111a* [45]	*C. burnetii* Nine-mile strain RSA 493 (Wageningen Bioveterinary Research, Wageningen, Netherlands)	n/a

Pathogenic *Leptospira* spp.	*lipL*32 [91]	*Leptospira interrogans* serovar Copenhagenii Strain Wijnberg (OIE Reference Laboratory for Leptospirosis, Amsterdam, Netherlands)	n/a
Neospora caninum	*ITS1* [92]	DNA extracted from tachyzoites of NC1 strain (MRI, UK)	n/a
Toxoplasma gondii	*ITS1* [93]	DNA extracted from tachyzoites of RH strain (MRI, UK)	n/a
Bluetongue virus (BTV)	n/a	RNA provided in kit	Virotype BTV pan/8 RT-PCR Kit (Indical® Bioscience GmbH)
Pestiviruses (BVDV/ BDV)	n/a	RNA provided in kit	Virotype BVDV RT-PCR Kit (Indical® Biosciences GmbH)
Rift Valley fever virus (RVFV)	RV [94]	RVF strain MP12 (MRC-University of Glasgow Centre for Virus Research, Glasgow, UK)	

Previously published qPCR assays designed and validated for the diagnosis of

infection with the respective pathogens were used to detect the presence of *Brucella*

spp. (two gene targets [88, 89]), *C. burnetii* [45], *C. abortus* [90], and pathogenic *Leptospira*

spp. [91]. These qPCRs were performed in single well reactions using a QuantiNova®

Probe kit (QIAGEN) with 5 μl of DNA template (or 2.5 μl per target for *Brucella* spp.).

Nested conventional PCR assays for the presence of *Neospora* spp. and

Toxoplasma gondii were performed targeting the internal transcribed spacer (ITS1)

region between the 18S and 5.8S rRNA genes [95]. The primary (1°) nested PCR was

performed using 2 μl of DNA template. The secondary (2°) nested PCR was

performed using with 2 μl of a 1:5 diluted primary PCR product. Both 1° and 2°

assays were performed with 2x PCR master mix (Promega Corporation, Wisconsin,

USA). The final PCR product from the 2° nested PCR was checked by

electrophoresis in a 1.5% agarose gel. Samples were considered positive if the size

of bands visible under UV light were congruous with those expected of the *Neospora*

(249 bp) and *Toxoplasma* (227 bp) targets.

The presence of BTV and pestivirus RNA in samples was determined by RT-qPCR

using commercial kit-based assays validated for the detection of the respective

pathogens (Table 4). Single samples were tested, as per kit instructions, using 5 μl

of RNA template. Finally, the presence of RVFV RNA was assessed using the

QuantiNova® RT-PCR probe kit (QIAGEN). Samples were tested in duplicate using

5 μl of RNA template per well.

<u>Pestivirus speciation</u>

As commercial kit-based RT-qPCR assays for pestiviruses do not distinguish

between the different pestivirus species, genotyping was performed at MRI to

determine the infecting pestivirus species. The RNA extracts that were positive in

pan-pestivirus RT-PCR screening assays were subjected to RT-PCR and then

nested PCR using four primer pairs developed for genotyping of livestock

pestiviruses: 324-326 (5'UTR); BD1-BD3/BD4 (N^{pro}); BD1-BD2 (N^{pro}-C); and 17f-

1400r (UTR-Erns) to determine the infecting virus type [96, 97, 98]. RT-PCR products of

the expected sizes were purified from agarose gels and submitted for sequencing

using the relevant PCR primers (Eurofins-MWG). Sequence traces for each sample

were assembled using SeqMan Pro (DNASTAR Lasergene) and consensus

sequences that excluded the terminal primer sites were generated. The 504 nt N^{pro}

coding sequences were aligned with a set of N^{Pro} reference sequences that included

representatives of pestivirus species A-H plus K, as described previously [36].

Alignment was performed using MAFFT (https://mafft.cbrc.jp/alignment/). Maximum

likelihood (ML) phylogenetic analysis was performed using MEGA version X. Model

selection was used to identify the best evolutionary model for the dataset. The ML

analysis was bootstrapped 500 times to estimate the reliability of the resulting

phylogeny.

Additional pathogen genotyping

T. gondii genotyping was also performed to determine the infecting sub-type. The

methods of this genotyping work are detailed in the supplementary materials S1; D.

Pathology and immunohistochemistry (IHC)

Histopathology was performed on placental tissue samples where available. For

samples that tested positive for *C. burnetii* by qPCR, *C. burnetii* IHC was also

performed. The methods for the histopathology and IHC are given in supplementary

materials S1; B.

Diagnosis of abortion aetiology

For the purposes of our study, we determined a set of case definitions that we used

to attribute livestock abortion events to a specific aetiology. These case definitions

are shown in Table 5. Case definitions were based on OIE guidelines for the

diagnosis of each pathogen. Due to the challenges in systematically collecting tissue

samples for histopathology from abortion cases (e.g. lack of available tissue in many

cases), we designed these diagnostic criteria (Table 5) to be independent of

histopathological examination but used evidence from histopathological examination

to support our diagnosis where samples existed to support this diagnostic approach.

For *N. caninum*, for which no OIE diagnostic guidelines exist, the case definition for

was based on criteria used by MRI, UK [99], a specialist diagnostic veterinary

laboratory. For *C. burnetii,* OIE guidelines indicate that definitive diagnosis of

infectious abortions resulting from *C. burnetii* infection can only be robustly achieved

at the flock or herd level, but also that bacterial load is predictive of causality in

individual animals [79, 80]. Therefore, to determine the *C. burnetii* case definitions, we

used a two-step approach. First, we used the minimum bacterial load associated

with abortions as stated by the OIE ($> 10^4$ bacteria per swab or per gram of tissue,

equivalent to Ct values of 37.5 and 42.7 respectively on our PCR platform).

Secondly, using a histogram of the Ct results at the study population-level, we

identified a bimodal distribution in the results, suggesting two distinct biological

processes within our sampled population (see supplementary material S1; E). We

used this frequency distribution, supported by histopathology and IHC results to

select a conservative threshold value using a cut-off Ct value of ≤ 27 for attributing

abortion events to *C. burnetii* in this study.

Confidence in this study's diagnostic case definition was determined to be either

'definitive', 'presumptive' or 'suggestive', depending on how closely our diagnostic

approach aligned with the OIE guidelines for diagnosis of abortions caused by each

pathogen. Where our study diagnostic approach met OIE definitions for a definitive

diagnosis, we classified our diagnosis as 'definitive' and had the highest confidence

in our aetiologic diagnosis. Where our study diagnostic approach met OIE definitions

for a presumptive diagnosis or for *N. caninum* where no OIE guidelines exist, we

classified our diagnosis as 'presumptive'. Where our study showed evidence of

recent dam infection by serology, but did not detect the presence of the pathogen

directly in any samples (e.g., for BHV-1 in cattle, or *C. abortus* in small ruminants),

we classified the diagnosis as 'suggestive' indicating a lower level of confidence in

these aetiologic diagnoses.

Table 5: Diagnostic criteria used by this study to attribute the aetiology of abortions in cattle, sheep, and goats reported to the abortion surveillance platform. Diagnostic case definitions apply to all three ruminant species unless otherwise specified.

Pathogen	Were OIE diagnostic guidelines available?	Summary of OIE diagnostic criteria (presuming recent history of abortion in all cases)	Diagnostic case definition applied in this study	Confidence in diagnostic case definition
Brucella spp.	Yes - for aetiological diagnosis of abortion [100]	*Definitive diagnosis (dx): Isolation of* Brucella *from abortion material, udder secretions or from tissues removed at post-mortem* *Presumptive diagnosis (dx): Evidence of* Brucella *by the demonstration of* Brucella-*like organisms in abortion material or vaginal discharge using modified acid-fast staining, if supported by serological tests. Polymerase chain reaction* (*PCR*) *methods are additional means for detection of the presence of Brucella DNA in a sample.*	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR against both *IS711* (Ct <35) and bcsp31 (Ct <40) targets **AND** Evidence of seropositivity in acute and/or convalescent serum samples from dam	Presumptive
Chlamydia abortus	Yes - for aetiological diagnosis of abortion in small ruminants [101]	*Definitive dx: Evidence of purulent to necrotising placentitis with vasculitis by histopathology, and demonstration of large number of the organism in placentae or vaginal excretions by PCR or antigen tests or stained smears.* *Presumptive dx: In the absence of samples for direct pathogen detection, a rise in*	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR (Ct < 40)	Definitive
			OR In the absence of a vaginal swab or placental sample, evidence of seroconversion	Suggestive

			between acute and convalescent serum samples from dam (goats and sheep only)	
Coxiella burnetii	Yes - for aetiological diagnosis of abortion at the herd or flock level only [80]	<u>*Presumptive dx (individual animal):* *Positive identification of high levels of C. burnetii >10⁴ per vaginal swab or per gram of placental tissue [79, 102]] by PCR in placenta or vaginal discharges.*</u>	Placental tissue, vaginal swab or foetal swab positive by qPCR by *IS1111a* qPCR (Ct ≤ 27)*	Presumptive
Pathogenic *Leptospira* spp.	Yes - for diagnosis of dam infection only [103]	<u>*Definitive dx:* Isolation *of* Leptospira *(recommended) or demonstration by PCR (suitable method) from blood or milk*</u> <u>*Presumptive dx:* demonstration of agent (isolation or PCR) in the genital tract or vaginal fluids in association with compatible clinical signs and evidence of serological exposure</u>	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR (Ct < 40) **OR** Evidence of seroconversion between acute and convalescent serum samples from dam (cattle only) [59]	Presumptive
Neospora caninum	No	N/A	Vaginal swab, foetal swab and/or placental tissue samples positive by PCR [104]	Presumptive
Toxoplasma gondii	Yes - for aetiological diagnosis of abortion [105]	<u>*Definitive dx:* *Demonstration of* Toxoplasma *in the placenta of an aborted foetus by either histopathology or PCR*</u>	Vaginal swab, foetal swab and/or placental tissue samples positive by PCR [53]	Definitive

Bovine herpes virus-1 (BHV-1)	Yes - for aetiological diagnosis of abortion [106]	*Definitive dx: Demonstration of BHV-1 by PCR or virus isolation from abortion material.* *Presumptive dx: Demonstration of BHV-1 by PCR or virus isolation from swabs from the genital tract or evidence of seroconversion between acute and convalescent samples confirming acute infection demonstrated by ELISA or virus neutralisation.*	Evidence of seroconversion between acute and convalescent serum samples from dam (cattle only)	Suggestive
Bluetongue virus (BTV)	Yes - for aetiological diagnosis of abortion [107, 108]	*Definitive dx: Detection of pathogen by RT-PCR or classical virus isolation in blood or tissue samples including from aborted or stillborn animals.*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40)	Definitive
Pestiviruses (BVDV or BDV)	Yes - for aetiological diagnosis of abortion [109]	*Definitive dx: Detection of pathogen by RT-PCR, virus isolation or immunohistochemistry in foetal tissues or fluids.*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40)	Definitive
Rift Valley fever virus (RVFV)	Yes - for aetiological diagnosis of abortion [110]	*Definitive dx: Requires a combination of at least two positive test results from two diagnostic test methods including detection of pathogen by virus isolation or RT-PCR performed on blood (from dam), tissue from aborted foetuses or other abortion products; positive detection of antibodies (serum); or*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40) **AND** Detection of RVFV IgG in acute and/or convalescent serum from dam by RVFV	Definitive

		positive for IgM or IgG ELISA with demonstration of rising titres between paired serum samples collected 2-4 weeks apart.	IgG ELISA [110]	

756   *See supplementary materials S1; E for further details

758 **Supplementary materials:**
759 **A: Table A**: Cross-sectional and abortion cohort acute serology
760
761 **B:** Histopathology and immunohistochemistry for abortion cohort study including **Table S1B**
762 (Summary of results from histopathologic examination of placental samples collected from
763 aborting ruminant livestock as part of the abortion cohort survey, northern Tanzania) and
764 interpretation and conclusions.
765
766 **C: Table C**: Target pathogen serologic (ELISA) and molecular (PCR/RT-PCR) assay
767 information, including primer/probe sequences and cycling conditions for samples from
768 various livestock species.
769
770 **D:** Genetic characterisation of *T. gondii* isolated from an abortion case
771
772 **E:** Determining qPCR cut-off values for diagnosing *Coxiella burnetii* as the cause of abortion
773 in ruminant livestock in Tanzania.

REFERENCES

1. De Vries, A. Economic Value of Pregnancy in Dairy Cattle. *J. Dairy Sci.* **89**, 3876-3885 (2006).

2. Givens, M.D. & Marley, M.S.D. Infectious causes of embryonic and fetal mortality. *Theriogenol.* **70**, 270-285 (2008).

3. FAO. World Livestock 2011 – Livestock in food security. Rome: FAO; 2011.

4. FAO. World Livestock: Transforming the livestock sector through the Sustainable Development Goals. Rome: Food and Agriculture Organization of the United Nations; 2018. Report No.: 222 pp. Licence: CC BY-NC-SA 3.0 IGO.

5. Cabell, E. Bovine abortion: aetiology and investigations. **29**, 455-463 (2007).

6. Campero, C.M., Moore, D.P., Odeón, A.C., Cipolla, A.L. & Odriozola, E. Aetiology of bovine abortion in Argentina. *Vet. Res. Comm.* **27**, 359-369 (2003).

7. Wolf-Jäckel, G.A. *et al.* Diagnostic studies of abortion in Danish cattle 2015–2017. *Acta Vet. Scand.* **62**, 1 (2020).

8. Kirkbride, C.A. Bacterial agents detected in a 10-year study of bovine abortions and stillbirths. *J. Vet. Diagn. Invest.* **5**, 64-68 (1993).

9. Buxton, D. & Henderson, D. Infectious abortion in sheep. *In Pract.* **21**, 360-368 (1999).

10. Hazlett, M.J. *et al.* A prospective study of sheep and goat abortion using real-time polymerase chain reaction and cut point estimation shows *Coxiella burnetii* and *Chlamydophila abortus* infection concurrently with other major pathogens. *J. Vet. Diagn. Invest.* **25**, 359-368 (2013).

11. van den Brom, R. *et al.* Abortion in small ruminants in the Netherlands between 2006 and 2011. *Tijdschrift Diergeneesk* **137**, 450-457 (2012).

12. Oliveira, R.D. *et al.* Domestic sheep show average *Coxiella burnetii* seropositivity generations after a sheep-associated human Q fever outbreak and lack detectable shedding by placental, vaginal, and fecal routes. *PLoS One* **12**, e0188054 (2017).

13. Loftis, A.D., Reeves, W.K., Miller, M.M. & Massung, R.F. *Coxiella burnetii*, the agent of Q fever, in domestic sheep flocks from Wyoming, United States. *Vector Borne Zoonotic Dis.* **12**, 189-191 (2012).

14. Velasova, M. *et al.* Herd-level prevalence of selected endemic infectious diseases of dairy cows in Great Britain. *J. Dairy Sci.* **100**, 9215-9233 (2017).

15. Essig, A. & Longbottom, D. *Chlamydia abortus*: New aspects of infectious
abortion in sheep and potential risk for pregnant women. *Curr. Clin. Microbiol.*
Rep. **2**, 22-34 (2015).

16. Benkirane, A., Jabli, N. & Rodolakis, A. [Frequency of abortion and
seroprevalence of the principal diseases causing ovine infectious abortion in
the area of Rabat (Morocco)]. *Ann. Vet. Res.* **21**, 267-273 (1990).

17. Hireche, S., Ababneh, M.M., Bouaziz, O. & Boussena, S. Seroprevalence and
molecular characterization of *Chlamydia abortus* in frozen fetal and placental
tissues of aborting ewes in northeastern Algeria. *Trop. Anim. Health Prod.* **48**,
255-262 (2016).

18. Rekiki, A. *et al.* Isolation and characterisation of local strains of
Chlamydophila abortus (*Chlamydia psittaci* serotype 1) from Tunisia. *Vet.*
Res. **33**, 215-222 (2002).

19. Seth-Smith, H.M.B. *et al.* European *Chlamydia abortus* livestock isolate
genomes reveal unusual stability and limited diversity, reflected in
geographical signatures. *BMC Genomics* **18**, 344 (2017).

20. Budasha, N.H., Gonzalez, J.P., Sebhatu, T.T. & Arnold, E. Rift Valley fever
seroprevalence and abortion frequency among livestock of Kisoro district,
South Western Uganda (2016): a prerequisite for zoonotic infection. *BMC Vet.*
Res. **14**, 271 (2018).

21. Bodenham, R.F. *et al.* Prevalence and speciation of brucellosis in febrile
patients from a pastoralist community of Tanzania. *Sci. Rep.* **10**, 7081 (2020).

22. Mboera, L.E.G., Kishamawe, C., Kimario, E. & Rumisha, S.F. Mortality
patterns of toxoplasmosis and its comorbidities in Tanzania: a 10-year
retrospective hospital-based survey. *Front. Public Health* **7**, 25 (2019).

23. Alemayehu, G. *et al.* Causes and flock level risk factors of sheep and goat
abortion in three agroecology zones in Ethiopia. *Front. Vet. Sci.* **8** (2021).

24. Okumu, T.A., John, N.M., Wabacha, J.K., Tsuma, V. & VanLeeuwen, J.
Seroprevalence of antibodies for bovine viral diarrhoea virus, *Brucella abortus*
and *Neospora caninum*, and their roles in the incidence of abortion/foetal loss
in dairy cattle herds in Nakuru District, Kenya. *BMC Vet. Res.* **15**, 95 (2019).

25. Tolosa, T., Bezabih, D. & Regassa, F. Study on seroprevalence of bovine
brucellosis and abortion and associated risk factor. *Bull. Anim. Health Prod.*
Afr. **58** (2010).

26. Reichel, M.P., Wahl, L.C. & Hill, F.I. Review of diagnostic procedures and
approaches to infectious causes of reproductive failures of cattle in Australia
and New Zealand. *Front. Vet. Sci.* **5** (2018).
872

873 27. Vidal, S. *et al.* Neglected zoonotic agents in cattle abortion: tackling the
874 difficult to grow bacteria. *BMC Vet. Res.* **13**, 373 (2017).
875

876 28. Maziku, M., Gebru, G. & Stapleton, J. Livestock health priorities in the
877 Tanzania livestock master plan. Tanzania Livestock Master Plan Brief 4.
878 Nairobi, Kenya: IRLI; 2017.
879

880 29. CDC. Workshop summary. One health zoonotic disease prioritization for
881 multisectoral engagement in Tanzania. In: Centers for Disease Control and
882 Prevention, editor.; 2017.
883

884 30. Crump, J.A. *et al.* Invasive bacterial and fungal infections among hospitalized
885 HIV-infected and HIV-uninfected children and infants in northern Tanzania.
886 *Trop. Med. Int. Health* **16**, 830-837 (2011).
887

888 31. Nyarobi, J. The epidemiology of Rift Valley fever in northern Tanzania. PhD
889 thesis, University of Glasgow, 2020.
890

891 32. Sindato, C. *et al.* A spatial analysis of Rift Valley fever virus seropositivity in
892 domestic ruminants in Tanzania. *PLoS One* **10**, e0131873-e0131873 (2015).
893

894 33. de Glanville, W.A. *et al.* Classification and characterisation of livestock
895 production systems in northern Tanzania. *PLoS One* **15**, e0229478 (2020).
896

897 34. Tamura, K. & Nei, M. Estimation of the number of nucleotide substitutions in
898 the control region of mitochondrial DNA in humans and chimpanzees.
899 *Molecular Biol. Evol.* **10**, 512-526 (1993).
900

901 35. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
902 evolutionary genetics analysis across computing platforms. *Molecular Biol.*
903 *Evol.* **35**, 1547-1549 (2018).
904

905 36. Braun, U., Hilbe, M., Peterhans, E. & Schweizer, M. Border disease in cattle.
906 *Vet. J.* **246**, 12-20 (2019).
907

908 37. Munyua, P. *et al.* Rift Valley fever outbreak in livestock in Kenya, 2006-2007.
909 *Am. J. Trop. Med. Hyg.* **83**, 58-64 (2010).
910

911 38. Woods, C.W. *et al.* An outbreak of Rift Valley fever in Northeastern Kenya,
912 1997-98. *Emerg. Infect. Dis.* **8**, 138-144 (2002).
913

914 39. Prabhu, M. *et al.* Q fever, spotted fever group, and typhus group rickettsioses
915 among hospitalized febrile patients in northern Tanzania. *Clin. Infect. Dis.* **53**,
916 e8-e15 (2011).
917

918 40. Anyamba, A. *et al.* Rift Valley fever outbreak in East Africa: signature of
919 climate extremes. American Geophysical Union, Fall Meeting 2018, abstract
920 #GH14A-07; 2018. pp. GH14A-07.
921

41. Mohamed, M. *et al.* Epidemiologic and clinical aspects of a Rift Valley fever outbreak in humans in Tanzania, 2007. *Am. J. Trop. Med. Hyg.* **83**, 22-27 (2010).

42. Mbotha, D. *et al.* Inter-epidemic Rift Valley fever virus seroconversions in an irrigation scheme in Bura, south-east Kenya. *Transbound Emerg Dis* **65**, e55-e62 (2018).

43. Hassan, A. *et al.* Epidemiological investigation of a Rift Valley fever outbreak in humans and livestock in Kenya, 2018. *Am. J. Trop. Med. Hyg.* **103**, 1649-1655 (2020).

44. ProMED-mail. Rift Valley fever - Eastern Africa: (Tanzania) alert. ProMED-mail. Archive Number: 20180619.5863493 2018 [cited 2021 March 19]Available from: http://www.promedmail.org

45. Roest, H.I. *et al.* Molecular epidemiology of *Coxiella burnetii* from ruminants in Q fever outbreak, the Netherlands. *Emerg. Infect. Dis.* **17**, 668-675 (2011).

46. Salifu, S.P., Bukari, A.A., Frangoulidis, D. & Wheelhouse, N. Current perspectives on the transmission of Q fever: Highlighting the need for a systematic molecular approach for a neglected disease in Africa. *Acta Tropica* **193**, 99-105 (2019).

47. Vanderburg, S. *et al.* Epidemiology of *Coxiella burnetii* infection in Africa: a OneHealth systematic review. *PLoS Negl. Trop. Dis.* **8**, e2787-e2787 (2014).

48. Crump, J.A. *et al.* Etiology of severe non-malaria febrile illness in Northern Tanzania: a prospective cohort study. *PLoS Negl. Trop. Dis.* **7**, e2324-e2324 (2013).

49. Galal, L. *et al.* *Toxoplasma* and Africa: One parasite, two opposite population structures. *Trends Parasitol.* **34**, 140-154 (2018).

50. Swai, E.S. & Kaaya, J.E. A survey of *Toxoplasma gondii* antibodies by latex agglutination assay in dairy goats in northern Tanzania. *Trop. Anim. Health Prod.* **45**, 211-217 (2013).

51. Tonouhewa, A.B. *et al. Toxoplasma gondii* infection in meat animals from Africa: systematic review and meta-analysis of sero-epidemiological studies. *Vet. World* **10**, 194-208 (2017).

52. Ferreira, S.C.M. *et al.* Evidence of high exposure to *Toxoplasma gondii* in free-ranging and captive African carnivores. *Int. J. Parasitol.* **8**, 111-117 (2019).

53. Innes, E.A., Bartley, P.M., Buxton, D. & Katzer, F. Ovine toxoplasmosis. *Parasitol.* **136**, 1887-1894 (2009).

971   54.   Barkallah, M. *et al.* Molecular prevalence of *Chlamydia* and *Chlamydia*-like
972   bacteria in Tunisian domestic ruminant farms and their influencing risk factors.
973   *Transbound Emerg Dis* **65**, e329-e338 (2018).
974

975   55.   Bhandi, S. *et al.* Brucellosis and chlamydiosis seroprevalence in goats at
976   livestock-wildlife interface areas of Zimbabwe. *Onderstepoort J. Vet. Res.* **86**,
977   e1-e9 (2019).
978

979   56.   Samkange, A., Katsande, T.C., Tjipura-Zaire, G. & Crafford, J.E.
980   Seroprevalence survey of *Chlamydophila abortus* infection in breeding goats
981   on commercial farms in the Otavi Veterinary District, northern Namibia.
982   *Onderstepoort J. Vet. Res.* **77**, E1-5 (2010).
983

984   57.   Allan, K.J. *et al.* Assessment of animal hosts of pathogenic *Leptospira* in
985   northern Tanzania. *PLoS Negl. Trop. Dis.* **12**, e0006444-e0006444 (2018).
986

987   58.   Bouley, A.J. *et al.* Brucellosis among hospitalized febrile patients in northern
988   Tanzania. *Am. J. Trop. Med. Hyg.* **87**, 1105-1111 (2012).
989

990   59.   Ellis, W.A. Animal Leptospirosis. In: Adler, B. (ed). *Leptospira and*
991   *Leptospirosis*, vol. 387. Springer: Berlin, 2015.
992

993   60.   Semango, G. *et al.* The sero-epidemiology of *Neospora caninum* in cattle in
994   northern Tanzania. *Front. Vet. Sci.* **6** (2019).
995

996   61.   Eleni, C. *et al.* Detection of *Neospora caninum* in an aborted goat foetus. *Vet.*
997   *Parasitol.* **123**, 271-274 (2004).
998

999   62.   Varaschin, M.S. *et al.* Congenital neosporosis in goats from the State of
1000   Minas Gerais, Brazil. *Korean J. Parasitol.* **50**, 63-67 (2012).
1001

1002   63.   González-Warleta, M. *et al.* Endogenous transplacental transmission of
1003   *Neospora caninum* during successive pregnancies across three generations
1004   of naturally infected sheep. *Vet. Res.* **49**, 106 (2018).
1005

1006   64.   Wodajo, H.D. *et al.* Contribution of small ruminants to food security for
1007   Ethiopian smallholder farmers. *Small Rumin. Res.* **184**, 106064 (2020).
1008

1009   65.   Smith, T., Godfrey, S. H., Buttery, P. J. The contribution of sheep and goats in
1010   alleviating poverty: communicating messages from research: Proceedings of
1011   the third DFID Livestock Production Programme link project (R7798)
1012   workshop for small ruminant keepers. Izaak Walton Inn, Embu, Kenya, 4-7
1013   February 2003
1014

1015   66.   Main, K. Seroprevalence and risk factors associated with bovine herpesvirus-
1016   1 infection in cattle in northern Tanzania. MSc thesis, University of Glasgow,
1017   2019.
1018

1019   67.   Newcomer, B.W. & Givens, D. Diagnosis and control of viral diseases of
1020   reproductive importance: Infectious bovine rhinotracheitis and bovine
1021   viral diarrhea. *Food Anim. Pract.* **32**, 425-441 (2016).
1022

1023   68.   Barkallah, M. *et al.* Survey of infectious etiologies of bovine abortion during
1024   mid- to late gestation in dairy herds. *PLoS One* **9**, e91549 (2014).
1025

1026   69.   Gethmann, J., Probst , C. & Conraths, F.J. Economic impact of a Bluetongue
1027   serotype 8 epidemic in Germany. *Front. Vet. Sci.* **7** (2020).
1028

1029   70.   Nusinovici, S., Souty, C., Seegers, H., Beaudeau, F. & Fourichon, C.
1030   Decrease in milk yield associated with exposure to bluetongue virus serotype
1031   8 in cattle herds. *J. Dairy Sci.* **96**, 877-888 (2013).
1032

1033   71.   Abera, T. *et al.* Bluetongue disease in small ruminants in south western
1034   Ethiopia: cross-sectional sero-epidemiological study. *BMC Res. Notes* **11**, 112
1035   (2018).
1036

1037   72.   Mulabbi, E.N., Ayebazibwe, C., Majalija, S., Batten, C.A. & Oura, C.A.
1038   Circulation of bluetongue virus in goats in the Karamoja region of Uganda. *J.*
1039   *S. Afr. Vet. Assoc.* **84**, E1-3 (2013).
1040

1041   73.   Toye, P.G. *et al.* Bluetongue and epizootic haemorrhagic disease virus in
1042   local breeds of cattle in Kenya. *Res. Vet. Sci.* **94**, 769-773 (2013).
1043

1044   74.   Coetzee, P., Stokstad, M., Venter, E.H., Myrmel, M. & Van Vuuren, M.
1045   Bluetongue: a historical and epidemiological perspective with the emphasis on
1046   South Africa. *Virolog. J.* **9**, 198 (2012).
1047

1048   75.   Maclachlan, N.J., Drew, C.P., Darpel, K.E. & Worwa, G. The pathology and
1049   pathogenesis of bluetongue. *J. Comp. Pathol.* **141**, 1-16 (2009).
1050

1051   76.   Knoll, M.D. *et al.* Bayesian estimation of pneumonia etiology: epidemiologic
1052   considerations and applications to the pneumonia etiology research for child
1053   health study. *Clin. Infect. Dis.* **64**, S213-s227 (2017).
1054

1055   77.   Hilbert, A. *et al.* Prevalence of *Coxiella burnetii* in clinically healthy German
1056   sheep flocks. *BMC Res. Notes* **5**, 152 (2012).
1057

1058   78.   Borel, N. *et al.* Laboratory diagnosis of ruminant abortion in Europe. *Vet. J.*
1059   **200**, 218-229 (2014).
1060

1061   79.   Sidi-Boumedine, K. *et al.* Development of harmonised schemes for the
1062   monitoring and reporting of Q‑fever in animals in the European Union. *EFSA*
1063   *Supporting Publication* **7** (2010).
1064

1065   80.   World Organisation for Animal Health (OIE). Q Fever. In: OIE, editor. *OIE*
1066   *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*; 2018.
1067

1068   81.   Herzog, C.M. *et al.* Pastoral production is associated with increased peste des
1069   petits ruminants seroprevalence in northern Tanzania across sheep, goats
1070   and cattle. *Epidemiol. Infect.* **147**, e242 (2019).
1071
1072   82.   Kadohira, M., McDermott, J.J., Shoukri, M.M. & Kyule, M.N. Variations in the
1073   prevalence of antibody to *Brucella* infection in cattle by farm, area and district
1074   in Kenya. *Epidemiol. Infect.* **118**, 35-41 (1997).
1075
1076   83.   Kairu-Wanyoike, S. *et al.* Positive association between *Brucella* spp.
1077   seroprevalences in livestock and humans from a cross-sectional study in
1078   Garissa and Tana River Counties, Kenya. *PLoS Negl. Trop. Dis.* **13**,
1079   e0007506 (2019).
1080
1081   84.   Bodenham, R.F. Brucellosis in northern Tanzania: Investigating the
1082   epidemiology of human infection and evaluating diagnostic test performance
1083   in animal hosts. PhD thesis, University of Glasgow, 2020.
1084
1085   85.   Dorai-Raj, S. binom: binomial confidence intervals for several
1086   parameterizations. *R package version 1.1-1*; 2014.
1087
1088   86.   Wilson, K., Livingstone, M. & Longbottom, D. Comparative evaluation of eight
1089   serological assays for diagnosing *Chlamydophila abortus* infection in sheep.
1090   *Vet. Microbiol.* **135**, 38-45 (2009).
1091
1092   87.   Hamilton, C.M. *et al. Toxoplasma gondii* in livestock in St. Kitts and Nevis,
1093   West Indies. *Parasit Vectors* **8**, 166 (2015).
1094
1095   88.   Matero, P. *et al.* Rapid field detection assays for *Bacillus anthracis*, *Brucella*
1096   spp., *Francisella tularensis* and *Yersinia pestis. Clin. Microbiol. Infect. Dis.* **17**,
1097   34-43 (2011).
1098
1099   89.   Probert, W.S., Schrader, K.N., Khuong, N.Y., Bystrom, S.L. & Graves, M.H.
1100   Real-time multiplex PCR assay for detection of *Brucella* spp., *B. abortus*, and
1101   *B. melitensis. J. Clin. Microbiol.* **42**, 1290-1293 (2004).
1102
1103   90.   Livingstone, M., Wheelhouse, N., Maley, S.W. & Longbottom, D. Molecular
1104   detection of *Chlamydophila abortus* in post-abortion sheep at oestrus and
1105   subsequent lambing. *Vet. Microbiol* **135**, 134-141 (2009).
1106
1107   91.   Stoddard, R.A., Gee, J.E., Wilkins, P.P., McCaustland, K. & Hoffmaster, A.R.
1108   Detection of pathogenic *Leptospira* spp. through TaqMan polymerase chain
1109   reaction targeting the *LipL*32 gene. *Diagn. microbiol. Infect. Dis.* **64**, 247-255
1110   (2009).
1111
1112   92.   Buxton, D. *et al.* The pathogenesis of experimental neosporosis in pregnant
1113   sheep. *J. Comp. Pathol.* **118**, 267-279 (1998).
1114
1115   93.   Hurtado, A., Aduriz, G., Moreno, B., Barandika, J. & García-Pérez, A.L. Single
1116   tube nested PCR for the detection of *Toxoplasma gondii* in fetal tissues from
1117   naturally aborted ewes. *Vet. Parasitol.* **102**, 17-27 (2001).

1118
1119 94. Drosten, C. *et al.* Rapid detection and quantification of RNA of Ebola and
1120   Marburg viruses, Lassa virus, Crimean-Congo hemorrhagic fever virus, Rift
1121   Valley fever virus, dengue virus, and yellow fever virus by real-time reverse
1122   transcription-PCR. *J. Clin. Microbiol.* **40**, 2323-2330 (2002).
1123
1124 95. Burrells, A. *et al.* Evidence of the three main clonal *Toxoplasma gondii*
1125   lineages from wild mammalian carnivores in the UK *Parasitol.* **140**, 1768-1776
1126   (2013).
1127
1128 96. Becher, P. *et al.* Phylogenetic analysis of pestiviruses from domestic and wild
1129   ruminants. *J. Gen. Virol.* **78 (Pt 6)**, 1357-1366 (1997).
1130
1131 97. Vilcek, S., Nettleton, P.F., Paton, D.J. & Belák, S. Molecular characterization
1132   of ovine pestiviruses. *J. Gen. Virol.* **78**, 725-735 (1997).
1133
1134 98. Vilcek, S. *et al.* Bovine viral diarrhoea virus genotype 1 can be separated into
1135   at least eleven genetic groups. *Arch. Virol.* **146**, 99-115 (2001).
1136
1137 99. Dubey, J.P. & Schares, G. Neosporosis in animals - the last five years. *Vet.*
1138   *Parasitol.* **180**, 90-108 (2011).
1139
1140 100. World Organisation for Animal Health (OIE). Brucellosis (*Brucella abortus*, *B.*
1141   *melitensis* and *B. suis*) (infection with *B. abortus, B. melitensis* and *B. suis*).
1142   In: OIE, editor. *Manual of diagnostic tests and vaccines for terrestrial animals*;
1143   2018.
1144
1145 101. World Organisation for Animal Health (OIE). Enzootic Abortion of Ewes
1146   (Ovine Chlamydiosis) (Infection with *Chlamydia abortus*). In: OIE, editor.
1147   *Manual of diagnostic tests and vaccines for terrestrial animals*; 2018.
1148
1149 102. Touratier, A. *et al.* Comment faire le diagno stic d'un élevage cliniquement
1150   atteint de fièvre Q? *Recueil des Journées Nationales des GTV, Nantes*, 147-
1151   155 (2007).
1152
1153 103. World Organisation for Animal Health (OIE). Leptospirosis In: OIE, editor. *OIE*
1154   *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*; 2018.
1155
1156 104. Dubey, J.P. & Schares, G. Diagnosis of bovine neosporosis. *Vet. Parasitol.*
1157   **140**, 1-34 (2006).
1158
1159 105. World Organisation for Animal Health (OIE). Toxoplasmosis. In: OIE, editor.
1160   *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*; 2018.
1161
1162 106. World Organisation for Animal Health (OIE). Infectious Bovine
1163   Rhinotracheitis/Infectious Pustular Vulvovaginits. In: OIE, editor. *Manual of*
1164   *diagnostic tests and vaccines for terrestrial animals*; 2018.
1165

1166 107. World Organisation for Animal Health (OIE). Bluetongue (infection with
1167 bluetongue virus). In: OIE, editor. *Manual of diagnostic tests and vaccines for*
1168 *terrestrial animals*; 2018.
1169
1170 108. World Organisation for Animal Health (OIE). Bluetongue. Aetiology,
1171 epidemiology, diagnosis, prevention and control references. 2021.
1172
1173 109. World Organisation for Animal Health (OIE). Bovine Viral Diarrhoea. In: OIE,
1174 editor. *Manual of diagnostic tests and vaccines for terrestrial animals*; 2018.
1175
1176 110. World Organisation for Animal Health (OIE). Rift Valley Fever (infection with
1177 Rift Valley fever virus). In: OIE, editor. *OIE Manual of Diagnostic Tests and*
1178 *Vaccines for Terrestrial Animals*; 2018.
1179
1180

Acknowledgements

The abortion cohort study was supported by the Supporting Evidence Based

Interventions project, University of Edinburgh (grant number R83537 CH). The cross-

sectional studies were supported by US National Institutes of Health-National (NIH)

Science Foundation Ecology and Evolution of Infectious Disease program (R01

TW009237) and the UK Biotechnology and Biological Sciences Research Council

(BBSRC) (grant numbers: BB/J010367/1, BB/L018926/1). MD was funded by the

Scottish Government. CH, CH, EI, FK, ML, GR and DL are supported by funding

from the Scottish Government Rural and Environment Science and Analytical

Services Division (RESAS). BW is funded by the BBSRC. We would like to thank

Rosanne de Jong, Divine Ekwem, Erin Hodgkinson, Karen Main and Jen Jen Yu for

assistance with serological testing at the University of Glasgow and Clare

Underwood (MRI) for expert immunohistological preparations. We appreciate the

huge efforts of our field team, Rigobert Tarimo, Fadhili Mshana, and Hassan Hussein

as well as the assistance from District Veterinary Officers and LFOs. Thanks to Paul

Johnson (University of Glasgow) for support with data management and curation,

and Dassa Nkini (NM-AIST) and Elizabeth Kussaga (KCRI) for administrative

support.

1199

Author contributions

Developed the study concept and design of the work: KT, JRC, WdG, FL, JB, JAC,

MD, JH, CH, EAI, FK, DL, BM, ON, NW, SC, KA; Collected samples and metadata:

TK, WdG, FL, GS; Sample processing KT, NA, RC, GC, MD, CH, EI, FK, ML, CM,

VM, JN, GR, NW, BW, SC, KA; Data analysis and interpretation: KT, JRC, WdG, FL,

NA, RC, GC, JAC, MD, JH, CH, EI, FK, ML, DL, CM, VM, JN, GR, NW, BW, SC, KA;

Initial drafting the article: KT, WdG, FL, JH, SC, KA; All authors provided critical

revision of the article and provided approval for submission.

1208

1209 **Competing interests**

1210 The authors declare no competing interests.