Prospective cohort study reveals unexpected aetiologies of livestock abortion

in northern Tanzania

Kate M. Thomas[1,2*$], Tito Kibona[3], John R. Claxton[4], William A. de Glanville[4], Felix

Lankester[5,6], Nelson Amani[2], Joram J. Buza[3], Ryan W. Carter[4], Gail E. Chapman[7],

John A. Crump[1,8], Mark P. Dagleish[9], Jo E. B. Halliday[4], Clare M. Hamilton[9],

Elisabeth A. Innes[9], Frank Katzer[9], Morag Livingstone[9], David Longbottom[9], Caroline

Millins[7#], Blandina T. Mmbaga[2,8], Victor Mosha[2], James Nyarobi[4], Obed M.

Nyasebwa[10], George C. Russell[9], Paul N. Sanka[11], George Semango[3,], Nick

Wheelhouse[12], Brian J. Willett[13], Sarah Cleaveland[4], and Kathryn J. Allan[4,7]

11
[1]Centre for International Health, Dunedin School of Medicine, University of Otago,
Dunedin, New Zealand
[2]Kilimanjaro Clinical Research Institute, Good Samaritan Foundation, Moshi, United
Republic of Tanzania
[3]Nelson Mandela African Institution of Science and Technology (NM-AIST), Tengeru,
United Republic of Tanzania
[4]Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical
Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom
[5]Paul G. Allen School for Global Animal Health, Washington State University, WA,
United States of America
[6]Global Animal Health Tanzania, Arusha, United Republic of Tanzania
[7]School of Veterinary Medicine, University of Glasgow, Glasgow, United Kingdom
[8]Kilimanjaro Christian Medical University College, Moshi, United Republic of
Tanzania
[9]Moredun Research Institute, Midlothian, United Kingdom
[10]Zonal Veterinary Centre – Arusha, Ministry of Livestock and Fisheries, Arusha,
United Republic of Tanzania
[11]Tanzania Veterinary Laboratory Agency, Arusha, United Republic of Tanzania
[12]School of Applied Sciences, Edinburgh Napier University, Edinburgh, United
Kingdom
[13]Medical Research Council, University of Glasgow Centre for Virus Research,
Glasgow, United Kingdom
34
35 * Corresponding author
36 E-mail: katethomas144@gmail.com OR Kate.Thomas@mpi.govt.nz (KMT)
37
38 $ Current address: New Zealand Food Safety, Ministry for Primary Industries,
39 Wellington, New Zealand
40 # Current address: Institute of Infection, Veterinary and Ecological

41   Sciences, University of Liverpool, United Kingdom
42

Abstract

Livestock abortion is an important cause of productivity losses worldwide and many

infectious causes of abortion are zoonotic pathogens that impact on human health.

Little is known about the relative importance of infectious causes of livestock

abortion in Africa, including in subsistence farming communities that are critically

dependent on livestock for food, income, and wellbeing. We conducted a prospective

cohort study of livestock abortion, supported by cross-sectional serosurveillance, to

determine aetiologies of livestock abortions in livestock in Tanzania. This approach

generated several important findings including detection of a Rift Valley fever virus

outbreak in cattle; high prevalence of *C. burnetii* infection in livestock; and the first

report of *Neospora caninum*, *Toxoplasma gondii,* and pestiviruses associated with

livestock abortion in Tanzania. Our approach provides a model for abortion

surveillance in resource-limited settings. Our findings add substantially to current

knowledge in sub-Saharan Africa, providing important evidence from which to

prioritise disease interventions.

Introduction

Infectious causes of abortion, including early foetal loss and stillbirth, affect livestock production worldwide with the potential to cause major economic losses (De Vries, 2006). Many abortions are caused by zoonotic pathogens that also pose a risk of infection to people (Givens & Marley, 2008). Although the economic losses can be substantial in high-production, intensive agricultural systems (De Vries, 2006), abortion-associated livestock production losses have marked negative impacts for the poorest communities in the world, many millions of whom are critically dependent on livestock for economic livelihoods, food security, health, and wellbeing (FAO, 2011, 2018).

Livestock abortions can be caused by a wide range of pathogens as well as non-infectious causes such as nutritional deficiencies and trauma (Givens & Marley, 2008). While many infectious causes of abortions occur worldwide, aetiological data based on robust laboratory-confirmed diagnoses are dominated by studies from farming systems in high- and middle-income countries (Cabell, 2007; Campero, Moore, Odeón, Cipolla, & Odriozola, 2003; Wolf-Jäckel et al., 2020) with data lacking from low-income settings. Comparison of aetiologies across geographic areas is challenging as studies vary in methodologies and in the specific pathogens that are investigated. Furthermore, the ability to identify a specific aetiology of a livestock abortion is often relatively low, with less than 35% cases attributed to specific diagnoses (Cabell, 2007; Kirkbride, 1993; Wolf-Jäckel et al., 2020).

Considerable surveillance and diagnostic infrastructure is required for robust case attribution of livestock abortions (Buxton & Henderson, 1999; Cabell, 2007), which

typically limits the data available from low resource settings. For countries where

data do exist, geographic patterns appear to vary quite widely. For example, *Coxiella*

burnetii has been determined to be a common cause of abortion in sheep and goats

in Canada (Hazlett et al., 2013), and in goats in the Netherlands (van den Brom et

al., 2012), but only rarely in sheep in the United States of America (USA) (Kirkbride,

1993; Oliveira et al., 2017) or the United Kingdom (UK) (Buxton & Henderson, 1999),

despite evidence of widespread seropositivity in both countries (Loftis, Reeves,

Miller, & Massung, 2012; Velasova et al., 2017). *Chlamydia abortus* is one of the

most common causes of infectious ovine abortion worldwide, including the UK, the

USA and most of northern Europe but has not been found to be an abortion issue in

sheep in Australia or New Zealand (Essig & Longbottom, 2015) and there is little

available information from the African continent, other than in a few select countries

(e.g. Tunisia, Algeria, Namibia, Zimbabwe, and Morocco (Benkirane, Jabli, &

Rodolakis, 1990; Hireche, Ababneh, Bouaziz, & Boussena, 2016; Rekiki et al., 2002;

Seth-Smith et al., 2017). In tropical areas, epidemic-prone pathogens, such as Rift

Valley fever virus (RVFV) which poses a severe health threat to both animals and

people, are also a concern (Budasha, Gonzalez, Sebhatu, & Arnold, 2018). No less

important are the endemic zoonoses such as *Brucella abortus* and *B. melitensis*, and

zoonotic protozoal agents such as *Toxoplasma gondii,* which are of substantial

concern in subsistence farming settings where people live closely with their animals

(Bodenham et al., 2020; FAO, 2011) or in communities where HIV predisposes the

human population to severe infection e.g., with toxoplasmosis (Mboera, Kishamawe,

Kimario, & Rumisha, 2019).

In Africa, inferences about the potential role and relative importance of abortigenic

agents are often made from serological studies of livestock (Alemayehu et al., 2021;

Okumu, John, Wabacha, Tsuma, & VanLeeuwen, 2019; Tolosa, Bezabih, &

Regassa, 2010). This is due to a number of factors including a lack of national-level

surveillance for syndromes such as abortions in livestock; restricted access to

veterinary care for many subsistence farmers, who are therefore unable to

investigate individual abortion events; and limitations in diagnostic laboratory

capacity (e.g., little to no access to histopathology services or molecular diagnostic

tests that could help to generate data on different aetiologies of abortion). While

serological data is valuable to identify the presence of abortigenic agents circulating

within a livestock population, there is also a need to recognise the limitations of such

data and the inferences that can be made based on it (Reichel, Wahl, & Hill, 2018;

Vidal et al., 2017). Livestock seropositivity indicates only that an animal has been

exposed to a pathogen and does not necessarily indicate current disease status or

causality of an abortion event. Additionally, chronic low-level carriage of some

abortigenic agents (e.g., *C. burnetii*) in healthy animals complicates attribution of

abortion. To date, in sub-Saharan Africa, there remains an almost complete absence

of published data on livestock abortion aetiologies determined through confirmatory

diagnostic testing, resulting in a very weak evidence base on which to prioritise

animal health interventions.

In Tanzania, livestock diseases are prioritised for prevention and control largely on

the basis of the potential for transboundary spread and zoonotic impacts (Maziku,

Gebru, & Stapleton, 2017). A recent exercise to prioritise zoonotic diseases ranked

two abortigenic agents, *Brucella* and RVFV amongst the top six zoonoses for

prevention and control (CDC, 2017). While there is clear evidence that brucellosis

and Rift Valley fever (RVF) are both diseases of public health concern in Tanzania

(Crump et al., 2011; Nyarobi, 2020; Sindato et al., 2015), no data currently exist on

the relative importance of these or other zoonotic pathogens, such as *C. burnetii*,

Leptospira spp., *C. abortus* or *T. gondii*, as causes of livestock abortion. Nor are

there any data available on the contribution of other abortigenic pathogens of global

importance, including *N. caninum*, bluetongue virus and pestiviruses such as bovine

viral diarrhoea virus (BVDV) and border disease virus (BDV). Without robust

aetiological data, it is unlikely that current approaches to livestock disease

prioritisation will reflect the true impact of abortigenic pathogens, with the risk that

scarce resources for disease control and prevention will be targeted ineffectively and

achieve sub-optimal productivity and public health outcomes.

We sought to address knowledge gaps around the aetiology of livestock abortion in

Tanzania through a prospective cohort study conducted in three regions spanning a

range of agro-ecological settings (de Glanville et al., 2020). We used a combination

of field investigation and laboratory diagnosis of abortion events in cattle, goats, and

sheep to determine aetiologies of livestock abortions for a targeted selection of

known abortigenic agents. For these selected agents, we developed a set of case

definitions to attribute abortion events to a specific aetiological agent based on

molecular diagnostic assays in the absence of comprehensive histopathology data.

In the same study area, we also carried out serological analyses of a large sample of

livestock sera collected by cross-sectional studies in the same area to determine

seroprevalence to the same range of abortigenic pathogens at a regional level.

Results

Figure 1 shows the location of sampling for the serological cross-sectional and

abortion cohort studies.

Figure 1: Distribution of villages in cross-sectional studies for serologic

baseline testing (2013-2016; red dots) and abortion cohort sampling wards

(2017-2019; yellow shading) with number of abortion events investigated,

Arusha, Kilimanjaro, and Manyara Regions, northern Tanzania.

<u>Cross-sectional sample serology</u>

Samples were available from 3,629 cattle, 3,344 goats, and 2,628 sheep from 567

households in 43 villages. Figure 2 summarises the cross-sectional seroprevalence

of pathogens in ruminant livestock sampled in northern Tanzania. Seroprevalence at

individual level for cross sectional livestock with Clopper Pearson 95% confidence

intervals is detailed in supplementary materials (S1, A).

Figure 2: Seroprevalence of pathogens from cross-sectional studies (2013-2015 and 2016) in cattle, goats, and sheep, northern Tanzania.
Where x = not tested and error bars represent Clopper-Pearson 95% confidence intervals

Abortion cohort study

During the prospective cohort study period, a total of 215 abortion events were investigated, 71 (33%) in cattle, 100 (46.5%) in goats, and 44 (20.5%) in sheep. The numbers of each sample type collected for abortion events are detailed in Table 1.

Table 1: Number and type of samples collected from abortion events for each livestock animal species, northern Tanzania, 2017-2019

Sample types collected	Cattle (n=71)	Goat (n=100)	Sheep (n=44)
Vaginal swabs	71 (100%)	98 (98.0%)	44 (100%)
Exterior foetal swabs	37 (52.1%)	25 (25%)	5 (11.4%)
Placental tissue	34 (47.9%)	25 (25%)	1 (2.3%)
Acute sera	71 (100%)	98 (98.0%)	43 (97.7%)

Convalescent sera	49 (69.0%)	56 (56.0%)	29 (65.9%)

Abortion cohort acute serology

The acute seroprevalence results by pathogen in the abortion cohort was broadly similar to the cross-sectional survey. A table showing these results is available in the supplementary materials (S1; A).

Diagnosis of abortion aetiology

We attributed 42 (19.5%) of 215 abortion events to one or more of the abortigenic pathogens included in our diagnostic panel. Table 2 summarises the number of abortion events with positive diagnostic test results that met the study case definitions. Histopathology and immunohistochemistry (IHC) results are shown in supplementary materials S1, B.

Table 2: Number of abortion cases (cattle, goats and sheep) attributed to abortigenic agents using study case definitions, northern Tanzania 2017-2019

Pathogen	Case definition	Number of cases attributed by animal species				Strength of evidence of abortion cause
		Cattle n/n (%)	Goats n/n (%)	Sheep n/n (%)	Total (n)	
Brucella spp.	qPCR	1/71 (1.4)	0/100 (0.0)	0/44 (0.0)	1	Presumptive
Chlamydia abortus	qPCR	0/71 (0.0)	0/100 (0.0)	0/44 (0.0)	0	Definitive
	Sero-conversion	n/a	1/56[‡] (0.0)	0/29 (0.0)		Suggestive
Coxiella burnetii	qPCR	0/71 (0.0)	5/100 (5.0)	1/44 (2.3)	6	Presumptive
Leptospira spp.	qPCR	0/71 (0.0)	0/100 (0.0)	0/44 (0.0)	0	Presumptive
	Sero-conversion	0/49 (0.0)	n/a	n/a		
Neospora caninum	PCR	9/71[#] (12.7)	1/100 (1.0)	0/44 (0.0)	10[#]	Presumptive
Toxoplasma gondii	PCR	0/49 (0.0)	0/100 (0.0)	1/44 (2.3)	1	Definitive

Bovine herpes virus (BHV-1)	Sero-conversion	2/49[#]	(4.1)	n/a		n/a		2[#]	Suggestive
Bluetongue virus (BTV)	RT-qPCR	0/71	(0.0)	0/100	(0.0)	0/44	(0.0)	0	Definitive
Pestiviruses	RT-qPCR	2/71	(2.8)	1/100	(1.0)	6/44	(13.6)	9	Definitive
Rift Valley fever virus (RVFV)	RT-qPCR	14/71	(19.7)	0/100	(0.0)	0/44	(0.0)	14	Definitive
	Sero-conversion	1/49[*]	(1.4)	0/56	(0.0)	0/29	(0.0)		
None	Not detected	44/71	(62.0)	92/100	(92.0)	36/44	(81.8)	173	n/a
TOTAL		**27/71**	**(38.0)**	**8/100**	**(8.0)**	**8/44**	**(18.2)**	**215**	

[#] One case in a cow met both BHV-1 (seroconversion) and *Neospora* (PCR) case definition and was
thus included in both
[*] One case in a cow met both RT-qPCR and seroconversion case definitions for RVFV and was thus
included in both
[‡] Seroconversion was observed in one goat with a qPCR-negative swab sample for *C. abortus* and
therefore this case did not meet the case definition.
n/a = not applicable

For some bacterial pathogens, despite evidence of infection in the dam, laboratory

data was not considered sufficient to support a diagnosis of abortion associated with

the pathogen. This was particularly true for *C. burnetii*, which was detected by qPCR

with a threshold cycle (Ct) value <40 in 16 (22.5%) of 71 cattle, 24 (24.5%) of 98

goats, and 12 (27.3%) of 44 sheep, however only six (11.5%) of these 52 animals

met the study case definition (Ct ≤ 27) for *C. burnetii* as a presumptive aetiology of

abortion. For *C. abortus*, although seroconversion between acute and convalescent

samples to *C. abortus* was detected in one (1.8%) of 56 goats with paired sera

available for testing, this animal had a negative vaginal swab qPCR result and

therefore did not meet our case definition for *C. abortus*-associated abortion.

Of the protozoal pathogens, *Neospora* DNA was diagnosed by nested conventional

polymerase chain reaction (PCR) in nine (12.7%) of 71 cattle and one (1.0%) of 98

goat abortion events. In one cattle case, a co-infection with BHV-1 was also detected

suggesting a multi-factorial aetiology. *Toxoplasma* DNA was detected by nested

conventional PCR in one (2.3%) of 44 sheep abortion events. The single *T. gondii*

PCR positive sheep sample was genotyped at 7/10 marker and was found to have

type II alleles at all of them, indicative of a type II clonal genotype infection.

Of the viral pathogens, RVF was most commonly diagnosed, with RVFV RNA

detected by reverse transcriptase qPCR (RT-qPCR) in 14 (19.7%) of 71 cattle

abortion events. Evidence of seroconversion was detected in one animal that was

positive by RT-qPCR. The other 13 RT-qPCR positive animals were positive in both

acute and convalescent sera, meeting World Organisation for Animal Health (OIE)

guidelines for definitive diagnosis of infection. Pestiviruses were also commonly

detected with two (2.8%) of 71 cattle, six (13.6%) of 44 sheep, and one (1.0%) of

100 goat abortion events positive by RT-qPCR.

<u>Pestivirus speciation results</u>

Among the nine animals that tested positive for pestiviruses by RT-qPCR, N^{Pro}

coding sequence was obtained from six samples for phylogenetic analysis (Figure

3). BVDV infection was confirmed in two animals (one cow and one sheep) with a

distinct genotype detected in each case. The cattle sample (SF1-190) was most

similar to reference sequences from BVDV genotype 1d whereas the sheep sample

(SD1-147) was most similar to reference sequences from BVDV genotype 1b.

Sequences obtained from the remaining four samples (SD1-144, SD1-145, SD1-155,

SD1-160), all obtained from sheep, were similar to each other but appeared to be

distinct from all of the reference sequences. These sequences represent a

potentially novel, previously unreported pestivirus species and formed a separate

clade on phylogenetic analysis.

Figure 3: Results of evolutionary analysis of the N^{Pro} coding region from pestivirus sequences obtained from cattle and sheep sampled in the abortion cohort study, northern Tanzania.

The evolutionary history was inferred using the Maximum Likelihood method and Tamura-Nei model of nucleotide substitution (Tamura & Nei, 1993). The tree with the highest log likelihood is shown. The percentage of trees from 500 Bootstrap repeats in which the associated taxa clustered together is shown next to the branches. Evolutionary analyses were conducted in MEGA X (Kumar, Stecher, Li, Knyaz, & Tamura, 2018). Reference sequences representing nine pestivirus species were also included (Braun, Hilbe, Peterhans, & Schweizer, 2019) and are labelled with virus species (A-H, K), genotype and strain name and Genbank accession numbers. Pestivirus species are also shown by bracket labels on the right of the figure. Samples analysed in this study are shown in bold and labelled with unique identifiers (SD/F1-XXX), where SD denotes vaginal swab taken from aborting dam; SF denotes swab taken from aborted foetus. Abbreviations: APPV = atypical porcine pestivirus; BDV =

264 border disease virus; BVDV = bovine viral diarrhoea virus; CSFV = classical swine
265 fever virus.
266

Discussion

To our knowledge, this is the first multi-pathogen study of the aetiologies of livestock

abortion in Tanzania. We developed and applied an approach designed to optimise

abortion investigations in resource limited settings, complemented by traditional

cross-sectional surveillance to generate a more comprehensive understanding of the

distribution and impact of several important abortigenic pathogens. This approach

generated several new insights for Tanzania including: a) detection of an RVF

outbreak in cattle during an inter-epidemic period; b) relatively high prevalence of *C.*

burnetii infection in cattle, goats and sheep; c) the first demonstration of *N. caninum*,

T. gondii, and pestiviruses, including novel pestivirus species, associated with

livestock abortion in Tanzania, including detection in less typical hosts (for example,

N. caninum in goats); but, contrary to prior expectations, d) limited or no detection of

Brucella spp., *C. abortus*, and *Leptospira* spp. in cases of livestock abortion. Our

findings provide important information on the health and production impacts of a

range of livestock pathogens in East Africa with relevance for prioritising disease

interventions in the region.

Our study demonstrates the importance of zoonotic pathogens as causes of abortion

in ruminants in Tanzania, with implications for human as well as animal health. Two

important zoonotic pathogens were diagnosed in multiple abortion events by our

study: RVFV, a virus that can cause devastating outbreaks of human and animal

disease (Munyua et al., 2010; Woods et al., 2002), and *C. burnetii*, an important

bacterial cause of human febrile illness in Tanzania (Prabhu et al., 2011). RVFV is a

high priority zoonosis in East Africa and outbreaks are frequently associated with

livestock abortion storms (Budasha et al., 2018). Direct animal to human

transmission of infection can occur following inhalation of aerosolized virus particles

or during close contact with tissues and bodily fluids of infected animals, for

example, when assisting with parturition or handling aborted foetuses (Anyamba et

al., 2018; Mohamed et al., 2010). Although RVF usually occurs as large outbreaks

occurring every 10-20 years, there is increasing serological evidence of transmission

in the period between these epidemics (Mbotha et al., 2018). This study detected the

occurrence of an apparent outbreak in cattle that was previously unreported in

Tanzania, despite RVF outbreaks being detected in Kenya, Uganda, and Rwanda at

the same time (Anyamba et al., 2018; Hassan et al., 2020). Although an alert of RVF

was issued in Tanzania in June 2018 at the time of this study (ProMED-mail, 2018),

to our knowledge, no human cases were reported. Our finding highlights the value of

livestock abortion surveillance in early detection of this epidemic-prone pathogen

and also demonstrates the value of molecular diagnostic capacity for detection of

RVF livestock cases.

Coxiella burnetii is a pathogen of concern in Europe, where outbreaks of infection in

small ruminant livestock have also led to outbreaks of human disease (Roest et al.,

2011). *C. burnetii* has also been detected in humans and in a wide range of animal

species across Africa (Salifu, Bukari, Frangoulidis, & Wheelhouse, 2019; Vanderburg

et al., 2014), and accounts for notable proportions of severe human febrile illness in

Africa (Vanderburg et al., 2014), including in northern Tanzania (Crump et al., 2013;

Prabhu et al., 2011). Our results provide important evidence of the impact of *C.*

burnetii as a cause of livestock reproductive losses, particularly in small ruminants,

revealing a dual human and animal health burden of this zoonotic pathogen. In our

abortion cohort study, we applied a stringent case definition that exceeded the

minimum guidelines stated by the OIE for diagnosis of *C. burnetii*-associated

abortion. Using this case definition, we diagnosed six cases of *C. burnetii*-associated

abortion, but it is also worth noting that *C. burnetii* DNA was detected in a total of 52

(24.4%) abortion events indicating that shedding of this zoonotic bacterium is even

more prevalent than the number of attributed cases would suggest. Based on our

findings, we suggest that the development of preventive and control measures

against *C. burnetii* should be a priority for future work that could have substantial

benefits for human and animal health.

Our study confirmed *T. gondii,* another zoonotic pathogen, is associated with

livestock abortion in Tanzania. Toxoplasmosis is also an important, sometimes fatal,

zoonosis in Tanzania, particularly among immunocompromised people (Mboera et

al., 2019). There are limited data on genotypes present in Africa but the Type II

clonal lineage found in the single PCR positive sheep abortion event is known to be

dominant in North and East Africa so aligns with previous studies (Galal et al., 2018).

The cross-sectional serological data showing widespread exposure of *T. gondii* in

sheep and goats are consistent with previous published findings of high

seroprevalence in people and animals, including livestock (Swai & Kaaya, 2013;

Tonouhewa et al., 2017) and wild felid species in East Africa (Ferreira et al., 2019).

Given that toxoplasmosis is one of the most important abortigenic agents of sheep

worldwide, it is perhaps surprising that we did not detect more cases of *T. gondii* as

a cause of sheep abortion in this study. However, high flock seroprevalences are

typically associated with low rates of abortion due to the development of protective

immunity in ewes infected with *T. gondii* prior to mating (Innes, Bartley, Buxton, &

Katzer, 2009).

Other zoonotic pathogens were detected infrequently in the abortion cohort study,

with only a single case attributed to *Brucella* spp. in a cow, a single detection of

seroconversion to *C. abortus* in a goat, and no evidence of abortions associated with

Leptospira spp. in any livestock species. Although *C. abortus* has been reported as a

cause of abortion in sheep elsewhere in northern and southern Africa (Barkallah et

al., 2018; Bhandi et al., 2019; Samkange, Katsande, Tjipura-Zaire, & Crafford, 2010),

our study generated little evidence that *C. abortus* was contributing to the burden of

production losses in sheep in northern Tanzania at the time of our study. Our cross-

sectional serological findings also support the conclusion that levels of exposure in

small ruminants are very low across northern Tanzania. However, these findings do

raise concerns about the potential impact should infection be introduced to this

predominantly naïve population in the future.

Both *Brucella* spp. and *Leptospira* spp. are recognised as important zoonoses in

northern Tanzania, causing large proportions of severe human febrile illness (Allan

et al., 2018; Bodenham et al., 2020; Bouley et al., 2012; Crump et al., 2013) and are

known to be circulating in livestock in these areas of Tanzania. Seroprevalence of

Brucella spp. was low in all three livestock species (Fig. 2), which is consistent with

the low frequency of detection in abortion cases. In contrast, although *Leptospira*

was not detected by qPCR in any abortion event investigated, evidence of exposure

to *Leptospira borgpetersenii* serovar Hardjo, and *Leptospira interrogans* serovar

Hardjo (subsequently referred to as *Leptospira* Hardjo serovars) was relatively high

in cattle in the cross-sectional study. Several possible explanations may explain this

discrepancy. Similar to *T. gondii,* prior exposure to *Leptospira* serovars may lead to

protective immunity prior to animals becoming pregnant. Additionally, livestock

abortions associated with *Leptospira* are often seen in outbreaks or 'abortion storms'

following either a new introduction into a herd or a seasonal outbreak where flooding

or wet weather conditions support transmission of novel serovars (Ellis, 2015). The

timespan of our project meant that we may have missed intermittent or infrequent

abortion storms associated with *Leptospira* in livestock. However, even accounting

for these limitations, our study suggests that in these agricultural settings, the

proportion of abortions attributable to *Brucella* spp. and *Leptospira* spp. may be

lower than is widely perceived. Alongside other constraints to the uptake and use of

livestock brucellosis vaccines these data raise questions about the likely

effectiveness and sustainability of livestock vaccination-based control strategies in

this context if based on a farmer pays model only. To effectively mitigate both the

animal and human impacts of brucellosis a diversity of strategies may be needed to

achieve the vaccination coverage required to reduce animal production impacts and

also to reduce the substantial human health burden from this disease.

In addition to zoonotic pathogens, our study provides also provides valuable data on

several production-limiting infections of ruminants that are often over-looked in sub-

Saharan Africa. In particular, while *N. caninum* and pestiviruses are considered

important causes of abortions and productivity loss in Europe and the USA, there are

few data available describing their impact in African livestock. Our results provide the

first direct evidence of livestock abortion associated with *N. caninum* infection in

Tanzania, a conclusion which is supported by a positive relationship between *N*

caninum seroprevalence and in-herd abortion rates in cattle detected by earlier work

in our study area (Semango et al., 2019). In this study, *N. caninum* was

predominately diagnosed among cattle abortions, but notably also in one goat

abortion event. Serological evidence of low-level exposure to *N. caninum* in goats

(1.7%) and sheep (1.5%) was also detected across the region. Although abortion

and foetal pathology associated with *N. caninum* has been described in goats and

sheep (Eleni et al., 2004; González-Warleta et al., 2018; Varaschin et al., 2012),

natural infection of *N. caninum* in small ruminants is not widely reported. Given the

importance of small ruminants within livestock economies and for food security in

Africa (Smith, 2004; Wodajo et al., 2020), further investigation of the patterns and

impact of *N. caninum* infection in small ruminants are warranted.

Pestiviruses were the most commonly diagnosed pathogen associated with abortion

in sheep in this study, but pestiviruses are rarely discussed as priority production-

limiting pathogens in livestock in Africa. Analysis of the six pestivirus sequences

obtained in this study identified at least three different viral lineages associated with

abortion in livestock in northern Tanzania including a novel lineage in sheep. Based

on our analysis of the N^{Pro} coding region, this novel lineage appears to be distinct

from other pestiviruses and is likely to be a new, previously unreported pestivirus

species. Notably, several of the most similar pestivirus species were detected in

sympatric wildlife species including giraffe in Kenya (Genbank accession numbers

U80907 and AY163647). As wildlife interactions with livestock are common in many

areas of Tanzania, we hypothesise that spill-over from wildlife may also be involved

in the epidemiology of this novel pestivirus lineage, which appears to be a

contributing cause of ovine reproductive loss in northern Tanzania.

Serological data from the cross-sectional study confirmed widespread exposure to

BHV-1 in the study area (Fig. 2), and previous studies in this area have

demonstrated an association between seropositivity and cattle abortion (Main, 2019).

However, the evidence for BHV-1 as a cause of abortion in this study is relatively

weak. As latent BHV-1 infection may lead to false positive results by molecular

detection (Newcomer & Givens, 2016), we relied on serology to diagnose infection.

Two cattle showed evidence of BHV-1 seroconversion suggesting recent infection

which may be implicated in the aetiology of abortion in these cases but further data

are needed to evaluate the impact of BHV-1 pathogen on livestock reproduction in

Tanzania and our findings highlight the challenges of attribution in settings where

endemic pathogens may co-exist in cattle herds (Barkallah et al., 2014).

BTV, which causes production losses in livestock in Europe (Gethmann, Probst , &

Conraths, 2020; Nusinovici, Souty, Seegers, Beaudeau, & Fourichon, 2013), was not

detected by RT-qPCR in any samples from livestock abortion events, despite high

BTV seroprevalences detected in all species in the cross-sectional study population.

Serological evidence of infection is widespread East Africa (Abera et al., 2018;

Mulabbi, Ayebazibwe, Majalija, Batten, & Oura, 2013; Toye et al., 2013) and several

studies have also demonstrated viraemia using RT-qPCR assays, reporting infection

prevalence as high as 88.9% in cattle in Kenya (Toye et al., 2013) and 56.0% in

goats in Uganda (Mulabbi et al., 2013). Collectively, these studies suggest that BTV

infection is endemic in livestock in East Africa. However, research indicates that

African breeds of livestock may be relatively resistant to clinical disease associated

with BTV. Studies from southern Africa have indicated that indigenous African

breeds of livestock are less susceptible to clinical disease from BTV infection than

European breeds (Coetzee, Stokstad, Venter, Myrmel, & Van Vuuren, 2012).

Furthermore, transplacental foetal infection is only associated with certain BTV

strains, especially attenuated vaccine strains (Maclachlan, Drew, Darpel, & Worwa,

2009). Our findings supported by these trends in BTV epidemiology and

pathogenesis suggest that BTV is not a major cause of livestock abortions in

northern Tanzania and underscores the need for caution in making inferences about

livestock diseases and causes of abortion in particular based solely on

seroprevalence data.

The approach described in our abortion cohort study enabled us to attribute

approximately 20% of reported abortion events to a specific aetiology. Our success

rate was not dissimilar to the quoted rate of aetiological abortion diagnosis in high

income settings (35%) (Cabell, 2007; Kirkbride, 1993; Wolf-Jäckel et al., 2020)

despite working in a challenging and resource-limited environment. Our approach

was specifically designed to optimise abortion investigations in a resource-limited

setting, where veterinary surveillance may be limited. Our focus on the use of

molecular diagnostics to detect a range of selected pathogens proved to be valuable,

particularly for high priority pathogens such as RVFV. While molecular diagnostic

capability was established for several known abortigenic agents, it was not possible

for our diagnostic panel to be fully comprehensive of all possible aetiologies of

abortion and did not include agents such as *Campylobacter spp.*, *Listeria spp.*, and

Salmonella enterica serovars that are known to be important causes of livestock

abortion worldwide. Non-infectious causes were also not considered for investigation

and these two factors may account for some of the undiagnosed cases.

Although histopathology is conventionally considered the most robust approach for attributing causality in livestock abortion, histopathology was of limited value in this study due to the limited availability and poor quality of tissues samples obtained through field investigations that were often conducted in remote settings. Additional challenges to inference of causality were encountered due to the nature of our study site. In areas such as rural Tanzania, where remote locations, high ambient temperatures and scavenging animals mean that the type and quality of diagnostic samples is often sub-optimal, inference about the cause can be particularly challenging. For example, placental tissues were not available for sampling from many small ruminants, and even where placental tissues were available for collection, most tissue samples showed evidence of significant autolysis, which precluded a definitive diagnosis in some cases. However, despite these sampling limitations, molecular diagnostics provided a wealth of information regarding the presence of abortigenic agents in swab and tissue samples, and therefore we recommend that molecular diagnostics should be included as a core element of abortion surveillance studies to infer aetiology in resource-limited settings.

In considering appropriate criteria for attributing the aetiology of abortion events reported to this study, we became aware of variations in case definitions and the limitations of the evidence base underpinning attribution criteria for livestock abortions. While methods for assigning attribution in situations of imperfect diagnostic testing are well-developed and widely adopted in aetiological studies of human diseases (Knoll et al., 2017), these questions are rarely addressed in veterinary investigations. Attribution was particularly problematic for *C. burnetii*, as

the pathogen is frequently detected in vaginal fluids at parturition in clinically healthy

animals (Hilbert et al., 2012) and environmental contamination of placental material

or foetal surfaces may also lead to false positives in high risk environments (Borel et

al., 2014). In this study, we developed a case definition for *C. burnetii* based on OIE

guidelines (Sidi-Boumedine et al., 2010; World Organisation for Animal Health (OIE),

2018g) but also driven by population-level data generated by this study. Further

development of this case definition would be beneficial to understand how clinical

history, gross pathology (where available), measured bacterial load and other

adjunct diagnostics could be used to more confidently attribute abortion cases both

within and outside the Tanzanian context.

Substantial differences in reporting rates between wards further limits the

representativeness of our sample. Important differences in livestock seroprevalence

between production systems have been observed in northern Tanzania (Herzog et

al., 2019) as well as in other areas of East Africa (Kadohira, McDermott, Shoukri, &

Kyule, 1997). Spatial heterogeneity in livestock exposure risk, with clustering in

seropositivity at the village-level, have also been described for pathogens such as

Brucella (Kairu-Wanyoike et al., 2019). It is therefore possible that sample collection

in more administrative areas, with improved representation of pastoral and

agropastoral livestock keepers, would reveal different frequencies of pathogen

detection than described here. Our data should therefore be considered as only

indicative of the relative importance of different pathogens as causes of livestock

abortion in northern Tanzania as a whole.

A key insight from our study is that, apart from RVF, none of the abortigenic

pathogens identified by this study have been considered important pathogens in

Tanzania, or prioritised for interventions. Gaps remain in our understanding of the

causes of livestock abortion in Tanzania. However, even with testing a limited range

of pathogens, our study has been able to generate data that adds substantially to the

existing evidence base for the development of interventions to reduce the burden of

production losses associated with livestock abortions. Our study further highlights

the value and importance of establishing laboratory-based molecular diagnostics and

developing clear case definitions and diagnostic criteria for use in resource-limited

settings. With laboratory capacity and technical capability expanding rapidly in Africa,

there is enormous potential and an urgent need for the veterinary sector to shift

towards more evidence-based policy and decision-making, with the potential for this

transition to lead to substantial improvements in animal and human health.

Methods and Materials

<u>Research ethics</u>

Ethics approval for this research was granted by Kilimanjaro Christian Medical

Centre (KCMC) Ethics Committee (No. 535 & No. 832); National Institute of Medical

Research (NIMR), Tanzania (NIMR/HQ/R.8a/Vol.IX/1522 &

NIMR/HQ/R.8a/Vol.IX/2028); Research Ethics Coordinating Committee and the

Institutional Review Board for Clinical Investigations of Duke University Health

System in the United States (Pro00037356), University of Otago Ethics Committee

(H15/069 & H17/069); and University of Glasgow College of Medical Veterinary and

Life Sciences Ethics Committee (200140152 & 200170006). Sample shipments

between the UK and Tanzania were performed in accordance with the Nagoya

Protocol on Access to Genetic Resources under Material Transfer Agreements

between KCMC and University of Glasgow, and between KCMC and Moredun

Research Institute (MRI) with authorisation for export and import, respectively, from

the Tanzanian Ministry of Livestock and Fisheries (permit number VIC/ZIS/8582),

and The Scottish Government's Agriculture, Food and Rural Communities

Directorate (TARP(S) 2019/07).

<u>Cross-sectional serological study</u>

Two cross-sectional livestock serological studies were performed from 2013 through

2016, to provide a baseline of exposure to a range of abortigenic pathogens.

Livestock were selected and sampled as previously described (Bodenham, 2020;

Semango et al., 2019) from six districts in Arusha Region, three districts in

Kilimanjaro Region, and four districts in Manyara Region. Serum was separated from

coagulated whole blood samples by centrifugation (≤1,300 x g for 10 mins at room

temperature) and aliquoted to create replicate sets. Samples were transported to the

Zoonoses Laboratory at Kilimanjaro Clinical Research Institute (KCRI), Moshi,

Tanzania, for archiving at -80°C prior to testing. Cattle, goat, and sheep sera were

tested for exposure to *Brucella* spp., *C. burnetii,* and RVFV using commercial ELISA,

as per manufacturer's instructions (Table 3). Cattle sera were additionally tested for

exposure to *Leptospira* Hardjo serovars, BHV-1, and BVDV (Table 3). Absorbance

was measured using a MultiSkan FC plate reader (Fisher Scientific, Leicestershire,

UK). For each sample, one of the duplicate aliquots was heat treated at 56°C for two

hours and archived at -80°C until it was shipped, on dry ice, to MRI or University of

Glasgow, UK, for testing. All or a random selection of available samples were tested

by indirect ELISA for *N. caninum* and *T. gondii*, and *C. abortus* (goats and sheep

only) as described previously (Helmick et al., 2002, Marques et al., 2012, Wilson et

al., 2009). A random selection of cross-sectional sera was also tested for exposure

BTV (Table 3). Clopper-Pearson 95% confidence intervals were calculated using the

binom package (Dorai-Raj, 2014) in R statistical software (version 3.6.0).

Table 3: Cross-sectional and abortion cohort serological tests (ELISA) for

abortigenic pathogens used for various livestock species, northern Tanzania,

2013-2019.

Target pathogen	Livestock species tested			ELISA kit name (manufacturer or source)
	Cattle	Goat	Sheep	
Brucella spp.	✓	✓	✓	*Brucella* COMPELISA (Animal and Plant Health Agency, UK)
Chlamydia abortus	NA	✓	✓	In-house MRI ELISA (Wilson, Livingstone, & Longbottom, 2009)
Coxiella burnetii	✓	✓	✓	PrioCHECK Ruminant Q Fever Ab Plate § (Thermofisher®)
Leptospira Hardjo	✓	NA	NA	Bovine Leptospira Hardjo 5 Plate Solid ELISA kit (Linnodee®)
Neospora caninum	✓	✓	✓	In-house MRI ELISA (Semango et al., 2019) **OR** *Neospora caninum* Indirect ELISA (ID Screen®)
Toxoplasma gondii	✓	✓	✓	In-house MRI ELISA (Hamilton et al., 2015) **OR** Toxoplasmosis Indirect Multi-species ELISA (ID Screen®)

Bluetongue virus (BTV)	✓	✓	✓	Bluetongue competition kit (ID Screen®)
Bovine herpesvirus (BHV-1)	✓	NA	NA	IBR Individual Ab Test (IDEXX®)
Pestiviruses (BVDV/ BDV)	✓	NA	NA	BVDV p80 Ab ELISA (IDEXX®)
Rift Valley fever virus (RVFV)	✓	✓	✓	Rift Valley Fever Competition Multispecies ELISA kit (ID Screen®) **OR** In-house nucleocapsid-based ELISA pre-screen of goat and sheep CS sera (University of Glasgow) (Nyarobi, 2020)

§: Formerly known as LSIVet Ruminant Q fever Serum/Milk
KCRI = Kilimanjaro Clinical Research Institute, Moshi, Tanzania; MRI = Moredun Research
Institute, UK
NA = not applicable, test not done. For *Leptospira,* BHV-1 and BVDV, serological testing of
goat and sheep sera was not performed because the ELISA kits were not validated for small
ruminant sera.
578

<u>Abortion cohort study</u>

A prospective abortion cohort study was undertaken from October 2017 through

September 2019 in 13 wards in Arusha, Kilimanjaro, and Manyara Regions of

northern Tanzania (Fig. 1). Study wards were selected from those included in the

cross-sectional exposure studies. Two wards were selected purposively from among

those included in these cross-sectional studies because of existing relationships with

the livestock-keeping community that were expected to promote participation. The

remaining eleven wards were selected randomly from the wards targeted in the

cross-sectional studies.

Livestock field officers (LFOs), animal health professionals working for the

Tanzanian Ministry of Livestock and Fisheries, were recruited to participate in the

study with one LFO responsible for a single ward. At the start of the study the LFOs attended a training course on causes of livestock abortion, as well as the practical investigation and management of abortion events. Training included the safe collection, storage, and transport of biological samples from the aborting dam, foetus, and placental material, the use of personal protective equipment for the safe handling and disposal of aborted material and methods for data collection. Following the training course, LFOs were asked to disseminate information about the study to the community living within their respective wards, requesting livestock owners to report any instances of livestock abortion to them so that an investigation could be carried out. Following receipt of a report, the LFO was asked to pass on the information to the research coordinator, a veterinarian, for investigation. For the purposes of this study, an abortion event was defined as: farmer reported visual evidence of premature foetal loss or stillbirth in cattle, goats, or sheep. The abortion was considered eligible for inclusion in this study if the research field team or LFO could attend within 72 hours of the event occurring. Two investigation options were available. Where LFOs carried out the investigation, vaginal swabs and 10 ml blood samples were collected in red-top vacutainer tubes (Becton, Dickinson and Company, New Jersey, USA) from the dam(s) that had aborted, as identified by the livestock keeper. Where the veterinarian-led research team was able to join the LFO's within 72 hours, additional samples were collected, including swabs from the surface of the foetus exterior and placental (cotyledon) tissues, when available. Duplicate swab samples (breakpoint shaft, viscose tip; Technical Service Consultants Ltd, Lancashire, UK) and duplicate tissue samples were stored in 1 ml DNA/RNA shield (Zymo Research, California, USA) to inactivate infectious agents and preserve nucleic acid integrity prior to molecular testing. Due to practical

limitations, samples comprehensive sets of tissues including foetal fluids for bacterial

culture were not collected in this study. Tissue samples were also stored in duplicate

in 10% buffered formal saline to preserve tissue structure for histopathology and

IHC. Samples were transported to the Zoonoses Laboratory at KCRI, for archiving at

-80°C prior to testing against a panel of abortigenic agents.

<u>Abortion cohort serological testing</u>

Sera from dams that aborted were tested using the range of host and pathogen

specific assays described in Table 3. For each sample, one of these duplicate tubes

was heat treated at 56°C for two hours prior to export to the UK for testing. The

remaining tube was retained for testing at KCRI and stored at -80°C. Households

were re-visited four to six weeks later to collect a convalescent blood sample from

the same dam(s). Seroconversion was defined as a negative acute serum with a

positive convalescent serum.

<u>DNA and RNA extraction</u>

RNA was extracted using a RNeasy® Mini kit (QIAGEN, Hilden, Germany) following

manufacturer's instructions. For RNA extraction from the placental tissues, a sterile

scalpel was used to add approximately 25 mg of tissue to 600 µl RLT buffer in a

sterile bead beating tube containing ceramic beads (CK28, Precellys, Bertin

Instruments, France). Bead beating was performed for one minute at 3,450 rpm

using a MiniBeadBeater-16 (BioSpec Products, Inc., Oklahoma, USA). The liquid

was aliquoted into a microcentrifuge tube with 590 µl nuclease-free water and 10 µl

Proteinase K (QIAGEN, Hilden, Germany). For RNA extraction from swabs, 200 µl

sample liquid was added to 600 µl RLT buffer, and 10 µl Proteinase K. All tubes

were then incubated at 55°C for 10 min. Once cooled to room temperature, tubes

were centrifuged at 10,000 x g for three minutes. The supernatant was transferred to

a new microcentrifuge tube where half the volume of ethanol (96-99%) was added to

the tube (e.g., 450 µl ethanol to 900 µl supernatant) and mixed by pipetting up and

down gently. RNA was extracted from 700 µl of the lysate/ethanol mix using a final

elution volume of 50 µl.

DNA was extracted using a DNeasy® blood & tissue kit (QIAGEN), following

manufacturer's instructions. For DNA extraction from placental tissues, a sterile

scalpel was used to add approximately 25 mg of placenta into a microcentrifuge tube

with 180 µl ATL buffer and 20 µl Proteinase K. The tubes were incubated at 56°C,

vortexing occasionally, until the tissue was lysed, after which, 200 µl AL buffer was

added, and the tube was vortexed briefly. For DNA extraction from swabs, 200 µl

sample liquid was added to 200 µl AL buffer with 20 µl Proteinase K and vortexed.

All tubes were incubated at 56°C for 10 min. When the buffer/sample mix had cooled

to room temperature, 200 µl ethanol (96-99%) was added, and the tube was

vortexed. DNA was extracted from the entire volume of the lysate/ethanol mix. Final

elution volumes for swabs and placental tissue samples were 100 µl and 200 µl,

respectively.

All batches of nucleic acid extractions included a negative extraction control where

sample was replaced by nuclease-free water and the run alongside the nucleic acid

extraction process.

<u>Molecular testing</u>

The molecular targets for all pathogens are listed in Table 4. Primer and probe

sequences and cycling conditions for each PCR, qPCR, and RT-qPCR assay, are

detailed in supplementary materials (S1; Table C). DNA or RNA extracts from

positive controls listed in Table 4 were used for individual PCR, qPCR, and RT-

qPCR assays. All available vaginal swab, foetal swab, and placental tissue samples

for all three livestock species were for all pathogens included in this study. Samples

were tested in duplicate except for pathogens tested using commercial kits (BTV and

pestiviruses) where single samples were tested. Nuclease-free water and extraction

controls were included in each PCR, qPCR, and RT-PCR run as negative controls.

Samples were considered positive only if negative extraction controls and negative

controls showed no evidence of amplification and the internal and external positive

controls also amplified. Where single wells amplified for samples tested in duplicate,

the assay was repeated. Unless specified, testing was performed at KCRI.

Table 4: Abortion cohort study molecular assays (PCR, qPCR and RT-qPCR)

for cattle, goat and sheep samples for each target pathogen, northern

Tanzania, 2017-2019.

Target pathogen	PCR/qPCR/RT-PCR testing		
	Target (source)	Positive control strain (source)	Commercial kit (manufacturer)
Brucella spp.	*IS711* (Matero et al., 2011) **AND** *bcsp31* (Probert, Schrader, Khuong, Bystrom, & Graves, 2004)	*Brucella abortus* strain 16M (Animal and Plant Health Agency, UK)	n/a
Chlamydia abortus	*Omp1*	*C. abortus* strain S26/3 (Moredun	n/a

		Research Institute (MRI), UK	
Coxiella burnetii	*IS1111a* (Roest et al., 2011)	*C. burnetii* Nine-mile strain RSA 493 (Wageningen Bioveterinary Research, Wageningen, Netherlands)	n/a
Pathogenic *Leptospira* spp.	*lipL*32 (Stoddard, Gee, Wilkins, McCaustland, & Hoffmaster, 2009)	*Leptospira interrogans* serovar Copenhagenii Strain Wijnberg (OIE Reference Laboratory for Leptospirosis, Amsterdam, Netherlands)	n/a
Neospora caninum	*ITS1* (Buxton et al., 1998)	DNA extracted from tachyzoites of NC1 strain (MRI, UK)	n/a
Toxoplasma gondii	*ITS1* (Hurtado, Aduriz, Moreno, Barandika, & García-Pérez, 2001)	DNA extracted from tachyzoites of RH strain (MRI, UK)	n/a
Bluetongue virus (BTV)	n/a	RNA provided in kit	Virotype BTV pan/8 RT-PCR Kit (Indical® Bioscience GmbH)
Pestiviruses (BVDV/ BDV)	n/a	RNA provided in kit	Virotype BVDV RT-PCR Kit (Indical® Biosciences GmbH)

Rift Valley fever virus (RVFV)	RV (Drosten et al., 2002)	RVF strain MP12 (MRC-University of Glasgow Centre for Virus Research, Glasgow, UK)	

Previously published qPCR assays designed and validated for the diagnosis of

infection with the respective pathogens were used to detect the presence of *Brucella*

spp. (two gene targets (Matero et al., 2011; Probert et al., 2004)), *C. burnetii* (Roest

et al., 2011), *C. abortus* (Livingstone et al., 2009), and pathogenic *Leptospira* spp.

(Stoddard et al., 2009). These qPCRs were performed in single well reactions using

a QuantiNova® Probe kit (QIAGEN) with 5 μl of DNA template (or 2.5 μl per target

for *Brucella* spp.).

Nested conventional PCR assays for the presence of *Neospora* spp. and

Toxoplasma gondii were performed targeting the internal transcribed spacer (ITS1)

region between the 18S and 5.8S rRNA genes (Burrells et al., 2013). The primary

(1°) nested PCR was performed using 2 μl of DNA template. The secondary (2°)

nested PCR was performed using with 2 μl of a 1:5 diluted primary PCR product.

Both 1° and 2° assays were performed with 2x PCR master mix (Promega

Corporation, Wisconsin, USA). The final PCR product from the 2° nested PCR was

checked by electrophoresis in a 1.5% agarose gel. Samples were considered

positive if the size of bands visible under UV light were congruous with those

expected of the *Neospora* (249 bp) and *Toxoplasma* (227 bp) targets.

The presence of BTV and pestivirus RNA in samples was determined by RT-qPCR

using commercial kit-based assays validated for the detection of the respective

pathogens (Table 4). Single samples were tested, as per kit instructions, using 5 μl

of RNA template. Finally, the presence of RVFV RNA was assessed using the

QuantiNova® RT-PCR probe kit (QIAGEN). Samples were tested in duplicate using

5 μl of RNA template per well.

<u>Pestivirus speciation</u>

As commercial kit-based RT-qPCR assays for pestiviruses do not distinguish

between the different pestivirus species, genotyping was performed at MRI to

determine the infecting pestivirus species. The RNA extracts that were positive in

pan-pestivirus RT-PCR screening assays were subjected to RT-PCR and then

nested PCR using four primer pairs developed for genotyping of livestock

pestiviruses: 324-326 (5'UTR); BD1-BD3/BD4 (N^{pro}); BD1-BD2 (N^{pro}-C); and 17f-

1400r (UTR-Erns) to determine the infecting virus type (Becher et al., 1997; Vilcek,

Nettleton, Paton, & Belák, 1997; Vilcek et al., 2001). RT-PCR products of the

expected sizes were purified from agarose gels and submitted for sequencing using

the relevant PCR primers (Eurofins-MWG). Sequence traces for each sample were

assembled using SeqMan Pro (DNASTAR Lasergene) and consensus sequences

that excluded the terminal primer sites were generated. The 504 nt N^{pro} coding

sequences were aligned with a set of N^{Pro} reference sequences that included

representatives of pestivirus species A-H plus K, as described previously (Braun et

al., 2019). Alignment was performed using MAFFT (https://mafft.cbrc.jp/alignment/).

Maximum likelihood (ML) phylogenetic analysis was performed using MEGA version

X. Model selection was used to identify the best evolutionary model for the dataset.

The ML analysis was bootstrapped 500 times to estimate the reliability of the

resulting phylogeny.

<u>Additional pathogen genotyping</u>

T. gondii genotyping was also performed to determine the infecting sub-type. The

methods of this genotyping work are detailed in the supplementary materials S1; D.

<u>Pathology and immunohistochemistry (IHC)</u>

Histopathology was performed on placental tissue samples where available. For

samples that tested positive for *C. burnetii* by qPCR, *C. burnetii* IHC was also

performed. The methods for the histopathology and IHC are given in supplementary

materials S1; B.

<u>Diagnosis of abortion aetiology</u>

For the purposes of our study, we determined a set of case definitions that we used

to attribute livestock abortion events to a specific aetiology. These case definitions

are shown in Table 5. Case definitions were based on OIE guidelines for the

diagnosis of each pathogen. Due to the challenges in systematically collecting tissue

samples for histopathology from abortion cases (e.g. lack of available tissue in many

cases), we designed these diagnostic criteria (Table 5) to be independent of

histopathological examination but used evidence from histopathological examination

to support our diagnosis where samples existed to support this diagnostic approach.

For *N. caninum*, for which no OIE diagnostic guidelines exist, the case definition for

was based on criteria used by MRI, UK (Dubey & Schares, 2011), a specialist

diagnostic veterinary laboratory. For *C. burnetii,* OIE guidelines indicate that

definitive diagnosis of infectious abortions resulting from *C. burnetii* infection can

only be robustly achieved at the flock or herd level, but also that bacterial load is

predictive of causality in individual animals (Sidi-Boumedine et al., 2010; World

Organisation for Animal Health (OIE), 2018g). Therefore, to determine the *C. burnetii*

case definitions, we used a two-step approach. First, we used the minimum bacterial

load associated with abortions as stated by the OIE (> 10^4 bacteria per swab or per

gram of tissue, equivalent to Ct values of 37.5 and 42.7 respectively on our PCR

platform). Secondly, using a histogram of the Ct results at the study population-level,

we identified a bimodal distribution in the results, suggesting two distinct biological

processes within our sampled population (see supplementary material S1; E). We

used this frequency distribution, supported by histopathology and IHC results to

select a conservative threshold value using a cut-off Ct value of ≤ 27 for attributing

abortion events to *C. burnetii* in this study.

Confidence in this study's diagnostic case definition was determined to be either

'definitive', 'presumptive' or 'suggestive', depending on how closely our diagnostic

approach aligned with the OIE guidelines for diagnosis of abortions caused by each

pathogen. Where our study diagnostic approach met OIE definitions for a definitive

diagnosis, we classified our diagnosis as 'definitive' and had the highest confidence

in our aetiologic diagnosis. Where our study diagnostic approach met OIE definitions

for a presumptive diagnosis or for *N. caninum* where no OIE guidelines exist, we

classified our diagnosis as 'presumptive'. Where our study showed evidence of

recent dam infection by serology, but did not detect the presence of the pathogen

directly in any samples (e.g., for BHV-1 in cattle, or *C. abortus* in small ruminants),

we classified the diagnosis as 'suggestive' indicating a lower level of confidence in

these aetiologic diagnoses.

Table 5: Diagnostic criteria used by this study to attribute the aetiology of abortions in cattle, sheep, and goats reported

to the abortion surveillance platform. Diagnostic case definitions apply to all three ruminant species unless otherwise specified.

Pathogen	Were OIE diagnostic guidelines available?	Summary of OIE diagnostic criteria (presuming recent history of abortion in all cases)	Diagnostic case definition applied in this study	Confidence in diagnostic case definition
Brucella spp.	Yes - for aetiological diagnosis of abortion (World Organisation for Animal Health (OIE), 2018c)	*Definitive diagnosis (dx): Isolation of* Brucella *from abortion material, udder secretions or from tissues removed at post-mortem* *Presumptive diagnosis (dx): Evidence of* Brucella *by the demonstration of* Brucella-*like organisms in abortion material or vaginal discharge using modified acid-fast staining, if supported by serological tests. Polymerase chain reaction (PCR) methods are additional means for detection of the presence of Brucella DNA in a sample.*	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR against both *IS711* (Ct <35) and bcsp31 (Ct <40) targets **AND** Evidence of seropositivity in acute and/or convalescent serum samples from dam	Presumptive
Chlamydia abortus	Yes - for aetiological diagnosis of abortion in small	*Definitive dx: Evidence of purulent to necrotising placentitis with vasculitis by histopathology, and demonstration of large number of the organism in*	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR (Ct < 40)	Definitive

	ruminants (World Organisation for Animal Health (OIE), 2018d)	*placentae or vaginal excretions by PCR or antigen tests or stained smears.* *Presumptive dx: In the absence of samples for direct pathogen detection, a rise in antibody titre (detected by ELISA) but does not occur in every case.*	**OR** In the absence of a vaginal swab or placental sample, evidence of seroconversion between acute and convalescent serum samples from dam (goats and sheep only)	Suggestive
Coxiella burnetii	Yes - for aetiological diagnosis of abortion at the herd or flock level only (World Organisation for Animal Health (OIE), 2018g)	*Presumptive dx (individual animal):* *Positive identification of high levels of C. burnetii >10^4 per vaginal swab or per gram of placental tissue (Sidi-Boumedine et al., 2010; Touratier et al., 2007)] by PCR in placenta or vaginal discharges.*	Placental tissue, vaginal swab or foetal swab positive by qPCR by *IS1111a* qPCR (Ct ≤ 27)*	Presumptive
Pathogenic *Leptospira* spp.	Yes - for diagnosis of dam infection only (World Organisation for Animal Health (OIE), 2018f)	*Definitive dx: Isolation of* Leptospira *(recommended) or demonstration by PCR (suitable method) from blood or milk* *Presumptive dx: demonstration of agent (isolation or PCR) in the genital tract or vaginal fluids in association with compatible clinical signs and evidence of serological exposure*	Vaginal swab, foetal swab and/or placental tissue samples positive by qPCR (Ct < 40) **OR** Evidence of seroconversion between acute and convalescent serum samples from dam (cattle only) (Ellis, 2015)	Presumptive

Neospora caninum	No	N/A	Vaginal swab, foetal swab and/or placental tissue samples positive by PCR (Dubey & Schares, 2006)	Presumptive
Toxoplasma gondii	Yes - for aetiological diagnosis of abortion (World Organisation for Animal Health (OIE), 2018i)	*Definitive dx: Demonstration of* Toxoplasma *in the placenta of an aborted foetus by either histopathology or PCR*	Vaginal swab, foetal swab and/or placental tissue samples positive by PCR (Innes et al., 2009)	Definitive
Bovine herpes virus-1 (BHV-1)	Yes - for aetiological diagnosis of abortion (World Organisation for Animal Health (OIE), 2018e)	*Definitive dx: Demonstration of BHV-1 by PCR or virus isolation from abortion material.* *Presumptive dx: Demonstration of BHV-1 by PCR or virus isolation from swabs from the genital tract or evidence of seroconversion between acute and convalescent samples confirming acute infection demonstrated by ELISA or virus neutralisation.*	Evidence of seroconversion between acute and convalescent serum samples from dam (cattle only)	Suggestive
Bluetongue virus (BTV)	Yes - for aetiological diagnosis of abortion (World Organisation for Animal	*Definitive dx: Detection of pathogen by RT-PCR or classical virus isolation in blood or tissue samples including from aborted or stillborn animals.*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40)	Definitive

	Health (OIE), 2018a, 2021)			
Pestiviruses (BVDV or BDV)	Yes - for aetiological diagnosis of abortion (World Organisation for Animal Health (OIE), 2018b)	*Definitive dx: Detection of pathogen by RT-PCR, virus isolation or immunohistochemistry in foetal tissues or fluids.*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40)	Definitive
Rift Valley fever virus (RVFV)	Yes - for aetiological diagnosis of abortion (World Organisation for Animal Health (OIE), 2018h)	*Definitive dx: Requires a combination of at least two positive test results from two diagnostic test methods including detection of pathogen by virus isolation or RT-PCR performed on blood (from dam), tissue from aborted foetuses or other abortion products; positive detection of antibodies (serum); or positive for IgM or IgG ELISA with demonstration of rising titres between paired serum samples collected 2-4 weeks apart.*	Vaginal swab, foetal swab and/or placental tissue samples positive by RT-qPCR (Ct < 40) **AND** Detection of RVFV IgG in acute and/or convalescent serum from dam by RVFV IgG ELISA (World Organisation for Animal Health (OIE), 2018h)	Definitive

782   *See supplementary materials S1; E for further details

Acknowledgements

We would like to thank Rosanne de Jong, Divine Ekwem, Erin Hodgkinson, Karen Main and Jen Jen Yu for assistance with serological testing at the University of Glasgow and Clare Underwood (MRI) for expert immunohistological preparations. We appreciate the huge efforts of our field team, Rigobert Tarimo, Fadhili Mshana, and Hassan Hussein as well as the assistance from District Veterinary Officers and LFOs. Thanks to Paul Johnson (University of Glasgow) for support with data management and curation, and Dassa Nkini (NM-AIST) and Elizabeth Kussaga (KCRI) for administrative support.

Financial support

The abortion cohort study was supported by the Supporting Evidence Based Interventions project, University of Edinburgh (grant number R83537 CH). The cross-sectional studies were supported by US National Institutes of Health-National (NIH) Science Foundation Ecology and Evolution of Infectious Disease program (R01 TW009237) and the UK Biotechnology and Biological Sciences Research Council (BBSRC) (grant numbers: BB/J010367/1, BB/L018926/1). MPD was funded by the Scottish Government. CMH, CH, EAI, FK, ML, GCR and DL are supported by funding from the Scottish Government Rural and Environment Science and Analytical Services Division (RESAS). BJW is funded by the BBSRC.

Supplementary materials:

A: Table A: Cross-sectional and abortion cohort acute serology

B: Histopathology and immunohistochemistry for abortion cohort study including **Table S1B** (Summary of results from histopathologic examination of placental samples collected from aborting ruminant livestock as part of the abortion cohort survey, northern Tanzania) and interpretation and conclusions.

C: Table C: Target pathogen serologic (ELISA) and molecular (PCR/RT-PCR) assay information, including primer/probe sequences and cycling conditions for samples from various livestock species.

D: Genetic characterisation of *T. gondii* isolated from an abortion case

E: Determining qPCR cut-off values for diagnosing *Coxiella burnetii* as the cause of abortion in ruminant livestock in Tanzania.

REFERENCES

Abera, T., Bitew, M., Gebre, D., Mamo, Y., Deneke, Y., & Nandi, S. (2018). Bluetongue disease in small ruminants in south western Ethiopia: cross-sectional sero-epidemiological study. *BMC Res Notes, 11*(1), 112. doi:10.1186/s13104-018-3222-z

Alemayehu, G., Mamo, G., Alemu, B., Desta, H., Tadesse, B., Benti, T., . . . Wieland, B. (2021). Causes and flock level risk factors of sheep and goat abortion in three agroecology zones in Ethiopia. *Front Vet Sci, 8*(27). doi:10.3389/fvets.2021.615310

Allan, K. J., Halliday, J. E. B., Moseley, M., Carter, R. W., Ahmed, A., Goris, M. G. A., . . . Cleaveland, S. (2018). Assessment of animal hosts of pathogenic *Leptospira* in northern Tanzania. *PLoS Negl Trop Dis, 12*(6), e0006444-e0006444. doi:10.1371/journal.pntd.0006444

Anyamba, A., Soebiyanto, R. P., Small, J. L., Linthicum, K. J., Forshey, B., Toolin, C. F., . . . Tucker, C. J. (2018). Rift Valley fever outbreak in East Africa: signature of climate extremes. In: Vol. 2018, pp. GH14A-07. American Geophysical Union, Fall Meeting 2018, abstract #GH14A-07.

Barkallah, M., Gharbi, Y., Hassena, A. B., Slima, A. B., Mallek, Z., Gautier, M., . . . Fendri, I. (2014). Survey of infectious etiologies of bovine abortion during mid- to late gestation in dairy herds. *PLoS ONE, 9*(3), e91549. doi:10.1371/journal.pone.0091549

Barkallah, M., Jribi, H., Ben Slima, A., Gharbi, Y., Mallek, Z., Gautier, M., . . . Gdoura, R. (2018). Molecular prevalence of *Chlamydia* and *Chlamydia*-like bacteria in Tunisian domestic ruminant farms and their influencing risk factors. *Transbound Emerg Dis, 65*(2), e329-e338. doi:10.1111/tbed.12757

Becher, P., Orlich, M., Shannon, A. D., Horner, G., König, M., & Thiel, H. J. (1997). Phylogenetic analysis of pestiviruses from domestic and wild ruminants. *J Gen Virol, 78 (Pt 6)*, 1357-1366. doi:10.1099/0022-1317-78-6-1357

Benkirane, A., Jabli, N., & Rodolakis, A. (1990). [Frequency of abortion and seroprevalence of the principal diseases causing ovine infectious abortion in the area of Rabat (Morocco)]. *Ann Rech Vet, 21*(4), 267-273.

Bhandi, S., Pfukenyi, D. M., Matope, G., Murondoti, A., Tivapasi, M., Ndengu, M., . . . De Garine-Wichatitsky, M. (2019). Brucellosis and chlamydiosis seroprevalence in goats at livestock-wildlife interface areas of Zimbabwe. *Onderstepoort J Vet Res, 86*(1), e1-e9. doi:10.4102/ojvr.v86i1.1670

Bodenham, R. F. (2020). Brucellosis in northern Tanzania: Investigating the epidemiology of human infection and evaluating diagnostic test performance in animal hosts. (PhD), University of Glasgow

Bodenham, R. F., Lukambagire, A.-H. S., Ashford, R. T., Buza, J. J., Cash-Goldwasser, S., Crump, J. A., . . . Halliday, J. E. B. (2020). Prevalence and speciation of brucellosis in febrile patients from a pastoralist community of Tanzania. *Sci Rep, 10*(1), 7081. doi:10.1038/s41598-020-62849-4

Borel, N., Frey, C. F., Gottstein, B., Hilbe, M., Pospischil, A., Franzoso, F. D., & Waldvogel, A. (2014). Laboratory diagnosis of ruminant abortion in Europe. *Vet J, 200*(2), 218-229. doi:10.1016/j.tvjl.2014.03.015

Bouley, A. J., Biggs, H. M., Stoddard, R. A., Morrissey, A. B., Bartlett, J. A., Afwamba, I. A., . . . Crump, J. A. (2012). Brucellosis among hospitalized febrile patients in northern Tanzania. *Am J Trop Med Hyg, 87*(6), 1105-1111. doi:10.4269/ajtmh.2012.12-0327

Braun, U., Hilbe, M., Peterhans, E., & Schweizer, M. (2019). Border disease in cattle. *Vet J, 246*, 12-20. doi:10.1016/j.tvjl.2019.01.006

Budasha, N. H., Gonzalez, J. P., Sebhatu, T. T., & Arnold, E. (2018). Rift Valley fever seroprevalence and abortion frequency among livestock of Kisoro district, South Western Uganda (2016): a prerequisite for zoonotic infection. *BMC Vet Res, 14*(1), 271. doi:10.1186/s12917-018-1596-8

Burrells, A., Bartley, P. M., Zimmer, I. A., Roy, S., Kitchener, A. C., Meredith, A., . . . Katzer, F. (2013). Evidence of the three main clonal *Toxoplasma gondii* lineages from wild mammalian carnivores in the UK *Parasitol, 140*(14), 1768-1776. doi:10.1017/S0031182013001169

Buxton, D., & Henderson, D. (1999). Infectious abortion in sheep. *In Pract, 21*(7), 360-368. doi:10.1136/inpract.21.7.360

Buxton, D., Maley, S. W., Wright, S., Thomson, K. M., Rae, A. G., & Innes, E. A. (1998). The pathogenesis of experimental neosporosis in pregnant sheep. *J Comp Pathol, 118*(4), 267-279. doi:10.1016/S0021-9975(07)80003-X

Cabell, E. (2007). Bovine abortion: aetiology and investigations. *In Pract 29*(8), 455-463. doi:10.1136/inpract.29.8.455 %J

Campero, C. M., Moore, D. P., Odeón, A. C., Cipolla, A. L., & Odriozola, E. (2003). Aetiology of bovine abortion in Argentina. *Vet Res Commun, 27*(5), 359-369. doi:10.1023/A:1024754003432

CDC. (2017). Workshop summary. One health zoonotic disease prioritization for multisectoral engagement in Tanzania. https://www.cdc.gov/onehealth/pdfs/tanzania-report-508.pdf

Coetzee, P., Stokstad, M., Venter, E. H., Myrmel, M., & Van Vuuren, M. (2012). Bluetongue: a historical and epidemiological perspective with the emphasis on South Africa. *Virolog J, 9*(1), 198. doi:10.1186/1743-422X-9-198

Crump, J. A., Morrissey, A. B., Nicholson, W. L., Massung, R. F., Stoddard, R. A., Galloway, R. L., . . . Bartlett, J. A. (2013). Etiology of severe non-malaria febrile illness in Northern Tanzania: a prospective cohort study. *PLoS Negl Trop Dis, 7*(7), e2324-e2324. doi:10.1371/journal.pntd.0002324

Crump, J. A., Ramadhani, H. O., Morrissey, A. B., Msuya, L. J., Yang, L. Y., Chow, S. C., . . . Kinabo, G. D. (2011). Invasive bacterial and fungal infections among hospitalized HIV-infected and HIV-uninfected children and infants in northern Tanzania. *Trop Med Int Health, 16*(7), 830-837. doi:10.1111/j.1365-3156.2011.02774.x

de Glanville, W. A., Davis, A., Allan, K. J., Buza, J., Claxton, J. R., Crump, J. A., . . . Cleaveland, S. (2020). Classification and characterisation of livestock production systems in northern Tanzania. *PLoS ONE, 15*(12), e0229478. doi:10.1371/journal.pone.0229478

De Vries, A. (2006). Economic Value of Pregnancy in Dairy Cattle. *J Dairy Sci, 89*(10), 3876-3885. doi:10.3168/jds.S0022-0302(06)72430-4

Dorai-Raj, S. (2014). binom: binomial confidence intervals for several parameterizations. *R package version 1.1-1*. https://CRAN.R-project.org/package=binom

Drosten, C., Göttig, S., Schilling, S., Asper, M., Panning, M., Schmitz, H., & Günther, S. (2002). Rapid detection and quantification of RNA of Ebola and Marburg viruses, Lassa virus, Crimean-Congo hemorrhagic fever virus, Rift Valley fever virus, dengue virus, and yellow fever virus by real-time reverse transcription-PCR. *J Clin Microbiol, 40*(7), 2323-2330. doi:10.1128/jcm.40.7.2323-2330.2002

Dubey, J. P., & Schares, G. (2006). Diagnosis of bovine neosporosis. *Vet Parasitol,*
140, 1-34.
Dubey, J. P., & Schares, G. (2011). Neosporosis in animals - the last five years. *Vet*
Parasitol, 180(1-2), 90-108. doi:10.1016/j.vetpar.2011.05.031
Eleni, C., Crotti, S., Manuali, E., Costarelli, S., Filippini, G., Moscati, L., & Magnino,
S. (2004). Detection of *Neospora caninum* in an aborted goat foetus. *Vet*
Parasitol, 123(3-4), 271-274. doi:10.1016/j.vetpar.2004.06.017
Ellis, W. A. (2015). Animal Leptospirosis. In B. Adler (Ed.), *Leptospira and*
Leptospirosis (Vol. 387). Berlin: Springer.
Essig, A., & Longbottom, D. (2015). *Chlamydia abortus*: New aspects of infectious
abortion in sheep and potential risk for pregnant women. *Curr Clin Microbiol*
Rep, 2(1), 22-34. doi:10.1007/s40588-015-0014-2
FAO. (2011). *World Livestock 2011 – Livestock in food security*. Retrieved from
Rome: http://www.fao.org/3/i2373e/i2373e.pdf
FAO. (2018). *World Livestock: Transforming the livestock sector through the*
Sustainable Development Goals (222 pp. Licence: CC BY-NC-SA 3.0 IGO).
Retrieved from Rome: http://www.fao.org/3/CA1201EN/ca1201en.pdf
Ferreira, S. C. M., Torelli, F., Klein, S., Fyumagwa, R., Karesh, W. B., Hofer, H., . . .
East, M. L. (2019). Evidence of high exposure to *Toxoplasma gondii* in free-
ranging and captive African carnivores. *Int J Parasitol Parasites Wildl, 8*, 111-
117. doi:10.1016/j.ijppaw.2018.12.007
Galal, L., Ajzenberg, D., Hamidović, A., Durieux, M.-F., Dardé, M.-L., & Mercier, A.
(2018). *Toxoplasma* and Africa: One parasite, two opposite population
structures. *Trends Parasitol, 34*(2), 140-154. doi:10.1016/j.pt.2017.10.010
Gethmann, J., Probst , C., & Conraths, F. J. (2020). Economic impact of a
Bluetongue serotype 8 epidemic in Germany. *Front Vet Sci, 7*(65).
doi:10.3389/fvets.2020.00065
Givens, M. D., & Marley, M. S. D. (2008). Infectious causes of embryonic and fetal
mortality. *Theriogenol, 70*(3), 270-285.
doi:10.1016/j.theriogenology.2008.04.018
González-Warleta, M., Castro-Hermida, J. A., Calvo, C., Pérez, V., Gutiérrez-
Expósito, D., Regidor-Cerrillo, J., . . . Mezo, M. (2018). Endogenous
transplacental transmission of *Neospora caninum* during successive
pregnancies across three generations of naturally infected sheep. *Vet Res,*
49(1), 106. doi:10.1186/s13567-018-0601-3
Hamilton, C. M., Kelly, P. J., Bartley, P. M., Burrells, A., Porco, A., Metzler, D., . . .
Katzer, F. (2015). *Toxoplasma gondii* in livestock in St. Kitts and Nevis, West
Indies. *Parasit Vectors, 8*(1), 166. doi:10.1186/s13071-015-0776-7
Hassan, A., Muturi, M., Mwatondo, A., Omolo, J., Bett, B., Gikundi, S., . . . Munyua,
P. (2020). Epidemiological investigation of a Rift Valley fever outbreak in
humans and livestock in Kenya, 2018. *Am J Trop Med Hyg, 103*(4), 1649-
1655. doi:10.4269/ajtmh.20-0387
Hazlett, M. J., McDowall, R., DeLay, J., Stalker, M., McEwen, B., van Dreumel, T., . .
. Cai, H. Y. (2013). A prospective study of sheep and goat abortion using real-
time polymerase chain reaction and cut point estimation shows *Coxiella*
burnetii and *Chlamydophila abortus* infection concurrently with other major
pathogens. *J Vet Diagn Invest, 25*(3), 359-368.
doi:10.1177/1040638713484729
Herzog, C. M., de Glanville, W. A., Willett, B. J., Kibona, T. J., Cattadori, I. M., Kapur,
V., . . . Bjornstad, O. N. (2019). Pastoral production is associated with

increased peste des petits ruminants seroprevalence in northern Tanzania
across sheep, goats and cattle. *Epidemiol Infect, 147*, e242.
doi:10.1017/S0950268819001262

Hilbert, A., Schmoock, G., Lenzko, H., Moog, U., Diller, R., Fröhlich, A., . . . Sprague,
L. D. (2012). Prevalence of *Coxiella burnetii* in clinically healthy German
sheep flocks. *BMC Res Notes, 5*(1), 152. doi:10.1186/1756-0500-5-152

Hireche, S., Ababneh, M. M., Bouaziz, O., & Boussena, S. (2016). Seroprevalence
and molecular characterization of *Chlamydia abortus* in frozen fetal and
placental tissues of aborting ewes in northeastern Algeria. *Trop Anim Health
Prod, 48*(2), 255-262. doi:10.1007/s11250-015-0944-y

Hurtado, A., Aduriz, G., Moreno, B., Barandika, J., & García-Pérez, A. L. (2001).
Single tube nested PCR for the detection of *Toxoplasma gondii* in fetal tissues
from naturally aborted ewes. *Vet Parasitol, 102*(1-2), 17-27.
doi:10.1016/s0304-4017(01)00526-x

Innes, E. A., Bartley, P. M., Buxton, D., & Katzer, F. (2009). Ovine toxoplasmosis.
Parasitology, 136(14), 1887-1894. doi:10.1017/s0031182009991636

Kadohira, M., McDermott, J. J., Shoukri, M. M., & Kyule, M. N. (1997). Variations in
the prevalence of antibody to *Brucella* infection in cattle by farm, area and
district in Kenya. *Epidemiol Infect, 118*(1), 35-41.
doi:10.1017/s0950268896007005

Kairu-Wanyoike, S., Nyamwaya, D., Wainaina, M., Lindahl, J., Ontiri, E., Bukachi, S.,
. . . Bett, B. (2019). Positive association between *Brucella* spp.
seroprevalences in livestock and humans from a cross-sectional study in
Garissa and Tana River Counties, Kenya. *PLoS Negl Trop Dis, 13*(10),
e0007506. doi:10.1371/journal.pntd.0007506

Kirkbride, C. A. (1993). Bacterial agents detected in a 10-year study of bovine
abortions and stillbirths. *J Vet Diagn Invest, 5*(1), 64-68.
doi:10.1177/104063879300500114

Knoll, M. D., Fu, W., Shi, Q., Prosperi, C., Wu, Z., Hammitt, L. L., . . . Zeger, S. L.
(2017). Bayesian estimation of pneumonia etiology: epidemiologic
considerations and applications to the pneumonia etiology research for child
health study. *Clin Infect Dis, 64*(suppl_3), S213-s227. doi:10.1093/cid/cix144

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular
evolutionary genetics analysis across computing platforms. *Mol Biol Evol,
35*(6), 1547-1549. doi:10.1093/molbev/msy096

Livingstone, M., Wheelhouse, N., Maley, S. W., & Longbottom, D. (2009). Molecular
detection of *Chlamydophila abortus* in post-abortion sheep at oestrus and
subsequent lambing. *Vet Microbiol, 135*(1-2), 134-141.
doi:10.1016/j.vetmic.2008.09.033

Loftis, A. D., Reeves, W. K., Miller, M. M., & Massung, R. F. (2012). *Coxiella burnetii*,
the agent of Q fever, in domestic sheep flocks from Wyoming, United States.
Vector Borne Zoonotic Dis, 12(3), 189-191. doi:10.1089/vbz.2011.0760

Maclachlan, N. J., Drew, C. P., Darpel, K. E., & Worwa, G. (2009). The pathology
and pathogenesis of bluetongue. *J Comp Pathol, 141*(1), 1-16.
doi:10.1016/j.jcpa.2009.04.003

Main, K. (2019). *Seroprevalence and risk factors associated with bovine herpesvirus-
1 infection in cattle in northern Tanzania.* (MSc), University of Glasgow,

Matero, P., Hemmilä, H., Tomaso, H., Piiparinen, H., Rantakokko-Jalava, K., Nuotio,
L., & Nikkari, S. (2011). Rapid field detection assays for *Bacillus anthracis*,

Brucella spp., *Francisella tularensis* and *Yersinia pestis. Clin Microbiol Infect,*
1021   *17*(1), 34-43. doi:10.1111/j.1469-0691.2010.03178.x
1022   Maziku, M., Gebru, G., & Stapleton, J. (2017). Livestock health priorities in the
1023   Tanzania livestock master plan. *Tanzania Livestock Master Plan Brief 4.*
1024   Mboera, L. E. G., Kishamawe, C., Kimario, E., & Rumisha, S. F. (2019). Mortality
1025   patterns of toxoplasmosis and its comorbidities in Tanzania: a 10-year
1026   retrospective hospital-based survey. *Front Public Health, 7,* 25.
1027   doi:10.3389/fpubh.2019.00025
1028   Mbotha, D., Bett, B., Kairu-Wanyoike, S., Grace, D., Kihara, A., Wainaina, M., . . .
1029   Lindahl, J. (2018). Inter-epidemic Rift Valley fever virus seroconversions in an
1030   irrigation scheme in Bura, south-east Kenya. *Transbound Emerg Dis, 65*(1),
1031   e55-e62. doi:10.1111/tbed.12674
1032   Mohamed, M., Mosha, F., Mghamba, J., Zaki, S. R., Shieh, W. J., Paweska, J., . . .
1033   Njenga, M. K. (2010). Epidemiologic and clinical aspects of a Rift Valley fever
1034   outbreak in humans in Tanzania, 2007. *Am J Trop Med Hyg, 83*(2 Suppl), 22-
1035   27. doi:10.4269/ajtmh.2010.09-0318
1036   Mulabbi, E. N., Ayebazibwe, C., Majalija, S., Batten, C. A., & Oura, C. A. (2013).
1037   Circulation of bluetongue virus in goats in the Karamoja region of Uganda. *J S*
1038   *Afr Vet Assoc, 84*(1), E1-3. doi:10.4102/jsava.v84i1.922
1039   Munyua, P., Murithi, R. M., Wainwright, S., Githinji, J., Hightower, A., Mutonga, D., . .
1040   . Njenga, M. K. (2010). Rift Valley fever outbreak in livestock in Kenya, 2006-
1041   2007. *Am J Trop Med Hyg, 83*(2 Suppl), 58-64. doi:10.4269/ajtmh.2010.09-
1042   0292
1043   Newcomer, B. W., & Givens, D. (2016). Diagnosis and control of viral diseases of
1044   reproductive importance: Infectious bovine rhinotracheitis and bovine
1045   viral diarrhea. *Vet Clin North Am Food Anim Pract, 32*(2), 425-441.
1046   doi:10.1016/j.cvfa.2016.01.011
1047   Nusinovici, S., Souty, C., Seegers, H., Beaudeau, F., & Fourichon, C. (2013).
1048   Decrease in milk yield associated with exposure to bluetongue virus serotype
1049   8 in cattle herds. *J Dairy Sci, 96*(2), 877-888. doi:10.3168/jds.2012-5800
1050   Nyarobi, J. (2020). The epidemiology of Rift Valley fever in northern Tanzania. (PhD
1051   PhD), University of Glasgow.
1052   Okumu, T. A., John, N. M., Wabacha, J. K., Tsuma, V., & VanLeeuwen, J. (2019).
1053   Seroprevalence of antibodies for bovine viral diarrhoea virus, *Brucella abortus*
1054   and *Neospora caninum*, and their roles in the incidence of abortion/foetal loss
1055   in dairy cattle herds in Nakuru District, Kenya. *BMC Vet Res, 15*(1), 95.
1056   doi:10.1186/s12917-019-1842-8
1057   Oliveira, R. D., Mousel, M. R., Pabilonia, K. L., Highland, M. A., Taylor, J. B.,
1058   Knowles, D. P., & White, S. N. (2017). Domestic sheep show average
1059   *Coxiella burnetii* seropositivity generations after a sheep-associated human Q
1060   fever outbreak and lack detectable shedding by placental, vaginal, and fecal
1061   routes. *PloS one, 12*(11), e0188054. doi:10.1371/journal.pone.0188054
1062   Prabhu, M., Nicholson, W. L., Roche, A. J., Kersh, G. J., Fitzpatrick, K. A., Oliver, L.
1063   D., . . . Crump, J. A. (2011). Q fever, spotted fever group, and typhus group
1064   rickettsioses among hospitalized febrile patients in northern Tanzania. *Clin*
1065   *Infect Dis, 53*(4), e8-e15. doi:10.1093/cid/cir411
1066   Probert, W. S., Schrader, K. N., Khuong, N. Y., Bystrom, S. L., & Graves, M. H.
1067   (2004). Real-time multiplex PCR assay for detection of *Brucella* spp., *B.*
1068   *abortus*, and *B. melitensis. J Clin Microbiol, 42*(3), 1290-1293.
1069   doi:10.1128/jcm.42.3.1290-1293.2004

ProMED-mail. (2018). Rift Valley fever - Eastern Africa: (Tanzania) alert. *ProMED-mail.* Archive Number: 20180619.5863493 Retrieved from http://www.promedmail.org

Reichel, M. P., Wahl, L. C., & Hill, F. I. (2018). Review of diagnostic procedures and approaches to infectious causes of reproductive failures of cattle in Australia and New Zealand. *Front Vet Sci, 5*(222). doi:10.3389/fvets.2018.00222

Rekiki, A., Sidi-Boumedine, K., Souriau, A., Jemli, J., Hammami, S., & Rodolakis, A. (2002). Isolation and characterisation of local strains of *Chlamydophila abortus* (*Chlamydia psittaci* serotype 1) from Tunisia. *Vet Res, 33*(2), 215-222. doi:10.1051/vetres:2002009

Roest, H. I., Ruuls, R. C., Tilburg, J. J., Nabuurs-Franssen, M. H., Klaassen, C. H., Vellema, P., . . . van Zijderveld, F. G. (2011). Molecular epidemiology of *Coxiella burnetii* from ruminants in Q fever outbreak, the Netherlands. *Emerg Infect Dis, 17*(4), 668-675. doi:10.3201/eid1704.101562

Salifu, S. P., Bukari, A. A., Frangoulidis, D., & Wheelhouse, N. (2019). Current perspectives on the transmission of Q fever: Highlighting the need for a systematic molecular approach for a neglected disease in Africa. *Acta Trop, 193*, 99-105. doi:10.1016/j.actatropica.2019.02.032

Samkange, A., Katsande, T. C., Tjipura-Zaire, G., & Crafford, J. E. (2010). Seroprevalence survey of *Chlamydophila abortus* infection in breeding goats on commercial farms in the Otavi Veterinary District, northern Namibia. *Onderstepoort J Vet Res, 77*(1), E1-5. doi:10.4102/ojvr.v77i1.1

Semango, G., Hamilton, C. M., Kreppel, K., Katzer, F., Kibona, T., Lankester, F., . . . de Glanville, W. A. (2019). The sero-epidemiology of *Neospora caninum* in cattle in northern Tanzania. *Front Vet Sci, 6*(327). doi:10.3389/fvets.2019.00327

Seth-Smith, H. M. B., Buso, L. S., Livingstone, M., Sait, M., Harris, S. R., Aitchison, K. D., . . . Thomson, N. R. (2017). European *Chlamydia abortus* livestock isolate genomes reveal unusual stability and limited diversity, reflected in geographical signatures. *BMC Genomics, 18*(1), 344. doi:10.1186/s12864-017-3657-y

Sidi-Boumedine, K., Rousset, E., Henning, K., Ziller, M., Niemczuck, K., Roest, H. I., & Thiéry, R. (2010). Development of harmonised schemes for the monitoring and reporting of Q-fever in animals in the European Union. *EFSA Supporting Publication, 7*(5).

Sindato, C., Pfeiffer, D. U., Karimuribo, E. D., Mboera, L. E. G., Rweyemamu, M. M., & Paweska, J. T. (2015). A spatial analysis of Rift Valley fever virus seropositivity in domestic ruminants in Tanzania. *PLoS ONE, 10*(7), e0131873-e0131873. doi:10.1371/journal.pone.0131873

Smith, T., Godfrey, S. H., Buttery, P. J. (2004). The contribution of sheep and goats in alleviating poverty: communicating messages from research. *Proceedings of the third DFID Livestock Production Programme link project (R7798) workshop for small ruminant keepers*. Izaak Walton Inn, Embu, Kenya, 4-7 February 2003.

Stoddard, R. A., Gee, J. E., Wilkins, P. P., McCaustland, K., & Hoffmaster, A. R. (2009). Detection of pathogenic *Leptospira* spp. through TaqMan polymerase chain reaction targeting the *LipL*32 gene. *Diagn Microbiol Infect Dis, 64*(3), 247-255. doi:10.1016/j.diagmicrobio.2009.03.014

Swai, E. S., & Kaaya, J. E. (2013). A survey of *Toxoplasma gondii* antibodies by latex agglutination assay in dairy goats in northern Tanzania. *Trop Anim Health Prod, 45*(1), 211-217. doi:10.1007/s11250-012-0193-2

Tamura, K., & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol, 10*(3), 512-526. doi:10.1093/oxfordjournals.molbev.a040023

Tolosa, T., Bezabih, D., & Regassa, F. (2010). Study on seroprevalence of bovine brucellosis and abortion and associated risk factor. *Bull Anim Health Prod Afr, 58*(3). doi:10.4314/bahpa.v58i3.64212

Tonouhewa, A. B., Akpo, Y., Sessou, P., Adoligbe, C., Yessinou, E., Hounmanou, Y. G., . . . Farougou, S. (2017). *Toxoplasma gondii* infection in meat animals from Africa: systematic review and meta-analysis of sero-epidemiological studies. *Vet World, 10*(2), 194-208. doi:10.14202/vetworld.2017.194-208

Touratier, A., Baurier, F., Beaudeau, F., Bendali, F., Buret, Y., DeCremoux, R., . . . Dufour, A. (2007). Comment faire le diagno stic d'un élevage cliniquement atteint de fièvre Q? *Recueil des Journées Nationales des GTV, Nantes*, 147-155.

Toye, P. G., Batten, C. A., Kiara, H., Henstock, M. R., Edwards, L., Thumbi, S., . . . Oura, C. A. (2013). Bluetongue and epizootic haemorrhagic disease virus in local breeds of cattle in Kenya. *Res Vet Sci, 94*(3), 769-773. doi:10.1016/j.rvsc.2012.11.001

van den Brom, R., Lievaart-Peterson, K., Luttikholt, S., Peperkamp, K., Wouda, W., & Vellema, P. (2012). Abortion in small ruminants in the Netherlands between 2006 and 2011. *Tijdschrift Diergeneesk, 137*(7), 450-457.

Vanderburg, S., Rubach, M. P., Halliday, J. E. B., Cleaveland, S., Reddy, E. A., & Crump, J. A. (2014). Epidemiology of *Coxiella burnetii* infection in Africa: a OneHealth systematic review. *PLoS Negl Trop Dis, 8*(4), e2787-e2787. doi:10.1371/journal.pntd.0002787

Varaschin, M. S., Hirsch, C., Wouters, F., Nakagaki, K. Y., Guimarães, A. M., Santos, D. S., . . . Langohr, I. M. (2012). Congenital neosporosis in goats from the State of Minas Gerais, Brazil. *Korean J Parasitol, 50*(1), 63-67. doi:10.3347/kjp.2012.50.1.63

Velasova, M., Damaso, A., Prakashbabu, B. C., Gibbons, J., Wheelhouse, N., Longbottom, D., . . . Guitian, J. (2017). Herd-level prevalence of selected endemic infectious diseases of dairy cows in Great Britain. *J Dairy Sci, 100*(11), 9215-9233. doi:10.3168/jds.2016-11863

Vidal, S., Kegler, K., Greub, G., Aeby, S., Borel, N., Dagleish, M. P., . . . Rodriguez-Campos, S. (2017). Neglected zoonotic agents in cattle abortion: tackling the difficult to grow bacteria. *BMC Vet Res, 13*(1), 373. doi:10.1186/s12917-017-1294-y

Vilcek, S., Nettleton, P. F., Paton, D. J., & Belák, S. (1997). Molecular characterization of ovine pestiviruses. *J Gen Virol., 78*(4), 725-735. doi:10.1099/0022-1317-78-4-725

Vilcek, S., Paton, D. J., Durkovic, B., Strojny, L., Ibata, G., Moussa, A., . . . Paifi, V. (2001). Bovine viral diarrhoea virus genotype 1 can be separated into at least eleven genetic groups. *Arch Virol, 146*(1), 99-115. doi:10.1007/s007050170194

Wilson, K., Livingstone, M., & Longbottom, D. (2009). Comparative evaluation of eight serological assays for diagnosing *Chlamydophila abortus* infection in sheep. *Vet Microbiol, 135*(1-2), 38-45. doi:10.1016/j.vetmic.2008.09.043

Wodajo, H. D., Gemeda, B. A., Kinati, W., Mulem, A. A., van Eerdewijk, A., & Wieland, B. (2020). Contribution of small ruminants to food security for Ethiopian smallholder farmers. *Small Rumin Res, 184*, 106064. doi:https://doi.org/10.1016/j.smallrumres.2020.106064

Wolf-Jäckel, G. A., Hansen, M. S., Larsen, G., Holm, E., Agerholm, J. S., & Jensen, T. K. (2020). Diagnostic studies of abortion in Danish cattle 2015–2017. *Acta Vet Scand, 62*(1), 1. doi:10.1186/s13028-019-0499-4

Woods, C. W., Karpati, A. M., Grein, T., McCarthy, N., Gaturuku, P., Muchiri, E., . . . Peters, C. J. (2002). An outbreak of Rift Valley fever in Northeastern Kenya, 1997-98. *Emerg Infect Dis, 8*(2), 138-144. doi:10.3201/eid0802.010023

World Organisation for Animal Health (OIE). (2018a). Bluetongue (infection with bluetongue virus). In OIE (Ed.), Manual of diagnostic tests and vaccines for terrestrial animals.

World Organisation for Animal Health (OIE). (2018b). Bovine Viral Diarrhoea. In OIE (Ed.), Manual of diagnostic tests and vaccines for terrestrial animals.

World Organisation for Animal Health (OIE). (2018c). Brucellosis (*Brucella abortus*, *B. melitensis* and *B. suis*) (infection with *B. abortus, B. melitensis* and *B. suis*). In OIE (Ed.), Manual of diagnostic tests and vaccines for terrestrial animals.

World Organisation for Animal Health (OIE). (2018d). Enzootic Abortion of Ewes (Ovine Chlamydiosis) (Infection with *Chlamydia abortus*). In OIE (Ed.), Manual of diagnostic tests and vaccines for terrestrial animals.

World Organisation for Animal Health (OIE). (2018e). Infectious Bovine Rhinotracheitis/Infectious Pustular Vulvovaginits. In OIE (Ed.), Manual of diagnostic tests and vaccines for terrestrial animals.

World Organisation for Animal Health (OIE). (2018f). Leptospirosis In OIE (Ed.), OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals.

World Organisation for Animal Health (OIE). (2018g). Q Fever. In OIE (Ed.), OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals.

World Organisation for Animal Health (OIE). (2018h). Rift Valley Fever (infection with Rift Valley fever virus). In OIE (Ed.), OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals.

World Organisation for Animal Health (OIE). (2018i). Toxoplasmosis. In OIE (Ed.), Manual of Diagnostic Tests and Vaccines for Terrestrial Animals.

World Organisation for Animal Health (OIE). (2021). Bluetongue. Aetiology, epidemiology, diagnosis, prevention and control references. https://www.oie.int/app/uploads/2021/03/bluetongue.pdf