Information Flow in the Fibroblast Growth Factor Receptor
Communication Channel

José Díaz[1*], Gustavo Martínez-Mekler[2]

[1]Laboratorio de Dinámica de Redes Genéticas, Centro de Investigación en Dinámica Celular,
Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, México.

[2]Instituto de Ciencias Físicas, Universidad Nacional Autónoma de México, Campus Cuernavaca,
Cuernavaca, Morelos, México.

[3]Centro de Ciencias de la Complejidad, Universidad Nacional Autónoma de México, Ciudad de
México, México

* Correspondence:
José Díaz
biofisica@yahoo.com

Keywords: MAPK, C-myc, Fibroblast Growth Factor, Nonlinear dynamics, Information Theory,
Gene Regulatory Networks

Abstract

In this work we analyze the flow of information through the Fibroblast Growth Factor Receptor
(FGFR) communication channel when different types of signals are transmitted by the MAPK
cascade to the gene regulatory network (GRN) formed by the genes *C-Myc*, *DUSP*, and *Cdc25A*,
which control fibroblast proliferation. We used the canonical mathematical model of the MAPK
cascade coupled to a stochastic model for the activation of the gene regulatory network, subject to
different types of FGF inputs (step, quadratic pulses, Dirac delta, and white noise), in order to
analyze the response of the gene regulatory network to each type of signal, and determine the
temporal variation of the value of its Shannon entropy in each case. Our model suggests that the
sustained activation of the FGFR communication channel with a step of FGF > 1 nM is required for
cell cycle progression and that during the G1/S transition the amount of uncertainty of the GRN
remains at a steady value of ~ 2.75 bits. In this case, the average bit rate of transmission of
information between the GNR and the dimer is zero, indicating that while the fibroblast stimulation
with FGF continues, the G1/S transition does not require an additional interchange of information
between the emitter and the gene regulatory network to be completed. We also found that either low
frequency pulses of FGF or low frequency noise, both with a frequency $f \leq 2.77$ Hz, are not filtered
by the MAPK cascade and can modify the output of the communication channel, i.e., the amount of
the effector proteins c-myc, cdc25A and DUSP. An additional effect of low frequency periodic
signals and noise suggested by our model is the possible blockage of the cell cycle progression
because the threshold value concentration of cdc25A for the G1/S transition is not sustained in the in
the nucleus during the 10 hours that this process lasts. Finally, from our model we can estimate the
capacity of this communication channel in 3 bits, which indicates that is not binary.

1   Introduction

There are three subgroups of the Fibroblast Growth Factor (FGF) family: the canonical FGFs, the intracellular FGFs and the hormonal FGFs. The canonical FGF comprises about 20 isoforms; they exert their action by short range diffusion in the extracellular medium, and can act in a dose-dependent fashion (Böttcher, and Niehrs, 2005). Canonical FGFs are secreted ligands that bind to their specific cell surface receptor belonging to the tyrosine kinase family of FGF receptors (FGFRs). FGFs bind to their receptors with different affinity (k_d) values that ranges from 6.19 x 10^{-8} M to 1.65 x 10^{-7} M (Mohammadi et al., 2005; Gillespie et al, 1989). The four FGFR isoforms are very similar in their global structure with an extracellular-ligand-binding domain, a single transmembranal domain and an intercellular tyrosine kinase domain (Figure 1a). The extracellular part of the receptor has two or three immunoglobulin (Ig)-like domains, and a heparin-binding domain which is important for the interaction with the ligand. This extracellular domain suffers multiple alternatives splicing, which modulates the affinity of the receptor to its ligand. Between the extracellular domains D1 and D2 (Figure 1a) there is a consecutive file of acidic amino acids building up the acidic box, which is an exclusive feature of these receptors (Böttcher and Niehrs, 2005; Mohammadi, et al., 2005). The acidic box is connected to the ligand binding portion of the receptor, which consists of two loops formed by domains D2 and D3. The heparin-binding site is located in de D2 loop. The tyrosine kinase domain of the receptor is located in the cytoplasm (Figure 1a).

Figure 1.- FGFR communication channel. a) The four FGFR isoforms are very similar in their global structure with an extracellular-ligand-binding domain, a single transmembranal domain and an intercellular tyrosine kinase domain. b) After the cross-phosphorylation of the FGF-FGFR complexes, the chemical signal generated by the external FGF concentration is transmitted to the small G-protein Ras through the docking proteins Shc, Grb2 and SOS. The activation of Ras is regulated by GAP, which is activated by the (FGF-FGFR)$_2$ complex. Ras transmits the signal emitted by dimeric FGFR downstream through a MAPK cascade in which the MAPKKK (MAP-Kinase-Kinase-Kinase) is Raf1. Activated Raf specifically phosphorylates either MEK1 or MEK2, which is a MAPKK (MAP-Kinase-Kinase). MEK phosphorylates tyrosine and threonine residues of a MAPK (MAP-Kinase), which can be either ERK1 or ERK2; a fraction of ERK is translocated into the nucleus and activates genes such as *C-Myc*, *DUSP*, and *Cdc25A* that control the cell cycle. This series of coupled chemical processes form the FGFR communication channel in which FGF is the input signal, the (FGF-FGFR)$_2$ dimer is the emitter and encoder, the MAPK is the transmitter, the GRN is the receiver and decoder, and the new synthesized proteins are the effectors or output.

When FGF binds to its specific receptor (FGFR) with tyrosine-kinase activity the dimeric complex
$(FGF-FGFR)_2$ is formed. Figure 1b shows that $(FGF-FGFR)_2$ phosphorylates the docking Sh2-Sh3
adaptor protein Shc, which binds to the phosphorylation sites Y653 and Y654 of the dimeric
complex. Also, $(FGF-FGFR)_2$ phosphorylates GTP-ase Activating Protein (GAP), which binds to the
phosphorylation sites Y653 or Y654 of the dimeric complex (Klint, and Claesson-Welsh, 1999).
After the cross-phosphorylation of the FGF-FGFR complexes, the chemical signal generated by the
external FGF concentration is transmitted to the small G-protein Ras through the docking proteins
Shc, Grb2 and SOS (Figure 1b). The activated state of Ras is regulated by GAP, which is activated
by the $(FGF-FGFR)_2$ complex. Once Ras is activated, the signal is transmitted downstream through
the MAPK cascade in which the MAPKKK (MAP-Kinase-Kinase-Kinase) is Raf1. This
serine/threonine kinase is translocated to the inner side of the cell membrane where is activated by
Ras, in presence of phosphatidic acid. Activated Raf specifically phosphorylates either MEK1 or
MEK2, which is a MAPKK (MAP-Kinase-Kinase). MEK phosphorylates tyrosine and threonine
residues of a MAPK (MAP-Kinase), which can be either ERK1 or ERK2; a fraction of ERK is
translocated into the nucleus and activates genes such as *C-Myc* (Garrington and Johnson,1999;
Arkun and Yasemi, 2018), *DUSP* (Chappell et al., 2013) and *Cdc25A* (Dang, 1999).

C-Myc gene belongs to the family of *Myc* genes that includes *B-Myc*, *L-Myc*, *N-Myc*, and *S-myc*. *C-*
Myc has a central role in the regulation of cell proliferation. This gene is located on chromosome 8 in
humans and is composed by three exons. The transcription factor c-myc, product of *C-Myc*, has a
transactivation domain within its first N-terminal 143 amino acids, and two canonical DNA binding
sites or E-boxes (5'-CACGTG-3'). Serine 62 of protein c-myc is phosphorylated by ERK to form a
heterodimer with Max protein in the fibroblast nucleus (Sears, 2004). The heterodimer c-myc/Max is
also capable of binding to the DNA E-box motif, competing with the monomer of c-myc for the
target genes binding sites (Dang, 1999). The heterodimer c-myc/Max bins to the promoter of *DUSP*
gene whose protein dephosphorylates ERK, inhibiting its catalytic activity and closing a negative
feedback loop (Chappell et al., 2013) (Figure 1b).

Another target gene of the c-myc/Max heterodimer is the gene *Cdc25A* whose protein product is the
phosphatase cdc25A. *Cdc25A* is required for the transition from G_1 phase to S phase during the cell
cycle. In this process, cdc25A removes the inhibitory phosphate groups from threonine and tyrosine
residues of G1/S cyclin-dependent kinases CDK4, CDK2 and Cdk1, activating them and allowing
cell cycle progression (Figure 1b).

A central problem in the analysis of the codification and transmission of information in this kind of
signaling networks *is how the external inputs are correctly coded and decoded to produce specific*
gene responses (Díaz and Álvarez-Buylla, 2009). The solution to this question is quit complex
because external inputs generate information that flows through the intricate signaling network of
target cells. Furthermore, this signaling network (SN) modulates the set of signals that are transmitted
to the gene regulatory network (GRN) in a nonlinear form. Both GRN and SN constitute the
nonlinear communication channel (CC) of the cell (Zaynab et al, 2015). In this particular case the CC
is formed by FGFR, the set of docking proteins Sh2-Sh3, the GAP-Ras circuit, the MAPK cascade,
and *C-Myc* with its targets genes. The input to this CC is the extracellular concentration of FGF
(Figure 1b).

The FGF CC can be represented with a directed graph formed by *nodes* (genes and proteins that form
the CC), and *links* that are the regulatory interactions between the nodes. In a GRN, genes interact
between them by regulatory proteins named transcription factors (TFs), which bind to the promoter
sites of target genes to promote or inhibit their transcription. The GRN associated to the FGFR is
shown in Figure 1b, in which the arrow represents activating interactions and the bars represent

inhibitory interactions. The current state of expression of a gene at time *t* depends on the combination
of activating and inhibitory interactions that act on it. The flow of information between the cell and
its environment determines the combination of activated and inhibited genes of a GRN at a particular
time *t*, which is the network state in response to a specific stimulus or *input*.

Although a great number of theoretical studies about the dynamics of the MAPK cascade has been
published (Kholodenko and Birtwistle, 2009), and the response of ERK to different kind of signals
has been tested (Grabowski et al., 2019), there is a lack in our knowledge of how the GRN of the
fibroblast activated by the FGFR CC respond to different classes of time-dependent input signals,
and how the information flow occurs in each case. Furthermore, cell cycle regulation and cell
differentiation show high sensitivity to the amplitude and duration of input signals from growth
factors (Kholodenko and Birtwistle, 2009), indicating that gene activation can show a broad range of
dynamical responses to time-dependent signals, i.e., the FGFR CC is not only a system of
information relay but also a system of information processing used by cells to compute the most
accurate gene response to a specific noisy input signal (Díaz and Martínez-Mekler, 2005).

In the present theoretical work, we analyze the response of the GRN formed by the genes *C-Myc*,
DUSP and *Cdc25A* to different kinds of FGF input signals (step, quadratic pulses, Dirac delta, and
noise) using the tools from the information theory in order to understand how the FGFR CC
processes the information from the extracellular medium. To reach this objective, we calculate the
Shannon entropy of the emitter and the GRN, and the mutual information between the emitter and the
GRN. This analysis also clarifies some important clues on how the FGF regulates the cell cycle
progression under noisy conditions.

2   Model

2.1   Stochastic model of a GRN

We define $p_i^{on}(t)$ as the probability that the gene *i* of a GRN of *r* interacting genes ($i = 1, 2,…, r$) is
expressed (state *on*), and $p_i^{off}(t)$ as the probability that the gene *i* is not expressed (state *off*) in
response to an input. We assume that the expression or non-expression of the gene *i* under the action
of an input *X* is a Bernoulli process and $p_i^{on}(t) + p_i^{off}(t) = 1$.

We propose that the time evolution of $p_i^{on}(t)$ is given in terms of the following master equation
(Mousaviana et al, 2016):

172
$$\frac{dp_i^{on}(t)}{dt} = k_i^{on} c_i p_i^{off}(t) - k_i^{off} p_i^{on}(t) \qquad 0 \le p_i^{on}(t) \le 1, \quad t \in [0,\infty)$$
$$p_i^{off}(t) = 1 - p_i^{on}(t)$$
(1)

where c_i is the concentration of TF molecules in nucleus at time *t*, k_i^{on} and k_i^{off} are the rate constants
for the transition between the expression states.

In consequence, the Shannon entropy at time *t* due the activation of gene *i* is:

$H_i(t) = -p_i^{on}(t)\log_2 p_i^{on}(t) - p_i^{off}(t)\log_2 p_i^{off}(t) \quad 0{<}p_i^{on}(t) \le 1 \quad t \in [0,\infty)$ (2)

and:

$H_i(t) = 0$ when $p_i^{on}(t) = 0$ or $p_i^{off}(t) = 0$ (3)

In this form the total Shannon entropy at time t due to the current state of activation of the GNR is:

$$H_Y(t) = \sum_{i=1}^{r} H_i(t) = - \sum_{j \in \{on,off\}} \sum_{i=1}^{r} p_i^j(t) \log_2 p_i^j(t) \quad 0 < p_i^{on}(t) \leq 1 \quad t \in [0,\infty)$$ (4)

which is the *Shannon entropy* of the receiver *Y*, at time *t*.

In similar way, if $\alpha(t)$ is the concentration of active receptors at the cell membrane and α_T is the total
concentration of receptors, the probability that a receptor molecule is in its active state at time *t* can
be approximated for a high number of molecules as:

$$p_{Emitter}^{on}(t) \approx \frac{\alpha(t)}{\alpha_T}$$ (5)

where the total number of receptors is constant at any time.

Thus, the Shannon entropy at the emitter *X* at time *t* is:

188
$$H_X(t) = -p_{Emitter}^{on}(t) \log_2 p_{Emitter}^{on}(t) - p_{Emitter}^{off}(t) \log_2 p_{Emitter}^{off}(t) \quad 0 < p_{Emitter}^{on}(t) \leq 1 \quad t \in [0,\infty)$$
                                                                                                    (6)
$$H_X(t) = 0 \text{ when } p_{Emitter}^{on}(t) = 0 \text{ or } p_{Emitter}^{off}(t) = 0$$

189   2.2   Model of the MAPK cascade

The model presented in this section corresponds to the canonical MAPK cascade without feedback
between the cascade components, shown in Figure 1b, and without crosstalk with other components
of the SN.

The first step in the cascade activation occurs as an effect of agonist stimulation with external FGF.
The binding of FGF to its receptor form the complex FGF-FGFR, and the cross phosphorylation of
two FGF-FGFR forms the dimeric complex (FGF-FGFR)$_2$:

196
$$FGF + FGFR \underset{k_{-1}}{\overset{k_1}{\rightleftharpoons}} FGF - FGFR$$
                                                                              (7)
$$2(FGF - FGFR) \overset{k_2}{\longrightarrow} (FGF - FGFR)_2 \overset{k_3}{\longrightarrow} \phi$$

where ϕ indicates the internalization of the (FGF-FGFR)$_2$ complex for degradation.

Denoting the amount of FGF as *F*, the amount of free FGFR as *R*, the amount of FGF-FGFR
complexes as *C*, and the amount of (FGF-FGFR)$_2$ complexes as C_2 we obtain the differential
equations:

$$\frac{dC}{dt} = k_1 F \cdot R - k_{-1} C \qquad (8)$$

$$\frac{dC_2}{dt} = k_2 C^2 - k_3 C_2 \qquad (9)$$

where k_1, k_{-1}, k_2, and k_3 are rate constants (see Table 1),

In this set of equations, we denote the total amount of FGF applied to the cell at time t as F_T. We
assume that the total amount of FGFRs (denoted by R_T) is a constant, i.e., that the process of FGFR
recycling is in a stationary state during the time of simulation. Thus, we can write the mass balance
equations:

$$F_T = F + C$$
208 $$\qquad (10)$$
$$R_T = R + C$$

Substituting Eq. (10) in Eq. (8) we finally obtain:

$$\frac{dC}{dt} = k_1 \left(F_T - C\right)\left(R_T - C\right) - k_{-1} C \qquad (11)$$

The formation of the (FGF-FGFR)$_2$ dimeric complex leads to the activation of SOS through the
activation of the docking proteins Shc and Grb2 (see Figure 1). We denote the total amount of SOS
as SOS_T, the amount of the activated complex Shc-Grb2 as C_{Grb2}, the amount of free SOS as sos, and
the amount of activated SOS as $sos*$. If we assume that $C_{Grb2} \propto C_2$, we obtain the new equation:

$$\frac{dsos^*}{dt} = k_4 C_2 \cdot \left(SOS_T - sos^*\right) - k_{-4} sos^* \qquad (12)$$

where k_4, and k_{-4} are rate constants (see Table 1).

GAP and Ras proteins form a phosphorylation cycle in which activated SOS promotes the exchange
of the GDP bonded to the small G protein Ras for a GTP molecule, while the GAP protein
hydrolyses the GTP to GDP. GAP protein is activated by the (FGF-FGFR)$_2$ dimeric complex
according to the equation:

$$\frac{dgap^*}{dt} = k_5 C_2 \cdot \left(GAP_T - gap^*\right) - k_{-5} gap^* \qquad (13)$$

where $gap*$ is the amount of activated GAP, GAP_T is the total of GAP in the cell, k_5 and k_{-5} are rate
constants (see Table 1).

In this form, the rate of activation of Ras is given by the equation:

$$\frac{dras^*}{dt} = \frac{k_7 sos^* \cdot (ras_T - ras^*)}{(ras_T - ras^*) + 0.1} - \frac{k_{-7} gap^* \cdot ras^*}{ras^* + 0.1} \qquad (14)$$

where *ras** is the amount of activated Ras, *ras$_T$* is the total amount of Ras in the cell, k_7 and k_{-7} are
rate constants (Klipp et al., 2009)(see Table 1).

Activated Ras targets a MAPKKK that in this case is Raf, the corresponding equation for the rate of
activation of Raf is:

$$230 \quad \frac{draf*}{dt} = V_{a1}ras* \cdot \left(raf_T - raf*\right) - \frac{V_{a2}raf*}{K_{a2} + raf*} \quad (15)$$

231 where *raf** is the amount of phosphorylated Raf, *raf$_T$* is the total amount of Raf in cell, k_8, k_{-8} and V_{a2}
232 are rate constants, and K_{a2} is also a constant (see Table 1) (Kholodenko, 2000).

233 Activated Raf initiates the sequence of events of the MAPK cascade that ends with the double
234 phosphorylation of Erk (Kholodenko, 2000) (Figure 1):

$$235 \quad \frac{dmek*}{dt} = v_3 + v_5 - v_4 - v_6 \quad (16)$$

$$236 \quad \frac{dmek**}{dt} = v_4 - v_5 \quad (17)$$

$$237 \quad \frac{derk*}{dt} = v_7 + v_9 - v_8 - v_{10} \quad (18)$$

$$238 \quad \frac{derk**}{dt} = v_8 - v_9 - D \cdot erk** \quad (19)$$

where *mek** and *mek*** are the amount of phosphorylated and double-phosphorylated Mek,
respectively. In Eqs. 18 and 19, *erk** and *erk*** are the amount of phosphorylated and double-
phosphorylated cytoplasmic Erk, respectively. *D* is the constant of transport of double-
phosphorylated Erk into the nucleus.

The variables *v$_3$* to *v$_{10}$* in Eqs. 16-19 are the rates of the phosphorylation processes of the kinases in
the cascade (Kholodenko, 2000):

$$245 \quad v_3 = \frac{k_8 \cdot raf* \cdot mek}{K_{a3} + mek}$$

$$246 \quad v_4 = \frac{k_9 \cdot raf* \cdot mek*}{K_{a4} + mek*}$$

$$247 \quad v_5 = \frac{V_{a5} \cdot mek**}{K_{a5} + mek**}$$

$$248 \quad v_6 = \frac{V_{a6} \cdot mek*}{K_{a6} + mek*}$$

$\quad v_7 = \dfrac{k_{10} \cdot mek** \cdot erk}{K_{a7} + erk}$

$\quad v_8 = \dfrac{k_{11} \cdot mek** \cdot erk*}{K_{a8} + erk*}$

$\quad v_9 = \dfrac{V_{a9} \cdot erk**}{K_{a9} + erk**}$

$\quad v_{10} = \dfrac{V_{a10} \cdot erk*}{K_{a10} + erk*}$

the value of the parameters and rate constants of these equations are the reported in Table 1.

Finally, the mass-balance equations for Eqs. 16-19 are:

$\quad mek = mek_T - mek* - mek** \quad (20)$

$\quad erk = erk_T - erk* - erk** \quad (21)$

where *mek* is the amount of non phosphorylated Mek, and *mek$_T$* is the total amount of Mek in the
cell; similarly, *erk* is the amount of non phosphorylated Erk, and *erk$_T$* is the total amount of Erk in the
cell.

The concentration of Erk in the nucleus must be adjusted because the volume of the nucleus (V_n) is
lower than the volume of the cytoplasm (V_{cyt}). In consequence de differential equation for the rate of
variation of the concentration of Erk in the fibroblast nucleus is:

$\quad \dfrac{derkn}{dt} = \left(\dfrac{V_{cyt}}{V_n}\right) D \cdot erk** - \dfrac{k_{12} erkn \cdot dusp}{dusp + 0.1} - k_{13} erkn \cdot cmyc - k_{14} erkn \quad (22)$

where *erkn* is the amount of doubled phosphorylated Erk in the nucleus, *cmyc* is the amount of
protein c-myc, *dusp* is the amount of protein DUSP, k_{12}, k_{13} and k_{14} are rate constants (see Table 1).
This equation indicates that the rate of variation of the concentration of nuclear Erk is the balance
between the rate of transport of phosphorylated Erk from the cytoplasm, minus the rate of interaction
of the nuclear Erk with proteins DUSP and c-myc, and the rate of inactivation of nuclear Erk. In this
equation, DUSP inhibits the activity of Erk in the nucleus closing a negative feedback circuit (Figure
1).

The rate of variation of the concentration of the protein c-myc in the nucleus is the balance between
the rate of translation of the gene *C-Myc* and its rate of degradation:

$\quad \dfrac{dcmyc}{dt} = V_{Cmyc}^{T\max} \cdot p_{Cmyc}^{on}(t) - k_{15} cmyc \quad (23)$

where $V_{Cmyc}^{T\max}$ is the maximum rate of translation of the gene *C-Myc*, $p_{Cmyc}^{on}(t)$ is the probability that
the gene *C-Myc* is active at time *t*, and k_{15} is a rate constant (see Table 1).

The rate of variation of phosphorylated c-myc (*cmyc**) in the nucleus is the balance between its rate
of activation by nuclear Erk, minus its rate of degradation and its rate of interaction with protein Mix:

278
$$\frac{dcmyc*}{dt} = k_{13}erkn \cdot cmyc - k_{16}cmyc* - k_{17}mix \cdot cmyc* \qquad (24)$$

279 where *mix* is the amount of Mix protein (that we assume is a constant in the model), k_{16} and k_{17} are
280 rate constants (see Table 1).

281 The rate of variation of the concentration of the complex (c-myc)-mix (*ft*) is:

282
$$\frac{dft}{dt} = k_{17}cmyc* \cdot mix - k_{-17}ft \qquad (25)$$

283 where k_{17} and k_{-17} are rate constants (see Table 1).

284 In this case, the genes *C-Myc*, *DUSP* and *Cdc25A* form the GRN. The probability of activation of
285 gene *C-Myc* according to Eq. 1 is:

286
$$\frac{dp_{Cmyc}^{on}(t)}{dt} = k_{18}ft\left(1 - p_{Cmyc}^{on}(t)\right) - k_{-18}p_{Cmyc}^{on}(t) \qquad (26)$$

287 where k_{18}, and k_{-18} are rate constants.

288 In a similar form, the probability of activation of the genes *DUSP* and *Cdc25A* are:

289
$$\frac{dp_{DUSP}^{on}(t)}{dt} = k_{19}ft\left(1 - p_{DUSP}^{on}(t)\right) - k_{-19}p_{DUSP}^{on}(t) \qquad (27)$$

290
$$\frac{dp_{Cdc25A}^{on}(t)}{dt} = k_{20}ft\left(1 - p_{Cdc25A}^{on}(t)\right) - k_{-20}p_{Cdc25A}^{on}(t) \qquad (28)$$

291 where k_{19}, k_{-19}, k_{20} and k_{-20} are rate constants.

292 The rate of production of protein DUSP (whose concentration is denoted *dusp*) is given by the
293 balance between the rate of translation of the gene *DUSP* minus the rate of inhibition of nuclear Erk
294 by DUSP and its rate of degradation:

295
$$\frac{d(dusp)}{dt} = V_{DUSP}^{T\max}(t)\, p_{DUSP}^{on}(t) - \frac{k_{12}erkn \cdot dusp}{erkn + 10} - k_{21}dusp \qquad (29)$$

where $V_{DUSP}^{T\max}$ is the maximum rate of translation of the gene *DUSP*, and k_{21} is a rate constant (see Eq.
16 and Table 1)

Finally, the rate of production of protein cdc25A (whose concentration is denoted *cdc25A*) is given
by the balance between the rate of translation of the gene *Cdc25A* minus its rate of degradation:

$\quad \dfrac{d\left(cdc25A\right)}{dt}=V_{Cdc25A}^{T\max}\left(t\right)p_{Cdc25A}^{on}\left(t\right)-k_{22}cdc25A \qquad (30)$

where $V_{Cdc25A}^{T\max}$ is the maximum rate of translation of the gene $Cdc25A$, and k_{22} is a rate constant (see
Table 1).

In this form, the Shannon entropy of the GRN at time t can be calculated as:

$\quad H_Y\left(t\right)=-\displaystyle\sum_{j\in\{on,off\}}\sum_{i\in S}p_i^j\left(t\right)\log_2 p_i^j\left(t\right) \quad \text{where } S=\left\{Cmyc,DUSP,Cdc25A\right\} \qquad (31)$

and the total amount of Shannon entropy of the GRN in the time interval $(0, \tau)$ of simulation can be
calculated as:

$\quad H_Y^T\left(\tau\right)=-\displaystyle\int_0^\tau H_Y\left(t\right)dt \qquad (32)$

In a similar form, the Shannon entropy of the emitter at time t can be calculated as:

$\quad H_X\left(t\right)=-p_{Emitter}^{on}\left(t\right)\log_2 p_{Emitter}^{on}\left(t\right)-p_{Emitter}^{off}\left(t\right)\log_2 p_{Emitter}^{off}\left(t\right) \qquad (33)$

where $p_{Emitter}^{on}\left(t\right)=\dfrac{C_2\left(t\right)}{R_T}$. The total amount of Shannon entropy of the emitter in the time interval

$(0,\tau)$ of simulation can be calculated as:

$\quad H_X^T\left(\tau\right)=-\displaystyle\int_0^\tau H_X\left(t\right)dt \qquad (34)$

The model was solved using the Euler predictor-corrector method with a time step of 0.05 s. We
simulated different time intervals from 120 to 600 minutes to analyze the dynamics of the respond of
the cascade subject to different inputs as described below. The initial conditions were set to zero for
all the variables of the model, except Raf, Mek y Erk whose initial values were set to 300 nM
(Kholodenko, 2000). Protein Mix was held at the constant value of 5 nM during the time of
simulation. We used the parameter values of Table 1.

3   Results

3.1   Dynamics of the FGFR CC in response to a step of FGF

We analyzed the dynamics of the CC of Figure 1 in response to a step input signal of 4 nM of
amplitude given by the function:

$\quad FGFT=\begin{cases}4 & 0\le t<\infty \\ 0 & t<0\end{cases} \qquad (35)$

Figure 2a shows that FGF produces a steady level of ~ 1 nM of the (FGF-FGFR)$_2$ dimer, which is the
emitter of the signal transmitted by the MAPK cascade. Figure 2b shows that, in turn, the signal from
the emitter produces a steady level of ~ 6 nM of double phosphorylated Erk in the cytoplasm, a

steady concentration of ~ 8 nM of c-myc in the nucleus, and a peak of nuclear double phosphorylated
Erk concentration of 2 minutes of duration with amplitude of 25 nM. This peak is followed by a
steady low concentration of nuclear double phosphorylated Erk of ~ 0.3 nM. Figure 2c shows that all
the genes of the GRN have reach a probability of activation higher than 0.9 2 minutes after the signal
from the emitter is sent. The probability of activation of the emitter reaches a steady value of 0.3 five
minutes after FGF is applied to the cell. The output of the FGFR CC is the temporal distribution of
the concentrations of the proteins c-myc, DUSP and cdc25A. Figure 2d show that under FGF
saturating conditions the output consist of ~ 8.5 nM of c-myc, ~ 7.5 nM of cdc25A and low
concentration of ~ 4.5 nM of DUSP, indicating a probable downregulation of DUSP by double
phosphorylated nuclear Erk. Thus, we expect high amounts of proteins c-myc and cdc25A and a low
concentration of DUSP in the nucleus.

Figure 2.- Dynamic of the GRN in response to a step of FGF. (a) 4 nM applied to the extracellular medium during 600
min produces a steady concentration of ~ 1 nM of the (FGF-FGFR)₂ complex (emitter). (b) The signal from the emitter is
transmitted through the MAPK cascade producing the accumulation of ~ 6 nM of double phosphorylated Erk in the
cytoplasm and ~ 8 nM of c-myc in the nucleus. Nuclear double phosphorylated Erk initially shows a peak of ~ 25 nM of
concentration that last 2 min until its decay to a steady concentration of ~ 0.3 nM. (c) The signal from the emitter
produces a steady distribution of probabilities in which the three genes of the GRN has a probability of activation of ~ 1.
GRN activation shows a time lag of 2.5 min with respect to the start of the signal at the emitter. In contrast, the
probability of activation of the emitter reaches a steady value of ~ 0.3. (d) The output of the FGF CC in response to the
step of 4 nM of FGF consist of a steady distribution of ~ 7.5 nM of c-myc, ~ 7.5 nM of cdc25A and ~ 4 nM of DUSP. We
show only the first 15 min of a total of 600 min of simulation.

3.2 Flow of information in the FGFR CC in response to a step of FGF

Figure 3a shows that the Shannon entropy of the GRN or $H_Y(t)$ has a peak of ~ 2.75 bits, 2.5 minutes after the signal produced by FGF is emitted and decays to 0.5 bits in ~ 4.75 minutes. The peak measures the *maximum uncertainty* in the state of activation of the GRN in response to the signal emitted. The curve of $H_Y(t)$ follows the temporal course of the variation of the concentration of nuclear double phosphorylated Erk with a slower decay due the presence of an additional low rate process. As is shown in Figure 2c, the maximum uncertainty value is reached when the genes of the GRN are not still fully expressed and decay to its minimum value when they are fully activated. In contrast $H_X(t)$ smoothly reaches a steady value of ~ 1 bit five minutes after the activation of the FGF receptor. The total value of $H_Y(t)$ (Eq. 26) after 600 minutes is ~ 1.3×10^4 bits (average bit rate of 21.66 bits/min) while the total value of $H_X(t)$ (Eq. 28) is ~ 3.1×10^4 bits (average bit rate of 51.66 bits/min).

Figure 3.- Information flow in the FGFR CC in response to a step of FGF. a) A step **of** 4 nM applied to the extracellular medium during 600 min produces a peak of 2.75 bits of amplitude and 2.5 minutes of duration in the Shannon entropy of the GRN ($H_Y(t)$) after the signal from the emitter arrives to the nucleus. $H_Y(t)$ slowly decays to a minimum steady value of 0.5 bits. In contrast, $H_X(t)$ reaches a steady value of ~ 1 bit after the stimulation with FGF. In this set of Figures we only show the first 15 minutes of simulation. b) $H_Y(t)$ shows an accumulated value of 1.3×10^4 bits and $H_X(t)$ shows an accumulated value of 3.1 x 104 bits after 600 min of simulation.

3.3 Effect of the FGF concentration on the FGFR CC dynamics

We test the effect of the concentration of FGF (*FGFT*) on the dynamics of the complete FGFR CC. We found that the CC has not response for FGFT < 1.3 nM, showing a threshold value of 1.3 nM. Above this value, the response of the CC increases in intensity as *FGFT* increases in the interval from 1.3 to 1.6 nM, and saturates at 1.6 nM. The time lag of response to the stimulation with FGF decreases as the concentration of the agonist increases, reaching its minimum value when *FGFT* = 4 nM. Thus, the full response of the CC is obtained when FGFT = 4 nM.

Figure 4a shows that 1.5 nM of FGFT produces a steady state concentration of ~ 0.3 of the complex (FGF-FGFR)$_2$, which is the emitted signal (Figure 4a). As a result, the concentration of cytoplasmic double phosphorylated Erk and nuclear c-myc shown in Figure 4b are less than the values shown in Figure 2b, in particular the variation of nuclear double phosphorylated Erk is of less amplitude and more duration that the shown in Figure 2b. Thus, the decrease in the concentration of external FGF

decreases the intensity of the signal carried by the cytoplasmic double phosphorylated Erk into the
nucleus, producing a delay of 10 min in the activation of the GRN and in the time to the full steady
activation of the three genes (~ 15 min) (Figure 4c), which in this case have a low probability of
activation: $p_{Cmyc}^{on}(t) = 0.7$, $p_{Cdc25A}^{on}(t) = 0.5$, $p_{DUSP}^{on}(t) = 0.4$. In consequence, the $H_Y(t)$ curve reaches its
maximum value ~ 10 min post stimulation of the fibroblast cell, and its decaying is slower (Figure
4d) with respect to the curve shown in Figure 3a, reaching a steady value of ~ 2.3 bits 20 min post
stimulation. This result indicates *that the reduction of the concentration of the agonist increments the*
uncertainty in the state of activation of the GRN in the FGFR CC. $H_Y(t)$ has an accumulated value of
~ 7.8 x 10⁴ bits after 600 min of simulation (average bit rate of 130 bits/min), which is much higher
than in the case of a step of 4 nM of FGF (Figure 3b). As expected, the probability of activation of
the emitter is low ~ 0.1, and $H_X(t)$ has a lower steady amplitude than in Figure 3b with an
accumulated value of ~ 1 x 10⁴ bits after 600 min of stimulation (average bit rate of 16.6 bits/min)

393
Figure 4.- Effect of FGF external concentration on the dynamic of the CC. (a) 1.5 nM of FGF applied to the
extracellular medium during 600 min produces a steady concentration of ~ 0.3 nM of the (FGF-FGFR)₂ dimer (emitter).
(b) The signal from the emitter is transmitted through the MAPK cascade producing the accumulation of ~ 0.3 nM of
double phosphorylated Erk in the cytoplasm and ~ 4 nM of c-myc in the nucleus. Nuclear double phosphorylated Erk
initially shows a peak of ~ 0.8 nM of concentration that last 12 min until its decay to a steady concentration of ~ 0.05 nM.
(c) The signal from the emitter produces a steady distribution of probabilities in which the gene *DUSP* has a probability
of activation of ~ 0.4, *Cdc25A* has a probability of activation of ~ 0.5, and *C-Myc* has a probability of activation of ~ 0.7.
GRN activation shows a time lag of 10 min with respect to the start of the signal at the emitter. In contrast, the probability
of activation of the emitter reaches a steady value of ~ 0.1. (d) $H_Y(t)$ shows a peak of 2.75 bits of amplitude and 10
minutes of duration and slowly decays to a steady value of ~ 2.3 bits. In contrast, $H_X(t)$ reaches a steady value of ~ 0.5
bits after the stimulation with FGF. In this set of Figures, we only show the first 15 minutes of a total of 600 min of
simulation.

3.4 Response of the GRN to a quadratic pulse

407 We analyze the dynamics of the GRN subject to a quadratic pulse of external FGF (*FGFT*) given by
408 the function:

$$409 \quad FGFT(t) = \begin{cases} 0 & -\infty < t < 0 \\ 4 & 0 \le t < 60 \\ 0 & 60 \le t < 120 \end{cases} \quad (36)$$

where *FGFT*(*t*) is measured in nM.

Figure 5.- Effect of a quadratic pulse of FGF on the FGFR CC dynamics. (a) FGF applied to the extracellular
medium according to Eq. 36 during 600 min produces a quasi-quadratic pulse response of the (FGF-FGFR)$_2$ dimer
(emitter). (b) The signal from the emitter is transmitted through the MAPK cascade producing a quasi-quadratic pulse
response of cytoplasmic double phosphorylated Erk and c-myc, however the dynamics of the nuclear double
phosphorylated Erk is not altered. (c) The quadratic input signal from the emitter produces a quadratic response in the
probability of activation of the emitter, *DUSP* and *Cdc25A*, but a slow decay in the probability of activation of *C-Myc*. (d)
$H_Y(t)$ shows a peak of 2.75 bits of amplitude and 2 minutes of duration, and a peak of ~ 2.25 bits of amplitude and ~ 150
min of duration after the CC is switched *off*. In contrast, $H_X(t)$ shows a quasi-quadratic response to the input signal. In
Figures (a) and (b) we show only 120 minutes of simulation. In Figures (c) and (d) we show only 250 min of simulation.

In the time interval $t \in (-\infty, 60)$ the response of the CC is identical to the shown in Figures 2a - 2d.
However, when the signal generated by the emitter is switched *off* in the time interval $t \in [60, 120)$

the emitter, the cytoplasmic and nuclear double phosphorylated Erk, and c-myc are turned *off* in ~ 25
min (Figures 5a and 5b). The GNR presents a delay in its response because *Cdc25A* turns *off* in ~ 60
min, *DUSP* turns *off* in ~ 120 min, and *C-Myc* takes about 200 min to switch *off* (Figure 5c).
Unexpectedly, $H_Y(t)$ shows a peak of ~ 2.75 bits of amplitude and ~ 2 min of duration when the CC
is turned *on*, and an additional peak of ~ 2.25 bits of amplitude and ~ 150 min of duration after the
CC is switched *off*. After this second peak $H_Y(t)$ shows slowly decays to zero when t > 100 min.
When the FGF signal becomes zero, the probability of activation of *C-Myc* slowly becomes less than
1, and according to Eq. 31 this gene mainly contributes to the value of $H_Y(t)$ when the other genes of
the GRN shut *off*.

In this form, a quadratic input of FGF produces a quasi-quadratic response in the CC, except in
nuclear double phosphorylated Erk and in the probability of activation of *C-Myc*. The response of the
CC is *quasi-linear* with an output that has a similar form of the input. The presence of an additional
peak in the value of $H_Y(t)$ indicates the possibility of an increase in the amount of uncertainty of the
state of the GRN each time the CC is switched *on* or *off*, i.e., *each time that the CC changes its*
operational state.

3.5   Response of the CC to a train of quadratic pulses

We analyzed the dynamics of the CC subject to a train of quadratic pulses of external FGF with
amplitude of 4 nM, period *T* and duration *d* = *T*. In each pulse the CC is switched on during *T*/2 min
and then is completely switched off during *T*/2 min.

We found that quadratic pulses of FGF with a period $0 < T < 60$ min (high frequency pulses) are
filtered by the MAPK cascade having not effect on the components of the CC downstream Ras
(Figure 6a – 6d). However, quadratic pulses of FGF with a period $T \geq 60$ min produce a train of
quadratic pulses with the same period and a longer duration (Figure 7a – 7d). As $T \to \infty$ the response
converges to the case of a single quadratic pulse discussed in Section 3.4. In this form, the response
of the CC to a train of quadratic pulses is quasi linear for low frequency pulses, i.e., the MAPK
cascade coupled to the GRN behaves like a *low frequency pass filter*. Moreover, the input produces a
series of peaks in the value of $H_Y(t)$ with the same period of the input (Figure 7d). In contrast, $H_X(t)$
show periodical variations in its value of the same period that the input signal as expected.

3.6   Response of the CC to noise

We found that the FGFR CC functions like a *low frequency pass filter*. Traditionally, experiments
and theoretical works (Kholodenko and Birtwistle, 2009; Tka č ik and Walczak, 2011; Benary et al.,
2018) have shown that the ultra sensitivity and all-or-none activation of the MAPK cascade filters
noise. However, we have found that when the MAPK cascade is coupled to a GRN this is not
completely true because the CC allows the transmission of noisy signals of low rate of variation.

Figures 8a and 8b show that a random signal of high rate of variation of the external concentration of
FGF has no effect on the steady concentration of the proteins c-myc, cdc25A, and DUSP, In this
case, variation of external FGF concentration occur randomly every τ minutes, with $0 < \tau < 50$.

Figure 6.- Effect of a train of quadratic pulses with period $T < 60$ min on the CC. (a) Pulses of 4 nM of FGF applied to the extracellular medium every 40 min during 120 min produces a quasi-quadratic pulse periodic response of the (FGF-FGFR)$_2$ dimer (emitter), which does not completely turn *off* at the end of each time interval. (b) The signal from the emitter is transmitted through the MAPK cascade and does not produce any affect on cytoplasmic and nuclear double phosphorylated Erk neither on c-myc. (c) The signal from the emitter is transmitted through the MAPK cascade and does not produce any affect on probability of activation of the genes of the GRN. (d) The signal from the emitter is transmitted through the MAPK cascade and does not produce any affect on $H_Y(t)$. $H_X(t)$ shows a quasi-quadratic response to the input signal. In this set of Figures, we show only 300 of the 600 min of simulation.

However, Figures 8c and 8d show that a random signal of low rate of variation of the external concentration of FGF produces a random variation of the concentration of the proteins c-myc, cdc25A and DUSP, in intervals of random duration, which is a distorted version of the output shown in Figure 7c. In this case, variation of external FGF concentration occurs randomly every τ minutes with $\tau > 50$. As expected, the distribution of the probability of activation of the genes of the GRN varies randomly in response to the input signal, given rise to random fluctuations in the value of $H_Y(t)$ and $H_X(t)$ (Figure 9). Thus, low rate of variation noise can distort the response of the GNR with undesirable consequences to the cell differentiation process of the fibroblast.

Figure 7.- Effect of a train of quadratic pulses of FGF with period $T \geq 60$ min on the CC. (a) Pulses of 4 nM of FGF applied to the extracellular medium every 60 min during 600 min produces a quasi-quadratic pulse periodic response of the (FGF-FGFR)$_2$ dimer (emitter), which completely turns *off* at the end of each time interval. (b) The signal from the emitter is transmitted through the MAPK cascade and produces the periodic variation in the concentration of c-myc, cdc25A, and DUSP proteins, with the same period of the input signal. c) The signal from the emitter is transmitted through the MAPK cascade and produces the periodic variation of the probability of activation of the genes of the GRN, with the same period of the input signal. This Figure shows that the GRN is shut *off* 30 min after the emitter. (d) The signal from the emitter is transmitted through the MAPK cascade and produces a series of peaks in the value of $H_Y(t)$ with the same period of the input. $H_X(t)$ shows a quasi-quadratic response to the input signal. In this set of Figures, we show 300 minutes of simulation.

Figure 8. - Effect of noise on the CC dynamics. (a) Random signals of high rate of variation of the external concentration of FGF. The signal varies every 0.004 min. (b) Noise of high rate of variation produces a steady concentration of proteins c-myc, cdc25A and DUSP. (c) Random signals of low rate of variation of the concentration of FGF. The signal varies every 50 min. (d) Noise of low rate of variation produces random variations in the concentration of the proteins c-myc, cdc25A and DUSP.

500 **Figure 9.- Effect of noise of low rate of variation on the Shannon Entropy of the CC.** Random signals of external
501 concentration of FGF that vary every 50 min produces random variations in the value of $H_Y(t)$ and $H_X(t)$.

3.7   Response of the CC to a Dirac delta

We analyzed the response of the CC to a Dirac delta of FGF given by the function:

$$FGFT = \begin{cases} \infty & t = 0 \\ 0 & t \neq 0 \end{cases} \quad (31)$$

Figure 10a shows the Dirac delta used as input. Figure 10b shows that de response of the CC to this
transitory intense signal is the full transitory activation of the GRN giving rise to the transitory
production of the proteins c-myc, cdc25A and DUSP during ~ 40 min.

Figure 10.- Effect of a Dirac delta of FGF on the CC dynamics. (a) Dirac delta of external FGF. (b) The effect of the
Dirac delta (input) is the transitory production of c-myc, cdc25A and DUSP proteins (output) during ~ 40 min.

4 Discussion

Mathematical models in Biology are representations of reality, which implies the simplification of a
complex biological process into a set of central components, which we assume are the key
determinants of the spatio-temporal dynamics of the system under study (Eftimie, et al., 2016).
Generally, this set of main components form a molecular network, in which the number of nodes and
links, together with the distribution of links between nodes, determine the network dynamics
(Eftimie, et al., 2016; Díaz, 2019). In the particular case of the coupling of a SN with a GRN, the
resultant network can be considered as a molecular CC. The main function of this kind of CC is to
mediate in the transmission of information from the cell surface to the GRN, and generate a specific
genetic response. Because GNRs are control systems, the analysis of the flow of information in this
type of CCs deals with a problem of control and communication (Bi and Noel, 2020; Wiener, 1961).

In this sense, beside the extensive research that has clarified the main features of a great number of
manmade CCs, there are features of the biological CCs that are just beginning to be understood. As
example, in the FGFR CC the coding of information is made using chemical reactions (Tka č ik and
Walczak, 2011). Thus, the input signal is the temporal variation of the external FGF concentration,
which is coded in the number of $(FGF\text{-}FGFR)_2$ complexes formed for each concentration of external
FGF (Figures 3a and 4a). In this form the emitter of the original signal is the $(FGF\text{-}FGFR)_2$ dimer,
which is coupled to the MAPK or transmitter, and the information is coded in the number of
activated dimers at time *t*.

Although the MAPK cascade interacts with other SNs, and the cascade itself has feedback circuits
between its components, we decided not to take into consideration these interactions in our model in
order to know the function of the basic molecular structure of the cascade in the modulation of the
signal transmitted by it. Previous works (Kholodenko, and Birtwistle, 2009), have shown that the
MAPK cascade has ultrasensitivity and filtering properties that reduce the effects of noise on the
signal transmitted by the cascade, increasing the fidelity of the transmission. Thus, the practically all-
or-none activation of the cascade avoids the distortion of the transmitted signal by cytoplasmic noise
(Tka č ik and Walczak, 2011; Vaseghi, 2000). Furthermore, we found that the MAPK cascade also
filters external noisy signals of high rate of random variation (Figures 8a and 8b). However, we
unexpectedly found that the cascade allows the transmission of periodic signals of low frequency

(Figure 7) or signals with low rate of random variation (Figures 8c and 8d), acting like a *low*
frequency pass filter.

In the FGFR CC the receiver of the signal is the *C-Myc* GRN, which is activated by the double
phosphorylated Erk that enters the nucleus (Figures 1 and 3b). In this form, the signal is translated
into a specific duration of the activated state of the target genes *C-Myc*, *DUSP* and *Cdc25A*. Thus, we
proposed a stochastic model of the process of gene transcription instead of a deterministic one (Eqs.
26-28), which we considere a better approach to the form in which gene translation occurs in real
GRNs. In consequence, the input signal is *decoded* into a specific time-dependent probability
distribution of gene activation for each concentration of the emitter:

$$\Omega = \left\{ p_{Cmyc}^{on}\left(t\right), p_{DUSP}^{on}\left(t\right), p_{Cdc25A}^{on}\left(t\right) \right\} \qquad (32)$$

that give rise to a temporal distribution of their respective protein concentrations (Eqs. 23, 29 and
30). These proteins are the *effectors* of the information generated by the input. In resume: *in the*
FGFR CC each external concentration of FGF is coded into the concentration of the (FGF-FGFR)$_2$
dimer, which activates the cascade in an all-or-non manner, transmitting the information from the
dimmer to the nucleus where is decoded into a probability distribution of activation of the genes of
the GRN that finally gives rise to a specific distribution of concentration of the effector proteins
(Díaz and Alvarez-Buylla, 2009; Mousaviana et al., 2016).

A fundamental question arises in this point: is the specific distribution of concentration of these
proteins the correct output for each input signal? The stochastic nature of gene expression produces a
certain level of uncertainty about the real state of activation of the GRN at time *t* that we can measure
with the Shannon entropy $H_Y(t)$ (Eq. 31). This quantity measures the contribution of the lost of
information in the real state of activation of each gene of the GRN at time *t* to the state of activation
of the overall GNR in response to a given input. This lost of information or uncertainty is due to
noise and other molecular processes in which the genes take part. Thus, in our model *we will never*
know the exact state of activation of a GNR, and we propose that the output reflects the *most*
probable specific distribution of concentration of the proteins c-myc, DUSP, and cdc25A in response
to a particular input. This flow of information between the emitter and the GRN to produce a specific
response is measured with the mutual information $I_{(X;Y)}(t)$ between *X* and *Y*.

In order to know the mutual information between the emitter and the GNR in the CC, we assume in
Eq. (31) that:

$$p_i^{on}\left(t\right) = p\left(\frac{Y_i = on}{X = on} \right)\left(t\right)$$

571   (33)

$$p_i^{off}\left(t\right) = p\left(\frac{Y_i = off}{X = on} \right)\left(t\right)$$

where the random variables Y_i = state of activation of gene *i* at time *t* , and *X* = state of activation of
the emitter at time *t*, can take only the values *on* and *off*.

When FGF is applied to the CC, then the emitter is in state *on*, and:

$$H_{Y/X}\left(t\right) = p_{Emitter}^{on}\left(t\right) H_Y\left(t\right) \qquad (34)$$

We calculate the mutual information between the random variables X and Y under FGF stimulation
as:

$\quad I_{(X,Y)}(t) = H_Y^{max}(t) - H_{Y/X}(t) \qquad (35)$

where the value of $H_Y^{max}(t)$ is estimated from the output of the model.

We obtain from Eq. (33):

$\quad I_{(X,Y)}(t) = H_Y^{max}(t) - H_{Y/X}(t) = 1 - p_{Emitter}^{on}(t) H_Y(t) \qquad (36)$

The capacity of a CC measures the ability of the receiver to discriminate between two different
signals and produce the correct output in a noisy channel. The Epidermal Growth Factor Receptor
(EGFR) CC operates in a non binary fashion when a sequence of EGF pulses is applied ($\chi = 1.84$
bits/h), which indicates that this CC does not work in a binary form (Benary et al., 2017; Grabowski
et al., 2019). In similar form, the FGFR CC shows a biphasic response to stimulus of different
constant amplitude indicating that this CC can be binary (Kanodia et al., 2014). In order to
determinate if the FGFR CC is binary we estimate its capacity χ as:

$\quad \chi = \max I_{(X,Y)}(t) \qquad (37)$

Figure 11a shows the graphic of $I_{(X,Y)}(t)$ when a *step* of 4 nM of FGF is applied to the CC. The
maximum value of $I_{(X,Y)}(t)$ is ~ 2.9 bits, thus the capacity of the CC (Eq. (37) is estimated in a value
of ~ 2.75 bits, which means that the FGFR CC is not a binary channel. It is possible that an increase
in the complexity of the GRN coupled to the MAPK cascade enlarges the capacity of the FGFR CC
in this case.

Figure 11.- Mutual information of the FGFR CC. (a) Mutual information graphics of the FGFR CC in response to
step of 4 nM of FGF. The figure shows a maximum value of ~ 2.9 bits, a minimum value of ~ 2.25 bits and a steady value
of ~ 2.3 bits. (b) Mutual information graphics of the FGFR CC in response to a train of quadratic pulses. The figure
shows that in this case the mutual information oscillates between a maximum value of ~ 2.75 bits and a minimum value
of ~ 2.3 bits with a periodicity of $T = 60$ min.

Our results show that c-myc regulates cell cycle progression of the fibroblast (Bretones et al., 2014)
down regulating the nuclear concentration of the dual phosphatase DUSP and enhancing the nuclear
concentration of the phosphatase cdc25A. Although DUSP can dephosphorylate the threonine/serine
and tyrosine residues of nuclear Erk, down regulating its kinase activity, Erk can also phosphorylate
DUSP and down regulate its activity. For the set of parameter values of Table 1, *our model predicts*
the down regulation of DUSP protein activity in the nucleus, and a decrease in the probability of
activation of the DUSP gene (Figures 2, 3 and 4), when the fibroblast is stimulated with FGF.

In contrast, the enhanced concentration of the phosphatase cdc25A and the high probability of
activation of its gene *Cdc25A* (Figure 3) induced by c-myc indicates that the stimulation of the
fibroblast with FGF favours its proliferation. Protein cdc25A acts over the complex cyclin D-cdk4
and cyclin E-cdk2 allowing the G1/S phase transition, and also regulates G2/M transition before its
degradation (Lim and Kaldis, 2013; Shen and Huang, 2012). In this form, the model predicts that
any step signal of FGF above the threshold concentration of ~ 1.3 nM induces cell cycle progression
from phase G1 to phase S, and from phase G2 to M, when DUSP is down regulated by double
phosphorylated nuclear Erk.

The above scenario is supported by the results shown in Figure 5, in which a quadratic pulse of FGF
induces the transitory production of cdc25A and DUSP proteins that possibly stops the transition
from F1 to S due to the inactivation of the *Cdc25A* gene after 100 min. In this second scenario, *an*
intense transitory stimulation of the fibroblast with a quadratic pulse of low duration of FGF above
the threshold value of ~ 1.3 nM is not enough to overcome cell arrest in G1 phase. Experimental
results indicate that cdc25A protein has a half life of ~ 20 min, thus it must be produced continuously
during at least ~ 10 hrs to ensure G1/S transition and during ~ 20 hrs to allow the G2/M transition
(Shen and Huang, 2012; Timofeev et al., 2009). Additionally, an intense stimulation of the fibroblast
with a Dirac delta of FGF produces a transitory response of the GRN of ~ 40 min of duration (Figure
10), which is insufficient to overcome the fibroblast arrest in the G1 phase.

Therefore, the low frequency pass filtering properties of the FGFR CC ensures the continuity of the
cell cycle progression avoiding the effects of periodic regular variations of high frequency (T < 60
min) (Figure 6) or noise of high rate of random variation (Figures 8a and 8b) on the concentration of
c-myc and cdc25A. However, periodic regular variations of low frequency (Figure 7) or noise of low
rate of random variation (Figures 8b and 8c) could obstruct cell cycle progression by continuously
switching *off* cdc25A production.

The filtering property of the FGFR CC can be an important feature to avoid intense and high
frequency interferences on the signals carried by the MAPK cascade to the nucleus ensuring the
fidelity of the information transmitted to the GRN from the (FGFR-FGFR)$_2$ dimer complex.
However, we are using a simplified model of the FGFR CC and possibly there are complementary
molecular mechanisms that damp the effect of low frequency interferences on cell cycle progression,
but the presence of this type of response of the CC to this type of signals is an unexpected result that
opens the possibility of a real mechanism that regulates cell proliferation.

During G1/S progression the emitter and the GNR have a steady mutual information of ~ 2.3 bits
(Figure 11a) with an average bit rate of zero, i.e., the amount of information that the state of
activation of the GRN gives about the state of activation of the (FGFR-FGFR)$_2$ dimer is ~ 2.3 bits,
value that is constant during all the process and measures the mutual nonlinear dependence between
the random variables *X* e *Y*. Thus, once the GNR is activated with a step of 4 nM of FGF the CC
reaches a steady state with a constant level of uncertainty about the correct transmission and

interpretation of the message elicited by the FGF at the cell surface. Unexpectedly, the average bit rate of transmission of information between the GNR and the dimer is zero, indicating that while the fibroblast stimulation with a step of FGF continues the G1/S transition does not require of additional interchange of information between the emitter and GRN to complete the cell cycle progression to phase S.

This scenario changes when the CC is subject to a train of quadratic pulses of 4 nM of FGF with a period of $T = 60$ min. Figure 11b shows a periodic variation in the value of $I_{(X,Y)}(t)$ from 2.4 to 2.9 bits during the interval of time that the CC is switching *off*, while the peaks in the value of $H_Y(t)$ occur just at the moment in which the CC turns *off* (Figure 7d). In this case the average bit rate is ~ 0.2 bits/min that indicates that there is a transitory transference of information between the GRN and the emitter, which probably is the gain in the amount of information about the state of activation of the emitter obtained from the GRN during its turning *off*, these peaks of information probably coincide with the intervals of time in which the G1/S transition is interrupted (Figures 7 and 11b). Figures 8d and 9 clearly show that random noise of low rate of variation completely distorts the flow of information in the CC probably producing the arrest of the cell cycle or an irregular progression of the transition from G1 to S.

5   Conclusions

Cell cycle progression is the central process in cell proliferation and must be rigorously controlled to allow the correct fibroblast tissue replacement, its regeneration after an injury and to avoid cancer. In this process, extracellular FGF triggers the activation of the genes responsible of this control by the activation of the intracellular signaling network that carries on the signal generated by the (FGF-FGFR)2 dimmer, or emitter, to the nucleus. An essential component of this signaling network is the MAPK cascade, which in our work is the transmitter of the signal from the cell surface to the nucleus and closes the FGFR communication channel. The MAPK cascade has properties of ultrasensitivity that traditionally is related to its function as noise filter that ensures a high fidelity of the information transmitted by it. However, most of the works that sustain this affirmation are based on theoretical and experimental models in which the cascade is uncoupled to a GRN. In our work we analyze the flow of information in the FGFR CC when the signals are transmitted by the MAPK cascade to the *C-Myc-DUSP-Cdc25A* GRN, in absence of crosstalk with other components of the signaling network, and we found that low frequency noise (period $T \geq 60$ min) is not filtered and can modify the output of the CC, i.e., the observed amount of the effector proteins c-myc, cdc25A and DUSP is continuously distorted by the noise. An additional effect is that low frequency noise can stop cell cycle progression from G1 phase to S phase because the required threshold value concentration of cdc25A is not continuously sustained in the nucleus during the 10 hours that the G1/S transition last. In consequence, our model suggest that the sustained activation of the FGFR CC with a step of FGF > 1.3 nM, or with periodic pulses of FGF with a period T < 60 min, is required for cell cycle progression. During the G1/S transition the amount of uncertainty at the GRN in response to a step of 4 nM of FGF remains at a steady value of ~ 2.75 bits, value that increases every time that the state of the GRN varies. In this case, the average bit rate of transmission of information between the GNR and the dimer is zero, indicating that while the fibroblast stimulation with FGF continues the G1/S transition does not require of additional interchange of information between the emitter and GRN to complete the cell cycle progression to S phase. From our model, we could estimate the capacity of the FGFR CC in 2.75 bits, value that we think probably changes with the size of the GNR and that indicates that this is not a binary channel.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6   Author Contributions

Both authors contributed equally to this work.

7   Funding

Not funding

8   Acknowledgments

JD thanks Erika Juarez Luna for logistical support.

9   References

Arkun, Y., and Yasemi, M. (2018). Dynamics and control of the ERK signaling pathway: Sensitivity, bistability, and oscillations. *PLoS ONE* 13: e0195513. doi: https://doi.org/10.1371/journal.pone.0195513

Benary M., Nolis I., Blüthgen N., Loewer A. (2018) Information Flow in a Mammalian Signal Transduction Pathway. In: Bossert M. (eds) Information- and Communication Theory in Molecular Biology. Lecture Notes in Bioengineering. Springer, Cham. https://doi.org/10.1007/978-3-319-54729-9_3

Bi, D., and Noel, A. (2020). A Survey of Molecular Communication in Cell Biology: Establishing a New Hierarchy for Interdisciplinary Applications. *arXiv*:2009.00090v1

• Böttcher, R.T., and Niehrs, C. (2005). Fibroblast growth factor signaling during early vertebrate development. *Endocr Rev.* 26: 63-77. doi: 10.1210/er.2003-0040

Bretones, G., Delgado, M.D., and León, J. (2014). Myc and cell cycle control. *Biophys. Acta*, http://dx.doi.org/10.1016/j.bbagrm.2014.03.013

• Chappell, J., Sun, Y., Singh, A., and Dalton, S. (2013). MYC/MAX control ERK signaling and pluripotency by regulation of dual-specificity phosphatases 2 and 7. Genes Dev. 27:725–733. doi: 10.1101/gad.211300.112

Dang, C.V. (1999). c-Myc Target Genes Involved in Cell Growth, Apoptosis, and Metabolism. *Molecular and Cellular Biology* 19: 1-11. doi: 10.1128/mcb.19.1.1.

Díaz, J. (2019). SARS-Cov-2 Systems Biology. *Annals of Systems Biology* 3 (1), 029-032. doi: https://doi.org/10.17352/asb.000009

Díaz, J., and Alvarez-Buylla, E.R. (2009). Information flow during gene activation by signaling molecules: ethylene transduction in *Arabidopsis* cells as a study system. *BMC Systems Biology* 3: 48. doi: https://link.springer.com/article/10.1186/1752-0509-3-48

Eftimie, R. et al. Mathematical Models for Immunology: Current State of the Art and Future
Research Directions (2016). *Bull Math Biol* 78:2091–2134.   doi: 10.1007/s11538-016-0214-9

Garrington, T.P., and Johnson, G.L. (1999). Organization and regulation of mitogen-activated protein
kinase signaling pathways. *Curr Opin Cell Biol.* 11: 211-218. doi: 10.1016/s0955-0674(99)80028-3

Gillespie, L.L., Paterno, G.D., and Slack, J.M. (1989). Analysis of competence: receptors for
fibroblast growth factor in early *Xenopus* embryos. *Development* 106: 203-208. doi:
https://dev.biologists.org/content/106/1/203

Gómez-Cruz, N., and Maldonado, C. (2011). Biological Computation: A Road to Complex
Engineered Systems. *Unifying Themes in Complex Systems* VIII: 918-927. Cambridge, MA: NECSI
Knowledge Press. doi: https://ssrn.com/abstract=3663378

Grabowski, F., Czyż, P., Kochańczyk, M., and Lipniacki, T. (2019). Limits to the rate of information
transmission through the MAPK pathway. *J. R. Soc. Interface* 16: 20180792. doi:
http://dx.doi.org/10.1098/rsif.2018.0792

Huang,C.F.,and Ferrell, J.E.(1996). Ultrasensitivity in the mitogen-activated protein kinase cascade.
Proc. Natl. Acad. Sci. USA 93: 10078-1008. doi: https://doi.org/10.1073/pnas.93.19.10078

Kanodia, J., Chai, D., Vollmer, J., et al. (2014). Deciphering the mechanism behind Fibroblast
Growth Factor (FGF) induced biphasic signal-response profiles. *Cell Communication and Signaling*
12:34. doi: http://www.biosignaling.com/content/12/1/34

Kauffman S (1969). Metabolic stability and epigenesis in randomly constructed genetic nets. Journal
of Theoretical Biology. 22 (3): 437–467. doi:10.1016/0022-5193(69)90015-0

Kholodenko, B. N. (2000). Negative feedback and ultrasensitivity can bring about oscillations in the
mitogen-activated protein kinase cascades. *Eur. J. Biochem.* 267: 1583-1588 . doi: 10.1046/j.1432-
1327.2000.01197.x.

Kholodenko, B. N., and Birtwistle, M.R. (2009). Four-dimensional dynamics of MAPK information
processing systems. Wiley Interdiscip Rev Syst Biol Med. 1: 28–44. doi: 10.1002/wsbm.16.

Klint, P., and Claesson-Welsh, L. (1999). Signal transduction by fibroblast growth factor receptors.
Front Biosci. 15: D165-77. doi: 10.2741/klint

Klipp, E., et al. (2009). Systems Biology: A Text Book. WILEY-BLACKWELL. ISBN: 978-3-527-
31874-2.

Lemmon. M.A., and Schlessinger, J. (2010). Cell Signaling by Receptor Tyrosine Kinases. *Cell* 141:
117- 1134. doi: 10.1016/j.cell.2010.06.011

Lim, S., and Kaldis, P. (2013). Cdks, cyclins and CKIs: roles beyond cell cycle regulation.
Development 140: 3079-3093. doi:10.1242/dev.091744

Mohammadi, M., Olsen, S.K., and Ibrahimi O. A. (2005). Structural basis for fibroblast growth factor
receptor activation. *Cytokine Growth Factor Rev.* 16: 107-137. doi: 10.1016/j.cytogfr.2005.01.008

Mousaviana, Z., Díaz, J., and Masoudi-Nejad, A. (2016). Information theory in systems biology. Part II: protein–protein interaction and signaling networks. *Seminars in Cell & Developmental Biology* 51: 14-23. doi: https://doi.org/10.1016/j.semcdb.2015.12.006

Sears, R.C. (2004). The life cycle of c-myc: from synthesis to degradation. *Cell Cycle* 3:1133-1137. doi: 10.4161/cc.3.9.1145

Shen, T., and Huang, S. (2012). The role of Cdc25A in the regulation of cell proliferation andapoptosis. *Anticancer Agents Med Chem*. 12: 631–639. doi:10.2174/187152012800617678

Vaseghi, S.V. (2000). Advanced Digital Signal Processing and Noise Reduction, Second Edition John Wiley & Sons Ltd. ISBNs: 0-471-62692-9

Timofeev, O., Cizmecioglu, O., Hu, E., Orlik T., and Hoffmann. (2009). Human Cdc25A phosphatase has a non-redundant function in G2 phase by activating Cyclin A-dependent kinases. *FEBS Letters* 583: 841-847. doi: 10.1016/j.febslet.2009.01.044

Tka č ik, G., and Walczak, A.M. (2011). Information transmission in genetic regulatory networks: a review. *J Phys Condens Matter* 23(15):153102. doi: 10.1088/0953-8984/23/15/153102

Wiener, N. (1961). Cybernetics: Or Control and Communication in the Animal and the Machine. *Paris, (Hermann & Cie) & Camb. Mass. (MIT Press)* ISBN 978-0-262-73009-9; 1948, 2nd revised ed. 1961

789 **10   Tables**

790   **Table 1.- Parameters of the model.**

Parameter	Value	Parameter	Value
k_1	0.00476 nM^{-1}s^{-1}	SOS_T	90 nM
k_{-1}	0.0008 s^{-1}	GAP_T	60 nM
k_2	0.00625 nM^{-1}s^{-1}	ras_T	70 nM
k_3	0.059 s^{-1}	raf_T	300 nM
k_4	2 nMs^{-1}s^{-1}	mek_T	300 nM
k_{-4}	0.5 s^{-1}	erk_T	300 nM
k_5	10 nM^{-1}s^{-1}|	V_{a1}	0.01 s^{-1}
k_{-5}	1 s^{-1}	V_{a2}	0.25 nM^{-1}s^{-1}
k_7	0.7 s^{-1}	V_{a5}	0.75 nM^{-1}s^{-1}
k_{-7}	0.91 s^{-1}	V_{a6}	0.75 nM^{-1}s^{-1}
k_8	0.025 s^{-1}	V_{a9}	0.5 nM^{-1}s^{-1}
k_9	0.025 s^{-1}	V_{a10}	0.5 nM^{-1}s^{-1}
k_{11}	0.025 s^{-1}	K_{a2}	8 nM
k_{12}	3 s^{-1}	K_{a3}	15 nM
k_{13}	5 nM^{-1}s^{-1}	K_{a4}	15 nM
k_{14}	0.35 s^{-1}	K_{a5}	15 nM
k_{15}	1 s^{-1}	K_{a6}	15 nM
k_{16}	0.5 s^{-1}	K_{a7}	15 nM
k_{17}	0.2 nM^{-1}s^{-1}	K_{a8}	15 nM
k_{-17}	1 s^{-1}	K_{a9}	15 nM
k_{18}	0.01 s^{-1}	K_{a10}	15 nM
k_{-18}	0.00045 s^{-1}	D	1 s^{-1}
k_{19}	0.005 s^{-1}	$V_{Cmyc}^{T\max}$	8 nM^{-1}s^{-1}
k_{-19}	0.001 s^{-1}	$V_{DUSP}^{T\max}$	4 nM^{-1}s^{-1}
k_{20}	0.02 s^{-1}	$V_{CDC25A}^{T\max}$	4 nM^{-1}s^{-1}
k_{-20}	0.0075 s^{-1}	V_{cyt}	800 µm^3
k_{21}	1 s^{-1}	V_n	400 µm^3
k_{22}	0.5 s^{-1}	mix_T	5 nM

808   **References:** Kholodenko (2000), Klip et al. (2009), Huang and Ferrell (1996).