Loss of full-length dystrophin expression results in major cell-autonomous abnormalities in proliferating myoblasts.

Maxime R. F. Gosselin[1], Virginie Mournetas[2], Malgorzata Borczyk[3], Lukasz Bozycki[1,4], Michal Korostynski[3], Samuel Robson[1,5], Christian Pinset[6], Dariusz C. Górecki[1] *

[1]School of Pharmacy and Biomedical Sciences, University of Portsmouth, PO1 2DT, Portsmouth, UK

[2]INSERM UEVE UMR861, I-STEM, AFM, 28 rue Henri Desbruères, 91100 Corbeil-Essonnes, France

[3]Laboratory of Pharmacogenomics, Maj Institute of Pharmacology PAS, Krakow, Poland

[4]Laboratory of Biochemistry of Lipids, Nencki Institute of Experimental Biology, Warsaw, Poland.

[5]Centre for Enzyme Innovation, University of Portsmouth, PO1 2DT, Portsmouth, UK

[6]CNRS, I-STEM, AFM, 28 rue Henri Desbruères, 91100 Corbeil-Essonnes, France

*Corresponding author (darek.gorecki@port.ac.uk)

Keywords

DMD, dystrophin, mdx, myoblast, transcriptomics

Abstract

Background. Duchenne muscular dystrophy (DMD) is the most common inherited muscle disease that leads to severe disability and death in young men. DMD is caused by out-of-frame mutations in the largest known gene, which encodes dystrophin. The loss of DMD gene expression manifests in progressive degeneration and wasting of striated muscles aggravated by sterile inflammation. Current conventional treatments are palliative only, whereas experimental therapeutic approaches focus on the re-expression of dystrophin in myofibers. However, recent studies established that DMD pathology begins already in prenatal development prior to myofiber formation while, in adult muscle, it affects satellite (stem) cells and the proper development of myofibers. Regeneration defects that exacerbate muscle degeneration appear to be a good therapeutic target, as maintaining regeneration would counteract muscle wasting. It is also the only feasible treatment in advanced stages of the disease. Yet, it is unknown whether dystrophic

myoblasts, the intermediary between satellite cells and myofibers and effectors of muscle growth and repair, are also affected. Therefore, we investigated whether DMD myoblasts show a dystrophic phenotype.

Methods and Findings. Using a combination of transcriptomic, molecular, biochemical, and functional analyses we demonstrate, to our knowledge for the first time, convergent cell-autonomous abnormalities in primary mouse and human dystrophic myoblasts. In *Dmd*^{mdx} mouse myoblasts lacking full-length dystrophin transcripts, expression of 170 other genes was significantly altered. Myod1 (p=2.9e^{-21}) and key muscle genes controlled by MyoD (Myog, Mymk, Mymx, epigenetic regulators, ECM interactors, calcium signalling and fibrosis genes) were significantly downregulated. Gene ontology enrichment analysis indicated significant alterations in genes involved in muscle development and function. These transcriptomic abnormalities translated into increased proliferation (p=3.0e^{-3}), reduced migration towards both sera-rich (p=3.8e^{-2}) and cytokine-containing medium (p=1.0e^{-2}), and significantly accelerated differentiation in 3D organotypic cultures. These altered myoblast functions are essential for muscle regeneration. The defects were caused by the loss of expression of full-length dystrophin as strikingly similar and not exacerbated alterations were also observed in dystrophin-null Dmd$^{mdx-βgeo}$ myoblasts. Furthermore, corresponding abnormalities were identified in human DMD primary myoblasts and in an established dystrophic mouse muscle (SC5) cell line, confirming universal, cross-species and cell-autonomous nature of this defect.

Conclusions. These results, for the first time, demonstrate the disease continuum: DMD defects in satellite cells cause myoblast dysfunctions diminishing muscle regeneration, which is essential to counteract myofiber degeneration. Full-length dystrophins play a critical role in these processes. Contrary to the established belief, our data identify myoblasts as a novel and important therapeutic target for treatment of this lethal disease.

Introduction

Duchenne muscular dystrophy (DMD) is a debilitating and lethal neuromuscular disorder caused by mutations in the *DMD* gene located on the X chromosome [1]. Diagnosis is made between the age of 2 and 5, loss of ambulation occurs around 12 and young adults die due to respiratory and/or cardiac failure [2].

DMD is the largest human gene known [3]: Three full-length transcripts encode 427 kDa proteins, while further intragenic promoters drive expression of progressively truncated variants. Dp427 and the dystrophin-associated protein complex (DAPC) are important for the functional development of

differentiating myotubes [4] and subsequently prevent contraction-induced injury in the mature muscle[5].

Several studies showed that the ablation of dystrophin in fully differentiated myofibres did not trigger their degeneration [5,6], and even that myofibres can function entirely without dystrophin [7,8].

In fact, DMD pathology is active prior to diagnosis: delays in the attainment of motor and non-motor milestones are discernible in 2 months old DMD babies [9] and transcriptomes of muscles from asymptomatic DMD patients revealed typical dystrophic abnormalities [10]. Studies of human foetuses [11–13] and various animal DMD models, including GRMD dogs [14], Sapje zebrafish [15] and Dmd^mdx mouse embryos [16] revealed that pathology starts already in prenatal development.

Indeed, in skeletal muscle lineages modelled in human DMD pluripotent stem cells, we have recently demonstrated marked transcriptome and miRNA dysregulations identifiable even before muscle specialisation [17]. These data, combined with the existence of the specific embryonic dystrophin Dp412e [18], substantiate the early disease manifestations in muscle precursor cells.

Importantly, there is increasing evidence that dysregulation of myogenic cells is behind muscle pathology and disease progression in adult DMD muscle. However, the mechanism(s) remain(s) to be elucidated and, given the interaction of myogenic and inflammatory cells in muscle regeneration [19], distinguishing the primary and the secondary consequences of dystrophin deficiency is crucial for development of effective therapies.

Helen Blau proposed that DMD is intrinsic to the undifferentiated myoblast [20]. This hypothesis was initially discounted [21], but there is new evidence that DMD mutations produce a range of cell-autonomous abnormalities in both human and mouse myogenic cells from adult muscles [22–25].

If DMD directly affects myoblasts, cells that are key to muscle regeneration, a better definition of the consequences of the loss of DMD gene expression could help identifying early disease biomarkers and establishing better therapeutic targets.

Herein, using a combination of RNA Seq, molecular and functional approaches, we compared the dystrophic and healthy myoblasts isolated from skeletal muscles of the commonly used Dmd^mdx mouse model.

While Dmd^mdx represents the loss of full-length (Dp427) dystrophins expression, which reflects the molecular defect affecting majority of DMD patients, Dp71 is present in regenerating muscles and specifically in myoblasts [26]. Yet, little attention has been given to the DMD pathology in dystrophin-null individuals. We have recently described exacerbated pathology in Dmd^mdx-βgeo mice

lacking all dystrophins [27]. Therefore, we also investigated consequences of the total loss of *Dmd* expression in *Dmd*^mdx-βgeo^ myoblasts [28].

Finally, we compared transcriptomic alterations in mouse and human dystrophic myoblasts to identify defects occurring across species.

We report that the absence of expression of the full-length dystrophin triggers major transcriptomic and functional abnormalities in myoblasts. These abnormalities are cell autonomous, as they persist in myogenic cells maintained long-term in culture. Importantly, key alterations are common between mouse and human myoblasts.

Materials and methods

Animals

Male C57BL/10ScSn-*Dmd*^mdx^/J, C57BL/10ScSnJ, C57BL/6-DmdGt(ROSAbgeo)1Mpd/J (*Dmd*^mdx-βgeo^) and C57BL/6J eight week old mice were used in accordance with institutional Ethical Review Board and Home Office (UK) Approvals. All mice were maintained under pathogen-free conditions and in a controlled environment (12-hour light/dark cycle, 19-23°C ambient temperature, 45-65 % humidity). For environmental enrichment, tubes and nesting materials were routinely placed in cages. The genotypes of all experimental animals were confirmed by PCR.

The C57BL/10 and C57BL/6 strains derived from a common origin [29] and it has been demonstrated that the mdx mutation on the C57BL/6 background shows the same pathology as the original C57BL/10 strain [30]. Nevertheless, for RNA Seq experiments, mixed background animals were generated to facilitate the direct comparison between *Dmd*^mdx^ and *Dmd*^mdx-βgeo^ transcriptomes. For this, *Dmd*^mdx^ males were paired with C57BL/10 females and C57BL/6 males were paired with *Dmd*^mdx-βgeo^ females. The resulting *Dmd*^mdx/WT^ females and *Dmd*^mdx-βgeo^ males were then bred together. The resulting *Dmd*^WT^ males were used as controls and bred with the *Dmd*^mdx/mdx-βgeo^ females. The next generation *Dmd*^mdx^ and *Dmd*^mdx-βgeo^ males were used as the dystrophic groups. For the subsequent functional pairwise analyses *Dmd*^mdx^ were compared against C57BL/10 and *Dmd*^mdx-βgeo^ against C57BL/6 mice.

Primary myoblast extraction and culture

Gastrocnemii from 8-week-old male mice were used for myoblast extraction and culture. *Gastrocnemius* was chosen as it is considered a fast muscle with predominantly IIB fibres [31] and therefore more prone to dystrophic changes and producing a relatively homogenous population of myoblasts. Briefly, muscles were dissected free of fat and connective tissue before enzymatic

digestion by 0.2 % type IV collagenase (Sigma, C5138) in Dulbecco's Modified Eagles' Medium (DMEM, Gibco 31885-023) for 90 min at 37°C, 5 % CO_2. The digested muscles were then rinsed in plating medium (10 % horse serum, Gibco 26050-088; 0.5 % chicken embryo extract, Seralab CE-650-J; 100 units penicillin / 0.1 mg streptomycin per mL, Sigma P0781; in DMEM, Gibco 31885-023) incubated at 37°C, 5 % CO_2. Digested muscles were then disrupted by successive passages in 50 mL, 25 mL and 10 mL serological pipettes. The freed muscle fibres were transferred to a dish containing plating medium.

The dish was then placed under a stereo microscope and 200 live fibres were transferred to another dish containing plating medium. This procedure was repeated 2 more times until 150 fibres were transferred to a third dish. 120 live fibres were then transferred to a tube containing proliferating medium (20 % foetal bovine serum, Gibco 10500-064; 10 % horse serum, Gibco 26050-088; 0.5 % chicken embryo extract, Seralab CE-650-J; penicillin / streptomycin, Sigma P0781; in DMEM, Gibco 31885-023). The resulting suspension of muscle fibres was then disrupted by passing it through an 18-gauge needle 10 times before passing the resulting solution through a 40 μm cell strainer (BD Falcon 352340).

The strained cell suspension was transferred to a collagen I (Sigma C8919) coated cell culture dish and proliferating medium was added to obtain an appropriate volume for the dish. The cells were then cultured at 37°C, 5% CO_2 humidified atmosphere and expanded as needed for assays and experiments. Cells were lifted from dishes using Accutase (Biowest L0950).

Myoblasts cell lines

The SC5 (mdx) and IMO (WT) cell lines were derived from the leg muscle of the H2Kb-tsA58 line [32]. Cells were cultured in DMEM supplemented with 20% FCS, 4mM L-glutamine, 100 unit/ml penicillin, 100 μg/ml streptomycin and 20 unit/ml murine γ-interferon (Invitrogen) at 33°C, 5% CO_2 humidified atmosphere. When cells reached 95% confluence, they were cultured as primary cells in KnockOut DMEM (Invitrogen) supplemented with 10% v/v Knockout Serum Replacement (KSR, Invitrogen), 5% v/v Donor Horse Serum (DHS, Sera Labs) and 2mM L-glutamine at 37°C.

RNA extraction

Myoblasts isolated from gastrocnemii from 3 individual mice per experimental group were treated as biological replicates. Total RNA was extracted from Dmd^{mdx}, $Dmd^{mdx-\beta geo}$ and corresponding wild-type myoblasts using RNEasy Plus Universal mini kits (Qiagen 73404). Briefly, cells were washed 2 times with warm Dulbecco's phosphate buffered saline DPBS, Gibco 14190144) before lysis by applying QIAzol directly to the freshly washed cells. Lysate was then homogenised by passing it through a 25-gauge needle 20 times. Samples were then processed according to the kit

manufacturer's instructions. RNA quality and concentration were measured using a NanoDrop 1000 Spectrophotometer (Thermo Scientific). RNA integrity was assessed using electrophoresis of 100 ng of total RNA in a 1 % agarose gel (Sigma A4718) in TRIS-acetate-EDTA (TAE) buffer or using an automated capillary electrophoresis system (2100 Bioanalyzer Instrument G2939BA, Agilent) using a kit assay (Agilent RNA 6000 Nano Kit 5067-1511, Agilent).

RNA Sequencing

Total RNA samples (n=3 per group) were processed by TheragenEtex (Republic of Korea) using an Agilent Bioanalyzer 2100 for quality assessment of samples with a threshold of 7.0 for the RNA integrity number (RIN). An Illumina TruSeq stranded total RNA kit was used to generate libraries following a ribodepletion step using a Ribo-Zero Human/Mouse/Rat kit. Libraries were sequenced in a paired-end 100 bp run using an Illumina HiSeq 2500 sequencing platform.

Raw reads were quality assessed using fastQC [33] and reads were trimmed using trim-galore [34] with parameters "--illumina -q 20 --stringency 5 -e 0.1 --length 40 --trim-n" to remove adapter sequence and low-quality sequence tails. Reads were mapped to the GRCm38 *Mus musculus* genome from Ensembl using the STAR universal RNA Seq aligner [35] with parameters "--outSAMmultNmax 300 --outSAMstrandField intronMotif". Output mapping files were processed and filtered to remove non-mapping reads with mapping quality less than 20 using samtools [36].

The DESeq2 package [37] in R [38] was used to perform differential gene expression analysis. Gene models were taken from Ensembl version 91, and read counts over unique genes were quantified using the *summarizeOverlaps()* function in the GenomicAlignments package [39] using parameters 'mode = "Union", singleEnd = FALSE, ignore.strand = FALSE, fragments = FALSE, preprocess.reads = invertStrand'. *P* values were adjusted for multiple comparisons by using the Benjamini and Hochberg correction [40]. Significantly differentially expressed genes were identified based on a fold-change of 2-fold or greater (up- or downregulated) and an adjusted *p*-value < 5.0e-2. An additional filter was put in place to remove genes where the mean normalised Fragments Per Kilobase Mapped (FPKM) was < 1 for both conditions to avoid changes in low abundance transcripts. GO enrichment analysis was conducted using the clusterProfiler package [41] in R [38].

REVIGO [42] was used to reduce the redundancy of GO enrichment data to rationalise the categories being compared in figures. The default "SimRel" semantic similarity measure with a 0.4 threshold cut-off was used. When filtering GO categories for redundancy, the mouse UniProt database was used for all analyses, except when analysing the list of GO categories enriched in both human and mouse primary datasets, in which case the human UniProt database was used. STRING [43] was used

to explore and visualize interaction networks. Only interactions present in databases for relevant species or experimentally determined were used.

Regulatory regions of the differentially expressed genes (listed in Tables S2, S3, S5, and Table S3H), defined as the region 1 Kb upstream or downstream of the transcription start site (TSS) of the gene, were analysed for overrepresentation of binding sites of transcription factors (TFs). The binding signal was analysed using the seqinspector tool (http://seqinspector.cremag.org, [44] and publicly available ChIP-Seq data [45]. The overrepresentation was assessed by comparing the lists of tested genes with 1000 randomly selected gene promoters (Mus musculus reference genome mm9) Ensembl 75 gene symbols. TF signal was considered enriched if Bonferroni-corrected p-value returned by seqinspector was < 5.0e-2.

RNA Seq data have been deposited in the ArrayExpress database at EMBL-EBI under accession number E-MTAB-10322 for the primary mouse myoblast samples (https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-10322) and accession number E-MTAB7287 for SC5 and IMO myoblast cell line samples (https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-7287).

Human primary adult myoblast RNA Seq data

RNA Seq data from human dystrophic (duplication exons 3–26, deletion exons 8–43, and stop exon 7) and healthy primary myoblasts were generated and analysed as described [17]. Raw RNA Seq data have been deposited in the ArrayExpress database at EMBL-EBI under accession number E-MTAB-8321 (https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-8321).

cDNA synthesis and qPCR analysis

Total RNA samples were converted to cDNA using SuperScript VILO cDNA Synthesis Kit (Invitrogen 11754050) as per manufacturer instructions.

qPCR reactions were run in duplicates using 25 ng of cDNA per reaction with Precision Plus Mastermix (Primer Design PPLUS-LR), forward and reverse primers synthesized by Eurofins (Table S10) and DEPC treated water (Fisher Bioreagents BP561) as per manufacturer instructions, in 96 well plates using an Applied Biosystem ViiA7 RT-PCR instrument, and expression quantified using the ΔΔCT method.

Cell proliferation assay

Primary myoblasts were seeded in 10 cm diameter dishes (Sarstedt 83.3902) coated with Collagen I (Sigma C8919) at 30 % confluency in proliferation medium, as above and left to attach for 2 hours before adding bromodeoxyuridine / 5-bromo-2'-deoxyuridine (BrDU, Invitrogen B23151) to a final

concentration of 75 μM and left to proliferate for 6 hours. Cells were then rinsed with warm DPBS, detached using Accutase (Biowest L0950) and fixed using ice cold ethanol. Samples were left at -20°C over-night and then analysed by flow cytometry. Fixed cells were submitted to an acid wash (2M HCl, 0.5 % Triton X-100) for 30 min at room temperature, washed in PBS and the remaining acid was neutralized using a borate buffer (pH= 8.5). Cells were then stained in 200 μL of PBS (0.5 % Tween 20) with 1 μg of Anti-BrDU FITC conjugated antibody (BD 556028) or an FITC conjugated isotype control for 30 min at room temperature. Cells were then washed in HEPES buffer (pH= 7.4) and resuspended in PBS before processing using a FACScalibur system.

Cell migration and invasion assay

Primary myoblasts from dystrophic and WT mice were treated for 2 hours with proliferating medium complemented with 10 μg.mL^{-1} mitomycin C (Fisher Bioreagents BP2531) to prevent proliferation from confounding the results of the migration assay. Cells were then rinsed with warm DPBS twice to remove excess mitomycin C. Each Matrigel coated (Corning 354234) or uncoated insert (Sartstedt 83.3932.800) was seeded with 10 000 myoblasts in serum-free medium while the wells were filled with either proliferating medium or serum-free medium complemented with 10 ng.mL^{-1} FGF-b (R&D Systems 3139-FB), IGF-I (R&D Systems 791-MG) and HGF (R&D Systems 2207-HG). Cells were left to migrate and invade for 12 hours before fixation in buffered formalin (Sigma HT501128) and crystal violet staining for cell counting and spectrophotometer measurements.

Cell differentiation assay

Myoblasts were used to generate spheroids as such a culture provides an environment that facilitates myogenic differentiation due to a 3D structure improving physical interactions, differentiation and fusion of myoblasts [46–49]. One million myoblasts per genotype and per mouse were pelleted at 300g for 5 minutes in 15 mL centrifuge tubes before pellets were transferred into individual wells of a 6-well plate containing 50 % proliferation medium (20 % foetal bovine serum, Gibco 10500-064; 10 % horse serum, Gibco 26050-088; 0.5 % chicken embryo extract, Seralab CE-650-J; penicillin / streptomycin, Sigma P0781; in DMEM) and 50 % DMEM (Gibco 31885-023) and cultured for 24 hours. Then cells were placed in 25 % proliferation medium with 75 % DMEM for 24 hours and finally in the differentiation medium (2 % horse serum, Gibco 26050-088; penicillin / streptomycin, Sigma P0781; in DMEM). Introduction to differentiation medium was considered t=0 and differentiation was allowed for 6 days.

At the collection time, spheroids undergoing differentiation were rinsed in DPBS and collected in 1 mL of Qiazol solution before freezing for subsequent processing. On thawing, spheroids were disrupted using progressively smaller gauge needles (19, 21, 23 and 25 gauge) and then processed as

per the RNA extraction section. *Mymk* was chosen as a marker of myoblast fusion and therefore entry into the final stages of differentiation [50] and *Myog* and *Myh1* as robust markers of late stage myogenic differentiation [51].

Immunolocalisation with confocal microscopy

Muscles (tibialis anterior) frozen in liquid N_2 chilled-isopentane in cryo-embedding matrices (Cellpath KMA-0100-00A) were allowed to equilibrate to -20 °C in the cryostat chamber. Cryosections (10 μm) were cut and attached to poly-L-lysine coated slides (Thermo Scientific J2800AMNZ). Sections were allowed to air-dry before fixation and staining. Myoblasts were seeded at 10 000 per collagen-coated coverslip in a 24 well plate (Sarstedt 83.1840.002 and 83.3922) and left to adhere for 12 hours before processing. Spheroids having undergone the differentiation process were frozen in embedding medium and 10 μm sections were cut, as previously described.

Both muscle sections and cultured myoblasts were then processed identically. They were fixed in 10 % neutral buffered formalin (Sigma-Aldrich HT501128) at room temperature for 10 minutes and washed 3 times in PBS containing 0.1 % v/v Triton X-100 (Sigma-Aldrich X100), blocked for 30 min using 10 % v/v chicken serum (Gibco 16110082) in PBS-Triton then incubated for 90 minutes under agitation in primary antibody solution: anti dystrophin (Developmental Studies Hybridoma Bank MANDYS1 clone 3B7 cell culture supernatant) 25 % v/v antibody or anti Myosin-4 (Invitrogen, 14-6503-95) 1μg.mL^{-1}, 10 % v/v chicken serum in PBS-Triton. Samples were then washed one time in PBS-Triton then incubated in secondary antibody solution (Invitrogen A-21200) with 10 μg.mL^{-1} antibody, 10 % v/v chicken serum and DAPI 300 nM (Invitrogen D1306) in PBS-Triton for 1 hour followed by 2 washes in PBS-Triton then PBS and mounted using an anti-fading mounting medium (Thermo Scientific TA-006-FM). Omitting primary antibodies in the reaction sequence served to confirm the specificity of the signal.

Samples were examined with a confocal laser-scanning microscope (LSM 880; Zeiss, Oberkochen, Germany) using a Plan Apochromat, 63x DIC oil objective (NA 1.4) (pixel size 0.13 μm) objective. Images were acquired using sequential acquisition of the different channels to avoid crosstalk between fluorophores. Pinholes were adjusted to 1.0 Airy unit. In all cases where multiple images were captured from the same immunohistochemical reaction, laser power, pinhole, and exposure settings were captured once on sample from a representative control section and maintained throughout imaging. Images were processed with the software Zen (Zeiss) and exported into tiff images for figure generation or WMV for video exporting. Only brightness and contrast were adjusted for the whole frame, and no part of any frame was enhanced or modified in any way.

Statistical analysis.

Student's unpaired t-test was used for comparisons between two data groups. two-way ANOVA was used to determine if there was any statistically significant difference between groups and time points in the differentiation assay and Fisher's LSD was used to determine which time points exhibited statistically significant differences between genotypes. Pearson's correlation coefficient was calculated to quantify the level or correlation between datasets and Fisher's exact test was used to further investigate the similarities between human and mouse primary myoblasts gene expression data. All results which passed the threshold for statistical significance were tested for normal distribution using the Shapiro-Wilk normality test, all analysed data were normally distributed. A p-value of ≤ 5.0e-2 was considered statistically significant, and the values are reported as follows in figures: *p ≤ 5.0e-2, **p< 1.0e-2, ***p < 1.0e-3, ****p < 1.0e-4. All statistical analyses outside of differential gene expression and GO enrichment analyses were performed using GraphPad Prism 8.4.2.

Results

Transcriptomic alterations in proliferating Dmd^{mdx} myoblasts

Total RNA extracted from primary myoblasts isolated from gastrocnemii of 8 week old male Dmd^{mdx}, $Dmd^{mdxβgeo}$ and control mice was subjected to RNA Seq and analysed for the differential expression of genes between groups and the enrichment of GO categories in the generated lists of differentially expressed genes.

Bioinformatic analysis of the RNA Seq data showed the impact of the Dmd^{mdx} allele on the primary myoblast transcriptome. Sample-based hierarchical clustering clearly segregates genes into two groups corresponding to genotypes (Figure 1a) and a volcano plot illustrates the presence of a substantial number of significantly dysregulated genes (Figure 1b; Table S1).

Expression of 170 genes was found to be significantly up- or down-regulated more than 2-fold when comparing Dmd^{mdx} and WT primary mouse myoblasts (Table S1). Among these, *Myod1* and *Myog,* key coordinators of skeletal muscle development and repair, were found to be downregulated in dystrophic myoblasts and their downregulation was confirmed by qPCR (Figure S1). *Pax3* and *Pax7* transcriptional regulators [52] were also individually investigated, with Pax3 levels confirmed as significantly downregulated by qPCR (Figure S1). Analysis identified downregulation of further important regulators and effectors of the muscle program (Figure 1c; Table S1) such as *Mymx and Bex1,* known to be regulators of muscle repair [53]. *Des*, a structural component of myofilaments,

and *Itga7*, the primary laminin-1 receptor of myoblasts and mature fibres were also found to be significantly downregulated (Figure 1c; Table S1).

The *H19* gene, defined as promoting differentiation [54], was found to be significantly downregulated in proliferating *Dmd*[mdx] myoblasts (Figure 1c; Table S1), which combined with reduced expression of aforementioned muscle program markers, as well as myogenic differentiation markers such as *Acta1*, *Actc1*, *Actn3*, *Atp2a1* and upregulation of *Nov,* known to inhibit myogenic differentiation [55], imply an altered readiness of dystrophic myoblasts to differentiate. Moreover, numerous genes encoding histones and *Smyd1* [56] and *Hmga1* [57,58] regulators of chromatin organisation/chromatin interacting proteins, were found to be significantly dysregulated in dystrophic myoblasts (Figure 1c; Table S1). Taken together, these gene expression changes indicated an increased readiness of dystrophic myoblasts to proliferate, compared to their WT counterparts.

Several genes key to myoblast-extracellular matrix interactions and fibrosis were found to be significantly dysregulated in *Dmd*[mdx] myoblasts (Figure 1c; Table S1), including: *Mmp3*, *Mmp9* and *Mmp10* metalloproteases*,* collagen genes (*Col2a1*, *Col4a1*, *Col4a2* and *Col8a2*), *Lrrn1* a transmembrane protein interacting with fibronectin [59] as well as genes encoding fibronectin itself (*Fn1*) and decorin (*Dcn*). The latter promotes proliferation and differentiation in myoblasts [60] but also has a role in collagen assembly and mineralisation [61].

Finally, calsequestrins (*Casq1* and *Casq2*) (Table S1) were significantly downregulated, in line with previous observations in the dystrophic muscle [62].

No exacerbation of transcriptomic alterations in the dystrophin-null Dmd[mdx-βgeo] compared to Dmd[mdx] myoblasts

Corresponding bioinformatic comparison of *Dmd*[mdx-βgeo] vs. *Dmd*[mdx] myoblast transcriptomes found no substantial difference between them, as illustrated by sample clustering (Figure 2a) and volcano plot (Figure 2b), with no segregation between groups. Only 11 genes were found to be significantly differentially expressed between the two genotypes. Of these, 6 are pseudogenes, 1 is a processed transcript and only 4 are protein coding genes (*Igf1*, *Npr3*, *Postn* and *Capn6*). None of these genes have a higher log2 fold change than 1.7 and the adjusted *p*-values have an average of 1.0e-2 compared to $9.6.10^{-3}$ for the genes found to be significantly differentially expressed between *Dmd*[mdx] and WT myoblasts.

Interestingly, comparison of the *Dmd*[mdx-βgeo] to the WT transcriptome returned fewer (81) (Table S2) significantly differentially expressed genes than *Dmd*[mdx] versus WT (170) (Table S1). However, comparison of the fold changes between the two analyses (Figure 2c) revealed that the two models show strikingly similar alterations with a strong, significant correlation (r = 0.9418, *p*< 1.0e-4) in the

log2 fold changes for genes significantly altered in Dmd^{mdx} versus WT and/or $Dmd^{mdx-\beta geo}$ versus WT analyses (Figure 2c).

MyoD dependent downregulation of gene expression in dystrophic myoblasts

Lists of genes altered in dystrophic myoblasts were analysed for common patterns of transcriptional regulation. Overrepresented transcription factor binding sites (TFBS) on promoter regions of the dysregulated genes were examined using the seqinspector tool and available ChIP-Seq data [45]. Genes found downregulated in Dmd^{mdx} had increased ChIP-Seq signal at the TFBS for four transcription factors: MyoD (adjusted p-value: $2.9e^{-21}$, track GEO accession GSM915165), TCF12 ($1e^{-19}$, GSM915178), MYOG ($5.6e^{-19}$, GSM915164) and TCF3 (adjusted p-value 3.3e-3, GSM915177) (Figure 3; Table S3B). A range of genes, including *Mymk, Mymx, Chrna1* and *Acta1* showed a binding signal at their TFBS above the background of at least three out of four of these overrepresented TFs (Figure 3b; Table S3C). Furthermore, transcripts downregulated in $Dmd^{mdx-\beta geo}$ also exhibit overrepresentation of MyoD binding sites (Table S3E). In contrast, genes upregulated in any cell type under investigation did not show any statistically significant overrepresentation of the binding signal of any TF (Table S3A and D).

GO enrichment analysis indicates significant functional alterations in dystrophic myoblasts

Two broad GO categories: "muscle system process" and "regulation of muscle system process" were enriched in the downregulated genes found when comparing Dmd^{mdx} and $Dmd^{mdx-\beta geo}$ vs. WT primary mouse myoblasts (Figure 4a; Table S4). These categories include genes involved in muscle development and function. For a myoblast to fulfil its role in regenerating a damaged muscle, it must be able to proliferate, migrate and differentiate in response to relevant stimuli. GO enrichment analyses of downregulated genes (Figure 4; Table S4) imply phenotypic alterations in dystrophic myoblasts affecting all three functions.

Specifically, categories such as "chromatin assembly" and "muscle cell proliferation" (Figure 4b; Table S4), are enriched, indicating an altered proliferative state of dystrophic myoblasts. The enrichment of the GO categories "muscle cell migration" and "positive regulation of cell migration" in downregulated genes (Figure 4; Table S4) indicate a migrative phenotype in dystrophic myoblasts, while GO categories "muscle cell differentiation" and "striated muscle cell differentiation" (Figure 4a and b; Table S4) suggest that dystrophic myoblasts may have an altered ability to differentiate.

Moreover, several GO categories related to calcium ion transport, homeostasis and calcium-mediated signalling were found significantly enriched in Dmd^{mdx} vs. WT downregulated genes (Figure 4b; Table S4), in line with this well-established dystrophic myofibre abnormality [63].

These results suggested alterations in proliferation, migration, and differentiation in both Dmd^{mdx} and $Dmd^{mdx-\beta geo}$ cells. To further assess these processes, we performed functional analyses of these three key myoblast functions - proliferation, migration and differentiation – in primary myoblasts isolated from gastrocnemii of dystrophic and control mice.

Increased proliferative capacity of dystrophic myoblasts

The ability of myoblasts to proliferate is key to obtaining enough myogenic cells to repair muscle fibre damage. Given that the transcriptomic data indicated this mechanism as likely to be affected, we tested, in a BrDU incorporation assay, the dystrophic myoblasts capacity to respond to the proliferative stimulus of the exposure to a sera-rich medium (Figure 5a). After six hours, 57 % of Dmd^{mdx} myoblasts have incorporated BrDU compared to 26 % of WT cells (p=3.0e-3). 39.1 % of $Dmd^{mdx-\beta geo}$ cells were positive for BrDU compared to 24 % of the corresponding C57Bl/6 controls (p=1.0e-2). This statistically significant two-fold increase in BrDU incorporation denotes an exaggerated response to a proliferative stimulus in both dystrophic cells compared to their respective WT controls. Although Dp71 expression was described as enhancing myoblast proliferation [64], we did not observe any negative effect on proliferation in dystrophin-null $Dmd^{mdx-\beta geo}$ cells. On the other hand, this absence of differences between these two genotypes was consistent with the differential gene expression data described earlier.

Significantly decreased migration of dystrophic myoblasts in response to stimuli

In dystrophic myoblasts several GO categories associated with cell migration were found to be over-represented in the downregulated gene lists. We therefore assessed, using the Boyden chamber assay, the migration of dystrophic cells toward sera-rich medium and toward medium containing a cocktail of chemo-attractants (Figure 5b). Dmd^{mdx} myoblasts were found to migrate significantly less towards both sera-rich and cytokine-containing medium, with respectively 28 % (p=3.8e-2) and 21% (p=1.0e-2) cells migrating through the membrane compared to WT. $Dmd^{mdx-\beta geo}$ myoblasts also showed significantly reduced migration at 44 % (p=5.0e-6) and 54 %, (p=7.8e-4), compared to WT controls in serum-rich and cytokines media, respectively.

Altered differentiation of Dmd^{mdx} dystrophic myoblasts

Dystrophic myoblasts showed altered expression of several key genes involved in muscle differentiation (Figure 1c) and GO categories related to differentiation were enriched in downregulated gene lists (Figure4; Table S4). Even though these analyses were performed in proliferating myoblasts, such alterations could indicate a distorted capacity for differentiation in dystrophic cells.

To assess this, differentiation of Dmd^{mdx} myoblasts was compared against WT, using a three-dimensional (3D) culture, facilitating interactions closely resembling those occurring in muscles *in situ*.

Spheroids of dystrophic or WT cells (n = 3 per group), were placed in differentiation medium (t=0) and collected at days 0,2, 4 and 6 to monitor their differentiation. We first established that after 6 days in such condition's spheroids differentiated sufficiently to express myosin heavy chain (Figure S2; supplementary movie https://youtu.be/1axNWeK-Yb0). Changes in differentiation were assessed by qPCR measurement of the expression of *Myh1, Myog and Mymk,* gene markers of myoblast differentiation. Analysis of the time course of expression revealed a significantly altered pattern in Dmd^{mdx} cultures. Specifically, *Mymk* (Time: DF = 3, F = 7.520, *p* = 3.4e-2. Group: DF = 1, F = 50.40, *p* = 2.1e-3), *Myog* (Time: DF = 3, F = 9.811, *p* = 1.7e-2. Group: DF = 1, F = 11.72, *p* = 2.7e-2) and *Myh1* (Time: DF = 3, F = 24.97, *p* = 8.0e-4. Group: DF = 1, F = 66.95, *p* = 1.2e-3) were found to have their expression statistically significantly increased (two-way ANOVA) in dystrophic spheroids, when compared to WT (Figure 5c).

Also, in our hands, Dmd^{mdx} myogenic cells in 2D cultures formed myotubes within ≈ 5 days, while WT myoblasts required minimum 7 days in the differentiation medium (unpublished), which is in agreement with reports that dystrophic myoblasts differentiate faster [22].

It is worth noting that these increases in *Myh1*, *Myog* and *Mymk* expression in differentiating cells contrast with their expression profiles in proliferating myoblast, where dystrophic cells showed consistently lower expression levels of these markers (Table S1; Figure 1c and Figure S1).

Dystrophic myoblast cell line reproduces transcriptome anomalies found in primary cells

Although multiple replicates of primary myoblasts from the same genotype had similar transcriptome profiles (Figure 1), primary cultures might not be free from other cell types. Moreover, myoblasts isolated from Dmd^{mdx} muscle have been exposed to the dystrophic niche and some of the alterations might result from environmental factors. Therefore, we investigated whether alterations found in primary cells also occur in an established myoblast cell line. We used the SC5 (Dmd^{mdx}) and IMO (WT) cells, both derived from the H2Kb-tsA58 mice [32] and thus having an identical genetic background. Comparison of RNA Seq data showed a strong segregation of samples according to genotypes (Figure 6a) and a clear dysregulation in gene expression (Figure 5b, Table S5). The SC5 dystrophic cell line transcriptome had significantly more dysregulated genes compared to IMO controls (Table S5) than the primary Dmd^{mdx} myoblasts compared to WT cells (Table S1). However, despite this transcriptome drift expected in cells maintained long term in culture [65], the key alterations seen in the transcriptome of dystrophic primary cells were also

found in the established dystrophic cell line. *Myod1, Myog, Mymk, Des* and 69 other genes (Figure 6c; Table S5) were found significantly dysregulated in dystrophic cells and an overlap in GO (Figure 6c and d; Table S6) demonstrated that the impact of the mdx mutation is present, with 61 GO categories for biological processes being enriched in the downregulated genes lists in both primary and cell line myoblasts carrying the *Dmd*^mdx^ allele.

Further analysis of these 61 GO categories (Table S6) revealed that all those altered are among the most significantly dysregulated both in terms of adjusted *p*-values and fold enrichment in both primary and established cells, involving processes and functions, such as "striated muscle tissue development" and "striated muscle cell differentiation", very relevant to myoblasts (Figure 6c; Table S6).

Moreover, analysis of overrepresented TFBS on promoter regions of the dysregulated genes using the seqinspector tool found transcripts downregulated in the SC5 cells to have significantly higher than background signal for MyoD, MYOG, and TCF12 (Table S3G), in clear agreement with TFBS overrepresented in primary *Dmd*^mdx^ myoblasts. Likewise, genes upregulated in SC5 did not exhibit statistically significant overrepresentation of the binding signal of any TF (Table S3F).

Finally, in line with these transcriptomic similarities, the dystrophic myoblast cell line exhibited the impaired migrative phenotype (Figure S3), identical to that identified in primary dystrophic myoblasts (Figure 5b).

Human and mouse dystrophic primary myoblasts exhibit corresponding transcriptome changes

Comparison of differential gene expression data from human DMD and healthy primary myoblasts also showed significant transcriptomic alterations with a clear segregation between samples according to genotypes (Figure 7a) and a greater number of significantly downregulated genes (334) compared to upregulated ones (86) (Table S7, Figure 7b). This profile was very similar to the primary mouse myoblast data (Figure 1; Table S1).

Comparison of log2 fold changes of significantly dysregulated genes in human DMD myoblasts and their orthologous counterparts in mouse primary *Dmd*^mdx^ myoblasts, revealed that genes significantly downregulated in one species tend to also be downregulated in the other, as shown by more genes in the lower-left quadrant of Figure 7c, compared to other quadrants (Fisher's exact test: *p* = 2.0e-4). With *r* values ranging from 0.2357 to 0.4688, depending on which gene populations are compared (Figure 7c and d), the magnitude of gene expression changes between species was equivocal.

However, comparison of the human GO enrichment for biological processes in the significantly downregulated gene list to its mouse counterpart returned a strikingly similar result, with 61 overlapping categories (Figure 7d and e; Table S8). Of these, 49 also overlap with the categories in the mouse dystrophic myoblasts cell line. Again, the most significantly altered categories (in terms of adjusted *p*-value) that overlap in both species were associated with muscle cell development and differentiation (Figure 7c, Table S8).

When comparing the list of significantly downregulated genes in Dmd^{mdx} versus WT (Table S1) and human DMD versus healthy human myoblasts (Table S8), taking into account only one to one orthologs, 41 genes are present in both lists including key myogenic program actors such as *Myog*, *Mymk*, *Myod1*, *Des*, *Smyd1*, *Acta1*, *Actc1* and *Atp2A1*.

Interestingly, other genes of interest found to be dysregulated in primary mouse myoblasts were found to have non-orthologous counterparts dysregulated in human primary DMD myoblasts: *COL11A1*, *COL14A1* and *COL15A1* for *Col2a1*, *Col4a1*, *Col4a2* and *Col8a2* or *MMP2* for *Mmp3*, *Mmp9* and *Mmp10*.

Finally, a three-way comparison of the GO enrichment for biological processes in the downregulated genes for dystrophic primary human myoblasts, primary mouse myoblasts and the dystrophic mouse myoblast cell line showed an important overlap: 49 categories were found enriched in all three datasets, with further 12 enriched in human and mouse primary cells, 12 in mouse primary and mouse cell line datasets and finally 47 categories enriched in both primary human and mouse cell line dystrophic myoblasts (Figure 8a). In the 49 categories enriched in all 3 datasets, many are relevant to the disease and consistent with the cell function anomalies described earlier. Specifically, categories such as "muscle cell differentiation", "muscle system process", "regulation of myoblast differentiation" and "regulation of muscle system process" being very significantly enriched in the downregulated genes lists for all datasets (Figure 8b).

Alterations in transcriptomes of human and mouse dystrophic myoblasts are strikingly analogous, with downregulation of gene expression predominating in both, reduction in key markers of the muscle program and relevant common GO categories being enriched. Thus, these myoblast abnormalities are likely to the consequence of *DMD* gene mutations.

Discussion

DMD presents in muscle stem cells, where loss of dystrophin affects asymmetrical cell divisions [66]. Subsequently, absence of dystrophin during myotube differentiation causes the typical dystrophic abnormalities such as altered calcium homeostasis and creatine kinase leakage [4]. We hypothesised

that myoblasts originating from dystrophic stem cells which give rise to myotubes are also affected. Indeed, we demonstrate here for the first time that, in proliferating human and mouse myoblasts, absence of *DMD* gene expression results in major abnormalities. Our data combining global RNA Seq and functional analyses demonstrate that dystrophinopathy directly affects myoblasts.

The key abnormalities concerned proliferation, migration, and differentiation. Importantly, synchronisation of these three processes, involving exit from the cell cycle, migration to the site of damage, activation of the skeletal muscle specific genes and cell fusion, is essential for muscle development and regeneration. Ultimately, myoblasts are the effector cells of muscle growth and repair, and it is the failure of muscle regeneration that drives DMD progression.

These defects are cell autonomous rather than caused by the inflammatory environment of the dystrophic niche because they persist in the dystrophic cell line maintained long-term in culture: it reproduced the key transcriptomic anomalies and has been found before to have functional alterations found in primary cells [67].

Moreover, alterations are shared between mouse and human DMD myoblasts, despite the significant heterogeneity within human samples [17]. These dystrophic anomalies, therefore manifest themselves irrespective of differences in genetic backgrounds and across species, which implies their significance.

DMD pathology being intrinsic to myoblasts has been proposed previously [20] and there has been growing evidence that *DMD* mutations cause a range of cell-autonomous abnormalities in human and mouse myogenic cells and myoblasts that affect cell division, differentiation, energy metabolism and signalling [22–24,24,27,27,66,68–71]. Interestingly, Dmd^{mdx} cell migration found reduced under physiological stimuli (this work) can be increased in the inflammatory environment [27]. Furthermore, altered proliferation and migration resulting in highly metastatic phenotypes, has been associated with the loss of Dp427 in tumours featuring myogenic differentiation [72].

Our team recently demonstrated that *DMD* mutations evoke marked transcriptome and miRNome dysregulations early in muscle cell development [17]. In that study, expression of key coordinators of muscle differentiation was also dysregulated in still proliferating dystrophic myoblasts (Table S1, Figure 1c), the differentiation of which was subsequently found to be altered (Figure 5c).

A significant number of downregulated transcripts in dystrophic myoblasts are controlled by the same TFs, most notably MyoD and MYOG. Given that MYOG itself is regulated by MyoD, the significant downregulation of *Myod1* expression found here appears to be the key trigger mechanism of the dystrophic abnormality in myoblasts: the lower levels of *Myod1* transcripts strongly correlate with downregulated expression of genes controlled by this TF, and which are key

coordinators of skeletal muscle repair. Consequently, altered expression of myogenic regulatory factors in dystrophic myoblasts could have multiple consequences. Indeed, complete *Myod1* ablation in *Dmd*[mdx] has been shown to exacerbate the disease due to a diminished capability for muscle regeneration [73,74]. Downregulation of MyoD, which inhibits the cell cycle, could explain the increased proliferation in dystrophic cells. The *Myod1-Mymk-Mymx* (Myomaker/ Myomixer) axis altered here was found to determine myotube formation [75]. Given that *Pax3* inhibits myogenic differentiation of myoblasts [76], its downregulation in dystrophic cells could also contribute to this accelerated differentiation pattern observed here.

GO categories concerning calcium homeostasis and signalling, known to be altered in dystrophic muscles across species, were also found altered in dystrophic myoblasts: calsequestrins 1 and 2 levels being significantly downregulated, in line with previous findings in the dystrophic muscle [7,62] and cell lines [67,69].

Given that Dp71 dystrophin is found in undifferentiated myogenic cells [26], and that in development shorter dystrophins have been associated with proliferation and migration, and long isoforms with terminal commitment [77], we hypothesized that eliminating expression of Dp71 may exacerbate dysfunctions in myogenic cells. Surprisingly, we found that both *Dmd*[mdx] and *Dmd*[mdx-βgeo] cells had similar abnormalities and, contrary to Dp71 overexpression increasing myoblast proliferation [64], BrdU incorporation in dystrophin-null myoblasts was still significantly increased compared to WT. Interestingly, *Dmd*[mdx] myoblasts, having a point mutation in exon 23, express a significantly lower levels of Dp71 (Figure S6), whose expression is driven by a promoted located over 40 exons downstream of the mutation. But whether this downregulation is sufficient to mimic its total loss is unknown.

Combined, these data bring to light the significance of full-length dystrophin defects in undifferentiated muscle cells, the consequences of which correspond to the impaired muscle regeneration occurring in this disease.

Clearly, the important question is the mechanism altering the myoblast phenotype. Dp427 in satellite cells appears to control asymmetrical cell division *via* association with MARK2 and PARD3 [66], which might suggest an interaction problem also in myoblasts.

However, even healthy myoblasts do not express Dp427 protein at levels detectable by Western blotting: an observation that contributed to the belief that myoblasts are not affected by *DMD* mutations. Although our attempts at immunolocalization in unsynchronised primary myoblasts failed to detect Dp427 (Figure S4), a precise spatio-temporal requirement for small amounts of full-length dystrophin, analogous to its role in satellite cells [66] or at neuronal synapses [78] is a

possibility that requires further studies. Importantly, WT myoblasts express the full-length transcript, which is depleted by nonsense-mediated decay in dystrophic cells. It is worth noting that this 14 kb mRNA, which transcription time may be as long as 16 h [3], is produced in rapidly dividing myoblasts. Its expression is therefore unlikely to be meaningless.

Is it possible that loss of the full-length dystrophin expression may trigger a mechanism not involving its protein product? Considering the increasing number of human pathologies caused by RNA, including another dystrophy – *dystrophia myotonica* [79], it is not inconceivable that loss of a large 14 kb transcript made up of 79 exons may trigger a novel RNA-mediated disease process [80]. Abnormalities in Dmd^{mdx} cells, that harbour just a point mutation, indicate that the patho-mechanism does not require large *Dmd* gene rearrangements [81]. Yet, improvements resulting from the expression of mini/micro dystrophins do not support the requirement for the entire 14 kb transcript or a toxicity of mRNA breakdown products generated *via* nonsense-mediated decay. However, the impact of premature dystrophin transcript termination on the downregulation of other genes [82] cannot be excluded.

A more likely explanation involves alterations of epigenetic regulation, a mechanism of increasing importance in DMD: The absence of dystrophin/DAPC in satellite cells leads to aberrant epigenetic activation, which impairs functions of the newly generated cells. This dystrophic SC division defect has been linked to polarization problem due to MARK2 mis-interaction [83] combined with β-syntrophin abnormalities [84,85]. Downregulation of β-syntrophin, a DAPC member, results in impaired polarization of its interactor - p38γ. This, in turn, increases CARM1 phosphorylation and reduces *MYF5* activation by PAX7 in in the daughter cell due for myogenic differentiation [86]. Furthermore, dystrophin anchors nNOS [87], which becomes mis-localised in dystrophic cells. This results in decreased nitric oxide-dependent S-nitrosylation of HDAC2 [88] and higher deacetylase activity that has been shown to be involved in the progression of muscular dystrophy [89].

Thus, loss of the full-length dystrophin in muscle stem cells does not abrogate myogenic cell divisions but causes significant abnormalities. The descendant myoblasts appear to be harbouring somatically heritable epigenetic changes analogous to genomic imprinting [90] and themselves might manifest significant epigenotype abnormalities [91].

Indeed, our *DMD* knockdown and rescue experiments in skeletal muscle progenitor cells [17] showed that phenotypic changes at day 17 were the consequences of some earlier events that occurred at day 10 and that some abnormalities might not be reproduced nor alleviated after crossing a specific checkpoint.

The data here indicate that epigenetic changes in dystrophic primary myoblasts may be present, as amongst the significantly altered genes *Hist2h2ac*, *Hist1h2ag*, *Hist1h2ah*, *Hist1h1a*, *Smyd1 and Hmga1* gene products are involved in chromatin modifications or belong to the GO category "negative regulation of gene expression, epigenetic". Methylated histone H3 lysine 4 (H3K4) is a key epigenetic signal for gene transcription. H3K4 methylation is mediated by several methyltransferases, including muscle-active *Smyd1* [92], found among the top downregulated transcripts in dystrophic myoblasts. Histone methylation can be reversed by histone demethylases, of which Lysine-Specific Histone Demethylase 1A (LSD1 encoded by *Kdm1a*) is an important muscle enzyme [93]. The network of interactions between genes in this aforementioned GO category (Figure S5, Table S9) showed a clear interface between the *Kdm1a* node and significantly downregulated histone genes. LSD1 is involved in controlling *Myod1* expression [94], which we found significantly downregulated in dystrophic myoblasts. Importantly, treatment of *Dmd*[mdx] mice with histone deacetylase inhibitors (HDACi) promoted myogenesis [95]. Epigenetically, HDACi upregulate *Myod1* [96] and therefore would counteract its decreased expression and the resulting downregulated expressions of genes controlled by this TF, which we described here.

We postulate that, in dystrophic myoblasts, epigenetic dysregulation of *Myod1* expression causes a pathological cascade of downregulated transcriptions of a range of genes controlled by MyoD, with functional consequences for muscle regeneration.

If it is the absence of dystrophin in myogenic cells that determines their fate, developmental muscle abnormalities should occur. This is indeed the case: asymptomatic DMD patients already have transcriptomic alterations [10] and myogenesis in *Dmd*[mdx] embryos is severely disrupted, with myotube hypotrophy, reduced myotube numbers and displacement defects [16]. This developmental abnormality continues in adult muscles during regeneration.

Altogether, our data identify a continuum, where mutations disrupting expression of full-length dystrophin cause SC division abnormalities impacting myoblast generation but also imprinting these cells with functionalities further reducing muscle regeneration. And as dystrophin is needed at the myotube stage for the initial assembly of the DAPC and subsequent formation of viable myofibres [4], dystrophinopathy results in contraction-induced injury in mature muscle [5]. Thus, the vicious circle of disease is closed with both degenerative and regenerative abnormalities contributing to DMD progression (Figure 9).

Such a scenario explains the apparent contradiction, where removal of dystrophin and/or DAPC in fully developed myofibres does not result in dystrophic muscle damage [5,6]. It would be the case if,

once formed in a healthy muscle, DAPC remains stable even if dystrophin is subsequently lost, as indeed have been demonstrated [6].

Moreover, as the efficacy of exon skipping and gene targeting methods is much higher in proliferating cells than in myofibres, all outcomes of these treatments could arise from dystrophin re-expression in myoblasts, which later differentiate into myotubes with the functional DAPC.

Findings described here agree with the existing data that loss of dystrophin expression disrupts many downstream processes. These processes offer good targets for therapeutic interventions that are not constrained by the causative mutation [97–99]. Importantly, adjustments made in dystrophic myoblasts can reduce or even prevent occurrence of dystrophy. For example, upregulation of Jagged 1 can counteract damage in muscles without any dystrophin [8] and corrections to COUP-TFII [100], EGFR-Aurka and Ghrelin expressions [101] can reduce muscle damage.

Given that these pathways are active in myoblasts and the proteins involved are structurally incompatible with substituting the scaffolding properties of dystrophin, these findings further challenge the role of dystrophin in myofibre sarcolemma stabilization as the key pathological alteration in DMD.

Our findings, combined with these data, provide compelling foundation for the call to re-evaluate the established belief on the pathogenesis of DMD. Such reconsideration is important because most current therapeutic approaches aim at dystrophin restoration in differentiated myofibres and are often conducted in the advanced-stage disease, with very limited regenerative muscle potential. Despite decades of intensive research and numerous clinical trials, none of the candidate treatments delivered disease modifying results [102]. Part of the problem is immune responses to re-expressed dystrophin [103], that need to be tackled [104] or prevented. However, dystrophin re-expression using new technologies [105] allowing re-targeting dystrophin to myoblasts and satellite cells in younger patients or, better still, in new-borns able to develop neonatal tolerance to dystrophin, could produce effective therapies for this devastating disease.

Acknowledgements

The authors would like to thank S Arkle, Colyn Crane-Robinson and K. Zabłocki for the critical comments on the manuscript.

Author's contributions

MRFG contributed mouse experiments and transcriptomic analyses, VM and CP contributed comparative human data, MB analysed TFBS, LB performed immunolocalizations in spheroids, SR and MK supervised transcriptomic analyses and DCG conceptualised the project, supervised

experiments and wrote the final version of the manuscript. All authors contributed to data analysis and writing of the manuscript.

Conflicts of interest

None.

References

1. Hoffman EP, Fischbeck KH, Brown RH, Johnson M, Medori R, Loike JD, et al. Characterization of dystrophin in muscle-biopsy specimens from patients with Duchenne's or Becker's muscular dystrophy. N Engl J Med. 1988;318: 1363–1368. doi:10.1056/NEJM198805263182104

2. Koeks Z, Bladen CL, Salgado D, van Zwet E, Pogoryelova O, McMacken G, et al. Clinical Outcomes in Duchenne Muscular Dystrophy: A Study of 5345 Patients from the TREAT-NMD DMD Global Database. J Neuromuscul Dis. 2017;4: 293–306. doi:10.3233/JND-170280

3. Tennyson CN, Klamut HJ, Worton RG. The human dystrophin gene requires 16 hours to be transcribed and is cotranscriptionally spliced. Nat Genet. 1995;9: 184–190. doi:10.1038/ng0295-184

4. Shoji E, Sakurai H, Nishino T, Nakahata T, Heike T, Awaya T, et al. Early pathogenesis of Duchenne muscular dystrophy modelled in patient-derived human induced pluripotent stem cells. Sci Rep. 2015;5: 12831. doi:10.1038/srep12831

5. Rader EP, Turk R, Willer T, Beltrán D, Inamori K-I, Peterson TA, et al. Role of dystroglycan in limiting contraction-induced injury to the sarcomeric cytoskeleton of mature skeletal muscle. Proc Natl Acad Sci U S A. 2016. doi:10.1073/pnas.1605265113

6. Ghahramani Seno MM, Graham IR, Athanasopoulos T, Trollet C, Pohlschmidt M, Crompton MR, et al. RNAi-mediated knockdown of dystrophin expression in adult mice does not lead to overt muscular dystrophy pathology. Hum Mol Genet. 2008;17: 2622–2632. doi:10.1093/hmg/ddn162

7. Waugh TA, Horstick E, Hur J, Jackson SW, Davidson AE, Li X, et al. Fluoxetine prevents dystrophic changes in a zebrafish model of Duchenne muscular dystrophy. Hum Mol Genet. 2014;23: 4651–4662. doi:10.1093/hmg/ddu185

8. Vieira NM, Elvers I, Alexander MS, Moreira YB, Eran A, Gomes JP, et al. Jagged 1 Rescues the Duchenne Muscular Dystrophy Phenotype. Cell. 2015;163: 1204–1213. doi:10.1016/j.cell.2015.10.049

9. Dommelen P van, Dijk O van, Wilde JA de, Verkerk PH. Early developmental milestones in Duchenne muscular dystrophy. Dev Med Child Neurol. 2020;62: 1198–1204. doi:https://doi.org/10.1111/dmcn.14623

10. Pescatori M, Broccolini A, Minetti C, Bertini E, Bruno C, D'amico A, et al. Gene expression profiling in the early phases of DMD: a constant molecular signature characterizes DMD muscle from early postnatal life throughout disease progression. FASEB J Off Publ Fed Am Soc Exp Biol. 2007;21: 1210–1226. doi:10.1096/fj.06-7285com

11. Toop J, Emery AE. Muscle histology in fetuses at risk for Duchenne muscular dystrophy. Clin Genet. 1974;5: 230–233. doi:10.1111/j.1399-0004.1974.tb01687.x

12. Emery AE. Muscle histology and creatine kinase levels in the foetus in Duchenne muscular dystrophy. Nature. 1977;266: 472–473. doi:10.1038/266472a0

13. Vassilopoulos D, Emery AE. Muscle nuclear changes in fetuses at risk for Duchenne muscular dystrophy. J Med Genet. 1977;14: 13–15. doi:10.1136/jmg.14.1.13

14. Nguyen F, Cherel Y, Guigand L, Goubault-Leroux I, Wyers M. Muscle lesions associated with dystrophin deficiency in neonatal golden retriever puppies. J Comp Pathol. 2002;126: 100–108. doi:10.1053/jcpa.2001.0526

15. Bassett DI, Bryson-Richardson RJ, Daggett DF, Gautier P, Keenan DG, Currie PD. Dystrophin is required for the formation of stable muscle attachments in the zebrafish embryo. Dev Camb Engl. 2003;130: 5851–5860. doi:10.1242/dev.00799

16. Merrick D, Stadler LKJ, Larner D, Smith J. Muscular dystrophy begins early in embryonic development deriving from stem cell loss and disrupted skeletal muscle formation. Dis Model Mech. 2009;2: 374–388. doi:10.1242/dmm.001008

17. Mournetas V, Massouridès E, Dupont J-B, Kornobis E, Polvèche H, Jarrige M, et al. Myogenesis modelled by human pluripotent stem cells: a multi-omic study of Duchenne myopathy early onset. J Cachexia Sarcopenia Muscle. 2021. doi:10.1002/jcsm.12665

18. Massouridès E, Polentes J, Mangeot P-E, Mournetas V, Nectoux J, Deburgrave N, et al. Dp412e: a novel human embryonic dystrophin isoform induced by BMP4 in early differentiated cells. Skelet Muscle. 2015;5: 40. doi:10.1186/s13395-015-0062-6

19. Tidball JG. Regulation of muscle growth and regeneration by the immune system. Nat Rev Immunol. 2017;17: 165–178. doi:10.1038/nri.2016.150

20. Blau HM, Webster C, Pavlath GK. Defective myoblasts identified in Duchenne muscular dystrophy. Proc Natl Acad Sci U S A. 1983;80: 4856–4860. doi:10.1073/pnas.80.15.4856

21. Hurko O, McKee L, Zuurveld J, Swick HM. Comparison of Duchenne and normal myoblasts from a heterozygote. Neurology. 1987;37: 675–681. doi:10.1212/wnl.37.4.675

22. Yablonka-Reuveni Z, Anderson JE. Satellite cells from dystrophic (mdx) mice display accelerated differentiation in primary cultures and in isolated myofibers. Dev Dyn Off Publ Am Assoc Anat. 2006;235: 203–212. doi:10.1002/dvdy.20602

23. Yeung D, Zablocki K, Lien C-F, Jiang T, Arkle S, Brutkowski W, et al. Increased susceptibility to ATP via alteration of P2X receptor function in dystrophic mdx mouse muscle cells. FASEB J Off Publ Fed Am Soc Exp Biol. 2006;20: 610–620. doi:10.1096/fj.05-4022com

24. Sacco A, Mourkioti F, Tran R, Choi J, Llewellyn M, Kraft P, et al. Short telomeres and stem cell exhaustion model Duchenne muscular dystrophy in mdx/mTR mice. Cell. 2010;143: 1059–1071. doi:10.1016/j.cell.2010.11.039

25. Górecki DC. Dystrophin: The dead calm of a dogma. Rare Dis Austin Tex. 2016;4: e1153777. doi:10.1080/21675511.2016.1153777

26. Howard PL, Dally GY, Ditta SD, Austin RC, Worton RG, Klamut HJ, et al. Dystrophin isoforms DP71 and DP427 have distinct roles in myogenic cells. Muscle Nerve. 1999;22: 16–27. doi:10.1002/(sici)1097-4598(199901)22:1<16::aid-mus5>3.0.co;2-r

27. Young CNJ, Gosselin MRF, Rumney R, Oksiejuk A, Chira N, Bozycki L, et al. Total Absence of Dystrophin Expression Exacerbates Ectopic Myofiber Calcification and Fibrosis and Alters Macrophage Infiltration Patterns. Am J Pathol. 2019. doi:10.1016/j.ajpath.2019.09.021

28. Wertz K, Füchtbauer E-M. Dmdmdx-βgeo: A new allele for the mouse dystrophin gene. Dev Dyn. 1998;212: 229–241. doi:10.1002/(SICI)1097-0177(199806)212:2<229::AID-AJA7>3.0.CO;2-J

29. Petkov PM, Ding Y, Cassell MA, Zhang W, Wagner G, Sargent EE, et al. An efficient SNP system for mouse genome scanning and elucidating strain relationships. Genome Res. 2004;14: 1806–1811. doi:10.1101/gr.2825804

30. McGreevy JW, Hakim CH, McIntosh MA, Duan D. Animal models of Duchenne muscular dystrophy: from basic mechanisms to gene therapy. Dis Model Mech. 2015;8: 195–213. doi:10.1242/dmm.018424

31. Augusto V, Padovani C, Eduardo G, Campos R. Skeletal muscle fiber types in C57BL6J mice. J Morphol Sci. 2004;21: 89–94.

32. Morgan JE, Beauchamp JR, Pagel CN, Peckham M, Ataliotis P, Jat PS, et al. Myogenic cell lines derived from transgenic mice carrying a thermolabile T antigen: a model system for the derivation of tissue-specific and mutation-specific cell lines. Dev Biol. 1994;162: 486–498. doi:10.1006/dbio.1994.1103

33. S. Andrews. FastQC: A Quality Control Tool for High Throughput Sequence Data. 2010. Available: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

34. F. Krueger. Trim Galore! A Quality Trimming Tool for High Throughput Sequence Data. 2012. Available: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

35. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinforma Oxf Engl. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

36. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinforma Oxf Engl. 2009;25: 2078–2079. doi:10.1093/bioinformatics/btp352

37. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-014-0550-8

38. R Core Team. R: A Language and Environment for Statistical Computing. R Found Stat Comput Vienna Austria. 2017.

39. Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for computing and annotating genomic ranges. PLoS Comput Biol. 2013;9: e1003118. doi:10.1371/journal.pcbi.1003118

40.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J R Stat Soc Ser B Methodol. 1995;57: 289–300. doi:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

41.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R package for comparing biological themes among gene clusters. Omics J Integr Biol. 2012;16: 284–287. doi:10.1089/omi.2011.0118

42.   Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms. PLOS ONE. 2011;6: e21800. doi:10.1371/journal.pone.0021800

43.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res. 2019;47: D607–D613. doi:10.1093/nar/gky1131

44.   Piechota M, Korostynski M, Ficek J, Tomski A, Przewlocki R. Seqinspector: position-based navigation through the ChIP-seq data landscape to identify gene expression regulators. BMC Bioinformatics. 2016;17: 85. doi:10.1186/s12859-016-0938-4

45.   ENCODE Project Consortium. The ENCODE (ENCyclopedia Of DNA Elements) Project. Science. 2004;306: 636–640. doi:10.1126/science.1105136

46.   Chiron S, Tomczak C, Duperray A, Lainé J, Bonne G, Eder A, et al. Complex interactions between human myoblasts and the surrounding 3D fibrin-based matrix. PloS One. 2012;7: e36173. doi:10.1371/journal.pone.0036173

47.   Corona BT, Machingal MA, Criswell T, Vadhavkar M, Dannahower AC, Bergman C, et al. Further development of a tissue engineered muscle repair construct in vitro for enhanced functional recovery following implantation in vivo in a murine model of volumetric muscle loss injury. Tissue Eng Part A. 2012;18: 1213–1228. doi:10.1089/ten.TEA.2011.0614

48.   Mei C, Chao C-W, Lin C-W, Li ST, Wu K-H, Yang K-C, et al. Three-dimensional spherical gelatin bubble-based scaffold improves the myotube formation of H9c2 myoblasts. Biotechnol Bioeng. 2019;116: 1190–1200. doi:10.1002/bit.26917

49.   Kim W, Lee H, Lee J, Atala A, Yoo JJ, Lee SJ, et al. Efficient myotube formation in 3D bioprinted tissue construct by biochemical and topographical cues. Biomaterials. 2020;230: 119632. doi:10.1016/j.biomaterials.2019.119632

50.   Chen B, You W, Wang Y, Shan T. The regulatory role of Myomaker and Myomixer-Myomerger-Minion in muscle development and regeneration. Cell Mol Life Sci CMLS. 2020;77: 1551–1569. doi:10.1007/s00018-019-03341-9

51.   Stern-Straeter J, Bonaterra GA, Kassner SS, Zügel S, Hörmann K, Kinscherf R, et al. Characterization of human myoblast differentiation for tissue-engineering purposes by quantitative gene expression analysis. J Tissue Eng Regen Med. 2011;5: e197–e206. doi:https://doi.org/10.1002/term.417

52.   Magli A, Baik J, Mills LJ, Kwak I-Y, Dillon BS, Gonzalez RM, et al. Time-dependent Pax3-mediated chromatin remodeling and cooperation with Six4 and Tead2 specify the skeletal myogenic lineage in developing mesoderm. PLOS Biol. 2019;17: e3000153. doi:10.1371/journal.pbio.3000153

53. Koo JH, Smiley MA, Lovering RM, Margolis FL. Bex1 knock out mice show altered skeletal muscle regeneration. Biochem Biophys Res Commun. 2007;363: 405–410. doi:10.1016/j.bbrc.2007.08.186

54. Qin CY, Cai H, Qing HR, Li L, Zhang HP. Recent advances on the role of long non-coding RNA H19 in regulating mammalian muscle growth and development. Yi Chuan Hered. 2017;39: 1150–1157. doi:10.16288/j.yczz.17-193

55. Sakamoto K, Yamaguchi S, Ando R, Miyawaki A, Kabasawa Y, Takagi M, et al. The nephroblastoma overexpressed gene (NOV/ccn3) protein associates with Notch1 extracellular domain and inhibits myoblast differentiation via Notch signaling pathway. J Biol Chem. 2002;277: 29399–29405. doi:10.1074/jbc.M203727200

56. Li D, Niu Z, Yu W, Qian Y, Wang Q, Li Q, et al. SMYD1, the myogenic activator, is a direct target of serum response factor and myogenin. Nucleic Acids Res. 2009;37: 7059–7071. doi:10.1093/nar/gkp773

57. Harrer M, Lührs H, Bustin M, Scheer U, Hock R. Dynamic interaction of HMGA1a proteins with chromatin. J Cell Sci. 2004;117: 3459–3471. doi:10.1242/jcs.01160

58. Qiu H, Zhong J, Luo L, Tang Z, Liu N, Kang K, et al. Regulatory Axis of miR-195/497 and HMGA1-Id3 Governs Muscle Cell Proliferation and Differentiation. Int J Biol Sci. 2017;13: 157–166. doi:10.7150/ijbs.17440

59. Haines BP, Gupta R, Jones CM, Summerbell D, Rigby PWJ. The NLRR gene family and mouse development: Modified differential display PCR identifies NLRR-1 as a gene expressed in early somitic myoblasts. Dev Biol. 2005;281: 145–159. doi:10.1016/j.ydbio.2005.01.030

60. Kishioka Y, Thomas M, Wakamatsu J-I, Hattori A, Sharma M, Kambadur R, et al. Decorin enhances the proliferation and differentiation of myogenic cells through suppressing myostatin activity. J Cell Physiol. 2008;215: 856–867. doi:10.1002/jcp.21371

61. Mochida Y, Parisuthiman D, Pornprasertsuk-Damrongsri S, Atsawasuwan P, Sricholpech M, Boskey AL, et al. Decorin modulates collagen matrix assembly and mineralization. Matrix Biol J Int Soc Matrix Biol. 2009;28: 44–52. doi:10.1016/j.matbio.2008.11.003

62. Doran P, Dowling P, Lohan J, McDonnell K, Poetsch S, Ohlendieck K. Subproteomics analysis of Ca+-binding proteins demonstrates decreased calsequestrin expression in dystrophic mouse skeletal muscle. Eur J Biochem. 2004;271: 3943–3952. doi:10.1111/j.1432-1033.2004.04332.x

63. Vallejo-Illarramendi A, Toral-Ojeda I, Aldanondo G, López de Munain A. Dysregulation of calcium homeostasis in muscular dystrophies. Expert Rev Mol Med. 2014;16: e16. doi:10.1017/erm.2014.17

64. Farea M, Rani AQM, Maeta K, Nishio H, Matsuo M. Dystrophin Dp71ab is monoclonally expressed in human satellite cells and enhances proliferation of myoblast cells. Sci Rep. 2020;10: 17123. doi:10.1038/s41598-020-74157-y

65. Kim S, Lee M-J, Choi J-Y, Park D-H, Kwak H-B, Moon S, et al. Roles of Exosome-Like Vesicles Released from Inflammatory C2C12 Myotubes: Regulation of Myocyte Differentiation and Myokine Expression. Cell Physiol Biochem Int J Exp Cell Physiol Biochem Pharmacol. 2018;48: 1829–1842. doi:10.1159/000492505

66. Dumont NA, Wang YX, von Maltzahn J, Pasut A, Bentzinger CF, Brun CE, et al. Dystrophin expression in muscle stem cells regulates their polarity and asymmetric division. Nat Med. 2015;21: 1455–1463. doi:10.1038/nm.3990

67. Róg J, Oksiejuk A, Gosselin MRF, Brutkowski W, Dymkowska D, Nowak N, et al. Dystrophic mdx mouse myoblasts exhibit elevated ATP/UTP-evoked metabotropic purinergic responses and alterations in calcium signalling. Biochim Biophys Acta Mol Basis Dis. 2019;1865: 1138–1151. doi:10.1016/j.bbadis.2019.01.002

68. Onopiuk M, Brutkowski W, Wierzbicka K, Wojciechowska S, Szczepanowska J, Fronk J, et al. Mutation in dystrophin-encoding gene affects energy metabolism in mouse myoblasts. Biochem Biophys Res Commun. 2009;386: 463–466. doi:10.1016/j.bbrc.2009.06.053

69. Onopiuk M, Brutkowski W, Young C, Krasowska E, Róg J, Ritso M, et al. Store-operated calcium entry contributes to abnormal Ca^{2+} signalling in dystrophic mdx mouse myoblasts. Arch Biochem Biophys. 2015;569: 1–9. doi:10.1016/j.abb.2015.01.025

70. Dumont NA, Wang YX, Rudnicki MA. Intrinsic and extrinsic mechanisms regulating satellite cell function. Dev Camb Engl. 2015;142: 1572–1581. doi:10.1242/dev.114223

71. Young CNJ, Sinadinos A, Lefebvre A, Chan P, Arkle S, Vaudry D, et al. A novel mechanism of autophagic cell death in dystrophic muscle regulated by P2RX7 receptor large-pore formation and HSP90. Autophagy. 2015;11: 113–130. doi:10.4161/15548627.2014.994402

72. Wang Y, Marino-Enriquez A, Bennett RR, Zhu M, Shen Y, Eilers G, et al. Dystrophin is a tumor suppressor in human cancers with myogenic programs. Nat Genet. 2014;46: 601–606. doi:10.1038/ng.2974

73. Megeney LA, Kablar B, Perry RL, Ying C, May L, Rudnicki MA. Severe cardiomyopathy in mice lacking dystrophin and MyoD. Proc Natl Acad Sci U S A. 1999;96: 220–225. doi:10.1073/pnas.96.1.220

74. Schuierer MM, Mann CJ, Bildsoe H, Huxley C, Hughes SM. Analyses of the differentiation potential of satellite cells from myoD-/-, mdx, and PMP22 C22 mice. BMC Musculoskelet Disord. 2005;6: 15. doi:10.1186/1471-2474-6-15

75. Zhang H, Wen J, Bigot A, Chen J, Shang R, Mouly V, et al. Human myotube formation is determined by MyoD-Myomixer/Myomaker axis. Sci Adv. 2020;6. doi:10.1126/sciadv.abc4062

76. Epstein JA, Lam P, Jepeal L, Maas RL, Shapiro DN. Pax3 inhibits myogenic differentiation of cultured myoblast cells. J Biol Chem. 1995;270: 11719–11722. doi:10.1074/jbc.270.20.11719

77. Hildyard JCW, Crawford AH, Rawson F, Riddell DO, Harron RCM, Piercy RJ. Single-transcript multiplex in situ hybridisation reveals unique patterns of dystrophin isoform expression in the developing mammalian embryo. Wellcome Open Res. 2020;5: 76. doi:10.12688/wellcomeopenres.15762.2

78. Lidov HG, Byers TJ, Watkins SC, Kunkel LM. Localization of dystrophin to postsynaptic regions of central nervous system cortical neurons. Nature. 1990;348: 725–728. doi:10.1038/348725a0

79. Kino Y, Satoh J, Ishiura S. Molecular Mechanisms of Myotonic Dystrophy: RNA-Mediated Pathogenesis and RNA-Binding Proteins. In: Takahashi MP, Matsumura T, editors. Myotonic Dystrophy: Disease Mechanism, Current Management and Therapeutic Development. Singapore: Springer; 2018. pp. 19–43. doi:10.1007/978-981-13-0508-5_2

80. Chan LY, Mugler CF, Heinrich S, Vallotton P, Weis K. Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability. Hinnebusch AG, Manley JL, Parker R, editors. eLife. 2018;7: e32536. doi:10.7554/eLife.32536

81. Sudmant PH, Rausch T, Gardner EJ, Handsaker RE, Abyzov A, Huddleston J, et al. An integrated map of structural variation in 2,504 human genomes. Nature. 2015;526: 75–81. doi:10.1038/nature15394

82. Kamieniarz-Gdula K, Proudfoot NJ. Transcriptional Control by Premature Termination: A Forgotten Mechanism. Trends Genet TIG. 2019;35: 553–564. doi:10.1016/j.tig.2019.05.005

83. Chang NC, Chevalier FP, Rudnicki MA. Satellite Cells in Muscular Dystrophy - Lost in Polarity. Trends Mol Med. 2016;22: 479–496. doi:10.1016/j.molmed.2016.04.002

84. Dumont NA, Rudnicki MA. Targeting muscle stem cell intrinsic defects to treat Duchenne muscular dystrophy. NPJ Regen Med. 2016;1. doi:10.1038/npjregenmed.2016.6

85. Ribeiro AF, Souza LS, Almeida CF, Ishiba R, Fernandes SA, Guerrieri DA, et al. Muscle satellite cells and impaired late stage regeneration in different murine models for muscular dystrophies. Sci Rep. 2019;9: 11842. doi:10.1038/s41598-019-48156-7

86. Chang NC, Sincennes M-C, Chevalier FP, Brun CE, Lacaria M, Segalés J, et al. The Dystrophin Glycoprotein Complex Regulates the Epigenetic Activation of Muscle Stem Cell Commitment. Cell Stem Cell. 2018;22: 755-768.e6. doi:10.1016/j.stem.2018.03.022

87. Brenman JE, Chao DS, Xia H, Aldape K, Bredt DS. Nitric oxide synthase complexed with dystrophin and absent from skeletal muscle sarcolemma in Duchenne muscular dystrophy. Cell. 1995;82: 743–752. doi:10.1016/0092-8674(95)90471-9

88. Colussi C, Mozzetta C, Gurtner A, Illi B, Rosati J, Straino S, et al. HDAC2 blockade by nitric oxide and histone deacetylase inhibitors reveals a common target in Duchenne muscular dystrophy treatment. Proc Natl Acad Sci U S A. 2008;105: 19183–19187. doi:10.1073/pnas.0805514105

89. Minetti GC, Colussi C, Adami R, Serra C, Mozzetta C, Parente V, et al. Functional and morphological recovery of dystrophic muscles in mice treated with deacetylase inhibitors. Nat Med. 2006;12: 1147–1150. doi:10.1038/nm1479

90. Tucci V, Isles AR, Kelsey G, Ferguson-Smith AC, Erice Imprinting Group. Genomic Imprinting and Physiological Processes in Mammals. Cell. 2019;176: 952–965. doi:10.1016/j.cell.2019.01.043

91. Zoghbi HY, Beaudet AL. Epigenetics and Human Disease. Cold Spring Harb Perspect Biol. 2016;8: a019497. doi:10.1101/cshperspect.a019497

92. Tan X, Rotllant J, Li H, De Deyne P, DeDeyne P, Du SJ. SmyD1, a histone methyltransferase, is required for myofibril organization and muscle contraction in zebrafish embryos. Proc Natl Acad Sci U S A. 2006;103: 2713–2718. doi:10.1073/pnas.0509503103

93. Tosic M, Allen A, Willmann D, Lepper C, Kim J, Duteil D, et al. Lsd1 regulates skeletal muscle regeneration and directs the fate of satellite cells. Nat Commun. 2018;9: 366. doi:10.1038/s41467-017-02740-5

94. Scionti I, Hayashi S, Mouradian S, Girard E, Esteves de Lima J, Morel V, et al. LSD1 Controls Timely MyoD Expression via MyoD Core Enhancer Transcription. Cell Rep. 2017;18: 1996–2006. doi:10.1016/j.celrep.2017.01.078

95. Saccone V, Consalvi S, Giordani L, Mozzetta C, Barozzi I, Sandoná M, et al. HDAC-regulated myomiRs control BAF60 variant exchange and direct the functional phenotype of fibro-adipogenic progenitors in dystrophic muscles. Genes Dev. 2014;28: 841–857. doi:10.1101/gad.234468.113

96. Mal A, Sturniolo M, Schiltz RL, Ghosh MK, Harter ML. A role for histone deacetylase HDAC1 in modulating the transcriptional activity of MyoD: inhibition of the myogenic program. EMBO J. 2001;20: 1739–1753. doi:10.1093/emboj/20.7.1739

97. Percival JM, Whitehead NP, Adams ME, Adamo CM, Beavo JA, Froehner SC. Sildenafil reduces respiratory muscle weakness and fibrosis in the mdx mouse model of Duchenne muscular dystrophy. J Pathol. 2012;228: 77–87. doi:10.1002/path.4054

98. Vidal B, Ardite E, Suelves M, Ruiz-Bonilla V, Janué A, Flick MJ, et al. Amelioration of Duchenne muscular dystrophy in mdx mice by elimination of matrix-associated fibrin-driven inflammation coupled to the αMβ2 leukocyte integrin receptor. Hum Mol Genet. 2012;21: 1989–2004. doi:10.1093/hmg/dds012

99. Sinadinos A, Young CNJ, Al-Khalidi R, Teti A, Kalinski P, Mohamad S, et al. P2RX7 purinoceptor: a therapeutic target for ameliorating the symptoms of duchenne muscular dystrophy. PLoS Med. 2015;12: e1001888. doi:10.1371/journal.pmed.1001888

100. Xie X, Tsai SY, Tsai M-J. COUP-TFII regulates satellite cell function and muscular dystrophy. J Clin Invest. 2016;126: 3929–3941. doi:10.1172/JCI87414

101. Wang YX, Feige P, Brun CE, Hekmatnejad B, Dumont NA, Renaud J-M, et al. EGFR-Aurka Signaling Rescues Polarity and Regeneration Defects in Dystrophin-Deficient Muscle Stem Cells by Increasing Asymmetric Divisions. Cell Stem Cell. 2019;24: 419-432.e6. doi:10.1016/j.stem.2019.01.002

102. Rodrigues M, Yokota T. An Overview of Recent Advances and Clinical Applications of Exon Skipping and Splice Modulation for Muscular Dystrophy and Various Genetic Diseases. Methods Mol Biol Clifton NJ. 2018;1828: 31–55. doi:10.1007/978-1-4939-8651-4_2

103. Vila MC, Novak JS, Benny Klimek M, Li N, Morales M, Fritz AG, et al. Morpholino-induced exon skipping stimulates cell-mediated and humoral responses to dystrophin in mdx mice. J Pathol. 2019;248: 339–351. doi:10.1002/path.5263

104. Sharma D, Al-Khalidi R, Edgar S, An Q, Wang Y, Young C, et al. Co-delivery of indoleamine 2,3-dioxygenase prevents loss of expression of an antigenic transgene in dystrophic mouse muscles. Gene Ther. 2017;24: 113–119. doi:10.1038/gt.2016.82

105. Mangeot PE, Risson V, Fusil F, Marnef A, Laurent E, Blin J, et al. Genome editing in primary cells and in vivo using viral-derived Nanoblades loaded with Cas9-sgRNA ribonucleoproteins. Nat Commun. 2019;10: 45. doi:10.1038/s41467-018-07845-z

106.   Rapaport D, Fuchs O, Nudel U, Yaffe D. Expression of the Duchenne muscular dystrophy gene products in embryonic stem cells and their differentiated derivatives. J Biol Chem. 1992;267: 21289–21292.

107.   Gonzalez JP, Kyrychenko S, Kyrychenko V, Schneider JS, Granier CJ, Himelman E, et al. Small Fractions of Muscular Dystrophy Embryonic Stem Cells Yield Severe Cardiac and Skeletal Muscle Defects in Adult Mouse Chimeras. Stem Cells Dayt Ohio. 2017;35: 597–610. doi:10.1002/stem.2518

108.   Young CNJ, Brutkowski W, Lien C-F, Arkle S, Lochmüller H, Zabłocki K, et al. P2X7 purinoceptor alterations in dystrophic mdx mouse muscles: relationship to pathology and potential target for treatment. J Cell Mol Med. 2012;16: 1026–1037. doi:10.1111/j.1582-4934.2011.01397.x

109.   Young CNJ, Chira N, Róg J, Al-Khalidi R, Benard M, Galas L, et al. Sustained activation of P2X7 induces MMP-2-evoked cleavage and functional purinoceptor inhibition. J Mol Cell Biol. 2018;10: 229–242. doi:10.1093/jmcb/mjx030

110.   Howard ZM, Lowe J, Blatnik AJ, Roberts D, Burghes AHM, Bansal SS, et al. Early Inflammation in Muscular Dystrophy Differs between Limb and Respiratory Muscles and Increases with Dystrophic Severity. Am J Pathol. 2021;191: 730–747. doi:10.1016/j.ajpath.2021.01.008

111.   Al-Khalidi R, Panicucci C, Cox P, Chira N, Róg J, Young CNJ, et al. Zidovudine ameliorates pathology in the mouse model of Duchenne muscular dystrophy via P2RX7 purinoceptor antagonism. Acta Neuropathol Commun. 2018;6. doi:10.1186/s40478-018-0530-4

112.   Otto T, Candido SV, Pilarz MS, Sicinska E, Bronson RT, Bowden M, et al. Cell cycle-targeting microRNAs promote differentiation by enforcing cell-cycle exit. Proc Natl Acad Sci. 2017 [cited 1 Jan 2021]. doi:10.1073/pnas.1702914114

Figure Legends

Figure 1: **Differential gene expression between Dmd^{mdx} and WT mouse primary myoblasts**

a) Sample-based hierarchical clustering heatmap based on the top 500 genes with the highest standard deviation for Dmd^{mdx} vs. WT based on RNA Seq data. Colour represents dissimilarity between samples, based on the Euclidean distance, from dark blue (0) for identical samples to white (40) for the most distinct.

b) Volcano plot for Dmd^{mdx} versus WT primary mouse myoblast differential gene expression analysis. Circles represent individual genes, with colour representing significance and direction of dysregulation (blue-significantly downregulated; red-significantly upregulated; black-not significantly dysregulated). Circle position represents fold change on a log2 scale for the x-axis and adjusted p-value on a -log10 scale for the y-axis. Horizontal dotted line represents the threshold of an adjusted

p-value of 5.0e-25.0e-2 or lower while the vertical doted lines represent the threshold for the absolute log2 fold change of 1.

c) Clustered heatmap of genes of interest from the dysregulated gene list of *Dmd*^mdx vs. WT primary myoblasts. Colour represents the z-score such that each gene has a mean of 0 and standard deviation of 1 to allow direct comparisons.

Figure 2: **Correlation of significantly dysregulated gene expressions between *Dmd*^mdx and *Dmd*^mdx-βgeo vs. WT.**

a) Sample-based hierarchical clustering heatmap based on the top 500 genes with the highest standard deviation for *Dmd*^mdx-βgeo vs. *Dmd*^mdx Colour represents dissimilarity between samples based on the Euclidean distance, from dark blue (0) for identical samples to white (40) for the most distinct. Note the absence of sample segregation into two groups corresponding to genotypes.

b) Volcano plot for *Dmd*^mdx-βgeo vs. *Dmd*^mdx primary mouse myoblast differential gene expression analysis. Circles represent individual genes, with colour representing significance and direction of dysregulation (blue-significantly downregulated; red-significantly upregulated; black-not significantly dysregulated). Circle position represents fold change on a log2 scale for the x-axis and adjusted *p*-value on a -log10 scale for the y-axis. Horizontal dotted line represents the threshold of an adjusted *p*-value of 5.0e-2 or lower while the vertical doted lines represent the threshold for the absolute log2 fold change of 1.

c) log2 fold change values (*Dmd*^mdx versus WT on the x-axis and *Dmd*^mdx-βgeo versus WT on the y-axis) for genes significantly dysregulated in one or both dystrophic primary mouse myoblasts vs. WT. Pearson's correlation coefficients (r) and *p*-value are shown. Green circles represent genes significantly dysregulated in *Dmd*^mdx-βgeo vs. WT, blue circles represent genes significantly dysregulated in *Dmd*^mdx vs. WT and red circles represent gernes significantly dysregulated in both.

Figure 3: **Overrepresentation of TF binding sites in genes downregulated in *Dmd*^mdx myoblasts.**

Averaged ChIP-Seq signal histograms for each of the significantly overrepresented TFBS in downregulated genes. Histograms are centred around the transcription start site (TSS) of each gene. x-axis – distance upstream and downstream from the TSS of each gene, y-axis - ChIP-Seq signal for each location averaged for all gene promoters in the submitted list; red line - downregulated genes, blue line - upregulated genes, green line - reference signal (based on 1000 random gene promoters); Bonferroni-corrected p-value of TF signal overrepresentation vs reference: *** < 1.0e-3; ** < 1.0e-2.

a) MYOD (adjusted p-value: 2.9e^{-21}, track GEO accession GSM915165), b) TCF12 (1e^{-19}, GSM915178), c) MYOG (5.6e^{-19}, GSM915164); d) TCF3 (0.0033, GSM915177); e) Heatmap of normalised TF-binding signal for each of the TFs overrepresented in genes downregulated in Dmd^{mdx} myoblasts. Genes that did not show significant binding above background (as provided by the seqinspector tool) were assigned TF binding signal of 0.

Figure 4: **Results of GO category enrichment analysis for biological processes on significantly downregulated gene lists from Dmd^{mdx} and $Dmd^{mdx-\beta geo}$ versus WT differential gene expression analyses.**

a) Bubble plot of the overlapping GO categories enriched in Dmd^{mdx} and $Dmd^{mdx-\beta geo}$ versus WT downregulated genes lists. Bubble size represents the number of genes from the downregulated gene lists belonging to the GO category and the bubble colour depicts the adjusted p-value from red (0) to blue (8.0e-3). **b**) Bubble plot of GO categories enriched in Dmd^{mdx} versus WT downregulated genes lists following REVIGO redundancy filtering (See methods). Bubble size represents the number of genes from the downregulated gene lists belonging to the GO category and the bubble colour depicts the adjusted p-value from red (0) to blue (1.0e-2).

Figure 5: **Dystrophic primary myoblasts show functional abnormalities.**

a) Proliferation analysis in myoblasts from Dmd^{mdx}, $Dmd^{mdx-\beta geo}$, C57BL/10 and C57BL/6 mice using a BrdU incorporation assay. (Left) Representative examples of BrdU flow cytometric dot plots from C57BL/10 and Dmd^{mdx} cells after 6 hours of incorporation, with FL-1 channel corresponding to BrdU fluorescence intensity and FSC channel denoting the cell size. (Right) Graph showing significantly increased proliferation of dystrophic myoblasts presented as ratio of BrdU-positive dystrophic cells relative to the respective wild type controls. Errors bars represent mean ± SEM, n = 3 or 4, * = $p \leq$ 5.0e-2, ** = $p < 1.0e-2$ (Student's unpaired t-test). **b**) Cell migration analysis in myoblasts from Dmd^{mdx}, $Dmd^{mdx-\beta geo}$, and respective wild type controls. Cells were seeded on a trans-well insert and allowed to migrate towards the bottom well containing either proliferation medium or serum-free medium complemented with cytokines. Graph shows significantly reduced migration of dystrophic myoblasts represented as the relative number of cells present on the well-side of the membrane after 12 hours of migration. Error bars represent mean ± SEM, n = 3 or 4, * = $p < 5.0e-2$, ** = $p <$ 1.0e-2 and *** = $p < 1.0e-3$ (Student's unpaired t-test).

c) Altered expression of differentiation markers in *Dmd*^{mdx} compared to C57BL/10 myoblasts. Results of qPCR expression analyses of *Myh1, Myog and Mymk* markers at specific timepoints over the 6-day period of spheroid differentiation. Individual values for biological replicates normalised to *Gapdh* expression and relative to wild type values at the commencement of differentiation (t=0) are shown. Error bars represent mean ± SEM, n = 3, * = $p \leq$ 5.0e-2, ** = $p <$ 1.0e-2 and **** = $p <$ 1.0e-4 (two-way ANOVA was used to determine the statistical significance between groups and time-points over the 6-day period and Fisher's LSD was used to determine which timepoints exhibited a significant difference between groups).

Figure 6: **Correlation of significantly dysregulated gene expressions between mouse dystrophic and WT myoblast cell lines (SC5 vs. IMO) and of significantly enriched GO categories in the downregulated gene list for primary *Dmd*^{mdx} myoblasts and corresponding cell lines.**

a) Sample-based hierarchical clustering heatmap, colour represents dissimilarity between samples based on the Euclidean distance, from black for identical samples to white for more distinct samples.

b) Volcano plot, showing the results of differential expression analysis between SC5 and IMO. Circles represent individual genes, with their colour representing significance and direction of dysregulation (blue-significantly downregulated; red-significantly upregulated; black-not significantly dysregulated). Circle position represents fold change on a log2 scale for the x-axis and adjusted *p*-value on a -log10 scale for the y-axis. Horizontal dotted line represents the threshold of an adjusted *p*-value of 5.0e-2 while the vertical doted lines represent the threshold for the absolute log2 fold change of 1.

c) Venn diagrams representing the overlap between genes dysregulated in SC5 and primary *Dmd*^{mdx} myoblasts vs. WT as well as the overlap between GO categories for biological processes overrepresented in both downregulated gene lists. Number in the white circle enumerates dysregulated genes or enriched GO categories in SC5 vs. IMO, black circle those of *Dmd*^{mdx} vs. WT and grey area numbers represent overlap between these two sets. d) Bubble plot of the overlapping GO categories following redundancy filtering applied to the 61 categories identified (panel c). Bubble size represents the number of genes from the downregulated gene lists belonging to the category and bubble colour represents the adjusted *p*-value from red (0) to blue (1.0e-2).

Figure 7: **Correlation of significantly dysregulated gene expressions between human and mouse dystrophic and WT primary myoblasts and of the corresponding significantly enriched GO categories in the downregulated gene list for the human and Dmd^{mdx} myoblast data.**

a) Sample-based hierarchical clustering heatmap, represented as described before.

b) Volcano plot, showing the results of differential expression analysis between DMD and healthy myoblasts. Circles represent individual genes, with their colour representing significance and direction of dysregulation (blue-significantly downregulated; red-significantly upregulated; black-not significantly dysregulated). Circle position represents fold change on a log2 scale for the x-axis and adjusted *p*-value on a -log10 scale for the y-axis. Horizontal dotted line represents the threshold of an adjusted *p*-value of 5.0e-2 while the vertical doted lines represent the threshold for the absolute log2 fold change of 1.

c) log2 fold change values (*Dmd*^{mdx} versus WT on the x-axis and human DMD versus WT on the y-axis) for genes significantly dysregulated in one or both dystrophic vs. WT primary myoblasts. Pearson's correlation coefficients (r) and *p*-values are shown. Green circles represent genes significantly dysregulated in DMD, blue circles represent genes significantly dysregulated in *Dmd*^{mdx} vs. WT and red circles represent genes significantly dysregulated in both. The number of genes found in each quadrant is shown.

d) Venn diagrams representing the overlap between genes dysregulated in primary human DMD and *Dmd*^{mdx} myoblasts vs. WT as well as the overlap between GO categories for biological processes overrepresented in both downregulated gene lists. Number in the white circle enumerates dysregulated genes or enriched GO categories in human DMD vs. healthy, black circle those of *Dmd*^{mdx} vs. WT and grey area numbers represent overlaps between these two sets.

e) Bubble plot of the overlapping GO categories following redundancy filtering applied to the 61 categories identified (panel d). Bubble size represents the number of genes from the downregulated gene lists belonging to the GO category and the bubble colour depicts the adjusted *p*-value from red (0) to blue (7.0e-3).

Figure 8: **Correlation of enriched GO categories between primary *Dmd*^{mdx} mouse myoblasts, *Dmd*^{mdx} mouse myoblast cell line, and human primary DMD myoblasts.**

a) Venn diagram representing the overlap between GO categories for biological processes overrepresented in downregulated gene lists in primary human DMD, *Dmd*^{mdx} myoblasts and

established *Dmd*[mdx] myoblast cell line (SC5). Numbers in each area enumerate enriched GO categories in individual datasets and those overlapping (see the graphic legend).

b) Bubble plot of the overlapping GO categories following redundancy filtering applied to the 49 categories identified (panel a). Bubble size represents the number of genes from the downregulated gene lists belonging to the given GO category and bubble colour depicts the adjusted *p*-value from red (0) to blue (8.0e-3).

Abstract: **Natural history of DMD.**

(1) Embryonic stem cells (ES) express dystrophin Dp71 [106] but its loss is not linked with arrested development. Yet, minor admixture of ES lacking Dp427 results in severe dystrophy [107]: **Defects in myogenic stem cells** (SC) descendant from DMD ES must be responsible, because in myofibre syncytium, lack of dystrophin caused by small proportion of DMD nuclei would be compensated and damage prevented. Early mesoderm committed human cells express Dp412e [18] and (**2**)**.** DMD manifests before entry into the skeletal muscle compartment [17]. (**3**) DMD abnormalities are found in dystrophic human [11–13], dog [14], zebrafish [15] and mouse embryos, with stem cell dysfunctions (hyperproliferation and death), disrupted myotube formation [16] and fibrosis as intrinsic features in developing muscle [17]. Typical dystrophic abnormalities (e.g., high serum CK) occur in new-borns years before diagnosis [9,10].

In **adult muscle**, healthy myogenic cells, myotubes and myofibres express Dp427 and DAPC assembles in differentiating fibres, supporting their functions. Impaired muscle regeneration occurs due to: **(4)** Cell-autonomous defects in dystrophic satellite cells (SC) affecting asymmetric cell divisions [24,66,70], **(5)** hyperproliferation of dystrophic myoblasts (this work) and purinergic hypersensitivity [25,108], **(6)** anomalous migration of DMD myoblasts: reduced under physiological conditions (this work) but augmented in the inflammatory environment [109] and **(7)** accelerated dystrophic myoblast differentiation (this work and Yablonka-Reuveni and Anderson, 2006) with defects in differentiating myotubes lacking Dp427 [4]. **(8)** Resulting mature muscle lacking DAPC show contraction-induced injury but dystrophin ablation in fully differentiated myofibres does not trigger muscular dystrophy [5,6]. **(9)** Chronic inflammation exacerbates muscle damage [99,110,111], but specific inflammatory mediators are also needed for regeneration [19]. **(10)** Muscle fibrosis intensifies, as the progressing loss of regenerative potential results in myofibres being replaced by the connective tissue.

Figure S1: **Differential expression of muscle programme markers in primary myoblasts from _Dmd_^{mdx} and _Dmd_^{mdx-βgeo} mice.**

Results of qPCR expression analyses of indicated markers relative to wild type, normalized to _Gapdh,_ shown as individual datapoints for biological replicates. Error bars represent mean ± SEM, n = 3, * = p < 5.0e-2, ** = p ≤ 1.0e-2, *** = p ≤ 1.0e-3, **** = p ≤ 1.0e-4 (Student's unpaired t-test).

Figure S2: **Immunodetection of the sarcomeric myosin heavy chain 4 in _Dmd_^{mdx} and BL/10 in differentiated spheroids.**

Representative immunofluorescence micrographs of sections across spheroids differentiated for 7 days showing presence of MYH4 protein (green). Cell nuclei stained with DAPI (blue). Scale bar = 50µm).

Figure S3: **Decreased cell migration of dystrophic established mouse myoblasts.**

In the trans-well assay, cells were migrating towards the well containing proliferation medium. Graph shows relative values of individual replicates revealing significantly reduced migration of dystrophic cells compared to their WT equivalent. Error bars represent mean ± SEM, n = 4, ** = p < 1.0e-2 (Student's unpaired t-test).

Figure S4: **Immunodetection of Dp427 in WT primary mouse myoblasts.**

Representative immunofluorescence micrographs of sections across C57BL/10 _tibialis anterior_ muscle and primary _Dmd_^{mdx} myoblasts stained with an antibody detecting epitopes in exons 31/32 of Dp427. Note the expected immunofluorescence signal (green) under the sarcolemma of myofibers. and absence of any discernible dystrophin immunoreactivity in myoblasts in culture, reacted and imaged under conditions identical to those used for muscle sections. Negative (no primary antibody) control shown. I indicates the specificity of the staining. Cell nuclei stained with DAPI (blue). Scale bar = 50 µm.

Figure S5: **STRING interaction diagram of genes in the GO category: Negative regulation of gene expression, epigenetic.**

Interaction network representing genes in GO:0045814. Each node depicts a gene; edges between nodes indicate interactions between protein products of the corresponding genes and the

aggregated interaction scores from experimental data and interaction databases (see the Edge Confidence graphic legend for details). Genes significantly downregulated in dystrophic myoblasts are highlighted in red and the common interactor between all significantly downregulated genes (Kdma1) is highlighted in yellow. Isolated nodes are not shown.

Figure S6: **Decreased expression of Dp71 transcript in primary myoblasts from *Dmd*^{mdx} mice.**

Results of qPCR expression assay shown as normalised individual datapoints for biological replicates relative to wild type, normalised to *Gapdh*. Error bars represent mean ± SEM, n = 3, * = p<5.0e-2 (Student's unpaired t-test).

Table legends

Table S1: **Genes significantly dysregulated in *Dmd*^{mdx} vs. WT primary myoblasts.**

From left to right, columns show ENSEMBL gene id, gene symbol, log2 fold change between *Dmd*^{mdx} and WT primary myoblasts, adjusted p-value as output by DESeq2, average FPKM for *Dmd*^{mdx} and WT groups. Genes are ordered from most downregulated to most upregulated.

Table S2: **Tabulation of genes significantly dysregulated in *Dmd*$^{mdx-βgeo}$ vs. WT primary myoblasts.**

From left to right, columns show ENSEMBL gene id, gene symbol, log2 fold change between *Dmd*$^{mdx-βgeo}$ and WT primary myoblasts, adjusted p-value as output by DESeq2, average FPKM for *Dmd*$^{mdx-βgeo}$ and WT groups. Genes are ordered from most downregulated to most upregulated.

Table S3: **Identification of putative molecular regulators of genes altered in Dmd^{mdx} myoblasts.**

The overrepresentation of binding sites for transcriptional regulators in the promoter region of the altered transcripts was identified using the seqinspector online resource (see Materials and Methods).

A) TFBS analysis results for transcripts upregulated in *Dmd*^{mdx} myoblasts. Column legend: id - ChIP-Seq track id; name - TF name, followed by the index of ChIP-Seq track for this TF; p-value - t-test p-value for overrepresentation of ChIP-Seq signal; Bonferroni - Bonferroni-corrected p-value; foldchange - a ratio of ChIP-Seq signal averaged across all promoters in the gene list versus 1000 random gene promoters; backgroundmean - averaged signal from the track across 1000 random

gene promoters; querymean - averaged signal across all promoters from the submitted list. B. TFBS analysis results for transcripts downregulated in Dmd^{mdx} myoblasts. columns - as in A. **C)** top 10 genes for each of the significantly overrepresented TFs for transcripts downregulated in Dmd^{mdx}. **D-G)** TFBS analysis results (column legend as in A) for the following gene lists: upregulated in $Dmd^{mdx-\beta geo}$; downregulated in $Dmd^{mdx-\beta geo}$; upregulated in SC-5; downregulated in SC-5. **H)** gene lists as submitted to the seqinspector tool, including gene symbols that were translated for database compatibility.

Table S4: **GO categories significantly enriched in the downregulated gene lists from primary Dmd^{mdx} vs. WT and $Dmd^{md-\beta geo}$ vs. WT.**

Columns A and D show the GO category ID, columns B and E show GO category description and columns C and F show adjusted p-value for GO categories enriched in the significantly downregulated genes in primary Dmd^{mdx} and $Dmd^{mdx-\beta geo}$ myoblasts, respectively. GO categories are ordered from the lowest (top) to the highest (bottom) p-value for each analysis and GO categories significantly enriched in both analyses are highlighted in red.

Table S5: **Genes significantly dysregulated in SC5 (dystrophic) vs. IMO (WT) myoblast cell line.**

From left to right, columns show ENSEMBL gene id, gene symbol, log2 fold change between SC5 and IMO primary myoblasts, adjusted p-value as outputted by DESeq2, average FPKM for SC5 and IMO groups. Genes are ordered from most downregulated to most upregulated.

Table S6: **GO categories significantly enriched in the downregulated gene lists from primary Dmd^{mdx} vs. WT and SC5 vs. IMO cell lines.**

Columns A and D show the GO category ID, columns B and E show GO description and columns C and F show adjusted p-value for GO categories enriched in the significantly downregulated genes in SC5 myoblast cell line and primary Dmd^{mdx} myoblasts, respectively. GO categories are ordered from the lowest (top) to the highest (bottom) p-value for each analysis and GO categories significantly enriched in both analyses are highlighted in red.

Table S7: **Genes significantly dysregulated in DMD vs. healthy primary human myoblasts.**

From left to right, columns show ENSEMBL gene id, gene symbol, log2 fold change between DMD and healthy human primary myoblasts, adjusted *p*-value as outputted by DESeq2, average FPKM for human DMD and health groups. Genes are ordered from most downregulated to most upregulated.

Table S8: **GO categories significantly enriched in the downregulated gene lists from dystrophic vs. WT human primary myoblasts and *Dmd*^{mdx} vs. WT mouse primary myoblasts.**

Columns A and D show GO category ID, columns B and E show GO category description and columns C and F show adjusted *p*-values for GO categories enriched in the significantly downregulated genes in primary human DMD and mouse *Dmd*^{mdx} myoblasts, respectively. GO categories are ordered from the lowest (top) to the highest (bottom) *p*-value for each analysis and GO categories significantly enriched in both analyses are highlighted in red.

Table S9: **Interactions between genes in the GO category GO:0045814: "Negative regulation of gene expression, epigenetic" generated in STRING.**

Columns A and B show nodes connected by an edge, column C shows the interaction score if one has been determined experimentally, column D shows the interaction score if an interaction is present in a database feeding into STRING and column E shows the combined interaction score.

Table S10: **Primers used in real-time quantitative PCR analysis.**

Primers were designed specifically except *Gapdh* primer set, which was taken from [112].