Landscaping the behavioural ecology of primate stone tool use

Katarina Almeida-Warren[a,b*], Tetsuro Matsuzawa[c,d], Susana Carvalho[a,b,e,f]

[a]Primate Models for Behavioural Evolution Lab, Institute of Human Sciences, University of Oxford, Oxford, UK

[b]Interdisciplinary Center for Archaeology and Evolution of Human Behaviour (ICArEHB), Universidade do Algarve, Faro, Portugal

[c]Chubu Gakuin University, Japan

[d]Division of the Humanities and Social Sciences, California Institute of Technology, Pasadena, CA, USA.

[e]CEF (Centre for Functional Ecology), Coimbra, Portugal

[f]Gorongosa National Park, Sofala, Mozambique

[*]Corresponding Author

E-mail: katarina.almeida-warren@anthro.ox.ac.uk

Author contributions

KAW is the main author and contributor and was responsible for conceptualization, methodology, data collection, formal analysis, data curation, visualization and writing of the original manuscript. TM provided resources and editorial advice. SC provided supervision, data and resources and contributed towards conceptualization, methodology and revisions of the manuscript.

Abstract

Ecology is fundamental to the development, transmission, and perpetuity of primate technology. Previous studies on tool site selection have addressed the relevance of targeted resources and raw materials for tools, but few have considered the broader foraging landscape. In this first landscape-scale study of the ecological contexts of wild chimpanzee (*Pan troglodytes verus*) tool-use, we investigate the conditions required for nut-cracking to occur and persist over time at discrete locations in Bossou (Guinea). We examine this at three levels: selection, frequency of use, and inactivity. We find that, further to the presence of a nut tree and availability of raw materials, abundance of food-providing trees as well as proximity to nest sites were significant predictors of nut-cracking occurrence. This suggests that the spatial distribution of nut-cracking sites is mediated by the broader behavioural landscape and is influenced by non-extractive foraging of predictable resources, as well as non-foraging activities. Additionally, the number of functional tools was greater at sites with higher frequency of nut-cracking and was negatively correlated with site inactivity. Our findings indicate that the technological landscape of the Bossou chimpanzees shares affinities with the 'favoured places' model of hominin site formation and provides new insights for reconstructing ancient patterns of landscape use.

Key words: behavioural ecology, landscape-use, nut-cracking, primate archaeology, wild chimpanzees

Résumé

L'écologie est fondamentale pour le développement, la transmission et la pérennité de la technologie des primates. Des études antérieures ont identifié la disponibilité des ressources cibles ainsi que les matières premières pour les outils comme des facteurs

influents dans la sélection des emplacements pour les activités technologiques. Cependant,

il y a peu d'études qui abordent cette recherche à l'échelle du paysage et du comportement

fourrager. Dans cette première étude paysagère sur l'utilisation d'outils par le chimpanzé

sauvage (*Pan troglodytes verus*), nous recherchons les conditions écologiques qui influencent

la sélection, l'utilisation et l'inactivité des emplacements utilisés pour le cassage des noix en

Bossou, Guinée. Nos résultats montrent qu'en plus de la présence d'un noyer et de la

disponibilité des matières premières, l'abondance d'arbres nourriciers ainsi que la proximité

des sites de nidification étaient des prédicteurs significatifs de l'occurrence du cassage des

noix. Cela suggère que la distribution spatiale des sites de cassage de noix est influencée par

le paysage comportemental et est influencée par le fourrage non-extractive de ressources

prévisibles, ainsi que par des activités non-fourragers. Nos résultats indiquent que le paysage

technologique des chimpanzés de Bossou partage des affinités avec le modèle des « lieux

favoris » de la formation des sites hominidés et fournit de nouvelles perspectives pour

reconstruire les modes d'utilisation du paysage anciens.

Introduction

Ecology plays an important role in shaping non-human primate behaviour from foraging

strategies, to ranging patterns, and sociality (Robbins and Hohmann, 2006; Strier, 2011). Tool-

use, particularly for extractive foraging, is no exception. Recent studies have highlighted that

ecology is key in determining whether tool-use emerges in a population, how it manifests itself,

and how it is maintained once it is established (S. Carvalho et al., 2007, 2011; Grund et al.,

2019; Koops et al., 2013, 2014).

Stone tool-use is often recurrent in spatially discrete locations, frequently involves the

reuse of tools, and leaves a recognisable archaeological footprint that can be traced back

thousands of years (Falótico et al., 2019; Mercader et al., 2007). However, little is known about

the ecological factors influencing selection and repeated use of specific locations for these

activities and how they fit within the broader foraging landscape. Lithic-based foraging

technology has been recorded in several wild, non-human primate species including

chimpanzees (*Pan troglodytes ssp.*; Whiten *et al.*, 1999), bearded capuchin monkeys

(*Sapajus libidinosus;* Ottoni and Izar, 2008), Burmese long-tailed macaques (*Macaca*

fascicularis; Gumert, Kluck and Malaivijitnond, 2009), and, most recently, in white-faced

capuchins (*Cebus capuchinus*; Barrett *et al.*, 2018).

Chimpanzees are of particular interest because they are our closest living relatives

(Langergraber et al., 2012), and they present the largest, most diverse and ecologically

adaptable technological repertoire compared to any other non-human species, reflecting a

level of cognitive flexibility akin to the earliest hominin toolmakers (S. Carvalho et al., 2013;

Pascual-Garrido and Almeida-Warren, 2021; Rolian and Carvalho, 2017). Chimpanzee nut-

cracking assemblages have been found to have close similarities to the low-density

assemblages characteristic of the early hominin record (S. Carvalho et al., 2008; S. Carvalho

and McGrew, 2012). Thus, understanding how patterns of nut-cracking behaviour accumulate

across the landscape can provide valuable insights into the formation and spatial distribution

of early hominin assemblages and allow the modelling of ancient landscape use and resource

exploitation.

Previous research on chimpanzee nut-cracking has established that the spatial

availability of nut trees and raw materials for tools influences site location and reuse, as well

as frequency and distance of tool transport (S. Carvalho et al., 2007, 2011). Nevertheless,

nut-cracking assemblages are yet to be explored within the context of the broader ecological

and foraging landscape. This requires the study of nut-cracking sites not only in relation to

direct ecological correlates such as access to raw materials and nuts, but also in relation to

ecological requirements of other daily activities critical to survival such as food, water, and

shelter.

Chimpanzees living in forested environments spend approximately 50% of their waking

hours foraging and travelling between feeding locations (Pruetz and Bertolani, 2009). Recent

studies have shown that chimpanzee ranging patterns are dynamic and are influenced by the

spatial distribution and seasonality of food (Trapanese et al., 2019), which may also determine

where non-foraging activities, such as nesting, take place (Basabose and Yamagiwa, 2002;

Hernandez-Aguilar, 2009; Janmaat et al., 2014). Their diet mainly consists of fruit (Morgan

and Sanz, 2006), but high-energy foods such as insects, nuts and honey acquired through

tool-assisted foraging are also important staples or nutritional supplements for many

chimpanzee populations (Sanz and Morgan, 2013). Yet, little is known about how extractive

foraging interacts with other feeding activities and the broader behavioural landscape. This

study explores this question for the first time in the context of chimpanzee nut-cracking using

stone tools.

Water is essential to life (Popkin et al., 2010). For non-human primate species living in

extremely arid conditions, such as savannah-dwelling chimpanzees and baboons, water is a

critical resource that constrains movement patterns and landscape use (Barton et al., 1992;

Pruetz and Herzog, 2017; Wessling et al., 2018). It has also featured in many discussions

surrounding early hominin evolution and behaviour (Joordens et al., 2019), and has been

spatially linked with early stone tool sites (Rogers et al., 1994). Similarly, the location of

chimpanzee nut-cracking sites has been suggested to coincide with the proximity to

hydrological features, such as streams and rivers (S. Carvalho et al., 2007), possibly because

they are the most likely sources of eroded lithic raw materials for tools (S. Carvalho, 2011),

although this remains to be empirically tested.

Chimpanzees habitually make a sleeping nest at the end of every day, and sometimes

make day nests for resting (Koops et al., 2012). They have been an important focus of

research since Sept (1992) recognised that they form clusters of debris akin to early hominin

assemblages, and saw their potential for understanding patterns of early hominin landscape

use and the origins of human shelter (Mcgrew, 2021). Nesting locations have subsequently

been linked with a range of ecological parameters such as tree species and tree architecture,

as well as surrounding topography and vegetation types (Badji et al., 2018; J. S. Carvalho et

al., 2015; Hernandez-Aguilar, 2009; Hernandez-Aguilar et al., 2013; Koops et al., 2012;

Ndiaye et al., 2018; Stewart et al., 2011). Other studies have found additional links between

nest sites and proximity to areas of high fruit availability (Basabose and Yamagiwa, 2002;

Furuichi and Hashimoto, 2004; Goodall, 1962; Janmaat et al., 2014). While the exact

conditions determining nest site suitability appears to be population-specific rather than

universal, these findings demonstrate that nesting activities shape chimpanzee landscape

use, and, in turn, are shaped by resource distribution and the local environment. Nevertheless,

little is known about how nesting relates to other activities such as spatially discreet forms of

tool use (e.g., nut-cracking and termite-fishing).

This is the first landscape-scale investigation of the ecological drivers of chimpanzee

tool-use, where we examine the conditions required for nut-cracking to occur and persist over

time in discrete locations at the long-term field site of Bossou (Guinea). This is divided into

three points of enquiry: 1) tool site selection; 2) tool site use; 3) tool site inactivity.

Tool site selection explores the ecological conditions that determine where nut-cracking

sites are established within the chimpanzee home-range. This is critical to understanding how

technological activities occur within the broader behavioural landscape, and how these

activities produce locally discrete assemblages that can remain archaeologically identifiable

for thousands of years (e.g. Falótico et al., 2019; Mercader et al., 2007). There is a growing

body of research on habitat selection for daily activities such as foraging, travelling, socializing

and sleeping, particularly within the context of anthropogenic landscapes and the implications

for conservation (Bryson-Morrison et al., 2017; K. B. Potts et al., 2016). However, regarding

technological activities, although we are beginning to learn more about raw material selection

for tools (S. Carvalho et al., 2008; Pascual-Garrido and Almeida-Warren, 2021), the selection

of locations for tool use remains unexplored.

Tool site use focuses on the ecological factors that influence the frequency with which

established sites attract nut-cracking activity. How often a site is used can serve as a proxy

for inferring preference in relation to other sites. From an archaeological perspective, generally

the more a location is used for debris-generating activities, the larger and more conspicuous

an archaeological signature becomes. This has important implications for understanding the

spatial clustering of activities and material evidence over time and can offer many insights into

the formation and perpetuity of early hominin archaeological assemblages (S. Carvalho and

Almeida-Warren, 2019; McGrew, 2010). Ethoarchaeological studies of chimpanzee nests

have found that sleeping sites are frequently revisited and the nests themselves may be

reused (Hernandez-Aguilar, 2009; Sept, 1992; Stewart et al., 2011). However, comparable

literature on the use of tool sites is scarce, except for the reuse of stone tools (S. Carvalho et

al., 2009), and sources of perishable raw materials (Pascual-Garrido, 2018; Pascual-Garrido

and Almeida-Warren, 2021).

Tool site inactivity investigates the ecological conditions that may cause the cessation

of nut-cracking activity at an established tool site. Chimpanzees live in dynamic landscapes

shaped by environmental change. Whether natural or anthropogenic, these shifts often result

in changes in resource distributions that may in turn lead to the development of new

behavioural adaptations and ranging patterns (e.g. Kalan et al., 2020). New opportunities may

arise (e.g., foraging food from crops: Hockings et al., 2015, 2012; nesting in novel plant

species: McCarthy et al., 2017), while formerly habitual activities may shift to new locations or

become obsolete (Gruber et al., 2012; Kühl et al., 2019). Environmental changes have been

identified as important drivers of our own evolutionary history (Bobe et al., 2002; Bobe and

Carvalho, 2019; Joordens et al., 2019; R. Potts, 1998; R. Potts et al., 2020; Reed, 1997), but

few studies have addressed empirically how these changes may have affected patterns of

landscape use and the distribution of early hominin tool sites (e.g. Rogers et al., 1994). Thus,

investigating the conditions that influence the cessation of activity at chimpanzee tool sites,

can provide important clues as to the factors that lead to their temporary or long-term

abandonment.

Throughout these steps we assess the effect of ecological parameters that have been

found to correlate with nut-cracking activities (nut availability; abundance of raw materials;

distance to water) as well as variables that encapsulate two key aspects of chimpanzee activity

patterns: non-extractive foraging (abundance of food resources: wild food trees; wild fruit

trees; THV - terrestrial herbaceous vegetation) and sleeping (distance to nesting site).

 # Methods

 ## Study site and subjects

Bossou (7° 39' N, 8° 30' W) is located in the southeast of the Republic of Guinea (West

Africa), 6 km from the foothills of Mount Nimba Strict Nature Reserve (Figure 1) (Humle,

2011a; Yamakoshi and Sugiyama, 1995). The chimpanzee community has been studied

continuously in both natural (since 1976) and experimental (outdoor laboratory since 1988)

settings (Tetsuro Matsuzawa et al., 2011; Sugiyama and Koman, 1979). Between 1976 and

2003 the population size ranged from 18 to 23 individuals (Sugiyama, 2004), but has since

declined largely due to a catastrophic flu-like epidemic from which it never recovered (Humle,

Figure 1 – Map of the Bossou Forest and surrounding area. Mount Nimba Strict Nature Reserve shapefile adapted from the WDPA database (UNEP-WCMC and IUCN, 2019)

2011b; Sugiyama and Fujita, 2011). At the time of this study, the population consisted of seven

individuals. The Bossou chimpanzees habitually crack and consume oil-palm nuts (*Elaeis*

guineensis) and are currently the only population known to use portable stones as both

hammers and anvils (Carvalho, Matsuzawa and McGrew, 2013; but see Ohashi, 2015 for

recent discoveries in Liberia). Bossou has two seasons – a short dry season lasting from

November to February, and a long rainy season extending from March to October (Humle,

2011a; Yamakoshi, 1998). Nut-cracking occurs year-round, but is most prevalent during peak

wet season (June – August) and at the start of the dry season (November – December) when

fruit is less abundant (Yamakoshi, 1998).

The Bossou forest has an estimated area of 16 km^2 and is intersected by roads (Figure

2; Hockings, Anderson and Matsuzawa, 2006). Within this, the chimpanzees range a core

Figure 2 – Map of the study area and surrounding area, highlighting the locations of recorded nut-cracking sites, quadrats, nests, and watercourses.

207   area of approximately 7 km^2 (Hockings et al., 2006). The habitat is comprised of a composite

208   of primary, secondary and riverine forests, savanna and cultivated fields (Hockings et al.,

209   2012). The northern slope of Mont Gban, in the Eastern part of the forest, is considered a

210   sacred area – forêt sacrée – by the local *Manon* culture, and access is forbidden to outsiders.

211   Out of respect to this tradition, no research was conducted in this area.

Data collection

Data were collected over two field trips: 14DEC17-01MAY18 and 28OCT18-13DEC18,

encompassing 160 days of fieldwork. We employed a mixed-method approach that combined

direct behavioural observation through active group follows of the chimpanzee population, with

archaeological documentation of nut-cracking sites (indirect behavioural observations), and

ecological research using the transect and quadrat method. At an initial stage, we targeted

nut-cracking sites that had previously been documented by SC in 2006 and 2008-09. Further

nut-cracking sites were discovered during surveys and group follows throughout both research

seasons. For each nut-cracking site we established 1-km transects intersecting the site datum

at 500 metres. Nut-cracking sites within 100 metres of a pre-established transect were either

assigned to that transect or became the mid-point of a new perpendicular transect, to ensure

even forest coverage. All transects were oriented N-S or E-W, except for two that were

oriented NE-SW and NW-SE due to access difficulties (Figure 2). 5-metre radius survey

quadrats were established at every 100-metres along the transects starting from the midpoint

where the nut-cracking site was located. At each quadrat nut-cracking specific and general

ecological and vegetation data were collected (further details below).

We employed a fully digital method of data collection. Quadrat datums and all data

entries (food-providing vegetation, tools, raw materials) were georeferenced using an Arrow

Gold GNSS receiver (μ_{HRMS} = 2 metres; Almeida-Warren et al., 2021; EOS Positioning

Systems Inc., 2017). Coordinates were instantly downloaded via Bluetooth to the GeoGrafi-M

application (MGISS, 2019) on an android device where further data could be entered through custom made forms.

Oil palms

For each oil palm encountered during quadrat surveys we documented diameter at breast height (DBH), and number of fruit bunches (total and ripe). For 25 of the nut trees associated to nut-cracking sites we also collected information on nut availability and new traces of nut-cracking on a weekly basis during the first field season (22JAN18-03MAY18) and once at the beginning and end of the second season (weeks of 29OCT18; 10DEC18). Additional data was collected by Henry Camara during the weeks of 30SEP19, 27APR20, and 25MAY20. As per Koops *et al.* (2013), we scored presence of edible nuts on the ground within a 2-metre radius of the nut tree: (0) nuts absent; (1) 1-50 nuts; (2) 51-100 nuts; (3) > 100 nuts. With aid from field guides, nut suitability was determined by checking a sample of randomly collected nuts for whether the nuts contained an edible kernel or were rotten (following Koops *et al.*, 2013). Nuts were not opened so as not to affect future availability, but the local people also crack oil palm nuts and are able to identify whether or not they are edible (Humle and Matsuzawa, 2004).

Tools and raw materials

All lithic material was recorded for size, raw material type, and portability (whether loose or imbedded in the ground). Adapted from Koops *et al.* (2013), size was scored into six categories: (1) 1-2 cm; (2) 3-5 cm; (3) 6-10 cm; (4) 11-20 cm; (5) 21-30 cm; (6) >30 cm. Tools and bi-products of nut-cracking were defined as stones that showed at least one of the following: a) traces of wear from nut-cracking; b) nutshell remains on or around them; c) could be refitted with another stone with evidence of a) or b). For this study, the variable *Tools* included all lithic materials used for nut-cracking excluding fragments that no longer or could no longer be used for nut-cracking.

The collective tool assemblage was also scored for status of nut-cracking activity:

(Active) New signs of nut-cracking activity were recorded during the fieldwork period. Nut

powder or cracked nut kernels were visible on top of or around tools, and there was at least

one hammer and anvil pair with impact points that had not rusted over; (Inactive) There were

no signs of recent nut-cracking activity during the entire fieldwork period. Cracked nut kernels

were either absent or present but showed clear signs of decay. Iron oxide or moss developing

on tool impact points.

Vegetation

We recorded all wild non-THV plants with a DBH > 2cm of species known to be

consumed by the chimpanzees of Bossou. We then cross-referenced the recorded species

with the current list of chimpanzee food resources to identify those which were sources of fruit

– the preferred food-type of chimpanzees. To distinguish permanent food sources from

ephemeral food sources, THV was documented separately. Domesticated or crop species

were not included in this study.

Nests

Because few nests were documented at the quadrat level or along transects, additional

forest-wide surveys were conducted through random walks and strategic walks targeting

areas that were known nesting locations. When a nest was encountered, elevation and

direction of travel were maintained along topographical contours and all nests within 50-metres

of the nest and either side of the projected route were documented until no further nests were

visible over a 50-metre stretch (modified from Hernandez-Aguilar, 2009). For data analysis,

nests were divided into spatial clusters. A cluster was defined as a 50-metre radius area with

a minimum of 21 documented nests, representing a minimum of three sleep-events for the

collective chimpanzee population (N = 7) as a proxy for a habitual nesting location. This was

achieved on QGIS (Version 2.18.4), by creating a 50-metre buffer zone around each nest and

using the 'count points in polygon' function to identify zones with a minimum of 21 nests. The

distance to nearest nest cluster variable used in the following analyses was computed with

the 'distance to nearest hub' function and was defined as the distance from a quadrat datum

to the centre of the nearest nest cluster.

Watercourses

287   Watercourse data was collected in 2008/2009 by SC. With the help of field guides,

288   streams and rivers were traced on foot and recorded using the track feature on a Garmin

289   hand-held GPS device. The *distance to nearest river* variable used in the following analyses

290   was computed in QGIS and was defined as the distance between a quadrat datum to the

291   nearest point along the watercourse polylines.

Data analysis

Tool site selection

Initial inspection of the data revealed that no nut-cracking occurred in quadrats where

nut trees were absent. This is consistent with previous literature describing that nut-cracking

occurs in close proximity to a nut tree (S. Carvalho et al., 2008). Additionally, there was very

little variation in the number of oil palms in each quadrat with only 15% of oil palm quadrats

documented with more than one oil palm. For these reasons, we restricted data analysis to

quadrats where an oil palm was present and did not include the number of oil palms as a

predictor, as this would mask the potential effect of the other variables of interest. The final

dataset had a total of 82 quadrats, 40 of which had traces of nut-cracking activity. We used a

binomial generalized linear model (GLM; Zuur *et al.*, 2009a) with a logit link function to

investigate the effect of five main predictors: raw materials, wild food trees, wild food THV,

distance to nearest nest cluster, distance to nearest river, on the presence (1) versus absence

(0) of a tool site in a given quadrat. We also analysed three sub-models to determine whether

more restricted variables yielded a better model fit (Appendix, Table A2). The first sub-model

replaced raw materials with a subset of raw materials of size class corresponding to the three

most common tool size classes (95% of tools. Size class: 3, 4, 5; Appendix, Table A1). The

second sub-model replaced wild food-providing trees with a subset formed only of fruit-

providing trees. The third sub-model included raw materials of size class 3-5 and fruit-

providing trees. Akaike's Information Criterion for small sample sizes (AICc) was used to

compare models (Burnham and Anderson, 2004), whereby the model with the lowest AICc

was chosen as the final model.

Tool site use

We investigated whether the hypothesized ecological variables (i.e., nut availability, raw

materials, food trees, distance to nearest nest cluster, and distance to nearest river) influenced

the frequency a nut-cracking site was used. From a total of 361 monitoring observations, only

35 cases of recent nut-cracking events were identified for 17 out of the 25 monitored nut-

cracking sites, where frequency of recent activity ranged between 1 and 4. Because of single

(N = 1) sample sizes for 2 and 4 events, frequency of activity was recoded as "Low" (≤ 2

events; N = 10) and "High" (> 2 events; N = 7). The small sample size (N = 17) was deemed

too small to justify a GL(M)M, therefore we only discuss descriptive statistics for this question

using two-sample t-tests (or Man-Whitney U tests when assumptions of normality were not

met).

Tool site inactivity

We used a binomial GLM with 'logit' link to investigate the effect of mean nut availability,

raw materials, and food trees, on tool-site inactivity. The response variable included nut-

cracking sites that were classified as active (response = 0; N = 24) with those classified as

inactive (response = 1; N = 16). The final dataset included 40 tool sites. Akin the model for

tool site selection, we also investigated four sub-models with raw materials of size class 3-5,

tools, and fruit tree subsets (Appendix, Table A6). The model with the lowest AICc was then

chosen as the final model.

General considerations

All analyses were processed in R Studio (version 1.1.383; R Studio Team, 2016), using

R (version 4.1.0; R Core Team, 2021). Data exploration for each GLM, following the protocol

described in Zuur, Ieno & Elphick (2010), did not raise any concerns. Collinearity among the

explanatory variables was assessed by calculating the Variance inflation factors (VIF) using

the function 'vif' of the car package (Jon Fox and Weisberg, 2011). None of the models

indicated any multicollinearity issues (Maximum VIF = 1.39, Quinn and Keough, 2002). To

assess the significance of the full models and sub-models, we ran likelihood ratio tests (LRT)

using the 'anova' function which compared each model to a corresponding null model from

which all fixed effects were excluded (Dobson, 2002). We tested the significance of main

effects for each model by systematically dropping them one at a time and comparing the

resulting model with the full model using the 'drop1' function (Dobson, 2002). P-values for the

individual effects were based on the LRT results from the 'drop1' function. The AICc for model

selection was calculated using the 'MuMIn' package (Bartón, 2020). Model assumptions were

verified by plotting residuals versus fitted values and versus each covariate in the model (Zuur

and Ieno, 2016). Influential observations were assessed by calculating and plotting the Cook's

distance (Smith and Warren, 2019); all values were under the recommended threshold of 1,

suggesting no evidence of influential points (John Fox, 2002; Smith and Warren, 2019).

For tool site use, comparisons between low and high frequency of nut-cracking activity

were computed for each of the variables of interest using unpaired two-sample t-tests or the

non-parametric equivalent, Wilcoxon rank-sum test (i.e., Mann-Whitney U test). Normality

assumptions were assessed using the Shapiro-Wilk test. Threshold for statistical significance

was set to $p \leq 0.05$.

Data availability

The datasets generated during and/or analysed during the current study will be available from the corresponding author on reasonable request. Source code will be made available at https://github.com/katarinawarren/bossou-chimps-analysis [to be replaced with DOI].

Ethical statement

All tool site and ecological data were collected when chimpanzees were absent from the survey locations. Efforts to ensure minimal disturbance of nut-cracking sites included: keeping all tools in their original locations; not removing or cracking nuts; collecting stone samples from existing tool fragments whenever possible. Research was conducted in accordance with all the research requirements of Guinea, and the ethical protocols set out by The University of Oxford, the Kyoto University Primate Research Institute, and the Institut de Recherche Environmentale de Bossou (IREB).

Results

Tool site selection

The sub-model where raw materials were replaced by a subset of size class 3-5 was the best fitted model according to the AICc (Appendix, Table A3), and had a clear effect on the probability of a nut-cracking site occurring in a location where at least one oil palm was present (full-null model comparison, LRT: $df = 5$, $deviance = 56.52$, $p < 0.001$). Raw materials had a significant positive effect on tool site prediction, as did food trees, while distance to nest cluster had a significant negative effect (Table 1; Figure 3). All other fixed effects were non-significant (Table 1). The sub-model replacing wild food trees with the fruit trees subset yielded the worst model fit in which fruit trees were not a significant predictor (Appendix, Table A3, Table A4).

Figure 3 – Probability of tool site presence in response to: a) Raw materials of size class 3 – 5; b) Trees that are sourced by chimpanzees for food; c) Distance to the nearest nest cluster.

Table 1 – Results of the final GLM investigating potential predictors influencing tool site selection

Term	Estimate	SE	χ^2	p^a
Intercept	-1.139	0.472		NI[b]
Raw materials (size class 3-5)	0.358	1.298	21.929	<0.001
Wild food trees	0.313	0.407	5.758	0.016
THV	-0.103	0.383	0.392	0.531
Distance to nearest nest cluster	-0.007	0.485	6.109	0.013
Distance to nearest river	0.005	0.411	1.397	0.237

[a] Results from the likelihood ratio test using the 'drop1' function.
[b] Not indicated because it has a limited interpretation.

Tool site use

Over a total period of 15 weeks, only 33 cases of nut-cracking were recorded for 17 out of 25 monitored tool sites. 10 of the 17 sites recorded one or two nut-cracking events (low

frequency), with the remaining seven showing recent traces between three and four times

during the monitoring period (high frequency). In general, mean nut availability was

significantly higher at nut-cracking sites that registered a higher frequency of nut-cracking

activity (T-test: *p* = 0.03; Figure 4; Table A5). Furthermore, distance to nearest nest cluster

revealed a negative trend, whereby high frequency sites tended to be nearer to nesting

locations (Wilcoxon rank-sum test: *p* = 0.07). For all other variables of interest (raw materials,

mean fruit bunches, wild food trees and distance to nearest river) there were no significant

differences between the groups (*p* > 0.15; Figure 4; Table A5).

Figure 4 – Frequency of tool site use in relative to: a) Raw materials that have been used as
tools; b) Distance to nearest nest cluster (km); c) Raw materials of size class 3 to 5; d) Wild
trees that are sourced by chimpanzees for food; e) Wild trees that are sourced by
chimpanzees for fruit; f) Distance to nearest river (km). Grey circles represent individual points,
and means indicated by diamonds. ** p < 0.05; * p < 0.07.

Tool site inactivity

Out of the sub-models, the tool subset model yielded the best fit, although the AICc for the tool and fruit trees model was only marginally higher and produced comparable results (Appendix, Table A7, Table A8). Comparison of the tool subset model with the null model was significant (LRT: *df* = 3, *deviance* = 13.20, *p* < 0.01). Overall, we found that lower values of mean nut availability and a lower number of tools were both significant predictors of tool site inactivity, while wild food trees had no effect (Figure 6; Table 3). However, the data distributions shown in Figure 6 suggest that the model is not very robust.

Figure 6 – Probability of tool site inactivity in response to: a) Mean nut availability; b) Raw materials that have been used as tools.

Table 3 – Results of the final GLM model investigating potential predictors influencing tool site abandonment

Term	Estimate	SE	χ^2	p[a]
Intercept	2.957	1.263		NI[b]
Mean. nut availability	-2.186	0.851	8.947	0.003
Tools	-0.186	0.093	5.183	0.023
Wild food trees	0.093	0.109	0.752	0.386

[a] Results from the likelihood ratio test using the 'drop1' function.
[b] Not indicated because it has a limited interpretation.

Discussion

Tool site selection

From the initial inspection of the data, it is evident that a minimum of one oil palm, specifically an oil palm in close proximity (within 10 metres), is required for nut-cracking to occur in a given location. Further to this, our results show that the abundance of raw materials and food trees as well as proximity to the nearest nest cluster are also important predictors for whether a tool site is established at an oil palm location. This suggests that, in addition to the *ecological pre-requisites* of nut-cracking, i.e., a producing oil palm and raw materials for tools, other predictable resources that form part of the chimpanzee diet (wild food-providing trees), as well as non-food related activities (sleep sites), are influential in the spatial distribution of nut-cracking locations. In contrast, THV had a negative but non-significant effect on whether tool-sites occurred. While THV is a frequently consumed food item by the chimpanzees of Bossou (Humle, 2011a), it differs from other wild plant foods in that they have low calorific value and an individual plant often can only be sourced once, after which it is permanently depleted. In contrast, food trees are replenishable, often seasonally, and constitute reliable and predictable resources that can be returned to on a seasonal basis. The fact that THV does

not bear a clear effect on tool site occurrence suggests that unpredictable food sources are

not ecological drivers of tool site selection.

Previous research has found that many primate species have goal-orientated foraging

trajectories towards spatially permanent resources and that they likely use mental maps to

guide their resource exploitation strategies (Trapanese et al., 2019). Our results provide

tentative evidence that the chimpanzees of Bossou may behave in a similar way, whereby

nut-cracking activities take place within a foraging strategy that primarily targets predictable,

high-value foods, while low-energy unpredictable foods like THV act as part of an opportunistic

strategy during forage-on-the-go.

Distance to nearest nest cluster was a significant predictor in all models, whereby the

likelihood of a tool site occurring increased with proximity to nest locations. Previous research

has found that nests sites occur in areas of high food availability (Basabose and Yamagiwa,

2002; J. S. Carvalho et al., 2015; Furuichi and Hashimoto, 2004; Goodall, 1962; Janmaat et

al., 2014). Given that nut-cracking sites are also located in areas with a greater number of

food providing trees, and bearing in mind that the Bossou chimpanzees source oil palms for a

range or other resources (e.g. fruit, pith, palm heart) and also nest in their crowns (Humle and

Matsuzawa, 2001, 2004; Yamakoshi and Sugiyama, 1995), it is possible that the relationship

between nut-cracking sites, proximity to nest sites, and food availability, is indicative that these

areas are activity hotspots, rich in resources and with habitat characteristics that are suitable

for a range of core chimpanzee activities.

Distance to the nearest river was not a significant predictor in any of the models. This

contradicts previous research in the nearby forest of Diecké, that identified that nut-cracking

locations occurred near waterlines (S. Carvalho et al., 2007). Emerging research on the role

and importance of water in shaping primate behaviour, adaptations, and landscape use, is

providing increasing evidence that there are differences in water-dependence between

populations. Rainforest-dwelling apes can usually obtain their daily hydration requirements

from the food they consume, and can go several days without drinking (Pontzer et al., 2021).

However, for primates that live in year-round or seasonally arid landscapes water is a critical

resource that shapes movement patterns and landscape use (Barton et al., 1992; Pruetz and

Herzog, 2017; Wessling et al., 2018). Fongoli chimpanzees usually drink water at least once

a day and often spend time near water sources during dry months to stay cool (Wessling,

pers. comm.; Pruetz and Bertolani, 2009). Conversely, the Bossou forest is much more humid,

with a long-wet season. It's many streams, and small forest area provide a hydrological

landscape in which chimpanzees are rarely more than 300 metres away from water.

Furthermore, during a total of ~500 hours of focal follows, the Bossou chimpanzees were only

seen to drink water on eight occasions, suggesting that they can get most of their fluids from

the foods they consume, in line with the general trend for non-human apes (Pontzer et al.,

2021).

While water was not a significant factor for Bossou, we predict that it could be a major

ecological driver regarding the spatial distribution and reuse of tool sites by savannah-living

chimpanzees. The Fongoli chimpanzees do not crack nuts, but they engage in termite-fishing,

which is also a spatially discrete technological activity tethered to the location of termite

mounds (Bogart and Pruetz, 2008, 2011), much like nut-cracking.

Nevertheless, climatic or hydrological differences cannot explain the differences

between Bossou and Diecké. Given the proximity of both field sites (approx. 50 km) and similar

climates it is unlikely that this is due to differences in aridity or water availability. However, the

chimpanzees of Diecké crack different nut species, *Panda oleosa* and *Coula edulis*, which are

absent in Bossou and may be more water dependent than the oil palm. Thus, this discrepancy

may be connected to the different plant species exploited and their respective ecology and

distribution.

Tool site use

Number of tools was the only variable of interest that differed significantly between sites

with low or high frequency of nut-cracking events. This could indicate that the visible traces of

nut-cracking found on tools act as visual cues for stimulating further nut-cracking behaviour.

The repeated use of discrete locations through stigmergy has been suggested to have led the

emergence of persistent places during the Middle Pleistocene (Matthew Pope et al., 2006;

Matthew Pope, 2017; Shaw et al., 2016). Similar hypotheses featuring local and stimulus

enhancement in chimpanzees have also been discussed as processes of social learning (e.g.

in the development of technical skills; Musgrave et al., 2020; Tennie et al., 2020; Whiten,

2021) as well why some plants are sourced more intensively than others for the manufacture

of termite fishing tools (Almeida-Warren et al., 2017). Conversely, it could indicate that

chimpanzees prefer sites with material that they are already familiar with. Previous research

in Bossou has demonstrated that chimpanzees reuse hammer-anvil pairs (tool-sets) more

often than others, that there is both group- and individual-level preference for certain tool-sets

(S. Carvalho et al., 2009), and that chimpanzees are selective of the types of materials they

use for nut-cracking (S. Carvalho et al., 2008). Analogous studies on chimpanzee plant

technologies, suggests similar patterns in the selection of materials for termite-fishing, ant-

dipping, honey gathering and water extraction (Almeida-Warren et al., 2017; Lamon et al.,

2018; Pascual-Garrido et al., 2012; Pascual-Garrido and Almeida-Warren, 2021).

Distance to nearest nest cluster showed a weak yet noteworthy difference, whereby the

frequency of nut-cracking events was marginally greater at tool sites that were closer to nest

locations. These results mirror those found for tool-site selection and offer further tentative

support that active tool sites and their frequency of use is influenced by their distribution

relative to current activity hotspots.

The number of wild food and fruit trees was largely the same for all active nut-cracking

sites. This suggest that, while food providing trees are good indicators of tool-site selection,

they may not good predictors of site use because the data collected did not capture temporal

changes in food availability or frequency of foraging activity. On the other hand, nests are

temporary features that rarely preserve for longer than six months in non-savannah

environments (Ihobe, 2005; Kamgang et al., 2020; Zamma and Makelele, 2012). Therefore,

they are a better spatial proxy for recent ranging patterns and possibly explains why

differences were found for nests, but not for vegetation.

Some consideration needs to be given as to the low number of weekly traces of nut-

cracking events recorded per tool site during the 15-weeks of monitoring. This is partially due

to the fact that not all active tool sites were monitored, with a further seven traces were found

through indirect observations at non-monitored sites. Our data indicates that a minimum of 40

nut-cracking events took place at natural nut-cracking sites during the 15 week monitoring

period, averaging approximately three events per week, which may be sufficient for the

existing chimpanzee population.

Furthermore, out of the nine nut-cracking events witnessed during group follows, six

took place at the outdoor laboratory, where nuts and stones were being artificially provisioned

for another project. The outdoor laboratory is located at the intersection of several routes which

the chimpanzees frequently travel through to access different parts of the forest. As a location

that has always experienced a high degree of natural thoroughfare (Tetsuro Matsuzawa,

2011), it may represent a pre-existing activity hotspot that has been enhanced by the

guaranteed encounter of tools and edible nuts. This could explain why a comparatively higher

number of nut-cracking events were observed there, similar to patterns recorded by Hockings

et al. (2009).

Nevertheless, due to small sample sizes and the limitations of the statistics employed,

more data are needed to explore these hypotheses further. For future research, it would be

important to investigate over a longer timescale whether and how often chimpanzees visited

the part of the forest where the tool site is located. This could make the use of complementary

data from camera traps placed in strategic locations, as full-day focal follows are not permitted

in Bossou.

Tool site inactivity

528   Understanding the contexts of tool site inactivity is an important step in investigating the

529   conditions required for nut-cracking to occur and persist over time in a particular location, and

530   the factors that might lead to their abandonment. Our data suggests that mean nut availability,

used as a proxy for tree productivity, and a high abundance of tools are important in maintaining the active status of a nut-cracking site. However, there are clear exceptions that appear to not quite fit the model (Figure 6), suggesting that other factors that were not considered in the analysis may also be at play.

The Bossou forest suffers from a great deal of human activity, particularly slash-and-burn agriculture, which leads to frequent and rapid changes in the spatial distribution of resources and localized vegetation composition (Hockings, 2011). While oil palms are not cut down during this process and are highly resistant to fire (Yamakoshi, 2011), the changes in the surrounding landscape and the increase in human presence may deter chimpanzees from visiting those areas, especially if they are near the forest boundary. Conversely, cultivated land that contains desirable food items (e.g. banana, mango, papaya) can often attract chimpanzees (Hockings, 2011), and perhaps, under these conditions, the chimpanzees prioritize the prized fruit over nuts that can be found almost anywhere. Site inactivity could also be an artefact of population decline, whereby fewer resources are sufficient to sustain the entire population. Previous literature has suggested that the Bossou forest has a carrying capacity for around 20 chimpanzees (Sugiyama and Fujita, 2011), so it is possible that the current population may no longer need to depend as highly on nuts to supplement their diets. A future longitudinal comparison drawing from historical and contemporary data will help investigate and test this further.

Conclusions

Our results indicate that proximity to a nut tree, an abundance of raw materials and predictable resources, as well as proximity to a nesting site are important ecological parameters for the establishment of a nut-cracking site in a given location. Distance to nearest nest cluster was also correlated with frequency of nut-cracking, which could potentially indicate that nesting sites are important anchors for ranging and activity patterns. Similarly, tool availability was significantly correlated with tool site use, as well as tool site inactivity, suggests

that familiarity of materials used for tools or the visual cues of tool use could be important in

the persistence of nut-cracking activities once a site has been established. While there was

no significant difference in nut availability among oil palms at active sites, the odds of tool site

inactivity were greater when mean nut availability was low, potentially indicating that a decline

in oil palm productivity at nut-cracking sites is driver of sites disuse. Together, these results

postulate that nut-cracking in Bossou is not only tethered to locations that provide the

necessary resources for this activity but is also intimately connected to a broader foraging and

behavioural landscape that is mediated by the spatio-temporal availability of primary target

resources, such as predictable food-providing trees, as well as the distribution of frequently

used nesting locations.

Preliminary comparisons with other sites regarding the importance of ecological features

such as the effect of water on tool use and ranging patterns, suggests that the ecology of

chimpanzee technology is context-specific and should be examined with this in mind. Further

studies investigating the technological landscapes of other chimpanzee populations, as well

as the integration of long-term data, will help better understand the effect of different

environmental and demographic contexts on the factors driving the spatial distribution and

reuse of tool sites, adding further detail to this picture.

While current evidence suggests that early hominin and chimpanzee lithic technologies

differ in form and function (Arroyo and de la Torre, 2016; Toth et al., 2006), it is likely they had

similar plant-dominated diets supported by insects and sporadic meat consumption (Panger

et al., 2003). Thus, it is plausible that, like chimpanzees, early hominin tool-use operated within

behavioural landscapes conditioned by localized environmental parameters, where foraging

strategies were shaped by the distribution and availability of predictable food sources, the

dietary dependence on extractive foraging and the availability of the necessary raw materials,

as well as the location of safe places for sleeping. With the aid of primate archaeological

inference, visualizing the spatial distribution of hominin lithic assemblages within this

framework will be instrumental in providing crucial insights for reconstructing the patterns of

landscape use and resource exploitation.

The present study suggests that the technological landscape of the chimpanzees of
Bossou shares affinities with the 'favoured places' model (Schick and Toth, 1993; Shick,
1987), which proposed that hominin tool sites formed at the centre of foraging areas where
hominins would process and consume food, rest and socialize, with sites being used more
intensively in areas with higher resource abundance. Such 'activity hotspots' would have acted
as ecological tethers, shaping early hominin movement and foraging patterns, which, in turn,
would have led to the formation and repeated use of tool sites over time. As the most
conspicuous evidence of these locations, stone tool assemblages may hold important clues
for uncovering behaviours beyond those associated to lithic technology and serve as starting
points to search for traces of other activities, such as sleeping, foraging, and insectivory, that
are currently extremely rare in the archaeological record. This research draws upon the work
of Glynn Isaac (e.g. Isaac, 1981; Isaac et al., 1981; Isaac and Harris, 1980) who pioneered
the application of landscape-scale approaches to the study of hominin assemblages, from
which the first concrete models of hominin site formation were developed. Further studies will
help guide future human origins research and provide an empirical framework for modelling
and testing hypotheses of early hominin behaviour associated to the archaeological record of
our earliest ancestors.

602   Acknowledgements

We are grateful to Vincent Mamy, Lawe Goigbe, Boniface Zogbila, Gouanou Zogbila,
Jules Doré, Henry Camara, and Pascal Goumi for support in the field; Dr. Jesse Van der Grient
and Dr. Alexander Mielke for statistical advice; Dr. Dora Biro and Dr. Bronwyn Tarr for helpful
comments and feedback on the original manuscript. We also thank the Direction Nationale de
la Recherche Scientifique, the Institut de Recherche Environmentale de Bossou (Guinea), and
the Kyoto University Primate Research Institute, for research permissions and logistical
support. KAW was funded by the Fundação pela Ciência e Tecnologia (grant number:
SFRH/BD/115085/2016), supported by the Programa Operacional Capital Humano (POCH)

and the European Union; the Boise Trust Fund (University of Oxford); and the National

Geographic Society (grant number: EC-399R-18); SC thanks the Leverhulme Trust for the

PLP 2016-114 which supported part of the field equipment and logistics; TM thanks the Japan

Society for the Promotion of Science (JSPS) Leading Graduate Program (U04-PWS), JSPS

core-to-core CCSN, and the Ministry of Education, Culture, Sports, Science and Technology

of Japan (MEXT)/JSPS-KAKENHI (grant numbers: #07102010, #12002009, #16002001,

#20002001, #24000001, #16H06283).

Competing interests

619   The authors declare no competing interests.

Bibliography

Almeida-Warren, K., Braun, D. R., and Carvalho, S. (2021). The DistoX2: A methodological

solution to archaeological mapping in poorly accessible environments. *Journal of*

Archaeological Science: Reports, *35*, 102688.

https://doi.org/10.1016/J.JASREP.2020.102688

Almeida-Warren, K., Sommer, V., Piel, A. K., and Pascual-Garrido, A. (2017). Raw material

procurement for termite fishing tools by wild chimpanzees in the Issa valley, Western

Tanzania. *American Journal of Physical Anthropology*, *164*(2), 292–304.

https://doi.org/10.1002/ajpa.23269

Arroyo, A., and de la Torre, I. (2016). Assessing the function of pounding tools in the Early

Stone Age: A microscopic approach to the analysis of percussive artefacts from Beds I

and II, Olduvai Gorge (Tanzania). *Journal of Archaeological Science*, *74*, 23–34.

https://doi.org/10.1016/j.jas.2016.08.003

Badji, L., Ndiaye, P. I., Lindshield, S. M., Ba, C. T., and Pruetz, J. D. (2018). Savanna

chimpanzee (*Pan troglodytes verus*) nesting ecology at Bagnomba (Kedougou, Senegal).

636 *Primates*, *59*(3), 235–241. https://doi.org/10.1007/s10329-017-0647-2

637 Barrett, B. J., Monteza-Moreno, C. M., Dogandžic, T., Zwyns, N., Ibáñez, A., and Crofoot, M.

638  C. (2018). Habitual stone-tool-aided extractive foraging in white-faced capuchins, *Cebus*

639  *capucinus.* *Royal* *Society:* *Open* *Science,* *5*(181002).

640  https://doi.org/10.1098/rsos.181002

641 Bartón, K. (2020). *MuMIn: Multi-Model Inference. R package version 1.43.17.*

642 Barton, R. A., Whiten, A., Strum, S. C., Byrne, R. W., and Simpson, A. J. (1992). Habitat use

643  and resource availability in baboons. *Animal Behaviour*, *43*(5), 831–844.

644  https://doi.org/10.1016/S0003-3472(05)80206-4

645 Basabose, A. K., and Yamagiwa, J. (2002). Factors affecting nesting site choice in

646  chimpanzees at Tshibati, Kahuzi-Biega national park: Influence of sympatric gorillas.

647  *International* *Journal* *of* *Primatology*, *23*(2), 263–282.

648  https://doi.org/10.1023/A:1013879427335

649 Bobe, R., Behrensmeyer, A. K., and Chapman, R. E. (2002). Faunal change, environmental

650  variability and late Pliocene hominin evolution. *Journal of Human Evolution*, *42*(4), 475–

651  497. https://doi.org/10.1006/jhev.2001.0535

652 Bobe, R., and Carvalho, S. (2019). Hominin diversity and high environmental variability in the

653  Okote Member, Koobi Fora Formation, Kenya. *Journal of Human Evolution*, *126*, 91–105.

654  https://doi.org/10.1016/j.jhevol.2018.10.012

655 Bogart, S. L., and Pruetz, J. D. (2008). Ecological context of savanna chimpanzee (*Pan*

656  *troglodytes verus*) termite fishing at Fongoli, Senegal. *American Journal of Primatology*,

657  *70*(6), 605–612. https://doi.org/10.1002/ajp.20530

658 Bogart, S. L., and Pruetz, J. D. (2011). Insectivory of savanna chimpanzees (*Pan troglodytes*

659  *verus*) at Fongoli, Senegal. *American Journal of Physical Anthropology*, *145*(1), 11–20.

660  https://doi.org/10.1002/ajpa.21452

661 Bryson-Morrison, N., Tzanopoulos, J., Matsuzawa, T., and Humle, T. (2017). Activity and

662  Habitat Use of Chimpanzees (*Pan troglodytes verus*) in the Anthropogenic Landscape of

663  Bossou, Guinea, West Africa. *International Journal of Primatology*, *38*(2), 282–302.

https://doi.org/10.1007/s10764-016-9947-4

Burnham, K. P., and Anderson, D. R. (Eds.). (2004). *Model Selection and Multimodel Inference* (2nd ed). Springer. https://doi.org/10.1007/b97636

Carvalho, J. S., Meyer, C. F. J., Vicente, L., and Marques, T. A. (2015). Where to nest? Ecological determinants of chimpanzee nest abundance and distribution at the habitat and tree species scale. *American Journal of Primatology*, *77*(2), 186–199. https://doi.org/10.1002/ajp.22321

Carvalho, S. (2011). Diécké Forest, Guinea: Delving into Chimpanzee Behavior Using Stone Tool Surveys. In T Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The Chimpanzees of Bossou and Nimba* (pp. 301–312). Springer Japan. https://doi.org/10.1007/978-4-431-53921-6_31

Carvalho, S., and Almeida-Warren, K. (2019). Primate Archaeology. In J. Chun Choe (Ed.), *Encyclopedia of Animal Behavior* (2nd ed., Vol. 1, pp. 397–407). Elsevier. https://doi.org/10.1016/B978-0-12-809633-8.90156-0

Carvalho, S., Biro, D., McGrew, W. C., and Matsuzawa, T. (2009). Tool-composite reuse in wild chimpanzees (*Pan troglodytes*): Archaeologically invisible steps in the technological evolution of early hominins? *Animal Cognition*, *12*(1 SUPPL), S103–S114. https://doi.org/10.1007/s10071-009-0271-7

Carvalho, S., Cunha, E., Sousa, C., and Matsuzawa, T. (2008). Chaînes opératoires and resource-exploitation strategies in chimpanzee (*Pan troglodytes*) nut cracking. *Journal of Human Evolution*, *55*(1), 148–163. https://doi.org/10.1016/j.jhevol.2008.02.005

Carvalho, S., Cunha, E., Sousa, C., and Matsuzawa, T. (2011). Extensive surveys of chimpanzee stone tools: from the telescope to the magnifying glass. In Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The Chimpanzees of Bossou and Nimba* (1st ed., pp. 145–155). Springer Japan.

Carvalho, S., Matsuzawa, T., and McGrew, W. C. (2013). From pounding to knapping: How chimpanzees can help us to model hominin lithics. In C. Sanz, J. Call, and C. Boesch (Eds.), *Tool Use in Animals* (pp. 225–241). Cambridge University Press.

https://doi.org/10.1017/CBO9780511894800.015

Carvalho, S., and McGrew, W. C. (2012). The origins of the Oldowan: Why chimpanzees (*Pan*

troglodytes) still are good models for technological evolution in Africa. In M. Domínguez-

Rodrigo (Ed.), *Stone Tools and Fossil Bones: Debates in the Archaeology of Human*

Origins   (pp.   201–221).   Cambridge   University   Press.

https://doi.org/10.1017/CBO9781139149327.010

Carvalho, S., Sousa, C., and Matsuzawa, T. (2007). New Nut-Cracking Sites in Diecké Forest,

Guinea: An Overview of the Surveys. *Pan Africa News*, *14*(1).

Dobson, A. J. (2002). An introduction to generalized linear models. In *New York*. Chapman &

Hall/CRC.

EOS   Positioning   Systems   Inc.   (2017).   *Arrow   Gold*.   https://eos-gnss.com/wp-

content/uploads/2017/11/eos-arrow-gold-revC-v4.pdf

Falótico, T., Proffitt, T., Ottoni, E. B., Staff, R. A., and Haslam, M. (2019). Three thousand

years of wild capuchin stone tool use. *Nature Ecology & Evolution*, *3*(7), 1034–1038.

https://doi.org/10.1038/s41559-019-0904-4

Fox, John. (2002). *An R and S-Plus companion to applied regression*. Sage Publications.

Fox, Jon, and Weisberg, S. (2011). An R Companion to Applied Regression. In *Sage* (2nd

ed.). Sage.

Furuichi, T., and Hashimoto, C. (2004). Botanical and Topographical Factors Influencing

Nesting-Site Selection by Chimpanzees in Kalinzu Forest, Uganda. *International Journal*

of   Primatology   2004   25:4,   25(4),   755–765.

https://doi.org/10.1023/B:IJOP.0000029121.25284.7F

Goodall, J. M. (1962). Nest building behavior in the free ranging chimpanzee. *Annals of the*

New York Academy of Sciences, *102*(2), 455–467. https://doi.org/10.1111/j.1749-

6632.1962.tb13652.x

Gruber, T., Potts, K. B., Krupenye, C., Byrne, M. R., Mackworth-Young, C., McGrew, W. C.,

Reynolds, V., and Zuberbühler, K. (2012). The influence of ecology on chimpanzee (*Pan*

troglodytes) cultural behavior: A case study of five ugandan chimpanzee communities.

Journal of Comparative Psychology, *126*(4), 446–457. https://doi.org/10.1037/a0028702

Grund, C., Neumann, C., Zuberbühler, K., and Gruber, T. (2019). Necessity creates opportunities for chimpanzee tool use. *Behavioral Ecology*, *30*(4), 1136–1144. https://doi.org/10.1093/beheco/arz062

Gumert, M., Kluck, M., and Malaivijitnond, S. (2009). The Physical Characteristics and Usage Patterns of Stone Axe and Pounding Hammers Used by Long-Tailed Macaques in the Andaman Sea Region of Thailand. *American Journal of Primatology*, *608*(71), 594–608. https://doi.org/10.1002/ajp.20694

Hernandez-Aguilar, R. A. (2009). Chimpanzee nest distribution and site reuse in a dry habitat: implications for early hominin ranging. *J. Hum. Evol.*, *57*(4), 350–364. https://doi.org/10.1016/j.jhevol.2009.03.007

Hernandez-Aguilar, R. A., Moore, J., and Stanford, C. B. (2013). Chimpanzee nesting patterns in savanna habitat: Environmental influences and preferences. *American Journal of Primatology*, *75*(10), 979–994. https://doi.org/10.1002/ajp.22163

Hockings, K. J. (2011). The crop-raiders of the sacred hill. In Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The Chimpanzees of Bossou and Nimba* (pp. 211–220). Springer Japan.

Hockings, K. J., Anderson, J. R., and Matsuzawa, T. (2006). Road crossing in chimpanzees: A risky business. In *Current Biology* (Vol. 16, Issue 17). https://doi.org/10.1016/j.cub.2006.08.019

Hockings, K. J., Anderson, J. R., and Matsuzawa, T. (2009). Use of wild and cultivated foods by chimpanzees at Bossou, Republic of Guinea: feeding dynamics in a human-influenced environment. *American Journal of Primatology*, *71*(8), 636–646. https://doi.org/10.1002/ajp.20698

Hockings, K. J., Anderson, J. R., and Matsuzawa, T. (2012). Socioecological adaptations by chimpanzees, *Pan troglodytes verus*, inhabiting an anthropogenically impacted habitat. *Animal Behaviour*, *83*(3), 801–810. https://doi.org/10.1016/j.anbehav.2012.01.002

Hockings, K. J., McLennan, M. R., Carvalho, S., Ancrenaz, M., Bobe, R., Byrne, R. W.,

Dunbar, R. I. M., Matsuzawa, T., McGrew, W. C., Williamson, E. A., Wilson, M. L., Wood,

B., Wrangham, R. W., and Hill, C. M. (2015). Apes in the Anthropocene: Flexibility and

survival. In *Trends in Ecology and Evolution* (Vol. 30, Issue 4, pp. 215–222). Elsevier Ltd.

https://doi.org/10.1016/j.tree.2015.02.002

Humle, T. (2011a). Location and Ecology. In Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama

(Eds.), *The Chimpanzees of Bossou and Nimba* (1st ed., pp. 13–22). Springer Japan.

Humle, T. (2011b). The 2003 Epidemic of a Flu-Like Respiratory Disease at Bossou. In

Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The Chimpanzees of Bossou*

and Nimba (pp. 325–333). Springer Japan.

Humle, T., and Matsuzawa, T. (2001). Behavioural Diversity among the Wild Chimpanzee

Populations of Bossou and Neighbouring Areas, Guinea and Côte d'Ivoire, West Africa.

Folia Primatologica, *72*(2), 57–68. https://doi.org/10.1159/000049924

Humle, T., and Matsuzawa, T. (2004). Oil palm use by adjacent communities of chimpanzees

at Bossou and Nimba Mountains, West Africa. *International Journal of Primatology*, *25*(3),

551–581. https://doi.org/10.1023/B:IJOP.0000023575.93644.f4

Ihobe, H. (2005). Life Span of Chimpanzee Beds at the Mahale Mountains National Park,

Tanzania. *Pan Africa News*, *12*(1), 10–12.

Isaac, G. L. (1981). Stone Age visiting cards: approaches to the study of early land use

patterns. In I. Hodder, G. L. Isaac, and N. Hammond (Eds.), *Pattern of the past: studies*

in honour of David Clarke (pp. 131–155). Cambridge University Press.

Isaac, G. L., and Harris, J. W. K. (1980). A method for determining the characteristics of

artefacts between sites in the Upper Member of the Koobi Fora Formation, East Lake

Turkana. In R. E. F. Leakey and B. A. Ogot (Eds.), *Proceedings of the 8th Panafrican*

Congress on Prehistory and Quaternary Studies (pp. 19–22).

Isaac, G. L., Harris, J. W. K., and Marshall, F. (1981). Small is informative: the application of

the study of mini-sites and least effort criteria in the interpretation of the Early Pleistocene

archaeological record at Koobi Fora, Kenya. In B. Isaac (Ed.), *The Archaeology of Human*

Origins: Papers by Glynn Isaac (pp. 101–119). Cambridge University Press.

Janmaat, K. R. L., Polansky, L., Ban, S. D., and Boesch, C. (2014). Wild chimpanzees plan
their breakfast time, type, and location. *Proceedings of the National Academy of*
Sciences, *111*(46), 16343–16348. https://doi.org/10.1073/pnas.1407524111

Joordens, J. C. A., Feibel, C. S., Vonhof, H. B., Schulp, A. S., and Kroon, D. (2019). Relevance
of the eastern African coastal forest for early hominin biogeography. *Journal of Human*
Evolution, *131*, 176–202. https://doi.org/10.1016/J.JHEVOL.2019.03.012

Kalan, A. K., Kulik, L., Arandjelovic, M., Boesch, C., Haas, F., Dieguez, P., Barratt, C. D.,
Abwe, E. E., Agbor, A., Angedakin, S., Aubert, F., Ayimisin, E. A., Bailey, E., Bessone,
M., Brazzola, G., Buh, V. E., Chancellor, R., Cohen, H., Coupland, C., … Kühl, H. S.
(2020). Environmental variability supports chimpanzee behavioural diversity. *Nature*
Communications, *11*(1), 4451. https://doi.org/10.1038/s41467-020-18176-3

Kamgang, S. A., Carme, T. C., Bobo, K. S., Abwe, E. E., Gonder, M. K., and Sinsin, B. (2020).
Assessment of in situ nest decay rate for chimpanzees (*Pan troglodytes ellioti* Matschie,
1914) in Mbam-Djerem National Park, Cameroon: implications for long-term monitoring.
Primates, *61*(2), 189–200. https://doi.org/10.1007/s10329-019-00768-3

Koops, K., McGrew, W. C., de Vries, H., and Matsuzawa, T. (2012). Nest-Building by
Chimpanzees (*Pan troglodytes verus*) at Seringbara, Nimba Mountains: Antipredation,
Thermoregulation, and Antivector Hypotheses. *International Journal of Primatology*,
33(2), 356–380. https://doi.org/10.1007/s10764-012-9585-4

Koops, K., McGrew, W. C., and Matsuzawa, T. (2013). Ecology of culture: do environmental
factors influence foraging tool use in wild chimpanzees, *Pan troglodytes verus*? *Animal*
Behaviour, *85*(1), 175–185. https://doi.org/10.1016/j.anbehav.2012.10.022

Koops, K., Visalberghi, E., and van Schaik, C. P. (2014). The ecology of primate material
culture. *Biology Letters*, *10*(11), 20140508. https://doi.org/10.1098/rsbl.2014.0508

Kühl, H. S., Boesch, C., Kulik, L., Haas, F., Arandjelovic, M., Dieguez, P., Bocksberger, G.,
McElreath, M. B., Agbor, A., Angedakin, S., Ayimisin, E. A., Bailey, E., Barubiyo, D.,
Bessone, M., Brazzola, G., Chancellor, R., Cohen, H., Coupland, C., Danquah, E., …
Kalan, A. K. (2019). Human impact erodes chimpanzee behavioral diversity. *Science*,

363(6434), 1453–1455. https://doi.org/10.1126/science.aau4532

Lamon, N., Neumann, C., Gier, J., Zuberbühler, K., and Gruber, T. (2018). Wild chimpanzees

select tool material based on efficiency and knowledge. *Proceedings of the Royal Society*

B: Biological Sciences, *285*(1888), 20181715. https://doi.org/10.1098/rspb.2018.1715

Langergraber, K. E., Prufer, K., Rowney, C., Boesch, C., Crockford, C., Fawcett, K., Inoue, E.,

Inoue-Muruyama, M., Mitani, J. C., Muller, M. N., Robbins, M. M., Schubert, G., Stoinski,

T. S., Viola, B., Watts, D., Wittig, R. M., Wrangham, R. W., Zuberbuhler, K., Paabo, S.,

and Vigilant, L. (2012). Generation times in wild chimpanzees and gorillas suggest earlier

divergence times in great ape and human evolution. *Proceedings of the National*

Academy of Sciences, *109*(39), 15716–15721. https://doi.org/10.1073/pnas.1211740109

Matsuzawa, Tetsuro. (2011). Field Experiments of Tool-Use. In T Matsuzawa, T. Humle, and

Y. Sugiyama (Eds.), *The Chimpanzees of Bossou and Nimba* (pp. 157–164). Springer

Japan.

Matsuzawa, Tetsuro, Humle, T., and Sugiyama, Y. (Eds.). (2011). *The Chimpanzees of*

Bossou and Nimba. Springer Japan.

McCarthy, M. S., Lester, J. D., and Stanford, C. B. (2017). Chimpanzees (*Pan troglodytes*)

Flexibly Use Introduced Species for Nesting and Bark Feeding in a Human-Dominated

Habitat. *International Journal of Primatology*, *38*(2), 321–337.

https://doi.org/10.1007/s10764-016-9916-y

McGrew, W. C. (2010). In search of the last common ancestor: New findings on wild

chimpanzees. In *Philosophical Transactions of the Royal Society B: Biological Sciences*

(Vol. 365, Issue 1556, pp. 3267–3276). The Royal Society.

https://doi.org/10.1098/rstb.2010.0067

McGrew, W. C. (2021). Sheltering Chimpanzees. *Primates*, *1*, 3.

https://doi.org/10.1007/s10329-021-00903-z

Mercader, J., Barton, H., Gillespie, J., Harris, J., Kuhn, S., Tyler, R., and Boesch, C. (2007).

4,300-year-old chimpanzee sites and the origins of percussive stone technology.

Proceedings of the National Academy of Sciences of the United States of America,

104(9), 3043–3048. https://doi.org/10.1073/pnas.0607909104

MGISS. (2019). *GeoGrafi-M - MGISS*. https://mgiss.co.uk/product/geografi-m/

Morgan, D. B., and Sanz, C. (2006). Chimpanzee feeding ecology and comparisons with

sympatric gorillas in the Goualougo Triangle, Republic of Congo. In H. Gottfried, M. M.

Robbins, and C. Boesch (Eds.), *Feeding Ecology in Apes and Other Primates* (pp. 97–

122). Cambridge University Press.

Musgrave, S., Lonsdorf, E., Morgan, D., Prestipino, M., Bernstein-Kurtycz, L., Mundry, R., and

Sanz, C. (2020). Teaching varies with task complexity in wild chimpanzees. *Proceedings*

of   the   National   Academy   of   Sciences,   *117*(2),   969–976.

https://doi.org/10.1073/pnas.1907476116

Ndiaye, P. I., Badji, L., Lindshield, S. M., and Pruetz, J. D. (2018). Nest-Building Behaviour by

Chimpanzees (*Pan troglodytes verus*) in the Non-Protected Area of Diaguiri (Kedougou,

Senegal): Implications for Conservation. *Folia Primatologica*, *89*(5), 316–326.

https://doi.org/10.1159/000490945

Ohashi, G. (2015). Pestle-pounding and nut-cracking by wild chimpanzees at Kpala, Liberia.

Primates, *56*(2), 113–117. https://doi.org/10.1007/s10329-015-0459-1

Ottoni, E. B., and Izar, P. (2008). Capuchin monkey tool use: Overview and implications.

Evolutionary Anthropology, *17*(4), 171–178. https://doi.org/10.1002/evan.20185

Panger, M. A., Brooks, A. S., Richmond, B. G., and Wood, B. (2003). Older than the Oldowan?

Rethinking the emergence of hominin tool use. *Evolutionary Anthropology: Issues, News,*

and Reviews, *11*(6), 235–245. https://doi.org/10.1002/evan.10094

Pascual-Garrido, A. (2018). Scars on plants sourced for termite fishing tools by chimpanzees:

Towards an archaeology of the perishable. *American Journal of Primatology*, *80*(9),

e22921. https://doi.org/10.1002/ajp.22921

Pascual-Garrido, A., and Almeida-Warren, K. (2021). Archaeology of the Perishable. *Current*

Anthropology, *62*(3), 333–362. https://doi.org/10.1086/713766

Pascual-Garrido, A., Buba, U., Nodza, G., and Sommer, V. (2012). Obtaining raw material:

Plants as tool sources for Nigerian Chimpanzees. *Folia Primatologica*, *83*(1), 24–44.

Pontzer, H., Brown, M. H., Wood, B. M., Raichlen, D. A., Mabulla, A., Harris, J. A., Dunsworth,
H., Hare, B., Walker, K., Luke, A., Dugas, L. R., Schoeller, D., Plange-Rhule, J., Bovet,
P., Forrester, T. E., Thompson, M. E., Shumaker, R. W., Rothman, J. M., Vogel, E., …
Ross, S. R. (2021). Evolution of water conservation in humans. *Current Biology*, *31*, 1–
7. https://doi.org/10.1016/j.cub.2021.02.045

Pope, Matthew. (2017). Thresholds in behaviour, thresholds of visibility: landscape processes,
asymmetries in landscape records and niche construction in the formation of the
Palaeolithic record. In M Pope, C. Gamble, and J. McNabb (Eds.), *Crossing the Human*
Threshold Dynamic Transformation and Persistent Places During the Late Middle
Pleistocene. Routledge.

Pope, Matthew, Russel, K., and Watson, K. (2006). Biface form and structured behaviour in
the Acheulean. *Lithics: The Journal of the Lithic Studies Society*, *27*, 44–57.

Popkin, B. M., D'Anci, K. E., and Rosenberg, I. H. (2010). Water, hydration, and health.
Nutrition Reviews, *68*(8), 439–458. https://doi.org/10.1111/J.1753-4887.2010.00304.X

Potts, K. B., Baken, E., Levang, A., and Watts, D. P. (2016). Ecological factors influencing
habitat use by chimpanzees at Ngogo, Kibale National Park, Uganda. *American Journal*
of Primatology, *78*(4), 432–440. https://doi.org/10.1002/AJP.22513

Potts, R. (1998). Environmental hypotheses of hominin evolution. *American Journal of*
Physical Anthropology, *107*, 93–136. https://doi.org/10.1002/(SICI)1096-
8644(1998)107:27+<93::AID-AJPA5>3.0.CO;2-X

Potts, R., Dommain, R., Moerman, J. W., Behrensmeyer, A. K., Deino, A. L., Riedl, S., Beverly,
E. J., Brown, E. T., Deocampo, D., Kinyanjui, R., Lupien, R., Owen, R. B., Rabideaux,
N., Russell, J. M., Stockhecke, M., deMenocal, P., Faith, J. T., Garcin, Y., Noren, A., …
Uno, K. (2020). Increased ecological resource variability during a critical transition in
hominin evolution. *Science Advances*, *6*(43), eabc8975.
https://doi.org/10.1126/sciadv.abc8975

Pruetz, J. D., and Bertolani, P. (2009). Chimpanzee (Pan troglodytes verus) Behavioral
Responses to Stresses Associated with Living in a Savannah-Mosaic Environment:

Implications for Hominin Adaptations to Open Habitats. *PaleoAnthropology*, *2009*, 252–

262. https://doi.org/10.4207/pa.2009.art33

Pruetz, J. D., and Herzog, N. M. (2017). Savanna Chimpanzees at Fongoli, Senegal, Navigate

a Fire Landscape. *Current Anthropology*, *58*(16), S337–S350.

https://doi.org/10.1086/692112

Quinn, G. P., and Keough, M. J. (2002). *Experimental Design and Data Analysis for Biologists*.

R Studio Team. (2016). *RStudio: Integrated Development for R* (1.1.383). RStudio, Inc.

Reed, K. E. (1997). Early hominid evolution and ecological change through the African Plio-

Pleistocene. *Journal of Human Evolution*, *32*(2–3), 289–322.

https://doi.org/10.1006/jhev.1996.0106

Robbins, M. M., and Hohmann, G. (2006). Primate feeding ecology: an integrative approach.

In G. Hohmann, M. M. Robbins, and C. Boesch (Eds.), *Feeding Ecology in Apes and*

Other Primates (pp. 1–13). Cambridge University Press.

Rogers, M. J., Harris, J. W. K., and Feibel, C. S. (1994). Changing patterns of land use by

Plio-Pleistocene hominids in the Lake Turkana Basin. In *Journal of Human Evolution* (Vol.

27, Issues 1–3, pp. 139–158). https://doi.org/10.1006/jhev.1994.1039

Rolian, C., and Carvalho, S. (2017). Tool use and manufacture in the last common ancestor

of Pan and Homo. In M. Muller, R. Wrangham, and D. Pilbeam (Eds.), *Chimpanzees and*

Human Evolution (pp. 602– 644). Belknap Press of Harvard University Press.

Sanz, C., and Morgan, D. B. (2013). Ecological and social correlates of chimpanzee tool use.

Phil Trans R Soc B, *268*, 20120416. https://doi.org/10.1098/rstb.2012.0416

Schick, K. D., and Toth, N. (1993). *Making silent stones speak*. Simon and Schuster.

Sept, J. M. (1992). Was there no place like home?: A new perspective on early hominid

archaeological sites from the mapping of chimpanzee nests. *Current Anthropology*, *33*(2),

187–207. https://doi.org/10.1086/204050

Shaw, A., Bates, M., Conneller, C., Gamble, C., Julien, M.-A., McNabb, J., Pope, M., and

Scott, B. (2016). The archaeology of persistent places: the Palaeolithic case of La Cotte

de St Brelade, Jersey. *Antiquity*, *90*(354), 1437–1453.

916   https://doi.org/10.15184/aqy.2016.212

917 Shick, K. D. (1987). Modeling the formation of Early Stone Age artifact concentrations. *Journal*

918   *of Human Evolution*, *16*(7–8), 789–807. https://doi.org/10.1016/0047-2484(87)90024-8

919 Smith, C., and Warren, M. (2019). *GLMs in R for Ecology*. Amazon Publishing.

920 Stewart, F. A., Piel, A. K., and McGrew, W. C. (2011). Living archaeology: Artefacts of specific

921   nest site fidelity in wild chimpanzees. *Journal of Human Evolution*, *61*(4), 388–395.

922   https://doi.org/10.1016/j.jhevol.2011.05.005

923 Strier, K. B. (2011). *Primate behavioral ecology*. Allyn and Bacon.

924 Sugiyama, Y. (2004). Demographic parameters and life history of chimpanzees at Bossou,

925   Guinea. *American Journal of Physical Anthropology*, *124*(2), 154–165.

926   https://doi.org/10.1002/ajpa.10345

927 Sugiyama, Y., and Fujita, S. (2011). The Demography and Reproductive Parameters of

928   Bossou Chimpanzees. In Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The*

929   *Chimpanzees of Bossou and Nimba* (pp. 23–34). Springer Japan.

930 Sugiyama, Y., and Koman, J. (1979). Tool-using and -making behavior in wild chimpanzees

931   at Bossou, Guinea. *Primates*, *20*(4), 513–524. https://doi.org/10.1007/BF02373433

932 Team, R. C. (2021). *R: A language and environment for statistical computing* (4.1.0). R

933   Foundation for Statistical Computing.

934 Tennie, C., Bandini, E., van Schaik, C. P., and Hopper, L. M. (2020). The zone of latent

935   solutions and its relevance to understanding ape cultures. *Biology and Philosophy*, *35*(5),

936   55. https://doi.org/10.1007/s10539-020-09769-9

937 Toth, N., Schick, K. D., and Semaw, S. (2006). A comparative study of the stone tool-making

938   skills of Pan, Australopithecus, and Homo sapiens. In N. Toth and K. Schick (Eds.), *The*

939   *Oldowan: case studies into the earliest Stone Age* (pp. 155–222). Stone Age Institute

940   Press.

941 Trapanese, C., Meunier, H., and Masi, S. (2019). What, where and when: spatial foraging

942   decisions in primates. *Biological Reviews*, *94*(2), 483–502.

943   https://doi.org/10.1111/brv.12462

UNEP-WCMC, and IUCN. (2019). *Protected Planet: The World Database on Protected Areas*

(WDPA). www.protectedplanet.net

Wessling, E. G., Kühl, H. S., Mundry, R., Deschner, T., and Pruetz, J. D. (2018). The costs of

living at the edge: Seasonal stress in wild savanna-dwelling chimpanzees. *Journal of*

Human Evolution, *121*, 1–11. https://doi.org/10.1016/j.jhevol.2018.03.001

Whiten, A. (2021). The burgeoning reach of animal culture. *Science*, *372*(6537), eabe6514.

https://doi.org/10.1126/science.abe6514

Whiten, A., Goodall, J., McGrew, W. C., Nishida, T., Reynolds, V., Sugiyama, Y., Tutin, C. E.

G., Wrangham, R. W., and Boesch, C. (1999). Cultures in chimpanzees. *Nature*,

399(6737), 682–685. https://doi.org/10.1038/21415

Yamakoshi, G. (1998). Dietary responses to fruit scarcity of wild chimpanzees at Bossou,

Guinea: Possible implications for ecological importance of tool use. *American Journal of*

Physical   Anthropology,   *106*(3),   283–295.   https://doi.org/10.1002/(SICI)1096-

8644(199807)106:3<283::AID-AJPA2>3.0.CO;2-O

Yamakoshi, G. (2011). Pestle-pounding behavior: the key to the coexistence of humans and

chimpanzees. In Tetsuro Matsuzawa, T. Humle, and Y. Sugiyama (Eds.), *The*

Chimpanzees of Bossou and Nimba (pp. 107–115). Springer Japan.

Yamakoshi, G., and Sugiyama, Y. (1995). Pestle-pounding behavior of wild chimpanzees at

Bossou, Guinea: A newly observed tool-using behavior. *Primates*, *36*(October), 489–500.

https://doi.org/10.1007/BF02382871

Zamma, K., and Makelele, M. (2012). Comparison of the Longevity of Chimpanzee Beds

between Two Areas in the Mahale Mountains National Park, Tanzania. *Pan Africa News*,

19(2), 25–28. https://doi.org/10.5134/168176

Zuur, A. F., and Ieno, E. N. (2016). A protocol for conducting and presenting results of

regression-type analyses. *Methods in Ecology and Evolution*, *7*(6), 636–645.

https://doi.org/10.1111/2041-210X.12577

Zuur, A. F., Ieno, E. N., and Elphick, C. S. (2010). A protocol for data exploration to avoid

common statistical problems. *Methods in Ecology and Evolution*, *1*(1), 3–14.

https://doi.org/10.1111/j.2041-210X.2009.00001.x

Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., and Smith, G. M. (2009). *GLM and GAM*

for Absence–Presence and Proportional Data (pp. 245–259).

https://doi.org/10.1007/978-0-387-87458-6_10

Appendix

Table A1 – Summary of the response variables used in this study and their descriptions.

Variable	Description	Units	Range
Raw materials	Total number of stones in the survey quadrat	Counts	0-99
Raw materials (size class 3-5)	Number of stones with sizes within the 95% confidence interval of tools used of nut-cracking	Counts	0-81
Tools	Total number of stones used for cracking nuts; Excludes unused fragments	Counts	0-28
Food trees	Total number of plants know to provide food sources to Bossou chimpanzees; Excludes plants with diameter at breast hight (DBH) under 2 cm	Counts	0-12
Fruit trees	Total number of plants know to be sources of fruit to Bossou chimpanzees; Excludes plants with diameter at breast hight (DBH) under 2 cm	Counts	0-9
THV	Total number of individual terrestrial herbaceous vegetation (THV) plants that are part of the Bossou chimpanzee diet	Counts	0-10
Nut availability	Approximate amount of nuts with edible kernels; Mean nut availability used in analysis	Categorical	0 (none); 1 (< 50); 2 (50 – 100); 3 (> 100)
Distance to nearest nest cluster	Distance, in metres, to nearest cluster of 21 nests; Cluster defined as nests within a 50-metre radius	Metres	50.01-781.06
Distance to nearest river	Distance, in metres, to nearest water course	Metres	6.92-399.84

981 Table A2 – Likelihood ratio test results of the Full-Null model comparisons of the site selection

982 models.

Model	Resid. df	Resid. Dev	df	Deviance	p
~1 (Null model)	81	113.627			
~Raw materials + Wild food trees + THV + Nest +River	76	59.289	5	54.338	<0.001
~**Raw materials (size class 3-5)** + Wild food trees + THV + Nest +River	76	57.109	5	56.519	<0.001
~Raw materials + **Wild fruit trees** + THV + Nest +River	76	62.480	5	51.147	<0.001
~Raw materials (size class 3-5) + Wild fruit trees + THV + Nest +River	76	61.817	5	51.810	<0.001

984 Table A3 – Results of the AICc-Based Model Selection of the GLM for the predictors of tool

985 site selection. Subset variables highlighted in bold.

Model	df	AICc
~Raw materials + Wild food trees + THV + Nest +River	6	72.409
~**Raw materials (size class 3-5)** + Wild food trees + THV + Nest +River	6	70.229
~Raw materials + **Wild fruit trees** + THV + Nest +River	6	75.599
~Raw materials (size class 3-5) + Wild fruit trees + THV + Nest +River	6	74.937

991   Table A4 – Results of the additional GLMs investigating potential predictors of tool site

992   selection. Subset variables highlighted in bold.

Model	Term	Estimate	SE	χ^2	P [a]
~Raw materials + Wild food trees + THV + Nest +River	Intercept	-0.575	1.076		NI[b]
	Raw materials	0.195	0.057	19.748	<0.001
	Wild food trees	0.303	0.155	4.944	0.026
	THV	-0.101	0.159	0.419	0.518
	Distance to nearest nest cluster	-0.007	0.003	7.377	0.007
	Distance to nearest river	0.004	0.004	0.887	0.346
~Raw materials +**Wild fruit trees** + THV + Nest +River	Intercept	-0.305	1.052		NI[b]
	Raw materials	0.203	0.059	20.316	<0.001
	Wild fruit trees	0.241	0.193	1.753	0.185
	THV	-0.031	0.136	0.051	0.821
	Distance to nearest nest cluster	-0.008	0.003	8.469	0.004
		0.005	0.004	1.270	0.260
~**Raw materials (size class 3-5) + Wild fruit trees** + THV + Nest +River	Intercept	-0.700	1.079		
	Raw materials (size class 3-5)	0.351	0.102	20.978	0.000
	Wild fruit trees	0.199	0.202	1.049	0.306
	THV	-0.017	0.141	0.014	0.905
	Distance to nearest nest cluster	-0.007	0.003	7.715	0.005
	Distance to nearest river	0.006	0.004	1.967	0.161

[a] Results from the likelihood ratio test using the 'drop1' function.
[b] Not indicated because it has a limited interpretation.

998   Table A5 – Descriptive statistics and test results of the comparison between tool sites with low

999   frequency (N = 10) and high frequency (N = 7) of nut-cracking events.

	Low frequency		High frequency		Test results			
Variable	Mean	SD	Mean	SD	Test[a]	t	df	p
Tools	7.100	3.143	11.429	3.780	tt	2.487	11.429	0.029
Distance to nearest nest cluster (m)	245.660	132.760	148.714	64.047	tt	-2.000	13.707	0.066
Raw materials (size class 3-5)	8.000[b]	5.250[c]	11.000[b]	3.500[c]	wt	50.000[d]	-1.430[e]	0.154
Raw materials	8.000[b]	14.500[c]	14.000[b]	4.000[c]	wt	49.000[d]	-1.320[e]	0.186
Mean nut availability	1.296	0.489	1.410	0.371	tt	0.547	14.835	0.593
Wild fruit trees	3.000	3.300	3.571	2.760	tt	0.387	14.401	0.704
Distance to nearest river (m)	139.613	77.486	124.710	84.836	tt	-0.369	12.265	0.718
Mean fruit bunches	1.252	0.630	1.188	0.630	tt	-0.204	13.069	0.842
Wild food trees	4.000	3.916	4.286	2.690	tt	0.178	15.000	0.861

[a] Test used: tt =t-test; wt = Wilcoxon rank sum test
[d] Reported as median
[e] Reported as inter-quartile range (IQR)
[b] Reported as w-statistic for Wilcoxon rank sum tests
[c] Reported as z-score for the w-statistic

1000

1001

1002

1003   Table A6 – Likelihood ratio test results of the Full-Null model comparisons of the site inactivity

1004   models.

Model	Resid. df	Resid. Dev	df	Deviance	p
~1 (Null model)	81	113.627			
~Mean nut availability + Raw materials + Wild food trees	36	46.532	3	8.017	0.046
~Mean nut availability + **Raw materials (size class 3-5)** + Wild food trees	36	46.518	3	8.030	0.045
~Mean nut availability + **Tools** + Wild food trees	36	41.350	3	13.198	0.004
~Mean nut availability + **Raw materials (size class 3-5)** + **Wild fruit trees**	36	46.621	3	7.928	0.048
~Mean nut availability + **Tools** + **Wild fruit trees**	36	41.968	3	12.581	0.006

1005

1006   Table A7 – Results of AICc-Based Model Selection of the GLM for the predictors of tool site

1007   inactivity. Subset variables highlighted in bold.

Model	df	AICc
~Mean nut availability + Raw materials + Wild food trees	4	55.675
~Mean nut availability + **Raw materials (size class 3-5)** + Wild food trees	4	55.661
~Mean nut availability + **Tools** + Wild food trees	4	50.493
~Mean nut availability + Raw materials (size class 3-5) + **Wild fruit trees**	4	55.764
~Mean nut availability + **Tools** + **Wild fruit trees**	4	51.110

1013 Table A8 – Results of the additional GLM models investigating potential predictors of tool site

1014 inactivity. Subset variables highlighted in bold.

1015

Model	Term	Estimate	SE	χ^2	P [a]
~Mean nut availability + Raw materials + Wild food trees	Intercept	1.275	0.927		
	Mean nut availability	-1.909	0.788	7.757	0.005
	Raw materials (size class 3-5)	0.001	0.017	0.001	0.973
	Wild food trees	0.083	0.111	0.567	0.452
~Mean nut availability + **Raw materials (size class 3-5)** + Wild food trees	Intercept	1.310	0.923		NI[b]
	Mean nut availability	-1.911	0.787	7.796	0.005
	Raw materials (size class 3-5)	-0.003	0.022	0.015	0.903
	Wild food trees	0.086	0.109	0.625	0.429
~Mean nut availability + Raw materials (size class 3-5) + **Wild fruit trees**	Intercept	1.838	0.971		NI[b]
	Mean nut availability	-1.895	0.773	7.757	0.005
	Raw materials (size class 3-5)	0.004	0.022	0.036	0.849
	Wild fruit trees	-0.104	0.145	0.522	0.470
~Mean nut availability + **Tools** + **Wild fruit trees**	Intercept	3.282	1.268		NI[b]
	Mean nut availability	-2.077	0.812	8.491	0.004
	Tools	-0.170	0.088	4.689	0.030
	Wild fruit trees	-0.053	0.144	0.135	0.714

[a] Results from the likelihood ratio test using the 'drop1' function.
[b] Not indicated because it has a limited interpretation.

1016

1017

1018