Detection and phylogeny of *Wolbachia* in field-collected *Aedes albopictus* and *Aedes aegypti*

from Manila City, Philippines

Maria Angenica F. Regilme[1,2], Tatsuya Inukai[1], and Kozo Watanabe[1,2]

[1] Center for Marine Environmental Studies (CMES), Ehime University, Matsuyama, Ehime,

Japan

[2] Ehime University - De La Salle University International Collaborative Research Laboratory,

Laguna Province, Philippines

Corresponding author

E-mail: watanabe.kozo.mj@ehime-u.ac.jp

Abstract

Wolbachia is the most common bacterial endosymbiont of arthropods, such as the

medically important *Aedes albopictus* and *Aedes aegypti*. Both *Wolbachia* and mitochondrial

DNA are maternally inherited. This study collected 12 adult *Ae. albopictus* and 359 *Ae. aegypti*

from 183 households in a dengue-prone area, Manila, Philippines, between June and

September 2017. *Aedes* larvae (n = 509) were also collected from 17 water containers from 11

households. The DNA of the *Aedes* larvae and adults were screened for the presence of

Wolbachia using the *wsp* and 16S markers, following optimized polymerase chain reaction

(PCR) conditions, and sequenced. We also performed PCR using the mitochondrial DNA *cox1*

marker and sequenced the *Wolbachia*-positive samples. Our results showed that 3 out of 359

(0.84%) adult *Ae. aegypti* and 12 out of 12 (100%) adult *Ae. albopictus* were *Wolbachia*

positive, whereas all larvae tested negative for *Wolbachia* (0/509; 0%). The *wsp* marker

revealed six *Wolbachia*-positive *Ae. albopictus*, whereas the 16S marker showed *Wolbachia* in

three *Ae. aegypti* and ten *Ae. albopictus*. Three *Wolbachia*-positive *Ae. albopictus* from one household were found to have a single haplotype for all *Wolbachia* (*wsp* and 16S) and mitochondrial *cox1* markers. This result can be used to infer that maternal transmission of *Wolbachia* in the household because siblings of a *Wolbachia*-infected mother mosquito share the *Wolbachia* and mitochondrial genomes. Our results suggest that the combined sequencing analyses of *Wolbachia* (*wsp* and 16S) and mitochondrial DNA can infer maternal transmission in field-collected mosquitoes.

Keywords

Wolbachia, dengue, *Aedes sp.,* haplotype, mitochondrial DNA

Introduction

Wolbachia is a maternally inherited endosymbiotic bacteria, infecting 40% of arthropod species and some filarial nematodes (Werren et al., 2008; Zug et al., 2012). Maternal transmission permits the rapid spread of *Wolbachia* in mosquito populations (Fraser et al., 2017). It is naturally present in medically important mosquito species, including *Aedes albopictus* (Albuquerque et al., 2011; Joanne et al., 2015; Kitrayapong et al., 2002) and *Aedes aegypti* (Balaji et al., 20119;Carvajal et al., 2019; Kulkarni et al., 2019; Teo et al., 2017) inhibiting the replication of arboviral pathogens (Iturbe-Ormaetxe et al., 2011). Natural populations of *Ae. aegypti* were considered to be negative for *Wolbachia*, but recent studies have reported both positive (Coon et al., 2016; Teo et al., 2017;Thongsripong et al., 2018; Hegde et al., 2018; Balaji et al., 2019; Kulkarni et al., 2019; Carvajal et al., 2019 and Bennett et al., 2019) and negative (Gloria-Soria et al., 2018 and Ross et al., 2019) results for *Wolbachia* in field-collected *Ae. aegypti* among different countries. Due to these varying results, further validation of natural infection of *Wolbachia* in field-collected *Ae. aegypti* is needed.

In the field, inference of maternal transmission is usually made by collecting adult

mosquitoes (Hoffman et al., 1998) or placing ovitraps (Schmidt et al., 2018) for *Wolbachia*

detection using polymerase chain reaction (PCR). In a study by Schimdt et al., 2018, ovitraps

were placed in field-released areas in Australia and three mosquito eggs per ovitrap were then

collected. Using the fluorescent real-time PCR/high resolution melt assay, 60 of 161 (37.27%)

Ae. aegypti were found to be *Wolbachia*-positive, indicating that full-siblings from most

ovitraps were able to sustain *Wolbachia* transmission in the field. The authors inferred

maternal transmission of *Wolbachia* because three eggs were collected from the same ovitrap

and, therefore, are assumed to be from the same mother. Notably, this method can only be

applied to ovitrap-based samples. In addition, they tested maternal transmission in Wolbachia-

released areas. Thus, in the present study, we used adult mosquitoes collected from the field

to infer maternal transmission of *Wolbachia*. If we need to see maternal transmission in

naturally occurring strains, this approach cannot be applied.

Since *Wolbachia* and mitochondrial genomes are maternally inherited, siblings of a

Wolbachia-infected mother mosquito will share these genomes, showing close association

leading to linkage disequilibrium among *Wolbachia*-infected mosquitoes. Maternal

transmission can be inferred through the analysis of both the *Wolbachia* and mitochondrial

genomes, indicating whether two or more *Wolbachia*-infected mosquitoes living in the same

or close localities (e.g., households) carried *Wolbachia* from the same mother. Yeap et al.

(2016) reported different mitochondrial DNA haplotypes between *Wolbachia*-transfected

mosquitoes from laboratory colonies and field-caught *Wolbachia*-negative mosquitoes. Based

on this result, they confirmed that the maternal transmission of *Wolbachia* occurs only in

mosquitoes with the same mitochondrial lineage as the transfected mosquitoes used for field

release. However, no study to date has assessed mosquito mitochondrial DNA haplotypes

along with *Wolbachia* sequences nor have any studies inferred maternal transmission of

naturally occurring *Wolbachia* strains from field-collected mosquitoes in areas without prior

release of *Wolbachia*-transfected mosquitoes.

The presence of natural *Wolbachia* strains in field-collected *Ae. aegypti* is important as

it may affect future *Wolbachia* mass release programs, thereby influencing the successful

invasion of *Wolbachia* strains from transfected mosquitoes into the natural population. Most

studies have found that naturally occurring *Wolbachia* in *Ae. aegypti* are strains that are

phylogenetically close to the wAlbA strain, which also infects *Ae. albopictus* (Balaji et al.,

2019;Carvajal et al., 2019; Kulkarni et al., 2019; Coon et al., 2016). *Ae. albopictus* is known to

be infected with two *Wolbachia* strains, known as wAlbA and wAlbB, which belong to the

supergroups A and B, respectively (Zhou et al., 1998). However, previous studies detecting

Wolbachia in field-collected *Ae. aegypti* did not test *Ae. albopictus*. To our knowledge, there

have been no previous studies that found *Wolbachia* supergroups A and B from *Ae. aegypti*

and *Ae. albopictus* samples collected from the same localities at the same time.

In this study, we detected *Wolbachia* in field-caught *Ae. albopictus* and *Ae. aegypti*

using *Wolbachia*-specific DNA markers (*wsp* and 16S). Combining the *Wolbachia* sequence

information with the mitochondrial DNA (*cox1*) sequence information, we inferred the

maternal transmission of *Wolbachia* from a common female ancestor. We tested the

hypothesis that *Wolbachia*-positive *Ae. albopictus* and/or *Ae. aegypti* mosquitoes collected

from same household may have the same haplotype in the mitochondrial DNA (*cox1*) and

Wolbachia markers (*wsp* and 16S), using adult samples collected from Manila City, Philippines.

We also identified and compared the *Wolbachia* strains and supergroups found in *Ae.*

albopictus and *Ae. aegypti*.

Methods

Study site, adult mosquitoes, larvae collection, and identification

We selected the study site in Manila City, the capital city of the Philippines. It is a

highly populated and urbanized area that connects two major cities, Manila City and Quezon

City. It consists of residential, commercial, and industrial infrastructure. We collected adults

and larvae between June and September 2017, the rainy season, and the peak time for dengue

cases.

Adults of 12 *Ae. albopictus* and 359 *Ae. Aegypti* were collected using commercially

available mosquito light traps (Jocanima Corporation, Manila, Philippines) from 183 of 428

households that were randomly selected and surveyed. We set the mosquito trap for 48 hours

inside or outside the surveyed households. The mosquito trap attracts the host-seeking and

blood-fed mosquitoes to enter a capture net via the heat and CO_2 transmitted by a strong

current from the ventilator (Balingit et al., 2020; Regilme et al., 2021). We also surveyed water

containers in each household (n = 428) and found 17 containers with *Aedes* larvae (n = 509)

outside the premises of 14 households. We classified the breeding containers into four general

types following the World Health Organization guidelines (WHO, 2011) —water storage

containers, such as drums, pails, and buckets, (n = 8); plant pots (n = 2); water fountain (n = 2);

and others (namely, a discarded toilet bowl and a street canal; n = 2). The number of larval

samples collected per water container ranged from 2 to 287 (mean = 31). The adults (n = 371)

and larvae (n = 509) collected were morphologically identified as *Aedes sp.* using a

stereomicroscope and the keys published by Rueda et al. (2004). The adult and larval samples

were preserved in RNAlater (Ambion, Invitrogen, CA) solution to keep their RNA and DNA

intact and stored at −20⬚ before nucleic acid extraction.

Using microsatellite data from a recent study (Regilme et al., 2021) , we were also able

to confirm the species of 359 adult mosquitoes as *Ae. aegypti.* We had already used the adult

DNA of 359 *Ae. aegypti* for population genetics study (Regilme et al., 2021) and used the

remaining DNA for this study for *Wolbachia* detection. The microsatellite primers were

species-specific for *Ae. aegypti* (Chambers et al., 2007; Slotman et al., 2007). The remaining 12

adult samples that did not show PCR amplification with the microsatellite primers were

confirmed to be *Ae. albopictus*.

DNA extraction, molecular identification, PCR amplification, and sequencing

We extracted DNA using Qiagen AllPrep DNA/RNA micro kit and Qiagen DNA Blood

and Tissue DNEasy Kits© (Qiagen, Hilden, Germany) in adult (*n* = 371) and larval (*n* = 509)

samples, respectively. DNA concentration and quality were measured using a NanoDrop 2000

Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA).

Wolbachia was detected using two molecular markers: *wsp* (610 base pairs) using the

primer pairs *wsp* 81F (5′-TGGTCCAATAAGTGATGAAGAAAC-3′) and *wsp* 691R (5′-

AAAAATTAAACGCTACTCCA-3′) (Zhou et al., 1998) and 16S specific for *Wolbachia* (850 base

pairs) with the primer pairs *WolbF* (5′-GAAGATAATGACGGTACTCAC-3′) and *Wspecr* (5′-

AGCTTC GAGTGAAACCAATTC-3′) (Simoes et al., 2011).

For both *wsp* and 16S gene amplification, we followed the protocol published in

(Carvajal et al., 2019), in a final volume of 10 µl with 1 µl of the genomic DNA. We used the

following components for the PCR reaction for both markers: 10x Ex Taq buffer, 25 mM MgCl$_2$,

2.5 mM dNTP, 10 µM forward and reverse primers, 10% dimethyl sulfoxide (Sigma-Aldrich, St.

Louis, MO, USA), and 5 units/µl of Takara Ex Taq™ (Takara Bio Inc., Shiga, Japan). The *wsp* PCR

amplification was as follows: initial denaturation at 95◻ for 3 minutes, denaturation at 95◻ for

1 minute, annealing at 55◻ for 1 minute, an extension at 72◻ for 1 minute for 40 cycles, and a

final extension at 72◻ for 3 minutes. The 16S amplification followed these conditions: initial

denaturation at 95℃ for 2 minutes, denaturation at 95℃ for 2 minutes, annealing at 60℃ for 1

minute, an extension at 72℃ for 1 minute for two cycles, another 35 cycles of denaturation at

95℃ for 30 seconds, annealing at 60℃ for 1 minute, an extension at 72℃ for 45 seconds, and a

final extension at 72℃ for 10 minutes. We included a positive control of a *Wolbachia*-positive

Culex sp. and negative control of water in each PCR run.

PCR products were analyzed in 1.5% agarose gel stained with Midori Green Advance

DNA stain at 100 V for 30 minutes. To validate the presence of *Wolbachia* in each sample, we

performed PCR amplification twice per marker. The criteria for a positive *Wolbachia* test were

based on two successful amplifications per molecular marker, *wsp,* and 16S.

We also amplified the *cox1* mitochondrial gene of *Wolbachia*-positive samples using

the primer pairs LCO-1490 (5′-GGTCAACAAATCATAAAGATATTGG-3′) and HCO1-2198 (5′-

AAACTTCAGGGTGACCAAAAAATCA-3′) (Folmer et al., 1994). We used the following PCR

amplification profile: initial denaturation at 95℃ for 5 minutes, denaturation at 95℃ for 30

seconds, annealing at 48℃ for 45 seconds for 35 cycles, and a final extension at 72℃ for 7

minutes. The amplified PCR products were purified using QIAquick (Qiagen, Hilden, Germany)

PCR Purification kits and sequenced by Eurofin Genomics Inc. Tokyo, Japan.

Identification of *Wolbachia* strains, haplotypes, and phylogenetic analysis

We assembled the forward and reverse sequences for each marker using the

CodonCode Aligner version 1.2.4 software (https://www.codoncode.com/aligner/). We aligned

the sequences using the online program MAFFT version 7 with the default settings

(https://mafft.cbrc.jp/alignment/software/). We checked all generated sequences for

similarities with reference sequences from GenBank (NCBI, 2016) using Basic Local Alignment

Search   Tool–Nucleotide   BLAST

(https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch). Finally, we checked the

sequence quality of the aligned sequences in Mesquite version 3.5 (Maddison et al., 2019) by

confirming the absence of stop codons.

All *wsp,* 16S, and *cox1* sequences were separately analyzed using DNAsp version

6.12.03 (Rozas et al., 2017) to determine the number of haplotypes. We assessed the

relationship of the *Wolbachia* strains of our study with representative sequences from

different insect hosts by constructing a phylogenetic tree for the *wsp* and 16S sequences using

PhyML 3.1 (Guindon et al., 2003) using the default settings. We applied the GTR + G model for

wsp and the GTR + G + 1 model for 16S. The model per marker was selected using the SMART

model selection method (Lefort et al., 2017). We used *Brugia pahangi* (AY527207) and

Rickettsia sp. (U11021) as the outgroups for the *wsp* and 16S, respectively.

Results

Detection of *Wolbachia* infection in adult and larval mosquitoes

We screened all individual *Aedes sp.* larvae (n = 509) and adults (n = 371) for

Wolbachia using the *wsp* and 16S markers. Larval samples showed no evidence of *Wolbachia*

from either marker. However, *Wolbachia* was detected in 15 (4.04%) out of 371 *Aedes sp.*

adult mosquitoes based on the *wsp* and 16S markers. A total of three (0.84%) were *Wolbachia*

positive out of 359 *Ae. aegypti*, based on the *wsp* marker and 12 (100%) out of 12 *Ae.*

albopictus (Figure 1), based on the *wsp* and 16S markers. The *wsp* marker detected *Wolbachia*

in six *Ae. albopictus,* whereas the 16S marker detected it in three *Ae. aegypti* and ten *Ae.*

albopictus. Among these *Wolbachia*-infected *Ae. albopictus,* 4 (26.67%) out of 15 were positive

for both *wsp* and 16S markers. Confirmation of the species identification using the *cox1* marker

revealed that the 12 adult mosquitoes were *Ae. albopictus*, whereas the remaining three were

Ae. aegypti based on BLAST results with 100% identity match (Table 1).

The phylogenetic tree built using the *wsp* marker (Figure 2 A) revealed two major

clades: supergroup A and supergroup B (Zhou et al., 1998). Two *Ae. albopictus* were infected

with *Wolbachia* belonging to supergroup A and four with *Wolbachia* belonging to supergroup

B. The haplotype sequences from supergroup A were grouped with the *Wolbachia* type strain

A (wAlb A), identified in an *Ae. albopictus* host in the USA (Zhou et al., 1998). In contrast, the

sequences from supergroup B clustered with the *Wolbachia* type strains from *Ae. aegypti*

(*wAegB*) in the Philippines (Balaji et al., 2019; Carvajal et al., 2019) and *Ae. albopictus* (*wAlbB*)

in the USA (Zhou et al., 1998).

In the phylogenetic tree constructed using 16S sequences, 11 out of 13 sequences

(84.62%) found in *Ae. albopictus* (n = 10) and *Ae. aegypti* (n = 3) were clustered within

supergroup A (Figure 2 B). The remaining two sequences (15.38%) found in *Ae. albopictus* were

clustered together with the supergroups C, D, E, F, and J from insect hosts such as *Folsomia*

candida, Mansonella azzardi, and *Dirofilaria immitis.*

Wolbachia and mitochondrial cox1 haplotypes

We identified six *cox1,* three *wsp,* and nine 16S haplotypes among the 15 *Wolbachia*-

positive sequences (Table 1). We found that three *Ae. albopictus* samples collected from one

household (NH149) were *Wolbachia* positive, based on the *wsp* and 16S markers (Table 1). An

identical *cox1* haplotype (sequence), C1, was observed among all three *Ae. albopictus*. The

same haplotypes of *wsp* (W2) and 16S (S6) also occurred among these three *Ae. albopictus*

(Table 2). Another household (NH56) had three *Wolbachia*-positive *Ae. albopictus* with the

same *cox1* haplotype (C1). Two *Ae. albopictus* (Individual codes 177 and 178) were *Wolbachia*

positive according to the *wsp* sequences, while two *Ae. albopictus* (Individual codes 95 and

177) were positive, according to the 16S sequence. We observed different haplotypes in each

wsp (W1 and W2) and 16S (S8 and S9) marker. Other than these two households, no other

households had more than one *Wolbachia*-positive mosquito.

Discussion

The presence of the same haplotype of *Wolbachia* (*wsp* and 16S) and mitochondrial

DNA (*cox1*) among the three *Wolbachia*-positive *Ae. albopictus* collected from household

NH149 may infer the maternal transmission of the *Wolbachia*.

We assumed that our approach eliminates the bias of laboratory-controlled conditions,

including temperature, light, and feeding times, during mosquito rearing. The maternal

transmission of *Wolbachia* in mosquitoes is usually confirmed by detecting *Wolbachia* in the

offspring of infected mothers (Balaji et al., 2019; Kulkarni et al., 2019). For example,

Wolbachia-microinjected *Ae. aegypti* embryos were reared into adults and then tested using

PCR for the presence of *Wolbachia* (McMeniman et al., 2009). Severe environmental

conditions such as high temperature causing heat stress to mosquitoes may decrease

Wolbachia density, thus reducing the ability of *Wolbachia* to persist in the mosquito

population. The results of a previous study (Ross et al., 2019) suggest that high temperature

affects *Wolbachia*. For example, when breeding containers are exposed to more sunlight than

shade, the density of *wMel* declined. Previous sterile insect studies failed to successfully

deploy *Wolbachia* because of the poor performance of the mosquitoes released in the field,

possibly because of the extreme environmental conditions differing from those of the

laboratory (Benedict et al., 2003; Helinski et al., 2013). Thus, it is important to verify *Wolbachia*

transmission in the field. In a previous study (da Silva et al., 2019), the establishment of

Wolbachia in Australia during and after mass release was determined by collecting mosquitoes

in the field and then detecting *Wolbachia* using loop-mediated isothermal amplification. The

researchers successfully confirmed maternal transmission because the field-collected

mosquitoes had wMel strains, which were considered absent from the natural populations

before mass release. However, the inference of maternal transmission of naturally occurring

Wolbachia strains in mosquito populations is not as simple as the case after mass release. It is

difficult to determine whether the strains detected in field-collected mosquitoes resulted from

transmission from the mother or the naturally occurring *Wolbachia* populations in the area.

Our approach, the combined analysis of *Wolbachia* and mitochondrial DNA sequences, can be

used to infer maternal transmission in field-collected mosquitoes.

One of the limitations of this study is that we observed an unexpected pattern found

in one household, NH56, that did not show the same haplotype of *Wolbachia wsp* and 16S

sequences in three *Wolbachia*-infected individuals. However, the same mitochondrial DNA

haplotype appeared in the three individuals. One of the possible reasons is that the resolution

of the *cox1* marker might not be fine enough to tell whether the mosquitoes with the same

cox1 haplotype are from the same mother. Based on the number of *cox1* haplotypes [i.e., 4

(C1–C4)] found in the 12 *Wolbachia*-positive *Ae. albopictus*, we calculated a Poisson probability

of 0.149 for obtaining the same *cox1* haplotype among 3 randomly selected individuals. Our

study detected 6 *cox1* haplotypes from the 15 *Wolbachia*-positive individuals, and some

mosquitoes from the same or different households shared the same *cox1* haplotype, which

might imply that although they are not from the same household, they may have the same

maternal lineage. The mitochondrial genome reflects the long-term maternal lineage (Folmer

et al., 1994). Therefore, although the mothers of these three individuals were different, they

are considered to have a common ancestor in their maternal lineage, although how many

generations ago cannot be estimated. In future studies, we suggest increasing the sample size

to obtain more *Wolbachia*-positive mosquitoes from one household for a stochastically strong

validation of the maternal transmission through whole genome sequencing or double-digest

restriction site-associated DNA. The use of additional mitochondrial markers or whole genome

sequencing (e.g., double-digest restriction site-associated DNA) is also recommended for more

accurate verification.

262   Our results were negative for *Wolbachia* in all the collected *Aedes* larvae (n = 509).

One possible reason is that we found only 17 water containers during our field sampling. Since

Wolbachia is maternally transmitted, all of its offsprings would also be negative for *Wolbachia*

if the mother is negative for *Wolbachia*. More *Wolbachia* might be detectable in larval samples

if we increase the number of water containers than individuals per container. Although 428

households were surveyed in this study, only 17 water storage containers were found. Thus,

we suggest increasing the number of sampled water containers in future studies by setting up

ovitraps in households. We also recommend detecting *Wolbachia* in representative larval

samples from one water container.

271   Another possible reason for obtaining negative results for *Wolbachia* in larvae can be

linked to a lower *Wolbachia* density during the larval stage of mosquitoes. Previous studies

(Coon et al., 2016; Stevanovic et al., 2015) have found that the *Wolbachia* density is much

lower in the larval stage than in the adult stage. Wiwatanaratanabutr et al. (2009) discovered

that temperature and larval crowding may affect the relative *Wolbachia* densities in *Ae.*

aelbopictus. Stevanovic et al., (2015) found a lower *Wolbachia* density in *Drosophila*

melanogaster larvae than in adult females. We presume that this is due to the immature

organs of larva, such as the thoracic ganglia and muscles, ovaries, head, and salivary glands. In

the present study, we were unable to quantify the *Wolbachia* density in *Wolbachia*-positive

individuals. Thus, we recommend performing real-time quantitative PCR analysis alongside

conventional PCR in future studies.

We also found that the *Ae. albopictus* were infected with both supergroups A and B of

Wolbachia using the *Wolbachia* surface protein gene (*wsp*), widely used for *Wolbachia* strain

identification and phylogeny studies (Zhou et al., 1998). A previous study (Carvajal et al., 2019)

found that field-collected *Ae. aegypti* positive for *Wolbachia* from Manila, Philippines were

clustered into two supergroups: A and B. We found more *Wolbachia*-positive *Ae. albopictus* in

the *Wolbachia* supergroup B (4/6; 66.66%) than in supergroup A (2/6; 33.33%), as observed in

a previous study (Hu et al., 2020) (China; 631/693; 91.05%). Thus, our research results support

previous studies, and we observed two *Wolbachia* supergroups in *Ae. albopictus* collected

from the same area at the same time. However, the proportions of *Wolbachia* supergroups

might change in field conditions due to the influence of temperature and other abiotic factors.

We found a low prevalence of *Wolbachia* in adult *Ae. aegypti*, which is concordant

with the previous results of Kulkarni et al. (2016) obtained from field-collected *Ae. aegypti* in

Florida, USA and those of Carvajal et al.'s study (2019) conducted in Metro Manila, Philippines.

In contrast, a high (>50%) prevalence rate was observed in the studies of Teo et al. (2017)

conducted in Malaysia and Kulkarni et al. (2016) conducted in New Mexico, USA. The low

Wolbachia prevalence rate in *Ae. aegypti* may be due to the low density of the *Wolbachia*

endosymbiont in *Ae. aegypti* (Carvajal et al., 2019); this is supported by the previous

metabarcoding studies of Kulkarni et al.(2019), Coon et al. (2016), and Thongsripong et al.

(2018), which all reported a low number of sequence reads in the *Ae. aegypti* midgut,

indicating a low density of *Wolbachia*. Although our results are limited since we did not

quantitatively measure *Wolbachia* density, our 40-cycle PCR amplification method adapted

from Carvajal et al. (2019) could infer that *Wolbachia*-positive *Ae. aegypti* might be present in

Metro Manila, Philippines. Further *Wolbachia* detection studies are thus needed to affirm

natural *Wolbachia* infection in field *Ae. aegypti.*

We acknowledge the uncertainties of *Wolbachia*-positive *Ae. aegypti* due to the 16S

marker and the conventional PCR method, e.g., the false-positive rate. In light of this

information, we were careful to ascertain positive *Wolbachia* in adult *Ae. aegypti.* Thus, we

used the primers of Simoes et al. (2011), which are known to produce fewer false-positive

results and negative detections of *Wolbachia.* We also performed repeated PCR tests of our

Wolbachia-positive samples to ensure successful *Wolbachia* detection as defined in the

previous study by Carvajal et al. (2020). In addition to conventional PCR, tests such as

fluorescent *in situ* hybridization and whole genome sequencing can be used to provide strong

support for the presence of *Wolbachia* in natural *Ae. aegypti* populations. Despite these

limitations, our results provide additional support for *Wolbachia* detection in field-collected *Ae.*

aegypti.

Conclusions

An interesting approach to infer the maternal transmission of *Wolbachia* in field-

collected mosquitoes can be through the analysis of *Wolbachia wsp* and 16S sequences

combined with mitochondrial sequences. Since *Wolbachia* and mitochondria are both

maternally inherited, a non-random association of the mitochondrial genome with the

Wolbachia endosymbiont can be observed. The strategy of combining mitochondrial and

Wolbachia sequence information can infer the maternal transmission of *Wolbachia* in field-

collected *Ae. albopictus* and *Ae. aegypti.* In this study, we used mosquitoes naturally infected

with *Wolbachia* collected from the same households. We suggest that they might be from the

same maternal lineage (e.g. mother, grandmother etc.) due to the presence of the same

mitochondrial haplotype sequences in mosquitoes collected from the same household. This

may also imply that the other set of mosquitoes in this study with the same mitochondrial DNA

haplotype but a different *Wolbachia* strain came from the same maternal lineage. Therefore,

the results of *Wolbachia wsp* and 16S sequences combined with mitochondrial DNA sequences

can support the use of genome sequencing to infer maternal transmission in field-collected

mosquitoes. Using field-collected mosquitoes allows us to infer maternal transmission in the

environment, a phenomenon that is possible to be overestimated under laboratory conditions.

Our findings may provide additional support for the maternal transmission of *Wolbachia* in

naturally infected mosquitoes in the field and its ability to invade and persist in the population.

List of Abbreviations

PCR: polymerase chain reaction

DNA: deoxyribonucleic acid

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated and/or analyzed during this study are included in this published article and

its additional files. All newly generated sequences are available in the GenBank database under

the Accession Numbers _____.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported in part by the Japan Society for the Promotion of Science (JSPS)

Grant-in-Aid Fund for the Promotion of Joint International Research [Fostering Joint

International Research (B)] under grant number 19KK0107, JSPS Grant-in-Aid for Scientific

Research (B) under grant number 21H02206, the JSPS Core-to-Core Program B Asia-Africa

Science Platforms under grant number JPJSCCB20190008, and the Endowed Chair Program of

the Sumitomo Electric Industries Group Corporate Social Responsibility Foundation.

Author's contributions

MAFR and KW conceptualized and designed the experiment. MAFR and TI collected and

identified the adult and larval samples for this study. MAFR and TI performed the *Wolbachia*

detection and all the molecular experiments for this study. MAFR, TI and KW performed the

data analysis. MAFR and KW wrote the manuscript. All authors read and approved the final

manuscript.

Acknowledgments

We are thankful to all the households that participated in our sampling collection of

mosquitoes and larvae and to the village officials of Sampaloc, Manila, for their assistance and

support. We appreciate the help of Johanna Beulah T. Sornillo of the Research Institute for

Tropical Medicine, Philippines, for her technical assistance in the household sampling design.

We want to thank Katherine Viacrusis for her service in the fieldwork and Mr. Micanaldo

Francisco for constructing the sampling map. Our deepest thanks to Dr. Thaddeus M. Carvajal,

Dr. Divina Amalin, Dr. Mary Jane Flores, Dr. Joeselle M. Serrana for their valuable suggestions

and technical assistance. The authors would like to thank Enago (www.enago.jp) for the

Reference

Albuquerque AL, Magalhães T, Ayres CF. High prevalence and lack of diversity of Wolbachia

pipientis in *Aedes albopictus* populations from Northeast Brazil. Memórias do Instituto

Oswaldo Cruz. 2011 Sep;106(6):773-6.

Balaji S, Jayachandran S, Prabagaran SR. Evidence for the natural occurrence of Wolbachia in

Aedes aegypti mosquitoes. FEMS microbiology letters. 2019 Mar;366(6):fnz055.

Balingit JC, Carvajal TM, Saito-Obata M, Gamboa M, Nicolasora AD, Sy AK, Oshitani H,

Watanabe K. Surveillance of dengue virus in individual Aedes aegypti mosquitoes collected

concurrently with suspected human cases in Tarlac City, Philippines. Parasites & vectors. 2020

Dec;13(1):1-3.

Benedict MQ, Robinson AS. The first releases of transgenic mosquitoes: an argument for the

sterile insect technique. Trends in parasitology. 2003 Aug 1;19(8):349-55.

Bennett KL, Gómez-Martínez C, Chin Y, Saltonstall K, McMillan WO, Rovira JR, Loaiza JR.

Dynamics and diversity of bacteria associated with the disease vectors *Aedes aegypti* and

Aedes albopictus. Scientific reports. 2019 Aug 21;9(1):1-2.

Carvajal TM, Hashimoto K, Harnandika RK, Amalin DM, Watanabe K. Detection of Wolbachia in

field-collected *Aedes aegypti* mosquitoes in metropolitan Manila, Philippines. Parasites &

vectors. 2019 Dec;12(1):361.

Chambers EW, Meece JK, McGowan JA, Lovin DD, Hemme RR, Chadee DD, McAbee K, Brown

SE, Knudson DL, Severson DW. Microsatellite isolation and linkage group identification in the

yellow fever mosquito *Aedes aegypti*. Journal of Heredity. 2007 May 1;98(3):202-10.

Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are essential

for development but vary greatly between local habitats. Molecular ecology. 2016

Nov;25(22):5806-26.

da Silva Gonçalves D, Hooker DJ, Dong Y, Baran N, Kyrylos P, Iturbe-Ormaetxe I, Simmons CP,

O'Neill SL. Detecting wMel Wolbachia in field-collected *Aedes aegypti* mosquitoes using loop-

mediated isothermal amplification (LAMP). Parasites & vectors. 2019 Dec;12(1):1-5.

Fraser JE, De Bruyne JT, Iturbe-Ormaetxe I, Stepnell J, Burns RL, Flores HA, O'Neill SL. Novel

Wolbachia-transinfected *Aedes aegypti* mosquitoes possess diverse fitness and vector

competence phenotypes. PLoS pathogens. 2017 Dec 7;13(12):e1006751.

Folmer O. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from

diverse metazoan invertebrates. Mol Mar Biol Biotech. 1994;3:294-9.

Gloria-Soria A, Chiodo TG, Powell JR. Lack of evidence for natural Wolbachia infections in

Aedes aegypti (Diptera: Culicidae). Journal of medical entomology. 2018 Aug 29;55(5):1354-6.

Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by

maximum likelihood. Systematic biology. 2003 Oct 1;52(5):696-704.

Hegde S, Khanipov K, Albayrak L, Golovko G, Pimenova M, Saldaña MA, Rojas MM, Hornett EA,

Motl GC, Fredregill CL, Dennett JA. Microbiome interaction networks and community structure

from laboratory-reared and field-collected *Aedes aegypti*, *Aedes albopictus*, and *Culex*

quinquefasciatus mosquito vectors. Frontiers in microbiology. 2018 Sep 10;9:2160.

Helinski ME, Harrington LC. Considerations for male fitness in successful genetic vector control

programs. InEcology of parasite-vector interactions 2013 (pp. 221-244). Wageningen Academic

Publishers, Wageningen.

Hoffmann AA, Hercus M, Dagher H. Population dynamics of the Wolbachia infection causing

cytoplasmic incompatibility in Drosophila melanogaster. Genetics. 1998 Jan 1;148(1):221-31.

Hu Y, Xi Z, Liu X, Wang J, Guo Y, Ren D, Wu H, Wang X, Chen B, Liu Q. Identification and

molecular characterization of Wolbachia strains in natural populations of Aedes albopictus in

China. Parasites & vectors. 2020 Dec;13(1):1-4.

Iturbe‐Ormaetxe I, Walker T, O'Neill SL. Wolbachia and the biological control of mosquito‐

borne disease. EMBO reports. 2011 Jun;12(6):508-18.

Joanne S, Vythilingam I, Yugavathy N, Leong CS, Wong ML, AbuBakar S. Distribution and

dynamics of Wolbachia infection in Malaysian *Aedes albopictus*. Acta tropica. 2015 Aug

1;148:38-45.

Kitrayapong P, Baimai V, O'Neill SL. Field prevalence of Wolbachia in the mosquito vector

Aedes albopictus. The American journal of tropical medicine and hygiene. 2002 Jan

1;66(1):108-11.

Kulkarni A, Yu W, Jiang J, Sanchez C, Karna AK, Martinez KJ, Hanley KA, Buenemann M, Hansen

IA, Xue RD, Ettestad P. Wolbachia pipientis occurs in *Aedes aegypti* populations in New Mexico

and Florida, USA. Ecology and evolution. 2019 May;9(10):6148-56.

Lefort V, Longueville JE, Gascuel O. SMS: smart model selection in PhyML. Molecular biology

and evolution. 2017 Sep 1;34(9):2422-4.

Maddison, W. P. and D.R. Maddison. 2019. Mesquite: a modular system for evolutionary

analysis. Version 3.5 http://www.mesquiteproject.org

McMeniman CJ, Lane RV, Cass BN, Fong AW, Sidhu M, Wang YF, O'Neill SL. Stable introduction

of a life-shortening Wolbachia infection into the mosquito Aedes aegypti. Science. 2009 Jan

2;323(5910):141-4.

NCBI Resource Coordinators. Database resources of the National Center for Biotechnology

Information. Nucleic Acids Res. 2016;44(D1):D7-D19. doi:10.1093/nar/gkv1290

Regilme MA, Carvajal TM, Honnen AC, Amalin DM, Watanabe K. The influence of roads on the

fine-scale population genetic structure of the dengue vector *Aedes aegypti* (Linnaeus). PLoS

neglected tropical diseases. 2021 Feb 26;15(2):e0009139.

Ross PA, Ritchie SA, Axford JK, Hoffmann AA. Loss of cytoplasmic incompatibility in Wolbachia-

infected *Aedes aegypti* under field conditions. PLoS neglected tropical diseases. 2019 Apr

19;13(4):e0007357.

Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE,

Sánchez-Gracia A. DnaSP 6: DNA sequence polymorphism analysis of large data sets. Molecular

biology and evolution. 2017 Dec 1;34(12):3299-302.

Rueda LM. Pictorial keys for the identification of mosquitoes (Diptera: Culicidae) associated

with dengue virus transmission. Walter Reed Army Inst Of Research Washington Dc

Department Of Entomology; 2004 Aug 3.

Schmidt TL, Filipović I, Hoffmann AA, Rašić G. Fine-scale landscape genomics helps explain the

slow spatial spread of Wolbachia through the Aedes aegypti population in Cairns, Australia.

Heredity. 2018 May;120(5):386-95.

Simoes PM, Mialdea G, Reiss D, Sagot MF, Charlat S. Wolbachia detection: an assessment of

standard PCR protocols. Molecular Ecology Resources. 2011 May;11(3):567-72.

Slotman MA, Kelly NB, Harrington LC, Kitthawee S, Jones JW, Scott TW, Caccone A, Powell JR.

Polymorphic microsatellite markers for studies of *Aedes aegypti* (Diptera: Culicidae), the vector

of dengue and yellow fever. Molecular Ecology Notes. 2007 Jan;7(1):168-71.

Stevanovic AL, Arnold PA, Johnson KN. Wolbachia-mediated antiviral protection in *Drosophila*

larvae and adults following oral infection. Applied and Environmental Microbiology. 2015 Dec

1;81(23):8215-23.

Teo CH, Lim P, Voon K, Mak JJ. Detection of dengue viruses and Wolbachia in *Aedes aegypti*

and *Aedes albopictus* larvae from four urban localities in Kuala Lumpur, Malaysia. Trop Biomed.

2017 Sep 1;34:583-97.

Thongsripong P, Chandler JA, Green AB, Kittayapong P, Wilcox BA, Kapan DD, Bennett SN.

Mosquito vector-associated microbiota: Metabarcoding bacteria and eukaryotic symbionts

across habitat types in Thailand endemic for dengue and other arthropod-borne diseases.

Ecology and evolution. 2018 Jan;8(2):1352-68.

Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology. Nature

Reviews Microbiology. 2008 Oct;6(10):741-51.

Wiwatanaratanabutr I. Geographic distribution of wolbachial infections in mosquitoes from

Thailand. Journal of invertebrate pathology. 2013 Nov 1;114(3):337-40.

World Health Organization. Operational guide for assessing the productivity of *Aedes aegypti*

breeding sites. Geneva: TDR. 2011.

Yeap HL, Rašić G, Endersby-Harshman NM, Lee SF, Arguni E, Le Nguyen H, Hoffmann AA.

Mitochondrial DNA variants help monitor the dynamics of Wolbachia invasion into host

populations. Heredity. 2016 Mar;116(3):265-76.

Zhou W, Rousset F, O'Neill S. Phylogeny and PCR–based classification of Wolbachia strains

using *wsp* gene sequences. Proceedings of the Royal Society of London. Series B: Biological

Sciences. 1998 Mar 22;265(1395):509-15.

Zug R, Hammerstein P. Still a host of hosts for Wolbachia: analysis of recent data suggests that

40% of terrestrial arthropod species are infected. PloS one. 2012 Jun 7;7(6):e38544.

502 **Table 1.** Summary of sampling data and *Wolbachia* detection results

	Total
Surveyed households	428
Households with adult Aedes mosquitoes	183
Households with larval Aedes	11
Collected Aedes mosquitoes	371
Aedes albopictus	12
Aedes aegypti	359
Collected Aedes larvae	509
Wolbachia positive	
Larva	0
Adult Ae. albopictus	12
Adult Ae. aegypti	3
Households with Wolbachia-positive mosquitoes	11

 Table 2. A detailed summary of the results of the Wolbachia infection in selected Aedes

 mosquitoes using wsp and 16S markers

	Household	Molecular species identification	Individual code	Haplotypes			
				cox1	wsp		16S (n = 13)*
					Supergroup	wsp (n = 6)*	
1	SH9	*Ae. albopictus*	40	C1			S7
2	SH27	*Ae. albopictus*	306	C5			S3
3	SH134	*Ae. aegypti*	384	C6			S1
4	SH214	*Ae. albopictus*	425	C1			S6
5	NH43	*Ae. albopictus*	172	C4			S3
6	NH56	*Ae. albopictus* (n=3)	095, 177, 178	C1	A (177) and B (178)	W1 (177) and W2 (178)	S8 (095) and S9 (177)
7	NH107	*Ae. albopictus*	218	C5			S4
8	NH130	*Ae. aegypti*	116	C2			S1
9	NH131	*Ae. albopictus*	240	C5	A	W3	S5

10	NH149	*Ae. albopictus* (n = 3)	258, 259, 260	C1	B (258, 259, 260)	W2 (258, 259, 260)	S6 (258, 259)
11	NH181	*Ae. aegypti*	121	C3			S2
	Total number of haplotypes			**6**		**3**	**9**

523   * total number of *Wolbachia*-positive mosquitoes

Figure 1. Spatial distribution of *Wolbachia*-positive households (n = 11) out of 428 households

surveyed across a 2-kilometer road in highly urbanized area in Manila, Philippines.

537 **Figure 2.** The maximum likelihood phylogenetic tree of *Wolbachia*-infected urban mosquitoes

538 based on the **(A)** *WSP* sequences with reference sequences from supergroups A, B, and C and

539 an outgroup of *Brugia pahangi* **(B)** 16S sequences with reference sequences from supergroups

540 A, B, C, D, E, F, and J and an outgroup of *Rickettsia* sp.