Microbial succession on degrading micropollutants under recurrent exposure

Running title: Ecological feedback between microbe-micropollutants

Dandan Izabel-Shen[1#], Shuang Li[2,3,4#], Tingwei Luo[5], Jianjun Wang[6],

Yan Li[2,3], Qian Sun[2,3], Chang-Ping Yu[2,7], Anyi Hu[2,3*]

1. Department of Ecology, Environment and Plant Sciences, Stockholm

University, 106 91, Stockholm, Sweden

2. CAS Key Laboratory of Urban Pollutant Conversion, Institute of Urban

Environment, Chinese Academy of Sciences, Xiamen 361021, China

3. University of Chinese Academy of Sciences, Beijing 100049, China

4. Department of Environmental Microbiology, UFZ, Helmholtz Centre for

Environmental Research, Leipzig, Germany

5. Institute of Marine Microbes and Ecospheres, State Key Laboratory of Marine

Environmental Science, Xiamen University, Xiamen, China

6. State Key Laboratory of Lake Science and Environment, Nanjing Institute of

Geography and Limnology, Chinese Academy of Sciences, Nanjing, 210008,

China

7. Graduate Institute of Environmental Engineering, National Taiwan University,

Taipei 106, Taiwan

#The authors contributed equally to this work.

*Correspondence to:

Dr. Anyi Hu, ayhu@iue.ac.cn

CAS Key Laboratory of Urban Pollutant Conversion, Institute of Urban Environment,

Chinese Academy of Sciences, Xiamen 361021, China, Telephone & Fax: 86-592-

6190582.

Abstract

The increasing-volume release of micropollutants into natural surface waters has raised

great concern due to their environmental accumulation. Persisting micropollutants can

impact multiple generations of organisms via recurrent exposure, but their microbially-

mediated degradation and the influence on community assembly remain understudied.

Here, freshwater microbes were treated with four common micropollutants, alone or in

combination, and then transferred to fresh medium containing the same level of

micropollutants every 5-day to mimic the recurrent exposure of microbes for 35 days.

The degradation of micropollutants and the associated communities were surveyed for

each transfer. Even though the degradation of micropollutants did not accelerate over

time, the degradation *per se* altered the microbial community composition. Community

assembly was dominated by stochastic processes during early exposure, via random

community changes and emergence of seedbanks, and deterministic processes later in

the exposure, via advanced community succession. Early exposure stages were

characterized by the presence of sensitive microorganisms such as Actinobacteria and

Planctomycetes, which were then replaced by more tolerant bacteria such as

Bacteroidetes and Gammaproteobacteria. This succession was tightly linked to a high

degradation of bisphenol A and a lower one for triclosan. We hypothesize that microbes

are not robust to micropollutant persistence but have an intrinsic capacity to recover

from that disturbance by acquiring carbon and energy from micropollutant degradation.

Our findings have important implication for ecological feedback between microbe-

micropollutants under anthropogenic climate change scenarios.

Keywords: Micropollutant transformation; press disturbance; community turnover;

stochastic assembly; deterministic assembly

Introduction

The rapid exploitation and widespread use of emerging organic chemicals have resulted

in environmental damage and human health issues during the last several decades

(Deblonde et al., 2011; Schwarzenbach et al., 2006). These organic chemicals include

pharmaceuticals and personal care products, hormones, and industrial chemicals, and

are typically classified as micropollutants, because of their occurrence at a low to very

low concentration of μg and ng per litter in the receiving waters (Luo et al., 2014; Z.

Wang et al., 2015). For instance, the majority of micropollutants originate from direct

and indirect inputs of organic contaminants that are released via the effluents of

wastewater treatments plants (da Silva et al., 2020; Eggen et al., 2014; Schwarzenbach

et al., 2006; Vila-Costa et al., 2020). The micropollutants gradually accumulate at sites

of their release and reach concentrations high enough to exert disruptive effects on

aquatic organisms such as fish, invertebrates, and microbes (Kuroda et al., 2012; Liu et

al., 2015; Roberts et al., 2016; Rodea-Palomares et al., 2016; Yang et al., 2013).

Nonetheless, little is known about the mechanisms by which micropollutants impact

the turnover of aquatic microbial communities (Cerro-Gálvez et al., 2021; Martinez-

Varela et al., 2021; Vila-Costa et al., 2020), especially those of freshwater bacteria

(Adyari et al., 2020; Tlili et al., 2020), and the role of microbes on trace micropollutant

degradation.

How microbial assemblages respond to micropollutants can be classified as:

subsidy responses and stress responses. The subsidy responses is mainly via

biodegradation, in which the microbes are able to utilize the micropollutant compounds

as carbon sources, while the stress responses is relevant to the growth and survival of

microorganisms inhibited (Chalew & Halden, 2009; Zhang et al., 2013). For instance,

bisphenol A (BPA) and its substituted derivatives (e.g., bisphenol S, BPS) (Staples et

al., 1998) are typical subsidy responses, as their relatively simple chemical structure

and low toxicity allow both to be utilized as carbon sources by microbial isolates (Choi

& Lee, 2017; Zhang et al., 2013). Stress responses are elicited by pharmaceutical and

personal care micropollutants such as triclosan (TCS) and its substituted derivatives

(e.g., triclocarban, TCC) (McMurry et al., 1998), because both inhibit the growth of

several microbial taxa, consistent with their resemblance to antimicrobials (Cabana et

al., 2007; Chalew & Halden, 2009). Both BPA and TCS in the field strongly influence

microbial diversity, especially that of the central species and module communities

comprising the microbial co-occurrence network (Hu et al., 2017).

Microbial responses to chemical pollutants can be modified by the exposure legacy

of the community. For example, microbial communities with a history of containment

exposure may differ from contaminant-naïve communities in the rate and extent of

chemical degradation (Boyd et al., 2008; Coll et al., 2020; Okere et al., 2017). They

may also differ in their plasticity (Cerro-Gálvez et al., 2021) upon subsequent exposure

to similar contaminants. However, the ecological drivers of community succession

following repeated micropollutant exposure are not well-understood. Studies of

community dynamics over multiple generations and a time period relevant for

community stability could provide insights into the ecological traits that determine

taxon sensitivity and tolerance, and therefore the ability of microbial community

members to degrade micropollutants.

Press disturbances in natural environments can impact multiple generations of

organisms (Edward A. Bender, 1984) by altering the affected ecosystems beyond the

possibility of recovery or by causing a shift to a community with a new stable state

(Shade et al., 2012). Following a stress event such as recurrent micropollutant exposure,

communities rely on reassembly from the local seedbanks such that succession

eventually results in a new steady state. During the early stages of community dynamics,

no-longer-dormant local species undergo a secondary succession characterized by new

abiotic and biotic interactions. The emergence of resuscitation from dormancy is

considered as stochastic processes that lead to divergent communities (Buerger et al.,

2012; Lee et al., 2017; Lennon et al., 2021). When abiotic and biotic conditions allow

a stabilization of the initial communities, community formation at a later phase of a

press disturbance is expected to converge, through the succession of microbial species

(Philippot et al., 2021). In addition, the plasticity of microbial taxa in response to

recurrent micropollutant exposure has been shown to increase over multiple generations

(Cerro-Gálvez et al., 2021), with later-assembled communities shifting to a new stable

state that differs from the one before the exposure.

Here, a press disturbance was simulated in a microcosm experiment in which a

reservoir bacterial community was transferred into fresh medium containing trace

concentrations of typical micropollutants that were refreshed with recurrent

micropollutant exposure every 5 days. Several different micropollutants are typically

present simultaneously within aquatic environments, but whether their interactions lead

to effects on microbial communities that differ from those of single micropollutants is

unclear (Rodea-Palomares et al., 2016). Thus, we also experimentally manipulated

single and multiple micropollutants to determine their impacts on taxon coexistence, the regularity of community assembly, and micropollutant degradation ability. We hypothesized that because BPA elicits a subsidy response and TCS a stress response, the removal rate of BPA would be higher than that of TCS, with larger differences in the removal rates by microbial communities exposed to these micropollutants individually rather than in combination with other micropollutants. We also hypothesized that micropollutants degrade and alter the microbial community compositions after exposure. Accordingly, the microbial communities assembled at the initial and late phases of the micropollutant exposure would differ substantially and are primarily driven by either stochastic or deterministic assembly process.

Materials and Methods

Water collection

We selected the bacterial species pools from Shidou Reservoir, which is one of largest sources of drinking water in Xiamen, China (Hu et al., 2018) and is therefore under governmental protection from pollution originating from household drainages and a nearby dam (Fig. S1). Unlike other contaminated aquatic environments (Chalew & Halden, 2009; Im & Löffler, 2016; Sun, Li, Ma, et al., 2016), the concentrations of BPA, BPS, TCS, and TCC in Shidou Reservoir (China) are very low (< 1 ng/L), such that the reservoir's waters provide an ideal system to investigate the influence of persistent micropollutants on an aquatic microbiome. Surface water (100 L) from Shidou Reservoir (24°41'39"N, 118°0'37"E) was collected at 0.5 m depth on June 1, 2017 (Fig. S1). Temperature, pH, conductivity, dissolved oxygen (DO), and dissolved oxygen saturation, measured *in situ* using a HACH HQ40d multi-parameter meter (HACH, Loveland, CO, USA), were 29.0°C, 8.90, 54.4 µs/cm, 8.84 mg/L and 116.7%,

respectively. The concentrations of BPA, BPS, TCS, and TCC in the reservoir water

were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS)

(ABI 3200QTRAP, Framingham, MA, USA) with solid-phase extraction and all were

< 1 ng/L at the time of sampling (details of the measurements are provided below). The

collected water was transported from the field to the laboratory within 2 h after

sampling.

Experimental setup

BPA, BPS, TCS, and TCC were dissolved in acetone at concentrations of 800, 800, 400,

and 400 μg/L, respectively, and stored as stocks in glass bottles (HPLC grade). The

micropollutants were previously shown to be chemically stable when dissolved in an

organic solvent (Sun et al., 2017; Sun, Li, Ma, et al., 2016). From the 100 L of collected

water, 15 L were filtered through a 1.2-μm polycarbonate filter (Millipore, Ireland), to

minimize top-down regulation by planktonic grazers, and then used as the initial

inoculum. The remaining unfiltered water was filtered through a 0.22-μm

polycarbonate filter (Millipore, Cork, Ireland) to remove microorganisms and then

stored in the dark for later use in the preparation of incubation medium. Two treatments

consisting of single micropollutants (BPA and TCS) and two treatments consisting of

micropollutant mixtures (MI = BPA + TCS) and (MII = BPA + BPS + TCS + TCC)

were established.

Time-serial microcosms comprising seven inoculation batches from batch 1

(named hereafter 'B1') through batch 7 ('B7') were established to investigate the

microbial community response to a press disturbance of micropollutants resulting from

a single short-term event but maintained for 35 days (Fig. 1). Prior to the experimental

initiation, the caps of the bottles containing the micropollutants were loosened for > 12

h to volatilize the acetone to minimize its influence on the microbial responses. In

microcosm B1, 600 ml of the initial inoculum was added to bottles that already

contained the micropollutants and nutrients, such that the final concentration of each

micropollutant in the treatments was as follows: BPA: 1 μg/L BPA, TCS: 1 μg/L TCS,

MI: 1 μg/L BPA + 1 μg/L TCS, and MII: 1 μg/L BPA + 0.1 μg/L BPS + 1 μg/L TCS +

0.1 μg/L TCC. BPS and TCC are the derivatives of the BPA and TCS, respectively, and

the concentrations of those analogues in the environment are one order of magnitude

lower than their chief compounds in the seawater (Staples et al., 1998; McMurry et al.,

1998; Xie et al., 2022). The final concentration of each micropollutant in the

microcosms closely represented their ambient environmental conditions as reported in

literature (Chalew & Halden, 2009; Yamazaki et al., 2015). Further, given the low

nutrient concentrations (Hu et al., 2018; Kalyuzhny & Shnerb, 2017), we adjusted the

concentrations of nitrogen and phosphorus in the filtered water to 128.6 and 8.04 μM

to promote microbial growth by the addition of KNO_3 and KH_2PO_4, respectively (Wang

et al., 2016). After 5 days of incubation, 60 mL of the B1 inoculum was pipetted from

each of the replicates and transferred to a sterile bottle containing 540 mL of 0.22-μm-

filtered water and freshly added micropollutants. This newly inoculated microcosm

served as batch 2 (named hereafter 'B2'). The micropollutant treatments and controls

were amended with 128.6 μM N and 8.04 μM P after each inoculation to maintain

similar amounts of these nutrients throughout the experiment. The same inoculation

procedure was applied at each 5-day interval to initiate microcosms B3–B7, such that

the experimental period was 35 days (Fig. 1). Microbial cells were transferred between

pairwise replicates, with the remainder used in the downstream analyses described

below. An additional set of experimental bottles without micropollutant addition served

as the controls. In total, there were five treatments, each with three replicates, for a total

of 15 microcosm replicates per batch. All microcosms, i.e., 105 experimental bottles

during the 35 days, were incubated in the dark at 28°C with shaking at 100 rpm.

Additional experiments to test micropollutant degradation

Additional experiments were conducted to determine (i) micropollutant removal rates

in the absence of microorganisms, to rule out the gradual degradation of the

micropollutants in the treatments due to their chemical instability, and (ii) how the

ability of the microorganisms to transform micropollutants differed depending on

whether they have previous exposures or not. For the first additional experiment,

reservoir water was filtered through 0.22-μm filters and then amended with a mixture

consisting of 1 μg BPA/L, 0.1 μg BPS/L, 1 μg TCS/L, and 0.1 μg TCC/L as a blank

control. A second additional experiment was conducted to assess whether the ability of

bacterial assemblages to degrade BPA and TCS was enhanced by the evolving

community succession following micropollutant addition. Thus, 40 ml of the initial

inoculum and 40 ml of the bacterial assemblage at the end of the experiment (after

incubation of the B7 microcosms) were each amended with 160 ml of BPA, 160 ml of

TCS, or a mixture thereof, with each micropollutant present at a final concentration of

1 mg/L. The removal rates of BPA and TCS in these treatments after 7 days were

measured and the rates in the initial inocula vs. those in the B7 microcosms were

compared. The incubation conditions were the same for each treatment and the

experiment was conducted in triplicate.

Micropollutant analysis

For each microcosm batch, ~20 mL of the inoculum was removed from each bottle at

0, 2 and 5 days for micropollutant analysis. The four micropollutants BPA ($C_{15}H_{16}O_2$,

BPS ($C_{12}H_{10}O_4S$), TCS($C_{12}H_7Cl_{13}O_2$), and TCC ($C_{13}H_9Cl_3N_2O$) were concentrated

and purified by a 12-port Visiprep-DL SPE vacuum manifold (Supelco, Bellefonte, PA,

USA) with Oasis HLB cartridges (500 mg, 6 mL; Waters, Millford, MA, USA), and

then were measured using LC-MS/MS (ABI 3200QTRAP, Framingham, MA, USA),

as described in our previous work (Sun et al., 2017). Chromatographic separation was

performed by using a Kinetex C18 column (100 mm × 4.6 mm, 2.6 μm; Phenomenex,

Torrance, CA, USA) from Shimadzu LC system (Shimadzu, Japan).

DNA extraction and 16S rDNA amplicon high-throughput sequencing

The remaining inoculum (~470 mL) of each replicate was filtered onto 0.22-μm

polycarbonate filters (Millipore, Ireland), which were stored at −80°C until used for

DNA extraction. DNA from the 105 samples was extracted using the FastDNA for soil

kit (MP Biomedical, Santa Ana, CA, USA) according to the manufacturer's protocol,

except that the beating speed was set to 6.0 m/s for 80 s. The universal primer pair 515F

(5'-GTG YCA GCM GCC GCG GTA-3') and 907R (5'-CCG YCA ATT YMT TTR

AGT TT-3') was used to amplify the V4-V5 region of the bacterial 16S rRNA genes.

The PCR products were purified, mixed in equal amounts, and sequenced on an

Illumina MiSeq platform with the 2 × 300 bp paired-end protocol at the Majorbio Bio-

Pharm Technology Co., Ltd. (Shanghai, China).

Sequence processing

The raw sequencing data were processed using the LotuS pipeline (Hildebrand et al.,

2014) with the following criteria: i) average sequence quality > 27, ii) sequence length >

170 bp, iii) no ambiguous bases, iv) no mismatches with the barcodes or primers, and

v) homopolymer length < 8 bp. Operational taxonomic units (OTUs) were clustered at

a 97% similarity using UPARSE (Edgar, 2013). A representative sequence of each OTU

was taxonomically quantified using the RDP classifier with the Silva database v132

(Quast et al., 2013), at a bootstrap cutoff of 80%. All samples were rarified to 36, 000

sequence reads (the smallest library size). The subsampled OTU table contained 1375

OTUs. The α- and β-diversity indices of bacterial communities were calculated from

the rarified dataset using QIIME v1.9.1 (Caporaso et al., 2010) with the script

"core_diversity.py".

Inferring the community growth phase from the16S rRNA gene copy number

The 16S rRNA gene copy number among prokaryotes within a defined period indicates

the number of individuals within a community and reflects the growth phase of that

community (Klappenbach et al., 2000). In this study, the 16S rRNA gene copy number

of each OTU found in the samples was estimated using the rrnDB database v5.5

(Klappenbach et al., 2001) and the weighted mean rRNA gene copy number of all OTUs

found in each sample was then calculated as described previously (Wu et al., 2016).

The abundance-weighted average gene copy number of a community was plotted

against the sampling time for each treatment. The similar overarching pattern of the

average rRNA gene copy numbers of the bacterial communities in all microcosms over

time (Figs. S2A-F) indicated the similar growth trajectories of those communities. In

addition, a hierarchical cluster analysis with complete clustering criterion was used to

cluster the similarity of average gene copy numbers from all micropollutant-treated

microcosms belonging to the same batch type, with an exclusion of the controls (Fig.

S2G). Three growth progression phrases across all microcosm batches were identified

based on similar average gene copy numbers and the time continuity of the inoculation

(Fig. S2G): phase 1 (B1–B2), phase 2 (B3–B4), and phase 3 (B5–B7).

Relative contribution of ecological processes

According to a quantitative ecological framework (Stegen et al., 2013, 2015),

community assembly can be summarized by five major ecological processes: i)

homogenizing selection for a convergent composition between communities, ii)

variable selection for a divergent composition between communities, iii) dispersal

limitation for a divergent composition between communities due to limited dispersal

events, iv) homogenizing dispersal for a convergent composition between communities

due to unlimited dispersal, and v) ecological drift (referred to "undominated processes"

in (Stegen et al., 2015)). The phylogenetic turnover among the communities was

quantified by the weighted β-mean nearest taxon distance (βMNTD). The βNTI was

calculated as the difference between the observed βMNTD and the mean of the null

distribution (999 randomizations). The criteria determining the fraction of each process

were as follows: i) homogenizing and variable selection: higher (> 2) and lower (< -2)

βNTI, respectively; ii) dispersal limitation and homogenizing dispersal: non-significant

$|\beta NTI|$ (< 2) but $RC_{bray} > 0.95$ and < -0.95, respectively. iii) ecological drift: non-

significant $|\beta NTI|$ value (< 2) and non-significant $|RC_{bray}|$ value (< 0.95) (Stegen et al.,

2015). To infer the relative contributions of the different community assembly

processes to community turnover over time, for each pair consisting of a micropollutant

treatment and its control the βNTI and RC_{bray} values were calculated separately for each

of the three growth phrases. By comparing the community assembly process in the

treatment vs. the controls, we were able to minimize confounding effects that may have

obscured community variation and thus better predict the treatment effect. The relative

contribution was expressed as the portion of pairwise sample assembly that could be

attributed to each ecological process.

Ecological grouping of microbial responses to micropollutants

The responses of individual OTUs to micropollutant addition were identified by

analyzing the differences in the relative abundance of each OTU during the three

growth phases for each treatment, using the 'DESeq2' R package v 1.26.0 (Love et al.,

2014). Significant values were corrected for multiple tests using the Benjamini-

Hochberg correction with an adjusted α value of 0.05. The responses were categorized

as: i) sensitive, when relative OTU abundances were significantly higher in phase 1

than in phases 2 and 3; ii) opportunistic, when relative abundances were significantly

higher in phase 2 than in phases 1 and 3; iii) tolerant, when relative abundances were

significantly higher in phase 3 than in phases 1 and 2. The ecological categories

represented a group of bacterial members whose growth fitness optimal in response to

repeated exposure could be documented collectively. In addition, to visualize the

phylogenetic relationships of the OTUs assigned to the ecological groups,

representative sequences of those OTUs were used to construct a phylogenetic tree

using iTOL v5 (Letunic & Bork, 2019).

Micropollutant turnover and community analysis

A Wilcoxon's test was used to assess the removal rate of BPA, TCS, BPS, and TCC in

consecutive microcosm batches for the single-micropollutant (BPA and TCA)

treatments and the mixed-micropollutant (MI and MII) treatments. Additionally, a

Kruskal-Wallis test was used to analyze the difference in the removal rates between the

single micropollutant treatments and between any two micropollutants for the mixture

treatments. In this case, the removal rate of the micropollutants from each microcosm

batch was used as replicates for the analysis.

Differences in the bacterial communities in all microcosms were explored using

non-metric multidimensional scaling (NMDS) based on the Bray-Curtis dissimilarity

metric and by fitting removal rates of micropollutants to the ordination. This allowed

us to assess whether the degradation of micropollutants was tied to the microbial

succession. A permutational multivariate analysis of variance (PERMANOVA)

(Anderson, 2001) was conducted to test the effects of micropollutants, inoculation time

(as indicated by the microcosm batch), and their interactive effects on the taxonomic

composition of the bacterial communities. A pairwise PERMANOVA was performed

to test for significant differences in community composition between each treatment

pair. The multivariate analyses were performed using the R package 'vegan' (Oksanen

et al., 2019). Whether communities with the shared taxa exhibit similar dynamics was

assessed in a dissimilarity-overlap analysis (Bashan et al., 2016). Briefly, the relative

abundance of each OTU in each community was calculated to identify sets of OTUs

within each pair of communities (i.e., within a treatment over time). In the calculation

of dissimilarity, only the shared OTUs were considered, and the relative abundance of

shared taxa was transformed to add up to 1. The dissimilarity between renormalized

vectors X and Y can be calculated using root Jensen–Shannon divergence. Based on

the dissimilarity-overlap analysis, 15 pairwise comparisons were obtained in this study.

This method and the relevant R script were modified from Vila et al. (2020). Moreover,

the rate of community turnover in relative to the controls over time was quantified with

a linear regression model (see the Supplemental methods for details). A Wilcoxon test

was then applied to test for significant differences in community turnover between

microcosm B1 (day 5) and the other microcosm batches from later time points (B2–

B7), using the 'ggsignif' package (Ahlmann-Eltze & Patil, 2021). The data obtained in

this study were visualized using the R package 'ggplot2'.

Results

Transformation of trace micropollutants

There was no evidence that the removal rate of any of the micropollutants differed

between any of the consecutive microcosm batches (Fig. 2, Wilcoxon's test, all $P >$

0.05). In the microcosms treated with single micropollutant, most of BPA added to each

microcosm batch was degraded, achieving > 80% removal in the individual batches,

whereas the removal rate of TCS was less than 60% (Fig. 2). When comparing the

removal rates of BPA and TCS in the MI microcosms, there was no evidence that the

removal of the two micropollutants from the MI microcosms differed ($P > 0.05$) (17.86

± 15.17% for BPA and 25.11 ± 18.88 % for TCS; Fig. S3B). In the MII microcosms,

the percentage of removed BPA (65.83 ± 6.54%) was highest ($P < 0.05$), followed by

the TCS (13.76 ± 16.76%), BPS (5.93 ± 8.97%), and TCC (11.97 ± 14.08%), but the

differences were not significant (Fig. S3B).

The removal proportion of the four micropollutants in the bacteria-free controls

was stable during the 7-day incubation, as shown in the first additional experiment (Fig.

S4). This therefore suggests that transformation of the micropollutants is negligible in

the absence of microorganisms and light. The second experiment showed that

microorganisms pre-exposed to low concentrations of BPA and TCS fully degraded

higher concentrations of those micropollutants during subsequent exposures and

degradation occurred at a faster rate (Fig. S5), but only when BPA and TCS were

provided alone and not in combination.

Community succession following micropollutant exposure

The bacterial communities in the microcosms became taxonomically distinct both over

time and by micropollutant type (Fig. 3). There was very weak evidence that the

variation in the community composition among all microcosm batches correlated with

differences in the removal rates of BPA (P =0.06, Table S1), with high association in

its single treatments. However, there was strong evidence that the community variation

was correlated with TCS, with high association in TCS single and mixed treatments

(Fig. 3 and Table S1, P =0.001). The inoculation time (PERMANOVA, R^2 = 0.149),

micropollutant addition (R^2 = 0.11), and their interaction (R^2 = 0.055) significantly

influenced the community composition (all $P < 0.001$) (Table S2). Irrespective of the

inoculation time, the composition of the TCS, MI, and MII communities differed

significantly from that of the controls (pairwise PERMANOVA, $P < 0.01$; Table S2)

but not from each other. Conversely, the BPA communities significantly differed from

the other three treatments but not from the controls.

Overall, the dissimilarity in the micropollutant-treated vs. control communities

significantly increased across microcosm batches, as each inoculation was replenished

with micropollutants (linear regression, all $P < 0.001$, Fig. S6). Furthermore, the

communities of the first inoculation (B1) impacted the response of some communities

inoculated at later time points (Fig. 4A). There was no evidence for the impact of the

B1 community composition on later communities across B2–B6 in the BPA-treated

microcosms and B2–B5 in the MII-treated microcosms, because of their community

resemblances (Wilcoxon test, $P > 0.05$; Fig. 4A). In both BPA and MII communities,

the correlation between dissimilarity and overlap (i.e., the number of shared OTUs between the community pair) was negative ($P < 0.05$ Fig. 4B). This result was not unexpected since the dispersal of microbial cells from one condition to another similar environment may drive community dynamics towards convergence (Lindström & Östman, 2011; Shen et al., 2018). However, in the TCS and MI treatments, an impact of the initial communities on later ones occurred only in B3, which is supported by the fact that the B3–B7 communities were more dissimilar to B1 ($P < 0.05$; Fig. 4A). The negative correlation of the dissimilarity and overlap analysis was nonsignificant for the TCS and MI communities such that the high overlap did not result in a low dissimilarity ($P > 0.05$; Fig. 4B). This suggests that there were differences in the dynamics of the shared taxa in those TCS and MI communities over time compared to the communities in the other treatments.

OTUs with distinct responses to micropollutant addition

In all treatments, a greater number of sensitive OTUs (41–64), a moderate number of tolerant OTUs (6–15), and fewer opportunistic OTUs (0–3) were identified across phases 1 (B1−B2), 2 (B3−B4), and 3(B5−B7) (Table 1 and Fig. 5). The ecological groups replaced one another over time, together accounting for a minimal relative abundance of 0.5 % and a maximum of 60% in the individual samples (Fig. S7). Regardless of the micropollutant type, members of Actinobacteria, Planctomycetes, and Verrucomicrobia primarily consisted of sensitive OTUs (Table 1; Fig. 5). The tolerant OTUs were primarily comprised of members affiliated with Alphaproteobacteria, Gammaproteobacteria and Bacteroidetes. The majority of the tolerant OTUs were unique to one treatment, with only two OTUs (OTU_48 and OTU_78) tolerating micropollutant addition in all treatments (Fig. 5 and Table S4). By contrast, many

sensitive OTUs (51) were shared between treatments, and at least between two treatments (Table S4). Among the overlapping, 20 OTUs found in all treatments, maximal relative abundance of which was mainly detected in the initial inocula or the controls, and most were members of Actinobacteria (Table S4; Fig. 5). For instance, a strictly sensitive taxa Actinobacteria *ACK-M1*, found in both single and mixed micropollutant treatments, exhibited its maximal relative abundance (15.94%) in the initial inocula. Whereas a tolerant taxa Alphaproteobacteria *Novosphingobium*, shared between TCS and MI, had its maximal relative abundance (30.48%) in the BPA treatments at late exposure batches (B6), which has been reported as TCS-degrading bacteria in wastewater treatments and agricultural soil (Zhou et al., 2013; Thelusmond et al., 2019).

Partition of ecological processes on community variation

Overall, homogenizing selection played a minor role in influencing community variation across phases for each community pair (micropollutant-treated and control), as it accounted for less than 8.33% of the ecological processes (Fig. 6). Instead, variable selection (13.89–61.73%) and ecological drift (30.86–80.56%) dominated the ecological processes across the three phases. These two processes exhibited compensatory contributions, showing increasing variable selection but decreasing ecological drift along the phases (Fig. 6). In the case of dispersal limitation, its relative proportion was greater at phase 1 than the other phases for all treatments, especially BPA and MII.

Discussion

We experimentally investigated the effects of persistent micropollutants on microbial

community dynamics to gain insights into community assembly mechanisms. Our first

hypothesis that differences in the removal rates of BPA vs. TCS in environments

exposed to single vs. mixed micropollutants was supported. he energy required by the

microbial communities to counteract the opposing responses to BPA and TCS in the MI

treatment (Choi & Lee, 2017; Zhang et al., 2013) likely accounted for the profound

inhibition of BPA degradation. In MII, a significantly larger fraction of BPA than of the

three other micropollutants was removed, indicating that the presence of TCC reduced

the inhibitory effect of TCS on BPA transformation and thus degradation. The

mechanism underlying such effect is beyond the scope of this study. Phenolic

compounds (both BPA and BPS) act as a growth medium for microorganisms by

providing a source of carbon and energy. However, they differ in degradability in the

seawater (BPA > BPS, Danzl et al., 2009), as well as negative impacts on biochemical

and microbiological activity of soil (BPA < BPS), Zaborowska et al., 2020). These lines

of reasoning explained a greater degradation of BPA than its analogue in MII (Fig. 2).

Altogether, our findings obtained from all three experiments demonstrate that

degradation of micropollutants is a microbially-mediated rather than an ordinary

chemical process. Pre-exposure to micropollutants can enhance the degradation

capacity of microbes later confronted with those or similar chemicals, which is

consistent with previous literature (Cerro-Gálvez et al., 2021; Coll et al., 2020).

Given that the ability of bacterial communities to transform the four

micropollutants surveyed, the questions remain: whether micropollutants accelerate

degradation through changes in bacterial communities, or the micropollutant degraded

and then altered the community composition? We found no evidence that the

degradation of any micropollutants differed across microcosm batches. Rather, there

was strong evidence on the differences in the bacterial communities between the

micropollutant-added microcosms vs. the controls, and that community turnover increased over time following micropollutant exposure (Figs. 4A, S6). These results supported the second hypothesis that the degradation of micropollutants alter the microbial communities. Such community dynamics is mainly linked by the degradation of BPA and TCS in the microcosms. We identified ecological groups based on changes in the relative abundance of taxa following repeated micropollutant exposure, which integrates population growth, death, survival, and reproduction over 35 days. Both sensitive and tolerant taxa identified here may have affected micropollutant degradation either directly or indirectly, such as by providing enzymatic repertoires to other community members that enabled degradation, including interspecies interactions (Freilich et al., 2011; Pacheco et al., 2019). The larger proportion of sensitive than tolerant taxa in all treatments suggested that the microbial community was not robust to the persistence of micropollutants in its surroundings. Noteworthily, two strictly tolerant genera across treatments were *Sphingomonas* and *Hydrogenophaga* (Table S4), which have been shown to be enriched in the BPA or TCS-contaminated environments (Oh et al., 2019; Zaborowska et al., 2020; Hagberg et al., 2021) and to produce enzymes capable of degrading different pollutants (Gao et al., 2019).

Our results supported the third hypothesis, that communities assembled at the early stage of micropollutant exposure would be compositionally distinct from those that formed at the later stage. Our findings generally corroborated previous studies showing that the persistence of micropollutants even at low concentrations can influence microbial community membership and abundance over time (Coll et al., 2020; Hu et al., 2017; Wolff et al., 2018). Despite a substantial community variation between initial and late stages of the exposure, community trajectory in the BPA and MII were driven by a set of shared taxa that interacted similarly with the respective environments and

other community members. Conversely, the less pronounced negative dissimilarity-overlap relationships in the case of the TCS and MI treatments indicated that the microbial taxa shared among different inoculation batches within those treatments interacted differently with the same environment over time. Hence, microbial communities assembled in identical habitats can perceive those habitats differently such that their feedback on the surroundings and other, coexisting members will differ as well, as noted elsewhere (Bajić et al., 2018; Vila et al., 2020).

We observed a gradual transition from stochastic to deterministic processes towards the later stage of micropollutant exposure. In all treatments ecological drift played a major role during phases 1 and 2 of community growth progression. This can be explained by the random changes in abundances within microbial communities exposed to micropollutants in confined microcosms, where dispersal is limited and locally dormant cells are about to be awakened to colonize empty niches in the environments upon disturbances (Nemergut et al., 2013; Philippot et al., 2021; Zhou & Ning, 2017). In fact, sensitive taxa declined in the relative abundances while tolerant taxa had negligible abundance (Fig. S7) during the growth phases 1-2 (B1-B2 and B3-B4). Hence, tolerant taxa identified in this study were recruited from rare member pool at the early stage of exposure and regrew to a large population size (~ relative abundance of 10 %) at the later stage of exposure. However, the increasing contribution of variable selection to community assembly over time suggested that the high phylogenetic turnover of the communities was driven by heterogenous abiotic and biotic conditions (Stegen et al., 2015; Vellend, 2010). The increasing effects of the micropollutants on the communities over time led to higher species selection and an

elevated impact on the interactions among those species within the community, such that the drivers of community assembly no longer resembled those of the control and resulted in a high rate of community turnover. Our findings suggest that the repeated introduction of a micropollutant enhances succession in microbial communities (Jiao et al., 2017).

We also considered the potential role of priority effects on community assembly, independent of the ecological process model used in our study. Priority effects emerge when early colonizers exhaust or modify available resources, thus influencing the establishment success of later colonizers (Fukami, 2015). A strong priority effect in the case of TCS and MI was evidenced by the significant difference in the composition of the initially inoculated vs. any of the later inoculated communities (Fig. 4A). Once they were re-inoculated, taxa that had thrived in the B1 batch modified the environments of later batches, such that some taxa (presumably activated from a seed bank, given that no source of new taxa was externally introduced into the systems) were less or more favorable in later batches. However, for the BPA and MII treatments, the effects occurred at later stage of the press disturbance, suggesting the increasing strength of priority effects with increasing establishment time, as noted elsewhere (Svoboda et al., 2018).

Although our microcosms represent an ideal system which is removed from *in situ* conditions, our experimental findings are valuable in at least three aspects, that is : i) the prediction of microbial responses to persistent micropollutants, ii) the   harnessing of ecological principles and microbiomes in pollution management (Bernstein, 2019;

Borchert et al., 2021; Rocca et al., 2021), and (iii) the development of a conceptual

model to be used to predict when and how stochastic and deterministic processes

contribute to community assembly after the exposure of persistent micropollutants. Our

results suggest that strong ecological drift drives random community changes at the

early stage of the exposure (Fig. 7). Prolonged ecological drift, variable selection

resulting from the effects of micropollutants, and the abiotic and biotic conditions

experienced differently by different microbes then determine community succession at

the mid-stage of the exposure. Additionally, the high environmental heterogeneity

together with the low densities of the communities permitted the emergence of priority

effects, in which the early establishment of taxa in micropollutant-containing

environments allowed a faster community succession. This was supported by the

decrease in the average gene copy number at the intermediate phase (Fig. S2), when

community assembly was driven by various ecological drivers (Nemergut et al., 2016).

Finally, strong variable selection accompanied by increased priority effects likely set

the communities on distinct trajectories, with multiple equilibria at the late stage of the

exposure. In fact, more advanced community succession and multiple equilibria

characterized the end of the press disturbance, and both played a larger role in the TCS

and MII (and to a lesser extent in the BPA) communities than in the MI community

(Fig. 7), suggesting convergent recovery of those freshwater bacterial communities to

repeated micropollutant exposure.

Conclusions

In summary, we evidenced that micropollutants degrade and drive microbial succession following repeated exposure. This work can aid in microbe-pollution management by providing insights into micropollutant removal as well as microbial assembly mechanisms and dynamics. From the perspectives of microbiome engineering, this study demonstrates the importance of the microbiome's preexposure history in the design of long-term experiments that include artificial selection to direct the evolution (Borchert et al., 2021; Chang et al., 2021) of a target microbiome highly optimized for pollutant removal in an aquatic environment.

Availability of data and material

The sequencing data were deposited in the NCBI SRA database with the Bioproject number PRJNA497759. Our R scripts for statistics, data visualization and computing notes are available on a zenodo archive (https://doi.org/10.5281/zenodo.5242686) of the Github repository (https://github.com/AnyiHu/Impacts-of-micropollutants-on-freshwater-microbiome)

Acknowledgements

We thank Dr. Linwei Wu for her assistance in calculating the copy number of 16S rRNA genes, and Dr. Hongjie Wang for his assistance in the molecular experiments. This work was supported by the National Natural Science Foundation of China (31870475, U1805244 and 31470539), the International Partnership Program of Chinese Academy of Sciences (132C35KYSB20200012), and the Natural Science Foundation of Fujian, China (2019I0030).

Author contributions

Dandan Izabel-Shen: Conceptualization, Formal analysis, Validation, Software, Visualization, Writing - original draft. **Shuang Li**: Investigation, Methodology, Formal analysis, Data curation, Visualization, Writing - review & editing. **Tingwei Luo**: Methodology, Resources. **Jianjun Wang**: Writing - review & editing. **Yan Li:** Methodology. **Qian Sun**: Methodology. **Chang-Ping Yu**: Resources, Supervision, Funding acquisition. **Anyi Hu**: Conceptualization, Data curation, Software, Validation, Writing - review & editing, Project administration, Funding acquisition.

References

Adyari, B., Shen, D., Li, S., Zhang, L., Rashid, A., Sun, Q., Hu, A., Chen, N., & Yu, C. P. (2020). Strong impact of micropollutants on prokaryotic communities at the horizontal but not vertical scales in a subtropical reservoir, China. In Science of the Total Environment (Vol. 721). https://doi.org/10.1016/j.scitotenv.2020.137767

Ahlmann-Eltze, C., & Patil, I. (2021). *ggsignif: Significance Brackets for "ggplot2". R package version 0.6.1.* (Vol. 3, Issue 123). https://cran.r-project.org/package=ggsignif

Anderson, M. J. (2001). Permutation tests for univariate or multivariate analysis of variance and regression. *Canadian Journal of Fisheries and Aquatic Sciences*, *58*(3), 626–639. https://doi.org/10.1139/cjfas-58-3-626

Bajić, D., Vila, J. C. C., Blount, Z. D., & Sánchez, A. (2018). On the deformability of an empirical fitness landscape by microbial evolution. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(44), 11286–11291. https://doi.org/10.1073/pnas.1808485115

Bashan, A., Gibson, T. E., Friedman, J., Carey, V. J., Weiss, S. T., Hohmann, E. L., & Liu, Y. Y. (2016). Universality of human microbial dynamics. *Nature*, *534*(7606), 259–262. https://doi.org/10.1038/nature18301

Bernstein, H. C. (2019). Reconciling Ecological and Engineering Design Principles for Building Microbiomes. *MSystems*, *4*(3), 1–5. https://doi.org/10.1128/msystems.00106-19

Borchert, E., Hammerschmidt, K., Hentschel, U., & Deines, P. (2021). Enhancing

Microbial Pollutant Degradation by Integrating Eco-Evolutionary Principles with
Environmental Biotechnology. *Trends in Microbiology*, *29*(10), 908–918.
https://doi.org/10.1016/j.tim.2021.03.002

Boyd, T. J., Smith, D. C., Apple, J. K., Hamdan, L. J., Osburn, C. L., & Montgomery,
M. T. (2008). Evaluating PAH biodegradation relative to total bacterial carbon
demand in coastal ecosystems: Are PAHs truly recalcitrant? In *Microbial Ecology*
Research Trends.

Buerger, S., Spoering, A., Gavrish, E., Leslin, C., Ling, L., & Epstein, S. S. (2012).
Microbial scout hypothesis, stochastic exit from dormancy, and the nature of slow
growers. *Applied and Environmental Microbiology*, *78*(9), 3221–3228.
https://doi.org/10.1128/AEM.07307-11

Cabana, H., Jiwan, J. L. H., Rozenberg, R., Elisashvili, V., Penninckx, M., Agathos, S.
N., & Jones, J. P. (2007). Elimination of endocrine disrupting chemicals
nonylphenol and bisphenol A and personal care product ingredient triclosan using
enzyme preparation from the white rot fungus Coriolopsis polyzona. *Chemosphere*,
67(4), 770–778. https://doi.org/10.1016/j.chemosphere.2006.10.037

Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello,
E. K., Fierer, N., Peña, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley,
S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., Mcdonald, D.,
Muegge, B. D., Pirrung, M., … Knight, R. (2010). correspondence QIIME allows
analysis of high- throughput community sequencing data Intensity normalization
improves color calling in SOLiD sequencing. *Nature Publishing Group*, *7*(5),
335–336. https://doi.org/10.1038/nmeth0510-335

Cerro-Gálvez, E., Dachs, J., Lundin, D., Fernández-Pinos, M. C., Sebastián, M., &
Vila-Costa, M. (2021). Responses of Coastal Marine Microbiomes Exposed to
Anthropogenic Dissolved Organic Carbon. *Environmental Science and*
Technology, 55: 9609–9621. https://doi.org/10.1021/acs.est.0c07262

Chalew, T. E. A., & Halden, R. U. (2009). Environmental exposure of aquatic and
terrestrial biota to triclosan and triclocarban. *Journal of the American Water*
Resources Association, *45*(1), 4–13. https://doi.org/10.1111/j.1752-
1688.2008.00284.x

Chang, C. Y., Vila, J. C. C., Bender, M., Li, R., Mankowski, M. C., Bassette, M.,
Borden, J., Golfier, S., Sanchez, P. G. L., Waymack, R., Zhu, X., Diaz-Colunga,
J., Estrela, S., Rebolleda-Gomez, M., & Sanchez, A. (2021). Engineering complex
communities by directed evolution. *Nature Ecology and Evolution*, *5*(7), 1011–
1023. https://doi.org/10.1038/s41559-021-01457-5

Choi, Y. J., & Lee, L. S. (2017). Aerobic Soil Biodegradation of Bisphenol (BPA)
Alternatives Bisphenol S and Bisphenol AF Compared to BPA. *Environmental*
Science and Technology, *51*(23), 13698–13704.

https://doi.org/10.1021/acs.est.7b03889

Coll, C., Bier, R., Li, Z., Langenheder, S., Gorokhova, E., & Sobek, A. (2020). Association between Aquatic Micropollutant Dissipation and River Sediment Bacterial Communities. *Environmental Science and Technology*, *54*(22), 14380–14392. https://doi.org/10.1021/acs.est.0c04393

Danzl, E., Sei, K., Soda, S., Ike, M., & Fujita, M. (2009). Biodegradation of bisphenol A, bisphenol F and bisphenol S in seawater. *International journal of environmental research and public health*, 6(4), 1472-1484. https://doi.org/10.3390/ijerph6041472

da Silva, G. C. X., Medeiros de Abreu, C. H., Ward, N. D., Belúcio, L. P., Brito, D. C., Cunha, H. F. A., & da Cunha, A. C. (2020). Environmental Impacts of Dam Reservoir Filling in the East Amazon. *Frontiers in Water*, *2*(May). https://doi.org/10.3389/frwa.2020.00011

Deblonde, T., Cossu-Leguille, C., & Hartemann, P. (2011). Emerging pollutants in wastewater: A review of the literature. *International Journal of Hygiene and Environmental Health*, *214*(6), 442–448. https://doi.org/10.1016/j.ijheh.2011.08.002

Edgar, R. C. (2013). UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nature Methods*, *10*(10), 996–998. https://doi.org/10.1038/nmeth.2604

Edward A. Bender, T. J. C. and M. E. G. (1984). Perturbation Experiments in Community Ecology: Theory and Practice. *Ecological Society of Amenica*, 1–13.

Eggen, R. I. L., Hollender, J., Joss, A., Schärer, M., & Stamm, C. (2014). Reducing the discharge of micropollutants in the aquatic environment: The benefits of upgrading wastewater treatment plants. *Environmental Science and Technology*, *48*(14), 7683–7689. https://doi.org/10.1021/es500907n

Freilich, S., Zarecki, R., Eilam, O., Segal, E. S., Henry, C. S., Kupiec, M., Gophna, U., Sharan, R., & Ruppin, E. (2011). Competitive and cooperative metabolic interactions in bacterial communities. *Nature Communications*, *2*(1), 587–589. https://doi.org/10.1038/ncomms1597

Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches, Species Pools, and Priority Effects. *Annual Review of Ecology, Evolution, and Systematics*, *46*, 1–23. https://doi.org/10.1146/annurev-ecolsys-110411-160340

Gao, H., LaVergne, J. M., Carpenter, C. M., Desai, R., Zhang, X., Gray, K., ... & Wells, G. F. (2019). Exploring co-occurrence patterns between organic micropollutants and bacterial community structure in a mixed-use watershed. *Environmental Science: Processes & Impacts*, 21(5), 867-880. https://doi.org/10.1039/C8EM00588E

Hagberg, A., Gupta, S., Rzhepishevska, O., Fick, J., Burmølle, M., & Ramstedt, M. (2021). Do environmental pharmaceuticals affect the composition of bacterial communities in a freshwater stream? A case study of the Knivsta river in the south of Sweden. *Science of the Total Environment*, 763, 142991. https://doi.org/10.1016/j.scitotenv.2020.142991

Hu, A., Ju, F., Hou, L., Li, J., Yang, X., Wang, H., Mulla, S. I., Sun, Q., Bürgmann, H., & Yu, C. P. (2017). Strong impact of anthropogenic contamination on the co-occurrence patterns of a riverine microbial community. *Environmental Microbiology*, *19*(12), 4993–5009. https://doi.org/10.1111/1462-2920.13942

Hu, A., Li, S., Zhang, L., Wang, H., Yang, J., Luo, Z., Rashid, A., Chen, S., Huang, W., & Yu, C. P. (2018). Prokaryotic footprints in urban water ecosystems: A case study of urban landscape ponds in a coastal city, China. *Environmental Pollution*, *242*, 1729–1739. https://doi.org/10.1016/j.envpol.2018.07.097

Im, J., & Löffler, F. E. (2016). Fate of Bisphenol A in Terrestrial and Aquatic Environments. *Environmental Science and Technology*, *50*(16), 8403–8416. https://doi.org/10.1021/acs.est.6b00877

Jiao, S., Zhang, Z., Yang, F., Lin, Y., Chen, W., & Wei, G. (2017). Temporal dynamics of microbial communities in microcosms in response to pollutants. *Molecular Ecology*, *26*(3), 923–936. https://doi.org/10.1111/mec.13978

Kalyuzhny, M., & Shnerb, N. M. (2017). Dissimilarity-overlap analysis of community dynamics: Opportunities and pitfalls. *Methods in Ecology and Evolution*, *8*(12), 1764–1773. https://doi.org/10.1111/2041-210X.12809

Klappenbach, J. A., Dunbar, J. M., & Schmidt, T. M. (2000). rRNA operon copy number reflects ecological strategies of bacteria. *Applied and environmental microbiology*, 66(4), 1328–1333. https://doi.org/10.1128/AEM.66.4.1328-1333.2000

Klappenbach, J. A., Saxman, P. R., Cole, J. R., & Schmidt, T. M. (2001). Rrndb: The ribosomal RNA operon copy number database. *Nucleic Acids Research*, *29*(1), 181–184. https://doi.org/10.1093/nar/29.1.181

Kuroda, K., Murakami, M., Oguma, K., Muramatsu, Y., Takada, H., & Takizawa, S. (2012). Assessment of groundwater pollution in Tokyo using PPCPs as sewage markers. *Environmental Science and Technology*, *46*(3), 1455–1464. https://doi.org/10.1021/es202059g

Lee, S. H., Sorensen, J. W., Grady, K. L., Tobin, T. C., & Shade, A. (2017). Divergent extremes but convergent recovery of bacterial and archaeal soil communities to an ongoing subterranean coal mine fire. *ISME Journal*, *11*(6), 1447–1459. https://doi.org/10.1038/ismej.2017.1

Lennon, J. T., den Hollander, F., Wilke-Berenguer, M., & Blath, J. (2021). Principles of seed banks and the emergence of complexity from dormancy. *Nature*

Communications, *12*(1), 1–16. https://doi.org/10.1038/s41467-021-24733-1

Letunic, I., & Bork, P. (2019). Interactive Tree of Life (iTOL) v4: Recent updates and new developments. *Nucleic Acids Research*, *47*(W1), 2–5. https://doi.org/10.1093/nar/gkz239

Lindström, E. S., & Östman, Ö. (2011). The importance of dispersal for bacterial community composition and functioning. *PLoS ONE*, *6*(10). https://doi.org/10.1371/journal.pone.0025883

Liu, W. R., Zhao, J. L., Liu, Y. S., Chen, Z. F., Yang, Y. Y., Zhang, Q. Q., & Ying, G. G. (2015). Biocides in the Yangtze River of China: Spatiotemporal distribution, mass load and risk assessment. *Environmental Pollution*, *200*, 53–63. https://doi.org/10.1016/j.envpol.2015.02.013

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21. https://doi.org/10.1186/s13059-014-0550-8

Luo, Y., Guo, W., Ngo, H. H., Nghiem, L. D., Hai, F. I., Zhang, J., Liang, S., & Wang, X. C. (2014). A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. *Science of the Total Environment*, *473–474*, 619–641. https://doi.org/10.1016/j.scitotenv.2013.12.065

Martinez-Varela, A., Cerro-Gálvez, E., Auladell, A., Sharma, S., Moran, M. A., Kiene, R. P., Piña, B., Dachs, J., & Vila-Costa, M. (2021). Bacterial responses to background organic pollutants in the northeast subarctic Pacific Ocean. *Environmental Microbiology*, *23*(8), 4532–4546. https://doi.org/10.1111/1462-2920.15646

McMurry, L. M., Oethinger, M., & Levy, S. B. (1998). Triclosan targets lipid synthesis [4]. *Nature*, *394*(6693), 531–532. https://doi.org/10.1038/28970

Nemergut, D. R., Knelman, J. E., Ferrenberg, S., Bilinski, T., Melbourne, B., Jiang, L., Violle, C., Darcy, J. L., Prest, T., Schmidt, S. K., & Townsend, A. R. (2016). Decreases in average bacterial community rRNA operon copy number during succession. *ISME Journal*, *10*(5), 1147–1156. https://doi.org/10.1038/ismej.2015.191

Nemergut, D. R., Schmidt, S. K., Fukami, T., O'Neill, S. P., Bilinski, T. M., Stanish, L. F., Knelman, J. E., Darcy, J. L., Lynch, R. C., Wickey, P., & Ferrenberg, S. (2013). Patterns and Processes of Microbial Community Assembly. *Microbiology and Molecular Biology Reviews*, *77*(3), 342–356. https://doi.org/10.1128/mmbr.00051-12

Oh, S., Choi, D., & Cha, C. J. (2019). Ecological processes underpinning microbial community structure during exposure to subinhibitory level of triclosan. *Scientific reports*, 9(1), 1-12. https://doi.org/10.1038/s41598-019-40936-5

Okere, U. V., Cabrerizo, A., Dachs, J., Ogbonnaya, U. O., Jones, K. C., & Semple, K. T. (2017). Effects of pre-exposure on the indigenous biodegradation of 14C-phenanthrene in Antarctic soils. *International Biodeterioration and Biodegradation*, *125*, 189–199. https://doi.org/10.1016/j.ibiod.2017.09.013

Oksanen, A. J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., Minchin, P. R., Hara, R. B. O., Simpson, G. L., Solymos, P., Stevens, M. H. H., & Szoecs, E. (2019). Vegan: community ecology package. In *Encyclopedia of Food and Agricultural Ethics*. https://doi.org/10.1007/978-94-024-1179-9_301576

Pacheco, A. R., Moel, M., & Segrè, D. (2019). Costless metabolic secretions as drivers of interspecies interactions in microbial ecosystems. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-018-07946-9

Philippot, L., Griffiths, B. S., & Langenheder, S. (2021). Microbial Community Resilience across Ecosystems and Multiple Disturbances. *Microbiology and Molecular Biology Reviews*, *85*(2). https://doi.org/10.1128/mmbr.00026-20

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, *41*(D1), 590–596. https://doi.org/10.1093/nar/gks1219

Roberts, J., Kumar, A., Du, J., Hepplewhite, C., Ellis, D. J., Christy, A. G., & Beavis, S. G. (2016). Pharmaceuticals and personal care products (PPCPs) in Australia's largest inland sewage treatment plant, and its contribution to a major Australian river during high and low flow. *Science of the Total Environment*, *541*, 1625–1637. https://doi.org/10.1016/j.scitotenv.2015.03.145

Rocca, J. D., Muscarella, M. E., Peralta, A. L., Izabel-Shen, D., & Simonin, M. (2021). Guided by Microbes: Applying Community Coalescence Principles for Predictive Microbiome Engineering. *MSystems*, *6*(4). https://doi.org/10.1128/msystems.00538-21

Rodea-Palomares, I., Gonzalez-Pleiter, M., Gonzalo, S., Rosal, R., Leganes, F., Sabater, S., Casellas, M., Muñoz-Carpena, R., & Fernández-Piñas, F. (2016). Hidden drivers of low-dose pharmaceutical pollutant mixtures revealed by the novel GSA-QHTS screening method. *Science Advances*, *2*(9), 1–12. https://doi.org/10.1126/sciadv.1601272

Schwarzenbach, R. P., Escher, B. I., Fenner, K., Hofstetter, T. B., Johnson, C. A., Von Gunten, U., & Wehrli, B. (2006). The challenge of micropollutants in aquatic systems. *Science*, *313*(5790), 1072–1077. https://doi.org/10.1126/science.1127291

Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., Huber, D. H., Langenheder, S., Lennon, J. T., Martiny, J. B. H., Matulich, K. L., Schmidt, T.

M., & Handelsman, J. (2012). Fundamentals of microbial community resistance and resilience. *Frontiers in Microbiology*, *3*(DEC), 1–19. https://doi.org/10.3389/fmicb.2012.00417

Shen, D., Langenheder, S., & Jürgens, K. (2018). Dispersal modifies the diversity and composition of active bacterial communities in response to a salinity disturbance. *Frontiers in Microbiology*, *9*(SEP). https://doi.org/10.3389/fmicb.2018.02188

Staples, C. A., Dorn, P. B., Klecka, G. M., O'Block, S. T., & Harris, L. R. (1998). A review of the environmental fate, effects, and exposures of bisphenol A. *Chemosphere*, *36*(10), 2149–2173. https://doi.org/10.1016/S0045-6535(97)10133-3

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L., & Konopka, A. (2013). Quantifying community assembly processes and identifying features that impose them. *ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Stegen, J. C., Lin, X., Fredrickson, J. K., & Konopka, A. E. (2015). Estimating and mapping ecological processes influencing microbial community assembly. *Frontiers in Microbiology*, *6*(MAY), 1–15. https://doi.org/10.3389/fmicb.2015.00370

Sun, Q., Li, M., Ma, C., Chen, X., Xie, X., & Yu, C. P. (2016). Seasonal and spatial variations of PPCP occurrence, removal and mass loading in three wastewater treatment plants located in different urbanization areas in Xiamen, China. *Environmental Pollution*, *208*, 371–381.

Sun, Q., Wang, Y., Li, Y., Ashfaq, M., Dai, L., Xie, X., & Yu, C. P. (2017). Fate and mass balance of bisphenol analogues in wastewater treatment plants in Xiamen City, China. *Environmental Pollution*, *225*, 542–549. https://doi.org/10.1016/j.envpol.2017.03.018

Svoboda, P., Lindström, E. S., Ahmed Osman, O., & Langenheder, S. (2018). Dispersal timing determines the importance of priority effects in bacterial communities. *ISME Journal*, *12*(2), 644–646. https://doi.org/10.1038/ismej.2017.180

Thelusmond, J. R., Strathmann, T. J., & Cupples, A. M. (2019). Carbamazepine, triclocarban and triclosan biodegradation and the phylotypes and functional genes associated with xenobiotic degradation in four agricultural soils. *Science of the Total Environment*, 657, 1138-1149. https://doi.org/10.1016/j.scitotenv.2018.12.145

Tlili, A., Corcoll, N., Arrhenius, Å., Backhaus, T., Hollender, J., Creusot, N., Wagner, B., & Behra, R. (2020). Tolerance Patterns in Stream Biofilms Link Complex Chemical Pollution to Ecological Impacts. *Environmental Science and Technology*, *54*(17), 10745–10753. https://doi.org/10.1021/acs.est.0c02975

Vellend, M. (2010). Conceptual synthesis in community ecology. *Quarterly Review of*

Biology, *85*(2), 183–206. https://doi.org/10.1086/652373

Vila-Costa, M., Cerro-Gálvez, E., Martínez-Varela, A., Casas, G., & Dachs, J. (2020). Anthropogenic dissolved organic carbon and marine microbiomes. In *ISME Journal* (Vol. 14, Issue 10, pp. 2646–2648). Springer US. https://doi.org/10.1038/s41396-020-0712-5

Vila, J. C. C., Liu, Y. Y., & Sanchez, A. (2020). Dissimilarity–Overlap analysis of replicate enrichment communities. *ISME Journal*, *14*(10), 2505–2513. https://doi.org/10.1038/s41396-020-0702-7

Wang, J., Pan, F., Soininen, J., Heino, J., & Shen, J. (2016). Nutrient enrichment modifies temperature-biodiversity relationships in large-scale field experiments. *Nature Communications*, *7*, 1–9. https://doi.org/10.1038/ncomms13960

Wang, Z., Zhang, X. H., Huang, Y., & Wang, H. (2015). Comprehensive evaluation of pharmaceuticals and personal care products (PPCPs) in typical highly urbanized regions across China. *Environmental Pollution*, *204*, 223–232. https://doi.org/10.1016/j.envpol.2015.04.021

Wolff, D., Krah, D., Dötsch, A., Ghattas, A. K., Wick, A., & Ternes, T. A. (2018). Insights into the variability of microbial community composition and micropollutant degradation in diverse biological wastewater treatment systems. *Water Research*, *143*, 313–324. https://doi.org/10.1016/j.watres.2018.06.033

Wu, L., Yang, Y., Chen, S., Zhao, M., Zhu, Z., Yang, S., Qu, Y., Ma, Q., He, Z., Zhou, J., & He, Q. (2016). Long-term successional dynamics of microbial association networks in anaerobic digestion processes. *Water Research*, *104*, 1–10. https://doi.org/10.1016/j.watres.2016.07.072

Xie, J., Zhao, N., Zhang, Y., Hu, H., Zhao, M., & Jin, H. (2022). Occurrence and partitioning of bisphenol analogues, triclocarban, and triclosan in seawater and sediment from East China Sea. *Chemosphere*, 287, 132218. https://doi.org/10.1016/j.chemosphere.2021.132218

Yamazaki, E., Yamashita, N., Taniyasu, S., Lam, J., Lam, P. K. S., Moon, H. B., Jeong, Y., Kannan, P., Achyuthan, H., Munuswamy, N., & Kannan, K. (2015). Bisphenol A and other bisphenol analogues including BPS and BPF in surface water samples from Japan, China, Korea and India. *Ecotoxicology and Environmental Safety*, *122*, 565–572. https://doi.org/10.1016/j.ecoenv.2015.09.029

Yang, X., Chen, F., Meng, F., Xie, Y., Chen, H., Young, K., Luo, W., Ye, T., & Fu, W. (2013). Occurrence and fate of PPCPs and correlations with water quality parameters in urban riverine waters of the Pearl River Delta, South China. *Environmental Science and Pollution Research*, *20*(8), 5864–5875. https://doi.org/10.1007/s11356-013-1641-x

Zaborowska, M., Wyszkowska, J., & Borowik, A. (2020). Soil microbiome response to contamination with bisphenol A, bisphenol F and bisphenol S. International

Journal   of   Molecular   Sciences,   21(10),   3529.
https://doi.org/10.3390/ijms21103529

Zhang, W., Yin, K., & Chen, L. (2013). Bacteria-mediated bisphenol A degradation.
Applied   Microbiology   and   Biotechnology,   *97*(13),   5681–5689.
https://doi.org/10.1007/s00253-013-4949-z

Zhou, N. A., Lutovsky, A. C., Andaker, G. L., Gough, H. L., & Ferguson, J. F. (2013).
Cultivation and characterization of bacterial isolates capable of degrading
pharmaceutical and personal care products for improved removal in activated
sludge   wastewater   treatment.   *Biodegradation*,   24(6),   813-827.
https://doi.org/10.1007/s10532-013-9630-9

Zhou, J., & Ning, D. (2017). Stochastic Community Assembly: Does It Matter in
Microbial Ecology? *Microbiology and Molecular Biology Reviews*, *81*(4), 1–32.
https://doi.org/10.1128/mmbr.00002-17

Table 1. Number of OTUs assigned to the ecological categories opportunistic, sensitive,

and tolerant in the four micropollutant treatments: 'BPA' microcosms containing

bisphenol A, 'TCS' microcosms containing triclosan, 'MI' microcosms containing a

mixture of bisphenol A and triclosan, 'MII' microcosms containing a mixture of

bisphenol A, triclosan, bisphenol S and triclocarban.

| Treatment | Number of OTUs in each ecological category | | |
	Opportunistic	Sensitive	Tolerant
BPA	1	64	6
TCS	1	47	15
MI	3	51	12
MII	0	41	10

Figure 1. Experimental setup. In microcosm batch 1 (B1), 600 ml of the initial

inoculum was added to bottles containing the micropollutants. Starting from the B1, 60

mL (10%) of the inoculum was pipetted from each of the replicates and transferred to

fresh medium containing 540 mL of 0.22-μm-filtered water and freshly added

micropollutants every 5 days. This procedure was carried out in the seven batches of

microcosms from batch 1 (B1) through batch 7 (B7). Samples for bacterial community

characterization and micropollutant removal were collected prior to the next inoculation.

Treatments are color-coded: 'Control' is (without micropollutant addition, grey), 'BPA'

(microcosms added with bisphenol A, red), 'TCS' (microcosms added with triclosan,

blue), 'MI' (microcosms added with a mixture containing bisphenol A and triclosan,

purple), 'MII' (microcosms added with a mixture containing bisphenol A and triclosan,

bisphenol S and triclocarban, yellow).

Figure 2. Percent removal of bisphenol A (BPA), triclosan (TCS), bisphenol S (BPS),

and triclocarbon (TCC) in all microcosm batches for the single-micropollutant (BPA

and TCA) treatments and in the mixed-micropollutant (MI and MII) treatments.

Removal was defined as the decrease in the concentration of the chemical relative to

the total concentration within each batch of microcosms (differing by 5-day intervals).

'B1-B7' represent the number of microcosm batch that corresponds to every 5-day

inoculation.

Figure 3. Difference in the bacterial communities in all microcosms batches as determined by non-metric multidimensional scaling (NMDS) ordination. Different treatments are represented by different symbols, and microcosms batches are color-coded. Explanatory environmental variables fitting the ordination are shown with solid arrows. Explanatory values of the environmental variables to differences in the communities are displayed in Supplementary Table S1. Treatment IDs are: 'Initial' containing the starting bacterial communities before the experimental implementation; 'Con' microcosms containing no micropollutant additions and serve as the controls; 'BPA' microcosms containing bisphenol A; 'TCS' microcosms containing triclosan; 'MI' microcosms containing a mixture of bisphenol A and triclosan; 'MII' microcosms containing a mixture of bisphenol A, triclosan, bisphenol-S and triclocarban. 'B1-B7' represent the number of microcosm batch that corresponds to every 5-day inoculation. Explanatory values of the environmental variables to differences in the communities are presented in Supplementary Table S1.

Figure 4. Bacterial community dissimilarity in the micropollutant treatments vs. the

controls (i.e., 'Con.') over time (A) and the dynamic regularity within treatments (B).

Significant differences in the Bray-Curtis dissimilarity for each pairwise comparison of

the treatment and control (B1 microcosms vs. later microcosms in (A) were determined

in a Wilcoxon test: ***, $P < 0.001$; **, $0.001 < P < 0.01$; *, $0.01 < P < 0.05$. The

dissimilarity-overlap analysis within each treatment is shown in (B); out of total 210

sample pairs, only the sample pairs that had median overlap above cutoff = 0.913 were

shown in the plots. The gray lines indicate the linear regression with the slope (m), and

the P value (the fraction of bootstrap realizations in which the slope was negative).

Gray shaded areas indicate the 95% confidence intervals of the linear regression.

Significance level was determined at $P < 0.05$. 'B1-B7' represent the number of

microcosm batch that corresponds to every 5-day inoculation.

Figure 5. Changes in the relative abundances (rows of heatmap) of the OTUs assigned to the ecological categories tolerant and sensitive across the three growth phases (columns of heatmap) (A) and their phylogenetic distribution (B). In (A), color gradients of heatmap represent the relative abundances of individual by column with warm colors (toward red) indicating high abundance and cold colors (toward blue indicating low abundance within the sample. Column labels are batch IDs and the grouping of growth phases, and row labels the OTU IDs. Tolerant and sensitive OTUs

939   were denoted in green and blue colors, respectively. In (B), the innermost circle

940   indicates the taxonomy (phylum/class) of the OTUs. The outer circles indicate the

941   ecological group of OTUs in each treatment group (BPA, TCS, MI, MII from inside to

942   outside, respectively).

Figure 6. Relative proportion of the ecological processes (homogeneous selection, variable selection dispersal limitation, and ecological drift) contributing to the communities assembled in the micropollutant treatments (BPA, TCS, MI or MII) vs. the controls (i.e., 'Con.'). Each of the three phases (corresponding to the early, mid, and late stages of the press disturbance) was classified according to the similarity in the average 16S rRNA gene copy numbers as an indicator of community growth progression. Three growth phrases across microcosm batches were: phase 1 (B1–B2), phase 2 (B3–B4), and phase 3 (B5–B7), as shown in Fig. S2G.

Figure 7 Conceptual model of microbial community assembly processes following a press disturbance. Considering the time continuity of the batch inoculation, the three phases of community growth progression (1, 2, and 3) also correspond to the early, mid, and late stages of the 35-day disturbance. This conceptual model illustrates when and how stochastic/deterministic processes contribute to the dynamics of microbial communities repeatedly exposed to micropollutants. Changes in community trajectory over time in the TCS or the MII treatment are indicated by the black line, MI treatment by the red line, and the BPA treatment by the purple line. Overall, stochastic and deterministic processes act in concert during community dynamics: stochastic assembly dominates at the early stage of a press disturbance/community development, and deterministic assembly at the later stage. Major community assembly mechanisms are delineated for each phase. Community succession accelerates at phase 2 such that multiple equilibria occur at phase 3. Note that community succession and multiple equilibria are not exclusive, but community turnover is high towards the end of the disturbance.