Recurrent micropollutant exposure leads to divergent degradation abilities and community succession in a freshwater microbiome

Running title: Ecologically disturbing effects of trace micropollutants on a freshwater microbiome

Dandan Izabel-Shen[1#], Shuang Li[2,3,4#], Tingwei Luo[5], Jianjun Wang[6], Yan Li[2,3], Qian Sun[2,3], Chang-Ping Yu[2,7], Anyi Hu[2,3*]

1. Department of Ecology, Environment and Plant Sciences, Stockholm University, 106 91, Stockholm, Sweden
2. CAS Key Laboratory of Urban Pollutant Conversion, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China
3. University of Chinese Academy of Sciences, Beijing 100049, China
4. Department of Environmental Microbiology, UFZ, Helmholtz Centre for Environmental Research, Leipzig, Germany
5. Institute of Marine Microbes and Ecospheres, State Key Laboratory of Marine Environmental Science, Xiamen University, Xiamen, China
6. State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing, 210008, China
7. Graduate Institute of Environmental Engineering, National Taiwan University, Taipei 106, Taiwan

#The authors contributed equally to this work.

*Correspondence to:

Dr. Anyi Hu, ayhu@iue.ac.cn

CAS Key Laboratory of Urban Pollutant Conversion, Institute of Urban Environment,

Chinese Academy of Sciences, Xiamen 361021, China, Telephone & Fax: 86-592-

6190582.

Keywords: Micropollutant removal; preexposure history; community turnover;

stochastic assembly; deterministic assembly;

Abstract

The high-volume release of micropollutants into natural surface waters has raised

concern due to their environmental accumulation. Persisting micropollutants can

impact multiple generations of organisms via press disturbances, but their microbially-

mediated degradation and the influence on community assembly remain understudied.

Here, freshwater microbial communities in microcosms were treated with four common

micropollutants, alone or in combination, and then transferred in stages to fresh medium

containing the same level of micropollutants to mimic the recurrent exposure of

microbes under environmentally relevant conditions. Our results showed that the

degradation of micropollutants was closely linked to the community succession, and

that recurrent exposure to micropollutants enhanced the degradation capacity. The

partitioning analysis of ecological processes revealed that community assembly was

dominated by stochastic processes during early exposure, via random community

changes, and by deterministic processes later in the exposure. Analyzing individual taxa

abundances over time revealed two distinct bacterial responses, in which a larger proportion of sensitive than tolerant taxa was present in the disturbed communities, and the abundances of the most sensitive taxa were significantly associated with micropollutant degradation. This study clearly showed that microbial communities are generally vulnerable to persisting micropollutants in aquatic environments, which has important implications for pollution management, especially regarding microbial dynamics and ecosystem functioning in micropollutant removal.

Introduction

The rapid exploitation and widespread use of emerging organic chemicals have resulted in environmental damage and human health issues during the last several decades [1, 2]. These organic chemicals include pharmaceuticals and personal care products, hormones, and industrial chemicals, and are typically classified as micropollutants, because of their occurrence at a low to very low concentration of μg/L and ng/L in the receiving environments [3, 4]. In aquatic systems, for instance, the majority of micropollutants originate from direct and indirect inputs of organic contaminants that are released via the effluents of wastewater treatments plants [1, 5–7]. The micropollutants gradually accumulate at sites of their release and reach concentrations high enough to exert disruptive effects on aquatic organisms such as fish, invertebrates, and microbes [8–13]. Many studies have assessed the eco-toxicity of micropollutants to humans and animals [e.g., 14–17]. However, little is known about the mechanisms by which micropollutants impact the turnover of aquatic microbial communities [6, 18, 19], especially those of freshwater bacteria [20, 21], and the role of microbes on trace micropollutant degradation.

The responses of microbial assemblages to micropollutants can be classified as: subsidy responses, in which, via biodegradation, the microorganisms are able to utilize the micropollutant compounds as carbon sources, and stress responses, in which the growth and survival of microorganisms are inhibited [22, 23]. Among the compounds eliciting subsidy responses are bisphenol A (BPA) and its substituted derivatives (e.g., bisphenol S, BPS) [24], as their relatively simple chemical structure and low toxicity allow both to be utilized as carbon sources by microbial isolates [23, 25]. Stress responses are elicited by pharmaceutical and personal care micropollutants such as triclosan (TCS) and its substituted derivatives (e.g., triclocarban, TCC) [26], because

both inhibit the growth of several microbial taxa, consistent with their resemblance to antimicrobials [22, 27]. Both BPA and TCS in the field strongly influence microbial diversity, especially that of the central species and module communities comprising the microbial co-occurrence network [28].

Microbial responses to chemical pollutants can be modified by the exposure legacy of the community. For example, microbial communities with a history of containment exposure may differ from contaminant-naïve communities in the rate and extent of chemical degradation [29–31]. They may also differ in their plasticity [18] upon subsequent exposure to similar contaminants. However, the ecological drivers of community succession following repeated micropollutant exposure are not well-understood. Studies of community dynamics over multiple generations and a time period relevant for community stability could provide insights into the ecological traits that determine taxon sensitivity and tolerance, and therefore the ability of microbial community members to degrade micropollutants.

Press disturbances in natural environments can impact multiple generations of organisms [32] by altering the affected ecosystems beyond the possibility of recovery or by causing a shift to a community with a new stable state [33]. Following a stress event such as recurrent micropollutant exposure, communities rely on reassembly from the local seedbanks such that succession eventually results in a new steady state. During the early stages of community dynamics, no-longer-dormant local species undergo a secondary succession characterized by new abiotic and biotic interactions. The emergence of and resuscitation from dormancy is considered as stochastic processes that lead to divergent communities [34–36]. When abiotic and biotic conditions allow a stabilization of the initial communities, community formation at a later phase of a press disturbance is expected to converge, through the succession of microbial species

[37]. In addition, the plasticity of microbial taxa in response to recurrent micropollutant exposure has been shown to increase over multiple generations [18], with later-assembled communities shifting to a new stable state that differs from the one before the exposure.

In this study, a press disturbance was simulated in a microcosm experiment in which a reservoir bacterial community was transferred into fresh medium containing trace concentrations of typical micropollutants that were refreshed with recurrent micropollutant exposure every 5 days. Several different micropollutants are typically present simultaneously within aquatic environments, but whether their interactions lead to effects on microbial communities that differ from those of single micropollutants is unclear [12]. Thus, we also experimentally manipulated single and multiple micropollutants to determine their impacts on taxon coexistence, the regularity of community assembly, and micropollutant degradation ability. We hypothesized, that because BPA elicits a subsidy response and TCS a stress response, the removal rate of BPA would be higher than that of TCS, with larger differences in the removal rates by microbial communities exposed to these micropollutants individually rather than in combination with other micropollutants. Accordingly, the community composition assembled at the initial and late phases of exposure would differ substantially, with convergent dynamics more apparent towards the end of the press disturbance. Finally, we predicted that community dynamics would greatly differ following exposure to BPA vs. TCS whereas the dynamics of communities exposed to a mixture of micropollutants would resemble those of either the BPA- or the TCS-exposed assembly, depending on the number and abundance of the shared taxa.

Materials and Methods

Water collection

We selected the bacterial species pools from Shidou Reservoir, which is one of largest sources of drinking water in Xiamen, China (Hu et al., 2018) and is therefore under governmental protection from pollution originating from household drainages and a nearby dam (Fig. S1). Unlike other contaminated aquatic environments [22, 38, 39], the concentrations of BPA, BPS, TCS, and TCC in Shidou Reservoir (China) are very low (< 1 ng/L), such that the reservoir's waters provide an ideal system to investigate the influence of persistent micropollutants on an aquatic microbiome. Surface water (100 L) from Shidou Reservoir (24˚41'39"N, 118˚0'37"E) was collected at 0.5 m depth on June 1, 2017 (Fig. S1). Temperature, pH, conductivity, dissolved oxygen (DO), and dissolved oxygen saturation, measured *in situ* using a HACH HQ40d multi-parameter meter (HACH, Loveland, CO, USA), were 29.0°C, 8.90, 54.4 μs/cm, 8.84 mg/L and 116.7%, respectively. The concentrations of BPA, BPS, TCS, and TCC in the reservoir water were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS) (ABI 3200QTRAP, Framingham, MA, USA) with solid-phase extraction and all were < 1 ng/L at the time of sampling (details of the measurements are provided below). The collected water was transported from the field to the laboratory within 2 h after sampling.

Experimental setup

BPA, BPS, TCS, and TCC were dissolved in acetone at concentrations of 800, 800, 400, and 400 μg/L, respectively, and stored as stocks in glass bottles (HPLC grade). The micropollutants were previously shown to be chemically stable when dissolved in an organic solvent [39, 40]. From the 100 L of collected water, 15 L were filtered through

a 1.2-µm polycarbonate filter (Millipore, Ireland), to minimize top-down regulation by planktonic grazers, and then used as the initial inoculum. The remaining unfiltered water was filtered through a 0.22-µm polycarbonate filter (Millipore, Cork, Ireland) to remove microorganisms and then stored in the dark for later use in the preparation of incubation medium. Two treatments consisting of single micropollutants (BPA and TCS) and two treatments consisting of micropollutant mixtures (MI = BPA + TCS) and (MII = BPA + BPS + TCS + TCC) were established.

Time-serial microcosms comprising seven inoculation batches from batch 1 (named hereafter 'B1') through batch 7 ('B7') were established to investigate the microbial community response to a press disturbance of micropollutants resulting from a single short-term event but maintained for 35 days (Fig. 1). Prior to the experimental initiation, the caps of the bottles containing the micropollutants were loosened for > 12 h to volatilize the acetone to minimize its influence on the microbial responses. In microcosm B1, 600 ml of the initial inoculum was added to bottles that already contained the micropollutants and nutrients, such that the final concentration of each micropollutant in the treatments was as follows: BPA: 1 µg/L BPA, TCS: 1 µg/L TCS, MI: 1 µg/L BPA + 1 µg/L TCS, and MII: 1 µg/L BPA + 0.1 µg/L BPS + 1 µg/L TCS + 0.1 µg/L TCC. The final concentration of each micropollutant in the microcosms closely represented their ambient environmental conditions as reported in literature [22, 41]. Further, given the low nutrient concentrations [42], we adjusted the concentrations of nitrogen and phosphorus in the filtered water to 128.6 and 8.04 µM to promote microbial growth by the addition of KNO_3 and KH_2PO_4, respectively [43]. After 5 days of incubation, 60 mL of the B1 inoculum was pipetted from each of the replicates and transferred to a sterile bottle containing 540 mL of 0.22-µm-filtered water and freshly added micropollutants. This newly inoculated microcosm served as batch 2 (named

hereafter 'B2'). The micropollutant treatments and controls were amended with 128.6

µM N and 8.04 µM P after each inoculation to maintain similar amounts of these

nutrients throughout the experiment. The same inoculation procedure was applied at

each 5-day interval to initiate microcosms B3–B7, such that the experimental period

was 35 days (Fig. 1). Microbial cells were transferred between pairwise replicates, with

the remainder used in the downstream analyses described below. An additional set of

experimental bottles without micropollutant addition served as the controls. In total,

there were five treatments, each with three replicates, for a total of 15 microcosm

replicates per batch. All microcosms, i.e., 105 experimental bottles during the 35 days,

were incubated in the dark at 28°C with shaking at 100 rpm.

Additional experiments were conducted to determine (i) micropollutant removal

rates in the absence of microorganisms, to rule out the gradual degradation of the

micropollutants in the treatments due to their chemical instability, and (ii) how the

ability of the microorganisms to transform micropollutants differed depending on

whether they have previous exposures or not. Details on the methods and the results of

these two experiments are provided in the Supplemental information.

Micropollutant analysis

For each microcosm batch, ~20 mL of the inoculum was removed from each bottle at

0, 2 and 5 days for micropollutant analysis. The four micropollutants BPA, BPS, TCS,

and TCC were concentrated and purified by a 12-port Visiprep-DL SPE vacuum

manifold (Supelco, Bellefonte, PA, USA) with Oasis HLB cartridges (500 mg, 6 mL;

Waters, Millford, MA, USA), and then were measured using LC-MS/MS (ABI

3200QTRAP, Framingham, MA, USA), as described in our previous work (Sun et al.

2017). Chromatographic separation was performed by using a Kinetex C18 column

(100 mm × 4.6 mm, 2.6 μm; Phenomenex, Torrance, CA, USA) from Shimadzu LC

system (Shimadzu, Japan).

DNA extraction and 16S rDNA amplicon high-throughput sequencing

The remaining inoculum (~470 mL) of each replicate was filtered onto 0.22-μm

polycarbonate filters (Millipore, Ireland), which were stored at −80°C until used for

DNA extraction. DNA from the 105 samples was extracted using the FastDNA for soil

kit (MP Biomedical, Santa Ana, CA, USA) according to the manufacturer's protocol,

except that the beating speed was set to 6.0 m/s for 80 s. The universal primer pair 515F

(5'-GTG YCA GCM GCC GCG GTA-3') and 907R (5'-CCG YCA ATT YMT TTR

AGT TT-3') was used to amplify the V4-V5 region of the bacterial 16S rRNA genes.

The PCR products were purified, mixed in equal amounts, and sequenced on an

Illumina MiSeq platform with the 2 × 300 bp paired-end protocol at the Majorbio Bio-

Pharm Technology Co., Ltd. (Shanghai, China).

Sequence analysis

The raw sequencing data were processed using the LotuS pipeline [44] with the

following criteria: i) average sequence quality > 27, ii) sequence length > 170 bp, iii)

no ambiguous bases, iv) no mismatches with the barcodes or primers, and v)

homopolymer length < 8 bp. Operational taxonomic units (OTUs) were clustered at a

97% similarity using UPARSE [45]. A representative sequence of each OTU was

taxonomically quantified using the RDP classifier with the Silva database v132 [46], at

a bootstrap cutoff of 80%. All samples were rarified to 36, 000 sequence reads (the

smallest library size). The subsampled OTU table contained 1375 OTUs. The α- and β-

diversity indices of bacterial communities were calculated from the rarified dataset

220 using QIIME v1.9.1 [47] with the script "core_diversity.py".

Inferring the community growth phase from the16S rRNA gene copy number

222 The 16S rRNA gene copy number among prokaryotes within a defined period indicates

223 the number of individuals within a community and reflects the growth phase of that

224 community [48]. In this study, the 16S rRNA gene copy number of each OTU found in

225 the samples was estimated using the rrnDB database v5.5 [49] and the weighted mean

226 rRNA gene copy number of all OTUs found in each sample was then calculated as

227 described previously [50]. The abundance-weighted average gene copy number of a

228 community was plotted against the sampling time for each treatment. The similar

229 overarching pattern of the average rRNA gene copy numbers of the bacterial

230 communities in all microcosms over time (Figs. S2A-F) indicated the similar growth

231 trajectories of those communities. In addition, a hierarchical cluster analysis with

232 complete clustering criterion was used to cluster the similarity of average gene copy

233 numbers from all micropollutant-treated microcosms belonging to the same batch type,

234 with an exclusion of the controls (Fig. S2G). Three growth progression phrases across

235 all microcosm batches were identified based on similar average gene copy numbers and

236 the time continuity of the inoculation (Fig. S2G): phase 1 (B1–B2), phase 2 (B3–B4),

237 and phase 3 (B5–B7).

Relative contribution of ecological processes

239 According to a quantitative ecological framework [51, 52], community assembly can

240 be summarized by five major ecological processes: i) homogenizing selection for a

convergent composition between communities, ii) variable selection for a divergent

composition between communities, iii) dispersal limitation for a divergent composition

between communities due to limited dispersal events, iv) homogenizing dispersal for a

convergent composition between communities due to unlimited dispersal, and v)

ecological drift (referred to "undominated processes" in [52]). The phylogenetic

turnover among the communities was quantified by the weighted β-mean nearest taxon

distance (βMNTD). The βNTI was calculated as the difference between the observed

βMNTD and the mean of the null distribution (999 randomizations). The criteria

determining the fraction of each process were as follows: i) homogenizing and variable

selection: higher (> 2) and lower (< -2) βNTI, respectively; ii) dispersal limitation and

homogenizing dispersal: non-significant $|βNTI|$ (< 2) but $RC_{bray} > 0.95$ and < -0.95,

respectively. iii) ecological drift: non-significant $|βNTI|$ value (< 2) and non-significant

$|RC_{bray}|$ value (< 0.95) [52]. To infer the relative contributions of the different

community assembly processes to community turnover over time, for each pair

consisting of a micropollutant treatment and its control the βNTI and RC_{bray} values were

calculated separately for each of the three growth phrases. By comparing the

community assembly process in the treatment vs. the controls, we were able to

minimize confounding effects that may have obscured community variation and thus

better predict the treatment effect. The same analyses were applied to each pair of

communities from the different treatments. The relative contribution was expressed as

the portion of pairwise sample assembly that could be attributed to each ecological

process.

Ecological grouping of microbial responses to micropollutants

The responses of individual OTUs to micropollutant addition were identified by

analyzing the differences in the relative abundance of each OTU during the three growth phases for each treatment, using the 'DESeq2' R package v 1.26.0 [53]. Significant values were corrected for multiple tests using the Benjamini-Hochberg correction with an adjusted α value of 0.05. The responses were categorized as: i) sensitive, when relative OTU abundances were significantly higher in phase 1 than in phases 2 and 3; ii) opportunistic, when relative abundances were significantly higher in phase 2 than in phases 1 and 3; iii) tolerant, when relative abundances were significantly higher in phase 3 than in phases 1 and 2. The ecological categories represented a group of bacterial members whose responsiveness to the experimental conditions could be documented collectively. In addition, to visualize the phylogenetic relationships of the OTUs assigned to the ecological groups, representative sequences of those OTUs were used to construct a phylogenetic tree using iTOL v5 [54]. A Spearman correlation test was used to explore the relationships between the OTUs with ecological strategies and the micropollutant removal rates. The P-value was corrected using the Benjamini–Hochberg method.

Micropollutant turnover and community analysis

A Kruskal-Wallis test was used to analyze the difference in the removal rates between the single micropollutant treatments (BPA vs. TCS) and between any two micropollutants for the mixture treatments (either MI or MII).

Differences in the bacterial communities in all microcosms were explored using non-metric multidimensional scaling (NMDS) based on the Bray-Curtis dissimilarity metric and by fitting removal rates of micropollutants to the ordination. This allowed us to assess whether the degradation of micropollutants was tied to the microbial

succession. A permutational multivariate analysis of variance (PERMANOVA) [55] was conducted to test the effects of micropollutants, inoculation time (as indicated by the microcosm batch), and their interactive effects on the taxonomic composition of the bacterial communities. A pairwise PERMANOVA was performed to test for significant differences in community composition between each treatment pair. The multivariate analyses were performed using the R package 'vegan'[56]. Whether communities with the shared taxa exhibit similar dynamics was assessed in a dissimilarity-overlap analysis [57]. Briefly, the relative abundance of each OTU in each community was calculated to identify sets of OTUs within each pair of communities (i.e., within a treatment over time and between treatments). In the calculation of dissimilarity, only the shared OTUs were considered, and the relative abundance of shared taxa was transformed to add up to 1. The dissimilarity between renormalized vectors X and Y can be calculated using root Jensen–Shannon divergence. Based on the dissimilarity-overlap analysis, 15 pairwise comparisons were obtained in this study. This method and the R script for this analysis were modified from [58].

To better understand community turnover over time, we quantified the compositional variation between the micropollutant-treated and non-treated communities (i.e., control) at each time point using the Bray-Curtis dissimilarity metric. The rate of community turnover over time was evaluated with a linear regression model (see the Supplemental methods for details). A Wilcoxon test was then applied to test for significant differences in community turnover between microcosm B1 (day 5) and the

other microcosm batches from later time points (B2–B7), using the 'ggsignif' package

[59]. The data obtained in this study were visualized using the R package 'ggplot2'.

Results

Transformation of trace micropollutants

In the microcosms treated with single micropollutant, a higher proportion of BPA (96.84

± 7.90%) than TCS (21.83 ± 24.11%) was removed (Kruskal-Wallis test, $P < 0.05$; Fig.

2A) whereas the removal of the two micropollutants from the MI microcosms did not

significantly differ ($P > 0.05$) (17.86 ± 15.17% for BPA and 25.11 ± 18.88 % for TCS;

Fig. 2B). In the MII microcosms, the percentage of removed BPA (65.83 ± 6.54%) was

highest ($P < 0.05$), followed by the TCS (13.76 ± 16.76%), BPS (5.93 ± 8.97%), and

TCC (11.97 ± 14.08%), but the differences were not significant (Fig. 2C).

Community succession following micropollutant exposure

The bacterial communities in the microcosms became taxonomically distinct both over

time and by micropollutant type (Fig. S3). The variation in the community composition

among all microcosm batches correlated with differences in the removal rates of BPA,

with higher association in its treatments alone, but was also correlated with TCA, with

higher association in TCA single and mixed treatments (Fig. S3). The inoculation time

(PERMANOVA, $R^2 = 0.149$), micropollutant addition ($R^2 = 0.11$), and their interaction

($R^2 = 0.055$) significantly influenced the community composition (all $P < 0.001$) (Table

S1). Irrespective of the inoculation time, the composition of the TCS, MI, and MII

communities differed significantly from that of the controls (pairwise PERMANOVA,

$P < 0.01$; Table S1) but not from each other. Conversely, the BPA communities

significantly differed from the other three treatments but not from the controls.

The dissimilarity in the micropollutant-treated vs. control communities significantly increased over time, as each inoculation was replenished with micropollutants (linear regression, all $P < 0.001$, Fig. S4). Community dissimilarity was more pronounced at late (B6 or B7) than at early (B1 or B2) microcosm batches. Furthermore, the communities of the first inoculation (B1) impacted the response of some communities inoculated at later time points (Fig. 3A). The impact of the B1 community composition on later communities persisted across B2–B6 in the BPA-treated microcosms and B2–B5 in the MII-treated microcosms, as evidenced by their community resemblances (Wilcoxon test, $P > 0.05$; Fig. 3A). In both BPA and MII communities, the correlation between dissimilarity and overlap (i.e., the number of shared OTUs between the community pair) was negative ($P < 0.05$ Fig. 3B). This result was not unexpected since the dispersal of microbial cells from one condition to another similar environment may drive community dynamics towards convergence [60, 61]. However, in the TCS and MI treatments, an impact of the initial communities on later ones occurred only in B2, which is supported by the fact that the B3–B7 communities were more dissimilar to B1 ($P < 0.05$; Fig. 3A). The negative correlation of the dissimilarity and overlap analysis was nonsignificant for the TCS and MI communities such that the high overlap did not result in a low dissimilarity (Fig. 3B). This suggests that there were differences in the dynamics of the shared taxa in those TCS and MI communities over time compared to the communities in the other treatments.

Analyses of each pair of communities from the different treatments did not reveal a significant correlation between dissimilarity and overlap in the community pairs of BPA vs. TCS, BPA vs. MI, and TCS vs. MII (Wilcoxon test, $P > 0.05$, Fig. 4). However, similar population dynamics were determined for the communities assembled in the BPA vs. MII and TCS vs. MI pairs, as indicated by the significant negative correlation

357 between high overlap and low dissimilarity ($P < 0.05$, Fig. 4).

OTUs with distinct responses to micropollutant addition

In all treatments, a greater number of sensitive OTUs (41–64), a moderate number of

tolerant OTUs (6–15), and fewer opportunistic OTUs (0–3) were identified across

phases 1 (B1−B2), 2 (B3−B4), and 3(B5−B7) (Table 1 and Fig. 5). Regardless of the

micropollutant type, members of Actinobacteria, Planctomycetes, and Verrucomicrobia

primarily consisted of sensitive OTUs (Table 1; Fig. 5). Those of Alphaproteobacteria,

Gammaproteobacteria and Bacteroidetes contained more sensitive than tolerant OTUs.

The majority of the tolerant OTUs were unique to one treatment, with only two OTUs

tolerating micropollutant addition in all treatments. By contrast, many sensitive OTUs

(51) were shared between treatments, and at least between two treatments (Table S3).

Among the overlapping, 20 OTUs found in all treatments, most were members of

Actinobacteria (Table S3; Fig. 5).

Interestingly, the relative abundances of 17 of the 64 sensitive OTUs in the BPA

microcosms correlated negatively with the removal rate of BPA (Spearman correlation,

$P < 0.01$, Table S2), whereas for the two tolerant OTUs a positive correlation between

the two parameters was determined (Table S2). Among the 47 sensitive OTUs in the

TCS microcosms, 28 correlated positively with the removal rate of TCS (Table S2).

Partition of ecological processes on community variation

376 Overall, homogenizing selection played a minor role in influencing community

377 variation across phases for each community pair (micropollutant-treated and control),

378 as it accounted for less than 8.33% of the ecological processes (Fig. 6). Instead, variable

379 selection (13.89–61.73%) and ecological drift (30.86–80.56%) dominated the

ecological processes across the three phases. These two processes exhibited compensatory contributions, showing increasing variable selection but decreasing ecological drift along the phases (Fig. 6). In the case of dispersal limitation, its relative proportion was greater at phase 1 than the other phases for all treatments, especially BPA and MII.

The relative proportion of ecological processes attributable to dispersal limitation decreased across all three phases and was consistent for all between-treatment pairwise comparisons. The three other processes exhibited a compensatory relationship (Fig. S5). Specifically, variable selection and ecological drift prevailed at phases 2 and 3 and replaced one another in comparisons of BPA and each of the other three treatments. However, in comparisons of TCS vs MI or MII, ecological drift was over-represented in all phases (> 50%), while variable selection accounted for a small fraction of the ecological processes (Fig. S5). The trend for MI vs. MII was similar whereas there was no clear trend for the process of homogenizing selection across any of the phases in all comparisons.

Discussion

In the present study, the effects of persistent micropollutants on microbial community dynamics were experimentally investigated to gain insights into community assembly mechanisms. Our first hypothesis, that differences in the removal rates of BPA vs. TCS in environments exposed to single vs. mixed micropollutants was not supported by the results obtained for the MI and MII treatments. The energy required by the microbial communities to counteract the opposing responses to BPA and TCS in the MI treatment [23, 25] likely accounted for the profound inhibition of BPA degradation (Fig. 2B). In MII, a significantly larger fraction of BPA than of the three other micropollutants was

removed, consistent with the synergistic effects of BPS and TCC on the microbially

mediated degradation of BPA. We further demonstrated that degradation was a

biological rather than a chemical process and that bacterial taxa were responsible for

those transformations, which are supported by the chemical stability and negligible

transformation of the four micropollutants in the bacteria-free microcosms (Fig. S6).

We also found that pre-exposure to micropollutants can enhance the degradation

capacity and dissipation rate by microbes later confronted with those or similar

chemicals, which is consistent with previous literature [18, 31]. This was supported by

the fact that the taxa pre-exposed to low concentrations of BPA and TCS fully degraded

higher concentrations of those micropollutants during subsequent exposures and

degradation occurred at a faster rate (Fig. S7), but only when BPA and TCS were

provided alone and not in combination. Our results therefore revealed that microbial

communities can transform each of the four micropollutants surveyed, which is

especially true for BPA.

The larger proportion of sensitive than tolerant taxa in all treatments suggested that

the microbial community was not robust to the persistence of micropollutants in its

surroundings. Despite the strong effect of TCS on microbial dynamics in the MI and

MII treatments, in the latter BPA was removed at a much higher rate than any of the

other micropollutants. It is important to consider that the compositional response to a

disturbance is often slower than the functional response [62]. Furthermore, most of the

sensitive taxa tended to correlate negatively or positively with the removal of BPA or

TCS, respectively (Table S2). Those taxa may have affected micropollutant degradation

either directly or indirectly, such as by providing enzymatic repertoires to other

community members that enabled degradation, including interspecies interactions [63,

64]. Actinobacteria, Alphaproteobacteria, and Bacteroidetes made up the majority of

the sensitive taxa; their associations with TCS transformation were previously reported

[65, 66].

There was less support for the second hypothesis, that communities assembled at

the early stage of micropollutant exposure would be compositionally distinct from those

that formed at the later stage, when community dynamics converge. Our findings

generally corroborated previous studies showing that the persistence of micropollutants

even at low concentrations can influence microbial community membership and

abundance over time [28, 31, 67]. Another study found that microbial communities in

identical habitats share a large number of taxa and exhibit convergent abundance

dynamics over time [58]. This was the case for the communities assembled in the BPA

and MII treatments, as indicated by the significantly negative dissimilarity-overlap

relationships (Fig. 3B). The dynamics of the communities within those environments

were driven by a set of shared taxa that interacted similarly with both the respective

environments and other community members. However, the less pronounced negative

dissimilarity-overlap relationships in the case of the TCS and MI treatments indicated

that those communities had not reached equilibrium or were characterized by multiple

stable states [57]. Within those treatments, the microbial taxa shared among different

inoculation batches interacted differently with the same environment over time. Hence,

microbial communities assembled in identical habitats can perceive those habitats

differently such that their feedback on the surroundings and other, coexisting members

will differ as well, as noted elsewhere [58, 68].

The third hypothesis, that the dynamics of microbial communities exposed to

mixed micropollutants would resemble those of the single micropollutant treatments,

was supported in the case of TCS vs. MI and MII. Community dynamics in those

treatments showed a consistent negative dissimilarity-overlap correlation, suggesting

that TCS overpowered the other micropollutants in the mixed treatments. This explanation was supported by the significant correlation between the changes in 16S rRNA gene copy numbers (Fig. S2) and the resemblance of the communities in the TCS, MI, or MII treatments (Fig. S3).

Despite evidence of regularity in the community dynamics across treatments, as revealed by the dissimilarity-overlap analysis, there was a gradual transition from stochastic to deterministic processes towards the later stage of micropollutant exposure. Thus, in all treatments ecological drift played a major role during phases 1 and 2 of community growth progression (Fig. 6). This can be explained by the strong ecological drift and thus the random changes in abundances within microbial communities confined to microcosms, where dispersal is limited and locally dormant cells are awakened to colonize empty niches in the environments upon disturbances [37, 69, 70]. However, the increasing contribution of variable selection to community assembly over time suggested that the high phylogenetic turnover of the communities was driven by heterogenous abiotic and biotic conditions [52, 71]. The increasing effects of the micropollutants on the communities over time led to higher species selection and an elevated impact on the interactions among those species within the community, such that the drivers of community assembly no longer resembled those of the control and resulted in a high rate of community turnover (Fig. 3A; Fig. S4). Our findings suggest that the repeated introduction of a micropollutant enhances succession in microbial communities [72].

We also considered the potential role of priority effects on community assembly, independent of the ecological process model used in our study. Priority effects emerge when early colonizers exhaust or modify available resources, thus influencing the establishment success of later colonizers [73]. A strong priority effect in the case of TCS and MI was evidenced by the significant difference in the composition of the initially inoculated vs. any of the later inoculated communities (Fig. 3A). Hence, once they were re-inoculated, taxa that had thrived in the B1 batch modified the environments of later batches, such that some taxa (presumably activated from a seed bank, given that no source of new taxa was externally introduced into the systems) were less or more favorable in later batches. However, for the BPA and MII treatments, the effects occurred at later stage of the press disturbance, suggesting the increasing strength of priority effects with increasing establishment time, as noted elsewhere [74].

Overall, although our microcosms represent an ideal conditions which is removed from *in situ* conditions, our experimental findings are valuable in at least three aspects, that is : i) the prediction of microbial responses to persistent micropollutants, ii) the harnessing of ecological principles and microbiomes in pollution management [75, 76, 77], and (iii) the development of a conceptual model to be used to predict when and how stochastic and deterministic processes contribute to community assembly after the exposure of persistent micropollutants. Our results suggest that strong ecological drift drives random community changes at the early stage of the exposure. Prolonged ecological drift, variable selection resulting from the effects of micropollutants, and the abiotic and biotic conditions experienced differently by different microbes then

determine community succession at the mid-stage of the exposure. Additionally, the high environmental heterogeneity together with the low densities of the communities permitted the emergence of priority effects, in which the early establishment of taxa in micropollutant-containing environments allowed a faster community succession. This was supported by the decrease in the average gene copy number at the intermediate phase (Fig. S2), when community assembly was driven by various ecological drivers [78]. Finally, strong variable selection accompanied by increased priority effects likely set the communities on distinct trajectories, with multiple equilibria at the late stage of the exposure. In fact, more advanced community succession and multiple equilibria characterized the end of the press disturbance, and both played a larger role in the TCS and MII (and to a lesser extent in the BPA) communities than in the MI community (Fig. 7).

In summary, our experimental design and ecological approaches showed that, while microbial communities are generally vulnerable to micropollutant persistence, pre-exposure to micropollutants may enhance the microbial degradation capacity. This work can aid in microbe-pollution management by providing insights into micropollutant removal as well as microbial assembly mechanisms and dynamics. From the perspectives of microbiome engineering, this study demonstrates the importance of the microbiome's preexposure history in the design of long-term experiments that include artificial selection to direct the evolution [76, 79] of a target microbiome highly optimized for pollutant removal in an aquatic environment.

Data availability

The sequencing data were deposited in the NCBI SRA database with the Bioproject number PRJNA497759. Our R scripts for statistics, data visualization and computing notes are available on a zenodo archive (https://doi.org/10.5281/zenodo.5242686) of the Github repository (https://github.com/AnyiHu/Impacts-of-micropollutants-on-freshwater-microbiome).

Acknowledgements

We thank Dr. Linwei Wu for her assistance in calculating the copy number of 16S rRNA genes, and Dr. Hongjie Wang for his assistance in the molecular experiments. This work was supported by the National Natural Science Foundation of China (31870475, U1805244 and 31470539), the International Partnership Program of Chinese Academy of Sciences (132C35KYSB20200012), and the Natural Science Foundation of Fujian, China (2019I0030).

Author contributions

AH and CPY conceived and designed the study. LS and TL performed the experiments and measurement analyses. YL and QS provided the assistance for the LC-MS/MS analysis. DIS, LS, JW and AH performed the data analyses and wrote the manuscript. All authors discussed the results and commented on the manuscript.

Competing Interests

The authors declare no competing interests.

References

1. Schwarzenbach RP, Escher BI, Fenner K, Hofstetter TB, Johnson CA, von Gunten U, et al. The Challenge of Micropollutants in Aquatic Systems. Science. 2006; 313:1072–1077.

2. Deblonde T, Cossu-Leguille C, Hartemann P. Emerging pollutants in wastewater: A review of the literature. Int J Hyg Envir Heal. 2011; 214:442–448.

3. Luo YL, Guo WS, Ngo HH, Nghiem LD, Hai FI, Zhang J, et al. A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. Sci Total Environ. 2014; 473:619–641.

4. Wang, Z.; Zhang, X. H.; Huang, Y.; Wang, H., Comprehensive evaluation of pharmaceuticals and personal care products (PPCPs) in typical highly urbanized regions across China. Environ Pollut. 2015; 204:223–232.

5. Eggen RIL, Hollender J, Joss A, Schärer M, Stamm C. Reducing the discharge of micropollutants in the aquatic environment: The benefits of upgrading wastewater treatment plants. Environ Sci Technol. 2014; 48:7683–7689.

6. Vila-costa M. Anthropogenic dissolved organic carbon and marine microbiomes. ISME J. 2020; 14:2646–2648.

7. da Silva GCX, Medeiros de Abreu CH, Ward ND, Belúcio LP, Brito DC, Cunha HFA, et al. Environmental Impacts of Dam Reservoir Filling in the East Amazon. Front Water. 2020; 2:11.

8. Kuroda K, Murakami M, Oguma K, Muramatsu Y, Takada H, Takizawa S. Assessment of groundwater pollution in Tokyo using PPCPs as sewage markers. Environ Sci Technol. 2012; 46:1455–1464.

9. Yang X, Chen F, Meng F, Xie Y, Chen H, Young K, et al. Occurrence and fate of PPCPs and correlations with water quality parameters in urban riverine waters of the Pearl River Delta, South China. ESPR. 2013; 20:5864–5875.

10. Liu WR, Zhao JL, Liu YS, Chen ZF, Yang YY, Zhang QQ, et al.Biocides in the Yangtze River of China: spatiotemporal distribution, mass load and risk assessment. Environ Pollut. 2015; 200:53–63.

11. Roberts J, Kumar A, Du J, Hepplewhite C, Ellis DJ. Christy AG, et al. Pharmaceuticals and personal care products (PPCPs) in Australia's largest inland sewage treatment plant, and its contribution to a major Australian river during high and low flow. Sci Total Environ. 2016; 541:1625–1637.

12. Rodea-Palomares I, Gonzalez-Pleiter M, Gonzalo S, Rosal R, Leganes F, Sabater S, et al. Hidden drivers of low-dose pharmaceutical pollutant mixtures revealed by the novel GSA-QHTS screening method. Sci Adv. 2016; 2:e1601272.

13. Sun Q, Li Y, Li M, Ashfaq M, Lv M, Wang H, et al. PPCPs in Jiulong River estuary (China): Spatiotemporal distributions, fate, and their use as chemical markers of wastewater. Chemosphere. 2016; 150:596–604.

14. Adeoya-Osiguwa SA, Markoulaki S. Pocock V, Milligan SR, Fraser LR.17β-Estradiol and environmental estrogens significantly affect mammalian sperm function. Hum Reprod. 2003; 18:100–107.

15. Tatarazako N, Ishibashi H. Teshima K, Kishi K, Arizono K. Effects of triclosan on various aquatic organisms. Environ Sci. 2004; 11:133–140.

16. MacLusky NJ, Hajszan T, Leranth C. The environmental estrogen bisphenol a inhibits

estradiol-induced hippocampal synaptogenesis. Environ Health Persp. 2005; 113:675–679.

17.   Rosi-Marshall EJ, Kincaid DW, Bechtold HA, Royer TV, Rojas M, Kelly JJ.
Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial
communities in stream biofilms. Ecol Appl. 2013; 23:583–593.

18.   Cerro-Gálvez E, Dachs J, Lundin D, Fernández-Pinos MC, Sebastián M, Vila-Costa M.
Responses of Coastal Marine Microbiomes Exposed to Anthropogenic Dissolved Organic
Carbon. Environ Sci Technol. 2021; 55:9609–9621.

19.   Martinez-Varela A, Cerro-Gálvez E, Auladell A, Sharma S, Moran MA, Kiene RP, et al.
Bacterial responses to background organic pollutants in the northeast subarctic Pacific
Ocean. (In Press, Environ Microbiol).   https://doi.org/10.1111/1462-2920.15646

20.   Adyari B, Shen D, Li S, Zhang L, Rashid A, Sun Q, et al. Strong impact of
micropollutants on prokaryotic communities at the horizontal but not vertical
scales in a subtropical reservoir, China. Sci Total Environ. 2020; 721:137767.

21.   Tlili A, Corcoll N, Arrhenius Å, Backhaus T, Hollender J, Creusot N, et al. Tolerance
patterns in stream biofilms link complex chemical pollution to ecological impacts. Environ
Sci Technol. 2020; 54:10745–10753.

22.   Chalew TEA, Halden RU. Environmental Exposure of Aquatic and Terrestrial Biota to
Triclosan and Triclocarban. J Am Water Resour Assoc. 2009; 45:4–13.

23.   Zhang WW, Yin K, Chen LX. Bacteria-mediated bisphenol A degradation. Appl Microbiol
Biotechnol. 2013; 97:5681–5689.

24.   Staples CA, Dome PB, Klecka GM, Oblock ST, Harris LR. A review of the environmental
fate, effects, and exposures of bisphenol A. Chemosphere. 1998; 36:2149-2173.

25.   Choi YJ, Lee LS. Aerobic Soil Biodegradation of Bisphenol (BPA) Alternatives Bisphenol
S and Bisphenol AF Compared to BPA. Environ Sci Technol. 2017; 51:13698–13704.

26.   McMurry LM, Oethinger M, Levy SB. Triclosan targets lipid synthesis. Nature. 1998;
394:531–532.

27.   Cabana H, Jiwan JLH, Rozenberg R, Elisashvili V, Penninckx M, Agathos SN, et al.
Elimination of endocrine disrupting chemicals nonylphenol and bisphenol A and personal
care product ingredient triclosan using enzyme preparation from the white rot fungus
Coriolopsis polyzona. Chemosphere. 2007; 67:770–778.

28.   Hu A, Ju F, Hou L, Li J, Yang X, Wang H, et al. Strong impact of anthropogenic
contamination on the co-occurrence patterns of a riverine microbial community. Environ
Microbiol. 2017; 19;4993–5009.

29.   Boyd TJ, Smith DC, Apple JK, Hamdan LJ, Osburn CL, Montgomery MT. Evaluating PAH
biodegradation relative to total bacterial carbon demand in coastal ecosystems: Are PAHs
truly recalcitrant? Microbial Ecology Research Trends. 2008. 1–38 p.

30.   Okere UV, Cabrerizo A, Dachs J, Ogbonnaya UO, Jones KC, Semple KT. Effects of pre-
exposure on the indigenous biodegradation of ^{14}C-phenanthrene in Antarctic soils. Int
Biodeterior Biodegrad. 2017; 125:189–99.

31.   Coll C, Bier R, Li Z, Langenheder S, Gorokhova E, Sobek A. Association between Aquatic
Micropollutant Dissipation and River Sediment Bacterial Communities. Environ Sci
Technol. 2020; 54:14380–14392.

32.   Bender EA, Case TJ, Gilpin ME. Perturbation experiments in community ecology :
theory and practice. Ecology. 1984; 65:1–13.

33.   Shade A, Peter H, Allison SD, Baho DL, Berga M, Bürgmann H, et al. Fundamentals of
microbial community resistance and resilience. Front Microbiol. 2012;3:1–19.

34.   Buerger S, Spoering A, Gavrish E, Leslin C, Ling L, Epstein SS. Microbial scout hypothesis,
stochastic exit from dormancy, and the nature of slow growers. Appl Environ Microbiol.
2012; 78:3221–3228.

35.   Lee S, Sorensen JW, Grady KL, Tobin TC, Shade A. Divergent extremes but convergent
recovery of bacterial and archaeal soil communities to an ongoing subterranean coal mine
fire. ISME J. 2017; 11:1447–1459.

36.   Lennon JT, den Hollander F, Wilke-Berenguer M, Blatch J. Principles of seed banks and the
emergence of complexity from dormancy. Nat Commun. 2021; 12:4807.

37.   Philippot L, Griffiths BS, Langenheder S. Microbial Community Resilience across
Ecosystems and Multiple Disturbances. Microbiol Mol Biol Rev. 2021; 85:e00026-20.

38.   Im J, Loffler FE. Fate of Bisphenol A in Terrestrial and Aquatic Environments. Environ Sci
Technol. 2016; 50;8403–8416.

39.   Sun Q, Li M, Ma C, Chen X, Xie X, Yu CP. Seasonal and spatial variations of PPCP
occurrence, removal and mass loading in three wastewater treatment plants located in
different urbanization areas in Xiamen, China. Environ Pollut. 2016; 208:371–33

40.   Sun Q, Wang Y, Li Y, Ashfaq M, Dai L, Xie X, et al. Fate and mass balance of bisphenol
analogues in wastewater treatment plants in Xiamen City, China. Environ Pollution. 2017;
225:542–549.

41.   Yamazaki E, Yamashita N, Taniyasu S, Lam J, Lam PKS. Moon HB, et al. Bisphenol A and
other bisphenol analogues including BPS and BPF in surface water samples from Japan,
China, Korea and India. Ecotoxicol Environ Saf. 2015; 122:565–572.

42.   Hu A, Li S, Zhang L, Wang H, Yang J, Luo Z, et al. Prokaryotic footprints in urban water
ecosystems: A case study of urban landscape ponds in a coastal city, China. Environ Pollut.
2018; 242:1729–1739.

43.   Wang J, Pan F, Soininen J, Heino J, Shen J. Nutrient enrichment modifies temperature-
biodiversity relationships in large-scale field experiments. Nat Commun. 2016; 7:13960.

44.   Hildebrand F, Tito RY, Voigt AY, Bork P,Raes J. LotuS: an efficient and user-friendly OTU
processing pipeline. Microbiome. 2014; 2:30.

45.   Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat
Methods. 2013; 10:996–998.

46.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: Improved data processing and web- based tools. Nucleic Acids
Res. 2013; 41:D590– D596.

47.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al,
QIIME allows analysis of high-throughput community sequencing data. Nat Methods. 2010;
7:335–336.

48.   Roller BRK, Stoddard SF, Schmidt TM. Exploiting rRNA Operon Copy Number to
Investigate Bacterial Reproductive Strategies. Nat Microbiol. 2017; 1:16160.

49.   Klappenbach JA, Saxman PR, Cole JR, Schmidt TM. rrndb: the ribosomal RNA operon
copy number database. Nucleic Acids Res. 2001; 29:181–184.

50.   Wu LW, Yang YF, Chen S, Zhao MX, Zhu ZW, Yang SH, et al. Long-term successional
dynamics of microbial association networks in anaerobic digestion processes. Water Res.

2016; 104:1–10.

51.   Stegen JC, Lin X, Fredrickson JK, Chen X, Kennedy DW, Murray CJ, et al. Quantifying community assembly processes and identifying features that impose them. ISME J. 2013; 7:2069–2079.

52.   Stegen JC, Lin X, Fredrickson JK, Konopka AE. Estimating and mapping ecological processes influencing microbial community assembly. Front Microbiol. 2015; 6:370.

53.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014; 15:1–21.

54.   Letunic I, Bork P. Interactive tree of life (iTOL) v4: Recent updates and new developments. Nucleic Acids Res. 2019; 47:W256–W259.

55.   Anderson MJ. Permutation tests for univariate or multivarite analysis of variance and regression. Can J Fish Aquat Sci. 2001; 58:626–639.

56.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. Vegan: community ecology package. 2018, R package version 1.17-4. Available from: https://cran.r-project.org/package=vegan

57.   Bashan A, Gibson TE, Friedman J, Carey VJ, Weiss ST, Hohmann EL, et al. Universality of Human Microbial Dynamics. Nature. 2016; 534:259–262.

58.   Vila JCC, Liu YY, Sanchez A. Dissimilarity–Overlap analysis of replicate enrichment communities. ISME J. 2020; 14:2505–2513.

59.   Ahlmann-Eltze C, Patil I. ggsignif: Significance Brackets for "ggplot2". R package version 0.6.1. 2021;(123). Available from: https://cran.r-project.org/package=ggsignif

60.   Lindström ES, Östman Ö. The importance of dispersal for bacterial community composition and functioning. PLoS One. 2011; 6:e25883.

61.   Shen D, Langenheder S, Jürgens K. Dispersal modifies the diversity and composition of active bacterial communities in response to a salinity Disturbance. Front Microbiol. 2018; 9:2188.

62.   Berga M, Székely AJ, Langenheder S. Effects of disturbance intensity and frequency on bacterial community composition and function. PLoS One. 2012; 7(5).

63.   Freilich S, Zarecki R, Eilam O, Segal ES, Henry CS, Kupiec M, et al. Competitive and cooperative metabolic interactions in bacterial communities. Nat Commun. 2011; 2:589.

64.   Pacheco AR, Moel M, Segrè D. Costless metabolic secretions as drivers of interspecies interactions in microbial ecosystems. Nat Commun. 2019; 10:103.

65.   Lee DG, Zhao F, Rezenom YH, Russell DH, Chu KH. Biodegradation of triclosan by a wastewater microorganism. Water Res. 2012; 46:4226–4234.

66.   Martin EK, Kemal S, Henrik NR, Alexander E, Natalia C, Henrik JC, et al. Triclosan changes community composition and selects for specific bacterial taxa in marine periphyton biofilms in low nanomolar concentrations. Ecotoxicology. 2020; 29:1083–1094.

67.   Wolff D, Krah D, Dötsch A, Ghattas AK, Wick A, Ternes TA. Insights into the variability of microbial community composition and micropollutant degradation in diverse biological wastewater treatment systems. Water Res. 2018; 143:313–324.

68.   Bajić D, Vila JCC, Blount ZD, Sánchez A. On the deformability of an empirical fitness landscape by microbial evolution. Proc Natl Acad Sci U S A. 2018;115:11286–11291.

69.   Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, et al. Patterns and processes of microbial community assembly. Microbiol Mol Biol Rev. 2013; 77:342–

714   56.

715   70.   Zhou J, Ning D. Stochastic community assembly : does It matter in microbial ecology.
716   Microbiol Mol Biol Rev. 2017; 81:1–32.

717   71.   Vellend M. Conceptual synthesis in community ecology. Q Rev Biol. 2010; 85:183–206.

718   72.   Jiao S, Zhang Z, Yang F, Lin Y, Chen W, Wei G. Temporal dynamics of microbial
719   communities in microcosms in response to pollutants. Mol Ecol. 2017; 26:923–936.

720   73.   Fukami T. Historical contingency in community assembly: integrating Niches, species pools,
721   and priority effects. Annu Rev Ecol Evol Syst. 2015; 46:1–23.

722   74.   Svoboda P, Lindström ES, Osman OA, Langenheder S. Dispersal timing determines the
723   importance of priority effects in bacterial communities.ISME J.2017; 12:644–646.

724   75.   Bernstein HC. Reconciling Ecological and Engineering Design Principles for Building
725   Microbiomes. mSystems. 2019; 4:1–5.

726   76.   Borchert E, Hammerschmidt K, Hentschel U, Deines P. Enhancing Microbial Pollutant
727   Degradation by Integrating Eco-Evolutionary Principles with Environmental Biotechnology.
728   (In Press). https://doi.org/10.1016/j.tim.2021.03.002

729   77.   Rocca JD, Muscarella ME, Peralta AL, Izabel-Shen D, Simonin M. Guided by microbes:
730   Applying community coalescence principles for predictive microbiome engineering.
731   mSystems. 2021; 6:e00538-21.

732   78.   Nemergut DR, Knelman JE, Ferrenberg S, Bilinski T, Melbourne B, Jiang L, et al. Decreases
733   in average bacterial community rRNA operon copy number during succession. ISME J. 2016;
734   10:1147–1156.

735   79.   Change C, Vila JCC, Bender M, Li R, Mankowski MC, Bassette M, et al. Engineering
736   complex communities by directed evolution. Nat Ecol Evol. 2021; 5:1011-1023.

738 **Table 1.** Number of OTUs assigned to the ecological categories opportunistic, sensitive,

739 and tolerant in the four micropollutant treatments: 'BPA' microcosms containing

740 bisphenol A, 'TCS' microcosms containing triclosan, 'MI' microcosms containing a

741 mixture of bisphenol A and triclosan, 'MII' microcosms containing a mixture of

742 bisphenol A, triclosan, bisphenol S and triclocarban.

	Number of OTUs in each ecological category		
Treatment	Opportunistic	Sensitive	Tolerant
BPA	1	64	6
TCS	1	47	15
MI	3	51	12
MII	0	41	10

Figure 1. Experimental setup. In microcosm batch 1 (B1), 600 ml of the initial

inoculum was added to bottles containing the micropollutants. Starting from the B1, 60

mL (10%) of the inoculum was pipetted from each of the replicates and transferred to

fresh medium containing 540 mL of 0.22-μm-filtered water and freshly added

micropollutants every 5 days. This procedure was carried out in the seven batches of

microcosms from batch 1 (B1) through batch 7 (B7). Samples for bacterial community

characterization and micropollutant removal were collected prior to the next inoculation.

Treatments are color-coded: 'Control' is (without micropollutant addition, grey), 'BPA'

(microcosms added with bisphenol A, red), 'TCS' (microcosms added with triclosan,

blue), 'MI' (microcosms added with a mixture containing bisphenol A and triclosan,

purple), 'MII' (microcosms added with a mixture containing bisphenol A and triclosan,

bisphenol S and triclocarban, yellow).

Figure 2. Percent removal of bisphenol A (BPA), triclosan (TCS), bisphenol S (BPS),

and triclocarbon (TCC) from the microcosms. (A) Removal in the single-

micropollutant (BPA and TCA) treatments; (B) removal in the mixed-micropollutant

(MI and MII) treatments. Removal was defined as the decrease in the concentration of

the chemical relative to the total concentration within each batch of microcosms

(differing by 5-day intervals). The open circles represent the removal percentage

obtained from each microcosm batch, and the black circles represent the outliner.

Tukey's *post-hoc* test was used to determine the taxonomic level at which the

differences occurred. The corresponding homogeneous groups were determined at

significance level $P < 0.05$ and are indicated by a and b.

Figure 3. Bacterial community dissimilarity in the micropollutant treatments vs. the

controls (i.e., 'Con.') over time (A) and the dynamic regularity within treatments (B).

Significant differences in the Bray-Curtis dissimilarity for each pairwise comparison of

the treatment and control (B1 microcosms vs. later microcosms in (A) were determined

in a Wilcoxon test: ***, $P < 0.001$; **, $0.001 < P < 0.01$; *, $0.01 < P < 0.05$. The

dissimilarity-overlap analysis within each treatment is shown in (B); out of total 210

sample pairs, only the sample pairs that had median overlap above cutoff = 0.913 were

shown in the plots. The gray lines indicate the linear regression with the slope (m), and

the P value (the fraction of bootstrap realizations in which the slope was negative).

Gray shaded areas indicate the 95% confidence intervals of the linear regression.

Significance level was determined at $P < 0.05$.

	Figure 4. The Dissimilarity-Overlap analysis of pairwise treatments and in the

treatments vs. controls (i.e., 'Con.'). Out of total 441 sample pairs, only the dark gray

points that had median overlap above cutoff = 0.854 were shown in the plots. The gray

lines indicate the linear regression with the slope (m) and the *P* value (the fraction of

bootstrap realizations in which the slope was negative). Gray shaded areas indicate the

95% confidence intervals. Significance level was detected at *P* < 0.05.

Figure 5. Changes in the relative abundances (rows of heatmap) of the OTUs assigned

to the ecological categories tolerant and sensitive across the three growth phases

(columns of heatmap) (A) and their phylogenetic distribution (B). In (A), color

gradients of heatmap represent the relative abundances of individual by column with

warm colors (toward red) indicating high abundance and cold colors (toward blue

indicating low abundance within the sample. Column labels are batch IDs and the

grouping of growth phases, and row labels the OTU IDs. Tolerant and sensitive OTUs

were denoted in green and blue colors, respectively. In (B), the innermost circle

indicates the taxonomy (phylum/class) of the OTUs. The outer circles indicate the

ecological group of OTUs in each treatment group (BPA, TCS, MI, MII from inside to

outside, respectively).

Figure 6. Relative proportion of the ecological processes (homogeneous selection,

variable selection dispersal limitation, and ecological drift) contributing to the

communities assembled in the micropollutant treatments (BPA, TCS, MI or MII) vs.

the controls (i.e., 'Con.'). Each of the three phases (corresponding to the early, mid, and

late stages of the press disturbance) was classified according to the similarity in the

average 16S rRNA gene copy numbers as an indicator of community growth

progression. Three growth phrases across microcosm batches were: phase 1 (B1–B2),

phase 2 (B3–B4), and phase 3 (B5–B7), as shown in Fig. S2G.

Figure 7 Conceptual model of microbial community assembly processes following a
press disturbance (e.g., recurrent micropollutant exposure). Considering the time
continuity of the batch inoculation, the three phases of community growth progression
(1, 2, and 3) also correspond to the early, mid, and late stages of the 35-day disturbance.
This conceptual model illustrates when and how stochastic/deterministic processes
contribute to the dynamics of microbial communities repeatedly exposed to
micropollutants. Changes in community trajectory over time in the TCS or the MII
treatment are indicated by the black line, MI treatment by the red line, and the BPA
treatment by the purple line. Overall, stochastic and deterministic processes act in
concert during community dynamics: stochastic assembly dominates at the early stage
of a press disturbance/community development, and deterministic assembly at the later
stage. Major community assembly mechanisms are delineated for each phase.
Community succession accelerates at phase 2 such that multiple equilibria occur at
phase 3. Note that community succession and multiple equilibria are not exclusive, but
community turnover is high towards the end of the disturbance.