Systemic regulation of nitrogen acquisition and use in *Oryza longistaminata* ramets

under nitrogen heterogeneity[1]

Misato Kawai[a], Ryo Tabata[a], Miwa Ohashi[a], Haruno Honda[a], Takehiro Kamiya[b], Mikiko Kojima[c],

Yumiko Takebayashi[c], Shunsuke Oishi[d], Satoru Okamoto[a,e], Takushi Hachiya[a,f], Hitoshi

Sakakibara[a,c,2,3]

[a]Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya 464-8601,

Japan

[b]Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo, Tokyo

113-8657, Japan

[c]RIKEN Center for Sustainable Resource Science, Tsurumi, Yokohama 230-0045, Japan

[d]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Chikusa, Nagoya

464-8602, Japan

[e]Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan

[f]Department of Molecular and Function Genomics, Interdisciplinary Center for Science

Research, Shimane University, Matsue, 690-8504, Japan

One sentence summary: *Oryza longistaminata*, a rhizomatous wild rice, systemically regulates

nitrogen acquisition and use in response to spatially heterogeneous nitrogen availability via

inter-ramet communication.

Footnotes:

Author Contributions: M.Ka., H.H., M.O., S.Ok., T.H. and H.S. designed the experiments and

analyzed the data; M.Ka. and H.H. performed all experiments and M.O., S.Ok., T.H., and H.S.

supervised M.Ka. and H.H.; R.T. and T.K. helped with RNAseq data analysis; M.Ko. and Y.K.

provided technical support to quantify cytokinins; S.Oi. synthesized CEP peptides; H.S.

conceived the project and wrote the manuscript with contributions from all co-authors; H.S.

agrees to serve as the author responsible for contact and ensures communication.

Funding:

[1]This research was supported by the Advanced Low Carbon Technology Research and

Development Program from the Core Research for Evolutional Science and Technology (to

H.S.; no. JPMJCR13B1), PRESTO (to S.Ok.; no. JPMJPR17Q2) by JST, Leading Initiative for

Excellent Young Researchers (to S.Ok.), Grants-in-Aid from the Ministry of Education, Culture,

Sports, Science and Technology, Japan (to H.S.; 17H06473 and 19H00931, to R.T.; 18K05373

and 20H05501, to T.K.; 17H03782), and by the Program for Promoting the Enhancement of

Research Universities, Nagoya University.

[2]Author for contact: sakaki@agr.nagoya-u.ac.jp.

[3]Senior author.

Abstract

Oryza longistaminata, a wild rice, vegetatively reproduces and forms a networked clonal colony

consisting of ramets connected by rhizomes. Although water, nutrients, and other molecules can

be transferred between ramets via the rhizomes, inter-ramet communication in response to

spatially heterogeneous nitrogen availability is not well understood. We studied the response of

ramet pairs to heterogeneous nitrogen availability by using a split hydroponic system that

allowed each ramet root to be exposed to different conditions. Ammonium uptake was

compensatively enhanced in the sufficient-side root when roots of the ramet pairs were exposed

to ammonium-sufficient and deficient conditions. Comparative transcriptome analysis revealed

that a gene regulatory network for effective ammonium assimilation and amino acid biosynthesis

was activated in the sufficient-side roots. Allocation of absorbed nitrogen from the nitrogen-

sufficient to the deficient ramets was rather limited. Nitrogen was preferentially used for newly

growing axillary buds on the sufficient-side ramets. Biosynthesis of *trans*-zeatin, a cytokinin, was

up-regulated in response to the nitrogen supply, but *trans*-zeatin appears not to target the

compensatory regulation. Our results also implied that the *O. longistaminata* ortholog of

OsCEP1 plays a role as a nitrogen-deficient signal in inter-ramet communication, providing

compensatory up-regulation of nitrogen assimilatory genes. These results provide insights into

the molecular basis for efficient growth strategies of asexually proliferating plants growing in

areas where nitrogen distribution is spatially heterogeneous.

Introduction

Oryza longistaminata is a wild rice species that preferentially grows in wetlands and

vegetatively reproduces through rhizome growth that can be so vigorous as to occupy an area

completely (Vaughan, 1994). Rhizome growth is characterized by the underground horizontal

outgrowth of axillary buds to form new rhizomes that expand into new territory (Yoshida et al.,

2016; Fan et al., 2017; Kyozuka, 2017; Bessho-Uehara et al., 2018; Toriba et al., 2020;

Shibasaki et al., 2021). Rhizome tips developmentally transform into photosynthetic above-

ground organs by growing into a new plantlet called a ramet. Continuous rhizome growth and

subsequent transformation form a networked clonal colony. In addition to *O. longistaminata*,

many other plant species vegetatively reproduce through rhizome growth, such as

Phyllostachys edulis (Moso bamboo) and *Zoysia matrella* (Manila grass)(Guo et al., 2021), and

show vigorous fertility. Since ramets are connected via rhizomes, water, nutrients, and other

molecules can be transferred between ramets (De Kroon et al., 1998). Therefore, the growth

and metabolism of a ramet are not totally independent of the neighboring ramets but have some

influence on each other via the rhizomes.

Nitrogen is one of the most limiting macronutrients for plants, so nitrogen acquisition and use

efficiency significantly affect plant growth and development. In submerged and reductive soil

conditions where wetland rice grows, plants mainly use ammonium ions as the inorganic

nitrogen source (Yoshida, 1981). In *Oryza sativa*, ammonium ions in the soil are taken up by

ammonium transporters (AMTs)(Sonoda et al., 2003; Suenaga et al., 2003; Yuan et al., 2007; Li

et al., 2016; Jia and von Wirén, 2020; Lee et al., 2020a), and then are initially assimilated into

glutamine and glutamate by the glutamine synthetase (GS)/ glutamate synthetase (GOGAT)

cycle (Lea and Miflin, 1974). The cycle is mainly composed of cytosolic GS and plastidic NADH-

GOGAT isoforms in the root (Ishiyama et al., 2004; Tabuchi et al., 2005; Tabuchi et al., 2007;

Funayama et al., 2013; Yamaya and Kusano, 2014; Ji et al., 2019; Lee et al., 2020b). Several

amino acids are synthesized by amino-transfer reactions from glutamine and glutamate to

organic acids derived from glycolysis, the tricarboxylic acid cycle, and the pentose phosphate

pathway. Previous studies using *O. sativa*, *Arabidopsis thaliana*, and *Nicotiana tabacum*

revealed a wide and versatile regulatory network of gene expression in response to nitrogen

(nitrate or ammonium) nutrition, including genes for the processes of uptake, assimilation,

amino acid synthesis, carbon skeleton supply, and hormone signaling (Scheible et al., 1997;

Scheible et al., 2004; Wang et al., 2004; Sakakibara et al., 2006; Chandran et al., 2016; Yang et

al., 2017; Sun et al., 2020).

Plants optimize their growth and development according to the amount of available nitrogen,

and signaling molecules, including phytohormones and peptide hormones, play a key role in this

regulation (Nishida and Suzaki, 2018; Ruffel, 2018; Sakakibara, 2021; Wheeldon and Bennett,

2021). Cytokinin, especially *trans*-zeatin (tZ), promotes nitrogen-responsive shoot growth via

modulating shoot meristem activity (Kiba et al., 2013; Davière and Achard, 2017; Osugi et al.,

2017; Landrein et al., 2018). An abundant nitrate supply promotes *de novo* tZ biosynthesis via

the up-regulation of *ADENOSINE PHOSPHATE-ISOPENTENYLTRANSFERASE3* (*IPT3*) and

CYP735A2 in Arabidopsis (Takei et al., 2004; Maeda et al., 2018; Naulin et al., 2019;

Sakakibara, 2021). In *O. sativa*, a glutamine-related signal stimulates *OsIPT4* and *OsIPT5*

expression that are involved in axillary bud outgrowth from shoots (Kamada-Nobusada et al.,

2013; Ohashi et al., 2017). In *O. longistaminata*, the outgrowth of rhizome axillary buds in

response to nitrogen nutrition is regulated by a similar process to that of the axillary shoot bud in

O. sativa, despite differences in the physiological roles of these organs (Shibasaki et al., 2021).

In contrast, strigolactone is involved in the nutrient-responsive suppression of shoot axillary bud

outgrowth (Gomez-Roldan et al., 2008; Umehara et al., 2008; Minakuchi et al., 2010; Umehara

et al., 2010). Strigolactone biosynthetic genes are known to be up-regulated in the roots of *O.*

sativa and *Zea mays* under nitrogen-deficient (Sun et al., 2014; Xu et al., 2015; Ravazzolo et

al., 2019; Ravazzolo et al., 2021) and phosphorus-deficient (Umehara et al., 2010) conditions.

It is essential for plants to efficiently acquire nutrients for optimal growth and development

even though mineral nutrients are heterogeneously distributed in the soil. Recent studies using

a split-root system in Arabidopsis showed that when part of the root system is subjected to

nitrogen deficiency, C-TERMINALLY ENCODED PEPTIDE1 (CEP1), synthesized in the

deficient root, is translocated to shoots via the xylem (Ohyama et al., 2008; Tabata et al., 2014).

The perception of CEP1 by the CEP1 RECEPTOR (CEPR) in shoots triggers the expression of

CEP DOWNSTREAM (CEPD) proteins that are translocated to the root system via the phloem,

thereby promoting compensatory nitrogen uptake by inducing the expression of high-affinity

nitrate transporters, including *NITRATE TRANSPORTER2.1* (*NRT2.1*) in the nitrogen-ample

side (Tabata et al., 2014; Okamoto et al., 2016; Ohkubo et al., 2017; Ruffel and Gojon, 2017;

Ota et al., 2020; Ohkubo et al., 2021). In rhizomatous plants, ramets connected by rhizomes are

regarded as one individual; however, it remains unknown whether each ramet responds to

nitrogen availability independently or compensatively when growing in areas where nitrogen

distribution is spatially heterogeneous.

In this study, we characterized the response of *O. longistaminata* ramet pairs, connected by a

rhizome, to spatially heterogeneous nitrogen conditions and found a compensatory gene

regulatory network for effective acquisition and assimilation of nitrogen in the ample-side ramet

root. Cytokinin biosynthesis was up-regulated in response to nitrogen supply, but this increase

appears to be independent of the compensatory regulation. Our results also imply that an *O.*

longistaminata ortholog of the *OsCEP1* gene plays a role in inter-ramet communication. This

study provides valuable hints for understanding the molecular basis of efficient growth strategies

of vegetatively proliferating plants.

Results

Compensatory promotion of ammonium uptake in response to a spatially heterogeneous

nitrogen supply in *O. longistaminata* ramet pairs

To examine nitrogen-related inter-ramet communication via the rhizome, we established a

split-hydroponic experimental system for *O. longistaminata* ramet pairs (Supplemental Fig. S1).

After preparing young ramet pairs of comparable growth stages that developed at adjacent

rhizome nodes, we treated the roots of each ramet to different levels of nitrogen nutrition to

mimic a condition of spatially heterogeneous nitrogen availability. We set up three conditions: (i)

both ramet roots were exposed to 2.5 mM NH_4Cl (+N), (ii) one ramet root was exposed to 2.5

mM NH_4Cl (+N split), and the other was exposed to 0 mM NH_4Cl (-N split), (iii) both ramet roots

were exposed to 0 mM NH_4Cl (-N) (Fig. 1A and Supplemental Fig. S1F). We compared the

absorption activity of ammonium between the +N and +N split roots using ^{15}N-labeled NH_4Cl as

the tracer. The results showed that the ammonium absorption activity was significantly

increased by about 1.6-fold in the +N split ramets compared to the +N ramets (Fig. 1B).

Expression of the *OlAMT1;2* gene, an ortholog of *O. sativa AMT1;2* that encodes an

ammonium-inducible ammonium transporter, was approximately 3.6-fold higher in the roots of

+N split than those of +N (Fig. 1C). These results suggest that when *O. longistaminata* ramet

pairs are exposed to nitrogen that is spatially heterogeneous, ammonium uptake is

compensatively enhanced in the ramet exposed to the nitrogen-rich condition and is

accompanied by an up-regulation of *OlAMT1;2*.

Compensatory regulation of amino acid biosynthesis and related genes in response to a

spatially heterogeneous nitrogen supply

To explore the molecular basis underlying the compensatory response to a heterogeneous

nitrogen supply, we analyzed the transcriptome using RNA sequencing (RNAseq) of the roots at

24 h after the start of the +N, +N split, -N split, and -N treatments. Up- and down-regulated

genes were identified from the expression data of the +N split, -N split, and -N treatments by

comparison with that of the +N condition (false discovery rate [FDR] < 0.05). In comparison to

the +N condition, 416 genes were significantly up-regulated in +N split root (Figs. 2A and 2B,

Supplemental Table S1). Also, 2361 and 2776 genes were up-regulated in +N roots compared

to the -N and -N split treatments, respectively (Fig. 2B, Supplemental Tables S2 and S3).

Among the 416 genes, the identity of 295 genes overlapped with both the up-regulated genes in

the +N treatment compared to the -N treatment and those in the +N treatment compared to the -

N split treatment (110 genes), or only with those in the +N treatment compared to the -N split

treatment (179 genes) and to the -N treatment (6 genes) (Fig. 2B). Gene Ontology (GO)

analysis showed that genes associated with metabolic processes related to nitrogen and

carbon, such as glycolysis (p = 2.7e-8), monosaccharide metabolic processes (p = 1.8e-7),

oxoacid metabolic processes (p = 3.5e-7), and cellular amino acid metabolic processes (4.6e-

6), were enriched among the 416 genes (Fig. 2C). Specifically, the up-regulated genes included

those involved in nitrogen assimilation and amino acid synthesis, such as genes for a cytosolic

glutamine synthetase *OlGS1;2*, an NADH-dependent glutamate synthase *OlNADH-GOGAT2*,

an aspartate aminotransferase *OlAspAT*, an aspartate semialdehyde dehydrogenase

OlASADH, an aspartate kinase-homoserine dehydrogenase *OlAK-HSDH1*, an alanine

aminotransferase *OlAlaAT1*, and an aspartate kinase *OlAK* (Supplemental Table S1). Using

reverse transcription-quantitative PCR (RT-qPCR), we confirmed that these genes were

significantly up-regulated in +N split roots compared to +N (Fig. 3B). In addition, the expression

of glycolytic enzyme genes involved in supplying carbon skeletons for amino acid biosynthesis,

such as glucose-6-phosphate isomerase (*OlGPI*), triosephosphate isomerase (*OlTPI1*), fructose

1,6-bisphosphate aldolase (*OlALDC1*), glyceraldehyde 3-phosphate dehydrogenase

(*OlGAPC1*), phosphofructokinase (*OlPFK5*), enolase (*OlENO1*), a glycolysis-related pyruvate

orthophosphate dikinase (*OlPPDKA*), and phosphoenolpyruvate carboxylase (*OlPPC4*) were

also significantly up-regulated in the +N split root condition compared to the +N treatment (Fig.

3A、Supplemental Table S1). These results suggest that a compensatory gene regulatory

network to acquire and assimilate ammonium effectively is activated in the ramet roots on the

nitrogen-sufficient side of the ramet pairs.

To know the impact of transcriptome changes on the accumulation level of amino acids, we

analyzed the amino acid concentrations in roots from the +N, +N split, -N split, and -N

treatments at a longer treatment period (48 h). As a result, the accumulation level of glutamine,

asparagine, aspartate, alanine, glutamate, and arginine were significantly higher in the +N split

roots than in the -N split roots (Fig. 3C). Among these amino acids, aspartate and alanine levels

in roots from the +N split treatment were significantly higher than those in the +N treatment.

Other amino acids except for arginine showed a similar tendency. These results are in line with

the transcriptome changes in response to heterogeneous nitrogen availability.

On the other hand, 1848 and 2087 genes were up-regulated in the -N and -N split treatments

compared to the +N condition, respectively (Supplemental Table S4 and S5). Overall, the

expression of 1176 genes overlapped between the two conditions (Supplemental Fig. S2). One

hundred and forty-five genes were up-regulated in the +N treatment compared to the +N split

treatment (Supplemental Table S6), and 15 genes overlapped with the -N-up-regulated genes

(Supplemental Fig. S2).

Allocation of absorbed nitrogen between ramets via the rhizome when nitrogen

availability is spatially heterogeneous

To examine the allocation of absorbed nitrogen in ramet pairs when nitrogen availability is

spatially different, $^{15}NH_4Cl$ was fed to +N and +N split roots with the same conditions as shown

in Fig. 1B, and the roots were grown for another 7 days with a non-labeled nitrogen source. In

this experimental condition, ^{15}N was detected in both the shoots and roots of the systemic +N

and -N split ramets that had not been fed with $^{15}NH_4Cl$ (Fig. 4A), indicating that absorbed

nitrogen is allocated to neighboring ramets via the rhizome. The percentage distribution of ^{15}N

was higher in the shoots than in the roots. Interestingly, the allocation of absorbed nitrogen from

the +N split to the -N split ramets was less than that from the +N to the +N ramets.

Growth response of ramets in spatially heterogeneous nitrogen conditions

To know the long-term effects of nitrogen split treatment on the growth of ramets, we

exposed ramet pairs to different nitrogen conditions for 5 weeks, and growth parameters for

each ramet were monitored (Figs. 4B to E). Plant height, the number of fully developed leaves,

chlorophyll content, and the number of growing axillary buds were significantly higher in the +N

and +N split ramets than in the -N split and -N ramets. The chlorophyll content was lower in the

-N split and -N ramet shoots, but the decrease was significantly alleviated in the -N split ramets

compared to the -N ramets (Fig. 4D), suggesting that the allocated nitrogen was used to retain

photosynthetic function. In contrast, the change in the number of growing axillary buds was

almost zero for the -N and -N split ramets, whereas the axillary bud number increased

significantly in the +N split and +N ramets (Fig. 4E). In particular, the axillary bud number was

significantly higher in the +N split ramets than in the +N ramets. These results suggest that

when neighboring ramets are exposed to different levels of nitrogen availability, the allocation of

absorbed nitrogen from the N-sufficient ramet to the deficient ramet is rather limited and,

nitrogen is preferentially allocated for newly growing axillary buds.

Response of cytokinin and strigolactone biosynthetic genes in response to a

heterogeneous nitrogen supply

To gain insight into the growth promotion of axillary buds in the +N split ramets, we focused

on cytokinin and strigolactone, two phytohormone families that promote and inhibit axillary bud

outgrowth, respectively. We analyzed the expression levels of cytokinin and strigolactone

biosynthetic genes by RT-qPCR using the roots of ramet pairs that had been split-treated for 7

days. The expression levels of cytokinin biosynthetic genes, *OlIPT4*, *OlIPT5*, *OlCYP735A3*, and

OlCYP735A4 were higher in the roots of the +N and +N split ramets than in those of the -N and

-N split ramets (Fig. 5A), and there was no difference between the +N and +N split conditions.

We also analyzed the concentration of cytokinins in the roots. The level of *trans*-zeatin (tZ)-

type cytokinins, including the riboside and ribotide precursors, was higher in the +N and +N split

roots compared to the -N split and -N roots, whereas the levels of N^6-(Δ^2-isopentenyl)adenine

(iP)-type cytokinins were somewhat lower (Fig. 5B, Supplemental Table S7). The tZ-type

cytokinin content in the +N and +N split roots was comparable, suggesting that de novo tZ-type

cytokinin biosynthesis is enhanced in response to nitrogen supply but is not under the

compensatory regulation.

Although strigolactone species in *O. longistaminata* have not been well characterized yet, we

analyzed orthologs of *O. sativa D27*, *D17*, and *D10* (*OlD27*, *OlD17*, and *OlD10*, respectively)

encoding enzymes involved in the production of carlactone, an intermediate of strigolactone

biosynthesis (Alder et al., 2012). The expression level of *OlD10* was significantly higher in -N

ramet roots than in other conditions, although *OlD27* and *OlD17* expression levels were

essentially similar and slightly lower, respectively (Fig. 5C). This result implies that de novo

strigolactone biosynthesis might be up-regulated in nitrogen-deficient roots.

Possible involvement of a CEP1-type peptide in inter-ramet nitrogen-deficiency signaling

In Arabidopsis, CEP1 plays a key role in the systemic regulation of nitrate acquisition in

response to heterogeneous nitrogen conditions as a root-to-shoot signaling molecule (Tabata et

al., 2014; Okamoto et al., 2016). To investigate whether a CEP1-type peptide is involved in the

observed inter-ramet communication, we analyzed the response of *O. longistaminata* CEP gene

orthologs to spatially heterogeneous nitrogen availability. In *O. sativa*, 15 genes encode CEPs

(Sui et al., 2016). We searched for the orthologs in *O. longistaminata* and found 15 sequences

corresponding to each of the *O. sativa* genes (Supplemental Table S8), and eight of the genes

(*OlCEP5*, *OlCEP6.1*, *OlCEP9*, *OlCEP10*, *OlCEP11*, *OlCEP12*, *OlCEP14*, and *OlCEP15*) were

annotated in our RNAseq data. However, expression of these eight genes was not significantly

different in -N roots compared to +N roots. Next, we focused on *OlCEP1* and analyzed its

expression level by RT-qPCR because the peptide sequences encoded by *OsCEP1* (also

OlCEP1) belong to the same group (group I) as Arabidopsis CEP1 based on the structural

features (Delay et al., 2013; Sui et al., 2016). In the RT-qPCR analysis, the expression level of

OlCEP1 in -N roots was significantly higher than in the +N and +N split roots (Fig. 6A).

To gain further insight into the involvement of *OlCEP1* gene products in nitrogen-related

ramet-to-ramet signaling, we examined the effect of an exogenous application of OlCEP1

peptides on the expression of *OlAMT1;2* and *OlGS1;2* genes. In addition to *OsCEP1*, the

coding sequence of the *OlCEP1* gene contains multiple CEP sequences containing proline

residues. Thus, we synthesized peptides OlCEP1a to OlCEP1d (See Materials and Methods)

and used a mixture. We treated ramet roots with the peptide mixture under nitrogen-sufficient

conditions to repress the endogenous *OlCEP1* expression. When we applied OlCEP1 peptides

to the roots of one side of the ramets, the accumulation level of transcripts for *OlAMT1;2* and

OlGS1;2 increased in both the local side and systemic side of the ramet roots in 6 h. The

transcript level significance was maintained in the systemic side of the root even at 24 h after

treatment (Fig. 6B), suggesting that CEP1-related signaling is possibly involved in the systemic

regulation of the genes via the rhizomes.

Discussion

In this study, we demonstrated inter-ramet communication occurs in *O. longistaminata* via

rhizomes for a systemic response to spatially heterogeneous nitrogen availability. When ramet

pairs of *O. longistaminata* connected by a rhizome were exposed to different nitrogen nutrient

conditions, a series of gene networks that allow complementary absorption and assimilation of

ammonium was activated in the root of the ramet in the nitrogen-sufficient condition. This

network also included genes capable of supplying the carbon skeletons for amino acid

synthesis, such as those involved in glycolysis. The expression level of these genes in the

nitrogen-sufficient side of the heterogeneous condition was higher than in the nitrogen

homogeneously sufficient condition, suggesting that the nitrogen-deficient side ramet conveyed

some kind of nitrogen deficiency signal to the sufficient side ramet via the rhizome to trigger the

systemic response.

In our transcriptome analysis, a large part of the compensatively up-regulated genes

overlapped with ammonium-responsive genes (295 genes/416 genes, Fig. 2B), indicating that

the expression of the compensatory genes is up-regulated by an ammonium-signal *per se* but

further boosted by the input of a systemic nitrogen-deficient signal from the adjacent ramet. It is

not clear at present whether a de-repression or a more facilitative regulatory event underlies the

process.

A small part of the nitrogen absorbed in the ramet roots in the heterogeneous nitrogen-

sufficient condition (+N split) was distributed to the adjacent nitrogen-deficient ramets (-N split),

contributing to the maintenance of the chlorophyll content. However, no evidence was obtained

to suggest that the distributed nitrogen was used for growth of the nitrogen-deficient side ramet.

In a previous study in *Carex flacca* examining water and nitrogen transfer between ramets via

the rhizome, the direction of nitrogen nutrient transport depended on the direction of water

transport (De Kroon et al., 1998). In rice cultivars, the nitrogen-sufficient condition elevates the

leaf transpiration rate compared to the limited condition (Xiong et al., 2015). Thus, it is likely that

the translocation of nitrogen from the sufficient side to the deficient side against the water flow is

limited between rhizome-connected ramets experiencing different nitrogen conditions. Our

results suggest that most compensatively acquired nitrogen is locally used for growth on the

sufficient side. Given that an *O. longistaminata* cluster is a single clonal colony, this use of

nitrogen might be a strategy to ensure the colony's survival in limited and heterogeneous

nitrogen conditions.

Notably, genes for the biosynthesis of tZ-type cytokinins, *IPTs* and *CYP735As*, were up-

regulated by the nitrogen supply. Still, the expression levels and accumulation of tZ-type

cytokinins (tZ and its precursors) were comparable between the +N and +N split treatments in

our experiments (Fig. 5B). Therefore, we hypothesize that cytokinin synthesis is similarly up-

regulated in response to both homogeneous and heterogeneous nitrogen conditions.

Ammonium acquisition and amino acid synthesis, however, are compensatively enhanced in the

nitrogen-rich ramet roots under heterogeneous conditions, resulting in a greater supply of

nitrogen assimilates for axillary bud outgrowth. Both the cytokinin signal and the supply of

building blocks could contribute to the preferential axillary shoot growth.

In our analysis, iP and its precursors were in low abundance in the +N and +N split roots,

mainly due to the decreased levels of its ribotide precursor (iPRPs, Fig. 5B). At present, we do

not have a clear explanation for the opposite trend despite the up-regulation of its biosynthetic

genes, *OlIPTs*. The iP ribotide precursor might be over consumed by CYP735As to produce tZ-

type species.

Previous studies in *O. sativa* and *O. longistaminata* indicated that *NADH-GOGAT1* is under

the same control as *IPT4* in the local nitrogen response and that glutamine-related signaling is

involved in the regulation (Kamada-Nobusada et al., 2013; Ohashi et al., 2017; Shibasaki et al.,

2021). *OlNADH-GOGAT2*, an isogene, is compensatively up-regulated in our experimental

conditions, whereas *OlNADH-GOGAT1* had a similar expression pattern to *OlIPT4*

(Supplemental Fig. S3). In *O. sativa*, *NADH-GOGAT2* is mainly expressed in mature leaves and

plays a role in providing glutamate for the GS1;1 reaction in vascular tissues for nitrogen

remobilization and recycling (Tabuchi et al., 2005; Tabuchi et al., 2007; Yamaya and Kusano,

2014). Our RNAseq data show that *OlGS1;1* is up-regulated in nitrogen-deficient conditions

(Supplemental Tables S4 and S5) in a manner distinct from *OlNADH-GOGAT2*. Thus, the

physiological role of the cytosolic GS isoforms and NADH-GOGAT isoforms might be different

between *O. sativa* and *O. longistaminata*.

Systemic regulation of nitrate acquisition in response to heterogeneous nitrogen nutrient

conditions by CEP1-CEPR-CEPD has been identified in Arabidopsis (Tabata et al., 2014;

Okamoto et al., 2016; Ohkubo et al., 2017; Ruffel and Gojon, 2017; Ota et al., 2020; Ohkubo et

al., 2021). In our analysis, the *O. longistaminata* ortholog of *OsCEP1* was markedly induced by

nitrogen deprivation, and exogenous application of the synthetic CEP1 peptides increased the

expression level of *OlAMT1;2* and *OlGS1;2* on the systemic side as well as the local side. This

finding implies that CEP plays a role in inter-ramet communication as a nitrogen-deficient

signaling molecule via the rhizome. Also, our results suggest that both the CEP-derived signal

and nitrogen sufficiency are necessary to drive the compensatory up-regulation system, as is

the case for *NRT2.1* up-regulation in Arabidopsis (Tabata et al., 2014). The systemic response

mechanism responding to heterogeneous nitrogen conditions in monocots, including rice, is still

largely unexplored. Identifying and characterizing the CEP receptor and downstream factors,

including CEPD orthologs in *O. longistaminata*, is necessary to understand entirely the whole

system at the molecular level. The findings of this study will be useful in characterizing survival

strategies of rhizomatous plants in response to heterogeneous nitrogen environments.

Materials and methods

Plant materials and growth conditions

The perennial wild rice species *O. longistaminata* (IRGC10404) was hydroponically grown

under natural light conditions in a greenhouse with a nutrient solution described by Kamachi et

al. (1991), except that the solution contained 1 mM NH_4Cl as the sole nitrogen source and was

renewed once every 3 or 4 days. Pairs of young ramets grown on the proximal nodes of

rhizomes were excised and used in experiments. The growth stage of all ramet pairs was

similar; each ramet had two to three fully developed leaves (Supplemental Fig. S1).

Split treatment

The roots of each ramet pair, connected by rhizomes, were incubated in two separate pots

(Supplemental Fig. S1). One L of nutrient solution was used for each ramet. For the ±N split

treatment, one of the ramet pairs was treated with the nutrient solution containing 2.5 mM NH_4Cl

(+N split), and the other was treated with 0 mM NH_4Cl (-N split). For comparison, both ramet

pairs were treated with 2.5 mM NH_4Cl (+N) or 0 mM NH_4Cl (-N). Treatments for the growth

analysis, and cytokinin and strigolactone response analysis were conducted in a greenhouse.

All other treatments were conducted in a growth cabinet (LPH-411PFQDT-SPC, NK System,

Osaka, Japan) with the following environmental conditions: 16 h (28°C) light/8 h (24°C) dark at

400 μmol photons m^{-2} s^{-1}.

$^{15}NH_4$ tracer experiments

The ramet pairs were initially incubated with water for 3 days, followed by a 24-h split treatment

with stable isotope-free nitrogen nutrient solutions (2.5 mM NH_4Cl for the +N ramet and the +N

split ramet, and 0 mM NH_4Cl for the -N ramet). Roots of the ramets were soaked in 800 mL of 1

mM $CaSO_4$ for 1 min to remove the nitrogen treatment solution. The solution for the +N-treated

side was then replaced with a solution containing 2.5 mM $^{15}NH_4Cl$ (^{15}N 99 atom%, Shoko

Science Co., Ltd., Yokohama, Japan), and $^{15}NH_4^+$ was allowed to be absorbed for 5 min. The

roots of ramet pairs were then soaked in 600 mL of 1 mM $CaSO_4$ for 1 min to wash out ^{15}N and

then treated again with the stable isotope-free +N or -N hydroponic solutions for 22 h to

measure the ammonium absorption activity and for 7 days to measure the absorbed nitrogen

distribution with daily changing of the culture solution in a growth cabinet (LPH-411PFQDT-

SPC, NK System) with 16 h (30°C) light/8 h (30°C) dark at 400 μmol photons m^{-2} s^{-1}.

The above-ground tissues, rhizomes, and roots were separately harvested and dried in an

oven at 70°C for at least 5 days. All dried tissues were weighed and ground into fine powders.

The ^{15}N and total nitrogen contents were analyzed by Shoko Science Co. with an elemental

analyzer-isotope ratio mass spectrometer (Flash2000-DELTA plus Advantage ConFloIII

System, Thermo Fisher Scientific, Waltham, MA, United States).

The absorption activity of ammonium was calculated as follows. The increase in ^{15}N (μmol)

in each sample ($\Delta^{15}N_s$) was calculated from equation (1). W_s, dry weight of the sample (gDW);

N_s, the total nitrogen concentration of the sample (%); $^{15}N_s$, the ^{15}N concentration of the sample;

and $^{15}N_0$, the naturally occurring ^{15}N concentration (%) of the sample.

$$\Delta^{15}N_s = W_s \times 10^6 \times \frac{N_s}{100} \times \frac{^{15}N_s - {}^{15}N_0}{100} \times \frac{1}{15} \qquad (1)$$

The $\Delta^{15}N_s$ of each sample (above-ground tissues and roots) were summed to obtain the

increase of ^{15}N in the whole ramet pair ($\Delta^{15}N_w$). The absorption activity of ammonium (μmol

gDW^{-1} h^{-1}) in roots under +N or +N split conditions was calculated using equation (2). A, the

absorption activity of ammonium; W_r, the dry weight of the root that absorbed ^{15}N.

$$A = \Delta^{15}N_w \times \frac{1}{W_r} \times \frac{60}{5} \qquad (2)$$

The increases in $^{15}N_s$ in each part of the ramet pair ($\Delta^{15}N_s$) were summed to obtain the increase

of ^{15}N in the whole ramet pair ($\Delta^{15}N_w$), and the percentage (%) of ^{15}N distributed in each part of

the ramet pair was calculated using equation (3). P, the percentage of ^{15}N distributed in each

part.

$$P = \frac{\Delta^{15}N_s}{\Delta^{15}N_t} \times 100 \qquad (3)$$

Reverse transcription-quantitative PCR analysis

Total RNA was extracted from frozen and ground tissues using the RNeasy Plant Mini Kit

(Qiagen, Hilden, Germany) with the RNase-Free-DNase Set (Qiagen) according to the

supplier's protocols. One μg of total RNA was used to synthesize cDNA using the ReverTra Ace

qPCR Master Mix (TOYOBO) according to the supplier's protocol. Twenty-ng cDNA was used

for each qPCR reaction with the KAPA SYBR FAST qPCR Master Mix (2X) (KAPA Biosystems,

Wilmington, MA, United States) and a real-time PCR system (Applied Biosystems QuantStudio

3). Expression levels were estimated using the relative quantification method (Livak and

Schmittgen, 2001) with *OITBC*, a homolog of TBC1 domain family member 22A (Maksup et al.,

2013), the internal standard for normalization. Gene locus IDs and the specific primers used for

amplification are listed in Supplemental Table S9.

Determination of free amino acids

Free amino acids were extracted as described by Konishi et al. (2014). Derivatization of amino

acids was carried out using the AccQ-Tag Ultra Derivatization Kit (Waters Corp., Milford, MA,

United States). The resulting AccQ-Tag-labeled derivatives were separated and quantified using

an HPLC System (Alliance 2695 HPLC system/2475, Waters Corp.) with the AccQTag Column

(3.9×150 mm, Waters Corp.) as described in the instruction manual.

RNA-Seq analysis

Libraries were prepared with the TruSeq Standard mRNA Library Prep Kit (Illumina, San Diego,

CA, United States) using 1 μg of total RNA. Sequence analysis of 40 M reads was performed

with a NovaSeq 6000 Sequencing System (Illumina). Library preparation and sequencing were

conducted by Macrogen Japan, Inc. (Tokyo). The sequencing reads were mapped to the *O.*

longistaminata genome obtained from the *Oryza longistaminata* Information Resource

(http://olinfres.nig.ac.jp/)(Reuscher et al., 2018) using HISAT2 (Kim et al., 2019), followed by

featureCounts (Liao et al., 2014) for counting reads and edgeR (Robinson et al., 2009) for

differential expression analysis. Low expression genes were removed using filterByExpr

function of edgeR with the default setting. We obtained the presumed function and MAPMAN

BIN code for each gene and the presumed corresponding gene ID of *O. sativa* from

Supplemental data 2 of Reuscher et al. (2018). Gene ontology (GO) analysis was performed

using the corresponding *O. sativa* gene ID list with the analysis tool agriGO

(http://systemsbiology.cau.edu.cn/agriGOv2/) (Du et al., 2010; Tian et al., 2017).

Growth analysis

Ramet pairs at a similar growth stage were first grown hydroponically in water for 4 days and

subsequently exposed to nitrogen split conditions for 5 weeks in a greenhouse using the

hydroponic culture solution with (+N split, 2.5 mM NH_4Cl) or without (-N split, 0 mM NH_4Cl)

nitrogen. The hydroponic solution was renewed every 3 to 4 days. Growth changes were

analyzed at 7-day intervals by measuring plant height, the number of fully expanded leaves, the

number of axillary buds growing more than 1 cm, and the chlorophyll content. The chlorophyll

content (SPAD value) was measured with a SPAD-502 Plus Chlorophyll Meter (Konica Minolta,

Tokyo, Japan) at the tip, middle, and basal parts of leaves, and the average was taken as the

SPAD value. For the controls, both ramets of each pair were fed 2.5 mM NH_4Cl (+N) or 0 mM

NH_4Cl (-N)-containing hydroponic solution for 5 weeks.

Phytohormone quantification

Cytokinins were extracted and semi-purified from about 100 mg fresh weight of root tissues as

described previously (Kojima et al., 2009). Cytokinins were quantified using an ultra-

performance liquid chromatography (UPLC)-tandem quadrupole mass spectrometer (ACQUITY

UPLC System/XEVO-TQXS; Waters Corp.) with an octadecylsilyl (ODS) column (ACQUITY

UPLC HSS T3, 1.8 μm, 2.1 mm × 100 mm, Waters Corp.; Kojima et al., 2009).

Peptide synthesis

OsCEP1a (DVRHypTNPGHSHypGIGH), OsCEP1b (DVRHypTNHypGHSHypGIGH), OsCEP1c

(GVRHypTNPGHSHypGIGH), and OsCEP1d (GVRHypTNHypGHSHypGIGH) were

synthesized on a CS 136X synthesizer (CS Bio, Menlo Park, CA, United States) using Fmoc

solid phase peptide synthesis chemistry. Hydroxyproline (Hyp) was introduced with Fmoc-

Hyp(Boc)-OH purchased from Watanabe Chemical Industries, Ltd. (Hiroshima, Japan). The

obtained crude peptides were purified by reverse-phase HPLC on Jasco, Inc. (Tokyo, Japan)

preparative instruments with a Jupiter C18 column (5 μm, 300 Å pore size, 21.2 mm internal

diameter × 250 mm) (Phenomenex, Torrance, CA, United States), and lyophilized to yield pure

peptides.

Treatment with synthesized CEP1 peptides

Ramet pairs at a similar growth stage were first grown in hydroponic solution for 3 days in a

growth chamber (LPH-411PFQDT-SPC, NK System). Root tips (about 3 cm) from one ramet

were removed with a razor blade, and water on the root surface was wiped off with a Kim Towel

(Nippon Paper Cresia, Tokyo Japan). The roots were immediately submerged into 100 mL of an

ammonium solution containing CEP peptide (+CEPs local: 2.5 mM NH_4Cl, 30 μM of each

CEP1a-d). Simultaneously, the root on the other side was submerged in the ammonium solution

(systemic: 2.5 mM NH_4Cl) for the indicated period. For the mock control, the roots of each ramet

pair were treated with 2.5 mM NH_4Cl.

Statistical analysis

Graphs represent the means ± SE of biological-independent experiments. The statistical

significance was assayed using means a two-tailed Student's *t*-test or Tukey's honestly

significant difference test ($p < 0.05$). The choice of test and the replicate numbers are provided

in the corresponding figure legend.

Accession numbers

RNA-seq data were deposited to the Gene Expression Omnibus

(https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE182486.

Supplemental data

The following supplemental materials are available.

Supplemental Figure S1. The split hydroponic experimental system for *O. longistaminata*

ramet pairs.

Supplemental Figure S2. A Venn diagram showing the overlap among genes up-regulated in

the -N treatment compared to the +N treatment, those up-regulated in the -N split treatment

compared to +N treatment, and those up-regulated in the +N treatment compared to +N split

treatment.

Supplemental Figure S3. Transcript abundance of *OlNADH-GOGAT1* in the roots of ramet

pairs after a 24-h split treatment as measured by RT-qPCR.

Supplemental Table S1. Genes up-regulated in the +N split treatment compared to the +N

treatment

Supplemental Table S2. Genes up-regulated in the +N treatment compared to the -N treatment

Supplemental Table S3. Genes up-regulated in the +N treatment compared to the -N split

treatment

Supplemental Table S4. Genes up-regulated in the -N treatment compared to the +N treatment

Supplemental Table S5. Genes up-regulated in the -N split treatment compared to the +N

treatment

Supplemental Table S6. Genes up-regulated in the +N treatment compared to the +N split

treatment

526 **Acknowledgments**

527 We are grateful to Drs. Motoyuki Ashikari, Tomoyuki Furuta, Takatoshi Kiba, Fanny Bellegarde,

528 Mimi Hashimoto, Shoji Segami (Nagoya University), and Drs. Shinjiro Yamaguchi and Kiyoshi

529 Mashiguchi (Kyoto University) for their helpful support and discussion.

 Figure Legends

Figure 1. Compensatory up-regulation of ammonium ion uptake in response to spatially

heterogeneous nitrogen availability. A, A schematic representation of the split hydroponic

experiment system in which the roots of a ramet pair were separately exposed to independent

nitrogen conditions. Details of the experimental design are shown in Fig. S1. B, Compensatory

uptake of ammonium ion in +N split roots compared to +N. Ammonium ion uptake activity was

measured using $^{15}NH_4^+$ as the tracer. Error bars represent SE of values for biological replicates

(n = 3). *$p < 0.05$ (Student's t-test). DW, dry weight. C, Accumulation pattern of *OlAMT1;2*

transcripts in the roots of *O. longistaminata* ramet pairs after a 24-h split treatment. Transcript

abundance, normalized to *OlTBC*, is expressed relative to that of +N, defined as 1. Error bars

represent SE of values for biological replicates (n = 3 or 4). Different lowercase letters at the top

of each column denote statistically significant differences by Tukey's honestly significant

difference test (HSD) ($p < 0.05$).

Figure 2. Transcriptome changes in the roots of ramet pairs in response to spatially

heterogeneous nitrogen availability. A, A volcano plot showing differentially expressed genes

(FDR < 0.05) between +N and +N split roots. Significantly up-regulated or down-regulated

genes in roots from the +N split treatment are shown as red or blue dots, respectively. Numbers

in red or blue indicate the total number of up-regulated or down-regulated genes. B, A Venn

diagram showing the overlap among the genes up-regulated in the +N split treatment compared

to the +N treatment (Up in +N split to +N), those up-regulated in the +N treatment compared to

the -N treatment (Up in +N to -N), and those up-regulated in the +N treatment compared to the -

N split treatment (Up in +N to -N split). C, The top 20 enriched GO categories among genes up-

regulated in the +N split treatment compared to the +N treatment. Highly redundant GO

categories were manually removed.

Figure 3. Metabolic responses to spatially heterogeneous nitrogen availability in the roots of *O.*

longistaminata ramet pairs. A, The expression profile of genes involved in nitrogen assimilation,

glycolysis and amino acid biosynthesis. The heatmaps in the pathways were deduced from

RNAseq data. Fru1,6-BP, fructose 1,6-bisphosphate; PEP, phospho*enol*-pyruvate; OAA,

oxaloacetate; 2OG. 2-oxoglutarate. The details of gene names are provided in Supplemental

Table S1. B, Expression patterns of genes analyzed by RT-qPCR. The expression level of each

gene, normalized to *OlTBC*, is expressed relative to that of the +N treatment defined as 1. C,

Amino acid quantification. The concentrations of amino acids in the roots of each ramet pair

after 48 h of split treatment are shown in histograms. Error bars represent the SE of values for

biological replicates (n = 3 or 4 for RT-qPCR, and n = 4 for amino acid analysis). Different

lowercase letters at the top of each column denote statistically significant differences by Tukey's

honestly significant difference test (HSD) ($p < 0.05$).

Figure 4. Allocation of absorbed nitrogen and growth profiles of ramet pairs. A, Distribution of

^{15}N in roots and shoots 7 d after ^{15}NH$_4$Cl feeding. One of the ramets in each split-treated ramet

pair was treated with 2.5 mM ^{15}NH$_4$Cl for 5 min and then returned to the normal split treatment.

Error bars represent SE of values for biological replicates (n = 3). B to E, Changes in plant

height (B, ΔPlant height), number of leaves (C, ΔLeaf number), chlorophyll content (D, ΔSPAD

index), and number of growing axillary buds (E, ΔGrowing axl. bud no.) in each ramet after 5

weeks of split treatment. Error bars indicate the SE. Different lowercase letters at the top of

each column denote statistically significant differences by Tukey's honestly significant difference

test (HSD) ($p < 0.05$). n = 6 to 12.

Figure 5. Response of cytokinin and strigolactone biosynthesis to heterogeneous nitrogen

conditions. A, Relative expression level of cytokinin biosynthetic genes in ramet roots in

response to different nitrogen treatments. Transcript abundance in roots of each ramet after 1

week of the split treatment was analyzed by RT-qPCR. B, Cytokinin concentration in the roots of

each ramet after 1 week of the split treatment. The complete data set is presented in

Supplemental Table S7. tZ, *trans*-zeatin; tZR, tZ riboside; tZRPs, tZR 5'-phosphates; cZ, *cis*-

zeatin; cZR, cZ riboside; cZRPs, cZR 5'-phosphates; iP, N^6-(Δ^2-isopentenyl)adenine; iPR, iP

riboside; iPRPs, iPR 5'-phosphates; and FW, fresh weight. C, Transcript abundance of

strigolactone biosynthetic genes in ramet roots in response to different nitrogen conditions. In

(A) and (C), the expression level of each gene normalized by *TBC* is expressed relative to that

of the +N treatment defined as 1. Error bars represent the SE of values for biological replicates

(n = 4). Different lowercase letters at the top of each column denote statistically significant

differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$).

Figure 6. A, Expression pattern of the *O. longistaminata* ortholog of *OsCEP1* (*OlCEP1*) in the

roots of ramet pairs after a 24-h split treatment. The expression level of each gene normalized

by *TBC* is expressed relative to that of the +N treatment defined as 1. Error bars represent SE

of values for biological replicates (n = 3 or 4). Different lowercase letters at the top of each

column denote statistically significant differences by Tukey's honestly significant difference test

(HSD) ($p < 0.05$). B, Expression of *OlAMT1;2* and *OlGS1;2* genes in the ramet roots in

response to exogenously supplied CEP1 peptides. The CEP1 peptide mix was applied to the

roots of one of the ramet pairs, and gene expression in the local (+CEP local) and systemic side

(systemic) roots after 6 and 24 h treatment was measured. The expression level of each gene,

normalized by *TBC*, is expressed relative to that of the +N mock treatment defined as 1. Error

bars represent SE of values for biological replicates (n = 3 or 4). *$p < 0.05$ (Student's t-test)

compared to the corresponding mock treatment.

LITERATURE CITED

Alder A, Jamil M, Marzorati M, Bruno M, Vermathen M, Bigler P, Ghisla S, Bouwmeester H,
Beyer P, Al-Babili S (2012) The path from β-carotene to carlactone, a strigolactone-like
plant hormone. Science 335: 1348–1351

Bessho-Uehara K, Nugroho JE, Kondo H, Angeles-Shim RB, Ashikari M (2018) Sucrose affects
the developmental transition of rhizomes in Oryza longistaminata. J Plant Res 131: 693–
707

Chandran AKN, Priatama RA, Kumar V, Xuan Y, Je B II, Kim CM, Jung KH, Han C deok (2016)
Genome-wide transcriptome analysis of expression in rice seedling roots in response to
supplemental nitrogen. J Plant Physiol 200: 62–75

Davière J-M, Achard P (2017) Organ communication: Cytokinins on the move. Nat Plants 3:
17116

Delay C, Imin N, Djordjevic MA (2013) CEP genes regulate root and shoot development in
response to environmental cues and are specific to seed plants. J Exp Bot 64: 5383–5394

Du Z, Zhou X, Ling Y, Zhang Z, Su Z (2010) agriGO: A GO analysis toolkit for the agricultural
community. Nucleic Acids Res 38: 64–70

Fan Z, Cai Z, Shan J, Yang J (2017) Letter to the Editor: Bud Position and Carbohydrate Play a
More Significant Role than Light Condition in the Developmental Transition between
Rhizome Buds and Aerial Shoot Buds of Oryza longistaminata. Plant Cell Physiol 58:
1281–1282

Funayama K, Kojima S, Tabuchi-Kobayashi M, Sawa Y, Nakayama Y, Hayakawa T, Yamaya T
(2013) Cytosolic glutamine synthetase1;2 is responsible for the primary assimilation of
ammonium in rice roots. Plant Cell Physiol 54: 934–943

Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pagès V, Dun EA, Pillot JP, Letisse F,
Matusova R, Danoun S, Portais JC, et al (2008) Strigolactone inhibition of shoot
branching. Nature 455: 189–194

Guo L, Plunkert M, Luo X, Liu Z (2021) Developmental regulation of stolon and rhizome. Curr
Opin Plant Biol 59: 101970

Ishiyama K, Inoue E, Tabuchi M, Yamaya T, Takahashi H (2004) Biochemical background and
compartmentalized functions of cytosolic glutamine synthetase for active ammonium
assimilation in rice roots. Plant Cell Physiol 45: 1640–1647

Ji Y, Li Q, Liu G, Selvaraj G, Zheng Z, Zou J, Wei Y (2019) Roles of Cytosolic Glutamine
Synthetases in Arabidopsis Development and Stress Responses. Plant Cell Physiol 60:
657–671

Jia Z, von Wirén N (2020) Signaling pathways underlying nitrogen-dependent changes in root
system architecture: from model to crop species. J Exp Bot 71: 4393–4404

Kamada-Nobusada T, Makita N, Kojima M, Sakakibara H (2013) Nitrogen-dependent regulation
of de novo cytokinin biosynthesis in rice: The role of glutamine metabolism as an
additional signal. Plant Cell Physiol 54: 1881–1893

Kiba T, Takei K, Kojima M, Sakakibara H (2013) Side-Chain Modification of Cytokinins Controls
Shoot Growth in Arabidopsis. Dev Cell 27: 452–461

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37: 907–915

Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M, Ashikari M,
Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al (2009) Highly sensitive and high-
throughput analysis of plant hormones using ms-probe modification and liquid
chromatographytandem mass spectrometry: An application for hormone profiling in oryza
sativa. Plant Cell Physiol 50: 1201–1214

De Kroon H, Van Der Zalm E, Van Rheenen JWA, Van Dijk A, Kreulen R (1998) The interaction
between water and nitrogen translocation in a rhizomatous sedge (Carex flacca).
Oecologia 116: 38–49

Kyozuka J (2017) Letter to the Editor: Author Response - Analysis of Rhizome Development in
Oryza longistaminata, a Wild Rice Species. Plant Cell Physiol 58: 1283

Landrein B, Formosa-Jordan P, Malivert A, Schuster C, Melnyk CW, Yang W, Turnbull C,
Meyerowitz EM, Locke JCW, Jönsson H (2018) Nitrate modulates stem cell dynamics in
Arabidopsis shoot meristems through cytokinins. Proc Natl Acad Sci U S A 115: 1382–
1387

Lea PJ, Miflin BJ (1974) Alternative route for nitrogen assimilation in higher plants. 251: 614–
616

Lee S, Marmagne A, Park J, Fabien C, Yim Y, Kim S ji, Kim TH, Lim PO, Masclaux-Daubresse
C, Nam HG (2020a) Concurrent activation of OsAMT1;2 and OsGOGAT1 in rice leads to
enhanced nitrogen use efficiency under nitrogen limitation. Plant J 103: 7–20

Lee S, Park J, Lee J, Shin D, Marmagne A, Lim PO, Masclaux-Daubresse C, An G, Nam HG
(2020b) OsASN1 overexpression in rice increases grain protein content and yield under
nitrogen-limiting conditions. Plant Cell Physiol 61: 1309–1320

Li C, Tang Z, Wei J, Qu H, Xie Y, Xu G (2016) The OsAMT1.1 gene functions in ammonium
uptake and ammonium–potassium homeostasis over low and high ammonium
concentration ranges. J Genet Genomics 43: 639–649

Liao Y, Smyth GK, Shi W (2014) FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30: 923–930

Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2-$\Delta\Delta CT$ method. Methods 25: 402–408

Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H,
Yanagisawa S (2018) A NIGT1-centred transcriptional cascade regulates nitrate signalling
and incorporates phosphorus starvation signals in Arabidopsis. Nat Commun 9: 1376

Maksup S, Supaibulwatana K, Selvaraj G (2013) High-quality reference genes for quantifying
the transcriptional responses of Oryza sativa L. (ssp. indica and japonica) to abiotic stress
conditions. Chinese Sci Bull 58: 1919–1930

Minakuchi K, Kameoka H, Yasuno N, Umehara M, Luo L, Kobayashi K, Hanada A, Ueno K,
Asami T, Yamaguchi S, et al (2010) FINE CULM1 (FC1) works downstream of
strigolactones to inhibit the outgrowth of axillary buds in rice. Plant Cell Physiol 51: 1127–
1135

Naulin PA, Armijo GI, Vega AS, Tamayo KP, Gras DE, De La Cruz J, Gutiérrez RA (2019)
Nitrate Induction of Primary Root Growth Requires Cytokinin Signaling in Arabidopsis
thaliana. Plant Cell Physiol 61: 342–352

Nishida H, Suzaki T (2018) Two Negative Regulatory Systems of Root Nodule Symbiosis: How
Are Symbiotic Benefits and Costs Balanced? Plant Cell Physiol 59: 1733–1738

Ohashi M, Ishiyama K, Kojima S, Kojima M, Sakakibara H, Yamaya T, Hayakawa T (2017) Lack
of cytosolic glutamine synthetase1;2 activity reduces nitrogen-dependent biosynthesis of
cytokinin required for axillary bud outgrowth in rice seedlings. Plant Cell Physiol 58: 679–
690

Ohkubo Y, Kuwata K, Matsubayashi Y (2021) A type 2C protein phosphatase activates high-
affinity nitrate uptake by dephosphorylating NRT2.1. Nat Plants 7: 310–316

Ohkubo Y, Tanaka M, Tabata R, Ogawa-Ohnishi M, Matsubayashi Y (2017) Shoot-to-root
mobile polypeptides involved in systemic regulation of nitrogen acquisition. Nat Plants 3:
17029

Ohyama K, Ogawa M, Matsubayashi Y (2008) Identification of a biologically active, small,
secreted peptide in Arabidopsis by in silico gene screening, followed by LC-MS-based
structure analysis. Plant J 55: 152–160

Okamoto S, Tabata R, Matsubayashi Y (2016) Long-distance peptide signaling essential for
nutrient homeostasis in plants. Curr Opin Plant Biol 34: 35–40

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H (2017) Systemic transport of
trans-zeatin and its precursor have differing roles in Arabidopsis shoots. Nat Plants. doi:
10.1038/nplants.2017.112

Ota R, Ohkubo Y, Yamashita Y, Ogawa-Ohnishi M, Matsubayashi Y (2020) Shoot-to-root
mobile CEPD-like 2 integrates shoot nitrogen status to systemically regulate nitrate uptake
in Arabidopsis. Nat Commun 11: 641

Ravazzolo L, Boutet-Mercey S, Perreau F, Forestan C, Varotto S, Ruperti B, Quaggiotti S

(2021) Strigolactones and Auxin Cooperate to Regulate Maize Root Development and

Response to Nitrate. Plant Cell Physiol 0: 1–14

Ravazzolo L, Trevisan S, Manoli A, Boutet-Mercey S, Perreau F, Quaggiotti S (2019) The

Control of Zealactone Biosynthesis and Exudation is Involved in the Response to Nitrogen

in Maize Root. Plant Cell Physiol 60: 2100–2112

Reuscher S, Furuta T, Bessho-Uehara K, Cosi M, Jena KK, Toyoda A, Fujiyama A, Kurata N,

Ashikari M (2018) Assembling the genome of the African wild rice Oryza longistaminata by

exploiting synteny in closely related Oryza species. Commun Biol 1: 162

Robinson MD, McCarthy DJ, Smyth GK (2009) edgeR: A Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics 26: 139–140

Ruffel S (2018) Nutrient-Related Long-Distance Signals: Common Players and Possible Cross-

Talk. Plant Cell Physiol 59: 1723–1732

Ruffel S, Gojon A (2017) Systemic nutrient signalling: On the road for nitrate. Nat Plants 3: 1–2

Sakakibara H (2021) Cytokinin biosynthesis and transport for systemic nitrogen signaling. Plant

J 105: 421–430

Sakakibara H, Takei K, Hirose N (2006) Interactions between nitrogen and cytokinin in the

regulation of metabolism and development. Trends Plant Sci 11: 440–448

Scheible WR, González-Fontes A, Lauerer M, Müller-Röber B, Caboche M, Stitt M (1997)

Nitrate acts as a signal to induce organic acid metabolism and repress starch metabolism

in tobacco. Plant Cell 9: 783–798

Scheible WR, Morcuende R, Czechowski T, Fritz C, Osuna D, Palacios-Rojas N, Schindelasch

D, Thimm O, Udvardi MK, Stitt M (2004) Genome-wide reprogramming of primary and

secondary metabolism, protein synthesis, cellular growth processes, and the regulatory

infrastructure of arabidopsis in response to nitrogen. Plant Physiol 136: 2483–2499

Shibasaki K, Takebayashi A, Makita N, Kojima M, Takebayashi Y, Kawai M, Hachiya T,

Sakakibara H (2021) Nitrogen Nutrition Promotes Rhizome Bud Outgrowth via Regulation

of Cytokinin Biosynthesis Genes and an Oryza longistaminata Ortholog of FINE CULM 1.

Front Plant Sci 12: 670101

Sonoda Y, Ikeda A, Saiki S, Von Wirén N, Yamaya T, Yamaguchi J (2003) Distinct expression

and function of three ammonium transporter genes (OsAMT1;1 - 1;3) in rice. Plant Cell

Physiol 44: 726–734

Suenaga A, Moriya K, Sonoda Y, Ikeda A, Von Wirén N, Hayakawa T, Yamaguchi J, Yamaya T

(2003) Constitutive expression of a novel-type ammonium transporter OsAMT2 in rice

plants. Plant Cell Physiol 44: 206–211

Sui Z, Wang T, Li H, Zhang M, Li Y, Xu R, Xing G, Ni Z, Xin M (2016) Overexpression of
peptide-encoding OsCEP6.1 results in pleiotropic effects on growth in rice (O. sativa).
Front Plant Sci 7: 1–12

Sun H, Tao J, Liu S, Huang S, Chen S, Xie X, Yoneyama K, Zhang Y, Xu G (2014)
Strigolactones are involved in phosphate- and nitrate-deficiency-induced root development
and auxin transport in rice. J Exp Bot 65: 6735–6746

Sun L, Di DW, Li G, Li Y, Kronzucker HJ, Shi W (2020) Transcriptome analysis of rice (Oryza
sativa L.) in response to ammonium resupply reveals the involvement of phytohormone
signaling and the transcription factor OsJAZ9 in reprogramming of nitrogen uptake and
metabolism. J Plant Physiol 246–247: 153137

Tabata R, Sumida K, Yoshii T, Ohyama K, Shinohara H, Matsubayashi Y (2014) Perception of
root-derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. Science
(80-) 346: 343–346

Tabuchi M, Abiko T, Yamaya T (2007) Assimilation of ammonium ions and reutilization of
nitrogen in rice (Oryza sativa L.). J Exp Bot 58: 2319–2327

Tabuchi M, Sugiyama K, Ishiyama K, Inoue E, Sato T, Takahashi H, Yamaya T (2005) Severe
reduction in growth rate and grain filling of rice mutants lacking OsGS1;1, a cytosolic
glutamine synthetase1;1. Plant J 42: 641–651

Takei K, Ueda N, Aoki K, Kuromori T, Hirayama T, Shinozaki K, Yamaya T, Sakakibara H
(2004) AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in
Arabidopsis. Plant Cell Physiol 45: 1053–1062

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z (2017) AgriGO v2.0: A GO analysis toolkit
for the agricultural community, 2017 update. Nucleic Acids Res 45: W122–W129

Toriba T, Tokunaga H, Nagasawa K, Nie F, Yoshida A, Kyozuka J (2020) Suppression of Leaf
Blade Development by BLADE-ON-PETIOLE Orthologs Is a Common Strategy for
Underground Rhizome Growth. Curr Biol 30: 509-516.e3

Umehara M, Hanada A, Magome H, Takeda-Kamiya N, Yamaguchi S (2010) Contribution of
strigolactones to the inhibition of tiller bud outgrowth under phosphate deficiency in rice.
Plant Cell Physiol 51: 1118–1126

Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya
Y, Shirasu K, Yoneyama K, et al (2008) Inhibition of shoot branching by new terpenoid
plant hormones. Nature 455: 195–200

Vaughan DA (1994) The Wild Relatives of Rice: A Genetic Resources Handbook. IRRI, Los
Banos

Wang R, Tischner R, Gutiérrez RA, Hoffman M, Xing X, Chen M, Coruzzi G, Crawford NM
(2004) Genomic analysis of the nitrate response using a nitrate reductase-null mutant of
arabidopsis. Plant Physiol 136: 2512–2522

Wheeldon CD, Bennett T (2021) There and back again: An evolutionary perspective on long-
distance coordination of plant growth and development. Semin Cell Dev Biol 109: 55–67

Xiong D, Yu T, Ling X, Fahad S, Peng S, Li Y, Huang J (2015) Sufficient leaf transpiration and
nonstructural carbohydrates are beneficial for high-temperature tolerance in three rice
(Oryza sativa) cultivars and two nitrogen treatments. Funct Plant Biol 42: 347–356

Xu J, Zha M, Li Y, Ding Y, Chen L, Ding C, Wang S (2015) The interaction between nitrogen
availability and auxin, cytokinin, and strigolactone in the control of shoot branching in rice
(Oryza sativa L.). Plant Cell Rep 34: 1647–1662

Yamaya T, Kusano M (2014) Evidence supporting distinct functions of three cytosolic glutamine
synthetases and two NADH-glutamate synthases in rice. J Exp Bot 65: 5519–5525

Yang HC, Kan CC, Hung TH, Hsieh PH, Wang SY, Hsieh WY, Hsieh MH (2017) Identification of
early ammonium nitrate-responsive genes in rice roots. Sci Rep 7: 1–16

Yoshida A, Terada Y, Toriba T, Kose K, Ashikari M, Kyozuka J (2016) Analysis of rhizome
development in oryza longistaminata, a wild rice species. Plant Cell Physiol 57: 2213–
2220

Yoshida S (1981) Mineral nutrition of rice. Fundam. rice Crop Sci. International Rice Research
Institute, Los Banos, Philippines, pp 111–176

Yuan L, Loqué D, Kojima S, Rauch S, Ishiyama K, Inoue E, Takahashi H, Wirén, von N (2007)
The organization of high-affinity ammonium uptake in Arabidopsis roots depends on the
spatial arrangement and biochemical properties of AMT1-type transporters. Plant Cell 19:
2636–2652

Figure 1. Compensatory up-regulation of ammonium ion uptake in response to spatially heterogeneous nitrogen availability. A, A schematic representation of the split hydroponic experiment system in which the roots of a ramet pair were separately exposed to independent nitrogen conditions. Details of the experimental design are shown in Fig. S1. B, Compensatory uptake of ammonium ion in +N split roots compared to +N. Ammonium ion uptake activity was measured using $^{15}NH_4^+$ as the tracer. Error bars represent SE of values for biological replicates (n = 3). *$p < 0.05$ (Student's t-test). DW, dry weight. C, Accumulation pattern of *OlAMT1;2* transcripts in the roots of *O. longistaminata* ramet pairs after a 24-h split treatment. Transcript abundance, normalized to *OlTBC*, is expressed relative to that of +N, defined as 1. Error bars represent SE of values for biological replicates (n = 3 or 4). Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$).

Figure 2. Transcriptome changes in the roots of ramet pairs in response to spatially heterogeneous nitrogen availability. A, A volcano plot showing differentially expressed genes (FDR < 0.05) between +N and +N split roots. Significantly up-regulated or down-regulated genes in roots from the +N split treatment are shown as red or blue dots, respectively. Numbers in red or blue indicate the total number of up-regulated or down-regulated genes. B, A Venn diagram showing the overlap among the genes up-regulated in the +N split treatment compared to the +N treatment (Up in +N split to +N), those up-regulated in the +N treatment compared to the -N treatment (Up in +N to -N), and those up-regulated in the +N treatment compared to the -N split treatment (Up in +N to -N split). C, The top 20 enriched GO categories among genes up-regulated in the +N split treatment compared to the +N treatment. Highly redundant GO categories were manually removed.

Figure 3. Metabolic responses to spatially heterogeneous nitrogen availability in the roots of *O. longistaminata* ramet pairs. A, The expression profile of genes involved in nitrogen assimilation, glycolysis and amino acid biosynthesis. The heatmaps in the pathways were deduced from RNAseq data. Fru1,6-BP, fructose 1,6-bisphosphate; PEP, phospho*enol*-pyruvate; OAA, oxaloacetate; 2OG. 2-oxoglutarate. The details of gene names are provided in Supplemental Table S1. B, Expression patterns of genes analyzed by RT-qPCR. The expression level of each gene, normalized to *OlTBC*, is expressed relative to that of the +N treatment defined as 1. C, Amino acid quantification. The concentrations of amino acids in the roots of each ramet pair after 48 h of split treatment are shown in histograms. Error bars represent the SE of values for biological replicates (n = 3 or 4 for RT-qPCR, and n = 4 for amino acid analysis). Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$).

Figure 4. Allocation of absorbed nitrogen and growth profiles of ramet pairs. A, Distribution of ^{15}N in roots and shoots 7 d after $^{15}NH_4Cl$ feeding. One of the ramets in each split-treated ramet pair was treated with 2.5 mM $^{15}NH_4Cl$ for 5 min and then returned to the normal split treatment. Error bars represent SE of values for biological replicates (n = 3). B to E, Changes in plant height (B, ΔPlant height), number of leaves (C, ΔLeaf number), chlorophyll content (D, ΔSPAD index), and number of growing axillary buds (E, ΔGrowing axl. bud no.) in each ramet after 5 weeks of split treatment. Error bars indicate the SE. Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$). n = 6 to 12.

Figure 5. Response of cytokinin and strigolactone biosynthesis to heterogeneous nitrogen conditions. A, Relative expression level of cytokinin biosynthetic genes in ramet roots in response to different nitrogen treatments. Transcript abundance in roots of each ramet after 1 week of the split treatment was analyzed by RT-qPCR. B, Cytokinin concentration in the roots of each ramet after 1 week of the split treatment. The complete data set is presented in Supplemental Table S8. tZ, *trans*-zeatin; tZR, tZ riboside; tZRPs, tZR 5'-phosphates; cZ, *cis*-zeatin; cZR, cZ riboside; cZRPs, cZ 5'-phosphates; iP, N^6-(Δ^2-isopentenyl)adenine; iPR, iP riboside; iPRPs, iPR 5'-phosphates; and FW, fresh weight. C, Transcript abundance of strigolactone biosynthetic genes in ramet roots in response to different nitrogen conditions. In (A) and (C), the expression level of each gene normalized by *TBC* is expressed relative to that of the +N treatment defined as 1. Error bars represent the SE of values for biological replicates (n = 4). Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$).

Figure 6. A, Expression pattern of the *O. longistaminata* ortholog of *OsCEP1* (*OlCEP1*) in the roots of ramet pairs after a 24-h split treatment. The expression level of each gene normalized by *TBC* is expressed relative to that of the +N treatment defined as 1. Error bars represent SE of values for biological replicates (n = 3 or 4). Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$). B, Expression of *OlAMT1;2* and *OlGS1;2* genes in the ramet roots in response to exogenously supplied CEP1 peptides. The CEP1 peptide mix was applied to the roots of one of the ramet pairs, and gene expression in the local (+CEP local) and systemic side (systemic) roots after 6 and 24 h treatment was measured. The expression level of each gene, normalized by *TBC*, is expressed relative to that of the +N mock treatment defined as 1. Error bars represent SE of values for biological replicates (n = 3 or 4). *$p < 0.05$ (Student's t-test) compared to the corresponding mock treatment.

Supplemental Figure S1. The split hydroponic experimental system for *O. longistaminata* ramet pairs. A, *O. longistaminata* hydroponic culture. The plants were maintained in a 10 L-container. Scale bar, 10 cm. B, A clonal colony of *O. longistaminata*. Scale bar, 10 cm. C, A rhizome grown in the hydroponic culture. Ovals identify adjacent nodes. The tip of a primary rhizome was cut at the yellow line to induce the outgrowth of secondary rhizomes. D, Growth of secondary rhizomes at adjacent nodes (arrowheads). The pair was excised from the parental colony at the yellow line for further growth. E, A ramet pair grown from adjacent rhizome nodes. Scale bar, 15 cm. F, Split treatments with different nitrogen conditions. Each pot contains 1 L of nutrient solution.

Supplemental Figure S2. A Venn diagram showing the overlap among genes up-regulated in the -N treatment compared to the +N treatment (Up in -N to +N), those up-regulated in the -N split treatment compared to +N treatment (Up in -N split to +N), and those up-regulated in the +N treatment compared to +N split treatment (Up in +N to +N split).

Supplemental Figure S3. Transcript abundance of *OlNADH-GOGAT1* in the roots of ramet pairs after a 24-h split treatment as measured by RT-qPCR. The expression level of each gene, normalized by *TBC*, is expressed relative to that of the +N treatment defined as 1. Error bars represent SE of values for biological replicates (n = 3 or 4). Different lowercase letters at the top of each column denote statistically significant differences by Tukey's honestly significant difference test (HSD) ($p < 0.05$).