Sensory Modality and Spoken Language Shape Reading Network in

Blind Readers of Braille

Mengyu Tian[a,1], Elizabeth J. Saccone[a], Judy S. Kim[a,b], Shipra Kanjlia[a,c], Marina Bedny[a]

[a] Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, Maryland 21218

[b] Department of Psychology, Yale University, New Haven, Connecticut 06511

[c] Department of Psychology, Carnegie Mellon University, Pittsburgh, PA 15213

[1]To whom correspondence may be addressed: Email: mengyutian@jhu.edu

Author Contributions: J.K., S.K., and M.B. designed research; J.K., S.K., and M.B. performed research;

M.T. analyzed data; M.T., E.S. and M.B. wrote the paper.

Competing Interest Statement: The authors declare no competing interests.

Running title： Reading Modality Shapes Reading Network

Abstract

The neural basis of reading is highly consistent across many languages and scripts. Are there alternative

neural routes to reading? How does the sensory modality of symbols (tactile vs. visual) influence their

neural representations? We examined these questions by comparing reading of visual print (sighted

group, n=19) and tactile Braille (congenitally blind group, n=19). Blind and sighted readers were

presented with written (words, consonant strings, non-letter shapes) and spoken stimuli (words, backward

speech) that varied in word-likeness. Consistent with prior work, the ventral occipitotemporal cortex

(vOTC) was active during Braille and visual reading. A posterior/anterior vOTC word-form gradient was

observed only in sighted readers with more anterior regions preferring larger orthographic units (words).

No such gradient was observed in blind readers. Consistent with connectivity predictions, in blind

compared to sighted readers, posterior parietal cortices were recruited to a greater degree and contained

word-preferring patches. Lateralization of Braille in blind readers was predicted by laterality of spoken

language and reading hand. The effect of spoken language increased along a cortical hierarchy, whereas

effect of reading-hand waned. These results suggested that the neural basis of reading is influenced by

symbol modality and spoken language and support connectivity-based views of cortical function.

Keywords: Braille reading; blindness; plasticity; posterior parietal cortex; VWFA

Introduction

Written language is among the most impressive human cultural achievements. The capacity to record and

transmit information over space and time has enabled the accumulation of scientific, technological, and

historical knowledge across generations and continents. How does the human brain accommodate this

cultural invention, which emerged only approximately 5,000 years ago?

Despite being a recent cultural invention, the neural basis of reading is highly consistent across a

variety of languages and visual scripts, including alphabetic, logographic (e.g., Chinese), and syllabic

writing systems (e.g., Japanese Kana) (Bolger et al. 2005; Hu et al. 2010; Nakamura et al. 2012; Rueckl

et al. 2015; Krafnick et al. 2016; Feng et al. 2020; Tian et al. 2020). All of these reading systems engage

regions within the left lateral ventral occipitotemporal cortex (vOTC). A region in the left lateral vOTC has

been termed the 'visual word form area' (VWFA) because of its preferential response to written words and

letter combinations over non-linguistic visual stimuli and speech (Cohen et al. 2000; Baker et al. 2007;

Dehaene et al. 2010; Dehaene and Cohen 2011). The VWFA is situated within a posterior/anterior

processing gradient. During reading, visual symbols are first processed by early visual cortices and

posterior portions of vOTC, which represent simple visual features (e.g., line junctions) (Dehaene et al.

2005; DiCarlo and Cox 2007). By contrast, the middle and anterior potions of lateral vOTC are specialized

for progressively larger orthographic units, from written letters, letter combinations/bigrams, and finally

whole words (Dehaene et al. 2005; Binder et al. 2006; Vinckier et al. 2007).

An open question is whether the vOTC's posterior/anterior processing stream is the only way for

the brain to implement reading and, relatedly, why the neural basis of reading takes this particular form.

Comparing tactile Braille reading among congenitally blind individuals to print reading in sighted people

offers unique insights into the causal mechanisms that determine the neural basis of reading. Braille is

read by passing the fingers along raised dot patterns, with each Braille character a three-rows-by-two-

columns dot matrix (Millar 2003). This distinctive tactile reading system provides insight into whether and

how the sensory modality of symbols influences their neural representations. Do blind readers of Braille

likewise show an anterior/posterior orthographic gradient in the vOTC?

A prominent view holds that, past the initial sensory entry points in V1 (print) and S1 (Braille),

reading depends on the same neural mechanisms in blind Braille readers and sighted print readers

(Büchel et al. 1998; Reich et al. 2011; Debowska et al. 2016; Rączy et al. 2019). This view is based on

the observation that reading Braille elicits activation in the anatomical location of the 'VWFA' (Reich et al.

2011; Debowska et al. 2016; Siuda-Krzywicka et al. 2016; Bola et al. 2019; Rączy et al. 2019; Dzięgiel-

Fivet et al. 2021). At the same time, no prior study to our knowledge has tested for the presence of an

orthographic gradient in vOTC of blind readers. The available work has only looked at whether the

anatomical location of the VWFA responds to Braille. Unlike visual print, Braille does not enter the vOTC

from V1, but rather originates in primary somatosensory cortex (S1). Does a posterior/anterior

orthographic gradient emerge despite this difference in neural entry points?

The first goal of the current study was to test whether sighted print and blind Braille readers

recruit a similar posterior/anterior orthographic gradient within vOTC. To answer this question, we

compared responses during Braille (blind) and visual print (sighted) reading of analogous written stimuli of

different orthographic richness (words, consonant strings, and unfamiliar shapes (control)). The same

participants were also presented with spoken words and backward speech control stimuli, to enable

comparison of written and spoken language processing.

An alternative, though not a mutually exclusive possibility, is that regions closer to somatosensory

cortices, in parietal cortex, play a special role in Braille reading. In sighted readers, the vOTC is the

culmination of the visual object recognition stream and occupies a key connectivity position between

visual input on the one hand and linguistic representations on the other (Yeatman et al. 2013; Bouhali et

al. 2014; Hannagan et al. 2015; Saygin et al. 2016; Stevens et al. 2017; Barttfeld et al. 2018; Li et al.

2020). In the case of tactile reading, posterior parietal cortices (PPC) arguably occupy an analogous

connectivity-based position for blind readers of Braille. The PPC lies adjacent and posterior to early

somatosensory cortex (SMC) and plays a key role in tactile shape and texture perception and higher

order tactile processing (Hegner et al. 2010; Bauer et al. 2015). Like the 'VWFA', PPC is anatomically

connected to language and working memory systems (Duhamel et al. 1998; Lewis and Van Essen 2000;

Kaas 2012; Ruschel et al. 2014; Burks et al. 2017). Whether the PPC of proficient blind Braille readers

contains Braille specialization, akin to specialization for visual letters and words found in vOTC of sighted

readers, has not yet been systematically investigated.

Several previous neuroimaging studies have documented PPC activity during Braille reading tasks but have not explored whether these responses are Braille-specific or distinct from what is observed in sighted readers in analogous tasks (Sadato et al. 1998; Burton, Snyder, Conturo, et al. 2002; Burton et al. 2012; Siuda-Krzywicka et al. 2016; Dzięgiel-Fivet et al. 2021). Previous studies also find that early somatosensory cortices (SMC) show expanded finger representations in proficient Braille readers, however, preferences for Braille over matched non-Braille tactile stimuli have not been found in SMC (Pascual-Leone and Torres 1993; Pascual-Leone et al. 1993; Sadato et al. 1998; Burton, Snyder, Conturo, et al. 2002; Burton et al. 2004; Kupers et al. 2007). Thus, it is unknown if any cortical areas outside of vOTC acquires specialization for Braille letters and words in blind readers.

We used sensitive region of interest (ROI) analyses as well as complementary data-driven cortical gradient maps to test whether regions within PPC show preferential responses to Braille over non-linguistic but perceptually similar tactile stimuli on the one hand and preferential responses to Braille over spoken language on the other, akin to the functional profile of the 'VWFA' in sighted readers. We hypothesized that analogous to the vOTC gradient, word-preferring portions of PPC should be found further from, and consequently posterior to, primary somatosensory cortices. Such analysis approaches are particularly relevant in the context of understanding the neural basis of reading, since previous studies find that in sighted readers preferential responses to print in vOTC are 'islands' among swaths of cortex that respond to complex visual shapes (Cohen and Dehaene 2004; Hannagan et al. 2015; Behrmann and Plaut 2020). Conventional whole-brain analyses that average across participants might therefore miss Braille specialization in PPC.

Finally, our third goal was to test the hypothesis that neural localization of spoken language influences the localization of Braille, analogous to what is observed in readers of visual print. In sighted readers, strong connectivity to spoken language networks predicts the localization of reading networks within vOTC and the lateralization of written language follows that of spoken language (Saygin et al. 2016; Stevens et al. 2017; Li et al. 2020). Reading, like spoken language, is on average strongly left-lateralized in sighted people but those sighted readers whose spoken language responses are localized to the right hemisphere also show right-lateralized written language (Schlaggar and McCandliss 2007; Cai et al. 2010; Seghier and Price 2011; Van der Haegen et al. 2012; Ossowski and Behrmann 2015). It is unknown

whether reading and spoken language networks co-lateralize in blind readers as they do in sighted

people.

Unlike the strong left-lateralization of spoken language in sighted people, lateralization of spoken

language is highly variable across congenitally blind individuals and on average only weakly left-

lateralized (Röder et al. 2000, 2002; Lane et al. 2017). This variability enabled us to use individual

difference analyses to test whether lateralization of Braille follows that of spoken language across blind

individuals. We also tested the effect of reading hand on lateralization and predicted that cortical areas

situated at earlier stages of Braille recognition (i.e., S1) would show stronger effects of reading hand,

whereas the effect of spoken language laterality would emerge in orthographic and higher-order language

regions (PPC, vOTC and inferior frontal cortex).

Method

Participants

Nineteen congenitally blind (12 females, mean age = 39.05 years, SD = 13.12) and 19 sighted controls

(11 females, mean age = 36.36 years, SD = 19.45) participated in the current study (Table 1). All

participants were native English speakers, and none had suffered from any known cognitive or

neurological disabilities (screened through self-report). All of the blind participants reported at least some

college education, with most having completed a college degree. The blind and sighted groups were

matched on age ($t_{(36)}$ = 0.499, p = 0.621) and years of education ($t_{(36)}$ = 0.867, p = 0.392). Blind

participants had at most minimal light perception from birth. Blindness was caused by pathology anterior

to the optic chiasm (i.e., not due to brain damage). All blind participants were fluent, frequent Braille

readers who began learning Braille at an average age of 4.6 years (SD = 1.53), rated their reading ability

as proficient to expert (mean = 4.6, SD =0.69 on a scale of 1 to 5), and reported reading Braille for an

average of 19.94 hours per week. We obtained information on Braille-reading hand dominance through a

post-experimental survey conducted over the telephone with 17 of the 19 blind adult participants (Table 1).

All participants gave informed consent according to procedures approved by the Johns Hopkins

Institutional Review Board.

Table 1. Participant information

Participant no.	Age (y)	Gender	Handedness	Reading handedness	Levels of education	Education yeas	Cause of blindness	Age started reading Braille (y)	Reading hours per week	Self-reported Braille reading ability (1-5)
B1	21	M	L	Bi-R	SC	16	LCA	4	14	5
B2	64	F	R	Bi-R	BA	16	ROP	6	56	5
B3	53	M	R	Bi-R	JD	20	LCA	6	7	4
B4	34	M	R	L	SC	15	Born without optic nerve	3	21	5
B5	42	M	Am	L	BA	17	ROP	3	21	5
B6	29	M	R	Bi-L	SC	14	LCA	4	<1	4
B7	39	F	R	L	BA	16	ROP	4	2	5
B8	34	F	R	--	SC	15	Optic Nerve Detached	3	--	5
B9	49	M	R	Bi-R	BA	17	unknown	8	<1	3
B10	26	F	R	Bi-R	MA	19	ROP	3	56	3
B11	49	F	L	R	MA	17	LCA	7	14	5
B12	39	F	R	L	MA	21	ROP	5	14	5
B13	35	F	R	Bi-L	MA	19	LCA	4	14	5
B14	46	F	R	--	BA	17	ROP	4	--	5
B15	33	F	R	L	BA	17	ROP	4	14	4
B16	25	F	Am	Bi-R	MA	19	LCA	5	56	5
B17	23	M	R	Bi-R	BA	15	LCA	4	28	5
B18	33	F	A	R	BA	16	LCA	3	14	5
B19	68	F	R	Bi-R	MA	19	ROP	5	7	5
Average										
Blind (n=19)	39.05 (SD=13.12)	12F	2L/3Am	--	BA	17.11 (SD = 1.91)	--	4.6 (SD=1.53)	19.94 (SD=18.36)	4.6 (SD=0.69)
Sighted (n=19)	36.36 (SD=19.45)	11F	2 L	--	BA	16.53 (SD = 2.19)	--	--		--

Handedness: left (L), ambidextrous (Am), or right (R), based on Edinburgh Handedness

Inventory. BA = Bachelor of Arts; MA = Master of Arts; HS = High School; JD = Juris Doctor; SC

= Some College; ROP = Retinopathy of prematurity; LCA = Leber's congenital amaurosis. For

Braille ability, participants were asked: "On a scale of 1 to 5, how well are you able to read Braille,

where 1 is 'not at all', 2 is 'very little', 3 is 'reasonably well', 4 is 'proficiently', and 5 is 'expert'?"

3 **Stimuli and experimental procedures**

4 The fMRI task included three reading conditions: words, consonant strings, and non-letter shapes (Figure

5 1) and two listening conditions: words, backwards speech (Kim et al. 2017). Reading stimuli were visual

6 for sighted participants and tactile for blind participants. Braille words were written in Grade-II contracted

English Braille, which is the most common form of Braille in the United States and therefore the most

ecologically valid form of written material for this population. Visual words were matched to Braille words

on average character length. Each consonant string stimulus consisted of 4 visual/Braille consonants.

The tactile control stimuli consisted of 24 unique strings of 4 non-letter shapes made of raised Braille dots.

The shapes were chosen based on pilot testing with a blind proficient Braille reader who reported them to

be recognizable and not confusable with Braille letters. The piloting process revealed that it was critical to

create shapes that did not fit into a single Braille cell (2 x 3 array of pins), otherwise the shapes would be

temporarily confused with Braille letters and characters. We therefore created shapes/pin arrays ranging

in size from 4 × 5 to 7 × 7 pins. The visual control stimuli were 24 unique strings, each comprised of 4

characters, which were false fonts. These were matched to English consonants on the number of strokes,

ascenders, descenders and stroke thickness (Kim et al. 2017). The auditory words were recorded by a

female native English speaker. The average word length was 5 letters long (SD = 1.4 letters) and the

average playtime was 0.41 s (SD = 0.3 s). The control auditory stimuli comprised backward speech

sounds created by playing each audio word in reverse.

On each trial, participants were presented with 6 stimuli from a single condition (e.g., Braille

words) followed by a memory probe. Participants judged whether the probe had appeared among the

previous 6 stimuli. This task was used to ensure attentive processing. The experiment had a total of 5

runs, each with 20 trials. The blind participants were asked to read with their dominant hand and

responded with the other hand. Each condition was repeated 4 times per run, and the order of conditions

was counterbalanced across runs. There were 6 rest periods (16 s) throughout each run. One sighted

participant and two blind participants were excluded from behavioral analysis due to failure to record their

responses (see SI and Kim et al., 2017 for details).

fMRI data acquisition

Functional and structural images were acquired using a 3T Phillips scanner at the F. M. Kirby Research

Center. T1-weighted images were collected using a magnetization-prepared rapid gradient-echo (MP-

RAGE) in 150 axial slices with 1 mm isotropic voxels. Functional BOLD scans were collected T1-weighted

structural images were collected in 150 axial slices with 1 mm isotropic voxels. Functional BOLD scans

were collected in 36 sequential ascending axial slices. TR = 2 s, TE = 30 ms, flip angle = 70°, voxel size =

2.4 × 2.4 × 2.5 mm, inter-slice gap = 0.5 mm, field of view (FOV) = 192 × 172.8 × 107.5.

fMRI data analysis

Preprocessing and whole-cortex analysis

Analyses were performed using FSL (version 5.0.9), FreeSurfer (version 5.3.0), the Human Connectome

Project workbench (version 1.2.0), and custom in-house software. The cortical surface was created for

each participant using the standard FreeSurfer pipeline (Dale et al. 1999; Smith et al. 2004; Glasser et al.

2013). For task data, preprocessing of functional data included motion-correction, high-pass filtering (128

s cut-off), and resampling to the cortical surface. Cerebellar and subcortical structures were excluded. On

the surface, the task data were smoothed with a 6 mm FWHM Gaussian kernel.

Three conditions of the reading task and two conditions of the listening task were included in a

general linear model. Analysis focused on the time-period during the initial six stimuli of each trial. Probe

stimulus and response periods were modeled separately and are not reported. White matter signal, CSF

signal, as well as motion spikes, were included as the covariates of no interest.

Whole-cortex random-effects analysis were run using mixed-effects and thresholded at $p < 0.01$

vertex-wise, and $p < 0.05$ cluster-wise, Family Wise Error corrected for multiple comparisons across the

cortex.

fMRI ROI analysis

We used subject-specific functional ROI analyses to examine five regions of interest: vOTC, V1, inferior

frontal cortex (IFC), and posterior parietal cortex (PPC), and the hand region of the left primary

somatosensory-motor cortex (SMC) (see SI for the search space construction details). Individual-subject

functional ROIs were defined within each of the above search spaces using a leave-one-run-out cross-

validation procedure (e.g., Ellis et al., 2021; Kim et al., 2017; Saygin et al., 2016). This procedure ensures

that independent sets of data are used to define ROIs and to test hypotheses and effectively uses each

run as a 'localizer scan' for the other runs of data (Cohen et al., 2019; Ellis et al., 2021; Kanjlia et al.,

2016, 2021; Ratan Murty et al., 2020; Saygin et al., 2016). ROIs were defined based on data from all but

one run, then the percent signal change (PSC) was extracted from the left-out run. This procedure was

repeated iteratively across all runs and the PSC was averaged across iterations (see SI for details). Each

individual subject's ROI was defined as the top 5% of vertices activated for tactile/visual consonant

strings > tactile/visual shapes. The main text reports results for ROIs defined using the consonant string >

control contrasts so as to focus on orthographic as opposed to semantic responses. In the supplement

we report results for ROIs defined using the words > control contrasts, which produced highly similar

results (see Figure S3 and Figure S6). Repeated-measured ANOVAs were used to analyze the ROI data,

and two-tailed paired *t*-tests were used for pairwise comparisons. All *p* values were Bonferroni-corrected

(see SI for details).

A vector of ROI analysis were conducted in PPC mask (Konkle and Caramazza 2013). We

divided the PPC mask into 13 segments centered along an equal distanced spline and plotted average

activity for three tactile conditions in each segment (see SI for details).

Topographical preference map

To examine topographic gradients in bilateral vOTC and PPC with data-driven winner-take-all approaches.

Each vertex was color-coded according to which stimulus condition showed highest activity. We also

calculated the topographical preferences winner-take-all map for each participant, then created group

histogram plots where each vertex was color-coded according to the number of participants showing

preference for a certain condition (i.e., words, consonant strings or control).

Laterality index analysis

LI was calculated separately for the reading and listening tasks for each participant in the SMC, PPC,

vOTC, and IFC. For the reading task, LI was determined based on the tactile/visual words > rest contrast.

For the listening task, LI was determined using the audio words > rest contrast. The LI was calculated

using the standard formula: (L - R) / (L + R), where L and R refer to the sums of the *z* statistics from the

relevant contrast within the left and right hemispheres, respectively. LI ranges from -1 to 1, with a score of

1 indicating strong left lateralization and -1 strong right lateralization. The bootstrap/histogram method

was used to ensure that LIs were not overly influenced by arbitrary activation threshold choices or outlier

voxels (Wilke and Schmithorst 2006). Bootstrapped LIs were computed using 20 evenly spaced

thresholds ranging from $z = 1.28$ to $z = 4.26$ (corresponding to one-sided $p = 0.1$ to $p = 0.00001$,

uncorrected). The LI reported for each participant represents the average across all thresholds.

Participants were excluded from the LI analysis if they did not have suprathreshold activation in both

hemispheres (see SI for details).

Results

Behavioral Results

Sighted and blind readers alike were more accurate at remembering more word-like stimuli in both the

reading and listening tasks (reading: main effect of lexicality (words and consonant strings > control

shapes): $F_{(2, 64)} = 24.991$, $p < 0.001$; listening (words > control shapes): $F_{(1, 32)} = 46.137$, $p < 0.001$. That is,

both groups achieved higher accuracy rates when reading words (sighted group visual words: 91.2%,

blind group tactile words: 83.1%) and consonant strings (sighted group: 86.6%, blind group: 77.9%), than

control shapes strings (sighted group: 72.7%, blind group: 62.5%). Likewise, accuracy rates were higher

for audio words (sighted group: 91.3%, blind group: 88.4%) than backwards speech (sighted group:

75.9%, blind group: 64.2%) (see Figure S1). There was no lexicality by group interaction in either reading

($F_{(2, 64)} = 0.067$, $p = 0.935$) or listening tasks ($F_{(1, 32)} = 2.252$, $p = 0.143$). Sighted readers had slightly

higher accuracy rates on both reading and listening tasks compared to blind readers (reading main effect

of group: $F_{(1, 32)} = 8.21$, $p < 0.01$; listening main effect of group: $F_{(1, 32)} = 6.305$, $p < 0.05$).

Participants were also faster at responding to word-like stimuli across reading (words and

consonant strings < control shapes; $F_{(2, 64)} = 12.998$, $p < 0.001$) and listening tasks (words < control

shapes; $F_{(1, 32)} = 30.763$, $p < 0.001$). Sighted readers were faster than blind readers for the reading task

(main effect of group: $F_{(1, 32)} = 10.641$, $p < 0.01$), but not for the listening task (main effect of group: $F_{(1, 32)}$

$= 2.488$, $p = 0.125$). For the reading task, there was a group by lexicality interaction ($F_{(2, 64)} = 3.943$, $p <$

0.05). Pairwise comparisons showed shorter reaction times for words and consonant strings relative to

the control condition in the blind group (words vs. control shapes, $t_{(32)} = -4.208$, $p < 0.01$; consonant

strings vs. control shapes, $t_{(32)} = -3.722$, $p < 0.05$; words vs. consonant strings, $t_{(32)} = -1.177$, $p = 0.24$).

For the sighted group, the only difference was faster reaction times for words than consonant strings

(words vs. control shapes, $t_{(32)}$ = -1.573, p =0.377; consonant strings vs. control shapes, $t_{(32)}$ = -0.718, p >

0.99; words vs. consonant strings, $t_{(32)}$ = -3.084, p < 0.05; see Figure S1).

fMRI Results

Visual (sighted) but not tactile Braille reading (blind) elicits a posterior-to-anterior functional

gradient in left vOTC and shows left-lateralization

We divided the left and right vOTC into posterior, middle, and anterior subregions (ROIs) and observed a

different posterior-to-anterior shift and a different lateralization pattern across visual and Braille reading

(Figure 1A). A four-way hemisphere (left, right) by posterior/anterior subregion (posterior, middle, anterior)

by lexicality (words, consonant strings, control shapes) by group (sighted, blind) ANOVA on the reading

task revealed a four-way interaction (F (4, 144) = 2.954, p < 0.05), indicating that lexicality, hemisphere,

and posterior/anterior subregion interacts with group (see SI for a complete summary of all effects).

As predicted, the sighted group showed the previously documented posterior-to-anterior functional

gradient in left but not right vOTC: larger responses to words in anterior vOTC, to consonant strings in

middle vOTC and no differences between visual conditions in posterior vOTC (three-way interaction

between hemisphere (left, right), posterior/anterior subregion (posterior, middle, anterior) and lexicality

(words, consonant strings, control shapes): $F_{(4, 72)}$ = 4.344, p < 0.01; in left vOTC two-way interaction

between lexicality (words, consonant strings, control shapes) and posterior/anterior subregion (posterior,

middle, anterior): $F_{(4, 72)}$ = 8.237, p < 0.001) (see SI for pairwise comparisons within each subregion).

By contrast, no gradient or laterality differences were observed in the blind group. Rather, all subregions

of bilateral vOTC responded most to words, followed by consonant strings followed by control shapes

(Figure 1A) (three-way hemisphere (left, right) by posterior/anterior subregion (posterior, middle, anterior)

by lexicality (words, consonant strings, control shapes) ANOVA interaction: $F_{(4, 72)}$ = 0.747, p = 0.563)

(see SI for pairwise comparisons and listening task results). A similar word preference was observed in

V1 for blind readers (Sadato et al. 1996; Cohen et al. 1997)(see SI for details).

The group differences identified by ROI analyses (above) were confirmed by data-driven

topographical preference maps where each vertex is color-coded according to the number of participants

showing numerical preference for a particular condition (Figure 1B, also see SI Figure S10 for

topographical preference maps in each participant and Figure S4 for topographical preference maps in

group average). In the sighted group, the word-preferring vertices were concentrated most consistently

across participants in the left anterior vOTC. The location of the consonant-preferring vertices was more

distributed, but the left anterior and middle vOTC were still the areas with the largest concentration of

subjects. The location of the control shapes-preferring vertices were most often observed in left posterior

and medial vOTC in the sighted. In the blind group, most participants showed a preference for words

throughout the vOTC. The highest degree of cross-participant overlap was observed bilaterally at MNI

coordinates (-44, -43, -18) and (41, -48, -22).

Figure 1. (A) Responses in left vOTC across the posterior, middle, and anterior subregions for blind and

sighted groups during the reading tasks. Bars show results from consonant string > control shapes leave-

one-run out individual subject ROI analysis. Error bars denote standard errors +/- the mean. Asterisks (*)

denote significant Bonferroni-corrected pairwise comparisons ($p < 0.05$). (B) Group histogram plots for

topographical preferences map. Each vertex was color-coded according to the number of participants

showing preference for words, consonants, or control shapes. Silver circles mark previously reported

location of VWFA (MNI coordinate: -46, -53, -20)(McCandliss et al. 2003).

The posterior parietal cortex (PPC) but not S1 of blind readers shows a preference for written

Braille words and consonant strings

Consistent with the hypothesis of Braille specialization in PPC of congenitally blind readers, the

PPC of blind readers preferred tactile over auditory stimuli and among tactile stimuli it preferred Braille

words and consonant strings over tactile control stimuli. The two-way lexicality (words, consonant strings,

control shapes) by group (sighted, blind) ANOVA revealed a marginally different response profile across

blind Braille and sighted print readers in the reading task (lexicality by group interaction effect: $F_{(2, 72)}$ =

2.682, p = 0.075; main effect of lexicality: $F_{(2, 72)}$ = 12.482, $p < 0.001$; main effect of group: $F_{(1, 36)}$ =0.081,

p = 0.778).

In blind readers, the PPC responded more to Braille words and Braille consonant strings than

control shapes (words vs. control shapes: $t_{(36)}$ = 3.602, $p < 0.05$; consonant strings vs. control shapes: $t_{(36)}$

= 3.595, $p < 0.05$; words vs. consonant strings: $t_{(36)}$ = 1.288, $p > 0.99$; Bonferroni-corrected paired t-tests)

(Figure S5). In the sighted group, words were not different from control shapes but consonant strings

elicited higher responses than control shapes, and words and consonants did not differ from each other

(words vs. control shapes: $t_{(36)}$ = 1.161, $p > 0.99$; consonant strings vs. control shapes: $t_{(36)}$ = 3.282, $p <$

0.05; words vs. consonant strings: $t_{(36)}$ = -2.208, p = 0.75; Bonferroni-corrected paired t-tests).

The PPC also showed larger responses to tactile than auditory stimuli in the blind group (two-way

modality (tactile, auditory) by lexicality (word, control) ANOVA, main effect of modality: $F_{(1, 18)}$ = 11.919, p

< 0.01; main effect of lexicality: $F_{(1, 18)}$ =9.079, $p < 0.01$; modality by lexicality interaction: $F_{(1, 18)}$ = 4.414,

p = 0.05). Responses to visual and auditory stimuli in the sighted were not different (two-way modality

(visual, auditory) by lexicality (word, control) ANOVA, main effect of modality: $F_{(1, 18)}$ =0.478, p = 0.498;

main effect of lexicality: $F_{(1, 18)}$ = 3.762, p = 0.068; modality by lexicality interaction: $F_{(1, 18)}$ =2.308, p =

0.146).

There was a larger response to auditory words than auditory control stimuli in both groups in the

PPC (two-way lexicality (audio words, audio control) by group (sighted, blind) ANOVA**,** main effect of

lexicality: words > control, $F_{(1, 36)}$ = 7.624, $p < 0.01$; main effect of group: $F_{(1, 36)}$ =0.602, p = 0.443; group

by lexicality interaction: $F_{(1, 32)}$ = 2.493, p = 0.123, Figure S5).

1   A data-driven topographic winner-take-all map showed that preferential responses to Braille

2 words were located in the most posterior portion of PPC and extended into parieto-occipital and dorsal

3 occipital regions (Figure 2A). A vector-of-ROIs analysis along the anterior/posterior extent of PPC

4 revealed a position by reading condition interaction in blind readers (left PPC: $F_{(12, 24)}$ = 16.62, p < 0.001;

5 right PPC: $F_{(14, 28)}$ = 13.15, p < 0.001; see Figure S7 for right PPC). The most anterior portions of PPC,

6 immediately adjacent to S1, showed preferential responses to tactile shapes, whereas posterior PPC

7 showed a preference for Braille words. In the winner-take-all map, a small middle region in left and right

8 PPC showed the highest responses to Braille consonant strings. These results were confirmed by a

9 participant-number-histogram (Figure 2B). This pattern is suggestive of an anterior-to-posterior decoding

10 gradient in PPC in blind Braille readers analogous to the posterior-to-anterior gradient observed in vOTC

11 of sighted print readers.

Figure 2. (A) Posterior parietal cortex (PPC) winner-take-all map (upper) during the reading task: words,

consonant strings, and control stimuli. The black outline indicates the hand region of the primary sensory-

motor cortex (Neurosynth). Vector-of-ROIs results (lower) showed the mean response to each reading

condition along left anterior/posterior PPC extent in the blind group (13 segments). Segment centers are

marked by red dots (see Figure S7 for right PPC). (B) Group participant-number-histogram plots for

topographical preferences winner-take-all map. Each vertex was color-coded according to the number of

participants showing preference for words, consonants, or control shapes.

By contrast to the PPC, early somatosensory cortices (SMC hand region, Figure S5) did not show a

preferential response to Braille words and showed the same functional profile across blind and sighted

readers (two-way lexicality (words, consonant strings, control shapes) by group (sighted, blind) ANOVA,

main effect of lexicality: $F_{(2, 72)} = 7.946$, $p < 0.001$; main effect of group: $F_{(1, 36)} = 2.776$, $p = 0.104$; group

by condition interaction: $F_{(2, 72)} = 1.922$, $p = 0.156$). (See SI for details of listening task results.) Likewise,

the left IFC, a high-level language region, showed a similarly preferential response to linguistic stimuli (i.e.,

words) across groups and tasks (reading task, two-way lexicality (words, consonant strings, control

shapes) by group (sighted, blind) ANOVA, main effect of lexicality: $F_{(2, 72)} = 38.6$, $p < 0.001$; main effect of

group: $F_{(1, 36)} = 0.453$, $p = 0.505$; lexicality by group interaction: $F_{(2, 72)} = 0.546$, $p = 0.581$) (see SI for

details). In sum, the PPC and adjacent parieto-occipital cortices showed differential recruitment for Braille

reading in people born blind as compared to reading of visual print in sighted people.

Lateralization of Braille is predicted by spoken language lateralization late in cortical processing

hierarchy (i.e., LIFC) and by Braille-reading hand early in cortical processing hierarchy (i.e., S1)

Lateralization index (LI) analysis revealed left-lateralized responses to written and spoken words in vOTC

and IFC of sighted readers, consistent with prior studies (one-sample t tests of LI = 0, visual print words >

rest: vOTC: $t_{(18)} = 5.235$, $p < 0.001$; IFC: $t_{(17)} = 3.854$, $p < 0.001$; spoken words > rest: vOTC: $t_{(18)} = 2.868$,

$p < 0.01$; IFC: $t_{(17)} = 2.06$, $p = 0.055$; see SI for other ROIs). By contrast, the blind group did not show

systematic left-lateralization in any region (one-sample t tests of LI = 0, Braille words > rest: vOTC: $t_{(18)} =$

0.935, $p = 0.362$; IFC: $t_{(18)} = 0.19$, $p = 0.852$; spoken words > rest: vOTC: $t_{(18)} = 0.583$, $p = 0.567$; IFC: $t_{(18)}$

= -0.834, $p = 0.415$; see SI for other regions). Consistent with prior evidence, there was substantial

variability in lateralization of spoken and written language among blind participants, with some

participants showing strong left and others strong right lateralization (Lane et al. 2017) (Figure 3).

Individual difference analysis revealed a strong relationship between lateralization of spoken and written

language in the blind group in all regions except early somatosensory cortex (SMC), including IFC, vOTC

and PPC (multiple regression with the LI of spoken words in IFC and dominant reading hand entered as

the regressors in each region, LI of spoken words, IFC: $t_{(18)}$ = 6.551, p < 0.001; vOTC: $t_{(18)}$ = 5.23, p <

0.001; PPC: IFC: $t_{(18)}$ = 5.83, p < 0.001; SMC: $t_{(18)}$ = 1.684, p = 0.112). Conversely, relative to other

regions, the SMC showed a strong effect of reading hand on lateralization (multiple regression, for the

regressor dominant reading hand: $t_{(18)}$ = 3.034, p < 0.01). The effect of reading hand was also observed

in PPC and vOTC but not in the IFC (multiple regression, effect of reading hand: PPC $t_{(18)}$ = 3.814, p <

0.01; vOTC $t_{(18)}$ = 3.424, p < 0.01; IFC $t_{(18)}$ = -0.693, p = 0.498, see SI for details.) Correlations between

spoken and written language lateralization were weaker in the sighted group and only reached

significance in IFC and PPC (IFC: r = 0.865, p < 0.001; PPC: r = 0.60, p < 0.05). This is likely due to low

variability of laterality scores in the current sighted sample (i.e., uniformly strong left-lateralization, see SI

for details).

Figure 3. (A) Correlations between the Laterality Index (LI) of audio words in IFC and the LI of tactile

words in SMC, PPC, vOTC in blind Braille readers. Data points represent individual participants, left-

handed Braille readers: dark grey circles; right-handed readers: light gray circles; two subjects who did

not obtain reading hand information: white circles. LI 1 score indicates strong left lateralization and -1

indicates strong right lateralization. (B) LI of Braille reading with left (grey) and right (white) hand

separately. Asterisks (*) on the bar denotes significant difference from 0; asterisks (*) between two bars

denote significant difference between the LI of left- vs. right-hand reading (*p* <0.05).

Whole cortex analyses

Reading-related activity (relative to rest) was left-lateralized in the sighted and bilateral in the blind group.

For reading as compared to rest, both sighted (visual words) and blind (Braille words) readers activated

the bilateral vOTC (blind peak: -40, -57, -13; sighted peak: -35, -45, -20), including the location of the

classic VWFA (peak: -46, -53, -20), as well as early visual cortices, specifically the foveal confluence

(V1/V2/V3) (Figure 4). vOTC responses in the blind group extended medially and anteriorly, as well as

into lateral occipito-temporal cortex. Both groups also activated prefrontal cortices (inferior frontal gyrus

and middle frontal gyrus). (See Table S1 for complete list of foci.)

Reading Braille in the blind group (relative to rest) extensively activated posterior parietal cortices

(superior parietal lobule, supramarginal gyrus (SMG)), posterior to early sensory-motor hand

representations. This activity extended into parieto-occipital and dorsal occipital regions in the blind group.

The sighted group activated a small cluster within PPC. A lateral temporal region was active in the

sighted but not blind group.

Like responses to Braille, responses to spoken words were left-lateralized in the sighted group

and bilateral in the blind group. Listening to words (audio words > rest) activated bilateral vOTC (peak: -

42, -43, -16), including the location of the classic VWFA, and early 'visual' cortices, in the blind group only.

Both groups activated classic fronto-temporal language regions in inferior and prefrontal as well as lateral

temporal cortices (Figure 4).

When reading and listening to words was compared directly, for the sighted group, reading

induced greater activation in bilateral vOTC, including the typical location of the VWFA and regions

posterior to it, as well as bilateral early visual cortices. The blind group also activated a region in left

vOTC (fusiform; peak: -31, -58, -16), but this activation was medial to the typical VWFA location. A cluster

of activity was also observed posterior to the typical VWFA location in the blind group, in the inferior temporal/lateral occipital cortex (peak: -42 -74 -5) as well as in left foveal early 'visual' cortices. The blind group showed more extensive activation than sighted group for Braille words > spoken words in PPC, including the SMG and superior parietal lobule, extending into dorsal occipital/parieto-occipital cortices.

In sum, although both groups activated vOTC during reading, the peak location, distribution and functional profile of responses in vOTC were distinct across groups. The blind group activated more extensive posterior parietal, parieto-occipital, and dorsal occipital areas during reading (Braille) (see Figure S8 for other contrast and Figure S9 for group comparison for all the contrast).

For the words compared to control shapes, the sighted group's whole brain results showed stronger responses in the expected left-lateralized lateral temporal language areas, as well as left sensory-motor cortex, left medial superior frontal gyrus and bilateral early visual cortices and precuneus (see Figure S8b). The blind group's whole brain results also showed stronger response to words over control shapes in lateral temporal language areas, extending on the lateral surface into visual cortices, as well as inferior frontal cortices. In the blind group, words>control also elicited stronger activation in bilateral vOTC and left early visual cortices. Words also elicited larger responses than consonant strings in lateral temporal cortices in both groups. In the blind readers, these responses were more posterior (see Supplementary Figure S8b for details). These whole-brain results demonstrate that although in the sighted group only small portions of vOTC responded and no portion of PPC preferentially responded to words, this is not because such effects were precluded by the current stimuli or task. We observed robust preferential responses to words in large swaths of expected, left-lateralized temporal lobe language areas.

2   **Figure 4.** Whole-cortex results for blind (left column) and sighted (right column) *p* < 0.05 cluster-

3   corrected. Blue circles mark previously reported location of VWFA (MNI coordinate: -46, -53, -20)

4   (McCandliss et al. 2003). The yellow outline marks the hand S1/M1 region.

Discussion

No word-form gradient in vOTC of blind Braille readers

Consistent with past research, we observed a posterior-to-anterior functional gradient, progressively

higher responses to words anteriorly in the left vOTC of sighted readers. The posterior-most portion of

vOTC showed no preference for words, consonant string or false font, the middle for consonants and the

anterior-most showed an equivalently high response to words and consonants (Bruno et al. 2008; Cohen

et al. 2008; Dehaene and Cohen 2011; Ludersdorfer et al. 2013). By contrast, in the vOTC of blind

readers we found no evidence for left-lateralization, no evidence for a posterior-to-anterior functional

gradient, or posterior/anterior change in modality preference. The entire posterior/anterior extent of

bilateral vOTC, as well as V1, showed a preference for words in the blind group (Sadato et al. 1996;

Cohen et al. 1997; Kupers et al. 2007). There was also no change in preference for written as opposed to

spoken words along the posterior/anterior extent of vOTC in blind readers (see Figure S2 in SI). In sum,

we find no evidence for an orthographic gradient in vOTC of blind Braille readers.

Although we found no word-form gradient in vOTC of blind Braille readers, the canonical location

of the VWFA responded to words in both blind and sighted groups. Even the functional profile of the

VWFA location differed somewhat across groups, however. In sighted readers the canonical location of

the VWFA responded as much or more to consonant strings as to words in the current task. This is

consistent with what was reported in a previous study with a smaller subsample of the present

participants (Kim et al. 2017). Importantly, this pattern of VWFA response is also consistent with previous

studies with sighted participants using attentionally demanding reading tasks analogous to the one used

in the current study. Such tasks include longer presentation times or require attention to the orthographic

content of the stimuli, like one-back memory task or phonological task (Bruno et al. 2008; Dehaene and

Cohen 2011; Twomey et al. 2011; Ludersdorfer et al. 2013; Mano et al. 2013). Like these prior studies,

the current experiment used relatively long presentations and involved a working memory task. Under

such conditions, the VWFA responds as much or more to non-words and letter strings as to words. This

response profile has been interpreted to reflect the increased demands on orthographic processing for

letter strings and nonwords relative to words in tasks that require attention to orthography. By contrast, in

studies using tasks that are relatively quick and automatic, like passive reading or visual feature detection,

the VWFA responds more to words than consonant strings or pseudowords, putatively because more

anterior portions of the VWFA code letter combinations and whole words (Cohen et al. 2002; Binder et al.

2006; Vinckier et al. 2007).

In the current study, which used an attentionally demanding task and relatively slow presentation,

the VWFA of blind readers responded more to words than consonant strings, whereas that of sighted

readers responded equally or more to consonant strings. One interpretation of this difference in VWFA

profile across groups, is that the VWFA supports orthographic processing in the sighted but is sensitive to

high level linguistic information (semantic, syntactic) in people born blind. No amount of attention to

orthography can override this preference. Regardless of the interpretation of this particular group

difference, we find that the same conditions that are sufficient to elicit an orthographic gradient response

in the sighted do not do so in people who are blind.

Whole-cortex analysis also revealed a partially distinct distribution of reading responses in vOTC

across blind and sighted people. When Braille and spoken words were each compared to rest, a peak of

activation was observed in the VWFA location in both groups. However, when reading words was

compared to hearing words, activity peaked medial to the VWFA in the blind but not sighted group (peak:

-27, -61, -14). A similar medial vOTC region was recently found to be functionally connected with dorsal

parietal cortices in sighted people (Bouhali et al., 2019; Cohen et al., 2008; Corbetta & Shulman, 2002;

Henry et al., 2005; Saalmann et al., 2007). Previous studies have also found that in people who are blind,

the classic lateral VWFA location is sensitive to syntactic complexity of spoken sentences, shows

enhanced responses to spoken language and enhanced connectivity with fronto-temporal language

networks (Burton, Snyder, Diamond, et al. 2002; Striem-Amit et al. 2012; Lane et al. 2015; Kim et al. 2017;

Dzięgiel-Fivet et al. 2021). An intriguing possibility to be tested in future work is that medial vOTC is

responsive to Braille-specific input from PPC, whereas the classic, more lateral VWFA location, is driven

by linguistic information from fronto-temporal networks.

Parieto-occipital decoding stream in blind readers of Braille

We observed more extensive and different involvement of parietal and parieto-occipital regions in Braille

as opposed to visual print reading. Some past studies have also reported parietal activity during Braille

reading but had not explored whether these responses were Braille-specific or merely related to general

tactile perception (Sadato et al. 1998; Burton, Snyder, Conturo, et al. 2002; Burton et al. 2012; Siuda-Krzywicka et al. 2016; Dzięgiel-Fivet et al. 2021). Using sensitive individual-subject analyses, we found that in blind readers, subregions of posterior PPC and parieto-occipital cortices respond preferentially to Braille words relative to both tactile shapes and spoken words. This functional profile is consistent with its specific involvement in Braille reading. In contrast, anterior regions of PPC, adjacent to S1, responded most to unfamiliar tactile shapes comprised of Braille dots. The hand regions of S1 itself did not show robust or preferential responses to Braille (Burton, Snyder, Conturo, et al. 2002; Kupers et al. 2007). This response profile is suggestive of an anterior-to-posterior Braille reading processing stream, with anterior regions supporting recognition of tactile patterns and posterior regions performing Braille-specific, orthographic processing.

Further work is needed to uncover the precise contribution of PPC and parieto-occipital cortices to Braille reading. In sighted readers, the PPC shows sensitivity to phonological rather than orthographic information, during visual reading, and is involved in effortful letter-by-letter reading (e.g., when words are degraded) (Booth et al. 2003; Henry et al. 2005; Costanzo et al. 2012; Taylor et al. 2013; Ossmy et al. 2014; Bouhali et al. 2019) . Whether the same portions of PPC contribute to Braille and visual reading is not known and the precise contribution of PPC to Braille and visual reading remains to be fully characterized. It is worth noting that the Braille words used in the current study were written in Grade-II contracted Braille, which is the most used form of Braille in the US and therefore the ecologically valid choice. About 60% of the Braille word stimuli in the current study contained one or more contractions, i.e., a single Braille character that stands for multiple letters. We would argue that a full understanding of the neural basis of Braille reading requires investigating Braille reading as it is naturally done by proficient readers. How Braille contractions influence neural responses in the reading network is not known and should be examined in future research. It is possible that some of the differences between the neural basis of Braille and the neural basis of visual print relate to the contracted nature of Braille. For example, it would be worth testing the role of PPC in unpacking Braille contractions. Given previously documented similarities between the neural bases of various visual scripts, we think it is unlikely that most of the robust group differences we observed in the current study are related to the contracted nature of Braille alone. Nevertheless, this topic merits further investigation.

Lateralization of Braille reading: effects of spoken language laterality and reading hand

We observed reduced and variable lateralization of both written and spoken language networks in blind compared to the sighted individuals. This observation is consistent with prior studies using spoken sentences (Lane et al., 2017; also see Röder et al., 2002). The reason for reduced lateralization of spoken language individuals born blind is not known. One hypothesis is that in sighted individuals language is "forced" out of the right hemisphere by visuospatial functions and such pressure is different in people born blind (Levy 1969; Kosslyn 1987). Alternatively, changes in the timing of language acquisition could affect their lateralization patterns in blindness (Locke 1997; Bishop 2013).

Importantly, here we find that laterality of responses to written words are predicted by the laterality of spoken language across blind individuals and this effect increases along the processing hierarchy peaking in IFC. In other words, those blind individuals who show right-lateralized responses to spoken words also show right-lateralized responses to written words. The only region which did not show a relationship between laterality of spoken and written language is early somatosensory cortex, where laterality was predicted only by reading hand. Previous studies with sighted readers with right hemisphere spoken language responses have likewise observed co-lateralization of spoken and written language, although this relationship was weak in the current study, likely due to low laterality variability in our sighted sample (Cai et al. 2010; Van der Haegen et al. 2012). The striking relationship between lateralization of Braille and spoken language suggests that written and spoken language co-lateralize regardless of reading modality. This observation is consistent with the hypothesis that strong connectivity and proximity to spoken language networks is one of the determining factors of which regions become 'recycled' for reading (Hannagan et al. 2015; Saygin et al. 2016; Stevens et al. 2017).

In contrast to the effect of spoken language on laterality, the effect of reading hand was strongest at early stages of processing (in the primary somatosensory cortex), weaker at intermediate stages (in PPC and vOTC), and absent in a high-level language region (IFC). These laterality effects support the view that PPC and vOTC participate in reading related processes in blindness, rather than solely sensory recognition or high-level language processing.

It is worth noting that in the current study, the participants were asked to read single words with one hand. However, in naturalistic reading, where multiple lines of text need to be scanned,

readers often use both hands (Millar 2003). The precise contribution of each hands is not entirely

clear (Hermelin and O'connor 1971; Millar 2003). Typically one hand is the dominant reading hand

whereas the other is used to run ahead to track position along the page and gain a preview (Millar

1984, 2003). The neural basis of naturalistic two-handed reading remains to be investigated in

future research.

General conclusions

The neural basis of tactile Braille reading in congenitally blind individuals and visual print reading in

sighted people is governed by analogous connectivity principles but has distinct anatomical profiles.

While visual print reading recruits a posterior/anterior vOTC gradient, no such gradient is observed in the

vOTC of blind readers of Braille. Blind readers of Braille recruit posterior parietal cortices to a greater

degree and in a different way compared to visual print reading in sighted people. Only blind readers show

preferential responses to written words in PPC and parieto-occipital cortex. We observed suggestive

evidence for an anterior-to-posterior stream of processing in the parietal cortex of blind Braille readers,

with anterior parietal areas responsive to non-Braille tactile patterns and more posterior parietal, parieto-

occipital and dorsal occipital regions responsive to Braille words. In blind and sighted readers alike,

lateralization of spoken language predicts lateralization of written language. In blind readers of Braille, the

effect of spoken language on laterality of Braille becomes more pronounced along the cortical processing

hierarchy. Conversely, reading-hand affects lateralization more in lower stages of processing. Together

these results suggest that analogous connectivity principles govern neural instantiation of reading in

sighted and blind readers: the localization of reading depends jointly on connectivity to sensory input

regions (unilateral S1/ bilateral V1) and language networks. These principles, however, give rise to

distinct anatomical profiles depending on modality of reading and neural basis of spoken language in a

particular individual.

1 **Data and Code Availability Statement**

Credit Author Statement

Funding

This work was supported by grants from the Johns Hopkins Science of Learning Institute (80034917) and

the National Institutes of Health (National Eye Institute, R01, EY027352-01).

Acknowledgments

We would like to thank all our blind and sighted participants, the blind community and the National

Federation of the Blind. Without their support, this study would not be possible. We would also like to

thank the F. M. Kirby Research Center for Functional Brain Imaging at the Kennedy Krieger Institute for

their assistance in data collection.

References

Baker CI, Liu J, Wald LL, Kwong KK, Benner T, Kanwisher N. 2007. Visual word processing and

experiential origins of functional selectivity in human extrastriate cortex. Proc Natl Acad Sci.

104:9087–9092.

Barttfeld P, Abboud S, Lagercrantz H, Adén U, Padilla N, Edwards AD, Cohen L, Sigman M, Dehaene S,

Dehaene-Lambertz G. 2018. A lateral-to-mesial organization of human ventral visual cortex at

birth. Brain Struct Funct. 223:3107–3119.

Bauer C, Yazzolino L, Hirsch G, Cattaneo Z, Vecchi T, Merabet LB. 2015. Neural correlates associated

with superior tactile symmetry perception in the early blind. Cortex. 63:104–117.

Behrmann M, Plaut DC. 2020. Hemispheric Organization for Visual Object Recognition: A Theoretical

Account and Empirical Evidence*. Perception. 49:373–404.

Binder JR, Medler DA, Westbury CF, Liebenthal E, Buchanan L. 2006. Tuning of the human left fusiform

gyrus to sublexical orthographic structure. NeuroImage. 33:739–748.

Bishop DVM. 2013. Cerebral Asymmetry and Language Development: Cause, Correlate, or

Consequence? Science. 340:1230531.

Bola Ł, Matuszewski J, Szczepanik M, Droździel D, Sliwinska MW, Paplińska M, Jednoróg K, Szwed M,

Marchewka A. 2019. Functional hierarchy for tactile processing in the visual cortex of sighted

adults. NeuroImage. 202:116084.

Bolger DJ, Perfetti CA, Schneider W. 2005. Cross-cultural effect on the brain revisited: Universal

structures plus writing system variation. Hum Brain Mapp. 25:92–104.

Booth JR, Burman DD, Meyer JR, Gitelman DR, Parrish TB, Mesulam MM. 2003. Relation between brain

activation and lexical performance. Hum Brain Mapp. 19:155–169.

Bouhali F, Bézagu Z, Dehaene S, Cohen L. 2019. A mesial-to-lateral dissociation for orthographic

processing in the visual cortex. Proc Natl Acad Sci U S A. 116:21936–21946.

Bouhali F, de Schotten MT, Pinel P, Poupon C, Mangin JF, Dehaene S, Cohen L. 2014. Anatomical

connections of the visual word form area. J Neurosci. 34:15402–15414.

Bruno JL, Zumberge A, Manis FR, Lu ZL, Goldman JG. 2008. Sensitivity to orthographic familiarity in the

occipito-temporal region. NeuroImage. 39:1988–2001.

Büchel C, Price C, Friston K. 1998. A multimodal language region in the ventral visual pathway. Nature.

394:274–277.

Burks JD, Boettcher LB, Conner AK, Glenn CA, Bonney PA, Baker CM, Briggs RG, Pittman NA,

O'Donoghue DL, Wu DH, Sughrue ME. 2017. White matter connections of the inferior parietal

lobule: A study of surgical anatomy. Brain Behav. 7:1–12.

Burton H, Sinclair RJ, Agato A. 2012. Recognition memory for Braille or spoken words: An fMRI study in

early blind. Brain Res. 1438:22–34.

Burton H, Sinclair RJ, McLaren DG. 2004. Cortical activity to vibrotactile stimulation: An fMRI study in

blind and sighted individuals. Hum Brain Mapp. 23:210–228.

Burton H, Snyder AZ, Conturo TE, Akbudak E, Ollinger JM, Raichle ME. 2002. Adaptive Changes in Early

and Late Blind: A fMRI Study of Braille Reading. J Neurophysiol. 87:589–607.

Burton H, Snyder AZ, Diamond JB, Raichle ME. 2002. Adaptive Changes in Early and Late Blind: A fMRI

Study of Verb Generation to Heard Nouns. J Neurophysiol. 88:3359–3371.

Cai Q, Paulignan Y, Brysbaert M, Ibarrola D, Nazir TA. 2010. The left ventral occipito-temporal response

to words depends on language lateralization but not on visual familiarity. Cereb Cortex. 20:1153–

1163.

Cohen, Celnik P, Pascual-Leone A, Corwell B, Faiz L, Dambrosia J, Honda M, Sadato N, Gerloff C,

Catala´ MD, Hallett M. 1997. Functional relevance of cross-modal plasticity in blind humans.

Nature. 389:180–183.

Cohen, Dehaene S, Naccache L, Lehéricy S, Dehaene-Lambertz G, Hénaff MA, Michel F. 2000. The

visual word form area: spatial and temporal characterization of an initial stage of reading in

normal subjects and posterior split-brain patients. Brain J Neurol. 291–307.

Cohen L, Dehaene S. 2004. Specialization within the ventral stream: The case for the visual word form

area. NeuroImage. 22:466–476.

Cohen L, Dehaene S, Vinckier F, Jobert A, Montavont A. 2008. Reading normal and degraded words:

Contribution of the dorsal and ventral visual pathways. NeuroImage. 40:353–366.

Cohen L, Lehéricy S, Chochon F, Lemer C, Rivaud S, Dehaene S. 2002. Language-specific tuning of visual cortex? Functional properties of the Visual Word Form Area. Brain J Neurol. 125:1054–1069.

Cohen MA, Dilks DD, Koldewyn K, Weigelt S, Feather J, Kell AJE, Keil B, Fischl B, Zöllei L, Wald L, Saxe R, Kanwisher N. 2019. Representational similarity precedes category selectivity in the developing ventral visual pathway. NeuroImage. 197:565–574.

Corbetta M, Shulman GL. 2002. Control of goal-directed and stimulus-driven attention in the brain. Nat Rev Neurosci. 3:201–215.

Costanzo F, Menghini D, Caltagirone C, Oliveri M, Vicari S. 2012. High frequency rTMS over the left parietal lobule increases non-word reading accuracy. Neuropsychologia. 50:2645–2651.

Dale AM, Fischl B, Sereno MI. 1999. Cortical surface-based analysis: I. Segmentation and surface reconstruction. NeuroImage. 9:179–194.

Debowska W, Wolak T, Nowicka A, Kozak A, Szwed M, Kossut M. 2016. Functional and structural neuroplasticity induced by short-term tactile training based on braille reading. Front Neurosci. 10:1–13.

Dehaene S, Cohen L. 2011. The unique role of the visual word form area in reading. Trends Cogn Sci. 15:254–262.

Dehaene S, Cohen L, Sigman M, Vinckier F. 2005. The neural code for written words: A proposal. Trends Cogn Sci. 9:335–341.

Dehaene S, Pegado F, Braga LW, Ventura P, Nunes Filho G, Jobert A, Dehaene-Lambertz G, Kolinsky R, Morais J, Cohen L, Henrich J, Heine SJ, Norenzayan A, Dehaene S, Cohen L, Castro-Caldas A, Carreiras M, Petersson KM, Silva C, Castro-Caldas A, Ingvar M, Reis A, Li G, Petersson KM, Reis A, Askelöf S, Castro-Caldas A, Ingvar M, Castro-Caldas A, Petersson KM, Reis A, Stone-Elander S, Ingvar M, Petersson KM, Reis A, Castro-Caldas A, Ingvar M, Morais J, Bertelson P, Cary L, Alegria J, Cheung H, Chen HC, Perre L, Pattamadilok C, Montant M, Ziegler JC, Ventura P, Morais J, Kolinsky R, Ziegler JC, Petrova A, Ferrand L, Shaywitz BA, Cohen L, Dehaene S, Baker CI, Brem S, Maurer U, Polley DB, Steinberg EE, Merzenich MM, Li W, Piëch V, Gilbert CD, Sigman M, Gauthier I, Curran T, Curby KM, Collins D, Harel A, Gilaie-Dotan S, Malach R, Bentin

S, Rossion B, Kung CC, Tarr MJ, Bolger DJ, Perfetti CA, Schneider W, Golarai G, Gaillard R, Cohen L, Dehaene S, Vinckier F, Jobert A, Montavont A, Eger E, Schweinberger SR, Dolan RJ, Henson RN, Downing PE, Chan AW, Peelen M V, Dodds CM, Kanwisher N, Nazir TA, Ben-Boutayab N, Decoppet N, Deutsch A, Frost R, Rayner K, Jacquemot C, Pallier C, LeBihan D, Dehaene S, Dupoux E, van Atteveldt N, Formisano E, Goebel R, Blomert L, Blau V, Turkeltaub PE, Gareau L, Flowers DL, Zeffiro TA, Eden GF, Cohen L, Jobert A, Bihan D Le, Dehaene S, Yoncheva YN, Zevin JD, Maurer U, McCandliss BD, Desroches AS, Booth JR, Kherif F, Josse G, Price CJ, Longcamp M, Anton JL, Roth M, Velay JL. 2010. How learning to read changes the cortical networks for vision and language. Science. 330:1359–1364.

DiCarlo JJ, Cox DD. 2007. Untangling invariant object recognition. Trends Cogn Sci. 11:333–341.

Duhamel JR, Colby CL, Goldberg ME. 1998. Ventral intraparietal area of the macaque: Congruent visual and somatic response properties. J Neurophysiol. 79:126–136.

Dzięgiel-Fivet G, Plewko J, Szczerbiński M, Marchewka A, Szwed M, Jednoróg K. 2021. Neural network for Braille reading and the speech-reading convergence in the blind: Similarities and differences to visual reading. NeuroImage. 231.

Ellis CT, Skalaban LJ, Yates TS, Turk-Browne NB. 2021. Attention recruits frontal cortex in human infants. Proc Natl Acad Sci U S A. 118.

Feng X, Altarelli I, Monzalvo K, Ding G, Ramus F, Shu H, Dehaene S, Meng X, Dehaene-Lambertz G. 2020. A universal reading network and its modulation by writing system and reading ability in French and Chinese children. eLife. 9:1–51.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S, Webster M, Polimeni JR, Van Essen DC, Jenkinson M. 2013. The minimal preprocessing pipelines for the Human Connectome Project. NeuroImage. 80:105–124.

Hannagan T, Amedi A, Cohen L, Dehaene-Lambertz G, Dehaene S. 2015. Origins of the specialization for letters and numbers in ventral occipitotemporal cortex. Trends Cogn Sci. 19:374–382.

Hegner YL, Lee Y, Grodd W, Braun C. 2010. Comparing tactile pattern and vibrotactile frequency discrimination: A human fMRI study. J Neurophysiol. 103:3115–3122.

Henry C, Gaillard R, Volle E, Chiras J, Ferrieux S, Dehaene S, Cohen L. 2005. Brain activations during letter-by-letter reading: A follow-up study. Neuropsychologia. 43:1983–1989.

Hermelin B, O'connor N. 1971. Right and left handed reading of braille. Nature. 231:470.

Hu W, Lee HL, Zhang Q, Liu T, Geng LB, Seghier ML, Shakeshaft C, Twomey T, Green DW, Yang YM, Price CJ. 2010. Developmental dyslexia in Chinese and English populations: Dissociating the effect of dyslexia from language differences. Brain. 133:1694–1706.

Kaas JH. 2012. Somatosensory System. In: The Human Nervous System. Third Edit. ed. Elsevier. p. 1074–1109.

Kanjlia S, Lane C, Feigenson L, Bedny M. 2016. Absence of visual experience modifies the neural basis of numerical thinking. Proc Natl Acad Sci U S A. 113:11172–11177.

Kanjlia S, Loiotile RE, Harhen N, Bedny M. 2021. 'Visual' cortices of congenitally blind adults are sensitive to response selection demands in a go/no-go task. NeuroImage. 236:118023.

Kim JS, Kanjlia S, Merabet LB, Bedny M. 2017. Development of the visual word form area requires visual experience: Evidence from blind braille readers. J Neurosci. 37:11495–11504.

Konkle T, Caramazza A. 2013. Tripartite organization of the ventral stream by animacy and object size. J Neurosci. 33:10235–10242.

Kosslyn SM. 1987. Seeing and Imagining in the Cerebral Hemispheres: A Computational Approach. Psychol Rev. 94:148–175.

Krafnick AJ, Tan LH, Flowers DL, Luetje MM, Napoliello EM, Siok WT, Perfetti C, Eden GF. 2016. Chinese Character and English Word processing in children's ventral occipitotemporal cortex: FMRI evidence for script invariance. NeuroImage. 133:302–312.

Kupers R, Pappens M, De Noordhout AM, Schoenen J, Ptito M, Fumal A. 2007. rTMS of the occipital cortex abolishes Braille reading and repetition priming in blind subjects. Neurology. 68:691–693.

Lane C, Kanjlia S, Omaki A, Bedny M. 2015. "Visual" cortex of congenitally blind adults responds to syntactic movement. J Neurosci. 35:12859–12868.

Lane C, Kanjlia S, Richardson H, Fulton A, Omaki A, Bedny M. 2017. Reduced Left Lateralization of Language in Congenitally Blind Individuals. J Cogn Neurosci. 29:65–78.

Levy J. 1969. Possible basis for the evolution of lateral specialization of the human brain [36]. Nature. 224:614–615.

Lewis JW, Van Essen DC. 2000. Corticocortical connections of visual, sensorimotor, and multimodal processing areas in the parietal lobe of the macaque monkey. J Comp Neurol. 428:112–137.

Li J, Osher DE, Hansen HA, Saygin ZM. 2020. Innate connectivity patterns drive the development of the visual word form area. Sci Rep. 10:1–12.

Locke JL. 1997. A Theory of Neurolinguistic Development. Brain Lang. 58:265–326.

Ludersdorfer P, Schurz M, Richlan F, Kronbichler M, Wimmer H. 2013. Opposite effects of visual and auditory word-likeness on activity in the visual word form area. Front Hum Neurosci. 7:491.

Mano QR, Humphries C, Desai RH, Seidenberg MS, Osmon DC, Stengel BC, Binder JR. 2013. The role of left occipitotemporal cortex in reading: Reconciling stimulus, task, and lexicality effects. Cereb Cortex. 23:988–1001.

McCandliss BD, Cohen L, Dehaene S. 2003. The visual word form area: Expertise for reading in the fusiform gyrus. Trends Cogn Sci. 7:293–299.

Millar S. 1984. Is There a "Best Hand" for Braille? Cortex. 20:75–87.

Millar S. 1987. The Perceptual "Window" in Two-Handed Braille: Do the Left and Right Hands Process Text Simultaneously? Cortex. 23:111–122.

Millar S. 2003. Reading by Touch, Reading by Touch. Routledge.

Nakamura K, Kuo WJ, Pegado F, Cohen L, Tzeng OJL, Dehaene S. 2012. Universal brain systems for recognizing word shapes and handwriting gestures during reading. Proc Natl Acad Sci U S A. 109:20762–20767.

Ossmy O, Ben-Shachar M, Mukamel R. 2014. Decoding letter position in word reading. Cortex. 59:74–83.

Ossowski A, Behrmann M. 2015. Left hemisphere specialization for word reading potentially causes, rather than results from, a left lateralized bias for high spatial frequency visual information. Cortex. 72:27–39.

Pascual-Leone A, Cammarota A, Wassermann EM, Brasil-Neto JP, Cohen LG, Hallett M. 1993. Modulation of motor cortical outputs to the reading hand of braille readers. Ann Neurol. 34:33–37.

Pascual-Leone A, Torres F. 1993. Plasticity of the sensorimotor cortex representation of the reading

finger in braille readers. Brain. 116:39–52.

Rączy K, Urbańczyk A, Korczyk M, Szewczyk JM, Sumera E, Szwed M. 2019. Orthographic Priming in

Braille Reading as Evidence for Task-specific Reorganization in the Ventral Visual Cortex of the

Congenitally Blind. J Cogn Neurosci. 31:1065–1078.

Ratan Murty NA, Teng S, Beeler D, Mynick A, Oliva A, Kanwisher N. 2020. Visual experience is not

necessary for the development of face-selectivity in the lateral fusiform gyrus. Proc Natl Acad Sci.

117:23011–23020.

Reich L, Szwed M, Cohen L, Amedi A. 2011. A ventral visual stream reading center independent of visual

experience. Curr Biol. 21:363–368.

Röder B, Rösler F, Neville HJ. 2000. Event-related potentials during auditory language processing in

congenitally blind and sighted people. Neuropsychologia. 38:1482–1502.

Röder B, Stock O, Bien S, Neville H, Rösler F. 2002. Speech processing activates visual cortex in

congenitally blind humans. Eur J Neurosci. 16:930–936.

Rueckl JG, Paz-Alonso PM, Molfese PJ, Kuo WJ, Bick A, Frost SJ, Hancock R, Wu DH, Einar Mencl W,

Duñabeitia JA, Lee JR, Oliver M, Zevin JD, Hoeft F, Carreiras M, Tzeng OJL, Pugh KR, Frost R.

2015. Universal brain signature of proficient reading: Evidence from four contrasting languages.

Proc Natl Acad Sci U S A. 112:15510–15515.

Ruschel M, Knösche TR, Friederici AD, Turner R, Geyer S, Anwander A. 2014. Connectivity architecture

and subdivision of the human inferior parietal cortex revealed by diffusion MRI. Cereb Cortex.

24:2436–2448.

Saalmann YB, Pigarev IN, Vidyasagar TR. 2007. Neural Mechanisms of Visual Attention: How Top-Down

Feedback Highlights Relevant Locations. Science. 316:1612–1615.

Sadato N, Pascual-Leone A, Grafman J, Deiber MP, Ibañez V, Hallett M. 1998. Neural networks for

Braille reading by the blind. Brain. 121:1213–1229.

Sadato N, Pascual-Leone A, Grafman J, Ibañez V, Deiber M, Dold G, Hallett M. 1996. Activation of the

primary visual cortex by Braille reading in blind subjects. Nature. 380:526–528.

Saygin ZM, Osher DE, Norton ES, Youssoufian DA, Beach SD, Feather J, Gaab N, Gabrieli JDE,

2   Kanwisher N. 2016. Connectivity precedes function in the development of the visual word form

3   area. Nat Neurosci. 19:1250–1255.

4   Schlaggar BL, McCandliss BD. 2007. Development of neural systems for reading. Annu Rev Neurosci.

5   30:475–503.

6   Seghier ML, Price CJ. 2011. Explaining left lateralization for words in the ventral occipitotemporal cortex.

7   J Neurosci Off J Soc Neurosci. 31:14745–14753.

8   Siuda-Krzywicka K, Bola Ł, Paplińska M, Sumera E, Jednoróg K, Marchewka A, Śliwińska MW, Amedi A,

9   Szwed M. 2016. Massive cortical reorganization in sighted braille readers. eLife. 5:1–26.

10   Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, Bannister PR,

11   De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N,

12   Brady JM, Matthews PM. 2004. Advances in functional and structural MR image analysis and

13   implementation as FSL. NeuroImage. 23:208–219.

14   Stevens WD, Kravitz DJ, Peng CS, Tessler MH, Martin A. 2017. Privileged functional connectivity

15   between the visual word form area and the language system. J Neurosci. 37:5288–5297.

16   Striem-Amit E, Cohen L, Dehaene S, Amedi A. 2012. Reading with Sounds: Sensory Substitution

17   Selectively Activates the Visual Word Form Area in the Blind. Neuron. 76:640–652.

18   Taylor JSH, Rastle K, Davis MH. 2013. Can cognitive models explain brain activation during word and

19   pseudoword reading? A meta-analysis of 36 neuroimaging studies. Psychol Bull. 139:766–791.

20   Tian M, Li H, Chu M, Ding G. 2020. Functional organization of the ventral occipitotemporal regions for

21   Chinese orthographic processing. J Neurolinguistics. 55:100909.

22   Twomey T, Kawabata Duncan KJ, Price CJ, Devlin JT. 2011. Top-down modulation of ventral occipito-

23   temporal responses during visual word recognition. NeuroImage. 55:1242–1251.

24   Van der Haegen L, Cai Q, Brysbaert M. 2012. Colateralization of Broca's area and the visual word form

25   area in left-handers: FMRI evidence. Brain Lang. 122:171–178.

26   Vinckier F, Dehaene S, Jobert A, Dubus JP, Sigman M, Cohen L. 2007. Hierarchical Coding of Letter

27   Strings in the Ventral Stream: Dissecting the Inner Organization of the Visual Word-Form System.

28   Neuron. 55:143–156.

1 Wilke M, Schmithorst VJ. 2006. A combined bootstrap/histogram analysis approach for computing a

2   lateralization index from neuroimaging data. NeuroImage. 33:522–530.

3 Yeatman JD, Rauschecker AM, Wandell BA. 2013. Anatomy of the visual word form area: Adjacent

4   cortical circuits and long-range white matter connections. Brain Lang. 125:146–155.