Reading Modality Modifies Reading Network:

Insights from Neural basis of

Braille in Proficient Blind Readers

Mengyu Tian[1*], Elizabeth J. Saccone[1], Judy S. Kim[1,2], Shipra Kanjlia[1,3], Marina Bedny[1]

[1] Department of Psychological and Brain Sciences, Johns Hopkins University, Baltimore, Maryland 21218

[2] Department of Psychology, Yale University, New Haven, Connecticut 06511

[3] Department of Psychology, Carnegie Mellon University, Pittsburgh, PA 15213

[*]To whom correspondence may be addressed: Mengyu Tian, Email: mengyutian@jhu.edu

Abstract

The neural basis of reading is highly consistent across a variety of languages and visual scripts. An unanswered question is whether the sensory modality of symbols influences the neural basis of reading. According to the modality-invariant view, reading depends on the same neural mechanisms regardless of the sensory input modality. Consistent with this idea, previous studies find that the visual word form area (VWFA) within the ventral occipitotemporal cortex (vOTC) is active when blind individuals read Braille by touch. However, connectivity-based theories of brain function suggest that the neural entry point of written symbols (touch vs. vision) may influence the neural architecture of reading. We compared the neural basis of the visual print (sighted n=15) and tactile Braille (congenitally blind n=19) in proficient readers using analogous reading and listening tasks. Written stimuli varied in word-likeness from real words to consonant strings and non-letter shape strings. Auditory stimuli consisted of words and backward speech sounds. Consistent with prior work, vOTC was active during Braille and visual reading. However, in sighted readers, visual print elicited a posterior/anterior vOTC word-form gradient: anterior vOTC preferred larger orthographic units (words), middle vOTC preferring consonant strings, and posterior vOTC responded to shapes (i.e., lower-level physical features). No such gradient was observed in blind readers of Braille. Consistent with connectivity predictions, in blind Braille readers, posterior parietal cortices (PPC) and parieto-occipital areas were recruited to a greater degree and PPC contained word-preferring patches. Lateralization of Braille in blind readers was predicted by laterality of spoken language, as well as by reading hand. These results suggested that the neural basis of reading is influenced by symbol modality and support connectivity-based views of cortical function.

Highlights

1. Only sighted but not blind (Braille) readers show a posterior/anterior vOTC lexicality gradient
2. Posterior parietal cortex distinctively contributes to Braille reading.
3. Lateralization of spoken language and reading hand predict lateralization of Braille
4. The sensory modality of written symbols influences the neural basis of reading

Keywords

Braille reading; blindness; reading; VWFA; posterior parietal cortex

Introduction

Written language is among the most impressive human cultural achievements. The capacity to record and transmit information over space and time has enabled the accumulation of scientific, technological, and historical knowledge across generations and continents. How does the human brain accommodate this cultural invention, which emerged only approximately 5,000 years ago?

Despite being a recent cultural invention, the neural basis of reading is highly consistent across a variety of languages and visual scripts, including alphabetic, logographic (e.g., Chinese), and syllabic writing systems (e.g., Japanese Kana) (Bolger, Perfetti, & Schneider, 2005; Feng et al., 2020; Hu et al., 2010; Krafnick et al., 2016; Nakamura et al., 2012; Rueckl et al., 2015). All of these reading systems engage regions within the left lateral ventral occipitotemporal cortex (vOTC) (Baker et al., 2007; Cohen et al., 2000; Dehaene & Cohen, 2011; Dehaene et al., 2010). A region in the left lateral vOTC has been termed the 'visual word form area' (VWFA) because of its preferential response to written words and letter combinations over other visual stimuli. The VWFA is situated within a posterior/anterior processing gradient. During reading, visual symbols are first processed by early visual cortices and posterior portions of vOTC, which represent simple visual features (e.g., line junctions) (Dehaene, Cohen, Sigman, & Vinckier, 2005; DiCarlo & Cox, 2007). By contrast, the middle and anterior potions of lateral vOTC are specialized for progressively larger orthographic units, from written letters, letter combinations/bigrams, and finally whole words (Binder, Medler, Westbury, Liebenthal, & Buchanan, 2006; Cohen et al., 2000; Dehaene, Cohen, Sigman, & Vinckier, 2005; Dehaene et al., 2004; Glezer, Jiang, & Riesenhuber, 2009; Lerma-Usabiaga, Carreiras, & Paz-Alonso, 2018; Purcell, Shea, & Rapp, 2014; Vinckier et al., 2007).

An open question is whether the vOTC posterior/anterior processing stream is the only way for the brain to implement reading and, relatedly, why the neural basis of reading takes this particular form. Examining the neural basis of tactile Braille offers unique insights into these questions. Specifically, we can ask whether and how the sensory modality of written symbols influences the neural basis of reading.

Tactile Braille reading achieves similar behavioral goals for people who are blind as visual print

reading does for the sighted: rapid access to linguistic meaning from a temporally stable

symbolic record. Proficient blind readers can read upwards of 200 words per minute by passing

the fingers along lines of Braille text, in which words are written as patterns of raised dots

(Millar, 2003). Each Braille character consists of dots positioned in a three-rows-by-two-

columns matrix. A single Braille character can be used to represent a letter, number, or

punctuation mark. In the most commonly used form of English Braille (Grade 2 Braille), Braille

characters also stand for frequent letter combinations (e.g., EA ∴, OW ∵) and whole words (e.g.,

e = every, tm = tomorrow) (http://www.brl.org) (Millar, 2003).

Consistent with a modality-invariant view of reading, several recent studies have reported that

the neural basis of Braille reading and that of visual print reading depend on similar vOTC

mechanisms (Büchel, Price, & Friston, 1998a; Debowska et al., 2016; Rączy et al., 2019; Reich,

Szwed, Cohen, & Amedi, 2011). Visual print and tactile Braille reading elicit activation peaks at

the anatomical location of the 'VWFA' in both sighted and blind readers (Debowska et al., 2016;

Dzięgiel-Fivet et al., 2021; Kim, Kanjlia, Merabet, & Bedny, 2017; Rączy et al., 2019; Reich et

al., 2011; Siuda-Krzywicka et al., 2016). In sighted adults who are trained to recognized Braille

words, transcranial magnetic stimulation (TMS) to the VWFA disrupts reading accuracy (Bola et

al., 2019; Siuda-Krzywicka et al., 2016). A recent study also found similar repetition suppression

effects in vOTC for tactile (blind readers) and visual (sighted readers) pseudowords (Rączy et al.,

2019). This evidence supports the idea that reading depends on the same neural mechanisms in

vOTC, regardless of symbol modality (i.e., touch vs. vision).

At the same time, both theoretical considerations and empirical evidence suggest that the neural

basis of tactile Braille and visual print reading may differ in important ways that have not been

fully tested. In sighted readers, posterior portions of vOTC receive visual written forms from

early visual cortices and pass this information along the posterior/anterior orthographic gradient

(e.g., Dehaene et al., 2005). By contrast, in people who are blind, Braille information enters the

cortex at primary somatosensory cortex (S1), making a posterior/anterior gradient unlikely. A

number of imaging studies also find that Braille reading activates visual areas outside of vOTC

in people who are blind, including V1 and dorsal occipital areas (Cohen et al., 1997, 1999;

Gizewski, Gasser, De Greiff, Boehm, & Forsting, 2003; Kupers et al., 2007; Melzer et al., 2001;

Sadato et al., 1998, 1996). TMS to the occipital pole and midoccipital cortex disrupts Braille

reading (Cohen et al., 1997, 1999; Kupers et al., 2007). This suggests that vOTC may not make a

unique contribution to Braille reading in the same way that it does to visual reading.

Moreover, visual cortices of people who are born blind, including vOTC and early visual areas

(V1-V3), are recruited for non-visual functions apart from Braille (e.g., Amedi, Raz, Pianka,

Malach, & Zohary, 2003; Büchel et al., 1998b; Burton, Snyder, Diamond, & Raichle, 2002;

Gougoux, Zatorre, Lassonde, Voss, & Lepore, 2005; Kanjlia, Loiotile, Harhen, & Bedny, 2021;

Kujala, Alho, Paavilainen, Summala, & Naatanen, 1992; Sathian, 2005). Particularly relevant for

the neural basis of Braille, large swaths of blind 'visual' cortex, including portions of V1,

participate in processing spoken language, including high-level semantic and grammatical

information (Bedny, Pascual-Leone, Dodell-Feder, Fedorenko, & Saxe, 2011; Bedny,

Richardson, & Saxe, 2015; Burton, Snyder, Diamond, & Raichle, 2002; Noppeney, Friston, &

Price, 2003; Röder, Stock, Bien, Neville, & Rösler, 2002; Watkins et al., 2012). Indeed, there is

evidence that the anatomical location of the 'VWFA' shows larger responses to spoken language

and responds to the grammatical structure of spoken sentences in people who are blind, more so

than in people who are sighted (Dzięgiel-Fivet et al., 2021; Kim et al., 2017). This pattern

suggests possible involvement in high-order language processing, rather than a reading-specific

role in blindness. Furthermore, since the anatomical distribution of written language is believed

to be influenced by the anatomical distribution of spoken language (Behrmann & Plaut, 2013;

Hannagan & Grainger, 2013; Saygin et al., 2016; Stevens, Kravitz, Peng, Tessler, & Martin,

2017a), recruitment of the visual cortex for language processing could itself modify the neural

basis of Braille reading in blind people. For example, we might expect Braille to recruit occipital

regions that are connected to visual networks recruited for spoken language. Together this

evidence suggests that the anatomical distribution and function role of visual cortices in blind

Braille readers and sighted visual readers may not be equivalent and merits further investigation.

There are also reasons to hypothesize that tactile Braille reading may differentially recruit

networks outside of the visual system, specifically the posterior parietal cortex (PPC). The vOTC

occupies a key connectivity position in sighted readers, in that it is connected to visual input on

the one hand and linguistic representations on the other (Barttfeld et al., 2018; Bouhali et al.,

2014; Hannagan, Amedi, Cohen, Dehaene-Lambertz, & Dehaene, 2015; Li, Osher, Hansen, &

Saygin, 2020; Saygin et al., 2016; Stevens, Kravitz, Peng, Tessler, & Martin, 2017b; Yeatman,

Rauschecker, & Wandell, 2013). The PPC arguably occupies an analogous connectivity-based

position for tactile Braille. Not only is PPC anatomically proximal and densely connected to

early somatosensory cortices (SMC) but like anterior/lateral vOTC, it is connected to language

and working memory systems (Burks et al., 2017; Duhamel, Colby, & Goldberg, 1998; Kaas,

2012; Lewis & Van Essen, 2000; Ruschel et al., 2014). Analogous to the functional role of

vOTC in visual shape recognition, the PPC furthermore plays a key role in tactile shape and

texture perception, pertinent to Braille recognition (Bauer et al., 2015; Hegner, Lee, Grodd, &

Braun, 2010). For example, stronger PPC activity is observed during tactile pattern

discrimination compared to vibrotactile detection (Hegner et al., 2010). We therefore

hypothesized that portions of PPC may specialize for tactile Braille letter and word recognition,

analogous to specialization for visual word form recognition within the vOTC of sighted print

readers. To our knowledge, the hypothesis of selective responses to Braille words in PPC has not

previously been tested. Although previous studies have examined activity in early SMC and

found expanded finger representations in proficient Braille readers, there is no evidence that this

plasticity reflects specialization for Braille letters and words (Burton, Snyder, Conturo, et al.,

2002; Burton, Sinclair, & McLaren, 2004; Kupers et al., 2007; Pascual-Leone et al., 1993;

Pascual-Leone & Torres, 1993; Sadato et al., 1998). One goal of the current study was therefore

to test whether any portion of PPC shows preferential responses to Braille letters and words in

blind readers of Braille, akin to specialization for visual letters and words found in vOTC of

sighted readers.

Finally, we hypothesized that lateralization patterns of Braille (blind) and visual print (sighted)

reading would be analogous but distinct. The reading network is typically strongly left-

lateralized in sighted people, like the spoken language network (Behrmann & Plaut, 2020;

Ossowski & Behrmann, 2015; Schlaggar & McCandliss, 2007; Seghier & Price, 2011; Vinckier

et al., 2007). Studies with sighted people who have right-lateralized spoken language responses

find that reading 'follows' spoken language into the right hemisphere (Behrmann & Plaut, 2020;

Cai, Lavidor, Brysbaert, Paulignan, & Nazir, 2008; Cai, Paulignan, Brysbaert, Ibarrola, & Nazir,

2010; Cai & Van der Haegen, 2015; Van der Haegen, Cai, & Brysbaert, 2012). In people who

are blind, left-lateralization of spoken language is reduced and highly variable across individuals

(Lane et al., 2017; Röder, Rösler, & Neville, 2000; Röder et al., 2002) We therefore

hypothesized that responses to Braille would be likewise less left-lateralized in blind readers and

would show co-lateralization with spoken language across individuals.

A further potential determining factor of Braille lateralization that does not arise for visual print,

is reading hand. In visual reading, information typically enters through both eyes and is projected

to both hemispheres. By contrast, in the case of Braille, it is possible for the information to enter

the left or the right hemisphere first, depending on the reading hand. Reading hand preferences

and reading styles differ widely across proficient blind Braille readers (Millar, 1984, 2003).

Many blind readers use both hands during naturalistic reading, however, one hand is thought to

track position on the page, while the other is used for word recognition (Millar, 2003). We

hypothesized that during single hand Braille reading, lateralization in early somatosensory

cortices would depend on which hand was used during word recognition, but that the effect of

reading hand would weaken in posterior parietal reading regions and would disappear in

language regions (Lane et al., 2017).

To test these predictions, we compared the neural basis of reading in proficient congenitally

blind and sighted readers using analogous reading and spoken language tasks. In the reading

tasks, participants were presented with words, consonant strings, and non-letter shapes/false

fonts. Reading stimuli were visual (print) for the sighted participants and tactile (Braille) for the

blind participants. In the spoken language task, both groups listened to audio words and

backward speech sounds. First, we tested the prediction that there is a posterior-to-anterior

gradient in preference from false-fonts to consonant strings and finally words in the vOTC of

sighted but not blind readers. Previous studies find that posterior vOTC responds as much or

more to false fonts as to letters and words, with only a small lateral/anterior portion (so-called

VWFA) being selective to written words and letters (Vinckier et al., 2007). By contrast, we

predicted that in blind readers, the entire extent of vOTC would show a preference for words,

consistent with its involvement in language processing (Kim et al., 2017; Lane, Kanjlia, Omaki,

& Bedny, 2015; Röder et al., 2002; Watkins et al., 2012). Next, we tested the hypothesis that the

PPC of blind Braille readers shows a functional profile analogous to the vOTC of the sighted:

selective responses to written words as opposed to tactile shapes in a subset of PPC, surrounded

by equal or greater responses to tactile shapes. We compared responses in PPC with those of

early SMC, where we would expect larger or equal responses to tactile shapes. Moreover, we

hypothesized that regions of PPC most distal from S1 and posterior to it are most likely to show

specialization for Braille letters and words, an anterior/posterior gradient analogous to the

posterior/anterior gradient observed in the vOTC of sighted readers. We also examined responses

across groups in other cortical areas previously implicated in reading: left inferior frontal cortex

(IFC) and primary visual cortex (V1) and used whole-cortex analyses to quantify the anatomical

distribution of visual and Braille reading (Burton, Snyder, Conturo, et al., 2002; Harold Burton,

Sinclair, & Agato, 2012; Rueckl et al., 2015; Sadato et al., 1998). Finally, we used laterality

index analyses to compare lateralization patterns across written and spoken word comprehension

in the two groups. We tested the prediction that lateralization of reading would be driven by the

lateralization of spoken language in higher-order language regions (left IFC), by reading hand in

early SMC, and by both factors in reading-related areas (PPC).

Method

Participants

Nineteen congenitally blind (12 females, mean age = 40.36 years, SD = 14.82) and 15 sighted

control (9 females, mean age = 23 years, SD = 6) participants took part in the task-based fMRI

experiment (see Table 1 for participant characteristics). The data from 10 blind and 15 sighted

participants have been reported previously (Kim et al., 2017). All participants were native

English speakers, and none had suffered from any known cognitive or neurological disabilities

(screened through self-report). Sighted participants had normal or corrected to normal vision. All

the blind participants had at most minimal light perception from birth. Blindness was caused by

pathology anterior to the optic chiasm (i.e., not due to brain damage). All blind participants were

fluent Braille readers who began learning Braille at an average age of 4.6 years (SD = 1.49) and

rated their reading ability as proficient to expert (mean = 4.57, SD =0.69 on a scale of 1 to 5) and

reported reading on average 20 hours per week (SD=19). We obtained information on Braille-

reading hand dominance, whether they read bimanually, and reading frequency through a post-

experimental survey conducted over the telephone with 17 of the 19 blind adult participants

(Table 1). All participants gave informed consent according to procedures approved by the Johns

Hopkins Medicine Institutional Review Board.

Table 1 Participants information

Participant no.	Age (y)	Gender	Handedness	Reading handedness	Levels of education	Cause of blindness	Age started reading Braille (y)	Reading hours per week	Self-reported Braille reading ability (1-5)
B1	21	M	L	Bi-R	SC	LCA	4	14	5
B2	64	F	R	Bi-R	BA	ROP	6	56	5
B3	53	M	R	Bi-R	JD	LCA	6	7	4
B4	34	M	R	L	SC	Born without optic nerve	3	21	5
B5	42	M	Am	L	BA	ROP	3	21	5
B6	29	M	R	Bi-L	SC	LCA	4	<1	4
B7	39	F	R	L	BA	ROP	4	2	5
B8	34	F	R	--	SC	Optic Nerve Detached	3	--	5
B9	49	M	R	Bi-R	BA	unknown	8	<1	3
B10	26	F	R	Bi-R	MA	ROP	3	56	3
B11	49	F	L	R	MA	LCA	7	14	5
B12	39	F	R	L	MA	ROP	5	14	5
B13	35	F	R	Bi-L	MA	LCA	4	14	5
B14	46	F	R	--	BA	ROP	4	--	5
B15	33	F	R	L	BA	ROP	4	14	4
B16	25	F	Am	Bi-R	MA	LCA	5	56	5
B17	23	M	R	Bi-R	BA	LCA	4	28	5
B18	70	F	R	R	HS	ROP	7	7	4
B19	68	F	R	Bi-R	MA	ROP	5	7	5
Average									
Blind (n=19)	41 (SD=14.82)	12F	2L/2Am	--	BA	--	4.68 (SD=1.49)	19.47 (SD=18.97)	4.57 (SD=0.69)
Sighted (n=15)	23 (SD=6)	9F	1 L	--	BA	--	--	--	--

Handedness: left (L), ambidextrous (Am), or right (R), based on Edinburgh Handedness Inventory. BA = Bachelor of Arts; MA = Master of Arts; HS = High School; JD = Juris Doctor; SC = Some College; ROP = Retinopathy of prematurity; LCA = Leber's congenital amaurosis. For Braille ability, participants were asked: "On a scale of 1 to 5, how well are you able to read Braille, where 1 is 'not at all', 2 is 'very little', 3 is 'reasonably well', 4 is 'proficiently', and 5 is 'expert'?"

Stimuli

The fMRI task including reading and listening tasks (Figure 1). There were three stimulus conditions for the reading task: words, non-word consonant strings, and non-letter shapes (control condition). During the reading task, stimuli were visual for the sighted participants and

tactile for the blind participants. For the listening task, there were two conditions: words and

backward speech sounds (control condition).

The word stimuli consisted of 240 common nouns, verbs, and adjectives. For the tactile reading

task (blind group), the Braille words were written in Grade-II contracted English Braille, which

is the most common form of Braille in the United States. Braille characters contain between 1-6

raised pins in set positions within a 2 x 3 array (see Figure 1). In Grade-II contracted English

Braille, there are contractions such that single Braille characters represent frequent letter

combinations (e.g., "th") or frequent whole words (e.g., the "c" can stand for "can"). With

contractions, the Braille words were on average 4 Braille characters (range = 1-8 Braille

characters, SD = 2.1 characters) and 11 tactile pins per word. Note that each participant was

presented with 120 of the 240 words during the reading task; the other 120 words were presented

auditorily during the listening task (see below). The word lists were counterbalanced across

participants. In the tactile consonant string condition, there were 24 strings repeated 5 times

throughout the experiment. Each string stimulus consisted of 4 Braille letters, which were

created using 20 English consonants. Last, the tactile control stimuli consisted of 24 unique

strings of 4 non-letter shapes made of Braille pins (see Figure 1). Note that any dot array within a

2 x 3 grid could be part of a Braille character. Therefore, to prevent participants from processing

the shapes as Braille letters, the shapes varied in size and pin number within arrays ranging in

size from 4×5 to 7×7. The average number of Braille pins per string in the control condition

was 58.

For the sighted group, the word stimuli consisted of 240 common nouns, verbs, and adjectives

that were on average 4 letters long (range = 3-5 letters, SD = 0.7 letters). Visual word stimuli

consisted of a new set of words matched to the Braille words on average character length (i.e., 4

visual letters matched to 4 Braille characters), raw frequency per million, averaged frequency per

million of orthographic neighbors, and averaged bigram frequency (all comparisons $p > 0.4$,

obtained from the MCWord Orthographic Wordform Database; Medler & Binder, 2005).

Different groups of words were used for the visual and Braille experiment to enable character

length matching since Braille contractions represent two or more English letters with a single

Braille character. Like the blind participants, sighted participants encountered half (120) of the

words during reading trials and the other half during auditory trials, counterbalanced across

participants. The visual consonant strings were the same 24 consonant letter combinations from

the tactile consonant strings described above. Lastly, the control stimuli in the visual reading task

were 24 unique strings, each comprised of 4 characters, which were false fonts. There were 20

false font characters in total, which matched the 20 English consonants on the number of strokes,

presence of ascenders and descenders, and the stroke thickness.

The stimuli for the listening task were taken from each group's respective word list. For the

audio word condition, stimuli were 120 words taken from the reading task described above. For a

given word, half of the participants received it in the reading task and half received it in the

listening task. The auditory words were recorded by a female native English speaker. The

average word length was 5 letters long (SD = 1.4 letters). The average playtime for the auditory

stimuli was 0.41 s long (SD = 0.3 s). The control auditory stimuli comprised backward speech

sounds, which were created by playing each audio word in reverse.

Procedure

The experiment had a total of 5 runs, each with 20 task trials. In each trial, participants were

presented with a block of 6 stimuli from a single condition (e.g., tactile reading consonant strings

condition) and then performed a memory probe task. All stimulus conditions for both reading

and listening trials were presented in every run. Each condition was repeated 4 times per run, and

the order of conditions was counterbalanced across runs. There were 6 rest periods (16 s)

throughout each run. One sighted participant and two blind participants were excluded from

behavioral analysis due to failure to record their responses.

For the blind participants, each trial began with a 0.5 s auditory cue instructing participants to

"Touch' (reading trial), or "Listen" (listening trial). Then participants felt or heard blocks of 6

target items, one at a time. For 10 of the blind participants, tactile target stimuli were presented

on the Braille display for 2 s, followed by a 0.75 s inter-stimulus interval (ISI) (6-item list

duration: 16.5 s) (Kim et al., 2017). For the newly added 9 blind participants, the ISI was

lengthened to 1.75 s due to a coding error which caused the 6-item list duration to be prolonged

to 22.5 s. Control analyses revealed no effects of ISI duration on the results and the data are

henceforth combined. After the 6-item list had been presented, there was a short delay (0.2 s),

followed by a beep (0.5 s). Then a probe stimulus (2 s) was then presented, and participants

indicated with a key press whether or not the probe had been present in the list. Participants had

5.3 s to make a response. The participants were asked to read with their dominant hand and

responded with the other hand. The listening task was analogous in format to the reading task.

The audio words and backward speech were on average 0.41 s long. The timing and sequence of

events were identical for the listening task (6-item list duration 16.5 s).

For sighted participants, the trial event sequence (cue, 6-item block, beep, probe, response) was

analogous to above. Each trial began with an auditory cue instructing participants to "Look"

(reading trial) or "Listen" (listening trial). During reading trials, 6 visual stimuli appeared

centrally for 1 s each, followed by an ISI of 0.75, during which participants were asked to

maintain gaze on a black central fixation cross (total block duration: 10.5 s). Note that visual

reading blocks were shorter than tactile reading blocks for the blind participants because pilot

testing indicated that visual reading is faster under these conditions. Listening trials also had a

total stimulus block duration of 10.5 s, to be consistent with the reading trials within the sighted

group.

fMRI data acquisition

Functional and structural images were acquired using a 3T Phillips scanner at the F. M. Kirby

Research Center. T1-weighted images were collected using a magnetization-prepared rapid

gradient-echo (MP-RAGE) in 150 axial slices with 1 mm isotropic voxels. Functional BOLD

scans were collected T1-weighted structural images were collected in 150 axial slices with 1 mm

isotropic voxels. Functional BOLD scans were collected in 36 sequential ascending axial slices.

TR = 2 s, TE = 0.03 s, flip angle = 70°, voxel size = 2.4 × 2.4 × 2.5 mm, inter-slice gap = 0.5

mm, field of view (FOV) = 192 × 172.8 × 107.5. Acquisition parameters were identical for the

resting-state and task fMRI experiment.

fMRI data analysis

Preprocessing and whole-cortex analysis

Analyses were performed using FSL (version 5.0.9), FreeSurfer (version 5.3.0), the Human Connectome Project workbench (version 1.2.0), and custom in-house software. The cortical surface was created for each participant using the standard FreeSurfer pipeline (Dale, Fischl, & Sereno, 1999; Glasser et al., 2013; Smith et al., 2004). For task data, preprocessing of functional data included motion-correction, high-pass filtering (128 s cut-off), and resampling to the cortical surface. Cerebellar and subcortical structures were excluded. On the surface, the task data were smoothed with a 6 mm FWHM Gaussian kernel. Two runs for blind and three runs for sighted participants were dropped due to equipment failure.

The three conditions in the reading task and two conditions in the listening task were entered as covariates of interest into general linear models. Only the six-item period in each trial was entered into the model. Covariates of interest were convolved with a standard hemodynamic response function, with temporal derivatives included. Probe stimulus, response periods, and the trials in which participants failed to respond were entered as covariates of no interest. The effect of the mean signal of white matter and CSF, as well as the motion spike, were also included as the covariates of no interest. Runs were combined within subjects using fixed-effects models.

Data across participants were combined within groups using random-effects analysis. Reported whole-cortex contrasts were run thresholded at $p < 0.01$ vertexwise, and $p < 0.05$ cluster-corrected.

fMRI ROI analysis

Individual-subject functional regions of interest (ROIs) were defined within the vOTC and other regions previously implicated in Braille reading (V1), language (left inferior frontal cortex, IFC), and tactile perception (left posterior parietal cortex, PPC, and hand region of the left primary somatosensory-motor cortex, SMC). To construct the left vOTC search space, we first combined the left fusiform, inferior temporal, and lateral occipital parcels from Freesurfer's automated aparc parcellation and then excluded V1, V2 regions, and the vertices with y-axis greater than -30 (Lerma-Usabiaga et al., 2018). To test the posterior-to-anterior function gradient, the left

vOTC search space was divided to three portions: posterior (y < -64), middle (-48 > y > = -64),

and anterior portion (y > = -48). The search space in the right hemisphere was created by

flipping the left vOTC masks along the x-axis. The V1 search space was defined from a

previously published anatomical surface-based atlas (PALS-B12; Van Essen, 2005). The left

inferior frontal language (IFC) search space was defined by using a sentence vs. non-words

contrast (Fedorenko, Hsieh, Nieto-Castañón, Whitfield-Gabrieli, & Kanwisher, 2010). The

parietal search space was defined by the orthogonal contrast of all tactile conditions (words,

consonant strings, and control) > rest in whole-cortex analysis, excluding the occipital parcels

from Freesurfer's automated aparc parcellation. To look for lateralization effects in vOTC across

groups, we examined responses separately for the right and left hemispheres.

Individual-subject functional ROIs were defined in group-wise search spaces (described below).

Each individual subject's ROI was defined as the top 5% of vertices activated for the

tactile/visual consonant strings > tactile/visual controls contrast within the search spaces listed

above. We used this consonant string contrast for the primary analysis in order to focus on

orthographic as opposed to semantic responses. However, all analyses were also repeated using

the words > control contrast and results from these analyses are reported in the supplementary

material (Figure S3 and Figure S5). To avoid using the same data to define ROIs and to test

hypotheses, a leave-one-run-out cross-validation procedure was used. ROIs were defined based

on data from all but one run, then the percent signal change (PSC) was extracted from the left-

out run. This procedure was repeated iteratively across all runs and the PSC was averaged across

iterations.

Repeated-measured ANOVAs were used to analyze the ROI data, and two-tailed paired *t*-tests

were used for pairwise comparisons. All *p* values were Bonferroni-corrected for multiple

comparisons.

Topographical preference map

To explore the posterior-to-anterior gradient in left vOTC and in a data-driven way, we mapped

the topographical preference of the vOTC during reading using a winner-take-all approach. We

took the bilateral vOTC as the mask, and color-coded each vertex within the mask according to

which stimulus condition it responded most strongly. The topographical preference map of the

PPC and parieto-occipital/dorsal occipital cortex was created using the same winner-take-all

approach. The mask was defined by the orthogonal contrast of all tactile conditions (words,

consonant strings, and control) > rest in the whole-cortex analysis.

Laterality index analysis

To determine whether spoken and written language co-lateralize to the same hemisphere, we

performed laterality index (LI) analyses. LI was calculated separately for the reading and

listening tasks for each participant in the SMC, PPC, vOTC, V1, IFC, and also for the whole

cortex. For the reading task, LI was determined based on the tactile/visual words > rest contrast.

For the listening task, LI was determined using the audio words > rest contrast. The LI was

calculated using the standard formula: $(L - R) / (L + R)$, where L and R refer to the sums of the z

statistics from the relevant contrast within the left and right hemispheres, respectively. LI ranges

from -1 to 1, with a score of 1 indicating strong left lateralization and -1 strong right

lateralization.

The bootstrap/histogram method was used to ensure that LIs were not overly influenced by

arbitrary activation threshold choices or outlier voxels. Bootstrapped LIs were computed using

20 evenly spaced thresholds ranging from $z = 1.28$ to $z = 4.26$ (corresponding to one-sided $p =$

0.1 to $p = 0.00001$, uncorrected). For every threshold, each participant's z statistic map was

masked to only include the voxels exceeding the threshold within the search space. Then we

sampled the suprathreshold voxels 100 times with replacement in each hemisphere at a sampling

ratio k = 1.0. The LIs were then calculated using each pair of left and right hemisphere samples,

yielding a histogram of 10,000 threshold-specific LIs. Next, a single LI for each threshold was

calculated by averaging the values after removing the upper and lower 25% of the 10,000

threshold-specific values. Finally, the LI reported for each participant represents the average

across all thresholds.

A small number of participants were excluded from the LI analysis for a particular region if they

did not have suprathreshold activation in both hemispheres (listening task- SMC: 2 sighted, 2

blind participants excluded; PPC: 1 sighted; V1: 6 sighted; IFC: 1 sighted; reading task- SMC: 4

sighted; PPC: 1 sighted; IFC: 1 sighted).

393 To examine the effect of spoken language lateralization and Braille reading handedness on the

394 reading lateralization, a multiple regression was conducted for each region. The LI of spoken

395 words in IFC and dominant reading hand were entered as regressors and the LI of written words

396 was the dependent variable. Although some participants reported reading Braille bimanually, the

397 participants were asked to read tactile stimuli during the experiment only with their dominant

398 reading hand. There were 7 blind participants in the left Braille-reading handed group and 10 in

399 the right Braille-reading handed group.

Results

Behavioral Results

Higher accuracy and shorter reaction times for word-like stimuli

404 Because the two groups differed in age, we regressed out the effect of age on accuracy and

405 reaction times and performed analyses on the residuals (see Figure S1 in Supplementary

406 materials, results from raw data are also included in Figure S1). In the reading task, there was a

407 significant effect of age on accuracy (main effect of age, $F_{(1, 85)} = 5.681$, $p < 0.05$). A two-way

408 lexicality (words, consonant strings, control) by group (sighted, blind) ANOVA performed on

409 the residuals revealed higher accuracy on more word-like stimuli (words and consonant strings >

410 control) in both blind and sighted groups (main effect of lexicality: $F_{(2, 54)} = 13.963$, $p < 0.001$).

411 There was no lexicality by group interaction ($F_{(2, 54)} = 0.872$, $p = 0.737$). The group effect was

412 marginal (sighted > blind, $F_{(1, 27)} = 3.603$, $p = 0.068$). For the listening task, there was a trending

413 effect of age on accuracy ($F_{(1, 56)} = 2.907$, $p = 0.094$). A two-way lexicality (words, control) by

414 group (sighted, blind) ANOVA on the residuals revealed a lexicality effect (words > control; $F_{(1, 29)} = 50.944$, $p < 0.001$), no group effect ($F_{(1, 27)} = 0.843$, $p = 0.367$) or group by lexicality

416 interaction ($F_{(1, 27)} = 0.549$, $p = 0.465$).

417 Likewise, for reaction times during the reading task, there was a significant effect of age ($F_{(1, 85)}$

418 $= 39.089$, $p < 0.001$). A two-way lexicality (words, consonant strings, control) by group (sighted,

419 blind) ANOVA on the residuals revealed a lexicality effect (words and consonant strings <

control; $F_{(2, 54)} = 8.09$, $p < 0.001$). There was no group effect ($F_{(1, 27)} = 8.09$, $p = 0.297$). The

group by lexicality interaction effect was marginal ($F_{(2, 54)} = 2.763$, $p = 0.072$). Pairwise

comparisons showed the shorter reaction times on more word-like stimuli in blind group, but

there was no difference across stimuli in the sighted group (blind: words vs. control, $t_{(16)} = -2.91$,

$p < 0.01$; consonant strings vs. control, $t_{(16)} = -2.604$, $p < 0.01$; words vs. consonant strings, $t_{(16)} =$

-0.686, $p > 0.99$; sighted: all pairwise comparisons $p > 0.05$; the p-values were Bonferroni-

corrected).

During the listening task, the main effect of age on reaction time was significant ($F_{(1, 85)} =$

15.892, $p < 0.001$). A two-way lexicality (words, control) by group (sighted, blind) ANOVA on

the residuals revealed a lexicality effect (words < control; $F_{(1, 29)} = 50.944$, $p < 0.001$). There was

no group effect ($F_{(1, 27)} = 0.071$, $p = 0.792$) or group by lexicality interaction ($F_{(1, 29)} < 0.001$, $p >$

0.99).

fMRI Results

Visual (sighted) but not tactile Braille reading (blind) elicits a posterior-to-anterior functional

gradient in left vOTC and shows left-lateralization

Two signatures of visual reading responses in vOTC are 1) a posterior-to-anterior word form

gradient and 2) left-hemisphere lateralization. We asked whether Braille reading in blind

individuals shows similar posterior-to-anterior and laterality effects as visual reading in sighted

people. We divided the left and right vOTC each into the posterior, middle, and anterior

subregions (ROIs) and compared responses in these subregions across hemispheres and groups

(see Methods, Figure 1). We first conducted a four-way hemisphere (left, right) by

posterior/anterior subregion (posterior, middle, anterior) by lexicality (words, consonant strings,

control) by group (sighted, blind) ANOVA to examine reading responses across groups. This

ANOVA revealed a four-way interaction ($F_{(4, 128)} = 3.028$, $p < 0.05$), indicating that lexicality,

hemisphere, and posterior/anterior subregion interact with group. Next, we used separate

ANOVAs for each group to unpack the 4-way interaction. Because of the large number of factors

and to preserve readability, we report only hypothesis-relevant effects in this section. A complete

summary of all effects can be found in the Supplemental Materials.

For the sighted group, we found the expected three-way interaction between hemisphere (left,

right), posterior/anterior subregion (posterior, middle, anterior) and lexicality (words, consonant

strings, control; $F_{(4, 56)} = 4.287$, $p < 0.01$). Next, we looked at each hemisphere separately in the

sighted group.

In the left vOTC, there was a two-way interaction between lexicality (words, consonant strings,

control) and posterior/anterior subregion (posterior, middle, anterior; $F_{(4, 56)} = 9.69$, $p < 0.001$),

reflecting the expected posterior-to-anterior functional gradient. Pairwise comparisons revealed

that the posterior vOTC responded similarly to all visual stimuli (all pairwise comparisons $p >$

0.05). By contrast, in middle vOTC, consonant strings elicited higher responses than both words

and control stimuli (Bonferroni-corrected paired t-test for words vs. consonant strings: $t_{(14)} = -$

3.918, $p < 0.05$; consonant strings vs. control: $t_{(14)} = 4.106$, $p < 0.01$). In anterior vOTC,

responses to words and consonant strings were both higher than control and not different from

each other (Bonferroni-corrected paired t-test for words vs. control: $t_{(14)} = 3.461$, $p < 0.05$;

consonant strings vs. control: $t_{(14)} = 3.327$, $p < 0.05$, all other pairwise comparisons $p > 0.05$).

In the right vOTC of the sighted group, a two-way lexicality (words, consonant strings, control)

by posterior/anterior subregion (posterior, middle, anterior) ANOVA revealed no main effect of

lexicality ($F_{(2, 28)} = 0.448$, $p > 0.05$) and no interaction ($F_{(4, 56)} = 0.987$, $p > 0.05$). To

summarize, these results demonstrate that in the sighted group, there was a posterior-to-anterior

functional gradient for processing word form during reading in the left but not right vOTC.

Next, we examined these effects in the blind group. We conducted a three-way hemisphere (left,

right) by posterior/anterior subregion (posterior, middle, anterior) by lexicality (words,

consonant strings, control) ANOVA. Unlike in the sighted, there was no significant three-way

interaction ($F_{(4, 56)} = 0.877$, $p = 0.482$). Although there was no interaction, we conducted a

separate ANOVA testing for a lexicality effect across the posterior/anterior subregions for each

hemisphere separately in order to match the analysis of the sighted group.

In the left vOTC of the blind group, all three (posterior, middle, anterior) subregions responded

most to words, followed by consonant strings followed by tactile shapes (Figure 1). There was a

two-way interaction between lexicality (words, consonant strings, control) and posterior/anterior

subregion (posterior, middle, anterior; $F_{(4, 72)} = 3.198$, $p < 0.05$). However, the nature of this

interaction was different from that observed in the sighted group. All pairwise-comparisons

between conditions were significant in all three subregions (words > consonant strings >

control), except the difference between words and consonant strings did not reach significance in

the anterior vOTC (Bonferroni-corrected paired t-test for words vs. consonant strings: posterior

vOTC $t_{(18)} = 2.678$, $p < 0.05$; middle vOTC: $t_{(18)} = 3.166$, $p < 0.05$; anterior vOTC: $t_{(18)} = 2.016$, p

$= 0.177$; words vs. control: posterior vOTC: $t_{(18)} = 5.463$, $p < 0.001$; middle vOTC $t_{(18)} = 8.547$, p

< 0.001; anterior vOTC: $t_{(18)} = 5.874$, $p < 0.001$; consonant strings vs. control: posterior vOTC:

$t_{(18)} = 3.413$, $p < 0.01$; middle vOTC $t_{(18)} = 4.696$, $p < 0.01$; anterior vOTC: $t_{(18)} = 5.034$, $p <$

0.001).

Unlike in the sighted group, in the right hemisphere of the blind group, lexicality effects were

similar to the left hemisphere. All three (posterior, middle, anterior) subregions responded most

to words, followed by consonant strings followed by tactile shapes. There was also a two-way

interaction between lexicality (words, consonant strings, control) and subregion (posterior,

middle, anterior; $F_{(4, 72)} = 7.064$, $p < 0.001$). Pairwise comparisons showed that the posterior

right vOTC responded more to words than control ($t_{(18)} = 4.112$, $p < 0.01$); the middle vOTC

responded more to words than both consonant strings ($t_{(18)} = 4.011$, $p < 0.01$) and control ($t_{(18)} =$

4.819, $p < 0.001$); and the anterior vOTC responded most strongly to words and consonant

strings than control stimuli (words vs. consonant strings: $t_{(18)} = 2.429$, $p = 0.07$; words vs.

control: $t_{(18)} = 5.561$, $p < 0.001$; consonant strings vs. control, $t_{(18)} = 4.522$, $p < 0.01$). Other

pairwise comparisons did not reach significance (posterior vOTC: words vs. consonant strings,

$t_{(18)} = 2.349$, $p = 0.091$; consonant strings vs. control, $t_{(18)} = 2.16$, $p = 0.134$; middle vOTC:

consonant strings vs control, $t_{(18)} = 2.073$; $p = 0.159$).

In summary, in the blind group, the entire posterior/anterior extent of the vOTC responded more

to words than either consonant strings or tactile shapes. Unlike in the sighted, we did not observe

the posterior-to-anterior functional gradient or a left hemisphere dominance for written words.

For the listening task, similar to the reading task, we conducted a four-way hemisphere (left,

right) by subregion (posterior, middle, anterior) by lexicality (words, control) by group (sighted,

blind) ANOVA. The four-way interaction effect with group was marginal and we, therefore, did

not proceed to further analyses ($F_{(2, 64)} = 2.717$, $p = 0.074$). It is worth noting that in sighted

group, responses to auditory stimuli were below rest in posterior vOTC and above rest in the

more anterior regions. This pattern was not observed in the blind group (see Figure S2).

Topographical preference map of vOTC: gradient only in sighted readers

In order to explore the posterior-to-anterior gradient in a data-driven way, we mapped the

topographical preferences of the blind and sighted vOTC during reading using a winner-take-all

approach (Figure 1B). We coded the vertex-wise preferences in different colors for words,

consonant strings, and control stimuli (see Methods). In the sighted group, a clear posterior-to-

anterior gradient in the left vOTC was observed. The posterior section shows a preference for the

visual control false font stimuli whereas anteriorly, most vertices preferred consonant strings or

words. In the sighted group's right vOTC, almost all vertices responded most strongly to the

control stimuli. These patterns contrast starkly with the blind vOTC maps, which show a clear,

bilateral preference for tactile words throughout the entire extent of both left and right vOTC.

Figure 1. (A) Responses in left vOTC across the posterior, middle, and anterior subregions for
blind and sighted groups during the reading tasks (left column). Bars show results from
consonant string > false fonts leave-one-run out individual subject ROI analysis. Error bars
denote standard errors +/- the mean. Asterisks (*) denote significant Bonferroni-corrected
pairwise comparisons within the task ($p < 0.05$). The right upper row showed example stimuli for
the reading trials for the blind and sighted groups. (B) Maps of the blind and sighted vOTC
showing topographical preference to words, consonant strings, or control stimuli during the
reading task.

The posterior parietal cortex (PPC) but not S1 of blind readers shows a preference for written

Braille words and consonant strings

We tested the hypothesis that the PPC shows preferential involvement in Braille reading,

analogous to vOTC preference for visual print in the sighted group (Figure 2A). A two-way

lexicality (words, consonant strings, control) by group (sighted, blind) ANOVA in the reading

task showed a main effect of lexicality ($F_{(2, 64)} = 13.206$, $p < 0.001$) and a group by lexicality

interaction ($F_{(2, 64)} = 5.123$, $p < 0.01$; functional ROIs were defined using the words > controls

contrast). There was no main effect of group ($F_{(1, 32)} = 1.452$, $p = 0.237$). In the sighted group,

consonant strings elicited higher responses than both words and control stimuli (Bonferroni-

corrected paired t-test, words vs. consonant strings: $t_{(14)} = -3.805$, $p < 0.01$; consonant strings vs.

control: $t_{(14)} = 6.922$, $p < 0.001$; words vs. control: $t_{(14)} = 1.406$, $p > 0.99$). By contrast, in the

blind group the PPC responded more to both tactile words and consonant strings relative to

control stimuli (Bonferroni-corrected paired t-test, words vs. consonant strings: $t_{(18)} = 1.571$, $p =$

0.298; consonant strings vs. control: $t_{(18)} = 3.028$, $p < 0.01$; words vs. control: $t_{(18)} = 3.165$, $p <$

0.01). Note that when the posterior parietal ROI was defined instead using the words > controls

contrast, the blind group continued to show a larger lexicality preference than the sighted (see

Supplemental Materials for details; Figure S5). These results suggest a specific involvement of

the PPC in tactile Braille reading.

For the listening task, the two-way lexicality (words, control) by group (sighted, blind) ANOVA

revealed a significant main effect of lexicality in the PPC (words > control, $F_{(1, 32)} = 11.112$, $p <$

0.01; see Figure S4). There was no main effect of group ($F_{(1, 32)} = 3.275$, $p = 0.08$) and no

interaction between group and lexicality ($F_{(1, 32)} = 2.372$, $p = 0.133$).

We examined responses of the left SMC hand region to test whether it showed a similar

preference for Braille words and consonant strings as the PPC (Figure 2A). For the reading task,

the two-way lexicality (words, consonant strings, control) by group (sighted, blind) ANOVA

showed a main effect of lexicality ($F_{(2, 64)} = 7.265$, $p < 0.001$; functional ROIs were defined

using the words > controls contrast), with higher responses to the consonant strings than control

stimuli. Noting that the responses to all stimuli were below rest in SMC in blind group. There

was no main effect of group ($F_{(1, 32)} = 0.604$, $p = 0.443$) and no group by condition interaction (F

$(2, 64) = 1.501, p = 0.231$). For the listening task, the two-way lexicality (words, control) by group

(sighted, blind) ANOVA revealed a main effect of group ($F_{(1, 32)} = 15.622, p < 0.001$), with

overall greater responses in sighted group than blind group. There was no main effect of

lexicality ($F_{(1, 32)} = 1.933, p = 0.174$) and no interaction ($F_{(1, 32)} = 0.658, p = 0.423$). Results

were similar when the SMC ROIs were instead defined using the words > controls contrast. In

sum, unlike in the PPC, we found no evidence for specialization of SMC for Braille reading as

compared to perception of control tactile shapes.

Topographical preference map of parieto-occipital stream: shift in preference from shapes to

word-like Braille stimuli along anterior-to-posterior axis

Finally, we constructed a data-driven preference map in PPC and parieto-occipital/dorsal

occipital cortex analogous to the one created for vOTC (see Figure 2B). In the blind group, this

map shows preferential responses to tactile shapes in anterior portions of PPC, immediately

adjacent to S1. A small middle region in left and right PPC showed a preference for consonant

strings, whereas the most posterior portion of PPC, as well as parieto-occipital and dorsal

occipital regions responded preferentially to words. To summarize,   the overall pattern suggests

an anterior-to-posterior decoding pattern in the parieto-occipital stream in the blind group,

analogous to the posterior-to-anterior vOTC gradient observed in sighted readers.

Figure 2. (A) Responses in left PPC, and SMC ROIs for blind and sighted groups during the reading (blue colors) tasks. Error bars denote standard errors +/- the mean. Asterisks (*) denote significant Bonferroni-corrected pairwise comparisons within the task ($p < 0.05$). T = tactile, V = visual. (B) Maps of the blind and sighted vOTC showing topographical preference to words, consonant strings, or control stimuli during the reading task in the posterior parietal and parieto-occipital cortex. Black line: hand region of the primary sensory-motor cortex; yellow line: vertices showed the preference for consonant strings.

Left vOTC responds to linguistic stimuli in blind and sighted readers, but differently to words
and consonant strings across groups

We examined the effects of lexicality across groups on left vOTC responses during the reading

tasks using a two-way lexicality (words, consonant strings, control) by group (sighted, blind)

ANOVA (functional ROIs were defined using the words > controls contrast, Figure 3). We

observed a main effect of lexicality ($F_{(2, 64)} = 42.293$, $p < 0.001$) and no main effect of group (F

$_{(1, 32)} = 0.004$, $p = 0.948$). A lexicality by group interaction revealed different response patterns

across sighted and blind individuals ($F_{(2, 64)} = 10.272$, $p < 0.001$). While in the blind group

words elicited larger responses than consonant strings, responses to consonant strings were

numerically but not significantly larger than to words in the sighted group (Bonferroni-corrected

paired *t*-test words vs. consonant strings blind: $t_{(18)} = 3.027$, $p < 0.05$; sighted: $t_{(18)} = -1.317$, $p =$

0.614). In both groups, words and consonant strings showed larger responses than control stimuli

(all pairwise comparisons $p < 0.05$) (Figure 3).

For the listening task, a two-way lexicality (words, control) by group (sighted, blind) ANOVA

revealed greater overall responses to words than control stimuli (main effect of lexicality, $F_{(1, 32)}$

$= 35.919$, $p < 0.001$; functional ROIs were defined using the words > controls contrast). There

was no main effect of group ($F_{(1, 32)} = 1.362$, $p = 0.252$). The lexicality by group interaction was

marginal ($F_{(1, 32)} = 3.785$, $p = 0.061$), indicating a larger difference between audio words and

audio control stimuli in the blind group than in the sighted group. A similar pattern was observed

when the vOTC functional ROI was instead defined using the words > control contrast (see

Supplemental Materials; Figure S3).

The left inferior frontal cortex (IFC) prefers word-like written and spoken stimuli across blind
and sighted readers

We analyzed responses in the left IFC across groups with the prediction that this high-level

language region would show similar response patterns across blind and sighted readers.

Consistent with this prediction, responses were similar across groups for both tasks in the left

IFC. For the reading task, a two-way lexicality (words, consonant strings, control) by group

(sighted, blind) ANOVA revealed a significant main effect of lexicality, with larger responses

for words and consonant strings over the control condition ($F_{(2, 64)} = 46.313$, $p < 0.001$; Figure 3;

functional ROIs were defined using the words > controls contrast). Neither the main effect of

group ($F_{(1, 32)} = 0.004$, $p = 0.947$) nor the interaction ($F_{(2, 64)} = 1.017$, $p = 0.367$) were

significant. Likewise, for the listening task, the two-way lexicality (words, control) by group

(sighted, blind) ANOVA revealed the expected main effect of lexicality (words > control; $F_{(1, 32)}$

$= 23.778$, $p < 0.001$). There was no main effect of group ($F_{(1, 32)} = 0.753$, $p = 0.392$) and no

lexicality by group interaction ($F_{(1, 32)} = 0.357$, $p = 0.554$). There was also no group-by-

condition interaction when functional ROIs were defined using the words > controls contrast.

Both groups still showed a preference for words over control stimuli and in this case, there was

also a larger response to words over consonant strings in both groups (see Supplemental

Materials for details; Figure S5). These results are consistent with prior studies showing similar

responses to spoken and written language in the left inferior frontal cortex of blind and sighted

adults.

V1 shows a preference for words in blind readers

We investigated the effects of lexicality across groups in V1 (Figure 3), because it was

previously identified as relevant to Braille reading (Sadato et al., 1996; Cohen et al. 1997). As

with vOTC, we first examined responses in left V1 during the reading task using the consonant

strings > control functional ROIs. A two-way lexicality (words, consonant strings, control) by

group (sighted, blind) ANOVA revealed main effects of lexicality ($F_{(2, 64)} = 4.247$, $p < 0.05$) and

group (sighted > blind, $F_{(1, 32)} = 6.964$, $p < 0.05$). There was also a significant lexicality by group

interaction ($F_{(2, 64)} = 9.487$, $p < 0.001$). In the blind group, V1 responded most to words and

there was no difference between consonant strings and control (Bonferroni-corrected paired t-

test, words vs. consonant strings: $t_{(18)} = 2.641$, $p < 0.05$; words vs. control: $t_{(18)} = 3.691$, $p < 0.01$;

consonant strings vs. control: $t_{(18)} = 2.367$, $p = 0.214$). In the sighted group, V1 responded more

to control stimuli than consonant strings (Bonferroni-corrected paired t-test, $t_{(14)} = 2.652$, $p <$

0.01). There was no difference between other conditions (pairwise comparisons $p > 0.05$.) V1

responses in the blind group were similar when functional ROIs were defined using words >

control (see Supplemental Materials for details; Figure S5). In the sighted group, however, a

marginal preference for words over false fonts emerged in this alternative analysis (Bonferroni-

corrected paired t-test, $t_{(14)} = 2.573$, $p = 0.067$; Figure S5). This latter result is consistent with

some previous studies showing that V1/V2 responded more to words than non-letter control

stimuli like scrambled words (Szwed et al., 2011; Szwed, Qiao, Jobert, Dehaene, & Cohen,
2014).

For the listening task, the two-way lexicality (words, control) by group (sighted, blind) ANOVA
showed a main effect of group ($F_{(1, 32)} = 16.067$, $p < 0.001$), with overall greater activation seen
in blind than sighted V1. There was no main effect of lexicality ($F_{(1, 32)} = 2.344$, $p = 0.316$) and
no interaction between the factors ($F_{(1, 32)} = 1.589$, $p = 0.217$). Notably in the sighted but not
blind group, responses to both words and audio control were below rest. This pattern of results
was the same in words > control ROI (See Supplemental Materials for details; Figure S5).

Figure 3. Response of left vOTC (upper left), left IFC (upper right), left V1 (lower left) during
reading (blue colors) and listening (orange colors) tasks for blind (left) and sighted groups
(right). Bars show PSC for tactile stimuli (blind group), visual stimuli (sighted group), and audio
stimuli (both groups). Error bars denote standard errors +/- the mean. Asterisks (*) denote

653 significant Bonferroni-corrected pairwise comparisons within the task ($p < 0.05$). T = tactile, V =
654 visual.

Lateralization of Braille correlates with spoken language lateralization and Braille-reading
hand

We used a lateralization index (LI) analysis to investigate the lateralization of spoken and written
language across blind and sighted readers. First, we computed LIs separately for written
(tactile/visual words > rest) and spoken (audio words > rest) language in the SMC, PPC, vOTC,
V1, IFC and whole cortex in sighted and blind groups. On average, the blind group showed no
systematic lateralization for written or spoken words in any region (one-sample t tests of LI = 0,
reading: SMC: $t_{(18)} = 0.167$, $p = 0.869$; PPC: $t_{(18)} = -1.257$, $p = 0.225$; vOTC: $t_{(18)} = 0.799$, $p =$
0.435; V1: $t_{(18)} = 0.735$, $p = 0.472$; IFC: $t_{(18)} = -0.054$, $p = 0.958$; whole cortex: $t_{(18)} = -0.166$, $p =$
0.87; listening: SMC: $t_{(13)} = -1.332$, $p = 0.206$; PPC: $t_{(18)} = 0.051$, $p = 0.96$; vOTC: $t_{(18)} = 0.322$, p
= 0.751; V1: $t_{(18)} = -0.506$, $p = 0.619$; IFC: $t_{(18)} = -1.135$, $p = 0.271$; whole cortex: $t_{(18)} = 0.395$, p
= 0.697). For the sighted group, we found left-lateralized activation in vOTC, IFC and whole
cortex for written words (one-sample t tests of LI = 0, vOTC: $t_{(14)} = 5.31$, $p < 0.001$; IFC: $t_{(13)} =$
5.776, $p < 0.001$; whole cortex: $t_{(14)} = 5.748$, $p < 0.001$). The sighted group's SMC, PPC and V1
activity was not systematically lateralized for written words (one-sample t tests of LI = 0, SMC:
$t_{(10)} = 1.172$, $p = 0.268$; PPC: $t_{(13)} = 0.404$, $p = 0.692$; V1: $t_{(14)} = 1.614$, $p = 0.129$). For spoken
words, the sighted group's vOTC and IFC activity was left lateralized (one-sample t tests of LI =
0, vOTC: $t_{(14)} = 3.42$, $p < 0.01$; IFC: $t_{(13)} = 3.767$, $p < 0.01$). We found right-lateralized activation
in PPC and V1 for spoken words in the sighted group (one-sample t tests of LI = 0, PPC: $t_{(13)} = -$
3.161, $p < 0.01$; V1: $t_{(8)} = -3.872$, $p < 0.01$). There were no systematic lateralization in SMC and
whole cortex for the listening task (one-sample t tests of LI = 0, SMC: $t_{(13)} = -0.848$, $p = 0.412$;
whole cortex: $t_{(14)} = 1.449$, $p = 0.169$). To summarize, we found left-lateralized activity in vOTC
and IFC for written and spoken words in the sighted group. By contrast, the blind group did not
show systematic lateralization in any of the regions or the whole cortex for written or spoken
words. Among blind participants there was substantial variability in lateralization, with some
participants showing strong left and others strong right lateralization, consistent with previous
studies of lateralization of spoken language in this population (Figure 4, see also Lane et al.,
2017 and Roder et al., 2002).

Next, we determined if lateralization of the Braille reading network could be predicted by the

laterality of spoken language and Braille reading hand across blind individuals. A multiple

regression analysis was conducted in each region, with the LI of spoken words in IFC and

dominant reading hand entered as the regressors and the LI of written words as the dependent

variable. First, both the dominant reading hand and the LI of spoken words in IFC predicted the LI

of written words in PPC, vOTC and whole cortex (PPC: dominant reading hand: $\beta = 0.55$, $p <$

0.001; LI of spoken words in IFC: $\beta = 0.55$, $p < 0.001$; adjust $r^2 = 0.843$; vOTC: dominant reading

hand: $\beta = 0.468$, $p < 0.01$; LI of spoken words in IFC: $\beta = 0.611$, $p = 0.001$; adjust $r^2 = 0.727$; whole

cortex: dominant reading hand: $\beta = 0.399$, $p = 0.001$; LI of spoken words in IFC: $\beta = 0.534$, $p <$

0.001; adjust $r^2 = 0.761$). Second, in V1 and the IFC, only the LI of spoken words predicted the LI

of written words (V1: dominant reading hand: $\beta = 0.258$, $p = 0.144$; LI of spoken words in IFC: $\beta =$

0.734, $p = 0.001$; adjust $r^2 = 0.575$; IFC: dominant reading hand: $\beta = -0.112$, $p = 0.359$; LI of spoken

words in IFC: $\beta = 0.814$, $p < 0.001$; adjust $r^2 = 0.702$). Last, we found in SMC, only the dominant

reading hand predicted the LI of written words (dominant reading hand: $\beta = 1.624$, $p < 0.001$; LI

of spoken words in IFC: $\beta = 0.311$, $p = 0.261$; adjust $r^2 = 0.771$). To summarize, in blind

individuals, responses to Braille written words and spoken words were co-lateralized to the same

hemisphere across most of the Braille reading network, including the vOTC, V1, PPC, and the

IFC. Braille reading hand also had an effect on the lateralization of Braille written words in

vOTC, PPC, and SMC.

In the sighted group, we did not find the co-lateralization of spoken and written language to the

same hemisphere. The correlation between the LI of spoken words in IFC and the LI of written

words in vOTC was not significant ($r = -0.233$, $p = 0.423$). In addition, there were no

correlations between the LI of spoken words in IFC and the LI of written words in V1, SMC or

whole cortex (V1: $r = -0.301$, $p = 0.296$; SMC: $r = 0.169$, $p = 0.62$; whole cortex: $r = 0.12$, $p =$

0.683). However, the LI of spoken words in IFC was positively correlated with the LI of written

words in PPC and IFC (PPC: $r = 0.55$, $p < 0.05$; IFC: $r = 0.732$, $p < 0.01$).

Figure 4. (A) Correlations between the LIs of audio words in IFC and the LIs of tactile words in SMC (upper left), PPC (upper middle), vOTC (upper right), V1 (lower left), IFC (lower middle) and whole cortex (lower right) in blind individuals. Data points represent individual participants. LI 1 score indicates strong left lateralization and -1 indicates strong right lateralization. (B) Average of LI in left-handed blind readers of Braille and right-handed blind readers of Braille. Asterisks (*) on the bar denote significant difference from 0; asterisks (*) between two bars denote significant difference between the LI of left-handed blind readers and the LI of right-handed blind readers (p <0.05); Cross (†) on the bar denotes marginal difference from 0 (0.05< p <0.1); cross(†) between two bars denotes marginal difference between the LI of left-handed blind readers and the LI of right-handed blind readers (0.05< p <0.1).

Whole cortex analyses

Tactile Braille (blind) and visual print (sighted) reading activated both common and distinctive

cortical areas across groups. For reading as compared to rest, both sighted (visual words) and

blind (Braille words) readers activated the bilateral vOTC (blind peak: -41, -57, -13; sighted

peak: -41, -58, -12), including the location of the classic VWFA (peak: -46, -53, -20), as well as

early visual cortices, specifically the foveal confluence (V1/V2/V3) (Figure 5). Within vOTC,

responses in the blind group extended further medially and anteriorly and were more extensive in

the right hemisphere, relative to the sighted group. The vOTC activation in the blind group also

extended further laterally and superiorly, into lateral occipital, occipitotemporal, and inferior

temporal cortex. Both groups also activated posterior prefrontal cortices (inferior frontal gyrus

and middle frontal gyrus). Notably, visual cortex responses (e.g., V1) are likely driven at least in

part, by different processes across groups, since the sighed group is performing a visual task,

whereas the blind group is performing a tactile task.

In the blind but not sighted group, reading relative to rest produced extensive activation in

bilateral posterior parietal cortices, including superior parietal lobule and supramarginal gyrus

(SMG). This parietal activation was posterior to early sensory-motor hand representations. The

sighted group activated only a small cluster in parietal cortex, in the left superior parietal lobule.

The blind, but not sighted group, also activated parieto-occipital and dorsal occipital regions

(middle occipital gyrus). The sighted group additionally activated a lateral temporal region that

was not observed in the blind group. Finally, whereas responses to written words were left-

lateralized in the sighted group, they were bilateral in the blind group.

Listening to words (audio words > rest) likewise revealed partially overlapping responses across

groups. In the blind group only, listening to words activated the bilateral vOTC (peak: -42, -44, -

16), including the location of the classic VWFA, and early 'visual' cortices. Both groups

activated classic fronto-temporal language regions in inferior and lateral prefrontal as well as

lateral temporal cortices (Figure 5). Responses in frontal regions were left-lateralized in the

sighted group and bilateral in the blind group. The sighted but not blind group activated the left

sensorimotor cortex/postcentral gyrus.

Reading as compared to hearing words (tactile/visual words > audio words) also revealed

similarities and differences across groups (Figure 5). For the sighted group (visual words > audio

words), reading words induced greater activation in bilateral vOTC, including the typical

location of the VWFA and regions posterior to it, as well as bilateral early visual cortices. Like

the sighted, the blind group also activated a region in the left vOTC (fusiform; peak: -27, -61, -

14), but this activation was medial to the typical VWFA location. A cluster of activity was also

observed lateral to the typical VWFA location in the blind group, in the inferior temporal/lateral

occipital cortex (peak: -45 -67 -6). Outside vOTC, a cluster of activity was also observed in the

blind group in left foveal early 'visual' cortices. The blind but not sighted group also showed

extensive activation in posterior parietal cortices, including the SMG and superior parietal

lobule. Blind readers also activated dorsal occipital/parieto-occipital cortices during reading. The

blind group additionally activated the bilateral superior frontal gyrus and right precentral gyrus.

In the sighted group, a small cluster was observed in the left superior parietal lobule.

In sum, we observed the following pattern. First, although both groups activated vOTC during

reading, the peak location, distribution and functional profile of responses in vOTC were distinct

across groups. Only the blind group showed robust vOTC responses during spoken word

comprehension. Second, in contrast to the sighted group, the blind group activated extensive

posterior parietal, parieto-occipital, and dorsal occipital areas during (Braille) reading. This

parieto-occipital stream was not engaged by spoken word comprehension in blind readers.

Figure 5. Whole-cortex results for blind (left column) and sighted (right column) groups for the following contrasts: tactile/visual words > rest (top row), audio words > rest (middle row), tactile/visual words > audio words (bottom row). The blue circles marked the approximate location of the classic VWFA (MNI coordinate: -46, -53, -20). The yellow line marked the hand S1/M1 region. Whole cortex contrasts were thresholded at p < 0.01 vertex-wise, and at *p* < 0.05 cluster-corrected.

Discussion

Consistent with previous studies, we find that reading activates partially overlapping networks across blind readers of tactile Braille and sighted readers of visual print. In particular, we observed similar responses to written and spoken words and letters in the left IFC of sighted and blind people. We also observed partially overlapping responses in the vOTC across groups. In agreement with past findings, the highest peak of activation for Braille reading relative to rest and visual reading relative to rest was near the canonical 'VWFA' location (Braille words > rest: -41, -57, -13; sighted visual words > rest: -41, -58, -12) (Cohen et al., 2000; Dzięgiel-Fivet et al., 2021; Kim et al., 2017; Rączy et al., 2019; Reich et al., 2011). However, we also observed key differences in the neural bases of Braille and visual print reading, in vOTC, V1, and posterior parietal cortices, as well as in lateralization patterns.

vOTC of sighted but not blind readers contains a hierarchical word form gradient

Consistent with past research, in sighted readers, we observed a posterior-to-anterior functional gradient only in the left vOTC. The posterior portion of the left vOTC responded equally to all visual stimuli in the ROI analysis, the middle portion showed a preference for consonant strings, while the most anterior portion responded more to words and consonant strings than to false fonts. Preferential responses to consonant strings in the middle vOTC of sighted readers are consistent with prior literature showing stronger activation to non-word stimuli in the VWFA when longer presentation times are used or more attention is required (Bruno, Zumberge, Manis, Lu, & Goldman, 2008; Cohen, Dehaene, Vinckier, Jobert, & Montavont, 2008; Dehaene & Cohen, 2011; Ludersdorfer, Schurz, Richlan, Kronbichler, & Wimmer, 2013). A winner-take-all map revealed a similar pattern as the ROI analysis and further showed larger responses to false fonts than consonant strings or words in posterior portions of left vOTC. The larger responses to false fonts in posterior vOTC likely reflect greater attention to less familiar visual stimuli, as indicated by slower reaction times and poorer accuracy. This pattern is consistent with prior studies with sighted readers (Ludersdorfer et al., 2013; Wang, Yang, Shu, & Zevin, 2011). In addition, in sighted readers, the posterior vOTC showed a modality-specific response: above rest activity during visual reading and deactivation during listening, while the most anterior aspect of left vOTC responded equally to visual and auditory stimuli. These results are consistent with the

view that in sighted readers the middle and anterior portions of left lateral vOTC become

specialized for recognition of letters and words, constituting the so-called 'VWFA.'

By contrast, in the blind group, we found no evidence for left-lateralization, no evidence for a

posterior-to-anterior functional gradient, or posterior/anterior change in modality preference. In

blind readers, the entire posterior/anterior extent of bilateral vOTC showed a preference for

Braille words over consonant strings and tactile shapes during reading and a larger response to

spoken words than backward speech. Unlike in the sighted group, in the blind group, no portion

of the vOTC showed a consonant string preference over words and shapes, whereas in the

sighted group, the middle vOTC responded more to consonant strings than words. In addition,

there was no change in preference for written as opposed to spoken words along the

posterior/anterior extent of vOTC.

The Whole-cortex analysis also revealed differences in lateral/medial organization of vOTC

across groups. As noted above and previously documented, when Braille and spoken words were

compared to rest, a peak of activation was observed in the classic VWFA region along the

medial/lateral axis, although in blind readers, additional activity was also observed throughout

much of vOTC. By contrast, when Braille words were compared to spoken words in the blind

group, peak activity in the vOTC was medial to the classic VWFA location (peak: -27, -61, -14).

We did not observe such a medial peak for the same contrast in sighted readers. This medial

vOTC region has previously been shown to be functionally connected to dorsal parietal cortices,

which are involved in spatial attention and effortful letter-by-letter reading in sighted people

(Bouhali, Bézagu, Dehaene, & Cohen, 2019; Cohen et al., 2008; Corbetta & Shulman, 2002;

Henry et al., 2005; Saalmann, Pigarev, & Vidyasagar, 2007). As discussed in detail below, the

PPC appears to play an important role in Braille reading and may send information to medial

vOTC in blind readers.

Although the precise role of vOTC in Braille reading remains to be determined, the present

evidence suggests that although the vOTC is involved in both tactile reading and visual reading,

the anatomical distribution of responses within vOTC, the functional profile, and therefore likely

the cognitive contribution differs. In sighted readers, information reaches lateral vOTC from

early visual areas and is sent onward to fronto-temporal language regions, as well as receiving

top-down input from the language regions (Bouhali et al., 2014; Hannagan et al., 2015; Saygin et
al., 2016; Stevens et al., 2017b; Yeatman et al., 2013). Lateral vOTC thus contributes to
decoding linguistic information (phonological, semantic, and grammatical) from visual word
forms (Dehaene & Cohen, 2011; Price & Devlin, 2011).   By contrast, we hypothesize that in
blind readers of Braille, the classic VWFA location in lateral vOTC receives linguistic (i.e.,
semantic, grammatical) information from fronto-temporal language circuits and serves as one of
the entry points for language into posterior 'visual' circuits. This hypothesis is supported by prior
studies showing that in blind but not sighted people, the classic VWFA location is sensitive to
syntactic complexity of spoken sentences and shows enhanced responses to spoken language
(Burton, Snyder, Diamond, et al., 2002; Dzięgiel-Fivet et al., 2021; Kim et al., 2017; Lane et al.,
2015). At the same time, the current data and prior evidence suggest that other parts of the
'visual' cortex, including a medial portion of vOTC, may play a role in Braille reading. An
intriguing albeit speculative possibility is that medial vOTC receives Braille-relevant input from
PPC. Lacking connectivity data, the present study cannot test this hypothesis directly. One way
to test this possibility in future work would be to use online TMS in combination with fMRI to
disrupt information flow to the vOTC in blind readers of Braille by stimulating parietal cortices.

Parieto-occipital decoding stream in blind readers of Braille

We observed more extensive and different involvement of posterior parietal/parieto-occipital
cortices in Braille as opposed to visual print reading. Large segments of PPC were activated
during Braille reading relative to rest and spoken word comprehension. PPC activity in the blind
group extended inferiorly and anteriorly, into regions adjacent to and immediately posterior to
S1, including the supramarginal gyrus (SMG) and much of the superior parietal lobule. Notably,
the hand regions of S1 itself did not show robust responses during Braille reading or preferential
responses to Braille letters or words, consistent with prior studies (Burton, Snyder, Conturo, et
al., 2002; Kupers et al., 2007). Additionally, in the blind group only, parietal activation extended
posteriorly into parieto-occipital and dorsal occipital regions adjacent to parietal cortices and
ultimately into the foveal confluence. By contrast, visual print reading (relative to false fonts) by
sighted readers activated only a small region within the superior parietal lobule, consistent with
prior studies (Cohen et al., 2008; Martin, Schurz, Kronbichler, & Richlan, 2015; Reilhac, Peyrin,
Démonet, & Valdois, 2013).

The cognitive role of the wider parietal network in Braille reading is not known. The PPC has

strong connectivity with S1 and contains high-level tactile areas, as well as multimodal

representations of texture and shape (Bauer et al., 2015; Hegner et al., 2010; Kaas, 2012). Some

of the activation we observed likely reflects processes related to recognition of tactile patterns

that constitute Braille but are not specific to Braille letters or words (Boven, Hamilton,

Kauffman, Keenan, & Pascual-Leone, 2000; Wong, Gnanakumaran, & Goldreich, 2011), akin to

general responses to shapes, including false fonts, observed in vOTC of sighted readers (Grant,

Thiagarajah, & Sathian, 2000; Sathian & Stilla, 2010; Stilla et al., 2008). Consistent with this

possibility, much of the PPC, particularly its anterior portion, was more responsive to the more

tactilely complex and unfamiliar dot shapes than to Braille letters or words. Again, this paralleled

preferential responses to false fonts in posterior vOTC of sighted readers. Importantly, however,

within the larger swath of PPC activation, ROI analyses revealed word and letter preferring

subregions in the blind group, suggesting a specific involvement in Braille processing.

Word-specific activation in parieto-occipital areas extended posteriorly into dorsal occipital

cortices, only in the blind group. Unlike anterior portions of the PPC, parieto-occipital and dorsal

occipital areas showed larger responses to Braille words than Braille consonants or control

shapes. However, like anterior PPC, parieto-occipital and dorsal occipital regions responded

more to Braille words than to spoken words. This pattern suggests that parieto-occipital and

dorsal occipital areas are involved in reading-specific processing, rather than language

comprehension or tactile pattern recognition.

The winner-take-all map of PPC showed that the preference for words is located in the posterior

aspect of the PPC, adjacent to parieto-occipital and dorsal occipital areas. Interestingly, in the

blind group only, this map also revealed consonant preferring regions in an anatomically

intermediate position between shape preferring areas in anterior portions of PPC and word

preferring areas in parieto-occipital and dorsal occipital cortices. These regions did not emerge in

corrected whole-brain analyses and therefore should be interpreted with caution, requiring

investigation in future studies. However, the overall pattern suggests an anterior-to-posterior

parieto-occipital reading stream, analogous to the posterior-to-anterior vOTC gradient observed

in sighted readers. Within this gradient, parietal regions closer to S1, in anterior PPC represent

shape/texture information relevant to Braille, with posterior PPC and parieto-occipital regions

representing Braille orthography and still more posterior occipital areas representing linguistic
information.

As noted in the Introduction, involvement of the PPC in Braille reading is predicted by
connectivity-based theories of brain function (Bedny, 2017; Hannagan et al., 2015; Mahon &
Caramazza, 2011; Saygin et al., 2016). One hypothesis, therefore, is that the PPC, along with
adjacent parieto-occipital areas, plays an analogous role in Braille orthographic processing to the
role of the vOTC in orthographic processing of visual print: conversion of tactile patterns to
orthographic representations (Dehaene & Cohen, 2011; Dehaene et al., 2005).

Further work is needed to uncover the precise cognitive contribution of PPC and parieto-
occipital cortices to Braille reading. In sighted readers, the PPC also contributes to reading but
under different circumstances. The PPC is thought to participate in grapheme to phoneme
conversion, letter position decoding, as well as working memory processes and shows more
robust activity when effortful letter-by-letter reading is required (e.g., when words are degraded)
(Carreiras, Quiñones, Hernández-Cabrera, & Duñabeitia, 2015; Cohen et al., 2004; Costanzo,
Menghini, Caltagirone, Oliveri, & Vicari, 2012; Dehaene-Lambertz, Monzalvo, & Dehaene,
2018; Henry et al., 2005; Jonides et al., 1998; Koenigs, Barbey, Postle, & Grafman, 2009;
Ossmy, Ben-Shachar, & Mukamel, 2014; Taylor, Rastle, & Davis, 2013). Parietal cortex also
shows sensitivity to phonological rather than orthographic information during visual reading, in
contrast to the VWFA (Booth et al., 2003; Bouhali et al., 2019). In future studies, it would be
interesting to separate parietal responses to phonological as opposed to word-form information in
blind readers of Braille. In addition, further research is needed to explore the anatomical layout
of Braille-responsive parietal areas. For example, whether the parieto-occipital stream contains
punctate regions analogous to the VWFA, or more distributed responses to Braille letters and
words remains an open question. Likewise, in future studies, it will be important to test the
precise role of the PPC in Braille reading and to dissociate the functions of PPC, parieto-
occipital, and dorsal occipital regions.

Differential role of early visual cortex in Braille and visual print reading

We observed responses to reading in V1 in the blind but not sighted group. Like dorsal occipital
areas, V1 showed a preference for words over consonant strings and control shapes. The

involvement of V1 in Braille reading is consistent with previous studies (Cohen et al., 1997;

Kupers et al., 2007; Sadato et al., 1996). We further found that, in whole-cortex results, a portion

of V1 (foveal aspect of left V1) responded more to Braille reading than auditory word

comprehension, whereas other portions of V1 (right hemisphere, and peripheral) did not show

such a preference. This evidence is consistent with prior work suggesting that V1 does not have a

single, homogeneous function in people who are blind but rather contains multiple anatomically

separable functional subdivisions (Amedi et al., 2003; Bedny et al., 2011; Burton, Diamond, &

McDermott, 2003; Burton, Snyder, Diamond, et al., 2002; Kanjlia et al., 2021; Kanjlia, Pant, &

Bedny, 2019; Lane et al., 2015; Noppeney et al., 2003). Likewise, V1 may contain anatomically

separable Braille-specific and high-level language responses in blind readers of Braille.

Lateralization of Braille reading: effects of spoken language lateralization and reading

hand

With the exception of the primary somatosensory cortex, laterality of responses to written words

in the entire reading network (vOTC, PPC, V1, and IFC) is predicted by the laterality of spoken

word comprehension across blind individuals. On average congenitally blind individuals showed

reduced left-lateralization of responses to spoken and written words (see also Lane et al., 2017).

Those blind individuals who show right-lateralized responses to spoken words also show right-

lateralized responses to written words. Previous studies with sighted readers with right

hemisphere spoken language responses have likewise observed co-lateralization of spoken and

written language (Cai et al., 2010; Van der Haegen et al., 2012).   We did not observe this pattern

in the current sighted sample, possibly because all sighted participants in the current study had

strongly left-lateralized responses to spoken language and thus there was little interindividual

variability. Together, these data suggest that written and spoken language tend to co-lateralize in

blind and sighted readers alike. This observation is consistent with the hypothesis that strong

connectivity to spoken language networks is one of the determining factors of which regions

become 'recycled' for reading.

We also found a significant effect of reading hand on the lateralization of Braille reading that

was independent of the effect of spoken language lateralization. That is, right-hand Braille

readers showed more left-lateralized activation whereas left-hand Braille readers showed a

bilateral response to Braille. In contrast to the effect of spoken language on laterality, the effect

of reading hand was strongest in the primary somatosensory cortex, persisted in PPC and vOTC,

and was absent in IFC and V1. This observation is consistent with the idea that V1 occupies the

top of a processing hierarchy for people who are blind (Buechel, 2003). Effects of reading hand

thus persist past S1 but wane at higher stages of processing, whereas effects of language

lateralization are most prominent at higher processing stages and disappear in early sensory areas

(i.e., S1).

In sum, the lateralization of Braille reading is jointly determined by the lateralization of spoken

language and the input hand that receives the initial Braille stimulus. Although specific

lateralization patterns differ across sighted and blind groups, an analogous connectivity principle

appears to govern lateralization of reading in sighted and blind readers: lateralization depends

jointly on connectivity to sensory input regions (unilateral S1/ bilateral V1) and language

networks.

General conclusions

We find that the neural basis of Braille reading differs from that of visual print reading in several

ways. While visual print reading recruits a posterior/anterior vOTC gradient, no such gradient is

observed in the vOTC of blind readers of Braille. Blind readers of Braille recruit posterior

parietal cortices to a greater degree and in a different way compared to visual print reading in

sighted people. Only blind readers show preferential responses to written words in PPC and

parieto-occipital cortex. We observed suggestive evidence for an anterior-to-posterior stream of

processing in the parietal cortex of blind Braille readers, with anterior parietal areas involved in

tactile pattern perception and more posterior parietal, parieto-occipital and dorsal occipital

regions involved in word recognition. In blind and sighted readers alike, lateralization of spoken

language predicts lateralization of written language. However, on average, spoken word and

visual word recognition is highly left-lateralized in sighted people. By contrast, neither Braille

reading nor spoken word recognition is strongly left-lateralized in people who are born blind. In

blind readers of Braille, reading hand also affects lateralization of responses to Braille.

Comparing the neural basis of reading across blind and sighted people suggests that there is no

'standard reading brain.' The input modality of symbols influences the neural basis of their

recognition. At the same time, similar anatomical principles govern the localization of visual

print and tactile Braille. Connectivity patterns constrain the localization of visual print and tactile

Braille reading alike.

Credit Author Statement

J.K., S.K., and M.B. designed research; J.K., S.K., and M.B. performed research; M.T. analyzed data; M.T., E.S. and M.B. wrote the paper.

Declaration of Competing Interest

The authors declare no competing interests.

Acknowledgments

We would like to thank all our blind and sighted participants, the blind community and the National Federation of the Blind. Without their support, this study would not be possible. This work was supported by grants from the Johns Hopkins Science of Learning Institute (80034917) and the NIH/NEI (R01 EY027352-01). We would also like to thank the F. M. Kirby Research Center for Functional Brain Imaging at the Kennedy Krieger Institute for their assistance in data collection.

References

Amedi, A., Raz, N., Pianka, P., Malach, R., & Zohary, E. (2003). Early "visual" cortex activation
correlates with superior verbal memory performance in the blind. *Nature Neuroscience*,
6(7), 758–766. https://doi.org/10.1038/nn1072

Baker, C. I., Liu, J., Wald, L. L., Kwong, K. K., Benner, T., & Kanwisher, N. (2007). Visual
word processing and experiential origins of functional selectivity in human extrastriate
cortex. *Proceedings of the National Academy of Sciences*, *104*(21), 9087–9092.
https://doi.org/10.1073/pnas.0703300104

Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., … Dehaene-
Lambertz, G. (2018). A lateral-to-mesial organization of human ventral visual cortex at
birth. *Brain Structure and Function*, *223*(7), 3107–3119. https://doi.org/10.1007/s00429-
018-1676-3

Bauer, C., Yazzolino, L., Hirsch, G., Cattaneo, Z., Vecchi, T., & Merabet, L. B. (2015). Neural
correlates associated with superior tactile symmetry perception in the early blind. *Cortex*,
63, 104–117. https://doi.org/10.1016/j.cortex.2014.08.003

Bedny, M. (2017). Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends in*
Cognitive Sciences, *21*(9), 637–648. https://doi.org/10.1016/j.tics.2017.06.003

Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E., & Saxe, R. (2011). Language
processing in the occipital cortex of congenitally blind adults. *Proceedings of the National*
Academy of Sciences of the United States of America, *108*(11), 4429–4434.
https://doi.org/10.1073/pnas.1014818108

Bedny, M., Richardson, H., & Saxe, R. (2015). "Visual" cortex responds to spoken language in
blind children. *Journal of Neuroscience*, *35*(33), 11674–11681.
https://doi.org/10.1523/JNEUROSCI.0634-15.2015

Behrmann, M., & Plaut, D. C. (2013). Distributed circuits, not circumscribed centers, mediate
visual recognition. *Trends in Cognitive Sciences*, *17*(5), 210–219.

https://doi.org/10.1016/j.tics.2013.03.007

Behrmann, M., & Plaut, D. C. (2020). Hemispheric Organization for Visual Object Recognition:

A Theoretical Account and Empirical Evidence*. *Perception*, *49*(4), 373–404.

https://doi.org/10.1177/0301006619899049

Binder, J. R., Medler, D. A., Westbury, C. F., Liebenthal, E., & Buchanan, L. (2006). Tuning of

the human left fusiform gyrus to sublexical orthographic structure. *NeuroImage*, *33*(2),

739–748. https://doi.org/10.1016/j.neuroimage.2006.06.053

Bola, Ł., Matuszewski, J., Szczepanik, M., Droździel, D., Sliwinska, M. W., Paplińska, M., …

Marchewka, A. (2019). Functional hierarchy for tactile processing in the visual cortex of

sighted adults. *NeuroImage*, *202*, 116084.

https://doi.org/10.1016/j.neuroimage.2019.116084

Bolger, D. J., Perfetti, C. A., & Schneider, W. (2005). Cross-cultural effect on the brain revisited:

Universal structures plus writing system variation. *Human Brain Mapping*, *25*(1), 92–104.

https://doi.org/10.1002/hbm.20124

Booth, J. R., Burman, D. D., Meyer, J. R., Gitelman, D. R., Parrish, T. B., & Mesulam, M. M.

(2003). Relation between brain activation and lexical performance. *Human Brain Mapping*,

19(3), 155–169. https://doi.org/10.1002/hbm.10111

Bouhali, F., Bézagu, Z., Dehaene, S., & Cohen, L. (2019). A mesial-to-lateral dissociation for

orthographic processing in the visual cortex. *Proceedings of the National Academy of*

Sciences of the United States of America, *116*(43), 21936–21946.

https://doi.org/10.1073/pnas.1904184116

Bouhali, F., de Schotten, M. T., Pinel, P., Poupon, C., Mangin, J. F., Dehaene, S., & Cohen, L.

(2014). Anatomical connections of the visual word form area. *Journal of Neuroscience*,

34(46), 15402–15414. https://doi.org/10.1523/JNEUROSCI.4918-13.2014

Boven, R. W. V., Hamilton, R. H., Kauffman, T., Keenan, J. P., & Pascual-Leone, A. (2000).

Tactile spatial resolution in blind Braille readers. *Neurology*, *54*(12), 2230–2236.

https://doi.org/10.1212/WNL.54.12.2230

Bruno, J. L., Zumberge, A., Manis, F. R., Lu, Z. L., & Goldman, J. G. (2008). Sensitivity to
        orthographic familiarity in the occipito-temporal region. *NeuroImage*, *39*(4), 1988–2001.
        https://doi.org/10.1016/j.neuroimage.2007.10.044

Büchel, C., Price, C., & Friston, K. (1998a). A multimodal language region in the ventral visual
        pathway. *Nature*, *394*(6690), 274–277. https://doi.org/10.1038/28389

Büchel, C., Price, C., & Friston, K. (1998b). A multimodal language region in the ventral visual
        pathway. *Nature*, *394*(6690), 274–277. https://doi.org/10.1038/28389

Buechel, C. (2003). Cortical hierarchy turned on its head. *Nature Neuroscience*, *6*(7), 657–658.

Burks, J. D., Boettcher, L. B., Conner, A. K., Glenn, C. A., Bonney, P. A., Baker, C. M., …
        Sughrue, M. E. (2017). White matter connections of the inferior parietal lobule: A study of
        surgical anatomy. *Brain and Behavior*, *7*(4), 1–12. https://doi.org/10.1002/brb3.640

Burton, H., Diamond, J. B., & McDermott, K. B. (2003). Dissociating cortical regions activated
        by semantic and phonological tasks: A fMRI study in blind and sighted people. *Journal of
        Neurophysiology*, *90*(3), 1965–1982. https://doi.org/10.1152/jn.00279.2003

Burton, H., Snyder, A. Z., Conturo, T. E., Akbudak, E., Ollinger, J. M., & Raichle, M. E. (2002).
        Adaptive Changes in Early and Late Blind: A fMRI Study of Braille Reading. *Journal of
        Neurophysiology*, *87*(1), 589–607. https://doi.org/10.1152/jn.00285.2001

Burton, H., Snyder, A. Z., Diamond, J. B., & Raichle, M. E. (2002). Adaptive Changes in Early
        and Late Blind: A fMRI Study of Verb Generation to Heard Nouns. *Journal of
        Neurophysiology*, *88*(6), 3359–3371. https://doi.org/10.1152/jn.00129.2002

Burton, H, Snyder, A. Z., Diamond, J. B., & Raichle, M. E. (2002). Adaptive Changes in Early
        and Late Blind: A fMRI Study of Verb Generation to Heard Nouns. *Journal of
        Neurophysiology*, *88*(6), 3359–3371. https://doi.org/10.1152/jn.00129.2002

Burton, Harold, Sinclair, R. J., & Agato, A. (2012). Recognition memory for Braille or spoken
        words: An fMRI study in early blind. *Brain Research*, *1438*, 22–34.
        https://doi.org/10.1016/j.brainres.2011.12.032

Burton, Harold, Sinclair, R. J., & McLaren, D. G. (2004). Cortical activity to vibrotactile
stimulation: An fMRI study in blind and sighted individuals. *Human Brain Mapping*, *23*(4),
210–228. https://doi.org/10.1002/hbm.20064

Cai, Q., Lavidor, M., Brysbaert, M., Paulignan, Y., & Nazir, T. A. (2008). Cerebral lateralization
of frontal lobe language processes and lateralization of the posterior visual word processing
system. *Journal of Cognitive Neuroscience*, *20*(4), 672–681.
https://doi.org/10.1162/jocn.2008.20043

Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D., & Nazir, T. A. (2010). The left ventral
occipito-temporal response to words depends on language lateralization but not on visual
familiarity. *Cerebral Cortex*, *20*(5), 1153–1163. https://doi.org/10.1093/cercor/bhp175

Cai, Q., & Van der Haegen, L. (2015). What can atypical language hemispheric specialization
tell us about cognitive functions? *Neuroscience Bulletin*, *31*(2), 220–226.
https://doi.org/10.1007/s12264-014-1505-5

Carreiras, M., Quiñones, I., Hernández-Cabrera, J. A., & Duñabeitia, J. A. (2015). Orthographic
coding: Brain activation for letters, symbols, and digits. *Cerebral Cortex*, *25*(12), 4748–
4760. https://doi.org/10.1093/cercor/bhu163

Cohen, Celnik, P., Pascual-Leone, A., Corwell, B., Faiz, L., Dambrosia, J., … Hallett, M. (1997).
Functional relevance of cross-modal plasticity in blind humans. *Nature*, *389*(6647), 180–
183. https://doi.org/10.1038/38278

Cohen, Dehaene, S., Naccache, L., Lehéricy, S., Dehaene-Lambertz, G., Hénaff, M. A., &
Michel, F. (2000). The visual word form area: spatial and temporal characterization of an
initial stage of reading in normal subjects and posterior split-brain patients. *Brain : A*
Journal of Neurology, (2), 291–307. https://doi.org/10.1093/brain/123.2.291

Cohen, L., Dehaene, S., Vinckier, F., Jobert, A., & Montavont, A. (2008). Reading normal and
degraded words: Contribution of the dorsal and ventral visual pathways. *NeuroImage*,
40(1), 353–366. https://doi.org/10.1016/j.neuroimage.2007.11.036

Cohen, L., Henry, C., Dehaene, S., Martinaud, O., Lehéricy, S., Lemer, C., & Ferrieux, S.

(2004). The pathophysiology of letter-by-letter reading. *Neuropsychologia*, *42*(13), 1768–
1780. https://doi.org/10.1016/j.neuropsychologia.2004.04.018

Cohen, Weeks, R. A., Sadato, N., Celnik, P., Ishii, K., & Hallett, M. (1999). Period of
susceptibility for cross-modal plasticity in the blind. *Annals of Neurology*, *45*(4), 451–460.
https://doi.org/10.1002/1531-8249(199904)45:4<451::AID-ANA6>3.0.CO;2-B

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in
the brain. *Nature Reviews Neuroscience*, *3*(3), 201–215. https://doi.org/10.1038/nrn755

Costanzo, F., Menghini, D., Caltagirone, C., Oliveri, M., & Vicari, S. (2012). High frequency
rTMS over the left parietal lobule increases non-word reading accuracy. *Neuropsychologia*,
50(11), 2645–2651. https://doi.org/10.1016/j.neuropsychologia.2012.07.017

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I. Segmentation
and surface reconstruction. *NeuroImage*, *9*(2), 179–194.
https://doi.org/10.1006/nimg.1998.0395

Debowska, W., Wolak, T., Nowicka, A., Kozak, A., Szwed, M., & Kossut, M. (2016).
Functional and structural neuroplasticity induced by short-term tactile training based on
braille reading. *Frontiers in Neuroscience*, *10*(OCT), 1–13.
https://doi.org/10.3389/fnins.2016.00460

Dehaene-Lambertz, G., Monzalvo, K., & Dehaene, S. (2018). *The emergence of the visual word*
form: Longitudinal evolution of category-specific ventral visual areas during reading
acquisition. *PLoS Biology* (Vol. 16). https://doi.org/10.1371/journal.pbio.2004103

Dehaene, S., & Cohen, L. (2011). The unique role of the visual word form area in reading.
Trends in Cognitive Sciences, *15*(6), 254–262. https://doi.org/10.1016/j.tics.2011.04.003

Dehaene, S., Cohen, L., Sigman, M., & Vinckier, F. (2005). The neural code for written words:
A proposal. *Trends in Cognitive Sciences*, *9*(7), 335–341.
https://doi.org/10.1016/j.tics.2005.05.004

Dehaene, S., Jobert, A., Naccache, L., Ciuciu, P., Poline, J. B., Bihan, D. Le, & Cohen, L.

(2004). Letter binding and invariant recognition of masked words: Behavioral and neuroimaging evidence. *Psychological Science*, *15*(5), 307–313. https://doi.org/10.1111/j.0956-7976.2004.00674.x

Dehaene, S., Pegado, F., Braga, L. W., Ventura, P., Nunes Filho, G., Jobert, A., … Velay, J. L. (2010). How learning to read changes the cortical networks for vision and language. *Science*, *330*(6009), 1359–1364. https://doi.org/10.1126/science.1194140

DiCarlo, J. J., & Cox, D. D. (2007). Untangling invariant object recognition. *Trends in Cognitive Sciences*, *11*(8), 333–341. https://doi.org/10.1016/j.tics.2007.06.010

Duhamel, J. R., Colby, C. L., & Goldberg, M. E. (1998). Ventral intraparietal area of the macaque: Congruent visual and somatic response properties. *Journal of Neurophysiology*, *79*(1), 126–136. https://doi.org/10.1152/jn.1998.79.1.126

Dzięgiel-Fivet, G., Plewko, J., Szczerbiński, M., Marchewka, A., Szwed, M., & Jednoróg, K. (2021). Neural network for Braille reading and the speech-reading convergence in the blind: Similarities and differences to visual reading. *NeuroImage*, *231*(February). https://doi.org/10.1016/j.neuroimage.2021.117851

Fedorenko, E., Hsieh, P. J., Nieto-Castañón, A., Whitfield-Gabrieli, S., & Kanwisher, N. (2010). New method for fMRI investigations of language: Defining ROIs functionally in individual subjects. *Journal of Neurophysiology*, *104*(2), 1177–1194. https://doi.org/10.1152/jn.00032.2010

Feng, X., Altarelli, I., Monzalvo, K., Ding, G., Ramus, F., Shu, H., … Dehaene-Lambertz, G. (2020). A universal reading network and its modulation by writing system and reading ability in French and Chinese children. *ELife*, *9*, 1–51. https://doi.org/10.7554/eLife.54591

Gizewski, E. R., Gasser, T., De Greiff, A., Boehm, A., & Forsting, M. (2003). Cross-modal plasticity for sensory and motor activation patterns in blind subjects. *NeuroImage*, *19*(3), 968–975. https://doi.org/10.1016/S1053-8119(03)00114-9

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., … Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human Connectome

Project. *NeuroImage*, *80*, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Glezer, L. S., Jiang, X., & Riesenhuber, M. (2009). Evidence for Highly Selective Neuronal

Tuning to Whole Words in the "Visual Word Form Area." *Neuron*, *62*(2), 199–204.

https://doi.org/10.1016/j.neuron.2009.03.017

Gougoux, F., Zatorre, R. J., Lassonde, M., Voss, P., & Lepore, F. (2005). A functional

neuroimaging study of sound localization: Visual cortex activity predicts performance in

early-blind individuals. *PLoS Biology*, *3*(2), 0324–0333.

https://doi.org/10.1371/journal.pbio.0030027

Grant, A. C., Thiagarajah, M. C., & Sathian, K. (2000). Tactile perception in blind braille

readers: A psychophysical study of acuity and hyperacuity using gratings and dot patterns.

Perception and Psychophysics, *62*(2), 301–312. https://doi.org/10.3758/BF03205550

Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G., & Dehaene, S. (2015). Origins of

the specialization for letters and numbers in ventral occipitotemporal cortex. *Trends in*

Cognitive Sciences, *19*(7), 374–382. https://doi.org/10.1016/j.tics.2015.05.006

Hannagan, T., & Grainger, J. (2013). The Lazy Visual Word Form Area: Computational Insights

into Location-Sensitivity. *PLoS Computational Biology*, *9*(10).

https://doi.org/10.1371/journal.pcbi.1003250

Hegner, Y. L., Lee, Y., Grodd, W., & Braun, C. (2010). Comparing tactile pattern and

vibrotactile frequency discrimination: A human fMRI study. *Journal of Neurophysiology*,

103(6), 3115–3122. https://doi.org/10.1152/jn.00940.2009

Henry, C., Gaillard, R., Volle, E., Chiras, J., Ferrieux, S., Dehaene, S., & Cohen, L. (2005).

Brain activations during letter-by-letter reading: A follow-up study. *Neuropsychologia*,

43(14), 1983–1989. https://doi.org/10.1016/j.neuropsychologia.2005.04.007

Hu, W., Lee, H. L., Zhang, Q., Liu, T., Geng, L. B., Seghier, M. L., … Price, C. J. (2010).

Developmental dyslexia in Chinese and English populations: Dissociating the effect of

dyslexia from language differences. *Brain*, *133*(6), 1694–1706.

https://doi.org/10.1093/brain/awq106

Jonides, J., Schumacher, E. H., Smith, E. E., Koeppe, R. A., Awh, E., Reuter-Lorenz, P. A., …
Willis, C. R. (1998). The role of parietal cortex in verbal working memory. *Journal of Neuroscience*, *18*(13), 5026–5034. https://doi.org/10.1523/jneurosci.18-13-05026.1998

Kaas, J. H. (2012). Somatosensory System. In *The Human Nervous System* (Third Edit, pp. 1074–1109). Elsevier. https://doi.org/10.1016/B978-0-12-374236-0.10030-6

Kanjlia, S., Loiotile, R. E., Harhen, N., & Bedny, M. (2021). 'Visual' cortices of congenitally blind adults are sensitive to response selection demands in a go/no-go task. *NeuroImage*, *236*(April), 118023. https://doi.org/10.1016/j.neuroimage.2021.118023

Kanjlia, S., Pant, R., & Bedny, M. (2019). Sensitive Period for Cognitive Repurposing of Human Visual Cortex. *Cerebral Cortex*, *29*(9), 3993–4005. https://doi.org/10.1093/cercor/bhy280

Kim, J. S., Kanjlia, S., Merabet, L. B., & Bedny, M. (2017). Development of the visual word form area requires visual experience: Evidence from blind braille readers. *Journal of Neuroscience*, *37*(47), 11495–11504. https://doi.org/10.1523/JNEUROSCI.0997-17.2017

Koenigs, M., Barbey, A. K., Postle, B. R., & Grafman, J. (2009). Superior parietal cortex is critical for the manipulation of information in working memory. *Journal of Neuroscience*, *29*(47), 14980–14986. https://doi.org/10.1523/JNEUROSCI.3706-09.2009

Krafnick, A. J., Tan, L. H., Flowers, D. L., Luetje, M. M., Napoliello, E. M., Siok, W. T., …
Eden, G. F. (2016). Chinese Character and English Word processing in children's ventral occipitotemporal cortex: FMRI evidence for script invariance. *NeuroImage*, *133*, 302–312. https://doi.org/10.1016/j.neuroimage.2016.03.021

Kujala, T., Alho, K., Paavilainen, P., Summala, H., & Naatanen, R. (1992). Neural plasticity in processing of sound location by the early blind: an event-related potential study * Teija Kujala, Kimmo Alho, Petri Paavilainen, Heikki Summala and Risto N?i?it~inen. *Methods*, *84*(5), 469–472.

Kupers, R., Pappens, M., De Noordhout, A. M., Schoenen, J., Ptito, M., & Fumal, A. (2007). rTMS of the occipital cortex abolishes Braille reading and repetition priming in blind subjects. *Neurology*, *68*(9), 691–693. https://doi.org/10.1212/01.wnl.0000255958.60530.11

Lane, C., Kanjlia, S., Omaki, A., & Bedny, M. (2015). "Visual" cortex of congenitally blind
adults responds to syntactic movement. *Journal of Neuroscience*, *35*(37), 12859–12868.
https://doi.org/10.1523/JNEUROSCI.1256-15.2015

Lane, C., Kanjlia, S., Richardson, H., Fulton, A., Omaki, A., & Bedny, M. (2017). Reduced Left
Lateralization of Language in Congenitally Blind Individuals. *Journal of Cognitive*
Neuroscience, *29*(1), 65–78. https://doi.org/10.1162/jocn_a_01045

Lerma-Usabiaga, G., Carreiras, M., & Paz-Alonso, P. M. (2018). Converging evidence for
functional and structural segregation within the left ventral occipitotemporal cortex in
reading. *Proceedings of the National Academy of Sciences of the United States of America*,
115(42), E9981–E9990. https://doi.org/10.1073/pnas.1803003115

Lewis, J. W., & Van Essen, D. C. (2000). Corticocortical connections of visual, sensorimotor,
and multimodal processing areas in the parietal lobe of the macaque monkey. *Journal of*
Comparative Neurology, *428*(1), 112–137. https://doi.org/10.1002/1096-
9861(20001204)428:1<112::AID-CNE8>3.0.CO;2-9

Li, J., Osher, D. E., Hansen, H. A., & Saygin, Z. M. (2020). Innate connectivity patterns drive
the development of the visual word form area. *Scientific Reports*, *10*(1), 1–12.
https://doi.org/10.1038/s41598-020-75015-7

Ludersdorfer, P., Schurz, M., Richlan, F., Kronbichler, M., & Wimmer, H. (2013). Opposite
effects of visual and auditory word-likeness on activity in the visual word form area.
Frontiers in Human Neuroscience, *7*(August), 491.
https://doi.org/10.3389/fnhum.2013.00491

Mahon, B. Z., & Caramazza, A. (2011). What drives the organization of object knowledge in the
brain? *Trends in Cognitive Sciences*, *15*(3), 97–103.
https://doi.org/10.1016/j.tics.2011.01.004

Martin, A., Schurz, M., Kronbichler, M., & Richlan, F. (2015). Reading in the brain of children
and adults: A meta-analysis of 40 functional magnetic resonance imaging studies. *Human*
Brain Mapping, *36*(5), 1963–1981. https://doi.org/10.1002/hbm.22749

Medler, D. A., & Binder, J. R. (2005). An on-line orthographic database of the English language.

Http,//Www. Neuro, Mew. Edu/Meword.

Melzer, P., Morgan, V. L., Pickens, D. R., Price, R. R., Wall, R. S., & Ebner, F. F. (2001).

Cortical activation during Braille reading is influenced by early visual experience in

subjects with severe visual disability: A correlational fMRI study. *Human Brain Mapping*,

14(3), 186–195. https://doi.org/10.1002/hbm.1051

Millar, S. (1984). Is There a "Best Hand" for Braille? *Cortex*, *20*(1), 75–87.

https://doi.org/10.1016/S0010-9452(84)80025-8

Millar, S. (2003). *Reading by Touch*. *Reading by Touch*. Routledge.

https://doi.org/10.4324/9780203359440

Nakamura, K., Kuo, W. J., Pegado, F., Cohen, L., Tzeng, O. J. L., & Dehaene, S. (2012).

Universal brain systems for recognizing word shapes and handwriting gestures during

reading. *Proceedings of the National Academy of Sciences of the United States of America*,

109(50), 20762–20767. https://doi.org/10.1073/pnas.1217749109

Noppeney, U., Friston, K. J., & Price, C. J. (2003). Effects of visual deprivation on the

organization of the semantic system. *Brain*, *126*(7), 1620–1627.

https://doi.org/10.1093/brain/awg152

Ossmy, O., Ben-Shachar, M., & Mukamel, R. (2014). Decoding letter position in word reading.

Cortex, *59*, 74–83. https://doi.org/10.1016/j.cortex.2014.07.002

Ossowski, A., & Behrmann, M. (2015). Left hemisphere specialization for word reading

potentially causes, rather than results from, a left lateralized bias for high spatial frequency

visual information. *Cortex*, *72*, 27–39. https://doi.org/10.1016/j.cortex.2014.12.013

Pascual-Leone, A., Cammarota, A., Wassermann, E. M., Brasil-Neto, J. P., Cohen, L. G., &

Hallett, M. (1993). Modulation of motor cortical outputs to the reading hand of braille

readers. *Annals of Neurology*, *34*(1), 33–37. https://doi.org/10.1002/ana.410340108

Pascual-Leone, A., & Torres, F. (1993). Plasticity of the sensorimotor cortex representation of

the reading finger in braille readers. *Brain*, *116*(1), 39–52.

https://doi.org/10.1093/brain/116.1.39

Price, C. J., & Devlin, J. T. (2011). The Interactive Account of ventral occipitotemporal

contributions to reading. *Trends in Cognitive Sciences*, *15*(6), 246–253.

https://doi.org/10.1016/j.tics.2011.04.001

Purcell, J. J., Shea, J., & Rapp, B. (2014). Beyond the visual word form area: The orthography-

semantics interface in spelling and reading. *Cognitive Neuropsychology*, *31*(5–6), 482–510.

https://doi.org/10.1080/02643294.2014.909399

Rączy, K., Urbańczyk, A., Korczyk, M., Szewczyk, J. M., Sumera, E., & Szwed, M. (2019).

Orthographic Priming in Braille Reading as Evidence for Task-specific Reorganization in

the Ventral Visual Cortex of the Congenitally Blind. *Journal of Cognitive Neuroscience*,

31(7), 1065–1078. https://doi.org/10.1162/jocn_a_01407

Reich, L., Szwed, M., Cohen, L., & Amedi, A. (2011). A ventral visual stream reading center

independent of visual experience. *Current Biology*, *21*(5), 363–368.

https://doi.org/10.1016/j.cub.2011.01.040

Reilhac, C., Peyrin, C., Démonet, J. F., & Valdois, S. (2013). Role of the superior parietal

lobules in letter-identity processing within strings: FMRI evidence from skilled and

dyslexicreaders. *Neuropsychologia*, *51*(4), 601–612.

https://doi.org/10.1016/j.neuropsychologia.2012.12.010

Röder, B., Rösler, F., & Neville, H. J. (2000). Event-related potentials during auditory language

processing in congenitally blind and sighted people. *Neuropsychologia*, *38*(11), 1482–1502.

https://doi.org/10.1016/S0028-3932(00)00057-9

Röder, B., Stock, O., Bien, S., Neville, H., & Rösler, F. (2002). Speech processing activates

visual cortex in congenitally blind humans. *European Journal of Neuroscience*, *16*(5), 930–

936. https://doi.org/10.1046/j.1460-9568.2002.02147.x

Rueckl, J. G., Paz-Alonso, P. M., Molfese, P. J., Kuo, W. J., Bick, A., Frost, S. J., … Frost, R.

(2015). Universal brain signature of proficient reading: Evidence from four contrasting

languages. *Proceedings of the National Academy of Sciences of the United States of*
America, *112*(50), 15510–15515. https://doi.org/10.1073/pnas.1509321112

Ruschel, M., Knösche, T. R., Friederici, A. D., Turner, R., Geyer, S., & Anwander, A. (2014).
Connectivity architecture and subdivision of the human inferior parietal cortex revealed by
diffusion MRI. *Cerebral Cortex*, *24*(9), 2436–2448. https://doi.org/10.1093/cercor/bht098

Saalmann, Y. B., Pigarev, I. N., & Vidyasagar, T. R. (2007). Neural Mechanisms of Visual
Attention: How Top-Down Feedback Highlights Relevant Locations. *Science*, *316*(5831),
1612–1615. https://doi.org/10.1126/science.1139140

Sadato, N., Pascual-Leone, A., Grafman, J., Deiber, M. P., Ibañez, V., & Hallett, M. (1998).
Neural networks for Braille reading by the blind. *Brain*, *121*(7), 1213–1229.
https://doi.org/10.1093/brain/121.7.1213

Sadato, N., Pascual-Leone, A., Grafman, J., Ibañez, V., Deiber, M., Dold, G., & Hallett, M.
(1996). Activation of the primary visual cortex by Braille reading in blind subjects. *Nature*,
380(6574), 526–528. https://doi.org/10.1038/380526a0

Sathian, K. (2005). Visual cortical activity during tactile perception in the sighted and the
visually deprived. *Developmental Psychobiology*, *46*(3), 279–286.
https://doi.org/10.1002/dev.20056

Sathian, K., & Stilla, R. (2010). Cross-modal plasticity of tactile perception in blindness.
Restorative Neurology and Neuroscience, *28*(2), 271–281. https://doi.org/10.3233/RNN-
2010-0534

Saygin, Z. M., Osher, D. E., Norton, E. S., Youssoufian, D. A., Beach, S. D., Feather, J., …
Kanwisher, N. (2016). Connectivity precedes function in the development of the visual
word form area. *Nature Neuroscience*, *19*(9), 1250–1255. https://doi.org/10.1038/nn.4354

Schlaggar, B. L., & McCandliss, B. D. (2007). Development of neural systems for reading.
Annual Review of Neuroscience, *30*, 475–503.
https://doi.org/10.1146/annurev.neuro.28.061604.135645

Seghier, M. L., & Price, C. J. (2011). Explaining left lateralization for words in the ventral

occipitotemporal cortex. *The Journal of Neuroscience : The Official Journal of the Society*

for Neuroscience, *31*(41), 14745–14753. https://doi.org/10.1523/JNEUROSCI.2238-

11.2011

Siuda-Krzywicka, K., Bola, Ł., Paplińska, M., Sumera, E., Jednoróg, K., Marchewka, A., …

Szwed, M. (2016). Massive cortical reorganization in sighted braille readers. *ELife*,

5(MARCH2016), 1–26. https://doi.org/10.7554/eLife.10762

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-

Berg, H., … Matthews, P. M. (2004). Advances in functional and structural MR image

analysis and implementation as FSL. *NeuroImage*, *23*(SUPPL. 1), 208–219.

https://doi.org/10.1016/j.neuroimage.2004.07.051

Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017a). Privileged

functional connectivity between the visual word form area and the language system.

Journal of Neuroscience, *37*(21), 5288–5297. https://doi.org/10.1523/JNEUROSCI.0138-

17.2017

Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017b). Privileged

functional connectivity between the visual word form area and the language system.

Journal of Neuroscience, *37*(21), 5288–5297. https://doi.org/10.1523/JNEUROSCI.0138-

17.2017

Stilla, R., Hanna, R., Hu, X., Mariola, E., Deshpande, G., & Sathian, K. (2008). Neural

processing underlying tactile microspatial discrimination in the blind: A functional

magnetic resonance imaging study. *Journal of Vision*, *8*(10), 1–19.

https://doi.org/10.1167/8.10.13

Szwed, M., Dehaene, S., Kleinschmidt, A., Eger, E., Valabrègue, R., Amadon, A., & Cohen, L.

(2011). Specialization for written words over objects in the visual cortex. *NeuroImage*,

56(1), 330–344. https://doi.org/10.1016/j.neuroimage.2011.01.073

Szwed, M., Qiao, E., Jobert, A., Dehaene, S., & Cohen, L. (2014). Effects of Literacy in Early

1345  Visual and Occipitotemporal Areas of Chinese and French Readers. *Journal of Cognitive*

1346  *Neuroscience*, *26*(3), 459–475. https://doi.org/10.1162/jocn_a_00499

1347 Taylor, J. S. H., Rastle, K., & Davis, M. H. (2013). Can cognitive models explain brain

1348  activation during word and pseudoword reading? A meta-analysis of 36 neuroimaging

1349  studies. *Psychological Bulletin*, *139*(4), 766–791. https://doi.org/10.1037/a0030266

1350 Van der Haegen, L., Cai, Q., & Brysbaert, M. (2012). Colateralization of Broca's area and the

1351  visual word form area in left-handers: FMRI evidence. *Brain and Language*, *122*(3), 171–

1352  178. https://doi.org/10.1016/j.bandl.2011.11.004

1353 Van Essen, D. C. (2005). A Population-Average, Landmark- and Surface-based (PALS) atlas of

1354  human cerebral cortex. *NeuroImage*, *28*(3), 635–662.

1355  https://doi.org/10.1016/j.neuroimage.2005.06.058

1356 Vinckier, F., Dehaene, S., Jobert, A., Dubus, J. P., Sigman, M., & Cohen, L. (2007). Hierarchical

1357  Coding of Letter Strings in the Ventral Stream: Dissecting the Inner Organization of the

1358  Visual Word-Form System. *Neuron*, *55*(1), 143–156.

1359  https://doi.org/10.1016/j.neuron.2007.05.031

1360 Wang, X., Yang, J., Shu, H., & Zevin, J. D. (2011). Left fusiform BOLD responses are inversely

1361  related to word-likeness in a one-back task. *NeuroImage*, *55*(3), 1346–1356.

1362  https://doi.org/10.1016/j.neuroimage.2010.12.062

1363 Watkins, K. E., Cowey, A., Alexander, I., Filippini, N., Kennedy, J. M., Smith, S. M., … Bridge,

1364  H. (2012). Language networks in anophthalmia: Maintained hierarchy of processing in

1365  "visual" cortex. *Brain*, *135*(5), 1566–1577. https://doi.org/10.1093/brain/aws067

1366 Wong, M., Gnanakumaran, V., & Goldreich, D. (2011). Tactile Spatial Acuity Enhancement in

1367  Blindness: Evidence for Experience-Dependent Mechanisms. *Journal of Neuroscience*,

1368  *31*(19), 7028–7037. https://doi.org/10.1523/JNEUROSCI.6461-10.2011

1369 Yeatman, J. D., Rauschecker, A. M., & Wandell, B. A. (2013). Anatomy of the visual word form

1370  area: Adjacent cortical circuits and long-range white matter connections. *Brain and*

1371  *Language*, *125*(2), 146–155. https://doi.org/10.1016/j.bandl.2012.04.010

1372

1373