From Learning to Remembering: How Memory Consolidation Differs in Term and Preterm Born Children from Young Adults

Iryna Schommartz[1], Philip F. Lembcke[2], Henriette Schuetz[2], Nina Wald de Chamorro[3,4], Martin Bauer[2], Angela M. Kaindl[3,4,5], Claudia Buss[2,7]*, and Yee Lee Shing[1,8]*

* Yee Lee Shing and Claudia Buss should be considered joint senior author.

[1]Department of Psychology, Goethe University Frankfurt, Frankfurt, Germany

[2] Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Medical Psychology, Berlin, Germany

[3]Charité – Universitätsmedizin Berlin, Department of Pediatric Neurology, Berlin, Germany

[4]Charité – Universitätsmedizin Berlin, Center for Chronically Sick Children, Berlin, Germany

[5]Charité – Universitätsmedizin Berlin, Institute of Cell- and Neurobiology, Berlin, Germany

[6]Charité – Universitätsmedizin Berlin, Department of Pediatric Surgery, Berlin, Germany

[7]Development, Health and Disease Research Program, Department of Pediatrics, University of California Irvine, USA

[8]Center for Individual Development and Adaptive Education of Children at Risk (IDeA), Frankfurt, Germany

*Correspondence: Prof. Dr. Yee Lee Shing, Goethe University Frankfurt, Theodor-W.-Adorno-Platz, 6, 60323, Germany. Email: shing@psych.uni-frankfurt.de

Acknowledgements: We thank all the children and parents who participated in the study. This project was funded by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation, Project-ID 327654276, SFB 1315, "Mechanisms and Disturbances in Memory Consolidation: from Synapses to Systems"). The work of YLS was also supported by the European Union (ERC-2018-StG-PIVOTAL-758898).

Conflict of interest statement: We have no known conflict of interest to disclose.

Data availability statement: The datasets generated and analysed during the current study are available from the corresponding authors upon reasonable request.

Ethic approval statement: The study was approved by the ethics committee of the Goethe University Frankfurt am Main, Germany (Nr. E 145/18).

RESEARCH HIGHLIGHTS

- Short-delay memory consolidation of object-location associations does not differ between 6-year-old preterm born and term born children, as well as between children and young adults.

- Long-delay memory consolidation in 6-year-old children is less efficient than in young adults.

- Larger HC volume is associated with higher long-delay retention rates in 6-year-old children, while thinner medial OFC predicted higher memory retention rates in all groups.

- Thinner medial OFC is associated with higher final learning performance in 6-year-old children and adults.

Abstract

Human capacity to remember experienced episodes over a long period of time has its roots in childhood and develops throughout the lifespan. However, the neural regions supporting memory consolidation in the developing brain remain to be ascertained. The present study examined system-level memory consolidation of object-location associations after one night of sleep (short delay) and after two weeks (long delay), and its relation to structural brain measures in normally developing term born and preterm born 6-year-old children, as well as in young adults as a reference group of mature consolidation systems. We showed that final learning performance was reduced in preterm in comparison to term born children, who in turn were outperformed by young adults. There were no differences in short- and long-delay memory consolidation between term and preterm born children. Despite comparable short-delay memory consolidation in all groups, both term and preterm born children showed less efficient long-delay memory consolidation in comparison to young adults. Moreover, long-delay memory consolidation was positively associated with larger hippocampal volume in children, while a thinner medial orbitofrontal cortex was associated with better overall memory retention rates in all age groups. Thinner medial orbitofrontal cortex was furthermore associated with higher final learning performance in children. Taken together, the results suggest that temporal dynamics of memory consolidation and its association with structural brain measures in 6-year-old term born and preterm born children are comparable but differ from young adults.

Keywords: episodic memory, object-scene associations, memory consolidation, preterm birth, hippocampus, medial orbital prefrontal cortex

1   INTRODUCTION

Humans have an impressive capacity to store and retrieve memories of past experiences with unique temporal-spatial features for years and even decades (Squire et al., 2015; Tulving & Markowitsch, 1998; Tulving, 2002). This is possible due to memory consolidation, a dynamic and complex process through which acquired memory traces become long-lasting (Dudai, 2004; Dudai et al., 2015; Moscovitch & Gilboa, 2021). The interplay between the hippocampus (HC) and neocortical brain structures underlies the formation, maintenance, and retrieval of declarative memory (Moscovitch & Gilboa, 2021; Simons & Spiers, 2003; Squire et al., 2015b; Squire & Alvarez, 1995). Events are bound by the HC into a coherent memory trace containing contextually rich information that is distributed within neocortical areas (Eichenbaum, 2017; Nadel & Peterson, 2013). The ventromedial prefrontal cortex the (vmPFC) is involved in the evolvement of memory representations either by integrating new memory representations into pre-existing schemata or by forming new schemata (Frankland & Bontempi, 2005; Squire et al., 2015b; Tompary & Davachi, 2017; Tse et al., 2007, 2011). Taken together, memory consolidation involves an intricate, temporally evolving interplay of different brain regions, networks, and cognitive processes to enable long-term retrieval of past experiences.

1.1   Memory consolidation across development

From an ontogenetic perspective, the ability to retrieve long-term declarative memories emerges with the close of childhood amnesia, i.e., the inability to recollect early life events, around four to seven years of age (Alberini & Travaglia, 2017; Hayne, 2004). This indicates the kickoff of long-term memory stabilization (Bauer, 2006; Henke, 2010; Morgan & Hayne, 2011; Perner & Ruffman, 1995; Scarf et al., 2013; Tustin & Hayne, 2010). In Germany the transition from kindergarten to school occurs at the age of 6 years and is accompanied by dramatic increase in learning and knowledge accumulation, demands posed on children due to a structured primary school curriculum (Linberg et al., 2013; Mashburn et al., 2018). It challenges not only cognitive abilities but potentially also memory consolidation processes in this age group (Brod et al., 2017; Nolden et al., 2021). Although successful retrieval of complex memory representations starts to steadily improve at the age of 5 to 6 years (Drummey & Newcombe, 2002; Riggins, 2014; Sluzenski et al., 2006), the rate of memory retention across longer consolidation periods (Murre & Dros, 2015; Ebbinghaus, 1885; Müller & Pilzecker, 1900) is much less researched and may progress with different temporal dynamics in children in comparison to adults. For instance, it has been shown that short-delay memory consolidation rates (i.e., measured after one night of sleep)

were comparable between children aged 6 to 8 years and young adults for word-pair associates (Wilhelm et al., 2008), and were even more efficient in 7 to 12 years old children in comparison to young adults for non-objects and their functions (Peiffer et al., 2020a) and for "what-where-when" memories of faces (Wang et al., 2018). At the same time, findings show that in children, successful retrieval of events over a long time increases with increasing age (Boyer et al., 1994; Østby et al., 2012a; Peterson, 2002). However, little is known about memory consolidation of complex representations over shorter and longer delays in 6-year-old children who are about to start or are just starting school, as it has not yet been systematically investigated in the field of developmental neuroscience.

The core neural structures involved in memory consolidation undergo profound morphological changes in children. For instance, the HC reaches its relative maturity in middle childhood (Keresztes et al., 2017; Shing et al., 2010; Uematsu et al., 2012, but also see Lee et al. 2016), which is concomitant with an improvement in the ability to bind event features into a coherent representation (Lee et al., 2016; Shing et al., 2008, 2010; Sluzenski et al., 2006). On the other hand, memory control processes (e.g., strategy use, resolving mnemonic interferences) that benefit encoding, consolidation, and later retrieval of memories depends more on the ventrolateral prefrontal cortex (vlPFC; Badre & Wagner, 2007; Kuhl et al., 2012), while schema-integration processes that benefit long-term consolidation depend more on the vmPFC (Brod & Shing, 2018). Each of these regions shows more prolonged maturation and reaches its relative maturity only in late adolescence/young adulthood (Gogtay et al., 2004a; Muftuler et al., 2012; Shing et al., 2010a; Sousa et al., 2018; Uda et al., 2015; Waber et al., 2007). In an exceptional study that examined consolidation and brain correlates in participants aged 8 to 19 years, Østby et al. (2012a) showed that a thinner orbitofrontal cortex (OFC) was associated with a higher short-delay memory retention, while bigger HC volumes were related to higher memory retention rates after 1-week delay in a visuospatial task. Taken together, these findings indicate that extended developmental trajectories of HC and PFC maturation may affect the efficiency of memory consolidation processes in children, beyond their effects on encoding or retrieval (Shing et al., 2008b, 2010b).

1.2 Preterm birth and neurocognitive development

Prenatal and perinatal conditions can alter the developing brain (Buss et al., 2012), among which premature birth has been shown to greatly affect brain and neurocognitive development (Nosarti & Froudist-Walsh, 2016). Approximately 11% of children are born prematurely (i.e., born before the completion of 37 weeks of gestation; WHO, 2021) annually worldwide (Walani, 2020). Premature birth is associated with altered structural brain anatomy in regions critical to memory

consolidation – i.e., reduced grey matter (GM) and white matter (WM) volumes in temporal and frontal brain regions (Ball et al., 2012; Nosarti et al., 2014a; Nosarti & Froudist-Walsh, 2016). Reduced HC volume in preterm born individuals has been observed from infancy through childhood, adolescence, and into adulthood (Aanes et al., 2020; Arhan et al., 2017; Ball et al., 2012; de Kieviet et al., 2012). A growing body of research suggests that brain injury or altered neurodevelopmental trajectories due to premature birth are related to impaired cognitive functioning (e.g., information processing speed, verbal abilities; Linsell et al., 2015; Murray et al., 2014; Rose et al., 2011; Stålnacke et al., 2019), executive functioning (e.g., working memory, inhibition; Arpi et al., 2019; Beauchamp et al., 2008; Ford et al., 2011), and spatial learning (Cimadevilla et al., 2014). Such deviation in brain functioning may pose a special challenge for preterm born children during the transition from kindergarten to school.

Evidence for the effect of preterm birth on memory functioning in children is more mixed. For example, preterm born preschool children showed impaired verbal short-term memory in comparison to their term born peers (Baron et al., 2012; Dall'Oglio et al., 2010; Roberts et al., 2011). It was also reported that 7-year-old preterm born children showed imparied learning and immediate spatial and visual memory (Thompson et al., 2013a, 2014) and deficits in learning and immediate and delayed (20 minutes) visuo-spatial memory (Omizzolo et al., 2014). In addition, Quesada et al. (2014) reported poorer general and visual memory in preterm born children aged 6 to 10 years. On the other hand, no deficits in delayed (20 minutes) visuo-spatial recognition (Omizzolo et al., 2014a) for 7-year-old preterm born children was reported as well as no impairment in delayed verbal and visual recognition in preterm born children aged 6 to 10 years (Quesada et al., 2014). Taken together, current empirical evidence suggests inconsistent patterns of memory functioning in children born preterm. Of note is that preterm born cohorts examined in previous research are often very heterogenous, including individuals with various perinatal complications (e.g., insults, intraventricular hemmorage, etc.) and known learning disabilities. The majority of studies did not apply inclusion criteria for general cognitive functioning (e.g., IQ), which may confound the results comparing preterm born cohorts with typically developing term born children (Fernández de Gamarra-Oca et al., 2021). Importantly, temporal dynamics of long-term memory consolidation in 6-year old preterm born children has not yet been systematically characterized.

Therefore, thus far it is unclear how efficient long-term memory consolidation is in term born and preterm born children from early and middle childhood, a developmental stage during which brain regions underlying the ability to bind events into a coherent representation, to store memories , and to use strategies and exert control over memory retrieval, undergo profound

changes (Lenroot & Giedd, 2006; Mills et al., 2016; Ofen et al., 2007; Shing et al., 2010a). In this study, we examined the consolidation of object-location associations across a 2-week-period, comparing 6-year-old full term born and preterm born children to young adults serving as a reference group with a mature memory consolidation system. To counteract the sample heterogenity that may confound previous research on the long-term neurocognitive consequences of premature birth and increase sample comparability, we included only typically developing term born and preterm born children without further birth complications and with at least average IQ. Using an object-location association task, we tracked initial learning of new information up to a predetermined criterion, as well as memory consolidation thereafter following a short delay (after one night of sleep) and a long delay (after two weeks). To minimize variances attributed to encoding, we used a repeated retrieval-encoding cycles procedure to bring initial learning to a memory accuracy criterion of 83% in all participants, so that the comparison of subsequent memory consolidation could be made with starting points as similar as possible.

We hypothesized small or no differences in short-delay memory consolidation between children and adults (Peiffer et al., 2020a; Wang et al., 2018), but less efficient long-delay consolidation in children in comparison to young adults, as memory consolidation over time is particularly dependent on the interactions between the HC and neocortical regions that are still maturing in preschool children (Ghetti & Bunge, 2012; Lebel et al., 2012; Shing et al., 2010a). Based on previous reports of maturational differences in hippocampal-cortical circuitry between preterm born and term born children (Nosarti & Froudist-Walsh, 2016), we predicted less efficient memory consolidation in preterm born children in comparison to term born children and young adults. We furthermore expected that thinner medial OFC and IFG would be associated with better short-delay memory consolidation, reflecting schema-integration and strategic control over memory, respectively (Badre & Wagner, 2007; Kharitonova et al., 2013; Leon et al., 2010; Shing et al., 2010b), and that larger HC volumes would be associated with long-delay memory consolidation (Gilboa, 2004; Østby et al., 2012b; Steinvorth et al., 2005; Tompary & Davachi, 2017). We expected these neural-behavioural associations to be more pronounced in children compared to adults. While our focus is on memory consolidation, we also explored potential group differences during the initial learning phase. Here, we expected similar final learning performance but less efficient learning efficiency of object-location associations in preterm born children in comparison to term born children. We also expected that thinner OFC would be related to more efficient final learning performance (Botdorf & Riggins, 2018; Burzynska et al., 2012; Østby et al., 2012a; Shing et al., 2010b).

2   METHODS

2.1   Participants

Preterm born children (born before 34 weeks of gestation) were recruited, either from the existing well-characterized cohort at the Department of Neurology and Neonatology of the Center for Chronically Sick Children (Sozialpädiatrisches Zentrum), Charité – Universitätsmedizin Berlin, or through word-of-mouth. For the recruitment of term born children, 2000 general research invitation letters were sent to randomly selected families with 6-year-old children in Berlin, of whom 60 families expressed interest in participation. After screening, 44 typically developing term born children were recruited to participate in the study. 35 term born young adults were recruited to participate in the study through advertisement in the newspaper, on the university campus, and through word-of-mouth.

All participants had normal vision with or without correction, no history of psychological or neurological disorders or head trauma and had a normal cognitive ability with an IQ > 85 assessed using the K-ABC II (Kaufman & Kaufman, 2015). Exclusion criteria were severe perinatal incidents such as hypoxia-ischemia, intraventricular hemorrhage or infection, necrotic enterocolitis, and/or severe bronchopulmonary dysplasia. 14 term born children were excluded due to incomplete task execution and missing data (n=7) or technical issues during data acquisition (n=7). 10 preterm born participants were excluded due to incomplete task execution and missing data (n=8). Four young adult participants were excluded due to incomplete task execution and missing data (n=2), and identification as an outlier (n=2) due to chance performance. Thus, the final sample size consisted of 30 typically developing term born children (12 female, mean age 6.37 years, SD = 0.36), 15 preterm born children (7 female, mean age 6.25 years, SD = 0.61), and 33 term born young adults (15 female, mean age 25.45 years, SD = 2.80; see Table 1).

All participants or their legal guardians gave written informed consent prior to participation. The study was approved by the ethics committee of the Goethe University Frankfurt am Main (approval E 145/18). The participants were compensated for participation on the three days of the experiment with 100 Euro.

Table 1. Sample characteristics by age group (preterm born children, term born children, young adults) showing comparability between groups

	Preterm born Children (PC; N = 15)		Term born Children (TC; N = 30)		Young Adults (YA; N = 33)		Group effect	PC vs. TC	TC vs. YA	PC vs. YA
	M	SD	M	SD	M	SD	p-value	ω^2	p-value	
Demographic measures										

	M	SD	M	SD	M	SD					
Age	6.3	0.6	6.4	0.4	25.5	2.8	***	0.96	ns	***	***
Gestational Age	28.1	2.7	39.8	1.5	39.7	1.7	***	0.86	***	ns	***
Sex (M/F)	8/7		18/12		18/15						
IQ score	112.9	13.3	116.1	15.1	107.7	12.1	*	0.05	ns	*	ns
Socioeconomical Status							ns				
Income – Family	3.9	1.4	3.8	1.8	-	-	ns				
ISCED – Family	5.3	1.5	5.9	1.6	4.3	2.0	**	0.14	ns	**	*
Education Years - Mother	17.3	4.0	19.5	3.5	-	-	.	0.06			
Education Years - Father	16.1	4.0	19.2	5.1	-	-	.	0.07			

Notes. Income is based on a 1-7 Scale (1= less than 15.000€, 7 = more than 100.000€). TC = term born children; PC = preterm born children; YA = young adults; ISCED = International Standard Classification of Education (2011); IQ = Intelligence Quotient based K-ABC (Kaufman & Kaufman, 2012) for term born and preterm born children and WAIS-IV (Wechsler, 2012); M = mean; SD = standard deviation; *p < .05; ** <.01, ***<.001 (significant difference); ns: non-significant difference.

2.2 Task design and procedure

2.2.1 Stimulus material and task

The stimuli were chosen based on the curriculum in social studies and science for the first and second grade of the German primary school (see similar procedure in Brod & Shing, 2019). Sixty different semantic themes (e.g., forest, farm, etc.) were chosen according to the ratings provided by four primary school teachers that assessed the familiarity of first graders with the topics. For each semantic theme, four scene pictures were combined with four thematically congruent object pictures resulting in four unique object-location associations (see Fig. 1 for example). We identified 18 possible placement areas, one of which was assigned to each object-location association (for more detailed information on task description refer to Fig. 1 and to the Methods section in the Supplementary Materials). We presented the task using Psychtoolbox-3 (Kleiner et al., 2007) software in Matlab 9.5, R2018b (The MathWorks, Inc., MA, USA).

Figure 1. **Task design and procedure.** (**A**) *Initial Encoding Phase*. For a set of 60 object-location pairs, participants saw the object followed by the same object superimposed on the scene at a particular location. Participants were instructed to remember 60 object-location pairs in total, memorizing the exact location of the object within the scene by creating a story or making a "mental photo" of the scene as the such semantic organization strategies aid the recollection of the information (Craik & Lockhart, 1972); (**B**) *Learning Phase*. Participants learned the correct location of the object within the scene during retrieval-encoding cycles. They had to choose the correct location out of three choices and received feedback for their response. After the feedback, the correct object-location associations were shown again. The cycles ended when participants provided correct responses to 83% of the trials (minimum two cycles, maximum four cycles). (**C**) *Retrieval Phase.* Participants had to choose the correct location of the object in the scene out of three options without feedback. *Note:* The retrieval phase was carried out inside the MRI scanner with a functional sequence, of which the data is not included as we focus on characterizing the retention rate behaviourally, both in terms of group comparison and relations to structural integrity.

2.2.2   Assessment of demographic and cognitive covariates

In addition to the experimental paradigm (memory consolidation task described below), IQ scores were assessed using the German version of the "Kaufman Assessment Battery for Children – Second Edition" (K-ABC II; Kaufman & Kaufman, 2015) in children and the "Wechsler Adult Intelligence Scale – Fourth Edition" (WAIS-IV; Wechsler, 2012) in young adults. General socio-demographic questionnaires to assess socio-demographic characteristics of the participants were applied as well. Other cognitive covariate tasks were also assessed, such as cognitive switching and object-location memory with immediate test but were not included in the current paper to retain the focus on memory consolidation (see Fig. 2).

2.2.3   Experimental procedure

Testing took place across three days. At Day 0, the experimental procedure began with a short training to familiarize participants with the task. The experimental task started with the initial encoding, which was followed by retrieval-encoding cycles until the 83% accuracy criterion was

reached. On Day 1 (short delay) and Day 14, retrieval was conducted (see Fig. 2 for a detailed description of the experimental procedure).

Figure 2. **Experimental Procedure**. The testing took place across three days. (i) At Day 0 participants had to learn 60 object-location associations. The experimental task started with the initial encoding of the first 30 object-location associations. The initial encoding was followed by a brief distraction task in which participants listened to and had to recall the string of numbers. The distraction task was followed by the learning phase with retrieval-encoding cycles until the 83% criterion. After a short break, the same procedure was repeated with the other half of 30 object-location associations. After the experimental task, several subtests of the K-ABC II Test *(e.g., Atlantis, Rover, Rebus, Riddle and Atlantis delayed) were administered to children, while young adults were tested with the WAIS-IV Test. (ii) On Day 1 (short delay) the retrieval was conducted. Participants had to retrieve 30 object-scene associations learnt on Day 0. In addition, children performed several subtests of the K-ABC II Test *(e.g., Expressive Vocabulary, Triangles, Pattern Reasoning). (iii) On Day 14 (long delay) the retrieval was conducted. Participants had to retrieve another 30 object-scene associations learnt on Day 0. In addition, children performed several subtests of the K-ABC II Test *(e.g., Patterns, Verbal Knowledge, Word Order). *Note:* * denotes specific subtests of K-ABC II distributed across three testing days.

2.3   Magnetic resonance imaging

MR images were acquired on a 3-T SIEMENS PRISMA scanner (Siemens Medical Solutions, Erlangen, Germany) using a 64-channel head coil. A MPRAGE (magnetization prepared rapid gradient echo) T1-weighted sequence was applied with the following parameters: time repetition/time echo/time to inversion/Flip Angle = 2400 ms/2.22 ms/1000 ms/8°, matrix 256 x 256, field of view = 256. Each scan took 6 min 38 s. Each volume consisted of 320 sagittal slices with voxel sizes 0.8 x 0.8 x 0.8 mm.

2.4 Data Analysis

2.4.1 MRI data processing

Volumes and cortical thickness measures used in the current analyses were derived using the anatomical pipeline of fMRIprep (version 20.2.1; Esteban et al., 2019), based on Nipype 1.5.1 (Gorgolewski et al., 2011). Brain tissue segmentation of cerebrospinal fluid, white-matter, and grey-matter was performed on the brain-extracted T1w using FAST (FSL 5.0.9; Zhang et al., 2001). Brain surfaces were reconstructed using recon-all (FreeSurfer 6.0.1; Dale et al., 1999). Volume-based spatial normalization to two standard spaces was performed through nonlinear registration with antsRegistration (ANTs 2.3.3), using brain-extracted versions of both T1w reference and the T1w template. Intracranial volume (ICV) was derived by the surfaced-based measures. Thickness measures were calculated from the distance between the reconstructed white matter and grey matter surfaces as well as from the reconstructed grey matter surface and cerebrospinal fluid boundaries (for a more detailed description of the segmentation procedure, please refer to the Methods section in Supplementary Materials).

2.4.2 Behavioural Data Analysis

The analyses of all behavioural measures were performed with R packages (Version 4.0.4, R Core Team, 2021) in R Studio 1.4.1106 (RStudio, Inc.). Throughout the analyses, the α levels were set to $\alpha < .05$. We conducted a linear mixed-effect model for memory measures (accuracy defined as percentage of correct responses) using the lmer function from the lme4 package in R (Bates et al., 2015) and lmerTest (Kuznetsova et al., 2017). The linear mixed effect model was calculated with maximum-likelihood estimation and *Subject* as random intercept to account for between-subject variability in memory accuracy. As fixed factors, we included the within-subject factor of *Session* (final performance on Day0, short delay, and long delay) and the between-subject factor of *Group* (PC, TC, and YA). In addition, *IQ*, *Sex,* and *Handedness* were added as covariates into the model. The main effects were followed up with Tukey post-hoc multiple comparisons (Haynes, 2013). For group differences in learning measures, we conducted one-way independent analysis of variance (ANOVA). In case of violated assumptions of homogeneity of variances, a Games-Howell test was performed (Lee & Lee, 2018). Moreover, Games-Howell is especially suitable for the groups with unequal sample sizes. The effect size estimation was performed with omega squared (ω^2) as a less biased estimate for reporting practical significance of observed effects (Finch & French, 2012; Okada, 2013; Troncoso Skidmore & Thompson, 2013). To determine the amount of variance explained by the model, we used partR2 package in R (Stoffel et al., 2020) with bootstrapping to calculate confidence intervals.

2.4.3 Analysis brain-behavioural associations

Based on the literature, we focused on the following brain regions of interest: HC volume, IFG cortical thickness (consisting of pars opercularis and pars triangularis), and medial OFC cortical thickness (Dudai, 2004; Kharitonova et al., 2013; Leon et al., 2010; Moscovitch & Gilboa, 2021; Østby et al., 2012b; Petrides, 2007). To control for head size, hippocampal volumes were adjusted for ICV using a residual approach, which has been suggested to be less susceptible to bias (Bathelt et al., 2018; Mathalon et al., 1993; Sanfilipo et al., 2004; Voevodskaya et al., 2014). Cortical thickness was adjusted for sex but not head size because cortical thickness and head size are not associated (Barnes et al., 2010; Mills et al., 2016). We conducted a linear mixed-effect model analysis of memory measures (accuracy defined as percentage of correct responses) using the lmer function from the lme4 package in R (Bates et al., 2015) and lmerTest (Kuznetsova et al., 2017). The linear mixed effect models were calculated with maximum-likelihood estimation and *Subject* as a random intercept. As fixed factors, we included the within-subject factor of *Session* (final performance on Day0, short delay, and long delay), the between-subject factor of *Group* ((PC, TC, and YA) or (children and adults)), and structural measures (HC volume, IFG cortical thickness and medial OFC cortical thickness) separately for left and right brain hemisphere, expecting task-related lateralization (Bauer et al., 2019; Ezzati et al., 2016; Hopf et al., 2013). In addition, *Sex was* added as a covariate in the model to account for possible differences in brain volumes and cortical thicknesses (Mills et al., 2016). Regression analyses were performed to assess associations between final learning performance and brain structural measures (medial OFC cortical thickness). Group differences in brain structure were assessed using one-way independent ANOVA, which was followed by either post hoc Tukey comparisons or a Games-Howell test, depending on the homogeneity of variances.

3   RESULTS

3.1   Final Learning and Learning Efficiency at Day 0

In the following, we first tested for potential group-related differences in learning object-location associations during the learning phase. The one-way independent analysis of variance (ANOVA) revealed a significant effect of *Group* on final learning accuracy, $F_{(2,75)} = 45.73$, $p < 001$, $\omega^2 = .53$. Post hoc Games-Howell tests revealed that term born children learnt significantly more object-location associations than preterm born children, $p = .049$. Young adults showed significantly higher final learning accuracy than term born children, $p < .0001$, and preterm born children, $p < .0001$ (see Fig. 3). We also derived an index of learning efficiency, which is calculated as the

average of the standardized *z* scores of initial encoding, the learning slope, and final learning accuracy (Zerr et al., 2018). There was a significant effect of *Group* on learning efficiency, $F_{(2,75)} = 48.19$, $p < 0.001$, $\omega^2 = .55$. Post hoc Games-Howell tests revealed that learning efficiency did not differ between preterm born and term born children, $p = .27$, while the learning efficiency of young adults was significantly higher than that of term born children, $p < .0001$, and that of preterm born children, $p < .0001$ (see Fig. 3).

Figure 3. Learning accuracy and efficiency. (**A**) Final learning accuracy shows the percentage of correct responses after learning was completed in preterm born children, term born children and young adults. (**B**) Learning efficiency implies the learning rate and immediate retention rate for object-location associations as introduced by Zerr et al (2018). Error bars indicate $Q_{1/3} \pm 1.5 \cdot IQR$; *$p < .05$; **$p < .01$; ***$p < .001$(significant difference); ns: non-significant difference.

3.2 Memory Consolidation

In the following, group differences in short- and long-delay memory consolidation are presented. The linear mixed-effects model for retrieval accuracies of learned object-location pairs explained a significant amount of variance, $R^2 = .69$, 95% CI [.66, .76] (for a full overview, please see Table 2 in Supplementary Material). We observed a significant *Session* x *Group* interaction, $F_{(4,156)} = 4.71$, $p = .001$, $\omega^2 = .08$. As a follow-up to this interaction, we calculated model-based Tukey post hoc comparisons for short and long delay separately. The between group comparisons for short delay revealed that (i) retrieval accuracy in all groups after one night of sleep declined significantly (all $p < .05$); (ii) there was no difference in the slope of accuracy decline between groups (all $p > .12$), indicating a comparable efficiency of short-delay memory consolidation among 6-year-old children, irrespective of birth status, and young adults. For the long delay retrieval after a period of two weeks, Tukey post hoc between group comparisons

revealed (i) a significant decline in long delay retention rate in all groups (all p < .0001); (ii) no difference in long delay retention rates between preterm born and term born children, b = .003, $t_{(162)}$ = .08 p = .94; (iii) significantly higher long-delay retention rates in young adults than in both preterm born children, b = .105, $t_{(162)}$ = 3.14, p = .002, and term born children, b = .108, $t_{(162)}$ = 3.97 p = .0001, indicating more efficient long-term memory consolidation in young adults in comparison to 6-year-old children, irrespective of their birth status (see Fig. 4).

Figure 4. Memory consolidation and retention rates for short and long delay. (**A**) Memory consolidation over the course of two weeks operationalized by percentage of correctly retrieved object-location associations learned on day 0 (final accuracy after learning), on day 1 (after one night of sleep) and on day 14 (after two weeks). Shaded areas indicate 1 standard error of the mean. (**B**) Retention rate for short (after one night of sleep) and long (after two weeks) delay across preterm born children, term born children and young adults. Error bars indicate 1 standard error of the mean. *p < .05. **p < .01. ***p < .001(significant difference), ns: non-significant difference. P-values use Tuckey correction for multiple comparisons.

3.3   Brain-behavioural relationships

3.3.1   Hippocampal Volume and Memory Consolidation

The between group analyses showed that residualized left and right HC volumes were comparable between term born and preterm born children, while adults showed bigger left and right HC volumes in comparison to both groups of children (see Table 2). Based on this observation, term born and preterm born children were combined into one group for the subsequent analyses (referred to as the children group). Regarding memory consolidation,

separate linear mixed-effects models were conducted with left and right HC volumes as predictors of memory retention rate, respectively. Only right HC volumes, but not left HC, significantly predicted retention rates, irrespective of short or long delay and age group, R^2 = .70 95% CI [.66, .77], $F_{(1,75)}$ = 5.98, p = .016, ω^2 = .06 (see Fig. 5). Moreover, there was a significant *right HC x Group* interaction, $F_{(1,75)}$ = 6.01, p = .016, ω^2 = .06, indicating an age effect in the relation of right HC and memory consolidation. This interaction was probed by conducting two separate multilevel models for the group of children and the group of young adults with the same setup as above but without the *Group* factor, as in (Field et al., 2012). The analysis showed that for children, higher right HC volume was significantly associated with long-delay retention rates, b = .17, t(2.11), p = .037, but not short-delay retention rates, b = .08, t(1.04), p = .29, while such association was not significant in adults (all p > .65). An overview comprising the statistics of all effects of the applied models can be found in the supplementary materials (see Supplementary Results, Tab. S4-S7).

Table 2. *Results of the analyses on group differences for brain structural measures (HC volumes, medial OFC thickness, IFG thickness). Statistical values are shown for the ANOVA analyses and follow-up post hoc tests.*

Region	ANOVA			TC vs PC		TC vs YA		PC vs YA		Post hoc Test
	$F_{(2,72)}$	*p-value*	ω^2	β	*p-value*	β	*p-value*	β	*p-value*	
Right HC	9.84	***	.19	-1.62	ns	333.83	***	335.5	***	Games-Howell
Left HC	13.5	***	.25	11.4	ns	495.19	***	447.8	***	Games-Howell
Right mOFC	102.7	***	.73	-.14	*	-.42	***	-.56	***	Tukey
Left mOFC	148.9	***	.80	-.18	***	-.39	***	-.58	***	Tukey
Right IFG	76.8	***	.67	.03	ns	-.54	***	-.51	***	Tukey
Left IFG	43.6	***	.53	.01	ns	-.45	***	-.44	***	Tukey

Notes. ANOVA = analysis of variance; TC = term born children; PC = preterm born children; YA = young adults; HC = hippocampus; mOFC = medial orbitofrontal thickness; IFG= inferior frontal gyrus thickness (comprised of parsopercularis and parstriangularis); β = estimate showing difference in means; *p < .05; ** <.01, ***<.001 (significant difference); ns: non-significant difference.

3.3.2 Frontal Cortical Thickness and Memory Consolidation

The between group analyses showed that left and right medial OFC was thicker in preterm born children in comparison term born children, while adults showed thinner right medial OFC than both child groups. Furthermore, the between group analyses showed that left and right IFG cortical thickness (i.e., pars opercularis and pars triangularis) were comparable between term

born and preterm born children, while adults had thinner left and right IFG cortical thickness in comparison to both child groups. As we expect a similar relation of cortical thickness measures to memory measures in both groups of children, they were combined into one group for the subsequent analyses (referred to as the child group; Nam et al., 2015).

With respect to memory consolidation, linear mixed-effects models were conducted separately with left and right medial OFC thickness as predictors of memory retention rate. Only a main effect of right medial OFC thickness, $F_{(1,75)} = 4.64$, p = .03, $\omega^2 = .05$, was observed, indicating that thinner right medial OFC is related to higher overall retention rates irrespective of age group and delay. No such effect was observed for the left medial OFC (all p > .09). The linear mixed-effect models conducted with left and right IFG thickness as predictors of memory retention rate revealed no significant effects (all p > .19).

3.3.3 Exploratory analysis with final learning performance

Finally, we also explored brain-behavioural relationships for day 0 final learning performance. First, the association between the medial OFC thickness and final learning performance was investigated. As revealed by regression analyses, there was a marginally significant association between right medial OFC thickness and final learning performance, $F_{(1,71)} = 3.47$, p = .06, $\omega^2 = .03$, which differed by group, $F_{(1,71)} = 4.88$, p = .03, $\omega^2 = .05$. Moreover, left OFC thickness was also significantly associated with final learning performance, $F_{(1,71)} = 9.43$, p = .003, $\omega^2 = .10$, irrespective of age group, $F_{(1,71)} = 1.88$ p = .06, $\omega^2 = .02$. Follow up tests on observed groups differences revealed that in children, thinner right medial OFC thickness was predictive of final learning performance while thinner left OFC was associated with final learning performance in both groups of children and in adults (see Fig. 5). For completeness, we also explored the relationship between the IFG and HC with final learning performance, which did not yield any significant results (all p > .26). Therefore, only the medial OFC showed significant relationship with final learning performance.

Figure 5. Relation between hippocampus volumes, retention rate, and learning accuracy.
(**A**) Scatterplots depicting associations between short-delay and long-delay retention rates with residualized right HC volume. Retention rate on the y-axis is measured as a proportion between final accuracy on Day 0 and memory performance on Day 1 (short delay$_{day1/day0}$) and between final accuracy on Day 0 and memory performance on Day 14 (long delay$_{day14/day0}$). On the x-axis the residualized right HC volume was corrected for head size (using ICV) using the residual approach (Mathalon et al., 1993; Voevodskaya et al., 2014). (**B**) Scatterplot showing the relation between the final learning accuracy on Day 0 and right medial orbitofrontal cortical thickness (centered). HC: Hippocampus; ICV: intracranial volume.

4   DISCUSSION

In the present study, we investigated memory consolidation of object-location associations after both short delay (after one night of sleep) and long delay (after two weeks), in preterm born and term born 6-year-old children as well as young adults, and examined their associations with medial temporal and frontal structural brain measures. Given postulated differences among the three groups in terms of cognitive functioning and structural brain integrity in general, we took several steps to hone in on differences in memory consolidation by reducing group differences in general intellectual functioning or in the encoding of new information. All participants learned the object-location associations to a high degree of accuracy at day 0. Nevertheless, we found that preterm born children showed impaired final performance in learning object-locations associations when compared to term born children, who in turn were outperformed by young adults. We therefore took performance on day 0 into account in our statistical analysis of memory consolidation. Here, our main findings are: (i) preterm born children, term born children, and young adults showed comparable short-delay retention rates after one night of sleep; (ii) the two child groups showed comparable retention rates over long delay, but these rates were significantly

lower than those of young adults. In other words, children showed steeper forgetting of information over a longer time period.

Taking a step further, we tested for relationships between structural measures of key brain regions involved in memory consolidation and behavioural performance. In both children and young adults, a thinner medial orbitofrontal cortex correlated with higher retention rates, while no such association was found for IFG. Furthermore, we found that memory retention rates were positively associated with larger right HC volume uniquely in children. In addition, thinner medial OFC was associated with higher final learning performance in both children and adults. These results will be discussed in the following sections.

4.1 Short- and long-delay memory consolidation

4.1.1 Similar short- and long-delay memory consolidation in term born and preterm born children

Contrary to our hypothesis, there was no difference in short-delay and long-delay retention rates between preterm born and term born children, indicating similar memory consolidation performance despite premature birth. This finding may be attributed to a highly selected cohort of preterm born children that was comparable to the term born children in their general ability to learn and acquire knowledge as well as exhibiting comparable world knowledge according to subscales of K-ABC II (see Supplementary Materials for further information). These abilities benefit long-delay memory consolidation that relies on schemas or complex knowledge systems (Aghayan Golkashani et al., 2021; Bartlett, 1995). Interestingly, it has been shown that comparable memory performance can be achieved in adult preterm born individuals by engaging different brain circuits during encoding and successful retrieval (Brittain et al., 2014; Narberhaus et al., 2009; Salvan et al., 2014; Tseng et al., 2017). This may indicate a high degree of neuroplasticity in individuals born prematurely to already begin to compensate for alterations in the brain early in developmental, accounting for our findings of comparable memory consolidation between term born and preterm born children.

4.1.2 Similar short-delay but less efficient long-delay memory consolidation in term born and preterm born children in comparison to young adults

As expected, adults and children showed comparable short-delay retention rates, indicating that short-delay memory consolidation after one night of sleep in 6-year-old children was as efficient as in young adults irrespective of preterm born status. However, both groups of children showed steeper accuracy percentage change and thus lower long-delay retention rates in comparison to young adults, indicating reduced efficiency in retaining memory of complex associative information over longer time periods in children. This evidence is in line with the findings on

comparable or even more efficient short-delay memory retention rates in primary school children than in young adults (Peiffer et al., 2020; Wang et al., 2018), which may be due to a stronger effect of overnight sleep on short-delay memory consolidation in children in comparison to young adults, compensating for possible maturity-related mnemonic benefits in adults and also pointing at an early advantage of sleep for preschool children (Peiffer et al., 2020). On the other hand, superior long-delay memory consolidation in young adults may be attributed to their more extensive world knowledge and therefore more elaborate schemas that may be especially important for ensuring better accessible memory representations in long delay over short delay (Craik & Lockhart, 1972; van Kesteren et al., 2012). Moreover, these mnemonic benefits may be augmented by mature HC-vmPFC interaction that underlies long-term memory consolidation in young adults (Takashima et al., 2006; Yamashita et al., 2009). In children, the extended developmental trajectory into mid-childhood for the HC and into adulthood for the vmPFC (Gogtay et al., 2004b; Keresztes et al., 2017b; J. K. Lee et al., 2016b; Shing et al., 2010a) may have limited the efficiency of consolidating memory into neocortical regions over long delay.

4.1.3 Thinner right HC volume is associated with long-delay memory consolidation in children and thinner right mOFC thickness is associated with overall memory consolidation in all age groups

The premature birth-related brain structural differences previously reported in the literature (Nosarti et al., 2014b; Nosarti & Froudist-Walsh, 2016) could not be found in our study. Contrary to our expectations, there was no difference in HC volume or IFG cortical thickness between preterm born children and term born children. The only difference found within our ROIs was that preterm born children showed thicker medial OFC in comparison to term born children. Adults showed larger HC volumes and thinner IFG and medial OFC than both groups of children, which is in line with existing findings on developmental trajectories of brain morphology (see Hedman et al., 2012 for an overview).

Importantly, in children, higher right HC volumes were positively associated with long-delay memory retention, indicating that more mature right HC is associated with better memory consolidation over longer time periods in children irrespective of premature birth status. This finding is convergent with the established critical role of the HC in relational memory (Konkel, 2009). Concerning lateralization, in line with our findings, Hopf et al. (2013) could show that more adult-like right HC lateralization in children was associated with superior memory on a relational memory task. Moreover, activation in the right HC has been reported to be associated with spatial memory processing (Bohbot et al., 1998; Ezzati et al., 2016) and with memory for locations within an environment (Burgess et al., 2002). The lack of association between HC volumes and memory

retention rates in young adults may be attributed to the relative independence of memory consolidation processes on HC volumetric characteristics in adults compared to children, in which volumetric increase corresponds to maturation. It has been proposed that in young adults, HC involvement in memory consolidation is more pronounced in the functional dynamic interplay with other neocortical regions (Frankland & Bontempi, 2005).

On the other hand, thinner right medial OFC was related to higher overall retention rates irrespective of age group, in line with its role in schema-related memory integration through perceived congruency between to-be-learnt information and prior knowledge (Brod & Shing, 2018; Tse et al., 2007). This finding goes beyond previous functional evidence, showing that a structurally thinner medial OFC is also related to better memory formation and retention over time. Convergingly, Bauer et al. (2019) showed that right medial OFC thickness was related to memory of self-derived knowledge in 5 to 8 years old children. Memory recall has also been shown to be related to OFC thickness in 8 to 19 years old participants, irrespective of age (Østby et al., 2012a). Therefore, our results suggest that structural measures of OFC are related to memory consolidation as well, with thinner OFC indicating higher maturity and being associated with better memory integration with prior knowledge regardless of age (Schlichting & Preston, 2015).

Contrary to our expectations, IFG thickness was not associated with memory consolidation irrespective of age group and delay, despite functional brain imaging findings suggesting an association between the IFG and semantic organizational strategies for verbal learning (Savage et al., 2001) and monitoring strategic control processing over retrieval of memory (Badre & Wagner, 2007; Kuhl et al., 2012), as well as semantic elaborative strategies during memory retrieval that arise deliberately from prior encoding of distinctiveness (Han et al., 2012; Raposo et al., 2009). Therefore, structural morphology of IFG may not be associated with memory consolidation beyond its functional involvement in encoding and retrieval when the focus is not on distinctness or semantic elaborations aimed to capitalize on it for long-term memory retention.

4.2   Learning

4.2.1   *No difference in learning efficiency between preterm and term born children but reduced learning final performance in preterm in comparison to term born children*

While our focus was on memory consolidation, we also explored groups differences in Day0 learning. We found no difference between preterm born and term born children with respect to learning efficiency, indicating a comparable rate of learning. However, preterm born children showed reduced final learning accuracy in comparison to term born children. This finding may be related to the impaired visuo-spatial pattern recognition (Marlow et al., 2007; Schermann & Sedin,

2007) and visuo-spatial memory (Omizzolo et al., 2014b; Thompson et al., 2013b) reported for preterm born children. Indeed, in our sample, despite comparable general cognitive functioning, there was a small difference found in the simultaneous subscale of the K-ABC II Test (refer to Supplementary Materials for further information), suggesting a lower ability for simultaneous processing of visuospatial information in preterm born children, which may have limited the amount of information they could learn in our task.

4.2.2 *Less efficient learning in term born and preterm born children in comparison to young adults*

In line with our expectation, both groups of children were outperformed by young adults in final learning accuracy and learning efficiency. These findings indicate that young adults were able to retain more information after just a single exposure to the stimuli pairs, learn faster, and retrieve with higher final accuracy than both groups of children. The superior learning efficiency performance of young adults may be attributed to more efficient associative binding as well as better ability to exert strategic control over memory (Shing et al., 2008b, 2010b). The ongoing development of the PFC and HC in children as well as their lesser knowledge and poorer integration of newly learned information may have limited the efficiency of learning during the encoding-retrieval cycles (Shing et al., 2010c).

4.2.3 *Thinner medial orbitofrontal cortex is related to more efficient final learning performance in children and adults*

Final learning accuracy was positively related to thinner right medial OFC in children and to thinner left medial OFC across all participants. In line with this, a positive association between thinner frontal lobes with better immediate (30 minutes delay) memory was observed in visuospatial memory task in children aged 6 to 15 years (Sowell et al., 2001). In addition, Østby et al. (2012) showed that thinner left OFC was related to higher memory recall 30 minutes after learning independently of age in participants aged 8 to 19 years. Thus, these findings show that for to-be-remembered complex associations that are congruent with prior knowledge, thinner medial OFC is related to better memory formation (Brod & Shing, 2018). This association was already present bilaterally at the age of six, indicating more intense brain-behaviour interactions in the maturing brain, which continues to persist in mature brains in young adulthood as well.

5   LIMITATIONS

Several limitations of the current study should be noted. Despite our procedure of learning to the criterion to maximize comparability of retention rates, we nevertheless observed group differences in initial memory performance. Future studies may incorporate individualized item-

based approach of learning to criteria, excluding correctly remembered items from further learning cycles to ensure faster learning and lessening the overall task workload (Karpicke & Roedigeriii, 2007; McDermott & Zerr, 2019; Zerr et al., 2018). Moreover, due to the small sample of preterm born children, we may have been underpowered to detect memory consolidation differences in term and preterm born children. Therefore, future studies with a larger preterm born sample would help to draw more robust conclusions about potential differences in memory consolidation due to premature birth status (or gestational age) in well-characterized cohorts. In addition, the current findings concentrate mainly on associative memory of schema-congruent information. Future studies should investigate how violations of knowledge, namely schema-incongruent information, may impact the learning of associative information and their subsequent consolidation in short and long delay memory.

6 ORCHID

Iryna Schommartz https://orcid.org/0000-0001-8655-9259
Angela M. Kaindl https://orcid.org/0000-0001-9454-206X
Claudia Buß https://orcid.org/0000-0002-8738-3133
Yee Lee Shing https://orcid.org/0000-0001-8922-7292

Author contributions

Y.L.S, C.B., A.K secured funding. I.S and Y.L.S, C.B., A.K contributed to conception and design of the study. I.S., H.S., N.W.-C., and P.L. performed data collection and data curation. I.S., P.L., and M.B performed the statistical analysis. IS wrote the first draft of the manuscript, P.L. and M.B. wrote sections of the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

7   REFERENCES

Aanes, S., J. Bjuland, K., Lærum, A. M. W., Weider, S., Anne I. Evensen, K., S. Indredavik, M., Brubakk, A. M., Håberg, A., C. Løhaugen, G., & Skranes, J. (2020). Relationship between hippocampal subfield volumes and memory function in adults born preterm with very low birth weight (VLBW). *Clinical Obstetrics, Gynecology and Reproductive Medicine*, *6*(2). https://doi.org/10.15761/COGRM.1000281

Aghayan Golkashani, H., Leong, R. L. F., Wong, K. F., & Chee, M. W. L. (2021). Schema-driven memory benefits boost transitive inference in older adults. *Psychology and Aging*, *36*(4), 463–474. https://doi.org/10.1037/pag0000586

Alberini, C. M., & Travaglia, A. (2017). Infantile Amnesia: A Critical Period of Learning to Learn and Remember. *The Journal of Neuroscience*, *37*(24), 5783–5795. https://doi.org/10.1523/JNEUROSCI.0324-17.2017

Arhan, E., Gücüyener, K., Soysal, Ş., Şalvarlı, Ş., Gürses, M. A., Serdaroğlu, A., Demir, E., Ergenekon, E., Türkyılmaz, C., Önal, E., Koç, E., & Atalay, Y. (2017). Regional brain volume reduction and cognitive outcomes in preterm children at low risk at 9 years of age. *Child's Nervous System*, *33*(8), 1317–1326. https://doi.org/10.1007/s00381-017-3421-2

Arpi, E., D'Amico, R., Lucaccioni, L., Bedetti, L., Berardi, A., & Ferrari, F. (2019). Worse global intellectual and worse neuropsychological functioning in preterm-born children at preschool age: A meta-analysis. *Acta Paediatrica*, *108*(9), 1567–1579. https://doi.org/10.1111/apa.14836

Badre, D., & Wagner, A. D. (2007). Left ventrolateral prefrontal cortex and the cognitive control of memory. *Neuropsychologia*, *45*(13), 2883–2901. https://doi.org/10.1016/j.neuropsychologia.2007.06.015

Ball, G., Boardman, J. P., Rueckert, D., Aljabar, P., Arichi, T., Merchant, N., Gousias, I. S., Edwards, A. D., & Counsell, S. J. (2012). The Effect of Preterm Birth on Thalamic and Cortical Development. *Cerebral Cortex*, *22*(5), 1016–1024. https://doi.org/10.1093/cercor/bhr176

Barnes, J., Ridgway, G. R., Bartlett, J., Henley, S. M. D., Lehmann, M., Hobbs, N., Clarkson, M. J., MacManus, D. G., Ourselin, S., & Fox, N. C. (2010). Head size, age and gender adjustment in MRI studies: A necessary nuisance? *NeuroImage*, *53*(4), 1244–1255. https://doi.org/10.1016/j.neuroimage.2010.06.025

Baron, I. S., Kerns, K. A., Müller, U., Ahronovich, M. D., & Litman, F. R. (2012). Executive functions in extremely low birth weight and late-preterm preschoolers: Effects on working memory and response inhibition. *Child Neuropsychology*, *18*(6), 586–599. https://doi.org/10.1080/09297049.2011.631906

Bartlett, F. C. (1995). *Remembering: A study in experimental and social psychology*. Cambridge University Press.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Usinglme4. In *Journal of Statistical Software* (Vol. 67, Issue 1). https://doi.org/10.18637/jss.v067.i01

Bathelt, J., Gathercole, S. E., Johnson, A., & Astle, D. E. (2018). Differences in brain morphology and working memory capacity across childhood. *Developmental Science*, *21*(3), e12579. https://doi.org/10.1111/desc.12579

Bauer, P. J. (2006). Constructing a past in infancy: A neuro-developmental account. *Trends in Cognitive Sciences*, *10*(4), 175–181. https://doi.org/10.1016/j.tics.2006.02.009

Bauer, P. J., Dugan, J. A., Varga, N. L., & Riggins, T. (2019). Relations between neural structures and children's self-derivation of new knowledge through memory integration. *Developmental Cognitive Neuroscience*, *36*, 100611. https://doi.org/10.1016/j.dcn.2018.12.009

Bohbot, V. D., Kalina, M., Stepankova, K., Spackova, N., Petrides, M., & Nadel, L. (1998). Spatial memory deficits in patients with lesions to the right hippocampus and to the right parahippocampal cortex. *Neuropsychologia*, *36*(11), 1217–1238. https://doi.org/10.1016/S0028-3932(97)00161-9

Botdorf, M., & Riggins, T. (2018). When less is more: Thinner fronto-parietal cortices are associated with better forward digit span performance during early childhood. *Neuropsychologia*, *121*, 11–18. https://doi.org/10.1016/j.neuropsychologia.2018.10.020

Boyer, M. E., Barron, K. L., & Farrar, M. J. (1994). Three-year-olds Remember a Novel Event from 20 Months: Evidence for Long-term Memory in Children? *Memory*, *2*(4), 417–445. https://doi.org/10.1080/09658219408258957

Brittain, P. J., Froudist Walsh, S., Nam, K.-W., Giampietro, V., Karolis, V., Murray, R. M., Bhattacharyya, S., Kalpakidou, A., & Nosarti, C. (2014). Neural compensation in adulthood following very preterm birth demonstrated during a visual paired associates learning task. *NeuroImage: Clinical*, *6*, 54–63. https://doi.org/10.1016/j.nicl.2014.08.009

Brod, G., Bunge, S. A., & Shing, Y. L. (2017). Does One Year of Schooling Improve Children's Cognitive Control and Alter Associated Brain Activation? *Psychological Science*, *28*(7), 967–978. https://doi.org/10.1177/0956797617699838

Brod, G., & Shing, Y. L. (2018). Specifying the role of the ventromedial prefrontal cortex in memory formation. *Neuropsychologia*, *111*, 8–15. https://doi.org/10.1016/j.neuropsychologia.2018.01.005

Brod, G., & Shing, Y. L. (2019). A boon and a bane: Comparing the effects of prior knowledge on memory across the lifespan. *Developmental Psychology*, *55*(6), 1326–1337. https://doi.org/10.1037/dev0000712

Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The Human Hippocampus and Spatial and Episodic Memory. *Neuron*, *35*(4), 625–641. https://doi.org/10.1016/S0896-6273(02)00830-9

Burzynska, A. Z., Nagel, I. E., Preuschhof, C., Gluth, S., Bäckman, L., Li, S.-C., Lindenberger, U., & Heekeren, H. R. (2012). Cortical thickness is linked to executive functioning in adulthood and aging. *Human Brain Mapping*, *33*(7), 1607–1620. https://doi.org/10.1002/hbm.21311

Buss, C., Entringer, S., Swanson, J. M., & Wadhwa, P. D. (2012). The Role of Stress in Brain Development: The Gestational Environment's Long-Term Effects on the Brain. *Cerebrum: The Dana Forum on Brain Science*, *2012*, 4.

Cimadevilla, J. M., Roldán, L., París, M., Arnedo, M., & Roldán, S. (2014). Spatial learning in a virtual reality-based task is altered in very preterm children. *Journal of Clinical and Experimental Neuropsychology*, *36*(9), 1002–1008. https://doi.org/10.1080/13803395.2014.963520

Craik, F. I. M., & Lockhart, R. S. (1972). Levels of processing: A framework for memory research. *Journal of Verbal Learning and Verbal Behavior*, *11*(6), 671–684. https://doi.org/10.1016/S0022-5371(72)80001-X

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical Surface-Based Analysis. *NeuroImage*, *9*(2), 179–194. https://doi.org/10.1006/nimg.1998.0395

Dall'Oglio, A. M., Rossiello, B., Coletti, M. F., Bultrini, M., De Marchis, C., Ravà, L., Caselli, C., Paris, S., & Cuttini, M. (2010). Do healthy preterm children need neuropsychological follow-up? Preschool outcomes compared with term peers: Neuropsychological Follow-up of Healthy Preterm Children. *Developmental Medicine & Child Neurology*, *52*(10), 955–961. https://doi.org/10.1111/j.1469-8749.2010.03730.x

de Kieviet, J. F., Zoetebier, L., van ELBURG, R. M., Vermeulen, R. J., & Oosterlaan, J. (2012). Brain development of very preterm and very low-birthweight children in childhood and adolescence: A meta-analysis: Review. *Developmental Medicine & Child Neurology*, *54*(4), 313–323. https://doi.org/10.1111/j.1469-8749.2011.04216.x

Drummey, A. B., & Newcombe, N. S. (2002). Developmental changes in source memory. *Developmental Science*, *5*(4), 502–513. https://doi.org/10.1111/1467-7687.00243

Dudai, Y. (2004). The Neurobiology of Consolidations, Or, How Stable is the Engram? *Annual Review of Psychology*, *55*(1), 51–86. https://doi.org/10.1146/annurev.psych.55.090902.142050

Dudai, Y., Karni, A., & Born, J. (2015). The Consolidation and Transformation of Memory. *Neuron*, *88*(1), 20–32. https://doi.org/10.1016/j.neuron.2015.09.004

Eichenbaum, H. (2017). Prefrontal–hippocampal interactions in episodic memory. *Nature Reviews Neuroscience*, *18*(9), 547–558. https://doi.org/10.1038/nrn.2017.74

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: A robust preprocessing pipeline for functional MRI. *Nature Methods*, *16*(1), 111–116. https://doi.org/10.1038/s41592-018-0235-4

Ezzati, A., Katz, M. J., Zammit, A. R., Lipton, M. L., Zimmerman, M. E., Sliwinski, M. J., & Lipton, R. B. (2016). Differential association of left and right hippocampal volumes with verbal episodic and spatial memory in older adults. *Neuropsychologia*, *93*, 380–385. https://doi.org/10.1016/j.neuropsychologia.2016.08.016

Fernández de Gamarra-Oca, L., Zubiaurre-Elorza, L., Junqué, C., Solana, E., Soria-Pastor, S., Vázquez, É., Delgado, I., Macaya, A., Ojeda, N., & Poca, M. A. (2021). Reduced hippocampal subfield volumes and memory performance in preterm children with and without germinal matrix-intraventricular hemorrhage. *Scientific Reports*, *11*(1), 2420. https://doi.org/10.1038/s41598-021-81802-7

Field, A. P., Miles, J., & Field, Z. (2012). *Discovering statistics using R*. Sage.

Finch, W. H., & French, B. F. (2012). A Comparison of Methods for Estimating Confidence Intervals for Omega-Squared Effect Size. *Educational and Psychological Measurement*, *72*(1), 68–77. https://doi.org/10.1177/0013164411406533

Ford, R. M., Neulinger, K., O'Callaghan, M., Mohay, H., Gray, P., & Shum, D. (2011). Executive Function in 7-9-Year-Old Children Born Extremely Preterm or with Extremely Low Birth Weight: Effects of Biomedical History, Age at Assessment, and Socioeconomic Status.

Archives of Clinical Neuropsychology, 26(7), 632–644.
https://doi.org/10.1093/arclin/acr061

Frankland, P. W., & Bontempi, B. (2005). The organization of recent and remote memories. *Nature Reviews Neuroscience, 6*(2), 119–130. https://doi.org/10.1038/nrn1607

Ghetti, S., & Bunge, S. A. (2012). Neural changes underlying the development of episodic memory during middle childhood. *Developmental Cognitive Neuroscience, 2*(4), 381–395. https://doi.org/10.1016/j.dcn.2012.05.002

Gilboa, A. (2004). Remembering Our Past: Functional Neuroanatomy of Recollection of Recent and Very Remote Personal Events. *Cerebral Cortex, 14*(11), 1214–1225. https://doi.org/10.1093/cercor/bhh082

Gogtay, N., Giedd, J. N., Lusk, L., Hayashi, K. M., Greenstein, D., Vaituzis, A. C., Nugent, T. F., Herman, D. H., Clasen, L. S., Toga, A. W., Rapoport, J. L., & Thompson, P. M. (2004a). Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences, 101*(21), 8174–8179. https://doi.org/10.1073/pnas.0402680101

Gogtay, N., Giedd, J. N., Lusk, L., Hayashi, K. M., Greenstein, D., Vaituzis, A. C., Nugent, T. F., Herman, D. H., Clasen, L. S., Toga, A. W., Rapoport, J. L., & Thompson, P. M. (2004b). Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences, 101*(21), 8174–8179. https://doi.org/10.1073/pnas.0402680101

Gorgolewski, K., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., & Ghosh, S. S. (2011). Nipype: A Flexible, Lightweight and Extensible Neuroimaging Data Processing Framework in Python. *Frontiers in Neuroinformatics, 5.* https://doi.org/10.3389/fninf.2011.00013

Han, S., O'Connor, A. R., Eslick, A. N., & Dobbins, I. G. (2012). The Role of Left Ventrolateral Prefrontal Cortex during Episodic Decisions: Semantic Elaboration or Resolution of Episodic Interference? *Journal of Cognitive Neuroscience, 24*(1), 223–234. https://doi.org/10.1162/jocn_a_00133

Hayne, H. (2004). Infant memory development: Implications for childhood amnesia. *Developmental Review, 24*(1), 33–73. https://doi.org/10.1016/j.dr.2003.09.007

Haynes, W. (2013). Tukey's Test. In W. Dubitzky, O. Wolkenhauer, K.-H. Cho, & H. Yokota (Eds.), *Encyclopedia of Systems Biology* (pp. 2303–2304). Springer New York. https://doi.org/10.1007/978-1-4419-9863-7_1212

Hedman, A. M., van Haren, N. E. M., Schnack, H. G., Kahn, R. S., & Hulshoff Pol, H. E. (2012). Human brain changes across the life span: A review of 56 longitudinal magnetic resonance imaging studies. *Human Brain Mapping*, *33*(8), 1987–2002. https://doi.org/10.1002/hbm.21334

Henke, K. (2010). A model for memory systems based on processing modes rather than consciousness. *Nature Reviews Neuroscience*, *11*(7), 523–532. https://doi.org/10.1038/nrn2850

Hopf, L., Quraan, M. A., Cheung, M. J., Taylor, M. J., Ryan, J. D., & Moses, S. N. (2013). Hippocampal Lateralization and Memory in Children and Adults. *Journal of the International Neuropsychological Society*, *19*(10), 1042–1052. https://doi.org/10.1017/S1355617713000751

Karpicke, J., & Roedigeriii, H. (2007). Repeated retrieval during learning is the key to long-term retention☆. *Journal of Memory and Language*, *57*(2), 151–162. https://doi.org/10.1016/j.jml.2006.09.004

Keresztes, A., Bender, A. R., Bodammer, N. C., Lindenberger, U., Shing, Y. L., & Werkle-Bergner, M. (2017a). Hippocampal maturity promotes memory distinctiveness in childhood and adolescence. *Proceedings of the National Academy of Sciences*, *114*(34), 9212–9217. https://doi.org/10.1073/pnas.1710654114

Keresztes, A., Bender, A. R., Bodammer, N. C., Lindenberger, U., Shing, Y. L., & Werkle-Bergner, M. (2017b). Hippocampal maturity promotes memory distinctiveness in childhood and adolescence. *Proceedings of the National Academy of Sciences*, *114*(34), 9212–9217. https://doi.org/10.1073/pnas.1710654114

Kharitonova, M., Martin, R. E., Gabrieli, J. D. E., & Sheridan, M. A. (2013). Cortical gray-matter thinning is associated with age-related improvements on executive function tasks. *Developmental Cognitive Neuroscience*, *6*, 61–71. https://doi.org/10.1016/j.dcn.2013.07.002

Konkel, A. (2009). Relational memory and the hippocampus: Representations and methods. *Frontiers in Neuroscience*, *3*(2), 166–174. https://doi.org/10.3389/neuro.01.023.2009

Kuhl, B. A., Bainbridge, W. A., & Chun, M. M. (2012). Neural Reactivation Reveals Mechanisms for Updating Memory. *Journal of Neuroscience*, *32*(10), 3453–3461. https://doi.org/10.1523/JNEUROSCI.5846-11.2012

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). **lmerTest** Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13). https://doi.org/10.18637/jss.v082.i13

Lebel, C., Gee, M., Camicioli, R., Wieler, M., Martin, W., & Beaulieu, C. (2012). Diffusion tensor imaging of white matter tract evolution over the lifespan. *NeuroImage*, *60*(1), 340–352. https://doi.org/10.1016/j.neuroimage.2011.11.094

Lee, J. K., Wendelken, C., Bunge, S. A., & Ghetti, S. (2016a). A Time and Place for Everything: Developmental Differences in the Building Blocks of Episodic Memory. *Child Development*, *87*(1), 194–210. https://doi.org/10.1111/cdev.12447

Lee, J. K., Wendelken, C., Bunge, S. A., & Ghetti, S. (2016b). A Time and Place for Everything: Developmental Differences in the Building Blocks of Episodic Memory. *Child Development*, *87*(1), 194–210. https://doi.org/10.1111/cdev.12447

Lee, S., & Lee, D. K. (2018). What is the proper way to apply the multiple comparison test? *Korean Journal of Anesthesiology*, *71*(5), 353–360. https://doi.org/10.4097/kja.d.18.00242

Lenroot, R. K., & Giedd, J. N. (2006). Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience & Biobehavioral Reviews*, *30*(6), 718–729. https://doi.org/10.1016/j.neubiorev.2006.06.001

Leon, W. C., Bruno, M. A., Allard, S., Nader, K., & Cuello, A. C. (2010). Engagement of the PFC in consolidation and recall of recent spatial memory. *Learning & Memory*, *17*(6), 297–305. https://doi.org/10.1101/lm.1804410

Linberg, T., Baeumer, T., & Rossbach, H.-G. (2013). Data on Early Child Education and Care Learning Environments in Germany. *International Journal of Child Care and Education Policy*, *7*(1), 24–42. https://doi.org/10.1007/2288-6729-7-1-24

Linsell, L., Malouf, R., Morris, J., Kurinczuk, J. J., & Marlow, N. (2015). Prognostic Factors for Poor Cognitive Development in Children Born Very Preterm or With Very Low Birth Weight: A Systematic Review. *JAMA Pediatrics*, *169*(12), 1162. https://doi.org/10.1001/jamapediatrics.2015.2175

Marlow, N., Hennessy, E. M., Bracewell, M. A., Wolke, D., & for the EPICure Study Group. (2007). Motor and Executive Function at 6 Years of Age After Extremely Preterm Birth. *PEDIATRICS*, *120*(4), 793–804. https://doi.org/10.1542/peds.2007-0440

Mashburn, A. J., LoCasale-Crouch, J., & Pears, K. C. (Eds.). (2018). *Kindergarten Transition and Readiness*. Springer International Publishing. https://doi.org/10.1007/978-3-319-90200-5

Mathalon, D. H., Sullivan, E. V., Rawles, J. M., & Pfefferbaum, A. (1993). Correction for head size in brain-imaging measurements. *Psychiatry Research: Neuroimaging*, *50*(2), 121–139. https://doi.org/10.1016/0925-4927(93)90016-B

McDermott, K. B., & Zerr, C. L. (2019). Individual Differences in Learning Efficiency. *Current Directions in Psychological Science*, *28*(6), 607–613. https://doi.org/10.1177/0963721419869005

Mills, K. L., Goddings, A.-L., Herting, M. M., Meuwese, R., Blakemore, S.-J., Crone, E. A., Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., & Tamnes, C. K. (2016). Structural brain development between childhood and adulthood: Convergence across four longitudinal samples. *NeuroImage*, *141*, 273–281. https://doi.org/10.1016/j.neuroimage.2016.07.044

Morgan, K., & Hayne, H. (2011). Age-related changes in visual recognition memory during infancy and early childhood. *Developmental Psychobiology*, *53*(2), 157–165. https://doi.org/10.1002/dev.20503

Moscovitch, M., & Gilboa, A. (2021). *Systems consolidation, transformation and reorganization: Multiple Trace Theory, Trace Transformation Theory and their Competitors* [Preprint]. PsyArXiv. https://doi.org/10.31234/osf.io/yxbrs

Muftuler, L. T., Davis, E. P., Buss, C., Solodkin, A., Su, M. Y., Head, K. M., Hasso, A. N., & Sandman, C. A. (2012). Development of white matter pathways in typically developing preadolescent children. *Brain Research*, *1466*, 33–43. https://doi.org/10.1016/j.brainres.2012.05.035

Murray, A. L., Scratch, S. E., Thompson, D. K., Inder, T. E., Doyle, L. W., Anderson, J. F. I., & Anderson, P. J. (2014). Neonatal brain pathology predicts adverse attention and processing speed outcomes in very preterm and/or very low birth weight children. *Neuropsychology*, *28*(4), 552–562. https://doi.org/10.1037/neu0000071

Murre, J. M. J., & Dros, J. (2015). Replication and Analysis of Ebbinghaus' Forgetting Curve. *PLOS ONE*, *10*(7), e0120644. https://doi.org/10.1371/journal.pone.0120644

Nadel, L., & Peterson, M. A. (2013). The hippocampus: Part of an interactive posterior representational system spanning perceptual and memorial systems. *Journal of Experimental Psychology: General*, *142*(4), 1242–1254. https://doi.org/10.1037/a0033690

Nam, K. W., Castellanos, N., Simmons, A., Froudist-Walsh, S., Allin, M. P., Walshe, M., Murray, R. M., Evans, A., Muehlboeck, J.-S., & Nosarti, C. (2015). Alterations in cortical thickness development in preterm-born individuals: Implications for high-order cognitive functions. *NeuroImage*, *115*, 64–75. https://doi.org/10.1016/j.neuroimage.2015.04.015

Narberhaus, A., Lawrence, E., Allin, M. P., Walshe, M., McGuire, P., Rifkin, L., Murray, R., & Nosarti, C. (2009). Neural substrates of visual paired associates in young adults with a history of very preterm birth: Alterations in fronto-parieto-occipital networks and caudate

nucleus. *NeuroImage*, *47*(4), 1884–1893.
https://doi.org/10.1016/j.neuroimage.2009.04.036

Nolden, S., Brod, G., Meyer, A.-K., Fandakova, Y., & Shing, Y. L. (2021). Neural Correlates of Successful Memory Encoding in Kindergarten and Early Elementary School Children: Longitudinal Trends and Effects of Schooling. *Cerebral Cortex*, *31*(8), 3764–3779. https://doi.org/10.1093/cercor/bhab046

Nosarti, C., & Froudist-Walsh, S. (2016). Alterations in development of hippocampal and cortical memory mechanisms following very preterm birth. *Developmental Medicine & Child Neurology*, *58*, 35–45. https://doi.org/10.1111/dmcn.13042

Nosarti, C., Nam, K. W., Walshe, M., Murray, R. M., Cuddy, M., Rifkin, L., & Allin, M. P. G. (2014a). Preterm birth and structural brain alterations in early adulthood. *NeuroImage: Clinical*, *6*, 180–191. https://doi.org/10.1016/j.nicl.2014.08.005

Nosarti, C., Nam, K. W., Walshe, M., Murray, R. M., Cuddy, M., Rifkin, L., & Allin, M. P. G. (2014b). Preterm birth and structural brain alterations in early adulthood. *NeuroImage: Clinical*, *6*, 180–191. https://doi.org/10.1016/j.nicl.2014.08.005

Ofen, N., Kao, Y.-C., Sokol-Hessner, P., Kim, H., Whitfield-Gabrieli, S., & Gabrieli, J. D. E. (2007). Development of the declarative memory system in the human brain. *Nature Neuroscience*, *10*(9), 1198–1205. https://doi.org/10.1038/nn1950

Okada, K. (2013). Is Omega Squared Less Biased? A Comparison of Three Major Effect Size Indices in One-Way Anova. *Behaviormetrika*, *40*(2), 129–147. https://doi.org/10.2333/bhmk.40.129

Omizzolo, C., Scratch, S. E., Stargatt, R., Kidokoro, H., Thompson, D. K., Lee, K. J., Cheong, J., Neil, J., Inder, T. E., Doyle, L. W., & Anderson, P. J. (2014a). Neonatal brain abnormalities and memory and learning outcomes at 7 years in children born very preterm. *Memory*, *22*(6), 605–615. https://doi.org/10.1080/09658211.2013.809765

Omizzolo, C., Scratch, S. E., Stargatt, R., Kidokoro, H., Thompson, D. K., Lee, K. J., Cheong, J., Neil, J., Inder, T. E., Doyle, L. W., & Anderson, P. J. (2014b). Neonatal brain abnormalities and memory and learning outcomes at 7 years in children born very preterm. *Memory*, *22*(6), 605–615. https://doi.org/10.1080/09658211.2013.809765

Østby, Y., Tamnes, C. K., Fjell, A. M., & Walhovd, K. B. (2012a). Dissociating Memory Processes in the Developing Brain: The Role of Hippocampal Volume and Cortical Thickness in Recall after Minutes versus Days. *Cerebral Cortex*, *22*(2), 381–390. https://doi.org/10.1093/cercor/bhr116

Østby, Y., Tamnes, C. K., Fjell, A. M., & Walhovd, K. B. (2012b). Dissociating Memory
Processes in the Developing Brain: The Role of Hippocampal Volume and Cortical
Thickness in Recall after Minutes versus Days. *Cerebral Cortex*, *22*(2), 381–390.
https://doi.org/10.1093/cercor/bhr116

Peiffer, A., Brichet, M., De Tiège, X., Peigneux, P., & Urbain, C. (2020a). The power of
children's sleep—Improved declarative memory consolidation in children compared with
adults. *Scientific Reports*, *10*(1), 9979. https://doi.org/10.1038/s41598-020-66880-3

Peiffer, A., Brichet, M., De Tiège, X., Peigneux, P., & Urbain, C. (2020b). The power of
children's sleep—Improved declarative memory consolidation in children compared with
adults. *Scientific Reports*, *10*(1), 9979. https://doi.org/10.1038/s41598-020-66880-3

Perner, J., & Ruffman, T. (1995). Episodic Memory and Autonoetic Conciousness:
Developmental Evidence and a Theory of Childhood Amnesia. *Journal of Experimental
Child Psychology*, *59*(3), 516–548. https://doi.org/10.1006/jecp.1995.1024

Peterson, C. (2002). Children's long-term memory for autobiographical events. *Developmental
Review*, *22*(3), 370–402. https://doi.org/10.1016/S0273-2297(02)00007-2

Petrides, M. (2007). The Orbitofrontal Cortex: Novelty, Deviation from Expectation, and
Memory. *Annals of the New York Academy of Sciences*, *1121*(1), 33–53.
https://doi.org/10.1196/annals.1401.035

Quesada, A. A., Tristão, R. M., Pratesi, R., & Wolf, O. T. (2014). Hyper-responsiveness to acute
stress, emotional problems and poorer memory in former preterm children. *Stress*, *17*(5),
389–399. https://doi.org/10.3109/10253890.2014.949667

Raposo, A., Han, S., & Dobbins, I. G. (2009). Ventrolateral prefrontal cortex and self-initiated
semantic elaboration during memory retrieval. *Neuropsychologia*, *47*(11), 2261–2271.
https://doi.org/10.1016/j.neuropsychologia.2008.10.024

Riggins, T. (2014). Longitudinal investigation of source memory reveals different developmental
trajectories for item memory and binding. *Developmental Psychology*, *50*(2), 449–459.
https://doi.org/10.1037/a0033622

Roberts, G., Lim, J., Doyle, L. W., & Anderson, P. J. (2011). High Rates of School Readiness
Difficulties at 5 Years of Age in Very Preterm Infants Compared with Term Controls.
Journal of Developmental & Behavioral Pediatrics, *32*(2), 117–124.
https://doi.org/10.1097/DBP.0b013e318206d5c9

Rose, S. A., Feldman, J. F., Jankowski, J. J., & Van Rossem, R. (2011). Basic information
processing abilities at 11 years account for deficits in IQ associated with preterm birth.
Intelligence, *39*(4), 198–209. https://doi.org/10.1016/j.intell.2011.03.003

Salvan, P., Froudist Walsh, S., Allin, M. P. G., Walshe, M., Murray, R. M., Bhattacharyya, S., McGuire, P. K., Williams, S. C. R., & Nosarti, C. (2014). Road work on memory lane—Functional and structural alterations to the learning and memory circuit in adults born very preterm. *NeuroImage*, *102*, 152–161. https://doi.org/10.1016/j.neuroimage.2013.12.031

Sanfilipo, M. P., Benedict, R. H. B., Zivadinov, R., & Bakshi, R. (2004). Correction for intracranial volume in analysis of whole brain atrophy in multiple sclerosis: The proportion vs. residual method. *NeuroImage, 22*(4), 1732–1743. https://doi.org/10.1016/j.neuroimage.2004.03.037

Savage, C. R., Deckersbach, T., Heckers, S., Wagner, A. D., Schacter, D. L., Alpert, N. M., Fischman, A. J., & Rauch, S. L. (2001). Prefrontal regions supporting spontaneous and directed application of verbal learning strategies: Evidence from PET. *Brain*, *124*(1), 219–231. https://doi.org/10.1093/brain/124.1.219

Scarf, D., Gross, J., Colombo, M., & Hayne, H. (2013). To have and to hold: Episodic memory in 3- and 4-year-old children. *Developmental Psychobiology*, *55*(2), 125–132. https://doi.org/10.1002/dev.21004

Schermann, L., & Sedin, G. (2007). Cognitive function at 10 years of age in children who have required neonatal intensive care. *Acta Paediatrica*, *93*(12), 1619–1629. https://doi.org/10.1111/j.1651-2227.2004.tb00853.x

Schlichting, M. L., & Preston, A. R. (2015). Memory integration: Neural mechanisms and implications for behavior. *Current Opinion in Behavioral Sciences*, *1*, 1–8. https://doi.org/10.1016/j.cobeha.2014.07.005

Shing, Y. L., Werkle-Bergner, M., Brehmer, Y., Müller, V., Li, S.-C., & Lindenberger, U. (2010a). Episodic memory across the lifespan: The contributions of associative and strategic components. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1080–1091. https://doi.org/10.1016/j.neubiorev.2009.11.002

Shing, Y. L., Werkle-Bergner, M., Brehmer, Y., Müller, V., Li, S.-C., & Lindenberger, U. (2010b). Episodic memory across the lifespan: The contributions of associative and strategic components. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1080–1091. https://doi.org/10.1016/j.neubiorev.2009.11.002

Shing, Y. L., Werkle-Bergner, M., Brehmer, Y., Müller, V., Li, S.-C., & Lindenberger, U. (2010c). Episodic memory across the lifespan: The contributions of associative and strategic components. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1080–1091. https://doi.org/10.1016/j.neubiorev.2009.11.002

Shing, Y. L., Werkle-Bergner, M., Li, S.-C., & Lindenberger, U. (2008a). Associative and strategic components of episodic memory: A life-span dissociation. *Journal of Experimental Psychology. General*, *137*(3), 495–513. https://doi.org/10.1037/0096-3445.137.3.495

Shing, Y. L., Werkle-Bergner, M., Li, S.-C., & Lindenberger, U. (2008b). Associative and strategic components of episodic memory: A life-span dissociation. *Journal of Experimental Psychology. General*, *137*(3), 495–513. https://doi.org/10.1037/0096-3445.137.3.495

Simons, J. S., & Spiers, H. J. (2003). Prefrontal and medial temporal lobe interactions in long-term memory. *Nature Reviews Neuroscience*, *4*(8), 637–648. https://doi.org/10.1038/nrn1178

Sluzenski, J., Newcombe, N. S., & Kovacs, S. L. (2006). Binding, relational memory, and recall of naturalistic events: A developmental perspective. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *32*(1), 89–100. https://doi.org/10.1037/0278-7393.32.1.89

Sousa, S. S., Amaro, E., Crego, A., Gonçalves, Ó. F., & Sampaio, A. (2018). Developmental trajectory of the prefrontal cortex: A systematic review of diffusion tensor imaging studies. *Brain Imaging and Behavior*, *12*(4), 1197–1210. https://doi.org/10.1007/s11682-017-9761-4

Sowell, E. R., Delis, D., Stiles, J., & Jernigan, T. L. (2001). Improved memory functioning and frontal lobe maturation between childhood and adolescence: A structural MRI study. *Journal of the International Neuropsychological Society*, *7*(3), 312–322. https://doi.org/10.1017/S135561770173305X

Squire, L. R., & Alvarez, P. (1995). Retrograde amnesia and memory consolidation: A neurobiological perspective. *Current Opinion in Neurobiology*, *5*(2), 169–177. https://doi.org/10.1016/0959-4388(95)80023-9

Squire, L. R., Genzel, L., Wixted, J. T., & Morris, R. G. (2015a). Memory Consolidation. *Cold Spring Harbor Perspectives in Biology*, *7*(8), a021766. https://doi.org/10.1101/cshperspect.a021766

Squire, L. R., Genzel, L., Wixted, J. T., & Morris, R. G. (2015b). Memory Consolidation. *Cold Spring Harbor Perspectives in Biology*, *7*(8), a021766. https://doi.org/10.1101/cshperspect.a021766

Stålnacke, S. R., Tessma, M., Böhm, B., & Herlenius, E. (2019). Cognitive Development Trajectories in Preterm Children With Very Low Birth Weight Longitudinally Followed

Until 11 Years of Age. *Frontiers in Physiology*, *10*, 307.
https://doi.org/10.3389/fphys.2019.00307

Steinvorth, S., Levine, B., & Corkin, S. (2005). Medial temporal lobe structures are needed to re-experience remote autobiographical memories: Evidence from H.M. and W.R. *Neuropsychologia*, *43*(4), 479–496.
https://doi.org/10.1016/j.neuropsychologia.2005.01.001

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2020). *partR2: Partitioning R ^{2} in generalized linear mixed models* [Preprint]. Bioinformatics.
https://doi.org/10.1101/2020.07.26.221168

Takashima, A., Petersson, K. M., Rutters, F., Tendolkar, I., Jensen, O., Zwarts, M. J., McNaughton, B. L., & Fernandez, G. (2006). Declarative memory consolidation in humans: A prospective functional magnetic resonance imaging study. *Proceedings of the National Academy of Sciences*, *103*(3), 756–761.
https://doi.org/10.1073/pnas.0507774103

Thompson, D. K., Adamson, C., Roberts, G., Faggian, N., Wood, S. J., Warfield, S. K., Doyle, L. W., Anderson, P. J., Egan, G. F., & Inder, T. E. (2013a). Hippocampal shape variations at term equivalent age in very preterm infants compared with term controls: Perinatal predictors and functional significance at age 7. *NeuroImage*, *70*, 278–287.
https://doi.org/10.1016/j.neuroimage.2012.12.053

Thompson, D. K., Adamson, C., Roberts, G., Faggian, N., Wood, S. J., Warfield, S. K., Doyle, L. W., Anderson, P. J., Egan, G. F., & Inder, T. E. (2013b). Hippocampal shape variations at term equivalent age in very preterm infants compared with term controls: Perinatal predictors and functional significance at age 7. *NeuroImage*, *70*, 278–287.
https://doi.org/10.1016/j.neuroimage.2012.12.053

Thompson, D. K., Omizzolo, C., Adamson, C., Lee, K. J., Stargatt, R., Egan, G. F., Doyle, L. W., Inder, T. E., & Anderson, P. J. (2014). Longitudinal growth and morphology of the hippocampus through childhood: Impact of prematurity and implications for memory and learning. *Human Brain Mapping*, *35*(8), 4129–4139. https://doi.org/10.1002/hbm.22464

Tompary, A., & Davachi, L. (2017). Consolidation Promotes the Emergence of Representational Overlap in the Hippocampus and Medial Prefrontal Cortex. *Neuron*, *96*(1), 228-241.e5.
https://doi.org/10.1016/j.neuron.2017.09.005

Troncoso Skidmore, S., & Thompson, B. (2013). Bias and precision of some classical ANOVA effect sizes when assumptions are violated. *Behavior Research Methods*, *45*(2), 536–546. https://doi.org/10.3758/s13428-012-0257-2

Tse, D., Langston, R. F., Kakeyama, M., Bethus, I., Spooner, P. A., Wood, E. R., Witter, M. P., & Morris, R. G. M. (2007). Schemas and Memory Consolidation. *Science*, *316*(5821), 76–82. https://doi.org/10.1126/science.1135935

Tse, D., Takeuchi, T., Kakeyama, M., Kajii, Y., Okuno, H., Tohyama, C., Bito, H., & Morris, R. G. M. (2011). Schema-Dependent Gene Activation and Memory Encoding in Neocortex. *Science*, *333*(6044), 891–895. https://doi.org/10.1126/science.1205274

Tseng, C.-E. J., Froudist-Walsh, S., Brittain, P. J., Karolis, V., Caldinelli, C., Kroll, J., Counsell, S. J., Williams, S. C. R., Murray, R. M., & Nosarti, C. (2017). A multimodal imaging study of recognition memory in very preterm born adults: Imaging Recognition Memory in Very Preterm Born Adults. *Human Brain Mapping*, *38*(2), 644–655. https://doi.org/10.1002/hbm.23405

Tulving, E. (2002). Episodic Memory: From Mind to Brain. *Annual Review of Psychology*, *53*(1), 1–25. https://doi.org/10.1146/annurev.psych.53.100901.135114

Tulving, E., & Markowitsch, H. J. (1998). Episodic and declarative memory: Role of the hippocampus. *Hippocampus*, *8*(3), 198–204. https://doi.org/10.1002/(SICI)1098-1063(1998)8:3<198::AID-HIPO2>3.0.CO;2-G

Tustin, K., & Hayne, H. (2010). Defining the boundary: Age-related changes in childhood amnesia. *Developmental Psychology*, *46*(5), 1049–1061. https://doi.org/10.1037/a0020105

Uda, S., Matsui, M., Tanaka, C., Uematsu, A., Miura, K., Kawana, I., & Noguchi, K. (2015). Normal development of human brain white matter from infancy to early adulthood: A diffusion tensor imaging study. *Developmental Neuroscience*, *37*(2), 182–194. https://doi.org/10.1159/000373885

Uematsu, A., Matsui, M., Tanaka, C., Takahashi, T., Noguchi, K., Suzuki, M., & Nishijo, H. (2012). Developmental Trajectories of Amygdala and Hippocampus from Infancy to Early Adulthood in Healthy Individuals. *PLoS ONE*, *7*(10), e46970. https://doi.org/10.1371/journal.pone.0046970

van Kesteren, M. T. R., Ruiter, D. J., Fernández, G., & Henson, R. N. (2012). How schema and novelty augment memory formation. *Trends in Neurosciences*, *35*(4), 211–219. https://doi.org/10.1016/j.tins.2012.02.001

Voevodskaya, O., Simmons, A., Nordenskjöld, R., Kullberg, J., Ahlström, H., Lind, L., Wahlund, L.-O., Larsson, E.-M., Westman, E., & Alzheimer's Disease Neuroimaging Initiative. (2014). The effects of intracranial volume adjustment approaches on multiple

regional MRI volumes in healthy aging and Alzheimer's disease. *Frontiers in Aging Neuroscience*, *6*. https://doi.org/10.3389/fnagi.2014.00264

Waber, D. P., De Moor, C., Forbes, P. W., Almli, C. R., Botteron, K. N., Leonard, G., Milovan, D., Paus, T., Rumsey, J., & THE BRAIN DEVELOPMENT COOPERATIVE GROUP. (2007). The NIH MRI study of normal brain development: Performance of a population based sample of healthy children aged 6 to 18 years on a neuropsychological battery. *Journal of the International Neuropsychological Society*, *13*(05). https://doi.org/10.1017/S1355617707070841

Walani, S. R. (2020). Global burden of preterm birth. *International Journal of Gynecology & Obstetrics*, *150*(1), 31–33. https://doi.org/10.1002/ijgo.13195

Wang, J.-Y., Weber, F. D., Zinke, K., Inostroza, M., & Born, J. (2018). More Effective Consolidation of Episodic Long-Term Memory in Children Than Adults-Unrelated to Sleep. *Child Development*, *89*(5), 1720–1734. https://doi.org/10.1111/cdev.12839

Wilhelm, I., Diekelmann, S., & Born, J. (2008). Sleep in children improves memory performance on declarative but not procedural tasks. *Learning & Memory*, *15*(5), 373–377. https://doi.org/10.1101/lm.803708

Yamashita, K. -i., Hirose, S., Kunimatsu, A., Aoki, S., Chikazoe, J., Jimura, K., Masutani, Y., Abe, O., Ohtomo, K., Miyashita, Y., & Konishi, S. (2009). Formation of Long-Term Memory Representation in Human Temporal Cortex Related to Pictorial Paired Associates. *Journal of Neuroscience*, *29*(33), 10335–10340. https://doi.org/10.1523/JNEUROSCI.1328-09.2009

Zerr, C. L., Berg, J. J., Nelson, S. M., Fishell, A. K., Savalia, N. K., & McDermott, K. B. (2018). Learning Efficiency: Identifying Individual Differences in Learning Rate and Retention in Healthy Adults. *Psychological Science*, *29*(9), 1436–1450. https://doi.org/10.1177/0956797618772540

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical Imaging*, *20*(1), 45–57. https://doi.org/10.1109/42.906424

Kleiner, M., Brainard, D., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's new in psychtoolbox-3. Perception, 36(14), 1-16.

„Kaufman Assessment Battery for Children — Second Edition", Autoren: Alan S. Kaufman und Nadeen L. Kaufman; deutsche Adaptation: Dr. Peter Melchers und Martin Melchers © 2004 NCS Pearson, Inc.; deutsche Fassung © 2015 NCS Pearson, Inc.

"Wechsler Adult Intelligence Scale – Fourth Edition", Authors: D. Wechsler; deutsche
Adaptation: F. Petermann © 2004 NCS Pearson, Inc.; deutsche Fassung © 2015 NCS
Pearson, Inc.