Linking ecological specialization to its macroevolutionary consequences:

An example with passerine nest site

Rosana Zenil-Ferguson[1]*

Jay P. McEntee[2]

J. Gordon Burleigh[3]

Renée A. Duckworth[4]

[1] *School of Life Sciences, University of Hawaiʻi Mānoa, 1800 East-West Rd, Honolulu, HI 96822;*

[2] *Department of Biology, Missouri State University, 901 S. National Ave., Springfield, MO 65897 USA;*

[3] *Department of Biology, University of Florida, 220 Bartram Hall, P.O. Box 118525, Gainesville, FL 32611;*

[4] *Department of Ecology and Evolutionary Biology, University of Arizona, 1041 E. Lowell St., Tucson, AZ 85721 USA;*

* *E-mail: roszenil@hawaii.edu*

Abstract. – A long-standing hypothesis in evolutionary biology is that the evolution of

resource specialization can lead to an evolutionary dead end, where specialists have

low diversification rates and limited ability to evolve into generalists. In recent years,

advances in comparative methods investigating trait-based differences associated with

diversification have enabled more robust tests of this idea and have found mixed

support. We test the evolutionary dead end hypothesis by estimating net diversification

rate differences associated with nest site specialization among 3,224 species of

passerine birds. In particular, we test whether the adoption of hole-nesting, a nest site

specialization that decreases predation, results in reduced diversification rates relative

to nesting outside of holes. Further, we examine whether evolutionary transitions to the

specialist hole-nesting state have been more frequent than transitions out of hole-

nesting. Using diversification models that accounted for background rate heterogeneity

and different extinction rate scenarios, we found that hole-nesting specialization was not

associated with diversification rate differences. Furthermore, contrary to the assumption

that specialists rarely evolve into generalists, we found that transitions out of hole-

nesting occur more frequently than transitions into hole-nesting. These results suggest

that interspecific competition may limit adoption of hole-nesting, but that such

competition does not result in limited diversification of hole-nesters. In conjunction with

other recent studies using robust comparative methods, our results add to growing

evidence that evolutionary dead ends are not a typical outcome of resource

specialization.

Keywords: diversification, hidden-state models, cavity nesting, passerines, resource

specialization

Resource specialization, where species use a narrower range of resources compared to

related taxa (Futuyma and Moreno 1988), is a common phenomenon in evolution.

There are many reasons resource specialization (hereafter specialization) can evolve.

For example, specialists can benefit from reduced competition, or from the avoidance of

predators or parasites (Futuyma and Moreno 1988; Bernays 1989; Schluter 2000).

Once evolved, specialization can have diverse consequences for macroevolutionary

dynamics. Historically, specialization often has been considered an evolutionary dead

end, possibly resulting in both reduced net diversification relative to generalists, and

difficulty evolving from specialist to generalist. This difficulty is thought to arise when

adaptation to specialized resources occurs across numerous traits (Futuyma and

Moreno 1988), and, when coupled with changes in resource availability, leads to

elevated extinction rates in specialists. Additionally, reduced diversification of specialists

could result if the number of available niches in the specialized state is low.

Consistent with the evolutionary dead end view of specialists, the prevailing

historical viewpoint was that specialists evolve from generalist ancestors (Schluter

2000). Prior studies have found mixed support for this viewpoint, with inferences from

phylogenetic comparative methods showing that generalists regularly evolve from

specialists in many clades (Day et al. 2016; Sexton et al. 2017; Villastrigo et al. 2021).

These results indicate that the dominant direction of evolutionary transitions involving

specialization should be treated as an open question for any specialization scenario.

More recent studies have also noted the potential for specialization to increase

diversification through multiple mechanisms. First, the release of specialists from the

effects of competition and/or predation could trigger periods of niche-filling

diversification (Schluter 2000), leading to higher rates of speciation in specialists

compared to generalists. Further, in circumstances where specialization releases

specialists from competition or predation (Futuyma and Moreno 1988), specialists may

experience greater population persistence than generalists, which could increase

diversification rates by decreasing extinction rates or by increasing rates of allopatric

speciation via the longer survival of incipient lineages (Harvey et al. 2019), or both.

Finally, specialist lineages could also have higher diversification rates if more

specialized lineages have more fragmented distributions or lower rates of dispersal

(Gavrilets et al. 2000; Birand et al. 2012).

Recent diversification models and statistical developments have the potential to

alter our conclusions on the macroevolutionary consequences of specialization. A

growing awareness of type I error (Davis et al. 2013; Rabosky and Goldberg 2015), the

importance of large sample sizes for statistical power (Davis et al. 2013; Day et al.

2016), and the misspecification of the null hypothesis for the original state-dependent

diversification models (Beaulieu and O'Meara 2016; Rabosky and Goldberg 2017;

Caetano et al. 2018) have produced a wave of new studies using comprehensive

statistical approaches on the question of whether specialization is linked to

diversification. The majority of studies after Day et al.'s (2016) review on specialization

have found either no association between specialization and diversification (e.g.,

Alhajeri and Steppan 2018; Crouch and Ricklefs 2019; Villastrigo et al. 2020) or have

found that specialization is associated with higher diversification rates (e.g., Conway

and Olsen 2019; Otero et al. 2019; Tonini et al. 2020), with few studies indicating

specialization leads to an evolutionary dead end (e.g., Cyriac and Kodandaramaiah

2018; Day et al. 2016). While these newer studies have relatively large sample sizes

(compared to pre-2016 studies) and more transitions to specialist states, some of the

sample sizes are still too small to fit state-dependent diversification models with more

than two states (Davis et al. 2013) power to estimate specialization's consequences for

diversification (Davis et al. 2013). This has also been difficult to accomplish because, for

the majority of the software that fits state-dependent diversification models, estimating

intervals requires complex likelihood function maximizations (Zenil-Ferguson et al.

2018) or many bootstrap simulations to approximate confidence-likelihood intervals for

diversification parameters (FitzJohn 2012; Beaulieu et al. 2021). Moving forward it is

necessary to integrate models that incorporate more states and more parameters that

mathematically link multiple specialist and generalist states to the process of

diversification while accounting for the potential for heterogeneity in the diversification

process (Beaulieu and O'Meara 2016). Furthermore, the calculation of uncertainty for

the parameter estimates in more complex models is key to understanding the power of

the sample and testing whether specialization is linked to diversification.

Here, we highlight these methodological approaches with a large dataset on

nesting microhabitats for 3,224 species of passerine birds. Passerines provide a unique

opportunity to examine the macroevolutionary consequences of specialization.

Specifically, hole-nesting is a prime example of resource specialization which has

evolved multiple times across the clade of 6000 passerine taxa, providing the power

needed to estimate rates of diversification linked to trait evolution. Second, passerines

can be grouped into three general nesting habits: open-cup nesters, dome nesters and

hole nesters (e.g., Wallace 1868; Martin 1995; Collias 1997). Open-cup and dome

nesters are relatively unspecialized in their nesting substrate compared to hole-nesters;

however, similar to hole nesters, dome nesters have reduced predation rates compared

to open-cup nesters (Oniki 1979a; Linder and Bollinger 1995; Auer et al. 2007; Martin et

al. 2017). Therefore, our approach of comparing adoption of these three nest types

enables us to simultaneously assess the potential effects of nest site specialization and

escape from predation on diversification rates (Fig. 1).

Figure 1. The nest types of passerine taxa differ by how specialized they are in nest substrate

and in their predation risk. Hole is the most specialized nest type and has the smallest predation

risk, followed by dome with also a lower risk but relatively unspecialized in its nesting substrate.

Finally, open-cups have the highest predation risk but are also considered unspecialized. Nest

colors represent the colors of the states for the models presented.

We employ a graphical modeling approach (Jordan 2004) in Bayesian statistics

to infer point estimates and credible intervals of parameters from state-dependent

speciation and extinction models (SSE models), and diversification-free models (Mkn-

Markov models without state-specific diversification parameters) coded in RevBayes

software (Höhna et al. 2016). The large nest site dataset and phylogenetic tree are the

input to models that are estimated in a Bayesian framework and this enables us to avoid

a number of the common pitfalls discussed above. Additionally, we adopt a systematic

approach to specifying prior distributions to model enhanced extinction rates to assess

the consistency of the statistical inference across a spectrum of macroevolutionary

diversification scenarios. Finally, after determining the best model for the evolution of

nest site, we used it to reconstruct ancestral states of nest types to visualize the pattern

of evolutionary history for the trait and compare with previous ancestral state

reconstructions.

140   *Nest site evolution and its potential links to diversification*

Nest site selection is a critically important aspect of avian habitat because nest failure

rates are high in birds (Nice 1957) with predation as the main cause of nestling mortality

(Ricklefs 1969; Martin 1993). Adoption of hole-nesting has the advantage of protecting

offspring from predation, as non-excavating hole nesters show an approximate 43%

reduction in nest failure compared to non-hole nesters (Martin 1995). Thus, a major

ecological consequence of hole-nesting specialization is that it increases nesting

success by providing a release from predation. Hole-nesting represents resource

specialization for two reasons: the great majority of species that nest in holes do so

obligately, and exclusive use of holes greatly restricts the substrates that are suitable for

nest building.

Hole-nesting is associated with the evolution of a suite of traits, such as

increased nesting period, brighter egg coloration, and larger clutch sizes, that may

make it difficult to transition out of hole-nesting (Martin and Li 1992; Kilner 2006).

Moreover, the great majority of hole-nesting passerines cannot excavate their own nest

holes, and competition for this limited resource is intense (Newton 1994). While hole-

nesting likely evolves as a response to predation pressure, the intense competition for

nesting sites decreases nest hole availability (Cockle et al. 2011). This competition may

limit ecological opportunity for hole-nesting lineages despite the reduction in predation

pressure. Further, reliance on holes for nesting may increase extinction risk when

environmental changes reduce nest hole availability, particularly if transitioning from

hole-nesting is difficult. This combination of characteristics makes hole-nesting in

passerines a form of specialization that could be an evolutionary dead end. However, as

hole-nesting decreases nest predation, it is also possible that adopting hole-nesting

could lead to greater population persistence over time which, in turn, could lead to

reduced extinction rates or increased rates of allopatric speciation (Harvey et al. 2019).

Thus, it is also possible that hole-nesting lineages could experience greater

diversification rates than the more generalist non hole-nesting lineages.

The evolutionary dead end hypothesis assumes that it is easy to adopt hole-

nesting habits, but difficult to transition out of them because of secondary adaptations

that restrict evolutionary transitions (Collias and Collias 1984). In passerines, most hole

nesters are secondary (i.e., they cannot make new cavities, and thus they rely on what

is already in the environment), and so adoption of hole-nesting is mainly a behavioral

shift not requiring extensive morphological modification (although see above for

examples of life history traits associated with hole-nesting). This could make it relatively

easy to evolve hole-nesting behavior. However, the intense competitive environment

that hole nesters face could mean that hole-nesting niches are already saturated,

making it a more difficult habit to adopt. To examine these possibilities, we not only

assess the impact of nest type on diversification rates, but we also calculate transition

rates between states.

181   MATERIALS AND METHODS

182   *Data*

Our nesting data is a slightly adapted version of a dataset originally assembled

for McEntee et al. (2021). This dataset was generated by scoring nesting behavior of

passerine species using descriptions from the Handbook of the Birds of the World Alive,

(Del Hoyo et al. 2017, last accessed 30 June 2016, hereafter HBW). Specifically,

species' nests were scored as either open cup, dome, or hole. A few species show

flexibility in nest type use among these three types, and are capable of nesting in, for

example, either an open cup or a hole. We scored these examples accordingly as

nesting in either. "Hole" refers to any nest built inside a tree cavity, rock crevice, or

earthen bank. The small number of passerine brood parasites were excluded from

analyses. (McEntee et al. 2021) were able to score the nest types for approximately

three quarters of the 5,912 passerine species from the information in the HBW species

descriptions.

Next, we matched the passerine species in the nesting dataset to the species'

names on the tips of the avian supermatrix phylogeny of Burleigh et al. (2015), one of

the largest phylogenetic trees of birds constructed exclusively from molecular data. This

tree was time-calibrated using penalized likelihood in r8s (Sanderson 2003) with twenty

fossil calibrations from throughout the avian phylogeny (Baiser et al. 2018). We

identified all cases in which a species in the Burleigh et al. (2015) phylogenetic tree,

which used the Clements taxonomy, did not have a corresponding species with the

exact same name in the dataset from the HBW Alive/BirdLife International taxonomy.

This was due to either missing nesting data or a taxonomic mismatch. For taxonomic

mismatches, we examined the taxonomic history for these species in Avibase

(https://avibase.bsc-eoc.org/), and when appropriate, changed the species name in the

nest dataset to match the phylogenetic tree (Burleigh et al. 2015). Taxa treated as

subspecies in the HBW taxonomy (2015) but as taxa in (Burleigh et al. 2015) were not

included in our analyses because they did not occur as tips in the phylogenetic tree. We

then trimmed the phylogenetic tree to include only species in the nesting dataset from

HBW. The resulting data set has the nesting state for 3,224 passerine species. Of

these, 1943 species had dome nests (D), 722 had cup nests (C), and 458 had hole

nests (H). Among species with multiple nest types, 60 were cup or hole nesters (CH),

29 were dome or cup nesters (DC), and 13 were dome or hole nesters (DH). In the few

instances (less than 1% of internal nodes) where this tree was not bifurcating (required

for the diversification analyses described below) because nodes were collapsed in the

r8s analysis, we resolved bifurcations randomly using the function `multi2di` from the

R package `ape` (Paradis and Schliep 2019).

218   *Modeling nest site linked to diversification*

Using the three states assigned to the tips of the 3,225-taxon phylogenetic tree

(3,224 passerines plus a single species representing the parrot outgroup), we defined

four state-dependent speciation and extinction models (SSEs) and one Markov model

that we call diversification-free (because it does not contain speciation and extinction

parameters Fig. 2). The first multistate SSE model, called MuSSE-3 (Fig. 2a), uses

three main states: dome D, cup C, and hole H (Fig. 2a), each with their own speciation

and extinction rate. MuSSE-3 also has transition rates between each state (q_{ij} with

$i, j = D, C, H$, and $i \neq j$) governing the rate of evolution from one nest type to another.

All transitions between nest types are possible. Taxa that have multiple nest types are

coded in the input data as belonging to multiple states simultaneously. For example, a

bird taxon reported to have nested in both hole and dome nests is coded in the input

data as (H, D). For taxa (tips) with two nest types, the likelihood calculation includes the

product of two transition probabilities from the most recent common ancestor to each

potential state in the tips instead of a single transition probability from the most recent

common ancestor to one single state.

The second model (Fig. 2b) is the hidden state extension of the MuSSE-3 and it

is called multi and hidden state dependent speciation and extinction model (MuHiSSE-

3). The hidden state extension includes background heterogeneity in the diversification

rate that is not linked to the trait of interest (Beaulieu and O'Meara 2016), and (Caetano

et al. 2018). The MuHiSSE-3 model expanded from three single nest type states to six

with subscripts A or B (Fig. 2b).

The character independent model (CID-3) or null hypothesis model (Fig. 2c)

accommodates heterogeneity in the process of diversification completely unlinked from

nest type but due to some unmeasured factors called hidden states. CID-3 specifies a

null hypothesis that is comparable to MuSSE-3 (Beaulieu and O'Meara 2016; Caetano

et al. 2018) because both models have the same number of diversification rates. That

is, the CID-3 has six speciation and extinction parameters, but makes the net

diversification for main nest type states D, C, H equal while assuming that the source of

differences in diversification come from the hidden states A, B, C. In mathematical terms

the null hypothesis (H_0 or CID-3 model) for a state-dependent diversification model with

three states and six diversification rate parameters is:

250            $H_0: \lambda_{Dk} = \lambda_{Ck} = \lambda_{Hk} = \lambda_k$ for $k = A, B, C$, but $\lambda_A \neq \lambda_B \neq \lambda_C$;

251   and $\mu_{Dk} = \mu_{Ck} = \mu_{Hk} = \mu_k$ for $k = A, B, C$, but $\mu_A \neq \mu_B \neq \mu_C$.

Note that in the CID-3 notation the 3 refers to the number of hidden states that have

different diversification rates, and not necessarily the number of states of the main trait

as discussed in Caetano et al. (2018).

For all SSE models in Fig. 2 we defined a prior distribution for the log-

transformed speciation rates λ_i to be a normal distribution with hyperparameters

$\left(\mu_\lambda = ln \left(ln \left(\frac{ln(3225/2)}{Tree\ Age} \right) \right), \sigma_\lambda = 1.12 \right)$ with the passerine tree age as 64.51 million years

old. This choice of hyperparameter μ_λ for the log-transformed speciation rate makes the

expected average rate for speciation approximately $e^{\mu_\lambda} = \frac{ln(3225/2)}{Tree\ Age} = 0.11$, representing

an approximation to the current observed extant diversity of 3225 taxa in the tree

conditional on having two extant lineages at the root of a phylogeny that is 64.5 million

years old (FitzJohn et al. 2009; Freyman and Höhna 2019; Zenil-Ferguson et al. 2019).

The same prior distributions were used for log-transformed extinction rates. For all

models we assume a sampling fraction $\rho = 0.51$ since our tree contains about 51% of

all passerine taxa.

Figure 2. Models of nest type evolution. a) The multistate dependent speciation and extinction

(MuSSE-3) assumes transitions among each nest type (D, C, H) and diversification rates

(speciation and extinction) associated with each state. b) The multi- and hidden state dependent

speciation and extinction model (MuHiSSE-3) expands the model to consider hidden states A and

B (and transitions between them) to model the potential of other factors along with nest type in

the process of diversification. c) The character independent model (CID-3) allows us to assess

whether diversification is linked to some unmeasured factors that are independent from nest type.

d) MuSSE-3 with prior distributions that restrict the parametric space allows us to infer trait-

associated diversification differences where relative extinction is enhanced, a part of parameter

space that is typically poorly explored in SSE models. e) MuHiSSE-3 with prior distributions that

restrict the parametric space allows us to infer trait- associated diversification differences where

relative extinction is enhanced. f) The Markov model (Mk3) assumes that there are no

diversification rate differences among nest types, and transitions among nest types (D, C, H) are

allowed.

In the supplementary information we fit state-dependent diversification models

with six states including taxa with two types of nesting as a separate state (i.e., cup and

dome nesters get their own state DC) to investigate if different state-coding changes

diversification and/or transition rates (Fig. S5).

287   *Testing the role of diversification in nest type evolution*

To test if nest type evolution is linked to the diversification of passerines, we

calculated the marginal log-likelihoods for the MuSSE-3 and the character independent

model (CID-3). Using the log-likelihoods we calculated Bayes factors (Kass and Raftery

1995) to compare CID-3 against the MuSSE-3 model. To approximate the marginal log-

likelihood we calculated 18 stepping stones using the methodology from Xie et al.

(2011). We calculated the difference between the log-marginal likelihoods of CID-3 and

MuSSE-3 models using the statistic κ defined as $\kappa = ln(P(X|CID - 3)) -$

$ln(P(X|MuSSE - 3))$, where X represents our state sample and the phylogenetic tree. If

κ has a value larger than 1, then the CID-3 model is preferred. The MuSSE-3 model is

preferred when $\kappa < -1$. The test statistic is inconclusive when κ has a value in the

interval (-1, 1).

300   *Diversification models with different assumptions about extinction rates*

Fossil records for passerines suggest rapid turnover of diversity over time, with

high species diversity and diversification rates (Jetz et al. 2012; Marshall 2017) despite

the extinctions of entire early passerine lineages (Hieronymus et al. 2019; Ksepka et al.

2019), high turnover of passerines within Europe between the Oligocene and the

present (Manegold 2008; Bochenski et al. 2021), and late Pleistocene species

extinctions (Steadman et al. 2015; Oswald and Steadman 2018). Diversification models

using phylogenetic trees with only extant taxa often underestimate the rates of

extinction (Höhna et al. 2011; Louca and Pennell 2021), producing errors in many

diversification analyses (Stadler 2013). This presents a potential problem in the

passerines because of evidence from fossils and other macroevolutionary studies

(Greenberg et al. 2021) that turnover has been high (see also Marshall's third law of

paleobiology: the average extinction rate approximately equals the average origination

rate; (Marshall 2017). Furthermore, classical paleontological work in extinction rates

(Raup 1992; May et al. 1995; Sepkoski 1998) has suggested that speciation rates are

only slightly larger than extinctions, and state-dependent diversification models have not

accounted for this potential assumption.

In a Bayesian framework, one way to incorporate knowledge about higher

extinction rates is by using a prior distribution that accommodates independent

information about the magnitude of the rates. Therefore, in our three-state, and hidden-

state models, we defined prior distributions for the extinction rates with elevated lower

bounds, and we systematically increased these lower bounds across model fits (Table

A1). We specified these lower bounds by making extinction rates' lower bound

dependent on speciation rates. Hence, in mathematical form, we defined this process

by using the linear function for extinction rates as $\mu_i = A * \lambda_i + \delta$ where $A =$

$0.7, 0.8, 0.9, 1, 1.1$, for all the states $i = D, C, H$ (Fig. 2d). We also explored the effect of

enhanced extinction in the MuHiSSE-3 model by allowing $A = 0.8$ (Fig. 2e). Defining the

extinction rate prior distribution as a linear combination of random variables allowed us

to fix the value A as the lower bound for the extinction fraction $a_i = \frac{\mu_i}{\lambda_i}$ (using use the

same definition of extinction fraction as Beaulieu and O'Meara 2016) while permitting

extra variability in the diversification rates so that extinction rates are not fully a

deterministic function of speciation rates. For example, an extinction rate of dome-

nesters that is at least 80% of the speciation rate is defined using the equation $\mu_D =$

$0.8 * \lambda_D + \delta$. In this example, the posterior estimation of the extinction fraction for state

Dome $(a_D = \frac{\mu_D}{\lambda_D})$ is forced to be at least 80%, but the extinction fraction a_D can still be

larger than the lower bound of 0.8 thanks to random variable δ, that is $a_D = \frac{\mu_D}{\lambda_D} > 0.8$. In

most of our models, prior distributions for extinction rates were defined equally (i.e.

$\mu_D = \mu_C = \mu_H = A * \lambda_D + \delta$). In one model, we specified $\mu_H < \mu_D < \mu_C$ by making $A =$

$0.6, 0.7$, and 0.8 respectively to verify that: 1) our inferences weren't driven by this

specification of the prior; and 2) that likelihood is informative given our dataset

(supplementary Fig. S12).

Overall, this choice in prior distributions allows us to restrict the parameter space

and find estimates of diversification assuming the extinction fraction is high. Linking

speciation and extinction rates does not change the dynamics of speciation and

extinction in SSE models because speciation and extinction parameters in birth and

death models are functionally linked by definition (Kendall 1948; Feller and Feller 1968;

Stadler and Bokma 2013). An in-depth discussion on how speciation and extinction

parameters in SSE models are functionally linked can be found in Helmstetter et al.

(2021).

350   *Ancestral state reconstruction using diversification-free models*

We estimated the most probable state at the root, since previous studies have

found that hole-nesting is the most likely the state at the root. The results showed that

nest type evolution is not linked to the diversification process (see Results section).

Therefore, we defined a three-state (Mk3, Fig. 2f) Markov model without diversification

parameters (hereafter, "diversification-free model" referring to the lack of diversification

parameters in the model) to reconstruct the ancestral nest type in passerines. We

calculated the marginal ancestral state reconstruction at each of the internal nodes.

Since our reconstructions use a Bayesian framework, in each of the nodes we plot the

state with the maximum *a posteriori* value of the marginal posterior distribution (Fig. 5),

as previously done in Freyman and Höhna (2018), and Zenil-Ferguson et al. (2019),

and implemented in RevBayes (Höhna et al. 2016). For all our models in Figure 2, we

assume the root values are stochastic, and are modeled using a Dirichlet probability

distribution with frequency parameter $(1/3,1/3,1/3)$ as the prior distribution (see the

number of parameters in Fig. 2). We calculated the posterior distribution for this

stochastic vector via our MCMC algorithms.

367   *Implementation of models and inferences*

368   All the diversification and diversification-free models in Fig. 2 were implemented

as graphical models in RevBayes software (Höhna et al. 2016). We customized and ran

a Markov chain Monte Carlo (MCMC; (Metropolis et al. 1953; Hastings 1970) algorithm

to sample the posterior distribution of each model. For all the models in Figure 2, we

assumed that the state value at the root was unknown, and we estimated the posterior

distribution of the frequencies at the root. Convergence and effective sample sizes of at

least 250 for every parameter in the MCMC were assessed using Tracer (Rambaut et

al. 2018) and two chains were run per model to verify convergence. Each MCMC run

was performed in the HiPerGator cluster at the University of Florida, and took an

average of 240 hours to converge. All our implementations are available at

https://github.com/roszenil/nestdivmodel.

380   RESULTS

381   *Diversification is not linked to nest type*

382   While our MuSSE-3 model found associations between nest type and

diversification rates, our MuHiSSE-3 model showed that these associations were

spurious (Fig. 3). This is because the addition of hidden states indicates that

diversification is driven by unmeasured factors other than nest type. In particular, for the

MuSSE-3 model of nest evolution, we found that hole nesters had faster net

diversification (defined as speciation minus extinction) than cup nesters, and the

smallest net diversification was associated with dome nesters (Fig. 3a). This result was

also true when the prior of extinction fraction was faster ($a = \mu/\lambda = 0.8$, Fig. 3a), and for

all elevated extinction fractions ($0 << a = \mu/\lambda < 1$) (model in Fig. 2d and estimation in

Fig. A1). When fitting the hidden state model MuHiSSE-3 (Fig. 3b), we found that the

three-state posterior distributions of net diversification rates completely overlap within

hidden states A and B while being different between A and B, indicating that the

diversification rate differences across the phylogenetic tree are due to hidden states

(Fig. 3b). These results were consistent when we allowed MuHiSSE-3 to have an

elevated extinction fraction of at least 0.8 (model in Fig. 2e and estimation in Fig. 3b).

Furthermore, when calculating Bayes factors comparing the CID-3 (diversification rates

independent of nest type, or null hypothesis) model against the MuSSE-3

(diversification rates depend on nest type), we found that the CID-3 model is preferred

over the MuSSE-3 ($\kappa = -6033.72 - (-7772.40) = 1738.67 > 1$). Altogether, there is

strong evidence that the heterogeneity observed in passerine diversification rates is not

linked to nest type.

a) Net diversification rates for SSE with nest type states

b) Net diversification rates for SSE with nest type states and hidden states

Figure 3. Net diversification rate estimates (speciation and extinction) for a) multistate dependent

diversification models (MuSSE-3), and b) multistate and hidden state dependent diversification

models (MuHiSSE-3). In models in (a) we consistently estimated that hole nesters have faster net

diversification rates than dome nesters, and that cup nesters had intermediate rate estimates.

This finding was consistent when we assumed the extinction fraction to be at least 0.8 (MuSSE-

3 model + prior extinction fraction > 0.8). The most noticeable change is that increasing the

extinction fraction (MuSSE-3 model+ extinction fraction > 0.8) decreases net diversification

estimates (y-axis), but the relative differences between states are maintained. For models in (b)

we observe that all diversification differences are due to the differences in hidden state (A<B)

rather than nest type (Dome=Cup=Hole for A and B states), which is strong evidence against the

direct influence of nest type on the speciation and extinction of passerines. Even increasing the

extinction fraction (MuHiSSE-3 + prior extinction fraction > 0.8) yields the same results.

417   *Evolving out of holes is faster than evolving back into them*

Since nest type is not linked to the diversification process we assessed the dynamics

under the diversification-free Mk3 model. The posterior distribution of transition rates

from the Mk3 model shows that the rate of transitioning from hole to cup was the fastest

(mean $q_{32} = 9.1x10^{-3}$, 95% credible interval $(6.5x10^{-3}, 0.01)$, followed by the transition

from hole to dome (mean $q_{31} = 6.5x10^{-3}$, 95% credible interval $(3.7x10^{-3}, 8.2x10^{-3})$,

Fig. 4). Transition rates from either dome or cup to hole nests were similar, and both

were slower than the transition rates out of hole nests (dome to hole transition rates

have a mean $q_{13} = 1.6x10^{-3}$, and 95% credible interval $(6.6x10^{-4}, 2.8x10^{-3})$; cup to

hole transition rates have a mean $q_{23} = 1.2x10^{-3}$, credible interval $(7.3x10^{-4}, 2.5x10^{-3})$,

shown in Fig. 4.

Figure 4. Transition rates between nest states for diversification-free models in the model with

three states (Mk3). The posterior distribution for the rate from hole to cup (q_{32}) is faster than

any other rate to and from hole. We found that evolving out of hole is faster than into hole, with

hole to cup faster than hole to dome.

Hole-nesting is the most probable state at the root

The Mk3 (Fig. 2f) model reconstructs hole-nesting as the state with the highest

probability at the most recent common ancestor of the Passeriformes. We found that the

maximum *a posteriori* for the root's marginal posterior probability was 0.77 for hole-

nesting when the tree included the parrot *Strigops habroptila* as the outgroup. This

probability decreased to 0.51 when the tree only contained passerines and no outgroup

(states hole and cup make the other 0.49 of probability). The posterior distribution for

the stochastic vector at the root had as its maximum the vector (Dome=0.30, Cup=0.25,

Hole=0.45; Figure 5).

Figure 5. Ancestral state reconstruction for nest type using a Markov model with three states

(Mk3). The size of a node represents the maximum probability value of the marginal posterior

distribution of the node. The color of the node represents the nest type associated with the

maximum posterior probability. This tree includes the hole-nesting *Strigops habroptila* as the

outgroup. Figure made using RevGadgets (Tribble et al. 2022)**.**

450   DISCUSSION

Over the years, there have been diverse perspectives on the link between

ecological specialization and evolutionary diversification (Vamosi et al. 2014). On the

one hand, ecological specialization has been viewed as an evolutionary dead end

because it leads to evolutionary changes in traits that are difficult to reverse, which can

leave specialized taxa at higher risk of extinction when conditions change. On the other

hand, other studies have emphasized the links between specialization and increased

diversification (Losos et al. 1994; Schluter 2000) because a narrower niche could

increase diversification either directly, if niche shifts are associated with multiple

speciation events (Yoder et al. 2010, Velasco et al. 2016), or indirectly, because of

differences in dispersal, connectivity, population persistence, and/or range size of

specialist and generalist species (Greenberg and Mooers 2017; Harvey et al. 2019).

Testing the outcomes of specialization on diversification rates has been difficult

because macroevolutionary models require datasets with a large number of

independent origins of specialization (Day et al. 2016).

Here, using state-dependent diversification models in a large tree of passerines,

we found that hole-nesting specialization does not differ from cup and dome-nesting in

diversification rates. We also found that transition rates into hole-nesting were low

compared to transitions out of hole-nesting (Fig. 4), and that the most probable state for

the root of all passerines was hole-nesting. These three results highlight the lack of

support for the hypothesis that specialization leads to an evolutionary dead end. Our

results, combined with a number of recent studies addressing this hypothesis (Day et al.

2016; Cyriac and Kodandaramaiah 2018), raise the question of whether the link

between diversification rates and ecological specialization is truly variable across taxa

and traits, or whether newer phylogenetic comparative methods and larger data sets are

allowing us to test hypotheses more rigorously. Interestingly, in our study, which uses a

large dataset with numerous transitions to the specialized hole-nesting state, we found

that, in the absence of accounting for hidden states, adoption of hole-nesting behavior

led to higher, not lower, diversification rates (Fig. 3a). However, when we included

hidden states, and thus accounted for the possibility that other, unmeasured variables

are driving the relationship, we found no difference in diversification rates across the

three nesting types (Fig. 3b). This comparison emphasizes that phylogenetic studies of

specialization that do not account for hidden states may need to reevaluate their null

hypotheses of diversification. For example, a recent study by Medina et al. (2022)

concluded that there are differences in extinction rates between cup and dome nesters,

but hidden state models in this study were not presented due to failed convergence, and

extinction rates cannot be interpreted independently from speciation rates (Helmstetter

et al. 2021), so the link between nest type and diversification remained untested. Our

results, in combination with recent tests of the evolutionary dead end hypothesis, show

that there is growing evidence that specialization rarely leads to an evolutionary dead

end (Stern et al. 2017; Villastrigo et al. 2020).

492   *Modeling the diversification consequences of specialization*

State-dependent diversification models are a flexible tool to test whether

specialization is linked to diversification rates. A recent ongoing discussion in the field of

macroevolution centers on the impossibility of estimating speciation and extinction rates

from phylogenetic trees with only extant taxa using birth and death stochastic models

with time-dependent parameters (Louca and Pennell 2020). The non-identifiability of

parameters in the time-dependent diversification models, and the possibility of multiple

congruent likelihoods across time-dependent models may affect the state-dependent

diversification models presented here. State-dependent diversification models are

simply a special case of the time-dependent models explored in Louca and Pennell

(2020). However, taking into consideration the nest type, and using informative priors to

represent different extinction scenarios, shows the potential for recovering the direction

of differences between states, even if point estimates for speciation and extinction rates

are not the same across these scenarios (Fig. 4, and Fig. A1), as suggested by Louca

and Pennell (2020). Our main findings under the assumptions of enhanced extinction

suggest that the relative differences among states are possible to infer despite the non-

identifiability between different extinction histories. Therefore, it is important when

applying state-dependent diversification models to clarify whether the goal is to obtain

point estimates for speciation or extinction rates, or to find relative differences in the

history of diversification linked to the states. In this study, we were interested in the

latter, and our results suggest that finding relative differences of diversification between

states can be performed in a systematic fashion with consistent results.

Overall, we were able to rigorously test different hypotheses and extinction

scenarios because of the size of the tree, the number of transitions between nest types,

and the large proportion of passerines in the sample (> 51%). Studies of specialization

linked to diversification might find spurious associations when independent transitions to

and from the specialized state are too few (Uyeda et al. 2018), tree size is not large

enough (Davis et al. 2013), sampling fraction is small (Chang et al. 2020), or when

hidden state models are not incorporated to test for the significance of trait-dependent

diversification (Beaulieu and O'Meara 2016; Caetano et al. 2018). Historically,

standards for sample quality and interpretation of the parameters as a joint function of

speciation and extinction in SSE models have been lacking (Helmstetter et al. 2021), so

it is important that future studies of state-dependent diversification be aware of these

issues.

526   *Ecological and evolutionary consequences of specialization*

527   The ecological consequences of specialization can be diverse – specialization

can, in some cases, decrease competition for resources and, in other cases, increase it.

For example, many studies have focused on diet or host specialization, which frequently

reduce competition over food or breeding resources (Vamosi et al. 2014). However,

specialization can also enable species to escape predation (e.g., Singer et al. 2019)

and, in such cases, may actually increase competition for scarce, protective resources.

Such is the case for hole-nesting birds, where an important consequence of hole-

nesting is that it strongly reduces the risk of nest predation, an important driving force

behind life history evolution in birds (Martin 1995). In our study, we were able to

distinguish between the influence of ecological specialization per se versus one specific

ecological consequence of specialization – reduced predation. Dome-nesting birds also

show decreased predation rates (Oniki 1979b; Linder and Bollinger 1995; Auer et al.

2007; Martin et al. 2017), but they are relatively unspecialized in nesting substrates

compared to hole-nesters. However, neither hole-nesting nor dome-nesting were

associated with elevated diversification rates. This suggests that, in passerine birds,

reduced nest predation rates do not have a strong influence on diversification dynamics

and, while escaping predation can often lead to ecological release (Herrmann et al.

2021), this does not necessarily translate into increased ecological opportunity or

subsequent adaptive diversification. Moreover, reduced predation can also increase

species' ability to persist and, hence, decrease extinction risk. Yet, a change in

abundance of nesting holes, such as declines in tree cavities, has been shown to lead

to heightened risk of extinction (Duckworth and Badyaev 2007). Thus, the lack of a

relationship with diversification rates may reflect a balance between these various

ecological consequences of adopting hole-nesting.

Another factor that may account for inconsistency across studies in the links

between specialization and macroevolutionary dynamics is the extent to which

specialization results in further evolution of specific traits. Ecological specialization is a

narrowing of an organism's niche, and so results from a change in how an organism

interacts with its environment. Futuyma and Moreno (1988) point out that specialization,

at a minimum, only requires a behavioral shift, and specialized taxa vary in the extent

that there is subsequent evolution of other traits. Thus, it is possible that whether

specialization is an evolutionary dead end or not may be strongly linked to the extent of

secondary adaptation that follows it. In the case of hole-nesting birds, specialization is

largely a behavioral shift, although it does lead to evolutionary changes in many life

history and breeding traits (see below). Thus, it may be that hole-nesting specialization

does not lead to an evolutionary dead end because of the relative ease of reversing

these traits (Fig. 4). The low rate of evolutionary transitions into, relative to out of, hole-

nesting (Fig. 4) provides further evidence against the evolutionary dead end hypothesis

and instead suggests that evolving ecological specialization can be challenging. In the

case of hole-nesters, becoming ecologically specialized involves facing increased

interspecific competition, as holes are sought not only by other birds but also by other

vertebrates (Newton 1994). Thus, competition may prevent evolutionary transitions

towards specialization. Evolutionary transitions from hole-nesting to open-cup nesting,

which occur at a relatively high rate, involve different evolutionary challenges, such as

life history specialization. Open-cup nesting species have exceptionally high nestling

growth rates, which appear to be an adaptation to increased predation risk (Ricklefs

1979). Moving from the relaxed selection regime on growth rates associated with hole-

nesting to the strong selection regime on growth rates in open-cup nests would seem to

act as a filter that would limit this transition. However, our results show that, at least in

passerine birds, evolving such life history shifts is easier than evolving the ability to deal

with the intense competition of hole-nesting.

We also found evidence that it was easier for birds to transition from hole to cup

rather than hole to dome. Most hole nesters build an open-cup style nest, rather than a

dome, within the cavity (Price and Griffith 2017). Our finding that transitions to open-cup

nesting were easier compared to transitions to dome may simply reflect that transitions

from hole-nesting more often occurred in taxa that were already building open-cup nests

within their holes. Thus, for most hole nesters, transitioning to dome-nesting may be

more difficult because it would require two steps: leaving cavities and changing how the

nest is built. While in the minority, there are several clades, particularly at the base of

the passerine tree, that build dome style nests within cavities (Price and Griffith 2017).

Future studies comparing transitions to dome and open-cup nests based on whether

hole-nesting species are already building a dome or cup style nest within their cavity

would add further insight into the mechanisms behind these different transition rates.

590   *Ancestral State Reconstruction*

In all our ancestral state reconstructions we found that hole-nesting is the most

probable state at the root of the passerines. When including the parrot *Strigops*

habroptilus as the outgroup*,* the maximum of the marginal posterior distribution of the

node had a probability of 0.77 for hole-nesting. Without this outgroup, we still recovered

hole-nesting with a maximum a posteriori of 0.51 (see Figure S12 in supplementary

information). These two reconstructions represent significant evidence in favor of the

specialized state hole-nesting at the origin of passerines. Previous work has

emphasized that hole-nesting, dome-nesting, and open-cup nesting all appear to have

occurred early in the history of passerines (Collias 1997; Price and Griffith 2017; Fang

et al. 2018; McEntee et al. 2018), with Collias (1997) suggesting that the nest type of

the earliest passerine might be unknowable because of the apparent rapid evolution of

nest type in early passerines. Our ancestral state reconstruction, using an approach

where we included strong species-level sampling and assessed whether differing

diversification rates had to be accounted for (Maddison 2006), tips the balance in favor

of hole-nesting as the ancestral state for the common ancestor of extant passerines.

Notably, this result contrasts with the ancestral state reconstruction of Fang et al.

(2018), which included more sampling outside of passerines and less species-level

sampling within passerines. This result might seem counterintuitive if the definition of

specialization is confounded with the concept of a "derived state". In the original

definition of specialization by Futuyma and Moreno (1988) it is stated that specialization

is based on ecology and function in an ecosystem; therefore, we shouldn't expect that a

specialized state is a derived one. In our study, support for hole-nesting as the ancestral

state serves to underscore how radically the evolution of hole-nesting as a

specialization fails to meet the generalists-to-specialists view of evolution; rather than a

dead end, all passerine diversity emerged from a specialist ancestor.

617   CONCLUSIONS

Our macroevolutionary analysis failed to find any link between specialization of

nest type and diversification rates under different extinction scenarios, suggesting that

there is little support for the evolutionary dead end hypothesis in this case study.

Moreover, contrary to this hypothesis, we also found that transitions from the

specialized state were relatively easy compared to transitions toward the more

generalist states and that the root state of passerines is most probably hole-nesting.

Our results suggest that the ecological consequences of resource specialization,

whether due to escape from competition or predation, might be key to understanding its

macroevolutionary consequences. This work adds to other recent studies that have

found little support for the evolutionary dead end idea, suggesting that evolution of

resource specialization is more evolutionarily labile than previously thought. We suggest

that future studies of this question would benefit from explicit comparison of resource

specialization that vary in their ecological consequences, as well as the extent of trait

evolution necessary for specialization. Such studies would enable a greater

understanding of the mechanisms that underlie variable links between specialization

and macroevolutionary dynamics.

635   ACKNOWLEDGEMENTS

636   We are grateful for Dr. Sonal Singhal who offered feedback on an initial draft of this

637 work, Dr. Carrie Tribble who helped with RevGadgets visualizations, and Jenna McCullough for

638 the illustration of nests. This work started and concluded during the toughest time of the COVID-

639 19 emergency, slow progress was made during two years thanks to the support of many of our

640 colleagues and families. RZF is supported by NSF-DEB 2055466.

642   LITERATURE CITED

Alhajeri B.H., Steppan S.J. 2018. Ecological and ecomorphological specialization are
not associated with diversification rates in muroid rodents (Rodentia: Muroidea).
Evol. Biol. 45:268–286.

Auer S.K., Bassar R.D., Joseph J., Martin T.E. 2007b. Breeding biology of passerines in
a subtropical montane forest in Northwestern Argentina. Condor. 109:321–333.

Baiser B., Valle D., Zelazny Z., Gordon Burleigh J. 2018. Non-random patterns of
invasion and extinction reduce phylogenetic diversity in island bird assemblages.
Ecography. 41:361–374.

Beaulieu J.M., O'Meara B.C. 2016. Detecting Hidden Diversification Shifts in Models of
Trait-Dependent Speciation and Extinction. Syst. Biol. 65:583–601.

Beaulieu J., O'Meara B., Caetano D., Boyko J., Vasconcelos T. 2021. Package "hisse."
.

Bernays E.A. 1989. Host range in phytophagous insects: the potential role of generalist
predators. Evol. Ecol. 3:299–311.

Birand A., Vose A., Gavrilets S. 2012. Patterns of species ranges, speciation, and
extinction. Am. Nat. 179:1–21.

Bochenski Z.M., Tomek T., Bujoczek M., Salwa G. 2021. A new passeriform (Aves:
Passeriformes) from the early Oligocene of Poland sheds light on the beginnings of

Suboscines. J. Ornithol.

Burleigh J.G., Kimball R.T., Braun E.L. 2015. Building the avian tree of life using a
large-scale, sparse supermatrix. Mol. Phylogenet. Evol. 84:53–63.

Caetano D.S., O'Meara B.C., Beaulieu J.M. 2018. Hidden state models improve state-
dependent diversification approaches, including biogeographical models. Evolution.
72:2308–2324.

Chang J., Rabosky D.L., Alfaro M.E. 2020. Estimating Diversification Rates on
Incompletely Sampled Phylogenies: Theoretical Concerns and Practical Solutions.
Syst. Biol. 69:602–611.

Cockle K.L., Martin K., Wesołowski T. 2011. Woodpeckers, decay, and the future of
cavity-nesting vertebrate communities worldwide. Front. Ecol. Environ. 9:377–382.

Collias N.E. 1997. On the origin and evolution of nest building by passerine birds.
Condor. 99:253–270.

Collias N.E., Collias E.C. 1984. Nest building and bird behavior. Princeton University
Press.

Conway M., Olsen B.J. 2019. Contrasting drivers of diversification rates on islands and
continents across three passerine families. Proc. Biol. Sci. 286:20191757.

Crouch N.M.A., Ricklefs R.E. 2019. Speciation Rate Is Independent of the Rate of
Evolution of Morphological Size, Shape, and Absolute Morphological Specialization
in a Large Clade of Birds. Am. Nat. 193:E78–E91.

Cyriac V.P., Kodandaramaiah U. 2018. Digging their own macroevolutionary grave:
fossoriality as an evolutionary dead end in snakes. J. Evol. Biol. 31:587–598.

Davis M.P., Midford P.E., Maddison W. 2013. Exploring power and parameter
estimation of the BiSSE method for analyzing species diversification. BMC Evol.
Biol. 13:38.

Day E.H., Hua X., Bromham L. 2016. Is specialization an evolutionary dead end?
Testing for differences in speciation, extinction and trait transition rates across
diverse phylogenies of specialists and generalists. J. Evol. Biol. 29:1257–1267.

Del Hoyo J., Elliott A., Sargatal J., Christie D.A., de Juana E. 2017. Handbook of the
birds of the world alive. Lynx Edicions, Barcelona, Spain.

Duckworth R.A., Badyaev A.V. 2007. Coupling of dispersal and aggression facilitates
the rapid range expansion of a passerine bird. Proc. Natl. Acad. Sci. U. S. A.
104:15017–15022.

Fang Y.-T., Tuanmu M.-N., Hung C.-M. 2018. Asynchronous evolution of

interdependent nest characters across the avian phylogeny. Nat. Commun. 9:1863.

Feller W., Feller V. 1968. An Introduction to Probability Theory and Its Applications,
Volume 1. Wiley.

FitzJohn R.G. 2012. Diversitree : comparative phylogenetic analyses of diversification in
R. Methods Ecol. Evol. 3:1084–1092.

FitzJohn R.G., Maddison W.P., Otto S.P. 2009. Estimating trait-dependent speciation
and extinction rates from incompletely resolved phylogenies. Syst. Biol. 58:595–
611.

Freyman W.A., Höhna S. 2018. Cladogenetic and Anagenetic Models of Chromosome
Number Evolution: A Bayesian Model Averaging Approach. Syst. Biol. 67:195–215.

Freyman W.A., Höhna S. 2019. Stochastic Character Mapping of State-Dependent
Diversification Reveals the Tempo of Evolutionary Decline in Self-Compatible
Onagraceae Lineages. Syst. Biol. 68:505–519.

Futuyma D.J., Moreno G. 1988. The evolution of ecological specialization. Annu. Rev.
Ecol. Syst. 19:207–233.

Gavrilets S., Li H., Vose M.D. 2000. Patterns of parapatric speciation. Evolution.
54:1126–1134.

Greenberg D.A., Mooers A.Ø. 2017. Linking speciation to extinction: Diversification
raises contemporary extinction risk in amphibians. Evolution Letters. 1:40–48.

Greenberg D.A., Pyron R.A., Johnson L.G.W., Upham N.S., Jetz W., Mooers A.Ø. 2021.
Evolutionary legacies in contemporary tetrapod imperilment. Ecol. Lett. 24:2464–
2476.

Harvey M.G., Singhal S., Rabosky D.L. 2019. Beyond Reproductive Isolation:
Demographic Controls on the Speciation Process. Annual Review of Ecology,
Evolution, and Systematics. 50:75–95.

Hastings W.K. 1970. Monte Carlo sampling methods using Markov chains and their
applications. Biometrika. 57:97–109.

Helmstetter A.J., Glemin S., Käfer J., Zenil-Ferguson R., Sauquet H., de Boer H.,
Dagallier L.-P.M.J., Mazet N., Reboud E.L., Couvreur T.L.P., Condamine F.L. 2021.
Pulled Diversification Rates, Lineages-Through-Time Plots and Modern
Macroevolutionary Modelling. Syst. Biol. 71(3):758-763

Herrmann N.C., Stroud J.T., Losos J.B. 2021. The evolution of "ecological release" into
the 21st century. Trends Ecol. Evol. 36:206–215.

Hieronymus T.L., Waugh D.A., Clarke J.A. 2019. A new zygodactylid species indicates

the persistence of stem passerines into the early Oligocene in North America. BMC
Evol. Biol. 19:3.

Höhna S., Landis M.J., Heath T.A., Boussau B., Lartillot N., Moore B.R., Huelsenbeck
J.P., Ronquist F. 2016. RevBayes: Bayesian Phylogenetic Inference Using
Graphical Models and an Interactive Model-Specification Language. Syst. Biol.
65:726–736.

Höhna S., Stadler T., Ronquist F., Britton T. 2011. Inferring speciation and extinction
rates under different sampling schemes. Mol. Biol. Evol. 28:2577–2589.

Jetz W., Thomas G.H., Joy J.B., Hartmann K., Mooers A.O. 2012. The global diversity
of birds in space and time. Nature. 491:444–448.

Jordan M.I. 2004. Graphical Models. SSO Schweiz. Monatsschr. Zahnheilkd. 19:140–
155.

Kass R.E., Raftery A.E. 1995. Bayes Factors. J. Am. Stat. Assoc. 90:773–795.

Kendall D.G. 1948. On the Generalized "Birth-and-Death" Process. Ann. Math. Stat.
19:1–15.

Kilner R.M. 2006. The evolution of egg colour and patterning in birds. Biol. Rev. Camb.
Philos. Soc. 81:383–406.

Ksepka D.T., Grande L., Mayr G. 2019. Oldest Finch-Beaked Birds Reveal Parallel
Ecological Radiations in the Earliest Evolution of Passerines. Curr. Biol. 29:657–
663.e1.

Linder E.T., Bollinger E.K. 1995. Depredation of artificial Ovenbird nests in a forest
patch. Wilson Bull.

Losos J.B., Irschick D.J., Schoener T.W. 1994. ADAPTATION AND CONSTRAINT IN
THE EVOLUTION OF SPECIALIZATION OF BAHAMIAN ANOLIS LIZARDS.
Evolution. 48:1786–1798.

Louca S., Pennell M.W. 2020. Extant timetrees are consistent with a myriad of
diversification histories. Nature. 580:502–505.

Louca S., Pennell M.W. 2021. Why extinction estimates from extant phylogenies are so
often zero. Current Biology. 31(14): 3168-3173.

Maddison W.P. 2006. Confounding asymmetries in evolutionary diversification and
character change. Evolution. 60:1743–1746.

Manegold A. 2008. Passerine diversity in the late Oligocene of Germany: earliest
evidence for the sympatric coexistence of Suboscines and Oscines. Ibis . 150:377–
387.

Marshall C.R. 2017. Five palaeobiological laws needed to understand the evolution of
the living biota. Nature ecology & evolution. 1:165.

Martin T.E. 1993. Nest predation and nest sites. Bioscience. 43:523.

Martin T.E. 1995. Avian life history evolution in relation to nest sites, nest predation, and
food. Ecol. Monogr. 65:101–127.

Martin T.E., Boyce A.J., Fierro-Calderón K., Mitchell A.E., Armstad C.E., Mouton J.C.,
Bin Soudi E.E. 2017. Enclosed nests may provide greater thermal than nest
predation benefits compared with open nests across latitudes. Funct. Ecol.
31:1231–1240.

Martin T.E., Li P. 1992. Life History Traits of Open- vs. Cavity-Nesting Birds. Ecology.
73:579–592.

May, Lawton, Stork. 1995. Assessing extinction rates. Extinction rates.

McEntee J.P., Zelazny Z., Burleigh J.G. 2018. The macroecology of passerine nest
types, in the light of macroevolution. bioRxiv.

McEntee J.P., Zelazny Z., Burleigh J.G. 2021. On the relationship between nest
concealment and the conspicuous coloration of female passerines: a test of
Wallace's hypothesis. Revista Ecuatoriana de Medicina y Ciencias Biológicas:139–
156.

Medina I., M Perez D., Silva A.C.A., Cally J., León C., Maliet O., Quintero I. 2022. Nest
architecture is linked with ecological success in songbirds. Ecol. Lett. 25:1365-
1375.

Metropolis N., Rosenbluth A.W., Rosenbluth M.N., Teller A.H., Teller E. 1953. Equation
of State Calculations by Fast Computing Machines. J. Chem. Phys. 21:1087–1092.

Newton I. 1994. The role of nest sites in limiting the numbers of hole-nesting birds: a
review. Biol. Conserv. 70:265–276.

Nice M.M. 1957. Nesting success in altricial birds. Auk. 74:305–321.

Oniki Y. 1979a. Is nesting success of birds low in the tropics? Ornithol. Monogr. 11:60–
69.

Oniki Y. 1979b. Is Nesting Success of Birds Low in the Tropics? Biotropica. 11:60–69.

Oswald J.A., Steadman D.W. 2018. The late Quaternary bird community of New
Providence, Bahamas. The Auk: Ornithological Advances. 135:359–377.

Otero A., Jiménez-Mejías P., Valcárcel V., Vargas P. 2019. Being in the right place at
the right time? Parallel diversification bursts favored by the persistence of ancient

epizoochorous traits and hidden factors in Cynoglossoideae. Am. J. Bot. 106:438–
452.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics. 35:526–528.

Price J.J., Griffith S.C. 2017. Open cup nests evolved from roofed nests in the early
passerines. Proc. Biol. Sci. 284.

Rabosky D.L., Goldberg E.E. 2015. Model inadequacy and mistaken inferences of trait-
dependent speciation. Syst. Biol. 64:340–355.

Rabosky D.L., Goldberg E.E. 2017. FiSSE: A simple nonparametric test for the effects
of a binary character on lineage diversification rates. Evolution. 71:1432–1442.

Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior
Summarization in Bayesian Phylogenetics Using Tracer 1.7. Syst. Biol. 67:901–
904.

Raup D.M. 1992. Extinction: bad genes or bad luck?. WW Norton & Company.
810

Ricklefs R.E. 1969. An analysis of nesting mortality in birds. Smithson. Contrib. Zool.
9:1–48.

Ricklefs R.E. 1979. Adaptation, constraint, and compromise in avian postnatal
development. Biol. Rev. Camb. Philos. Soc. 54:269–290.

Sanderson M.J. 2003. r8s: inferring absolute rates of molecular evolution and
divergence times in the absence of a molecular clock. Bioinformatics. 19:301–302.

Schluter D. 2000. The Ecology of Adaptive Radiation. Oxford, UK: Oxford University
Press.

Sepkoski J.J. Jr. 1998. Rates of speciation in the fossil record. Philos. Trans. R. Soc.
Lond. B Biol. Sci. 353:315–326.

Sexton J.P., Montiel J., Shay J.E., Stephens M.R., Slatyer R.A. 2017. Evolution of
Ecological Niche Breadth. Annu. Rev. Ecol. Evol. Syst. 48:183–206.

Singer M.S., Clark R.E., Johnson E.R., Lichter-Marck I.H., Mooney K.A., Whitney K.D.
2019. Dietary specialization is conditionally associated with increased ant predation
risk in a temperate forest caterpillar community. Ecol. Evol. 9:12099–12112.

Stadler T. 2013. How can we improve accuracy of macroevolutionary rate estimates?
Syst. Biol. 62:321–329.

Stadler T., Bokma F. 2013. Estimating speciation and extinction rates for phylogenies of

higher taxa. Syst. Biol. 62:220–230.

Steadman D.W., Oswald J.A., Rincón A.D. 2015. The diversity and biogeography of late
Pleistocene birds from the lowland Neotropics. Quat. Res. 83:555–564.

Stern D.B., Breinholt J., Pedraza-Lara C., López-Mejía M., Owen C.L., Bracken-
Grissom H., Fetzner J.W. Jr, Crandall K.A. 2017. Phylogenetic evidence from
freshwater crayfishes that cave adaptation is not an evolutionary dead-end.
Evolution. 71:2522–2532.

Tonini J.F.R., Ferreira R.B., Pyron R.A. 2020. Specialized breeding in plants affects
diversification trajectories in Neotropical frogs. Evolution. 74:1815–1825.

Tribble C.M., Freyman W.A., Landis M.J., Lim J.Y., Barido-Sottani J., Kopperud B.T.,
Höhna S., May M.R. 2022. RevGadgets: An R package for visualizing Bayesian
phylogenetic analyses from RevBayes. Methods Ecol. Evol. 13:314–323.

Uyeda J.C., Zenil-Ferguson R., Pennell M.W. 2018. Rethinking phylogenetic
comparative methods. Syst. Biol. 67:1091–1109.

Vamosi J.C., Armbruster W.S., Renner S.S. 2014. Evolutionary ecology of
specialization: insights from phylogenetic analysis. Proc. Biol. Sci. 281:20142004.

Velasco, J. A., E. Martínez-Meyer, O. Flores-Villela, A. García, A. C. Algar, G. Köhler,
and J. M. Daza. 2016. Climatic niche attributes and diversification in Anolis lizards.
Journal of Biogeography **43**:134-144.
848
Villastrigo A., Abellán P., Ribera I. 2021. Habitat preference and diversification rates in a
speciose lineage of diving beetles. Mol. Phylogenet. Evol. 159:107087.

Villastrigo A., Arribas P., Ribera I. 2020. Irreversible habitat specialization does not
constrain diversification in hypersaline water beetles. Mol. Ecol. 29:3637–3648.

Wallace A.R. 1868. A theory of birds' nests: showing the relation of certain sexual
differences of colour in birds to their mode of nidification. Journal of Travel and
Natural History. 1:73–89.

Xie W., Lewis P.O., Fan Y., Kuo L., Chen M.-H. 2011. Improving marginal likelihood
estimation for Bayesian phylogenetic model selection. Syst. Biol. 60:150–160.

Yoder J.B., Clancey E., Des Roches S., Eastman J.M., Gentry L., Godsoe W., Hagey
T.J., Jochimsen D., Oswald B.P., Robertson J., Sarver B.A.J., Schenk J.J., Spear
S.F., Harmon L.J. 2010. Ecological opportunity and the origin of adaptive
radiations. J. Evol. Biol. 23:1581–1596.

Zenil-Ferguson R., Burleigh J.G., Freyman W.A., Igić B., Mayrose I., Goldberg E.E.
2019. Interaction among ploidy, breeding system and lineage diversification. New
Phytol. 224:1252–1265.

Zenil-Ferguson R., Burleigh J.G., Ponciano J.M. 2018. chromploid: An R package for
chromosome number evolution across the plant tree of life. Appl. Plant Sci.
6:e1037.

Table A1. Assumptions of extinction in different MuSSE-3 models.

Assumption Extinction fraction	Extinction as a function of speciation	Prior distributions used*	Interpretation
1. NA (Fig. 2a)	NA	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\mu_i \sim LogN(\mu_\mu, \sigma_\mu)$	Speciation and extinction freely estimated
2. $\mu_i/\lambda_i \geq 0.7$ with $i = C, D, H$ (Fig. 2d)	$\mu_i = 0.7\,\lambda_i + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	Extinction rate is at least 0.7 of speciation rate
3. $\mu_i/\lambda_i \geq 0.8$ with $i = C, D, H$ (Fig. 2d)	$\mu_i = 0.8\,\lambda_i + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	Extinction rate is at least 0.8 of speciation rate
4. $\mu_i/\lambda_i \geq 0.9$ with $i = C, D, H$ (Fig. 2d)	$\mu_i = 0.9\,\lambda_i + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	Extinction rate is at least 0.9 of speciation rate
5. $\mu_i/\lambda_i \geq 1.0$ with $i = C, D, H$ (Fig. 2d)	$\mu_i = \lambda_i + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	Extinction rate is greater or equal to speciation rate
6. $\mu_i/\lambda_i \geq 1.1$ with $i = C, D, H$ (Fig. 2d)	$\mu_i = 1.1\,\lambda_i + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	Extinction rate is at least 1.1 of speciation rate
7. $\mu_C/\lambda_C \geq 0.8$ $\mu_D/\lambda_D \geq 0.7$ $\mu_H/\lambda_H \geq 0.6$ (Results in supplementary information Fig. S12)	$\mu_C = 0.8\,\lambda_C + \delta$ $\mu_D = 0.7\,\lambda_D + \delta$ $\mu_H = 0.6\,\lambda_H + \delta$	$\lambda_i \sim LogN(\mu_\lambda, \sigma_\lambda)$ $\delta \sim Exp(2)$ $i = C, D, H$	For cup, extinction rate is at least 0.8, for dome, is at least 0.7, and for hole, extinction rate is at least 0.6 of speciation rate.

*Prior distributions for speciation parameters are log-Normal with parameters $\left(\mu_\lambda = ln\left(ln\left(\frac{ln(3225/2)}{Tree\ Age}\right)\right), \sigma_\lambda = 1.12\right)$. For parameter ($\delta$) we chose an exponential with rate 2. Other distributions were tested for δ and results were unaltered.

In Figure A1 we summarize the results of assumptions 1-6 from table A1. Here we can

observe that when it is assumed that the extinction fraction is greater than 0.7 but less

than 1, we recover the differences between states (Dome < Cup < Hole). When the

extinction fraction is at least 1 and the extinction rate is expected to be about as large

as speciation, we found no differences in the posterior distributions of the three states.

Finally, when the extinction rate is assumed to be larger than the speciation rate we

obtain the reverse effect (Hole < Cup < Dome). These results show that despite our

prior assumptions about extinction rate, the hole-nesting taxa will have a greater

influence on the direction of net diversification than cup or dome nesters. And finally, the

state dome always has the smallest effect on net diversification in absolute value.

These results might only be applicable to a trait that does not influence diversification,

as we have shown using hidden state and character independent models for nest type

(Figure 2). Whether this pattern of finding similar differences in diversification is

sustained in the face of enhanced extinction holds for all traits in SSE models remains

to be tested.

Figure A1. Different *a priori* assumptions about the minimum of extinction fraction ($a =$

$\frac{\mu}{\lambda}$). When speciation and extinction are estimated freely ($a > 0$), the posterior

distributions for net diversification show that the rate of diversification for hole is faster

than cup, and both are faster than dome. This inference is supported even when we

assume that extinction rate should be faster (in other words for cases where $a >$

$0.7, a > 0.8, a > 0.9$ for all states) but still smaller than speciation; however, this

assumption changes the overall scale leading to smaller net diversification values. The

inference from net diversification starts changing when we assumed $a > 1$, that is, that

extinction is at least as fast as speciation for every state, where we found that the

posterior distributions of net diversifications for every state overlap. Once we assumed

that extinction is much faster than speciation ($a > 1.1$), we found a mirror trend in the

inference with hole net diversification being the slowest, and dome being the fastest.